1 **EpigeneticAgePipeline: an R package for comprehensive assessment of epigenetic age metrics from methylation microarrays**

2 Stanislav Rayevskiy[1], Quinn Le[1], Julia Nguyen[1], Steven Q. Chen[1], Christina A. Castellani[1,2,3,4]*

3

4 [1] Department of Pathology and Laboratory Medicine, Schulich School of Medicine and Dentistry, Western University, London, Ontario, Canada

5 [2] Department of Epidemiology and Biostatistics, Schulich School of Medicine and Dentistry, Western University, London, Ontario, Canada

6 [3] McKusick-Nathans Institute, Department of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, United

7 States of America

8 [4] Children's Health Research Institute, Lawson Research Institute, London, Ontario, Canada

9 * Corresponding author

10 Email: christina.castellani@schulich.uwo.ca

11

12 **Abstract**

13 Epigenetic age is a biological age estimate based on nuclear DNA methylation patterns. Epigenetic clocks measure biological age by

14 analyzing predictable changes in DNA methylation sites associated with aging. This study introduces EpigeneticAgePipeline, an R package that

15 streamlines the estimation of epigenetic age metrics including Horvath, Horvath skin and blood, Hannum, PhenoAge/Levine, GrimAge (V1, and

16 V2), and DunedinPACE plus additional acceleration metrics based on all other clocks. Quality control includes detection p-value filtering

17 (sample- and probe-level), bead-count thresholds, and Illumina quality control intensity checks. EpigeneticAgePipeline supports Illumina

18 Infinium methylation microarrays (HumanMethylation27, HumanMethylation450, HumanMethylationEPIC/EPICv2, and Human Methylation

19 Screening Array). It offers functionalities including data preprocessing, normalization, cell count imputation, residual generation accounting for

20 principal components and batch effects, and extensive visualizations for improved interpretability. Validation was performed using GEO dataset

21 GSE237561, confirming the accuracy of the pipeline. EpigeneticAgePipeline provides an integrated workflow from raw data to advanced

22 statistical analyses and visualizations, improving usability over existing tools. In addition to the traditional clocks mentioned, the package also

23 integrates a set of additional epigenetic age clocks (PedBE, Wu, TL, BLUP, and EN). Future updates will include emerging epigenetic age

24 measures to maintain relevance in this evolving field.

25 **Keywords:** Epigenetic Age, DNA Methylation, Epigenetic Clocks, Biological Aging, Bioinformatics Pipeline, R Package,

26 Methylation Microarrays

27 **Background**

28 The field of epigenomics has allowed for the understanding that environmental factors can significantly influence our genome and

29 how particular regions of our genome are regulated and expressed. As the field has expanded, research into the association of epigenetic features,

30 human health, disease risk, as well as aging has emerged, leading to the concept of epigenetic age and the creation of epigenetic age clocks.

31 Epigenetic clocks are computational tools that estimate biological age by analyzing specific DNA methylation patterns that are associated with

32 cellular aging across the genome. These clocks use measures of DNA methylation, chemical modifications that regulate gene expression without

33 altering the underlying DNA sequence, to provide a look into the aging process. This is important because epigenetic age may be a better measure

of an individual's health status than chronological age alone. *EpigeneticAgePipeline* aims to build off the functionality of other well established R

packages such as dnaMethyAge [1], methylclock [2], and minfi [3] to provide additional functionality and significantly improve usability.

The aim of this manuscript is to showcase the complete workflow performed by the package, from data preprocessing to visualization.

This comprehensive approach emphasizes the seamless integration of functionalities, highlighting the distinct advantages of

EpigeneticAgePipeline compared to other tools that typically require multiple separate steps to achieve similar outcomes.

What makes *EpigeneticAgePipeline* unique is that it integrates the entire workflow, including data pre-processing, epigenetic age

calculations, and downstream visualization tasks/final analysis into one seamless pipeline. The package allows for direct input of raw intensity

data (IDAT) files (27K, 450K, EPICv1/v2 and MSA microarray), which are then processed via sample and probe-level detection p-value filtering,

bead-count thresholds at both the sample and probe levels, Illumina control intensity checks, and the removal of cross-reactive probes and sex

probes [4–8]. Alternatively, users can input their own preprocessed methylation beta values. The package also allows for residual generation to

account for cell count data and batch effects that arise during methylation data collection. Residual generation also includes the option to regress

out principal components generated from the methylation dataset to account for additional sources of variation in the data. Cell proportion

estimation is also included, allowing the user to specify tissue type and methodology.

Furthermore, *EpigeneticAgePipeline* provides a range of visualizations, including a variety of correlation and distribution plots that

enhance the data's interpretability. These features, combined with the foundational functionalities of other well-established R packages, allow

EpigeneticAgePipeline to fill a unique niche in computational epigenetics.

Implementation

EpigeneticAgePipeline generates five common epigenetic age clocks, which include Horvath clock [9], Horvath skin and blood clock

[10], Hannum clock [11], PhenoAge/Levine clock [12] and GrimAge (V1 and V2 based on their published surrogate panels) [13,14] . We've also

included the PC versions for each of these clocks [1]. In addition to these 5 common clocks, the pipeline also generates the PedBE clock [15],

Wu's clock [16], Telomere Length's (TL) clock [17], BLUP clock [18], and EN clock [18]. Each one analyzes the methylation pattern of a unique

set of CpG (Cytosine-phosphate-Guanine) sites, which allows for more specific estimations depending on the tissue type and/or phenotype. Along

with epigenetic age estimation, the package allows for the generation of DunedinPACE [19], which reflects the pace of biological aging, and age

acceleration for each of the generated clocks.

When raw IDATs are provided, the pipeline performs detection p-value filtering at two levels: (i) sample-level retention if the mean

detection p-value is < 0.05, and (ii) probe-level retention only if the detection p-value is < 0.01 across passing samples. Bead-count filters exclude

samples with > 0.05 low-bead observations (bead count < 3) and drop probes with > 0.05 low-bead observations among retained samples. Default

parameters concerning detection p and bead count filtering can also be modified by the user. Illumina control intensity metrics are assessed via

minfi, where samples with mean log2 intensity of mMed and uMed < 10.5 are removed [3]. The pipeline then removes known cross-reactive

probes and sex-chromosome probes. For each of the microarray formats accepted (27K, 450K, EPICv1/v2, MSA), a separate list of previously

identified cross reactive probes is removed during this process [20–23]. Currently there has been no formal study which compiled a list of cross-

reactive probes for the MSA platform, however due to its similarity to EPICv2, we used EPICv2's list.

To handle replicate probes in newer EPICv2/MSA array formats targeting the same legacy CpG, beta values are collapsed by probe-ID

prefix using betasCollapseToPfx from the sesame R package, yielding a single averaged beta per CpG prior to clock computation [24].

To calculate cell proportion estimates, the package supports Constrained Projection (CP) and Robust Partial Correlations (RPC) style

generation for adult blood, cord blood, saliva, placenta (first and third trimester), and frontal cortex cells [25–32]. Buccal cell proportions are also

supported using the HEpiDISH (from EpiDISH R Package) hierarchical approach by first deconvolving epithelial, fibroblast and immune

fractions with a solid-tissue reference, then partitioning the immune fraction into blood cell subtypes [26,33,34].

For all clocks except DunedinPACE, the package provides three variations of epigenetic age acceleration: (i) the difference between

Epigenetic Age and chronological age, (ii) residuals from Epigenetic Age ~ Age + estimated cell proportions, and (iii) residuals from Epigenetic

Age ~ Age. These are analogous to the age accelerations provided by methylclock [2].

The original Horvath clock was one of the first epigenetic clocks created and remains one of the "gold standard" epigenetic age

measure. The clock uses a set of 353 CpG sites trained on 27K/450K arrays and is designed to be used on a wide range of tissue types. The

second original epigenetic clock is the Hannum clock. This clock relies on a set of 71 CpG sites trained on 450K array and is primarily designed

for use on blood samples. Both the Horvath and Hannum clocks have been shown to be significant predictors of age- and age-related conditions.

After the creation of the Horvath clock, the Horvath skin and blood clock was developed using a set of 391 CpG sites trained on 450K/EPICv1

array types for better estimations from skin, blood, saliva, human fibroblasts, keratinocytes, buccal cell, endothelial cell and lymphoblastoid cell

inputs. Another epigenetic age clock included is the Levine/PhenoAge clock which was created as a potential outperformer of other clocks

concerning predictions of age-based phenotypic outcomes, it uses a set of 513 CpG sites trained on 27K/450K/EPICv1. GrimAge is a measure

primarily designed for blood samples, with version 1 and 2 based on a set of 1030 CpG, difference being that version 2 uses an additional 2

DNAm (DNA methylation) surrogates. The surrogates are DNAm log(C-reactive protein) and DNAm log(Hemoglobin A1C), which allows

GrimAge2 to outperform the first in mortality prediction [14]. GrimAge has been known to predict a high Mortality Hazard Ratio compared to

other common epigenetic age measures. The PedBE clock uses 94 CpGs and was trained primarily on buccal epithelial samples (children and

adolescents) profiled on 450K/EPIC platforms. It is optimized for pediatric tissues and has been used to track developmental timing and age

acceleration in childhood cohorts. Wu's clock uses 111 CpGs and was trained on pediatric blood using 27K and 450K arrays. It provides highly

accurate age estimates in children and has been applied to investigate developmental and clinical outcomes in pediatric populations. The TL clock

uses 140 CpGs trained on blood using 450K and EPIC arrays to estimate leukocyte telomere length. It has been used to identify associations

between shorter telomere length and advancing age and relates to aging-related morbidity. The BLUP clock uses 319,607 CpGs trained on blood

and saliva measured on 450K and EPIC arrays. It leverages genome-wide features to yield high-precision age prediction and robust performance

across cohorts and tissues. The EN clock uses 514 CpGs trained on the same 450K/EPIC blood and saliva datasets as BLUP. The elastic-net (EN)

clock provides a sparse, portable predictor with strong cross-platform accuracy and reliable performance in external data.

The DunedinPACE measure is a measure of the pace of biological aging, which has been known to show strong associations with

phenotypic outcomes like morbidity, disability, and mortality. DunedinPace was developed for use in blood samples and uses a set of 173 CpG

sites.

EpigeneticAgePipeline also incorporates residual generation to allow for correction of important confounding variables. Principal

component analysis is performed on beta values, and residuals are generated adjusting for the first 5 PCs (default) generated on the methylation

data. Samples that are beyond 3 times (default) the median absolute deviation in any of the first PCs are marked as outliers. Residual generation

also incorporates adjustments for technical effects that commonly arise from methylation array measurements. The user may provide data for

array variables such as Row, Column, Slide, and Batch. Depending on which are given, the model will dynamically construct a suitable formula

for the regression (see Figure 1). There is also a built-in option to compute and adjust for cell count proportion estimates. Residuals generated

when chronological age is set as a predictor for biological age effectively gives a measure of age acceleration.

A series of plots and visualizations are also included in the final output to present the correlations between epigenetic age measures

and chronological age as well as the distribution of epigenetic age estimates for every sample. Raw data is exported as .txt file which can be used

for further analysis.

Results

The package's *main* function is the primary entry point for epigenetic age analysis. Users provide the directory containing methylation

and covariate data, if available (e.g., age, sex, smoking status, batch effects, etc.), along with options for normalizing beta values, selecting the

methylation array type, and processing raw or precomputed data. The function performs preprocessing steps, cell count generation, and the

creation of a file containing beta values for downstream analysis. This function also supports parallel processing for efficient handling of large

datasets and allows users to specify whether to use adult or cord blood assumptions during cell count generation.

The *generateResiduals* function is designed for advanced analysis following epigenetic age estimation. This function builds a linear

model based on user-specified variables or principal components (PCs) and outputs the corresponding residuals, which can be used for further

investigation of epigenetic age-related patterns. Users can input custom formulas for the model, specify the type of methylation array used, and

include additional covariates such as cell counts in the analysis. The function also supports the detection and removal of highly correlated

variables (if enabled) and identifies outliers using median absolute deviation (MAD). As with the main function, this process can leverage parallel

processing for efficiency and accommodates both adult and cord blood sample assumptions during cell count generation. To view a complete

outline of the workflow for either the main or generateResiduals function, see Figure 2/3.

To ensure the robustness of our pipeline, we first validated its implementation using the publicly available Gene Expression Omnibus

(GEO) dataset GSE237561 [35]. This dataset contains nuclear DNA methylation data for 126 samples—92 measured using the 450K array and 34

using the EPIC platform. Since both methylation and phenotypic data are publicly available, we were able to compare our results against

previously published findings. Specifically, a prior study investigating the relationship between clozapine use and epigenetic changes utilized this

dataset to calculate PhenoAge/Levine and DunedinPACE epigenetic age measures [36]. These epigenetic markers were incorporated into linear

mixed models to assess whether clozapine use predicts epigenetic age or age acceleration. To validate our pipeline, we used all 126 samples and

compared our computed PhenoAge/Levine and DunedinPACE measures against the published results. Our package generates these estimates

using the methylclock and dnaMethyAge packages, respectively, this comparison therefore seeks to confirm the accuracy of our implementation.

The full validation results can be found in the *Supplementary Information*. Following this validation, we applied our pipeline specifically to the

34 EPIC-measured samples for visualization. The covariates included in this analysis were sex, age, smoking score and number of days on

clozapine. Figure 4, as well as all *Supplementary Figures* (except validation) were generated using our EpigeneticAgePipeline on the EPIC-

measured samples from the GSE237561 dataset.

Inputs: The package accepts methylation data either in the form of a .csv file containing methylation beta values for each respective

sample, or a directory containing raw IDAT files.

Outputs: The package outputs raw epigenetic age/acceleration values (.txt and .md, Table S1), raw residual results (.csv), age and

epigenetic age correlation plots (Figure S2), matrix correlation plots (.png, Figure 3), and an epigenetic age distribution plot (.png, Figure 3). A

clock-coverage summary table is also outputted, reporting for each clock/surrogate the total CpGs, missing CpGs, and percent missing (Table S2).

Examples of all the plots can be found in the *Supplementary Information* files.

Specific information on installation procedures and file specificities, as well as vignettes are located within the GitHub readme [37].

Conclusions

EpigeneticAgePipeline provides a comprehensive and user-friendly method to generate epigenetic age measures for those studying the

effects of biological aging. Our pipeline better accounts for potential batch effects and confounding variables to minimize the effect of potential

biases on the results in studies concerning biological aging. Our pipeline provides a complete workflow for epigenetic age estimation, starting

with raw methylation data all the way to the generation of epigenetic age measures, incorporating a variety of quality control methods and

visualizations.

EpigeneticAgePipeline serves as a valuable tool for investigating the relationships between epigenetic age measures and a wide range

of health outcomes. Future updates will include the addition of emerging epigenetic aging measures/methods to ensure the pipeline remains

relevant to this expanding field.

Availability and Requirements

Project Name: EpigeneticAgePipeline

Project Homepage: https://github.com/CastellaniLab/EpigeneticAgePipeline

Operating Systems: Platform independent

Programing Language: R

Other Requirements: R 4.2+

License: CC0 1.0

Any restrictions to use by non-academics: No

List of Abbreviations

GEO: Gene Expression Omnibus

IDAT: Intensity data file

PC: Principal Component

MAD: Median Absolute Deviation

CpG: Cytosine-phosphate-Guanine

Declarations

Ethics approval and consent to participate: Not applicable, this study used publicly available datasets and did not involve any human participants,
human tissue, or human data collected by the authors.

Consent for publication: Not applicable.

Availability of data and materials: The datasets generated and/or analyzed during the current study are available in the Gene Expression Omnibus
repository, accession code GSE237561, https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE237561

Competing Interests: The authors declare that they have no competing interests

Funding: This work was supported by start-up funds to CC from The Department of Pathology and Laboratory Medicine at Western University. It
was further supported by institutional summer support (USRI) from Western University to SR.

Authors' contributions: SR led the software development portion of the project and contributed to testing and manuscript preparation. QL was
involved in extensive software testing and co-wrote the manuscript. JN contributed to code development and provided critical feedback during
the early stages of the project. SC assisted with testing and provided additional validation support. CC conceptualized and supervised the project,
provided funding, and oversaw all aspects of development and manuscript preparation. All authors read and approved the final manuscript.

Acknowledgements: We acknowledge the Digital Research Alliance and the Canadian Foundation for Innovation J. Evans Leaders Fund grant
#43481 for the support of the computing infrastructure used throughout our analyses.

Figure 1: Flowchart illustrating formula construction for the linear model during residual generation (*generateResiduals* function). *Xi* represents all other user provided covariates. This schematic is based off presence of covariates "Row", "Column", "Slide", and "Batch".

Figure 2: Workflow diagram illustrating the primary functionalities of the main() function

Flowchart for the main() function, which outlines the step-by-step process from data input to epigenetic age estimation and visualization, highlighting options for IDAT or beta value input and downstream processing.

Figure 3: Workflow diagram illustrating the primary functionalities of the generateResiduals() function

Flowchart for the generateResiduals() function, showcasing the process for residual generation, including data preprocessing, principal component analysis, outlier detection, and regression model construction.

Figure 4: A correlation matrix generated for Horvath epigenetic age from the GSE237561 dataset. The boxes below the diagonal showcase the magnitude of the correlation between each respective pair of variables. The boxes above the diagonal showcase a scatterplot of pairwise variables with a linear trendline, giving the directionality and strength of the correlation. The *main* function will output an analogous plot for each measure of epigenetic age if covariates have been included.

[1] Wang Y, Grant OA, Zhai X, Mcdonald-Maier KD, Schalkwyk LC. Insights into ageing rates comparison across tissues from recalibrating
cerebellum DNA methylation clock. GeroScience 2024;46:39–56. https://doi.org/10.1007/s11357-023-00871-w.
[2] Pelegí-Sisó D, de Prado P, Ronkainen J, Bustamante M, González JR. methylclock: a Bioconductor package to estimate DNA methylation
age. Bioinformatics 2021;37:1759–60. https://doi.org/10.1093/bioinformatics/btaa825.
[3] Aryee MJ, Jaffe AE, Corrada-Bravo H, Ladd-Acosta C, Feinberg AP, Hansen KD, et al. Minfi: a flexible and comprehensive
Bioconductor package for the analysis of Infinium DNA methylation microarrays. Bioinformatics 2014;30:1363–9.
https://doi.org/10.1093/bioinformatics/btu049.

[4]   IlluminaHumanMethylationEPICanno.ilm10b4.hg19. Bioconductor n.d.

http://bioconductor.org/packages/IlluminaHumanMethylationEPICanno.ilm10b4.hg19/ (accessed September 19, 2025).

[5]   IlluminaHumanMethylation27kanno.ilmn12.hg19. Bioconductor n.d.

http://bioconductor.org/packages/IlluminaHumanMethylation27kanno.ilmn12.hg19/ (accessed September 19, 2025).

[6]   IlluminaHumanMethylationEPICv2anno.20a1.hg38. Bioconductor n.d.

http://bioconductor.org/packages/IlluminaHumanMethylationEPICv2anno.20a1.hg38/ (accessed September 19, 2025).

[7]   IlluminaHumanMethylation450kanno.ilmn12.hg19. Bioconductor n.d.

http://bioconductor.org/packages/IlluminaHumanMethylation450kanno.ilmn12.hg19/ (accessed September 19, 2025).

[8]   IlluminaHumanMethylationMSAanno.ilm10a1.hg38. Bioconductor n.d.

http://bioconductor.org/packages/IlluminaHumanMethylationMSAanno.ilm10a1.hg38/ (accessed September 19, 2025).

[9]   Horvath S. DNA methylation age of human tissues and cell types. Genome Biol 2013;14:3156. https://doi.org/10.1186/gb-2013-14-10-

r115.

[10]   Horvath S, Oshima J, Martin GM, Lu AT, Quach A, Cohen H, et al. Epigenetic clock for skin and blood cells applied to Hutchinson

Gilford Progeria Syndrome and *ex vivo* studies. Aging 2018;10:1758–75. https://doi.org/10.18632/aging.101508.

[11]   Hannum G, Guinney J, Zhao L, Zhang L, Hughes G, Sadda S, et al. Genome-wide Methylation Profiles Reveal Quantitative Views of

Human Aging Rates. Mol Cell 2013;49:359–67. https://doi.org/10.1016/j.molcel.2012.10.016.

[12]   Levine ME, Lu AT, Quach A, Chen BH, Assimes TL, Bandinelli S, et al. An epigenetic biomarker of aging for lifespan and healthspan.

Aging 2018;10:573–91. https://doi.org/10.18632/aging.101414.

[13]   Lu AT, Quach A, Wilson JG, Reiner AP, Aviv A, Raj K, et al. DNA methylation GrimAge strongly predicts lifespan and healthspan. Aging

2019;11:303–27. https://doi.org/10.18632/aging.101684.

[14]   Lu AT, Binder AM, Zhang J, Yan Q, Reiner AP, Cox SR, et al. DNA methylation GrimAge version 2. Aging 2022;14:9484–549.

https://doi.org/10.18632/aging.204434.

[15]   McEwen LM, O'Donnell KJ, McGill MG, Edgar RD, Jones MJ, MacIsaac JL, et al. The PedBE clock accurately estimates DNA

methylation age in pediatric buccal cells. Proc Natl Acad Sci 2020;117:23329–35. https://doi.org/10.1073/pnas.1820843116.

[16]   Wu X, Chen W, Lin F, Huang Q, Zhong J, Gao H, et al. DNA methylation profile is a quantitative measure of biological aging in children.

Aging 2019;11:10031–51. https://doi.org/10.18632/aging.102399.

[17]   Lu AT, Seeboth A, Tsai P-C, Sun D, Quach A, Reiner AP, et al. DNA methylation-based estimator of telomere length. Aging

2019;11:5895–923. https://doi.org/10.18632/aging.102173.

[18]   Zhang Q, Vallerga CL, Walker RM, Lin T, Henders AK, Montgomery GW, et al. Improved precision of epigenetic clock estimates across

tissues and its implication for biological ageing. Genome Med 2019;11:54. https://doi.org/10.1186/s13073-019-0667-1.

[19]   Belsky DW, Caspi A, Corcoran DL, Sugden K, Poulton R, Arseneault L, et al. DunedinPACE, a DNA methylation biomarker of the pace

of aging. eLife 2022;11:e73420. https://doi.org/10.7554/eLife.73420.

[20]   Chen Y, Choufani S, Grafodatskaya D, Butcher DT, Ferreira JC, Weksberg R. Cross-Reactive DNA Microarray Probes Lead to False

Discovery of Autosomal Sex-Associated DNA Methylation. Am J Hum Genet 2012;91:762–4. https://doi.org/10.1016/j.ajhg.2012.06.020.

223 [21] Chen Y, Lemire M, Choufani S, Butcher DT, Grafodatskaya D, Zanke BW, et al. Discovery of cross-reactive probes and polymorphic

224   CpGs in the Illumina Infinium HumanMethylation450 microarray. Epigenetics 2013. https://doi.org/10.4161/epi.23470.

225 [22] Pidsley R, Zotenko E, Peters TJ, Lawrence MG, Risbridger GP, Molloy P, et al. Critical evaluation of the Illumina MethylationEPIC

226   BeadChip microarray for whole-genome DNA methylation profiling. Genome Biol 2016;17:208. https://doi.org/10.1186/s13059-016-

227   1066-1.

228 [23] Peters TJ, Meyer B, Ryan L, Achinger-Kawecka J, Song J, Campbell EM, et al. Characterisation and reproducibility of the

229   HumanMethylationEPIC v2.0 BeadChip for DNA methylation profiling. BMC Genomics 2024;25:251. https://doi.org/10.1186/s12864-

230   024-10027-5.

231 [24] Zhou W, Triche TJ Jr, Laird PW, Shen H. SeSAMe: reducing artifactual detection of DNA methylation by Infinium BeadChips in

232   genomic deletions. Nucleic Acids Res 2018;46:e123. https://doi.org/10.1093/nar/gky691.

233 [25] Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH, et al. DNA methylation arrays as surrogate

234   measures of cell mixture distribution. BMC Bioinformatics 2012;13:86. https://doi.org/10.1186/1471-2105-13-86.

235 [26] Teschendorff AE, Breeze CE, Zheng SC, Beck S. A comparison of reference-based algorithms for correcting cell-type heterogeneity in

236   Epigenome-Wide Association Studies. BMC Bioinformatics 2017;18:105. https://doi.org/10.1186/s12859-017-1511-5.

237 [27] FlowSorted.Blood.450k. Bioconductor n.d. http://bioconductor.org/packages/FlowSorted.Blood.450k/ (accessed September 25, 2025).

238 [28] FlowSorted.Blood.EPIC. Bioconductor n.d. http://bioconductor.org/packages/FlowSorted.Blood.EPIC/ (accessed September 19, 2025).

239 [29] FlowSorted.CordBloodCombined.450k. Bioconductor n.d. http://bioconductor.org/packages/FlowSorted.CordBloodCombined.450k/

240   (accessed September 19, 2025).

241 [30] BeadSorted.Saliva.EPIC. Bioconductor n.d. http://bioconductor.org/packages/BeadSorted.Saliva.EPIC/ (accessed September 19, 2025).

242 [31] planet. Bioconductor n.d. http://bioconductor.org/packages/planet/ (accessed September 19, 2025).

243 [32] FlowSorted.DLPFC.450k. Bioconductor n.d. http://bioconductor.org/packages/FlowSorted.DLPFC.450k/ (accessed September 19, 2025).

244 [33] Zheng SC, Webster AP, Dong D, Feber A, Graham DG, Sullivan R, et al. A Novel Cell-Type Deconvolution Algorithm Reveals

245   Substantial Contamination by Immune Cells in Saliva, Buccal and Cervix. Epigenomics 2018. https://doi.org/10.2217/epi-2018-0037.

246 [34] Salas LA, Zhang Z, Koestler DC, Butler RA, Hansen HM, Molinaro AM, et al. Enhanced cell deconvolution of peripheral blood using

247   DNA methylation for high-resolution immune profiling. Nat Commun 2022;13:761. https://doi.org/10.1038/s41467-021-27864-7.

248 [35] GEO Accession viewer n.d. https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE237561 (accessed September 19, 2025).

249 [36] Gillespie AL, Walker EM, Hannon E, McQueen GA, Sendt K-V, Avila A, et al. Longitudinal changes in DNA methylation associated with

250   clozapine use in treatment-resistant schizophrenia from two international cohorts. Transl Psychiatry 2024;14:390.

251   https://doi.org/10.1038/s41398-024-03102-8.

252 [37] Castellani CA. CastellaniLab/EpigeneticAgePipeline 2025.

256