Lipid hydrogen isotope compositions primarily reflect growth water in the model
archaeon *Sulfolobus acidocaldarius*

Carolynn M. Harris[a]#, Sebastian Kopf[b], Jeemin H. Rhim[a,c], Alec Cobban[a], Felix J. Elling[d,e], Xiahong
Feng[a], Jamie McFarlin[b,f], Yuki Weber[,d^], Yujiao Zhang[a%], Alice Zhou[a&], Harpreet Batther[b], Ann
Pearson[d], William D. Leavitt [a,g]

[a] Department of Earth Sciences, Dartmouth College

[b] Department of Geological Sciences, University of Colorado at Boulder

[c] Department of Ecology, Evolution, and Marine Biology, U.C. Santa Barbara

[d] Department of Earth & Planetary Sciences, Harvard University

[e] Leibniz-Laboratory for Radiometric Dating and Isotope Research, Kiel University

[f] Department of Geology and Geophysics, University of Wyoming

[g] Department of Chemistry, Dartmouth College

Running head: Lipid hydrogen isotopes of S. acidocaldarius

Address correspondence to Carolynn M. Harris**,** carolynn.m.harris.gr@dartmouth.edu

[&] Current address: Connecticut Agricultural Experiment Station, New Haven, CT

[^] Current address: Chroma Medicine, Boston, MA

[%] Current address: National Center for Science & Technology Evaluation, P.R. China

Abstract

The stable hydrogen isotope composition (δ^2H) of lipid biomarkers can track environmental
processes and remain stable over geologically relevant time scales, enabling studies of past
climate, hydrology, and ecology. Most research has focused on lipids from the domain Eukarya (e.g.,
plant waxes, long-chain alkanes), and the potential of prokaryotic lipid biomarkers from the domain
Archaea to offer unique insights into environments not captured by eukaryotic lipids remains
unclear. Here, we investigate the H-isotope composition of biphytanes in *Sulfolobus acidocaldarius*,
a model thermoacidophile and obligate heterotroph. We conducted a series of experiments that
varied temperature, pH, shaking rate, electron acceptor availability, or electron donor flux. From
these experiments, we quantified the lipid/water H-isotope fractionation ($^2\varepsilon_{L/W}$) values for core
biphytane chains derived from tetraether lipids. The $^2\varepsilon_{L/W}$ values are consistently negative (-230 to -
180 ‰) and are relatively invariant across all experiments despite a 20-fold change in doubling times
and 2-fold change in lipid cyclization. The magnitude and relative invariance of $^2\varepsilon_{L/W}$ values are
consistent with studies on other heterotrophic archaea and suggests archaeal lipids may be faithful
recorders of the δ^2H composition of growth water. Our study highlights the potential of archaeal lipid
$^2\varepsilon_{L/W}$ as a hydrological proxy, offering new insights into environments where traditional proxies, such
as plant-derived lipids, are not available, including extreme environments and extraterrestrial
settings.

Importance

Reconstructing past climates is crucial for understanding Earth's environmental history and its
responses to changing conditions. This study examines *Sulfolobus acidocaldarius*, a
thermoacidophilic archaeon that thrives in extreme environments like hot springs. These
microorganisms incorporate hydrogen water in the growth environment into membrane lipids,
creating hydrogen isotope signatures that can reflect hydroclimate conditions. Our findings show
that these hydrogen isotope ratios remain consistent even under varying temperature, pH, oxygen
levels, and electron donor fluxes, indicating a stable fractionation between lipids and water. This
invariance suggests that *S. acidocaldarius* lipids could serve as a robust proxy for reconstructing
ancient water H-isotope values, especially in extreme environments where traditional proxies, such
as plant waxes, are absent. This research has broader implications for planetary-scale
reconstructions, including potential applications in studying past climates on other planets, such as
Mars, where similar microorganisms may have existed in hydrothermal conditions.

1. Introduction

The relative abundances of hydrogen isotopes ($^2H/^1H$) in water are set by physical, hydrological, and climatic factors (1, 2). Certain biomolecules incorporate these isotopes with a consistent offset (fractionation) relative to water, preserving a record of the isotopic composition (δ^2H values) of the water present during growth (3). Some lipids can retain their original H-isotope composition for up to 10^8 years and lipid hydrogen isotope compositions have been used to study hydrologic and environmental change over millennia to millions of years, dating as far back as the Paleoproterozoic (4–6). Most studies of biomarker H-isotope composition focus on lipids from photoautotrophic organisms (e.g., plant wax *n*-alkanes) or chemoautotrophic and heterotrophic bacteria (e.g., fatty acids) (3, 7–9) and the H-isotope compositions recorded in lipids from the Archaea are comparatively poorly understood. A small, but growing, body of literature, however, has begun to address this gap (10–15). Understanding the extent to which the δ^2H value of archaeal lipids records the δ^2H value of growth water requires a deeper examination of how environmental conditions impact this signal. A comprehensive investigation of the physicochemical factors affecting H-isotope fractionation between archaeal lipids and water ($^2\varepsilon_{L/W}$) is essential for employing archaeal lipid δ^2H composition as a proxy for the δ^2H composition of environmental water over space and time.

The $^2\varepsilon_{L/W}$ value for a biomolecule integrates both the composition of source water and the net biosynthetic fractionations resulting from the specific pathways used during lipid biosynthesis. Reported $^2\varepsilon_{L/W}$ values vary widely among domains of life. In phototrophic Eukaryotes (e.g., plants and algae), $^2\varepsilon_{L/W}$ values of waxes (e.g., n-alkanes) are consistently negative, ranging from -150 to -50 ‰ (8, 16). It is important to understand the causes of this variability when applying lipid δ^2H values to reconstruct water δ^2H values. Net biosynthetic fractionation is generally consistent across taxa, though genetic factors introduce some variance (17–22). Environmental factors such as temperature, salinity, light availability, and CO_2 concentration, also influence $^2\varepsilon_{L/W}$ values, possibly via their impact on growth and biosynthesis rates (16, 23–28). These effects are well-characterized in plants, and to a lesser extent in algae, and can be accounted for when the δ^2H values of plant waxes are used to reconstruct past hydroclimate (8, 29, 30).

In contrast to the coherent magnitude and direction of $^2\varepsilon_{L/W}$ values observed from phototrophic eukaryotes, the $^2\varepsilon_{L/W}$ of bacterial fatty acids varies widely in both direction and magnitude (-400 ‰ to +300 ‰) (31–38). Bacteria that use different central energy metabolisms (e.g., chemoautotrophy, photoautotrophy, or heterotrophy) exhibit distinct ranges of $^2\varepsilon_{L/W}$ values, which has been attributed to differences in metabolic pathways and the substrate used for growth (e.g., simple sugars vs. tricarboxylic acid (TCA) cycle intermediates). A recent study suggests that variations in $^2\varepsilon_{L/W}$ values from aerobic heterotrophic bacteria are explained by energy fluxes through the intracellular hydride carriers $NADP^+$ and NADPH (38). Given this metabolism-dependent heterogeneity, the δ^2H values of

bacterial lipids in sediment records may have more utility as proxies for the energetic state of past
microbial metabolisms (33, 36, 38).

Lipid-water ^{2}H-fractionation from the domain Archaea have been understudied to-date. From what
we do know, $^2\varepsilon_{L/W}$ values have been reported for diether lipids (archaeol) in a mesophilic halophile
(11) and a methanogen (13), tetraether-derived biphytanes in three thermoacidophiles and one
thermomesophile (10, 15), and one aerobic marine chemoautotroph (14), and whole tetraethers or
tetraether-derived biphytanes in several environmental sediment samples (10, 12). The observed
$^2\varepsilon_{L/W}$ values for these archaeal lipids are uniformly negative (i.e. -300 to -150 ‰) and similar in
magnitude to those reported for isoprenoid lipids produced by eukaryotic phototrophs (e.g. C_{15} and
C_{30} isoprenoids) (3, 25).

Our understanding of the factors controlling the H-isotope compositions of archaeal lipids are less
developed than for the Eukarya and Bacteria. Recent experiments demonstrate that $^2\varepsilon_{L/W}$ is nearly
constant over a range of growth rates in a marine autotroph (*Nitrosopumilus maritimus* (14)) and that
the impacts of carbon metabolism and growth phase on $^2\varepsilon_{L/W}$ are minor (*Archaeoglobus fulgidus*
(15)). This relative invariance in $^2\varepsilon_{L/W}$ values is perhaps surprising considering the taxonomic and
metabolic diversity of the investigated Archaea. The impact of environmental factors on this
fractionation, however, has only been investigated within a single archaeal strain (*Haloarcula*
marismortui (11)). Further investigation of the effects of multiple environmental factors on $^2\varepsilon_{L/W}$ in
Archaea is crucial, particularly since many archaea are adapted to extreme environments that can
experience both rapid and large changes to physicochemical conditions (39).

Isoprenoid glycerol dibiphytanyl glycerol tetraethers (iGDGTs) are archaeal lipids with particular
relevance to paleoenvironmental reconstructions (40). These tetraethers are structurally diagnostic,
diagenetically robust, and routinely analyzed for paleoclimate reconstruction based on the relative
abundances of specific structural moieties (40–42). Archaea can synthesize a unique series of
iGDGTs that contain from zero to eight cyclopentane rings (e.g., iGDGT-0 through iGDGT-8), though
the most highly ringed moieties (iGDGT-7 and -8) have only been recovered from hydrothermal spring
environments (43). Members of the Thaumarchaeota phyla can also produce crenarchaeol, an
iGDGT containing four cyclopentane rings and one cyclohexyl ring (43). The distribution of these
moieties in natural systems is related to growth temperature and other environmental factors (44–
48) and forms the basis of the TEX_{86} paleo sea surface temperature proxy (41). At low pH and high
temperatures, Archaea synthesize more highly-ringed moieties as an adaptation to maintain
membrane rigidity and reduce permeability. In hydrothermal systems, the ring distributions of
thermoacidophilic archaea record the interactions between temperature, pH, energy availability,
and other factors that influence growth rate (48–50).

To better understand the signals recorded by archaeal lipid δ^2H composition in natural systems, it is
necessary to determine the extent to which environmental conditions influence $^2\varepsilon_{L/W}$ values. In this

study, we determined the H-isotope fractionation between iGDGT-derived biphytanes and growth
water ($^2\varepsilon_{L/W}$) for the thermoacidophilic and heterotrophic archaeon, *Sulfolobus acidocaldarius* (DSM
639). *S. acidocaldarius* evolved to grow optimally under multiple extreme conditions, namely low
pH's, high temperatures, and low oxygen partial pressure (51, 52). For this work *S. acidocaldarius*
was selected as a model organism because it thrives in extreme environments devoid of traditional
eukaryotic lipid proxies (e.g., plant waxes), and because it can grow under a wide range of
physicochemical conditions, with rapid doubling times and high biomass yields (53).

Here we cultivated *S. acidocaldarius* over a range of physicochemical conditions that included shifts
in temperature, pH, shaking rate, electron acceptor or electron donor flux. Changes in these
environmental conditions are known to impact the growth rate and degree of iGDGT cyclization (47–
49, 54, 55), but their impact on lipid/water fractionation is unexplored. Understanding how these
environmental factors influence lipid/water fractionation is critical for interpreting archaeal lipid δ^2H
signatures in natural systems and refining their use as robust proxies for reconstructing past
hydroclimate conditions.

2. Methods

2.1 Culture strains and growth

Axenic cultures of *S. acidocaldarius* (DSM 639) were grown aerobically under different
environmental conditions (**Table 1**). Culture media was prepared following the Brock recipe (56) with
the following modifications (see *Substrate* column in Table 1). For all experiments, media contained
5.9 mmol L^{-1} (0.2% w/v) sucrose as the organic substrate. The electron donor flux experiments were
also supplemented with 4.0 mmol L^{-1} (0.1% w/v) NZ-amine in addition to sucrose.

Growth was monitored photometrically at regular intervals throughout each experiment by
measuring the optical density (i.e., absorbance at 1 cm pathlength) of a culture aliquot at 600 nm
(OD_{600}) on a spectrophotometer. All experiments were inoculated to an initial OD_{600} between 0.004
and 0.008, using cells derived from an active pre-culture (grown on the same medium) in mid-
exponential phase. Specific growth rates (division day^{-1}; μ) were calculated using OD_{600}
measurements during the early to mid exponential phase after Cobban et al., (2020). Doubling time
(T_D) was calculated as:

Eq. (1) $T_D \ (hours) \ = \ ln(2)/\mu$

2.2 Experimental conditions experiments

We examined how temperature, pH, shaking rate, O_2 availability, and electron donor (sucrose) flux
influence H-isotope fractionation in *S. acidocaldarius* through a series of batch, fed-batch, and
chemostat experiments.

Batch experiments: Three sub-experiments were performed in which one environmental condition
(temperature, pH, or shaking rate) was systematically varied while holding the other two conditions
constant (previously described in Cobban et al., 2020). The reference conditions were 70 °C, pH 3,
and 200 RPM shaking rate. Shaking rate is considered a proxy for aeration rate, where faster speeds
increase oxygen availability in the media, though it also increases physical agitation.

Batch experiments were performed in temperature-controlled, shaking incubators (Innova-42,
Eppendorf) in 250 mL Erlenmeyer flasks containing 125 mL media covered with loose plastic caps to
reduce evaporation. Evaporation was further reduced by maintaining a 2L tray of water in the
incubator to humidify the atmosphere. All experiments were performed in quintuplicate. Biomass
samples were collected in mid-exponential phase by centrifuging (3214 x *g*; Eppendorf 5810 R, S-4-
104 rotor) 50 mL of culture for 15 minutes at 4 °C, the resulting cell pellets were stored at -80 °C until
lipid extraction.

Fed-batch bioreactor experiments: To more precisely determine the impact of oxygen availability,
we performed gas-fed batch experiments in bioreactors (previously described in Cobban et al.,
2020). Briefly, *S. acidocaldarius* was grown in 300 mL of medium in 1 L glass vessels. To modulate
availability (as flux) of the terminal electron acceptor O_2 was introduced to bioreactors at four
different partial pressures: 0.2, 0.5, 2.0 or 20 % (balance N_2). The O_2 dry mixing ratios 0.2, 0.5, and
2% were generated by mixing pure N_2 mixed with 2% O_2 at 9:1, 3:1, and 0:1 ratios, respectively, which
was controlled via my-Control PID controllers (Applikon, Delft, Netherlands). The 20% experiment
used a separate gas feedstock of Ultra ZeroGrade Air (AirGas). Culture conditions were maintained
at 70 °C, pH 3, stirred at 200 RPM (Rushton-type impeller), and at a total gas flux of 200 mL min^{-1}.
Each experiment was performed in triplicate via three bioreactors operated in parallel. Biomass
samples were collected by removing culture directly from the vessel, centrifuged at 4 °C, to pellet
biomass, and then stored at -80 °C until lipid extraction.

Chemostat experiments: To determine the impact of electron donor flux, we performed chemostat
experiments in which the dilution rate of the growth vessel was set to yield cell doubling times (T_D) of
7.0, 21.3, and 40.0 hours (previously described in Zhou et al. 2020). Briefly, *S. acidocaldarius* was
grown in 300 mL of medium maintained at 70 °C, pH 2.25, stirred at 200 RPM (Rushton-type
impeller), and aerated with a constant flux of 200 mL min^{-1} Zero Air (ultra-high purity, UHP, AirGas/
AirLiquide, Inc.). Three chemostats were operated in parallel under continuous culture conditions.

When steady state was achieved (assessed via regular OD_{600} measurements), biomass samples
were harvested by continually collecting the outflow of each bioreactor into a chilled vessel (0 to 4
°C) for between 1 and 4 days, followed by centrifugation at 4 °C and storage at -80 °C prior to lipid
processing. Multiple extracts representing 2 to 3 sampling events at identical growth rates were
combined before lipid extraction.

2.3 Lipid extraction and derivatization

Core lipid fractions (iGDGTs) were extracted from lyophilized cell pellets via acidic hydrolysis-
methanolysis (45, 48). iGDGTs were chemically derivatized to biphytanes to permit the
determination of the δ^2H value of component biphytanes (10, 14). In brief, ether bonds were cleaved
via digestion in 57% hydroiodic acid (HI) at 125 °C for 4 hours. The resulting alkyl iodides were
hydrogenated (e.g., reduced to biphytanes, BPs) in H_2/PtO_2. An isotope dilution occurs when H is
added during hydrogenation, which is corrected for during data processing and reduction (see
Section 2.5).

2.4 Lipid identities & distributions

Isoprenoid BP chains were analyzed via GC-MS (Thermo ISQ LT with TRACE 1310) for compound
identification and via GC-flame ionization detector (GC-FID; Thermo TRACE 1310) for compound
quantification at the University of Colorado at Boulder Earth Systems Stable Isotope Lab (CUBES-
SIL, Boulder, CO) (**FIG 1**). Biphytane yields and recovery were calculated using GC peak areas and
co-injected n-alkane mixture as a quantification standard.

To describe the average amount of cyclopentyl rings in each BP distribution, we calculated a BP-Ring
Index based on the relative abundance of each BP moiety (**Eq. 2**). This metric is similar to the iGDGT-
based Ring Index (**Eq. 3**), which describes the average amount of cyclopentyl rings in a distribution
based on the relative abundance of each iGDGT moiety (48, 57).

Eq. (2)
$$RI_{BP} = \frac{BP-1 + (2 \cdot BP-2) + (3 \cdot BP-3)}{\sum_o^3 BP}$$

214   **Eq. (3)**
$$RI_{GDGT} = \frac{iGDGT-1 + (2 \cdot iGDGT-2) + (3 \cdot iGDGT-3) + \ldots + (8 \cdot iGDGT-8)}{\sum_o^8 iGDGT}$$

2.5 δ^2H analysis of media waters, substrates, and biphytane lipids

2.5.1 Media waters

Water samples for hydrogen isotope (δ^2H) analysis were collected concurrently with harvesting
biomass for lipid analyses. Aliquots of filter-sterilized growth medium water was measured for δ^2H
at the Stable Isotope Laboratory at Dartmouth College via a dual inlet isotope ratio mass
spectrometer (IRMS; Thermo Delta Plus 304 XL) coupled to an H-Device (pyrolyzed to H_2 gas) for
water reduction by chromium powder at 850°C (previously described in Kopec et al. (2018). The
δ^2H$_{Water}$ values were corrected using three laboratory standards spanning -161 to -7 ‰ vs. VSMOW,
which were run every 8-10 samples. Analytical precision was <0.5 ‰.

2.5.2 Substrates

The δ^2H composition of non-exchangeable C-bound H from substrates used in growth media was
analyzed in triplicate as trifluoroacetate derivatives via a Thermo Delta Plus XP coupled to a TC-EA
(58). The δ^2H composition of sucrose was -93.0 ± 1.2 ‰.

2.5.3 Biphytanes

For all experiments, we report $^2\varepsilon_{L/W}$ values for biphytane (BP) hydrocarbons derived via ether cleavage
from acid extractable iGDGTs. The δ^2H composition of individual BPs was determined by gas
chromatography pyrolysis isotope ratio mass spectrometry (GC-P-IRMS) on a GC IsoLink II IRMS
System (Thermo Scientific) at the University of Colorado at Boulder Earth Systems Stable Isotope
Lab (CUBES-SIL, Boulder, CO). The system comprised a Trace 1310 GC fitted with a programmable
temperature vaporization (PTV) injector and either a 30 m ZB5HT column (i.d. = 0.25 mm, 0.25 µm,
Phenomenex, Torrance, CA, USA) or a 60 m DB1 column (i.d. = 0.25 mm, 0.25 µm, Agilent, Santa
Clara, CA, USA), ConFlo IV interface, and 253 Plus mass spectrometer (Thermo Scientific).

2.5.4 Data reduction and notation

The δ^2H values of individual biphytanes (δ^2H$_{BP}$) were measured relative to H_2 reference gas (δ^2H$_{raw}$)
and calibrated to the international references scale (VSMOW) using a standard n-alkane mixture (A7,
containing C15 through C30 n-alkanes spanning -9 to -263 ‰ vs. VSMOW; A. Schimmelmann,
Indiana University). The A7 standard was combined with a C_{36} n-alkane (nC_{36}, -259.2 ‰ vs. VSMOW;
A. Schimmelmann, Indiana University) and measured throughout each analytical run at regular
intervals and at different concentrations. The BP hydrogen isotope calibration was performed as in
Leavitt et al., (2023) using the R packages *isoreader* (v 1.3.0) and *isoprocessor* (v 0.6.11) available at
github.com/isoverse (59). In brief, δ^2H$_{raw}$ values were corrected for offset, scale compression and
peak-size effects using an inverted multivariate linear regression which was applied to all standards
and samples to determine δ^2H$_{cal}$ values. Further details of calibration and data reduction are
available in Leavitt et al., 2023. The total analytical uncertainty of the corrected δ^2H$_{BP}$ values was
calculated using standard error propagation of the peak-size adjusted error estimates and
hydrogenation correction assuming all errors to be uncorrelated. The hydrogenation correction
ranged from 10.3 to 12.7 ‰ and increased analytical uncertainty by up to 1.9 ‰.

All ^{2}H/^{1}H ratios are reported in delta notation (δ^2H) in permil (‰) units relative to the international
seawater standard on the VSMOW-SLAP (Vienna Standard Mean Ocean Water, Standard Light
Antarctic Precipitation) scale.

Eq. (4) $\delta^2 H\ (‰)\ =\ (\dfrac{^2H/\ ^1H_{Sample}}{^2H/\ ^1H_{Standard}} - 1\ )$

All fractionation factors are reported in epsilon notation ($^2\varepsilon$) in permil (‰) units. The hydrogen
isotope fractionation between growth water and BP lipids ($^2\varepsilon_{L/W}$) is calculated as:

Eq. (5 a, b) $^2\alpha_{L/W}\ =\ \dfrac{^2H/\ ^1H_{BP}}{^2H/\ ^1H_{Water}}$

260 $^2\varepsilon_{L/W}\ (‰)\ =\ (\ ^2\alpha_{L/W} - 1\ )$

Corrected $\delta^2 H_{BP}$ values and resulting $^2\varepsilon_{L/W}$ fractionation factors from biological replicates (n ≥ 1) and
analytical replicates (n ≥ 3) were averaged for each experimental condition. All averages are
weighted means of individual measurements ($1/\sigma^2$ weights) to account for the amplitude-dependent
range in uncertainties. The reported error estimate of each average is the larger of the standard
deviation of all replicates or the propagated uncertainty from individual measurements. Abundance
weighted averages for $\delta^2 H_{BP}$ and $^2\varepsilon_{L/W}$ values include all BPs with relative abundance > 5%.

The ring difference ($\Delta\varepsilon$/ring), or changes in $^2\varepsilon_{L/W}$ with increasing number of ring structures is
calculated for all pairwise combinations of $^2\varepsilon_{L/W}$ values after Leavitt et al., (2023) where X and Y refer
to the ring number of BP moieties and X > Y:

Eq. (6) $\dfrac{\Delta\varepsilon}{ring}(‰)\ =\ \dfrac{^2\varepsilon_{L/W}(X) -\ ^2\varepsilon_{L/W}(Y)}{(X-Y)}$

Raw data archived on Zenodo (60); code for all data manipulations and statistical analyses is
available on GitHub (https://github.com/carolynnharris/Saci_Enviro_d2H).

3. Results

We examined how temperature, pH, shaking rate, O_2 availability, and electron donor (sucrose) flux
influence H-isotope fractionation in *S. acidocaldarius* through a series of batch, fed-batch, and
chemostat experiments. Details of the growth responses and lipid iGDGT profile responses to these
conditions can be found in Cobban et al., (2020) and Zhou et al., (2019). In brief, each environmental
condition independently influences growth rate, the relative abundance of many subsets of iGDGT
core lipids, and the RI-GDGT (47, 48) (**Table 1, FIG 2A-E, S1**).

Here, we show that each environmental condition also influences the relative abundance of BP lipids
and the RI-BP (**Table 2**, **FIG. 2F-O**). Among all samples, the relative abundance of biphytanes

averaged 13%, 30%, 49%, 7% for BP-0, -1, -2, and -3, respectively; BP-3 was the least abundant and

was not measurable in some samples. RI-BP ranged from 1.01 to 2.04 and, as expected, was

positively correlated with RI-GDGT ($R^2 = 0.52$, $p = 0.001$, **FIG. S2**). Among all samples, the abundance

weighted mean δ^2H_{BP} values range from -276 to -225 ‰ and were consistently depleted in ^2H relative

to growth water, corresponding to $^2\varepsilon_{L/W}$ values from -230 to -180 ‰ (**Table 2, FIG. 2P-T**).

Despite significant effects on growth rate and lipid cyclization, individual environmental parameters

had minimal impact on $^2\varepsilon_{L/W}$ values, which spanned a range of 44, 31, 15, 21, and 25 ‰ in the

temperature, pH, shaking rate, O_2 availability, and electron donor flux experiments, respectively. We

observed a weak linear relationship between environmental conditions and $^2\varepsilon_{L/W}$ only in the electron

donor flux experiment, where larger doubling times are associated with slightly less negative $^2\varepsilon_{L/W}$

values (smaller offset from water) ($R^2 = 0.89$, $p = 0.22$, slope = 0.72 ± 0.25 ‰/ hour) (**FIG. 2T**).

In the batch and fed-batch experiments, $^2\varepsilon_{L/W}$ values for individual BPs showed no consistent pattern

with increasing number of cyclic moieties and the ring difference ($\Delta\varepsilon$/ring) varied widely in direction

and magnitude among all treatments (-39 to +24 ‰) (**Table 2**, **FIG. 3A-D, S3A-D**). The variability in

mean $\Delta\varepsilon$/ring was smaller, though still not of uniform direction, with values ranging from -5.4 to +10.2

‰ (**Table 2, FIG. S4A-D**). In contrast, $^2\varepsilon_{L/W}$ values in the chemostat experiment generally increased

with the number of cyclic moieties, showing offsets of 3.7, 4.1, and 10.1 ‰ for BP-1, BP-2, and BP-3

relative to BP-0 (**Table 2, FIG. 3E, S3E**). Similarly, the mean ring difference ($\Delta\varepsilon$/ring) was uniformly

positive and showed little variation among doubling times, ranging from 4.9 to 9.2 ‰ (**Table 2, FIG.**

S4E). On average in continuous culture, each additional ring contributes a 7.4 ± 2.2 ‰ increase to

biphytane $^2\varepsilon_{L/W}$.

Correlation matrices show the linear relationship between each environmental condition, growth

parameters (doubling time and maximum OD_{600}), mean lipid cyclization indices (RI-BP and RI-GDGT),

$\Delta\varepsilon$/ring and abundance weighted $^2\varepsilon_{L/W}$ values for each environmental condition experiment (**FIG. 4**).

These plots further demonstrate that the minimal variation in $^2\varepsilon_{L/W}$ observed within each experiment

is not related to variation in the experimentally manipulated environmental conditions or any other

measured parameter.

To explore the possibility that lipid/water fractionation is related to optimal growth, we relate

abundance weighted mean $^2\varepsilon_{L/W}$ values to doubling time for each environmental condition (**FIG. 5A-**

E). To allow for comparisons among all environmental conditions, we also relate $^2\varepsilon_{L/W}$ to doubling

time for all experiments combined. In this pooled dataset, abundance weighted mean $^2\varepsilon_{L/W}$ values

range from -230 to -180 ‰ (mean = -204 ± 12 ‰, based on 22 biological replicates; **Table 2**) and

showed no correlation to doubling time (**FIG. 5F**, $R^2 = 0.0$, $p = 0.97$), indicating a decoupling between

lipid/water fractionation and growth rate.

4. Discussion

4.1 Effect of environmental conditions on archaeal lipid $^2\varepsilon_{L/W}$ values

The primary goal of this work was to investigate the impact of environmental conditions on lipid/water fractionation in archaeal lipids, which relates to their potential as archives of hydroclimate across space and time. The conditions tested in this study were chosen to encompass the natural conditions of thermoacidophilic archaea, which are globally distributed in hot, acidic features including fumaroles (steam saturated discharges) as geysers and pools in the terrestrial environment, and hydrothermal vents in the marine environment (61). Both realms experience steep and rapidly changing gradients in temperature, pH, oxygen, and dissolved solute concentrations, which can disturb ecological niches (62). As a result, Archaea have adapted to or evolved strategies to mitigate these challenges (61, 63).

In *S. acidocaldarius*, our experimental manipulations of temperature, pH, shaking rate, electron acceptor (O_2) availability, and electron donor (sucrose) flux elicited a 20-fold change in growth rate and 2-fold change in mean biphytane cyclization (RI-BP), but $^2\varepsilon_{L/W}$ values were largely invariant (-204 ± 12 ‰, n = 22). This finding is consistent with previous studies that show archaeal lipids are consistently ^{2}H-depleted relative to source water and display a narrow range of $^2\varepsilon_{L/W}$ values despite considerable variation in taxonomy, metabolic mode, and growth conditions (10–15) (**FIG 6**).

In heterotrophic archaea, lipid-bound H can originate from water or the organic substrate used for growth, with intracellular hydride carriers, such as NADPH, mediating the incorporation of H from both sources into lipids (11, 33, 38). In heterotrophs, the NADPH incorporated into lipid products inherits some H from the available organic substrate(s). While there are several potential mechanisms by which each environmental stressor tested here could influence lipid/water fractionation in archaea, our data suggest these processes either do not occur or are insufficient to cause a measurable effect. Temperature, pH, oxygen levels, and nutrient availability are known to trigger physiological responses in archaea, including changes in growth rates, membrane lipid composition, and enzyme activity (47, 48, 52, 64). Enzyme kinetics, which are influenced by growth temperature and oxidative stress (65), could affect lipid/water fractionation if the enzyme catalyzing hydrogen transfer is associated with a significant isotope effect (33). High temperatures and low pH reduce water exchange across archaeal membranes to prevent osmotic stress (64), which could enrich the intracellular water in ^{2}H and potentially reduce apparent lipid/water fractionation (66). Additionally, changes in electron donor availability can shift the pathway by which NADPH is reduced, which could influence the δ^2H composition of NADPH, which ultimately supplies protons to lipid products (67, 68). For example, methanogenic archaea switch between the reductive acetyl-CoA and F_{420}-dependent hydrogenase systems depending on if electron donors (H_2 or formate) are abundant or limiting (67). Despite these possible mechanisms, lipid/water fractionation in *S. acidocaldarius* remains largely invariant in each experiment, indicating that these processes either

do not occur or are not significant enough to alter fractionation under the tested conditions. An
alternative hypothesis is that *S. acidocaldarius* maintains a constant internal metabolic balance,
particularly in NADPH production and utilization, regardless of external conditions.

Physiological responses to stressors impact growth rate and examining $^2\varepsilon_{L/W}$ values in response to
optimal vs. suboptimal growth may provide a more coherent explanation for the non-linear patterns
we observe across the environmental conditions tested. The $^2\varepsilon_{L/W}$ values for *S. acidocaldarius*,
however, are not correlated to growth rate in any single environmental condition experiment or when
data for all conditions are pooled. This finding suggests that cells maintain stable internal metabolic
processes resulting in a consistent H-balance over a range of growth rates and is consistent with the
theory that a constant lipid/water fractionation is a feature of an imperturbable central energy
metabolism (14, 15). *S. acidocaldarius*, like most archaea, employs energy-efficient metabolic
strategies (69, 70), possibly as an adaptation to energy-limiting environments (63, 71). In tightly
controlled metabolic pathways, the balance between supply and use of NADPH reducing power
appears to remain constant, even under energy-stress conditions. Previous research suggests
archaea may exist in a permanent state of NADPH deficiency such that all NADPH is immediately
used for biosynthesis with little to no flux from NADPH back to NADH (14). This efficiency in NADPH
use may explain why changing growth rates do not impact kinetic isotope effects associated with
biosynthesis and hydride exchange. NADPH deficiency (deemed NADPH flux imbalance by Wijker et
al., 2019) is associated with energy limitation, and can provide a framework to relate patterns in $^2\varepsilon_{L/W}$
values to the fluxes of H through dehydrogenase and transhydrogenase enzymes. Indeed, NADPH
flux imbalance may explain why all archaea (to-date) and autotrophic and obligately anaerobic
bacteria display negative $^2\varepsilon_{L/W}$ values of similar magnitude (15, 33, 36, 38). This similarity is striking
given that fatty acids and isoprenoid GDGTs are formed via different biosynthetic pathways and
suggests that the ultimate control on $^2\varepsilon_{L/W}$ values may be related to free energy yields (15).

Both the O_2 availability and sucrose flux experiments test the hypothesis that energy limitation –
either terminal electron acceptor or donor – can alter $^2\varepsilon_{L/W}$ values in archaea. These experiments
produced the largest range in doubling times (2 to 50 hours and 7 to 44 hours in the O_2 availability
and electron donor experiments, respectively); yet $^2\varepsilon_{L/W}$ values remained fairly consistent (-217 to -
196 ‰ and -276 to -244 ‰, respectively). *N. maritimus* is the only other archaeon in which the
relationship between electron donor (ammonium) flux and lipid/water fractionation was examined;
this archaeon similarly exhibited no significant correlation between a three-fold change in doubling
time and $^2\varepsilon_{L/W}$ values ($R^2 = 0.22$, *p* = 0.07, slope = 0.18 ± 0.09 ‰/ hour) (14). The authors speculate
this strain maintains a consistent H-balance over a range of electron donor fluxes as an adaptation
that allows cells to perform nitrification with energy-efficient CO_2 fixation in oligotrophic
environments (14). These findings suggest that archaea with diverse physiologies (e.g., heterotrophy
vs. chemoautotrophy, extremophily vs. mesophily) maintain a constant lipid-to-water fractionation
that is minimally affected by free energy availability and may reflect the fact that Archaea are
adapted to chronic energy limitation (63, 71). This stability may reflect tightly regulated physiological

processes, such as isoenzyme switching and modulation of enzyme activities, which help maintain
metabolic homeostasis under both optimal and sub-optimal growth conditions (72). In contrast, in
bacteria and some planktonic eukaryotes faster growth is associated with larger lipid/water
fractionation (66, 73, 74) and $^2\varepsilon_{L/W}$ values for Bacteria are influenced by electron donor flux (33, 36,
38).

The only other study in archaea to examine the relationship between environmental conditions and
lipid/water fractionation cultured the halophilic heterotrophic archaeon, *H. marismortui* (which
produces archaeol diethers), under varying temperatures and salinities when grown on complex rich
medium (11). The resulting $^2\varepsilon_{Archaeol/W}$ values are relatively invariant (ranging from -126 to -143 ‰;
more enriched than our observations for *S. acidocaldarius*) and are decoupled from growth rate (**FIG.**
S5; R^2 = 0.07, *p* = 0.7, slope = 0.85 ± 1.84 ‰ / hour), providing more evidence that archaea maintain
a stable internal H-balance across a range of growth conditions (11).

The impact of growth phase on H-isotope dynamics has not been studied in depth. The batch and
fed-batch experiment techniques used here for *S. acidocaldarius* and in the *H. marismortui* study
integrate across growth stages, whereas the chemostat electron donor flux experiments for *S.*
acidocaldarius and *N. maritimus* achieve constant growth rates. A recent study grew *A. fulgidus* in
batch cultures and found that $^2\varepsilon_{L/W}$ values were more depleted (larger offset from water) during
exponential phase relative to stationary phase (15). This pattern was consistent between
heterotrophic and autotrophic experiments. The culture conditions in Rhim et al. resulted in a
threefold difference in growth rates (similar change to the *N. maritimus* study, but much smaller
range than the 20-fold change we report for *S. acidocaldarius* here) and a 50 ‰ change in $^2\varepsilon_{L/W}$ values
(-280 ‰ to -230 ‰). Rhim and coauthors conclude that both metabolic mode (heterotrophic vs.
autotrophic growth) and growth phase impact $^2\varepsilon_{L/W}$ values, though the total variance among all
treatments was small and overlaps which ranges reported for other archaea (15).

4.2 Effect of biphytane ring number on archaeal lipid $^2\varepsilon_{L/W}$ values

Stoichiometric modeling predicts that BPs with more rings incorporate less of the highly-
fractionated hydrogen donated through GGR reduction, a process where the enzyme geranyl-geranyl
reductase (GGR) catalyzes the reduction of double bonds in isoprenoid chains, thereby affecting the
isotopic composition of the resulting lipids (14). Interestingly, in our *S. acidocaldarius* work, this
stepwise enrichment is only seen in the chemostat experiments, where each additional ring
contributes 7.4 ± 2.2 ‰ to biphytane $^2\varepsilon_{L/W}$ values. In the batch and fed-batch experiments, there is
no consistent dependence of $^2\varepsilon_{L/W}$ values on the number of biphytane moieties. In particular, the
Δε/ring trends differ between the shaking rate and O_2 availability experiments, which likely reflects
the fact that shaking rate can also impart stress via physical agitation in addition to increasing
oxygenation of microbial media (47).

iGDGT-derived BPs extracted from marine sediments also lack a clear stepwise enrichment with
increasing ring moieties (10), but this likely occurs because sediments integrate lipids synthesized
by different populations of archaea that are either growing on multiple substrates or in different
water masses (10). It is less clear why the pure *S. acidocaldarius* batch and fed-batch cultures do
not display a ring-specific pattern, though it perhaps relates to variations in biosynthetic
fractionations accompanying changes in lipid biosynthesis across the growth phases (44).
Considering all the available data, this enrichment pattern appears only minimally affected by
differences in growth rate and metabolism, as the magnitude of per-ring increase in $^2\varepsilon_{L/W}$ value is
similar for *Sulfolobus* sp. (10), *S. acidocaldarius* (chemostat experiment in this study), and *N.*
maritimus (14), which collectively represent a 5-fold range of growth rates, and a variety of
metabolisms and carbon substrates (*Sulfolobus* sp. = yeast extract + O_2; *S. acidocaldarius* = sucrose
+ O_2; *N. maritimus* = chemoautotroph on NH_4 + O_2). The magnitude of this stepwise enrichment,
however, is small and would likely have a minor impact (usually < 10 ‰) on the abundance weighted
mean $^2\varepsilon_{L/W}$ value even in cases where the BP Ring Index was markedly different among samples.

4.3 Archaeal lipid $^2\varepsilon_{L/W}$ as a function of central metabolism

The *S. acidocaldarius* $^2\varepsilon_{L/W}$ values we report are broadly similar to results from previous experimental
investigations of the *Sulfolobus* genus cultured heterotrophically. When grown at 75 °C and pH 2 on
yeast extract, a *Sulfolobus* strain displayed $^2\varepsilon_{L/W}$ values from -213 to -161 ‰ for BP-0 to -4
(abundance weighted mean = -180 ‰ (10) vs. -204 ± 12 ‰ for *S. acidocaldarius* here). In another
study, *Sulfolobus solfataricus* grown at 76 °C and pH 3.5 on glutamate displayed more enriched $^2\varepsilon_{L/W}$
values, averaging -134 ‰ when intact iGDGT-1 and -2 were analyzed (12). The difference in $^2\varepsilon_{L/W}$
values may in part be due to the structures analyzed. Lengger and colleagues analyzed intact iGDGTs
including the glycerols which may enrich the δ^2H_{iGDGT} composition, though the different substrate
may also play a role (12). Glutamate, an amino acid, is a more complex substrate than simple sugars
(such as sucrose) and primarily enters the *Sulfolobus* central metabolism through the TCA cycle (75).
Simple sugars enter metabolism primarily through the Entner-Doudoroff pathway, which converts
the sugar to intermediates that feed into the TCA cycle (75).

When *S. acidocaldarius* is grown heterotrophically on a simple sugar (glucose), the weighted mean
δ^2H composition of biphytane lipids is strongly correlated with δ^2H_{Water}, indicating a strong first-order
control (see *Supplementary Materials*, **FIG. S6**). A simple isotope mass balance model (**Eq. S1**)
suggests water contributes at least 56 ± 1% of the total H flux to lipids (**FIG. S7**). Further experimental
work and complementary isotope flux models are needed to determine the specific pathways (and
associated isotope effects) used during iGDGT synthesis in *S. acidocaldarius* (76).

Autotrophic archaea that produce iGDGTs, however, source their entire H budget to lipids from water
in the growth environment and show more negative $^2\varepsilon_{L/W}$ values than those we observe for *S.*
acidocaldarius. In the marine chemoautotroph, *N. maritimus*, grown in continuous culture, BP/water

fraction ranged from -272 to -260 ‰ (14). Similarly, BP/water fractionations for a metabolically-
flexible archaeon, *Archaeoglobus fulgidus*, ranged from -280 to -226 ‰ when cultured with several
unique carbon substrates and electron donor/acceptor pairs (15). This difference may be driven by
the distinct metabolic modes used (e.g., autotrophy vs. heterotrophy) and/or by unique metabolic
pathways that employ various intracellular hydride carriers and reductants to bring H's to the lipid
sites, which could generate different net biosynthetic fractionations. Metabolic flux models suggest
that in autotrophic archaea (e.g., *N. maritimus*), H-isotope fractionation due to the electron
transport chain and NADPH generation result in more depleted lipids than for heterotrophic archaea,
such as *Sulfolobus* species, because the source of NADPH impacts $^2\varepsilon_{L/W}$ values (14). In both
heterotrophic and aerobic archaea, most NADPH is generated by glucose dehydrogenase and
potentially glyceraldehyde dehydrogenase from the Entner-Doudoroff pathway (72, 77).

Overall, the range of $^2\varepsilon_{L/W}$ values in Archaeal lipids is narrower than the ranges observed in Bacteria
or Eukarya (**FIG. 6**). While the sample size for Bacteria and Eukarya are two-fold and four-fold larger
than for Archaea, respectively, Bartlett's test for homogeneity of variances confirm that Archaeal
$^2\varepsilon_{L/W}$ values have smaller variance compared to Eukarya ($K^2_{(1)}$ = 16.4, p < 0.0001) and Bacteria ($K^2_{(1)}$ =
118.5, p < 0.0001) after accounting for sample size. Reported $^2\varepsilon_{L/W}$ values for bacterial fatty acids
vary over 600 ‰ and include large negative and large positive values (15, 33, 36, 38, 78). In Eukarya,
$^2\varepsilon_{L/W}$ values are uniformly negative and span over 300 ‰, though certain lipid classes group together
(3, 25, 79). Lipids that are formed via the same biosynthetic pathway (e.g., acetogenic or isoprenoid
pathway) tend to occupy distinct ranges of $^2\varepsilon_{L/W}$ (15). Isoprenoid lipids produced via the MVA pathway
in eukaryotes have similar $^2\varepsilon_{L/W}$ values to iGDGT-derived biphytanes in archaea (15). This similarity
suggests that the MVA pathway used for isoprenoid lipid biosynthesis has a large net fractionation in
both domains of life.

4.4 Applications and Conclusion

Our calibration of biphytane δ^2H composition as a proxy for the δ^2H of environmental water in *S.*
acidocaldarius has significant implications for paleohydroclimate and paleoecology
reconstructions. First, this calibration may enable hydroclimate factors like precipitation, elevation,
and seasonality to be reconstructed for environments that do not contain traditionally analyzed lipid
biomarkers (e.g., plant waxes) – this includes extreme environments that do not support plant growth
(e.g., terrestrial hydrothermal systems, salars, and hyperarid regions). Archaeal lipid δ^2H values will
be most powerful as a proxy in cases where a single-substrate source is a reasonable assumption.

Second, biphytane δ^2H composition could be applied to marine or lacustrine sedimentary archives
where multiple sources of organic matter are suspected, such as a mix of plant waxes representing
in situ marine/ aquatic production and allochthonous terrestrial production (e.g., aeolian
deposition). In this case, plant wax δ^2H values would reflect a mixture of growth waters (marine/lake
water and soil water). In contrast, iGDGTs are likely derived solely from archaea living in the overlying

water column with few potential terrestrial sources and thus the δ^2H composition of iGDGTs would
only reflect marine/lake water (42).

Here, we show that in the model thermoacidophile *S. acidocaldarius* grown on simple sugars, lipid-
water fractionation ($^2\varepsilon_{L/W}$) is not dependent on any of the environmental conditions tested or on
growth rate. This finding suggests that *S. acidocaldarius* may have evolved stable metabolisms that
exist in a constant NADPH deficit, possibly reflecting adaptation to energy-limiting environments.
We conclude that *S. acidocaldarius* will likely demonstrate reliably invariant lipid/water fractionation
regardless of growth rate, nutrient status, or energy availability, and that lipid/water fractionation
may be more constant for certain archaeons than for phototrophic eukaryotes. For both domains,
however, there is more variability in lipid/water fraction relative to the spatial and temporal variability
in precipitation δ^2H values, and more research is needed to understand the sources of this variation
so that it can be accounted for in proxy applications.

This calibration underscores the potential for archaeal lipid δ^2H composition as a proxy for past
hydroclimates that could complement or be used independently of eukaryotic lipid δ^2H
composition. Future research should focus on exploring the effects of growth phase and metabolic
pathways on lipid/water fractionation in *S. acidocaldarius* and other archaeal species. Expanding
these investigations to include more diverse environmental conditions, such as varying carbon
substrates or mixed microbial communities, could further validate the consistency of $^2\varepsilon_{L/W}$ values
and refine its utility as a hydrological proxy. Integrating isotope flux models with experimental data
such as those generated here, will allow us to better elucidate the specific biochemical pathways
responsible for the observed fractionation patterns in domain Archaea.

Acknowledgements

This research was supported by research grant NSF EAR #1928303940 (WDL, SHK); Simons
Foundation Award #623881 (WDL); Dartmouth College (WDL); National Aeronautics and Space
Administration (NASA) New Hampshire Space Grant (NASA-SSW-80NSSC19K0539); Deutsche
Forschungsge-meinschaft grant 441217575 (F.J.E.). The authors thank the CU Boulder Earth
Systems Stable Isotope Lab (CUBES-SIL) Core Facility (RRID:SCR_019300) for analytical
contributions; B. Chiu (then at Dartmouth), J. Blewett (then at Harvard), S. Carter (Harvard), and A.
Maloney (CU Boulder) for expert laboratory support; Simon D. Kelly from International Atomic Energy
Agency, Vienna for measuring the δ^2H composition of carbon substrates; and all members of the
Leavitt lab for all their support.

References

1.   Gat JR. 1996. Oxygen and hydrogen isotopes in the hydrologic cycle. Annual Review of Earth
and Planetary Sciences 24:225–262.

2.   Hayes JM. 2001. Fractionation of carbon and hydrogen isotopes in biosynthetic processes.
Stable Isotope Geochemistry 43:225–277.

3.   Sessions AL, Burgoyne TW, Schimmelmann A, Hayes JM. 1999. Fractionation of hydrogen
isotopes in lipid biosynthesis. Organic Geochemistry 30:1193–1200.

4.   Sessions AL, Sylva SP, Summons RE, Hayes JM. 2004. Isotopic exchange of carbon-bound
hydrogen over geologic timescales. Geochimica et Cosmochimica Acta 68:1545–1559.

5.   Dawson D, Grice K, Wang SX, Alexander R, Radke J. 2004. Stable hydrogen isotopic
composition of hydrocarbons in torbanites (Late Carboniferous to Late Permian) deposited
under various climatic conditions. Organic Geochemistry 35:189–197.

6.   Love GD, Zumberge JA. 2021. Emerging Patterns in Proterozoic Lipid Biomarker Records, 1st
ed. Cambridge University Press.
https://www.cambridge.org/core/product/identifier/9781108847117/type/element. Retrieved
22 March 2024.

7.   Liu J, An Z. 2019. Variations in hydrogen isotopic fractionation in higher plants and sediments
across different latitudes: Implications for paleohydrological reconstruction. Science of The
Total Environment 650:470–478.

8.   McFarlin JM, Axford Y, Masterson AL, Osburn MR. 2019. Calibration of modern sedimentary
δ2H plant wax-water relationships in Greenland lakes. Quaternary Science Reviews
225:105978.

9.   Ladd SN, Nelson DB, Bamberger I, Daber LE, Kreuzwieser J, Kahmen A, Werner C. 2021.
Metabolic exchange between pathways for isoprenoid synthesis and implications for
biosynthetic hydrogen isotope fractionation. New Phytologist 231:1708–1719.

10. Kaneko M, Kitajima F, Naraoka H. 2011. Stable hydrogen isotope measurement of archaeal
ether-bound hydrocarbons. Organic Geochemistry 42:166–172.

11. Dirghangi SS, Pagani M. 2013. Hydrogen isotope fractionation during lipid biosynthesis by
Haloarcula marismortui. Geochimica et Cosmochimica Acta 119:381–390.

12. Lengger SK, Weber Y, Taylor KWR, Kopf S, Berstan R, Bull I, Mayser J, Leavitt WD, Blewett J,
Pearson A, Pancost RD. 2020. Determination of the d2H values of high molecular weight lipids

by high temperature GC coupled to isotope ratio mass spectro. Rapid Communications in
Mass Spectrometry.

13.   Wu W, Meador TB, Könneke M, Elvert M, Wegener G, Hinrichs K. 2020. Substrate-dependent
incorporation of carbon and hydrogen for lipid biosynthesis by Methanosarcina barkeri.
Environmental Microbiology Reports https://doi.org/10.1111/1758-2229.12876.

14.   Leavitt WD, Kopf SH, Weber Y, Chiu B, McFarlin JM, Elling FJ, Hoeft-McCann S, Pearson A.
2023. Controls on the hydrogen isotope composition of tetraether lipids in an autotrophic
ammonia-oxidizing marine archaeon. Geochimica et Cosmochimica Acta 352:194–210.

15.   Rhim JH, Kopf S, McFarlin J, Maloney AE, Batther H, Harris CM, Zhou A, Feng X, Weber Y,
Hoeft-McCann S, Pearson A, Leavitt WD. 2024. Metabolic imprints in the hydrogen isotopes of
Archaeoglobus fulgidus tetraether lipids. Geochimica et Cosmochimica Acta 386:196–212.

16.   Sachs J. 2014. Hydrogen Isotope Signatures in the Lipids of Phytoplankton, p. 79–94. *In*
Treatise on Geochemistry. Elsevier.

17.   Gao L, Edwards EJ, Zeng Y, Huang Y. 2014. Major Evolutionary Trends in Hydrogen Isotope
Fractionation of Vascular Plant Leaf Waxes. PLoS ONE 9:e112610.

18.   Bender, Suess, Chitwood, Bradley. 2016. Investigating genetic loci that encode plant-derived
paleoclimate proxies, p. PP24B-03. *In* .

19.   Baan J, Holloway-Phillips M, Nelson DB, Kahmen A. 2023. Species and biosynthetic effects
cause uncorrelated variation in oxygen and hydrogen isotope compositions of plant organic
compounds. Geochimica et Cosmochimica Acta 352:1–13.

20.   Baan J. 2024. Understanding the metabolic signal in hydrogen isotope values of plant organic
compounds. Thesis der Universitat Basel https://doi.org/DOI 10.5451/unibas-ep96717.

21.   Ladd N, Nelson D, Matthews B, Dyer S, Limberger R, Klatt A, Narwani A, Dubois N, Schubert C.
2024. Taxon-specific hydrogen isotope signals in cultures and mesocosms facilitate
ecosystem and hydroclimate reconstruction. EarthArXiv https://doi.org/10.31223/X57T30.

22.   Pilecky M, Kainz MJ, Wassenaar LI. 2024. Exploring hydrogen isotope fractionation in lipid
biomolecules of freshwater algae: implications for ecological and paleoenvironmental
studies. Isotopes in Environmental and Health Studies 60:585–595.

23.   Sachs JP, Kawka OE. 2015. The influence of growth rate on 2H/1H fractionation in continuous
cultures of the coccolithophorid Emiliania huxleyi and the diatom Thalassiosira pseudonana.
PLoS ONE 10:1–27.

24. Van Der Meer MTJ, Benthien A, French KL, Epping E, Zondervan I, Reichart G-J, Bijma J,
591   Sinninghe Damsté JS, Schouten S. 2015. Large effect of irradiance on hydrogen isotope
592   fractionation of alkenones in Emiliania huxleyi. Geochimica et Cosmochimica Acta 160:16–
593   24.

594 25. Sachs JP, Maloney AE, Gregersen J, Paschall C. 2016. Effect of salinity on 2H/1H fractionation
595   in lipids from continuous cultures of the coccolithophorid Emiliania huxleyi. Geochimica et
596   Cosmochimica Acta 189:96–109.

597 26. Maloney AE, Shinneman ALC, Hemeon K, Sachs JP. 2016. Exploring lipid 2H/1H fractionation
598   mechanisms in response to salinity with continuous cultures of the diatom Thalassiosira
599   pseudonana. Organic Geochemistry 101:154–165.

600 27. Wolfshorndl M, Danford R, Sachs JP. 2019. 2H/1H fractionation in microalgal lipids from the
601   North Pacific Ocean: Growth rate and irradiance effects. Geochimica et Cosmochimica Acta
602   246:317–338.

603 28. Torres-Romero I, Zhang H, Wijker RS, Clark AJ, McLeod RE, Jaggi M, Stoll HM. 2024. Hydrogen
604   isotope fractionation is controlled by CO_2 in coccolithophore lipids. Proc Natl Acad Sci USA
605   121:e2318570121.

606 29. Sachse D, Billault I, Bowen GJ, Chikaraishi Y, Dawson TE, Feakins SJ, Freeman KH, Magill CR,
607   McInerney FA, Van Der Meer MTJ, Polissar P, Robins RJ, Sachs JP, Schmidt HL, Sessions AL,
608   White JWC, West JB, Kahmen A. 2012. Molecular paleohydrology: Interpreting the hydrogen-
609   isotopic composition of lipid biomarkers from photosynthesizing organisms. Annual Review of
610   Earth and Planetary Sciences 40:221–249.

611 30. Sessions AL. 2016. Factors controlling the deuterium contents of sedimentary hydrocarbons.
612   Organic Geochemistry 96:43–64.

613 31. Sessions AL, Jahnke LL, Schimmelmann A, Hayes JM. 2002. Hydrogen isotope fractionation in
614   lipids of the methane-oxidizing bacterium Methylococcus capsulatus. Geochimica et
615   Cosmochimica Acta 66:3955–3969.

616 32. Valentine DL. 2009. Isotopic remembrance of metabolism past. Proceedings of the National
617   Academy of Sciences of the United States of America 106:12565–12566.

618 33. Zhang X, Gillespie AL, Sessions AL. 2009. Large D/H variations in bacterial lipids reflect
619   central metabolic pathways. Proceedings of the National Academy of Sciences of the United
620   States of America 106:12580–12586.

621 34. Dawson KS, Osburn MR, Sessions AL, Orphan VJ. 2015. Metabolic associations with archaea
622   drive shifts in hydrogen isotope fractionation in sulfate-reducing bacterial lipids in cocultures
623   and methane seeps. Geobiology 13:462–477.

35.   Leavitt WD. 2016. The Inner Lives of Archaea: the hydrogen isotopic composition of Archaeal
lipids as a proxy of metabolic state.

36.   Osburn MR, Dawson KS, Fogel ML, Sessions AL. 2016. Fractionation of hydrogen isotopes by
sulfate- and nitrate-reducing bacteria. Frontiers in Microbiology 7:1–16.

37.   Leavitt WD, Murphy SJL, Lynd LR, Bradley AS. 2017. Hydrogen isotope composition of
Thermoanaerobacterium saccharolyticum lipids: Comparing wild type with a nfn-
transhydrogenase mutant. Organic Geochemistry 113:239–241.

38.   Wijker RS, Sessions AL, Fuhrer T, Phan M. 2019. 2H/1H variation in microbial lipids is
controlled by NADPH metabolism. Proceedings of the National Academy of Sciences of the
United States of America 116:12173–12182.

39.   Van de Vossenberg JLCM, Driessen AJM, Konings WN. 1998. The essence of being
extremophilic: The role of the unique archaeal membrane lipids. Extremophiles 2:163–170.

40.   Schouten S, Hopmans EC, Sinninghe Damsté JS. 2013. The organic geochemistry of glycerol
dialkyl glycerol tetraether lipids: A review. Organic Geochemistry 54:19–61.

41.   Schouten S, Hopmans EC, SchefuB E, Sinninghe Damsté JS. 2002. Distributional variations in
marine crenarchaeotal membrane lipids: a new tool for reconstructing ancient sea water
temperatures? Earth and Planetary Science Letters 265–274.

42.   Pearson A, Ingalls AE. 2013. Assessing the use of archaeal lipids as marine environmental
proxies. Annual Review of Earth and Planetary Sciences 41:359–384.

43.   De Rosa M, Gambacorta A. 1988. The lipids of archaebacteria. Progress in Lipid Research
27:153–175.

44.   Elling FJ, Könneke M, Lipp JS, Becker KW, Gagen EJ, Hinrichs K-U. 2014. Effects of growth
phase on the membrane lipid composition of the thaumarchaeon Nitrosopumilus maritimus
and their implications for archaeal lipid distributions in the marine environment. Geochimica
et Cosmochimica Acta 141:579–597.

45.   Elling FJ, Könneke M, Nicol GW, Stieglmeier M, Bayer B, Spieck E, de la Torre JR, Becker KW,
Thomm M, Prosser JI, Herndl GJ, Schleper C, Hinrichs KU. 2017. Chemotaxonomic
characterisation of the thaumarchaeal lipidome. Environmental Microbiology 19:2681–2700.

46.   Hurley SJ, Elling FJ, Könneke M, Buchwald C, Wankel SD, Santoro AE, Lipp JS, Hinrichs KU,
Pearson A. 2016. Influence of ammonia oxidation rate on thaumarchaeal lipid composition
and the TEX86 temperature proxy. Proceedings of the National Academy of Sciences of the
United States of America 113:7762–7767.

47. Cobban A, Zhang Y, Zhou A, Pearson A, Leavitt WD. 2020. Multiple environmental parameters
impact core lipid cyclization in Sulfolobus acidocaldarius. Environmental Microbiology 1–28.

48. Zhou A, Weber Y, Chiu BK, Elling FJ, Cobban AB, Pearson A, Leavitt WD. 2020. Energy flux
controls tetraether lipid cyclization in Sulfolobus acidocaldarius. Environmental Microbiology
22:343–353.

49. Quehenberger J, Pittenauer E, Allmaier G, Spadiut O. 2020. The influence of the specific
growth rate on the lipid composition of Sulfolobus acidocaldarius. Extremophiles 24:413–420.

50. Yang W, Chen H, Chen Y, Chen A, Feng X, Zhao B, Zheng F, Fang H, Zhang C, Zeng Z. 2023.
Thermophilic archaeon orchestrates temporal expression of GDGT ring synthases in response
to temperature and acidity stress. Environmental Microbiology 25:575–587.

51. Brock TD, Brock KM, Belly RT, Weiss RL. 1972. Sulfolobus: A new genus of sulfur-oxidizing
bacteria living at low pH and high temperature. Archiv Mikrobiol 84:54–68.

52. Bischof LF, Haurat MF, Hoffmann L, Albersmeier A, Wolf J, Neu A, Pham TK, Albaum SP, Jakobi
T, Schouten S, Neumann-Schaal M, Wright PC, Kalinowski J, Siebers B, Albers S-V. 2019. Early
Response of Sulfolobus acidocaldarius to Nutrient Limitation. Front Microbiol 9:3201.

53. Quehenberger J, Albersmeier A, Glatzel H, Hackl M, Kalinowski J, Spadiut O. 2017. Sulfolobus
– A Potential Key Organism in Future Biotechnology. Frontiers in Microbiology 301:56–67.

54. Boyd ES, Pearson A, Pi Y, Li W-J, Zhang YG, He L, Zhang CL, Geesey GG. 2011. Temperature
and pH controls on glycerol dibiphytanyl glycerol tetraether lipid composition in the
hyperthermophilic crenarchaeon Acidilobus sulfurireducens. Extremophiles 15:59–65.

55. Jensen SM, Brandl M, Treusch AH, Ejsing CS. 2015. Structural characterization of ether lipids
from the archaeon *Sulfolobus islandicus* by high-resolution shotgun lipidomics: Mapping the
lipidome of *Sulfolobus islandicus*. J Mass Spectrom 50:476–487.

56. Wagner M, Van Wolferen M, Wagner A, Lassak K, Meyer BH, Reimann J, Albers S-V. 2012.
Versatile Genetic Tool Box for the Crenarchaeote Sulfolobus acidocaldarius. Front Microbio 3.

57. Zhang YG, Pagani M, Wang Z. 2016. Ring Index: A new strategy to evaluate the integrity of TEX
$_{86}$ paleothermometry. Paleoceanography 31:220–232.

58. Abrahim A, Cannavan A, Kelly SD. 2020. Stable isotope analysis of non-exchangeable
hydrogen in carbohydrates derivatised with N-methyl-bis-trifluoroacetamide by gas
chromatography – Chromium silver reduction/High temperature Conversion-isotope ratio
mass spectrometry (GC-CrAg/HTC-IRMS). Food Chemistry 318:126413.

59.   Kopf S, Davidheiser-Kroll B, Kocken I. 2021. Isoreader: An R package to read stable isotope
data files for reproducible research. JOSS 6:2878.

60.   Carolynn Harris. 2025. carolynnharris/Saci_Enviro_d2H: Supplementary material for Harris et
al., 2025, AEM (v.1.0.0). Zenodo.

61.   Lewis AM, Recalde A, Bräsen C, Counts JA, Nussbaum P, Bost J, Schocke L, Shen L, Willard
DJ, Quax TEF, Peeters E, Siebers B, Albers S-V, Kelly RM. 2021. The biology of
thermoacidophilic archaea from the order Sulfolobales. FEMS Microbiology Reviews 1–60.

62.   Reysenbach A-L, Cady SL. 2001. Microbiology of ancient and modern hydrothermal systems.
Trends in Microbiology 9:79–86.

63.   Auernik KS, Cooper CR, Kelly RM. 2008. Life in hot acid: pathway analyses in extremely
thermoacidophilic archaea. Current Opinion in Biotechnology 19:445–453.

64.   Siliakus MF, Van Der Oost J, Kengen SWM. 2017. Adaptations of archaeal and bacterial
membranes to variations in temperature, pH and pressure. Extremophiles 21:651–670.

65.   Jia B, Liu J, Van Duyet L, Sun Y, Xuan YH, Cheong G-W. 2015. Proteome profiling of heat,
oxidative, and salt stress responses in Thermococcus kodakarensis KOD1. Front Microbiol 6.

66.   Sachs JP, Schwab VF. 2011. Hydrogen isotopes in dinosterol from the Chesapeake Bay
estuary. Geochimica et Cosmochimica Acta 75:444–459.

67.   Thauer RK, Kaster A-K, Seedorf H, Buckel W, Hedderich R. 2008. Methanogenic archaea:
ecologically relevant differences in energy conservation. Nat Rev Microbiol 6:579–591.

68.   Spaans SK, Weusthuis RA, Van Der Oost J, Kengen SWM. 2015. NADPH-generating systems in
bacteria and archaea. Front Microbiol 6.

69.   Wakagi T, Oshima T. 1987. Energy metabolism of a thermoacidophilic
archaebacterium,Sulfolobus acidocaldarius. Origins Life Evol Biosphere 17:391–399.

70.   Liu L-J, Jiang Z, Wang P, Qin Y-L, Xu W, Wang Y, Liu S-J, Jiang C-Y. 2021. Physiology, Taxonomy,
and Sulfur Metabolism of the Sulfolobales, an Order of Thermoacidophilic Archaea. Front
Microbiol 12:768283.

71.   Valentine DL. 2007. Adaptations to energy stress dictate the ecology and evolution of the
Archaea. Nature Reviews Microbiology 5:316–323.

72.   Bräsen C, Esser D, Rauch B, Siebers B. 2014. Carbohydrate Metabolism in Archaea: Current
Insights into Unusual Enzymes and Pathways and Their Regulation. Microbiol Mol Biol Rev
78:89–175.

73.   Schouten S, Ossebaar J, Schreiber K, Kienhuis MVM, Langer G, Benthien A, Bijma J. 2006. The
effect of temperature, salinity and growth rate on the stable hydrogen isotopic composition of
long chain alkenones produced by Emiliania huxleyi and Gephyrocapsa oceanica.
Biogeosciences 3:113–119.

74.   Zhang Z, Sachs JP, Marchetti A. 2009. Hydrogen isotope fractionation in freshwater and
marine algae: II. Temperature and nitrogen limited growth rate effects. Organic Geochemistry
40:428–439.

75.   Quehenberger J, Albersmeier A, Glatzel H, Hackl M, Kalinowski J, Spadiut O. 2019. A defined
cultivation medium for Sulfolobus acidocaldarius. Journal of Biotechnology 301:56–67.

76.   Harris C, Rhim J, Zhang Y, Cobban A, McFarlin J, Batther H, Kopf S, Leavitt W. 2022.
Determining controls on hydrogen isotope fractionation in archaeal tetraether lipids in a
thermoacidophilic archaeal heterotrophGoldschmidt2022 abstracts. European Association
of Geochemistry, Honolulu, HI, USA.

77.   Nunn CEM, Johnsen U, Schönheit P, Fuhrer T, Sauer U, Hough DW, Danson MJ. 2010.
Metabolism of Pentose Sugars in the Hyperthermophilic Archaea Sulfolobus solfataricus and
Sulfolobus acidocaldarius. Journal of Biological Chemistry 285:33701–33709.

78.   Campbell BJ, Li C, Sessions AL, Valentine DL. 2009. Hydrogen isotopic fractionation in lipid
biosynthesis by H2-consuming Desulfobacterium autotrophicum. Geochimica et
Cosmochimica Acta 73:2744–2757.

79.   Zhang Z, Sachs JP. 2007. Hydrogen isotope fractionation in freshwater algae: I. Variations
among lipids and species. Organic Geochemistry 38:582–608.

Tables

Table 1. Summary of culture conditions and descriptive growth statistics for S. acidocaldarius grown
over varying environmental conditions. δ^2H_W is the H-isotope composition of media water at the time
of lipid sampling. *Growth d*ata for batch and fed-batch environmental condition experiments were
initially reported in Cobban et al., (2020), *growth* data for chemostat experiment was initially
reported in Zhou et al., (2019). *For batch experiments, RPM is shaking rate; for bioreactor
experiments, RPM is impellor speed.

Table 2. Descriptive statistics for individual biphytanyl (BP) lipids derived from iGDGTs for *S.*
acidocaldarius grown under varying environmental conditions. Mean and sd include biological
replication (N ≥ 1) and technical replication (e.g., multiple injections, n ≥ 3). Abundance-weighted
means are shown for δ^2H_{BP}, $^2\varepsilon_{L/W}$, and $\Delta\varepsilon$/ring values with propagated error.

Figures

Fig 1. (A) Representative chromatogram of intensity vs. retention time for four iGDGT-derived
biphytane (BP) moieties identified in *S. acidocaldarius* biomass that are named for the number of
cyclopentane rings they contain. Although a BP-3 peak was detected, in many cases it is present in
too low quantities to perform isotopic measurements **(B)** The identity and structure of BP-0 to BP-3.

Fig. 2. Results for *S. acidocaldarius* grown under varying environmental conditions. Top row (**A-E**): Doubling time in response to each environmental condition. Second row (**F-J**): Relative abundance of biphytane (BP) lipids derived from iGDGTs. BP-3 was the least abundant and was below the detection limit in some samples. Bars represent the mean across replicates for each treatment level. Third row (**K-O**): Ring Index (RI-BP) of biphytane lipids derived from iGDGTs. Bottom row (**P-T**) The abundance weighted mean H-isotope fractionation ($^2\varepsilon_{L/W}$) between growth water and biphytanes. For all rows, points represent the mean and propagated error across biological and technical replicates for each treatment level. In many cases, error bars are smaller than symbols. Only statistically significant regressions are shown ($p < 0.05$).

Fig. 3. (**A-E**) Ring difference (Δε/ring, ‰) for individual biphytanes in each environmental condition
experiment. BP-0 has a Δε/ring of 0 by definition and is shown for reference. BP-3 was not
recovered from every treatment. Note the axes are scaled differently for each plot. Points represent
mean of biological and technical replicates for each treatment level. (**F**) Ring difference for
biphytanes examined in other studies. Blue diamonds are *N. maritimus* from Leavitt et al., 2023;
red squares are *Sulfolobus* sp. from Kaneko et al., 2011. Kaneko et al., also recovered BP-4 from
Sulfolobus spp. (data not shown); the Δε/ring for BP-4 is 11.7 ‰.

Fig. 4. Correlation matrices showing linear relationships among environmentally manipulated parameters, growth statistics (doubling time (T_D) and maximum OD_{600}), mean cyclization of lipids (RI-GDGT and RI-BP), ring difference, and abundance weighted mean $^2\varepsilon_{L/W}$ for (**A**) temperature, (**B**) pH, (**C**) aeration rate, (**D**) O_2 mixing ratio, and (**E**) electron donor flux. Notably, mean $^2\varepsilon_{L/W}$ values do not show a strong linear correlation to any other measured parameter, indicating that environmental conditions, growth statistics, and mean cyclization do not significantly influence the degree of lipid/water fractionation in *S. acidocaldarius*. Circle color and size indicate the strength and direction (blue = correlated, red = anticorrelated) of the relationship. Significance levels: * $p < 0.05$, ** $p < 0.001$, *** $p < 0.0001$. Circles without a star indicate the relationship is not statistically significant.

Fig. 5. The abundance weighted H-isotope fractionation ($^2\varepsilon_{L/W}$) between growth water and
biphytanes (BPs) in response to doubling time (hours) in each individual environmental condition
experiment (**A-E**) and pooled across all experiments (**F**). When data from all environmental control
experiments is pooled, there is no relationship between doubling time (hours) and abundance
weighted mean $^2\varepsilon_{L/W}$. Points represent mean of biological and technical replicates for each
treatment level. Error bars represent propagated error associated with BP abundance
measurements, BP-δ^2H measurements, and weighted mean $^2\varepsilon_{L/W}$ calculations; bars are smaller
than the symbols in many cases.

Fig. 6. Summary of lipid/water fractionation ($^2\varepsilon_{L/W}$) in culture studies across three domains of life:
eukaryotic phototrophs (green; n = 270), bacteria (blue; n = 140), and archaea (pink; n = 72). Tick
marks indicate individual $^2\varepsilon_{L/W}$ values; shaded curves indicate probability density functions.
Archaea data includes measurements of phytanes and biphytanes from *Sulfolobus* sp. (Kaneko et
al., 2011), *M. barkeri* (Wu et al., 2020), *N. maritimus* (Leavitt et al., 2022), *Acidianus* sp. DS80, *A.
fulgidus, and *Metallosphaera sedula* (Rhim et al., 2024), and *S. acidocaldarius* (this study; black
tick marks). Data for Eukarya and Bacteria is redrawn from Rhim et al., (2024), Figure 6. Eukarya
data includes measurements for acetogenic and isoprenoid lipids. Bacteria data includes fatty
acids synthesized by multiple metabolisms (e.g., aerobic and anaerobic heterotrophy, aerobic and
anaerobic chemoautotrophy, and photoautotrophy).