1 **A universal hippocampal memory code across animals and environments**

3 Hannah S Wirtshafter[1]*, Sara A Solla[1], John F Disterhoft[1]

9 *Lead contact: hsw@northwestern.edu (HSW)

17 1. Department of Neuroscience, Northwestern University Feinberg School of Medicine, Chicago,

18 IL, USA.

How learning is affected by context is a fundamental question of neuroscience, as the ability to generalize learning to different contexts is necessary for navigating the world. An example of swift contextual generalization is observed in conditioning tasks, where performance is quickly generalized from one context to another. A key question in identifying the neural substrate underlying this ability is how the hippocampus (HPC) represents task-related stimuli across different environments, given that HPC cells exhibit place-specific activity that changes across contexts (remapping). In this study, we used calcium imaging to monitor hippocampal neuron activity as rats performed a conditioning task across multiple spatial contexts. We investigated whether hippocampal cells, which encode both spatial locations (place cells) and task-related information, could maintain their task representation even when their spatial encoding remapped in a new spatial context. To assess the consistency of task representations, we used advanced dimensionality reduction techniques combined with machine learning to develop manifold representations of population level HPC activity. The results showed that task-related neural representations remained stable even as place cell representations of spatial context changed, thus demonstrating similar embedding geometries of neural representations of the task across different spatial contexts. Notably, these patterns were not only consistent within the same animal across different contexts but also significantly similar across different animals, suggesting a standardized neural encoding or 'neural syntax' in the hippocampus. These findings bridge a critical gap between memory and navigation research, revealing how the hippocampus maintains cognitive consistency across different spatial environments. These findings also suggest that hippocampal function is governed by a neural framework shared between animals, an observation that may have broad implications for understanding memory, learning, and related cognitive processes. Looking ahead, this work opens new avenues for exploring the fundamental principles underlying hippocampal encoding strategies.

Introduction

How can learning can be generalized across contexts as well as remaining localized to one

context? This question is fundamental to both neuroscience and philosophy[1-4]. Deficits in

generalization or inappropriate generalization are hallmarks of many disorders, including

autism[5,6], schizophrenia[7,8], and post-traumatic stress disorder[9,10]. In spite of their importance,

many questions related to generalization remain to be answered.

The hippocampus (HPC) is important for learning, memory, and navigation, and damage to

this region can disrupt contextual learning[11-17]. Many aspects of context, including an animal's

spatial location and the presence of local and distal cues, can be represented by 'place cells' in

the HPC[18-22]. The activity of many HPC cells therefore changes drastically in different

environments (i.e. place cells remap), even when a task can be generalized across these different

contexts[23-25]. A major open question is if and how representations of task-relevant stimuli, which

are also found in the HPC[26-31], can be maintained against the background of remapped place

cells. A further question is whether different animals solve this problem using the same, or

similar, neural strategies.

Until recently, the neural mechanisms behind contextual learning have been challenging to

investigate due to the need for tracking large numbers of cells across various environments and

learning stages— tasks which were unachievable prior to the development of calcium imaging[32-

37]. The use of calcium imaging in this study has allowed for a detailed exploration of

hippocampal neuron dynamics, bridging previous gaps between studies of hippocampal

cognitive maps and the hippocampal bases of memory and learning. The traditional separation

between memory and navigation fields of HPC research is thus in the process of narrowing[38,39].

This research addresses two fundamental questions of learning and memory. The first

question pertains to the persistence of learning across varying contexts. The modulation of

hippocampal cells by spatial variables results in substantial changes to cell activity in different

environments[24,25,40,41]; how can conditioning-related neural representations remain stable amid

such remapping? The second question is the extent to which neural representations in the

hippocampus are invariant and consistent, not only within an individual across diverse contexts,

but across different individuals; is there a standardized neural encoding or 'neural syntax' in the

for learning and memory in the hippocampus? A commonality of encoding across animals would

imply that the functionality of the HPC is informed not solely by individual experiences but also

by a standardized framework of neural algorithms. Such a finding would provide key insights

into the underlying neural mechanisms that govern learning and memory, helping to identify

specific brain circuits and algorithms that drive behavior.

To answer these questions, we trained rats on an HPC-dependent conditioning task which is

rapidly generalized between spatial contexts[42,43]. We examined the same task in disparate

contexts and looked for changes in a representation of spatial location provided by a population

of place cells while the representation of the task features remains unchanged. We found that the

representations of the conditioning task were maintained as the animal generalized learning from

one environment to another, even as the representation of place changed. Surprisingly, we also

found that the neural representation of the task was consistent across animals.

This study demonstrates that despite the well-known phenomenon of place cell

remapping[24,25,40,41], there exists a stable population-level neural representation of task features

that persists across diverse environments. These representations also show consistency across

different individuals, indicating a standardized neural encoding or 'neural syntax' for the

conditioning task within the hippocampus. This novel finding suggests the existence of a

universal coding mechanism for associative learning in the hippocampus, a principle that could

reorient our approach to studying memory and could challenge our current understanding of

cognitive processes.

Results

We trained a cohort of 5 freely moving rats in a conditioning task in one of two distinct

environments, labeled A and B; the rats had been previously familiarized with the environment

by the time the training began (Fig. 1a, Fig. S1). Environment A was an unscented rectangular

enclosure with wire floor and walls, and white lighting; environment B was a scented ovular

enclosure with white solid floor and walls, and red lighting (Fig. 1b). Both environments were

located at the same spot in the room relative to external cues (see Methods); animals could see

external cues out of the top of both environments, as well as out of the sides of environment A

(animals also reared often, allowing them to see out of the sides of environment B). During

training sessions, we recorded cellular activity in the hippocampal CA1 region via miniscopes,

using Gcamp8m for calcium imaging (CaImg). To perform data analysis, we used both calcium

events and calcium traces (see Methods), as indicated when applicable.

Animals easily transfer a conditioning task across environments

During the initial phase of our study, freely-moving rats (Fig. 1b) underwent training for

trace eyeblink conditioning (tEBC) (Fig. S2), a hippocampus-dependent classical conditioning

task that serves as a robust model for associative memory formation[44-47]. This paradigm involves

presenting a 250ms conditioned stimulus (CS, in the form of a tone) followed by a 500ms trace

interval, followed by the 100ms presentation of an unconditioned stimulus (US, an eyelid shock)

(Fig. 1c). Both shock and blinking were recorded with wires inserted into the muscle of the

eyelid (see Methods). As rats were trained, they exhibited a conditioned blink (CR) to the tone.

Animals were considered to have learned the task after reaching criterion (70% CRs in 50 trials)

on three consecutive training sessions (termed 'criterion sessions') or when the previous four

training sessions averaged over 70% (in this instance, only the final three of those sessions were

considered 'criterion sessions') (Fig. 1d). There was substantial variability in the number of

sessions it took to learn the task, for an average of 20±4.2 training sessions (note: number of

sessions always includes criterion sessions). The two rats that learned the fastest reached

criterion in 14 sessions, and the rat that learned the slowest reached criterion after 24 sessions

(Fig. S2). After reaching criterion, the rats were introduced to environment B, where their ability

to perform tEBC was assessed over a two-day period (one session per day). Comparative

analysis revealed no significant difference in performance (measured in % CRs) between the

criterion sessions in environment A and the testing phase in environment B (mean in

environment A criterion sessions was 74.75±6.49, mean in environment B test sessions was

77.70±11.68, two tailed t-test(24)=-0.83, p>0.05) (Fig. 1d), indicating the successful transfer of

tEBC learning to a new environment.

Calcium imaging (CaImg) enabled the longitudinal monitoring of the same hippocampal

cells over multiple sessions in both environments. For criterion and testing sessions, we observed

an average of 459.85±265.31 cells per session per animal, with no significant difference between

the number of cells recorded in environment A and environment B (two-tailed t-test(24)= -0.56,

p>0.05) On average, 132±95 cells were present in both the last criterion session in A and the first

testing session in B. This was not a significantly different numbers of cells that were present, on

average, in both the semi-final session in A, session A(n-1), and the final session in A, session

A(n) (155±115 cells).

Hippocampal place cell representations differ across environments

To identify place cells, we compared actual mutual information (MI) to MI computed

after bootstrapping circularly shifted position data 500 times. A cell was deemed a place cell if

its MI was above the 95th percentile of null provided by the bootstrapped MI scores. Using this

95% cutoff, on average, 9.3%±4.2% of cells across criterion sessions in A and testing sessions in

B qualified as place cells. There was not a significant difference between the percent of cells that

qualified as place cells in environment A and environment B (average in environment A was

8.1%±3.3%, average in environment B was 11.1%±4.9%, two-tailed t-test t(23) =-1.9, p>0.05)

(Fig. 2a, Fig. S3). The average mutual information scores across all criterion and testing sessions

was 1.08±0.18, with no difference in average MI between environments (environment A mean

MI was 1.08±0.17, environment B mean MI was 1.08±0.21, two-tailed t-test t(4805)=-0.5,

p>0.05). (See methods for more detailed analysis).

Our analysis revealed that individual hippocampal (HPC) cells exhibited distinct spatial

representations for environments A and B, altering their configurations of place cells and place

fields relative to distal cues — a process known as 'place cell remapping' (Fig. 2b). We

confirmed this remapping through several approaches. First, we quantified the shift in the

location of highest calcium event rate (putative place field centers) by comparing their distances

on the last two criterion sessions in environment A (sessions A(n-1) and A(n)) to the shift

observed when transitioning from environment A on session n to environment B on session 1

(the centers of both environments were aligned, see methods). The data indicated a significantly

greater change in these putative place field centers when the animals transitioned from A to B

than when remaining within environment A (medians tested with Wilcoxon rank sum test

p=0.002, means tested with double sided t-test t(1430)= -2.5, p=0.01, distributions tested with

two-sample Kolmogorov-Smirnov (KS) test, p=3.6*10^{-4}) (Fig. 2c). Second, to compensate for

any differences in environment size, we also compared the distances between field centers to the

distances expected if all centers were shuffled 100 times, and found that the median distance

between field centers when comparing session A(n) to session A(n-1) was less than all median

shuffled values (p=0), while the median distance between field centers when comparing session

A(n) to session B(1) was greater or equal than 56% of shuffled values (p=0.56) (Fig. 2d). Third,

accounting for the fact that cells may have multiple place fields, we computed the population

vector correlation (PVC[40,48,49], see methods) using calcium events for sessions A(n-1), A(n), and

B(1). When using cells that appeared in both sessions A(n-1) and A(n), we found a significant

positive correlation when computing the PVC for these two sessions (p = 0.0023, r = 0.11).

Conversely, when using cells that appeared in both sessions A(n) and B(1), we found no

correlation (p>0.05, r=-0.04). This result indicates significantly similar calcium event patterns

between sessions A(n-1) and session A(n), with no significant similarity in patterns between

session A(n) and B(1). (Fig. 2e).

We then used a machine learning algorithm to determine the variation in neural

embeddings between environments A and B. To do this, we applied the CEBRA algorithm[50] to

calcium trace imaging data labelled with spatial coordinates from environment A, session A(n)

(all were trained on 75% of data with 25% held out for verification). The choice of CEBRA was

motivated by its efficacy and interpretability in decoding neural activity patterns when compared

to alternative methods such as PCA[51] and Isomap[52] (see Methods for additional details). We then

tested this model's ability to decode the animals' position in environment A and environment B

when applied to neural data not used for training. The results showed that the model, when run

500 times, predicted the positions in session A(n-1) with significantly greater accuracy than what

would be expected by chance; the null was constructed as determined when compared to a model

trained on shuffled position data and also run 500x. (All double sided t-tests, for each rat: rat1:

t(998):-34.3 p=$5.0*10^{-171}$, rat2: t(998):-72.5 p=0, rat3: t(998)=-1.96 p=0.05, rat4: t(998):-53.7

p=$2.6*10^{-299}$, rat5: t(998):-20.0 p=$4.1*10^{-75}$) (Fig. 3a-b). In contrast, when a model trained on

data from A(n) was applied to environment B, the model's predictions were significantly below

the accuracy of a model trained on shuffled position data, implying that the place cell coding

across environments A and B are actually more different than would be expected by chance (all

double sided t-tests, for each rat: rat1: t(998):84.1 p=0, rat2: t(998):18.8 p=$7.4*10^{-68}$, rat3:

t(998)=74.7 p=0, rat4: t(998):13.1 p=$2.0*10^{-36}$, rat5: t(998):154.4 p=0) (Fig. 3c-d). Collectively,

these findings suggest a significant remapping of place cells when transitioning between

environments, and also that the neural embeddings for place coding in individual rats change

when the animal switches contexts.

The hippocampus represents the conditioning task in both environments, and representations

of the conditioning task are not spatial representations

We then investigated whether conditioning related data was represented equally in both

environments A and B. It was obvious on visual inspection that individual cells varied their

calcium event rate (Fig. 4a) and calcium trace (Fig. 4b) during the conditioning periods. To

quantify this variation, we devised the metric 'CSUS mutual information' (CSUS-MI), analogous

to spatial mutual information; this enabled us to assess the extent of task-related information

captured by the calcium activity of each cell. We calculated the CSUS-MI for each cell and

benchmarked it against a control distribution generated by shuffling CS and US periods and

recalculating the MI 500 times. Using calcium event data, we found that 10.7±4.9% of cells held

significant CSUS information related to whether the animal was in a CS or US period (termed

CSUS-MI2 as the conditioning period was divided into 2 bins; see Methods) (Fig. S4a). An even

stronger relationship was noted if using calcium traces: 19.9±8.2 percent of cells contained

significant information related to whether the animal was in a CS or US period (Fig. S4a).

Importantly, neither of these MI metrics were significantly different between environments A

and B (double sided t-tests, using calcium events, $t(23)=0.48$, $p>0.05$, using calcium traces,

$t(23)=-0.52$, $p>0.05$).

We then extended this analysis to determine if calcium events or traces of individual cells

contained information about what temporal segment portion of the conditioning task the animal

was in. To do this, we divided the CSUS period into 5 equal sized bins, computed CSUS-MI

using these bins (termed CSUS-MI5, see methods), then compared these mutual information

values to the controls provided by shuffled data. Using calcium event data, we found that

15.5±7.8% of cells contained this information, compared to 10.0±7.8% of cells when we

calculated the MI using trace information (Fig. S4b). Again, neither of these mutual information

metrics were significantly different between environments A and B (double sided t-tests, using

calcium event data, $t(23)=-0.32$, using calcium trace data, $t(23)=-1.1$, $p>0.05$). There was not a

significant difference in CSUS-MI2 values when comparing values in session A(n) to session

A(n-1), versus comparing values in session A(n) to session B(1) (Wilcoxon rank sum test

$p>0.05$, double sided t-test, $t(1431)=0.86$, $p>0.05$). In contrast, there was a small but significant

difference in CSUS-MI5 when comparing session A(n) to session A(n-1) versus comparing A(n)

to session B(1) (Wilcoxon rank sum test $p=0.049$, double sided t-test, $t(1431)=-2.2$, $p=0.03$) (Fig.

S4c).Collectively, these results demonstrate that the conditioning task is represented in both

environments A and B, and that the percentage of cells representing the conditioning task was

not different between the two environments.

We then examined the overlap between cells that contained spatial information and those

with CSUS information, identifying a significant positive correlation between spatial MI and

CSUS-MI2 (linear regression, $r^2 = 0.04$, p=2.1*10^{-104}) and spatial MI and CSUS-MI5 (linear

regression, $r^2 = 0.09$, p=1.7*10^{-237}; note that the statistics have been computed for unbinned data

but the graph presents binned data for visualization purposes due to the large number of points)

(Fig. 4c). Further analysis revealed that cells with significant spatial modulation had a

significantly higher likelihood of being significantly modulated by CSUS compared to cells

without spatial modulation: 1.35 times higher chance of having significant spatial MI if the cell

has a significant CSUS-MI2 (Fisher's exact test, p=0.001) and 1.27 times higher chance if the

cell has a significant CSUS-MI5 (Fisher's exact test, p=0.002). We then inquired whether the

calcium events that occurred during conditioning periods were confined to the 'firing' fields of

place cells. We thus calculated the average location of calcium events during conditioning

periods versus the average location during periods of movement that were not conditioning

periods. We analyzed this data using the Mantel test, which statistically evaluates the correlation

between two distance matrices to determine if the spatial patterns they represent are significantly

related. Across all sessions, we found a Mantel statistic of 534.58; we compared this statistic to

the result of 10,000 shuffles to determine the statistic was not significant (p>0.05); i.e. the spatial

firing patterns during conditioning periods are not generally similar to those that occur during

non-conditioning periods (Fig. 4d). In other words, the spatial distribution of firing during

conditioning differs fundamentally from that during non-conditioning periods, a result which

would not be expected if conditioning related responses were restricted to the cells' place fields

We next assessed the consistency of task representation across environments. When we

compared cells that appeared in both sessions A(n) and B(1), there was no difference in cell

responses to either CS or US (double sided t-tests for CS: $t(1174) = 0.68$ p>0.05 and ks test

p>0.05, for US: $t(1174) = 1.20$ p>0.05, and ks-test p>0.05) (Fig. 4e).

Conditioning task representations are consistent across environments

We then trained a CEBRA model using calcium imaging data and time-stamped CS/US

periods from environment A, using only cells that were recorded in both environment A and B.

We then used this trained model to decode if the animal was in a CS or US period during an

additional session in environment A, as well as in environment B. All models successfully

decoded CS and US periods the additional session in environment A, as compared to shuffled

data (all double sided t-tests, for each rat: rat1: $t(998):6.7$ $p=2.5*10^{-11}$, rat2: $t(998):16.1$

$p=7.4*10^{-52}$, rat3: $t(998)=83.5$ p=0, rat4: $t(998):80.1$ p=0, rat5: $t(998):61.0$ p=0) (Fig. 5a,c). All

five models significantly outperformed chance level in environment B as determined by shuffled

data (all double sided t-tests, for each rat: rat1: $t(998):10.4$ $p=2.6*10^{-24}$, rat2: $t(998):2.7$

$p=7.6*10^{-3}$, rat3: $t(998)=75.3$ p=0, rat4: $t(998):63.7$ p=0, rat5: $t(998):106.3$ p=0) (Fig. 5b-c).

We then trained an additional model on data from environment A during session A(n)to

ascertain whether it could decode the temporal order within the conditioning period (CSUS5),

both in an alternate session in environment A (session A(n-1)) and in environment B (session

B(1)). After being trained on session A(n), all models were able to decode environment A(n-1)

well better than chance levels (all double sided t-tests for accuracy, for each rat: rat1: t(998):41.5

p=3.3*10^{-220}, rat2: t(998):28.7 p=7.8*10^{-133}, rat3: t(998)=122.6 p=0, rat4: t(998):118.5 p=0, rat5:

t(998):62.6 p=0) (Fig. 5d,f,g). Remarkably, our results for decoding environment B(1) showed

that temporal aspects of CS/US temporal order were decodable across environments, suggesting

that a refined level of task encoding is stable across both environments (all double sided t-tests

for accuracy, for each rat: rat1: t(998):55.1 p=4.9*10^{-305}, rat2: t(998):9.3 p=1.1*10^{-19}, rat3:

t(998)=71.4 p=0, rat4: t(998):62.6 p=0, rat5: t(998):106.2 p=0; results were also significant

compared to those for shuffled data for precision, recall, F1 score, and area under the receiver

operating characteristic curve, data not shown, see Methods) (Fig. 5e-g).

Remarkably, for both CSUS2 and CSUS5, the model trained on session A(n) was no less

accurate decoding session B(1) than it was decoding session A(n-1) (double sided t-tests, for

CSUS2 t(8)=-0.13, p>0.05, for CSUS5, t(8)=0.32, p>0.05) (Fig. 5h).

We then used CEBRA to analyze the embedding geometries of cell representations during

CS/US periods in both environments. First, we examined the embedding geometries for the

conditioning task divided into CS and US periods (CSUS2), for 2, 3, 5, 7, and 10 latents. We

compared sessions A(n-1), A(n), B(1), and B(2) to each other, as well as shuffled versions of

each session.

For all 5 rats, the geometries displayed a high and significant degree of similarity as

compared to the shuffled control. This significance was maintained when examining up to 10

latents, the largest number of latents we utilized (averaging results from all animals, all double

sided t-tests, for 2 latents all comparisons were $p<1*10^{-5}$, for 3 latents $p<1*10^{-4}$, for 5 latents

$p<1*10^{-5}$, for 7 latents $p<1*10^{-5}$, and for 10 latents $p<1*10^{-4}$) (Fig. 6a-c). This high degree of

similarity was maintained when the CS/US periods were divided into 5 segments (CSUS5, (all

double sided t-tests, for 2 latents all comparisons were $p<1*10^{-12}$, for 3 latents $p<1*10^{-7}$, for 5

latents $p<1*10^{-6}$, for 7 latents $p<1*10^{-5}$, and for 10 latents $p<1*10^{-5}$) (Fig. 6d-f).

These highly significant similarity signifies that the neural representations of the task

were consistent between environments A and B.

Conditioning task representations are consistent across animals

Considering the similarity between representations of the task in environments A and B, we

wondered if there was a universal, inter-animal, representation of the conditioning task. To

answer this question, we investigated if there were coding similarities of the conditioning task

across subjects. For each animal, we developed a unique model based on calcium signal patterns

and the structure of the conditioning task. We then calculated a similarity score among all

animal-specific models. We observed a markedly significant consistency across these trained

models compared to those trained on shuffled data. Notably, this consistency was apparent in

models trained to differentiate between CS and US periods, as well as in more granular models

that recognized five discrete time segments during CS presentation, the trace interval, and US

delivery (akin to the models in Figures 6d-g). When the conditioning period is divided into 2

periods (CSUS2), the similarity across animal models is not significantly different than the

similarity between models in one animal for all tested number of latents (2 latents: ttest(188) = -

0.45 p>0.05, 3 latents: ttest(188) = -1.57 p>0.05, 5 latents: ttest(188) = -0.40 p>0.05, 7 latents:

ttest(188) = -0.22, p>0.05, 10 latents: ttest(188) = 0.40, p>0.05) (Fig. 7a-b). This relationship

also holds when the conditioning period is divided into 5 periods (CSUS5): the similarity across

animal models is not significantly different than the similarity between models in one animal (2

latents: ttest(188) = -0.79 p>0.05, 3 latents: ttest(188) = 0.25 p>0.05, 5 latents: ttest(188) = -0.42

p>0.05, 7 latents: ttest(188) = 0.70, p>0.05, 10 latents: ttest(188) = 0.30, p>0.05) (Fig. 7c-d).

Discussion

Our study provides significant insights into hippocampal function, as it demonstrates that

the hippocampus not only responds to environmental change with changes in neural coding but

also maintains consistent task-related information across varying contexts. These mechanisms

underpin cognitive flexibility and the ability to apply learned behaviors in new situations. Below

we will discuss this interplay between variability and consistency of representations through

several theoretical lenses, including predictive coding and cognitive mapping theories, while

exploring the stability of these processes within and across subjects.

Task abstraction across environments

A critical aspect of our study highlights that the hippocampus retains stable task

representations, such as those required for eyeblink conditioning, despite variations in

environmental contexts. This ability to generalize learned tasks across different settings supports

models that posit a cognitive map that extends beyond simple spatial navigation, such as the

Tolman-Eichenbaum Machine's (TEM)[53]. According to this model, "spatial" maps integrate

task-related information and enable the hippocampus to utilize learned behaviors in novel

environments that share cognitive demands but differ in sensory or environmental specifics. This

flexible functionality exemplifies the hippocampus' role in abstracting and applying learned

knowledge, a hallmark of high-dimensional cognitive mapping. This integrated framework

facilitates the adaptation of learned behaviors across diverse contexts, an essential capability for

navigating both physical and abstract environments.

Our findings further reveal that the hippocampus abstracts task-related information from

the surrounding sensory environment and suggests that the hippocampus connects experiences

across different contexts by recognizing underlying similarities; these features help generalize

learning and adjust behavior. Our results thus support the TEM's perspective that the

hippocampus can encode higher-order, abstract information crucial for task execution. The

ability to detach task representation from immediate sensory inputs allows for a generalized

version of learned information[54-56], enhancing the hippocampus's utility in supporting the

organism's application of learned skills and behaviors in new albeit similar situations[55,56]. This

capacity for abstraction is indicative of a sophisticated neural coding mechanism and of an

adaptable and extensive cognitive mapping system, as it provides a buffer against potential

interference that could arise from the myriad of sensory stimuli an organism encounters. By

maintaining a conceptual, generalized version of learned information, the hippocampus supports

the organism's ability to apply learned skills and behaviors in new situations that share

underlying similarities with previous experiences but differ in sensory or contextual details.

Pattern Separation vs. Completion

Evidence from various areas of neuroscience has led to the development of a theory of HPC

function holding that the HPC treats states that involve equivalent actions or relationships such

as similar tasks as equivalent, resulting in learning that is easily transferred between

environments[53-55,57,58]. This theory contrasts with the theory that the HPC acts primarily to

perform competitive "pattern separation"[59-63]. The prevailing theory as well as the cellular and

systems level bases for contextual memory remain to be elucidated. In our study, the distinct

coding of different environments by hippocampal place cells provides evidence for pattern

separation: the hippocampus differentiates between distinct contexts. This separation reduces

interference between memories, allowing for more accurate recall based on specific

environmental cues. On the other hand, consistent decoding of the eyeblink task across different

environments suggests pattern completion. This process allows the hippocampus to reconstruct a

complete memory or learned response from partial or generalized cues, enabling the execution of

the learned task even when contextual details change.

Our results complement previous work that identifies the HPC as both a pattern

completer and a pattern separator. Studies have demonstrated that place cells can differentially

represent the same environment when task demands change[61,64,65], yet show similar firing

patterns when locations have similar task demands[66]. In more recent work[67], rats were exposed to

two distinct environments while performing variations of the same task: approaching object A in

the first environment and object B in the second; the study revealed anticorrelated hippocampal

firing patterns for events in the two contexts. This suggests that the hippocampus encodes

context-specific associations between items and locations, rather than just specific behavior. This

study underscored the role of the hippocampus in robust pattern separation when environments

differ but require similar behaviors, showing that even minor task variations can lead to

significant neuronal pattern separation[67]. In contrast, our study used a task that remained

identical across both environments and found consistent hippocampal population-level task

representations in both contexts. This consistency likely reflects that the task could be

generalized between environments, without necessitating the hippocampus to differentiate

between task demands. Therefore, the hippocampus seems to balance pattern separation and

completion based on how similar or distinct task demands are across different contexts.

Non-spatial hippocampal representations

There is ongoing debate about whether hippocampal pyramidal cells encode both spatial

and non-spatial aspects of a context[39,68,69]. Our findings show that responses to the conditioning

task were independent of place field location, with a fraction of the recorded individual cells

found able to represent both spatial location and the conditioning task (Fig. 4). This contrasts

with studies where conditioning responses were more closely tied to specific spatial

locations[70,71]. The difference likely stems from task design: in previous studies, spatially

contingent rewards or freezing behavior after a shock made location highly salient, by

associating the place where the shock occurred with the aversive event. In our experiment,

spatial position was irrelevant; this allowed pyramidal cells to encode task-relevant features

independent of location. This finding aligns with previous work showing that hippocampal cells

often respond to non-spatial aspects like sensory cues or task demands, particularly in non-spatial

tasks[19,72-74].

The discovery that population level patterns in the hippocampus are organized into

manifolds provides an elegant solution to the problem of single cells representing both spatial

coordinates and task features. Previous hippocampal work has described distinct encoding for

spatial location along the center stem of a T maze vs accumulation of evidence for a left-right

turning decision at the end of this branch through two distinct, orthogonal directions in a two-

dimensional neural manifold[75]. Other brain regions, such as the prefrontal cortex and cingulate

cortex, appear to use a similar orthogonal coding strategy[76-78]; this strategy has also emerged in

neural network simulations of a context-dependent classification task[79].

Intra-Subject and Inter-Subject Consistency

The stability of neural representations within subjects across different testing sessions

indicates that once hippocampal circuits are trained, their functional architecture remains

remarkably consistent, even in varying contexts. This intra-subject consistency supports theories

suggesting that neural circuits are not just reactive but possess a robust, predefined role in

processing and responding to specific stimuli[80,81]. Furthermore, the observation of similar neural

encoding patterns across different animals performing the same task suggests a species specific,

possibly evolutionary conserved, neural code[80,81]. These findings highlight a generalized neural

processing strategy that may have been shaped by natural selection to optimize cognitive and

behavioral responses across environmental challenges faced by a species. Such a generalized

coding strategy may be indicative of evolutionary pressures that have favored neural mechanisms

promoting cognitive flexibility and rapid adaptation to environmental challenges[82-84].

The results of our study highlight the preservation of hippocampal task encoding across

different contexts and species; this presents a surprising parallel to recent findings in motor

cortex[85,86] and insular cortex[87]. Motor functions, especially those fundamental to survival and

interaction with the physical environment, are expected to exhibit conserved neural dynamics

due to their innate and reflexive nature; motor tasks typically involve stereotyped and predictable

patterns of behavior that are essential for immediate responses and interactions with the

environment[88-91]. Similarly, the motivational states preserved across animals in the insular cortex

are those central to basic biological needs, such as thirst and hunger. These states and tasks are

often highly conserved across individuals because they rely on well-established neural circuits

that perform specific, crucial functions necessary for survival[89,91-95].

In contrast, hippocampal tasks involve complex cognitive processes that include memory,

learning, and spatial navigation; these require a higher degree of cognitive flexibility[83,96-98]. and

are generally influenced by individual experiences (and, at the species level, specific ecological

and evolutionary pressures)[99-103]. Therefore, the conservation of hippocampal task encoding

across different individuals, as observed in our study, challenges these traditional views and

suggests a deeper, possibly adaptive significance to these cognitive functions.

The surprising conservation of the neural representation of tasks within the hippocampus

suggests that certain aspects of cognitive mapping and memory processing might be as

evolutionarily essential as motor functions. This conservation might reflect universal cognitive

strategies that are critical for survival across a range of environmental contexts, providing

individuals within a species with the ability to adapt behavior based on past experiences and

anticipated future conditions. Such a mechanism would not only enhance an organism's ability to

navigate complex environments but also facilitate learning and decision-making across

generational timescales.

Conclusion

The consistent decoding of eyeblink conditioning tasks across different environmental

contexts indicates that the hippocampus can maintain a stable representation of task-specific

information irrespective of the external sensory environment. This suggests an advanced

capability for abstract cognitive mapping, where the hippocampus constructs and utilizes

cognitive maps not only for physical locations but also for abstract tasks and concepts, allowing

for effective application in varying contexts. These findings expand our understanding of how

memories are formed, stored, and retrieved. They suggest that memories are not just static

recollections of past events but dynamic and adaptable representations that can be applied to new

situations.

Citations

474   **Citations**

475   1   Bouton, M. E. A learning theory perspective on lapse, relapse, and the maintenance of
476   behavior change. *Health psychology* **19**, 57 (2000).
477   2   Herszage, J. & Censor, N. Modulation of learning and memory: a shared framework for
478   interference and generalization. *Neuroscience* **392**, 270-280 (2018).
479   3   Linda, Q. Y., Wilson, R. C. & Nassar, M. R. Adaptive learning is structure learning in time.
480   *Neuroscience & Biobehavioral Reviews* **128**, 270-281 (2021).
481   4   Censor, N. Generalization of perceptual and motor learning: a causal link with memory
482   encoding and consolidation? *Neuroscience* **250**, 201-207 (2013).
483   5   Brown, S. & Bebko, J. Generalization, overselectivity, and discrimination in the autism
484   phenotype: A review. *Research in Autism Spectrum Disorders* **6**, 733-740 (2012).
485   6   Church, B. A. *et al.* Learning, plasticity, and atypical generalization in children with
486   autism. *Psychonomic Bulletin & Review* **22**, 1342-1348 (2015).
487   7   Weiler, J. A., Bellebaum, C., Brüne, M., Juckel, G. & Daum, I. Impairment of probabilistic
488   reward-based learning in schizophrenia. *Neuropsychology* **23**, 571 (2009).
489   8   Shohamy, D. *et al.* Learning and generalization in schizophrenia: effects of disease and
490   antipsychotic drug treatment. *Biological psychiatry* **67**, 926-932 (2010).
491   9   Anastasides, N. *et al.* Increased generalization of learned associations is related to re-
492   experiencing symptoms in veterans with symptoms of post-traumatic stress. *Stress* **18**,
493   484-489 (2015).
494   10   Lis, S. *et al.* Generalization of fear in post-traumatic stress disorder. *Psychophysiology*
495   **57**, e13422 (2020).
496   11   Calandreau, L., Desgranges, B., Jaffard, R. & Desmedt, A. Switching from contextual to
497   tone fear conditioning and vice versa: the key role of the glutamatergic hippocampal-
498   lateral septal neurotransmission. *Learn Mem* **17**, 440-443 (2010).
499   https://doi.org:10.1101/lm.1859810
500   12   Desmedt, A., Garcia, R. & Jaffard, R. Vasopressin in the lateral septum promotes
501   elemental conditioning to the detriment of contextual fear conditioning in mice. *Eur J*
502   *Neurosci* **11**, 3913-3921 (1999).
503   13   Vouimba, R. M., Garcia, R. & Jaffard, R. Opposite effects of lateral septal LTP and lateral
504   septal lesions on contextual fear conditioning in mice. *Behav Neurosci* **112**, 875-884
505   (1998).
506   14   Anagnostaras, S. G., Gale, G. D. & Fanselow, M. S. Hippocampus and contextual fear
507   conditioning: recent controversies and advances. *Hippocampus* **11**, 8-17 (2001).
508   https://doi.org:10.1002/1098-1063(2001)11:1<8::AID-HIPO1015>3.0.CO;2-7
509   15   Honey, R. C. & Good, M. Selective hippocampal lesions abolish the contextual specificity
510   of latent inhibition and conditioning. *Behav Neurosci* **107**, 23-33 (1993).
511   https://doi.org:10.1037//0735-7044.107.1.23
512   16   Plitt, M. H. & Giocomo, L. M. Experience-dependent contextual codes in the
513   hippocampus. *Nat Neurosci* **24**, 705-714 (2021). https://doi.org:10.1038/s41593-021-
514   00816-6

17 Devito, L. M., Kanter, B. R. & Eichenbaum, H. The hippocampus contributes to memory
expression during transitive inference in mice. *Hippocampus* **20**, 208-217 (2010).
https://doi.org:10.1002/hipo.20610
18 Schimanski, L. A., Lipa, P. & Barnes, C. A. Tracking the course of hippocampal
representations during learning: when is the map required? *J Neurosci* **33**, 3094-3106
(2013). https://doi.org:10.1523/JNEUROSCI.1348-12.2013
19 Gauthier, J. L. & Tank, D. W. A Dedicated Population for Reward Coding in the
Hippocampus. *Neuron* **99**, 179-193 e177 (2018).
https://doi.org:10.1016/j.neuron.2018.06.008
20 Mamad, O. *et al.* Place field assembly distribution encodes preferred locations. *PLoS Biol*
15, e2002365 (2017). https://doi.org:10.1371/journal.pbio.2002365
21 Hok, V. *et al.* Goal-related activity in hippocampal place cells. *J Neurosci* **27**, 472-482
(2007). https://doi.org:27/3/472 [pii] 10.1523/JNEUROSCI.2864-06.2007
22 Knierim, J. J. Dynamic interactions between local surface cues, distal landmarks, and
intrinsic circuitry in hippocampal place cells. *J Neurosci* **22**, 6254-6264 (2002).
https://doi.org:20026608
23 Allen, K., Rawlins, J. N., Bannerman, D. M. & Csicsvari, J. Hippocampal place cells can
encode multiple trial-dependent features through rate remapping. *J Neurosci* **32**, 14752-
14766 (2012). https://doi.org:10.1523/JNEUROSCI.6175-11.2012
24 Jeffery, K. J., Gilbert, A., Burton, S. & Strudwick, A. Preserved performance in a
hippocampal-dependent spatial task despite complete place cell remapping.
Hippocampus **13**, 175-189 (2003). https://doi.org:10.1002/hipo.10047
25 Colgin, L. L., Moser, E.I., May-Britt Moser. Understanding memory through hippocampal
remapping. *Cell* (2008).
26 Christian, K. M. & Thompson, R. F. Neural substrates of eyeblink conditioning:
acquisition and retention. *Learn Mem* **10**, 427-455 (2003).
https://doi.org:10.1101/lm.59603
27 McEchron, M. D., Tseng, W. & Disterhoft, J. F. Single neurons in CA1 hippocampus
encode trace interval duration during trace heart rate (fear) conditioning in rabbit. *J*
Neurosci **23**, 1535-1547 (2003).
28 Wirtshafter, H. S. & Wilson, M. A. Locomotor and Hippocampal Processing Converge in
the Lateral Septum. *Current biology : CB* **29**, 3177-3192 (2019).
https://doi.org:10.1016/j.cub.2019.07.089
29 Berger, T. W., Rinaldi, P. C., Weisz, D. J. & Thompson, R. F. Single-unit analysis of
different hippocampal cell types during classical conditioning of rabbit nictitating
membrane response. *J Neurophysiol* **50**, 1197-1219 (1983).
https://doi.org:10.1152/jn.1983.50.5.1197
30 Wirtshafter, H. S. & Wilson, M. A. Differences in reward biased spatial representations in
the lateral septum and hippocampus. *eLife* **9** (2020).
https://doi.org:10.7554/eLife.55252
31 Wirtshafter, H. S. & Disterhoft, J. F. Place cells are nonrandomly clustered by field
location in CA1 hippocampus. *Hippocampus* **33**, 65-84 (2023).
https://doi.org:10.1002/hipo.23489

32 Aharoni, D. & Hoogland, T. M. Circuit Investigations With Open-Source Miniaturized
Microscopes: Past, Present and Future. *Frontiers in cellular neuroscience* **13**, 141 (2019).
https://doi.org:10.3389/fncel.2019.00141
33 Aharoni, D., Khakh, B. S., Silva, A. J. & Golshani, P. All the light that we can see: a new
era in miniaturized microscopy. *Nat Methods* **16**, 11-13 (2019).
https://doi.org:10.1038/s41592-018-0266-x
34 Silva, A. J. Miniaturized two-photon microscope: seeing clearer and deeper into the
brain. *Light: Science & Applications* **6**, e17104-e17104 (2017).
35 Wirtshafter, H. S. & Disterhoft, J. F. In Vivo Multi-Day Calcium Imaging of CA1
Hippocampus in Freely Moving Rats Reveals a High Preponderance of Place Cells with
Consistent Place Fields. *J Neurosci* (2022). https://doi.org:doi: 10.1523/JNEUROSCI.1750-
21.2022
36 Cunningham, J. P. & Byron, M. Y. Dimensionality reduction for large-scale neural
recordings. *Nature neuroscience* **17**, 1500-1509 (2014).
37 Bassett, D. S. & Sporns, O. Network neuroscience. *Nature neuroscience* **20**, 353-364
(2017).
38 Ekstrom, A. D. & Ranganath, C. Space, time, and episodic memory: The hippocampus is
all over the cognitive map. *Hippocampus* **28**, 680-687 (2018).
39 Eichenbaum, H., Dudchenko, P., Wood, E., Shapiro, M. & Tanila, H. The hippocampus,
memory, and place cells: is it spatial memory or a memory space? *Neuron* **23**, 209-226
(1999). https://doi.org:S0896-6273(00)80773-4 [pii]
40 Fenton, A. A. Remapping revisited: how the hippocampus represents different spaces.
Nat Rev Neurosci **25**, 428-448 (2024). https://doi.org:10.1038/s41583-024-00817-x
41 Sanders, H., Wilson, M. A. & Gershman, S. J. Hippocampal remapping as hidden state
inference. *eLife* **9** (2020). https://doi.org:10.7554/eLife.51140
42 Bouton, M. E., Nelson, J. B. & Rosas, J. M. Stimulus generalization, context change, and
forgetting. *Psychological bulletin* **125**, 171-186 (1999). https://doi.org:10.1037/0033-
2909.125.2.171
43 Lovibond, P. F., Preston, G. & Mackintosh, N. Context specificity of conditioning,
extinction, and latent inhibition. *Journal of Experimental Psychology: Animal Behavior
Processes* **10**, 360 (1984).
44 Moyer, J. R., Jr., Deyo, R. A. & Disterhoft, J. F. Hippocampectomy disrupts trace eye-blink
conditioning in rabbits. *Behav Neurosci* **104**, 243-252 (1990).
45 McEchron, M. D. & Disterhoft, J. F. Hippocampal encoding of non-spatial trace
conditioning. *Hippocampus* **9**, 385-396 (1999). https://doi.org:10.1002/(SICI)1098-
1063(1999)9:4<385::AID-HIPO5>3.0.CO;2-K
46 Woodruff-Pak, D. S. & Disterhoft, J. F. Where is the trace in trace conditioning? *Trends
Neurosci* **31**, 105-112 (2008). https://doi.org:10.1016/j.tins.2007.11.006
47 Weiss, C. & Disterhoft, J. F. The impact of hippocampal lesions on trace-eyeblink
conditioning and forebrain-cerebellar interactions. *Behav Neurosci* **129**, 512-522 (2015).
https://doi.org:10.1037/bne0000061
48 Neymotin, S. A., Talbot, Z. N., Jung, J. Q., Fenton, A. A. & Lytton, W. W. Tracking
recurrence of correlation structure in neuronal recordings. *J Neurosci Methods* **275**, 1-9
(2017). https://doi.org:10.1016/j.jneumeth.2016.10.009

49   Brun, V. H. *et al.* Progressive increase in grid scale from dorsal to ventral medial
entorhinal cortex. *Hippocampus* **18**, 1200-1212 (2008).
https://doi.org:10.1002/hipo.20504
50   Schneider, S., Lee, J. H. & Mathis, M. W. Learnable latent embeddings for joint
behavioural and neural analysis. *Nature* **617**, 360-368 (2023).
https://doi.org:10.1038/s41586-023-06031-6
51   Pearson, K. LIII. On lines and planes of closest fit to systems of points in space. *The*
London, Edinburgh, and Dublin philosophical magazine and journal of science **2**, 559-572
(1901).
52   Tenenbaum, J. B., Silva, V. d. & Langford, J. C. A global geometric framework for
nonlinear dimensionality reduction. *science* **290**, 2319-2323 (2000).
53   Whittington, J. C. R. *et al.* The Tolman-Eichenbaum Machine: Unifying Space and
Relational Memory through Generalization in the Hippocampal Formation. *Cell* **183**,
1249-1263 e1223 (2020). https://doi.org:10.1016/j.cell.2020.10.024
54   Wirtshafter, H. S. & Wilson, M. A. Artificial intelligence insights into hippocampal
processing. *Front Comput Neurosci* **16**, 1044659 (2022).
https://doi.org:10.3389/fncom.2022.1044659
55   Lehnert, L., Littman, M. L. & Frank, M. J. Reward-predictive representations generalize
across tasks in reinforcement learning. *PLoS Comput Biol* **16**, e1008317 (2020).
https://doi.org:10.1371/journal.pcbi.1008317
56   Collins, A. G. E. & Frank, M. J. Neural signature of hierarchically structured expectations
predicts clustering and transfer of rule sets in reinforcement learning. *Cognition* **152**,
160-169 (2016).
57   Stachenfeld, K. L., Botvinick, M. M. & Gershman, S. J. The hippocampus as a predictive
map. *Nat Neurosci* **20**, 1643-1653 (2017). https://doi.org:10.1038/nn.4650
58   Eichenbaum, H. Hippocampus: cognitive processes and neural representations that
underlie declarative memory. *Neuron* **44**, 109-120 (2004).
59   Wills, T. J., Lever, C., Cacucci, F., Burgess, N. & O'Keefe, J. Attractor dynamics in the
hippocampal representation of the local environment. *Science* **308**, 873-876 (2005).
https://doi.org:10.1126/science.1108905
60   Colgin, L. L. *et al.* Attractor-map versus autoassociation based attractor dynamics in the
hippocampal network. *J Neurophysiol* **104**, 35-50 (2010).
https://doi.org:10.1152/jn.00202.2010
61   Smith, D. M. & Mizumori, S. J. Learning-related development of context-specific
neuronal responses to places and events: the hippocampal role in context processing. *J*
Neurosci **26**, 3154-3163 (2006). https://doi.org:10.1523/JNEUROSCI.3234-05.2006
62   Colgin, L. L. *et al.* Attractor-map versus autoassociation based attractor dynamics in the
hippocampal network. *Journal of neurophysiology* **104**, 35-50 (2010).
63   Lee, I., Yoganarasimha, D., Rao, G. & Knierim, J. J. Comparison of population coherence
of place cells in hippocampal subfields CA1 and CA3. *Nature* **430**, 456-459 (2004).
64   Bahar, A. S., Shirvalkar, P. R. & Shapiro, M. L. Memory-guided learning: CA1 and CA3
neuronal ensembles differentially encode the commonalities and differences between
situations. *J Neurosci* **31**, 12270-12281 (2011).
https://doi.org:10.1523/JNEUROSCI.1671-11.2011

65 Markus, E. J. *et al.* Interactions between location and task affect the spatial and
directional firing of hippocampal neurons. *J Neurosci* **15**, 7079-7094 (1995).
66 Singer, A. C., Karlsson, M. P., Nathe, A. R., Carr, M. F. & Frank, L. M. Experience-
dependent development of coordinated hippocampal spatial activity representing the
similarity of related locations. *J Neurosci* **30**, 11586-11604 (2010).
https://doi.org:10.1523/JNEUROSCI.0926-10.2010
67 McKenzie, S. *et al.* Hippocampal representation of related and opposing memories
develop within distinct, hierarchically organized neural schemas. *Neuron* **83**, 202-215
(2014). https://doi.org:10.1016/j.neuron.2014.05.019
68 O'Keefe, J. & Krupic, J. Do hippocampal pyramidal cells respond to nonspatial stimuli?
Physiol Rev **101**, 1427-1456 (2021). https://doi.org:10.1152/physrev.00014.2020
69 Knierim, J. J. From the GPS to HM: Place cells, grid cells, and memory. *Hippocampus* **25**,
719-725 (2015). https://doi.org:10.1002/hipo.22453
70 Moita, M. A., Rosis, S., Zhou, Y., LeDoux, J. E. & Blair, H. T. Hippocampal place cells
acquire location-specific responses to the conditioned stimulus during auditory fear
conditioning. *Neuron* **37**, 485-497 (2003). https://doi.org:10.1016/s0896-
6273(03)00033-3
71 Shan, K. Q., Lubenov, E. V., Papadopoulou, M. & Siapas, A. G. Spatial tuning and brain
state account for dorsal hippocampal CA1 activity in a non-spatial learning task. *eLife* **5**
(2016). https://doi.org:10.7554/eLife.14321
72 Wood, E. R., Dudchenko, P. A. & Eichenbaum, H. The global record of memory in
hippocampal neuronal activity. *Nature* **397**, 613-616 (1999).
https://doi.org:10.1038/17605
73 Kennedy, P. J. & Shapiro, M. L. Motivational states activate distinct hippocampal
representations to guide goal-directed behaviors. *Proc Natl Acad Sci U S A* **106**, 10805-
10810 (2009). https://doi.org:10.1073/pnas.0903259106
74 Holscher, C., Jacob, W. & Mallot, H. A. Reward modulates neuronal activity in the
hippocampus of the rat. *Behav Brain Res* **142**, 181-191 (2003).
75 Nieh, E. H. *et al.* Geometry of abstract learned knowledge in the hippocampus. *Nature*
595, 80-84 (2021).
76 Johnston, W. J., Fine, J. M., Yoo, S. B. M., Ebitz, R. B. & Hayden, B. Y. Semi-orthogonal
subspaces for value mediate a binding and generalization trade-off. *Nature*
Neuroscience, 1-13 (2024).
77 Machens, C. K., Romo, R. & Brody, C. D. Functional, but not anatomical, separation of
"what" and "when" in prefrontal cortex. *Journal of Neuroscience* **30**, 350-360 (2010).
78 Mante, V., Sussillo, D., Shenoy, K. V. & Newsome, W. T. Context-dependent computation
by recurrent dynamics in prefrontal cortex. *nature* **503**, 78-84 (2013).
79 Flesch, T., Juechems, K., Dumbalska, T., Saxe, A. & Summerfield, C. Orthogonal
representations for robust context-dependent task performance in brains and neural
networks. *Neuron* **110**, 1258-1270. e1211 (2022).
80 Mizumori, S. J., Canfield, J. G. & Yeshenko, O. Parallel and interrelated neural systems
underlying adaptive navigation. *Integrative and Comparative Biology* **45**, 547-554
(2005).

Mizumori, S. J., Cooper, B. G., Leutgeb, S. & Pratt, W. E. A neural systems analysis of adaptive navigation. *Mol Neurobiol* **21**, 57-82 (2000). https://doi.org:10.1385/MN:21:1-2:057

Mizumori, S. & Smith, D. Directing Neural Representations of Space. *Animal Spatial Cognition: Comparative, Neural, and Computational Approaches* (2006).

Buzsáki, G. & Moser, E. I. Memory, navigation and theta rhythm in the hippocampal-entorhinal system. *Nature neuroscience* **16**, 130-138 (2013).

Bunsey, M. & Eichenbaum, H. Conservation of hippocampal memory function in rats and humans. *Nature* **379**, 255-257 (1996). https://doi.org:10.1038/379255a0

Safaie, M. *et al.* Preserved neural dynamics across animals performing similar behaviour. *Nature* **623**, 765-771 (2023). https://doi.org:10.1038/s41586-023-06714-0

Gallego, J. A. *et al.* Cortical population activity within a preserved neural manifold underlies multiple motor behaviors. *Nature communications* **9**, 1-13 (2018).

Talpir, I. & Livneh, Y. Stereotyped goal-directed manifold dynamics in the insular cortex. *Cell reports* **43** (2024).

Kogan, E., Lu, J. & Zuo, Y. Cortical circuit dynamics underlying motor skill learning: from rodents to humans. *Frontiers in Molecular Neuroscience* **16**, 1292685 (2023).

Manoli, D. S., Meissner, G. W. & Baker, B. S. Blueprints for behavior: genetic specification of neural circuitry for innate behaviors. *Trends in neurosciences* **29**, 444-451 (2006).

LeDoux, J. & Daw, N. D. Surviving threats: neural circuit and computational implications of a new taxonomy of defensive behaviour. *Nature Reviews Neuroscience* **19**, 269-282 (2018).

LeDoux, J. E. As soon as there was life, there was danger: the deep history of survival behaviours and the shallower history of consciousness. *Philosophical Transactions of the Royal Society B* **377**, 20210292 (2022).

Harris, K. D. & Shepherd, G. M. The neocortical circuit: themes and variations. *Nature neuroscience* **18**, 170-181 (2015).

Shmuelof, L. & Krakauer, J. W. Are we ready for a natural history of motor learning? *Neuron* **72**, 469-476 (2011).

Tierney, A. Evolutionary implications of neural circuit structure and function. *Behavioural processes* **35**, 173-182 (1995).

Martínez-García, F. & Lanuza, E. Evolution of vertebrate survival circuits. *Current Opinion in Behavioral Sciences* **24**, 113-123 (2018).

Sweis, B. M., Mau, W., Rabinowitz, S. & Cai, D. J. Dynamic and heterogeneous neural ensembles contribute to a memory engram. *Curr Opin Neurobiol* **67**, 199-206 (2021). https://doi.org:10.1016/j.conb.2020.11.017

Davidson, T. J., Kloosterman, F. & Wilson, M. A. Hippocampal replay of extended experience. *Neuron* **63**, 497-507 (2009). https://doi.org:10.1016/j.neuron.2009.07.027 S0896-6273(09)00582-0 [pii]

Rubin, R. D., Watson, P. D., Duff, M. C. & Cohen, N. J. The role of the hippocampus in flexible cognition and social behavior. *Frontiers in human neuroscience* **8**, 742 (2014).

Poulter, S., Hartley, T. & Lever, C. The neurobiology of mammalian navigation. *Current Biology* **28**, R1023-R1042 (2018).

100   Tucker, M. A., Ord, T. J. & Rogers, T. L. Evolutionary predictors of mammalian home
range size: body mass, diet and the environment. *Global Ecology and Biogeography* **23**,
1105-1114 (2014).
101   Schilder, B. M., Petry, H. M. & Hof, P. R. Evolutionary shifts dramatically reorganized the
human hippocampal complex. *Journal of Comparative Neurology* **528**, 3143-3170
(2020).
102   Sherry, D. F., Jacobs, L. F. & Gaulin, S. J. Spatial memory and adaptive specialization of
the hippocampus. *Trends in neurosciences* **15**, 298-303 (1992).
103   Pravosudov, V. V. & Roth II, T. C. Cognitive ecology of food hoarding: the evolution of
spatial memory and the hippocampus. *Annual Review of Ecology, Evolution, and*
Systematics **44**, 173-193 (2013).
744

Fig 1

Fig 1. Experimental training paradigm and results.

a. Animals explored environment A for one session before undergoing eyeblink conditioning until they reached the learning criterion. Criterion was defined as achieving 70% conditioned responses (CRs) in 50 trials across three consecutive training sessions (A(n-2), A(n-1), and A(n)) or averaging over 70% CRs across the previous four training sessions. After meeting the criterion in environment A, animals were allowed one session of exploration in environment B, followed by two test sessions of trace eyeblink conditioning (B(1) and B(2)) in environment B.

b. (Top) Schematics of environments A and B. Environment A is a rectangular enclosure with wire walls, floor, and ceiling, lit with white light, and unscented. Environment B is oval-shaped with solid white floors and walls, without a ceiling, lit with red light, and scented with clove oil. Both environments provided distal cues visible from the top and

sides. (Bottom) Animal trajectories in environments A and B during a single session,

extracted using DeepLabCut.

c. Trace eyeblink conditioning (tEBC) paradigm. A 250 ms tone (conditioned stimulus, CS)

was followed by a 500 ms trace interval, then a 100 ms eyelid shock (unconditioned

stimulus, US). Eyelid activity was recorded using an EMG electrode implanted above the

eye. Untrained animals only blinked in response to the US (unconditioned response, UR);

trained animals began blinking during the trace interval after the CS and before the US

(conditioned response, CR).

d. Performance of animals (n=5) in the tEBC task. Animals learned tEBC while freely

moving in environment A and successfully transferred this learning to environment B.

The dotted line indicates the performance criterion. No significant difference was found

between performance in the criterion sessions in environment A (mean $74.75 \pm 6.49\%$)

and the test sessions in environment B (mean $77.70 \pm 11.68\%$; two-tailed t-test, $t(24) = -$

0.83, $p > 0.05$). Error bars represent standard error.

Fig 2. Place cells remap between environment A and environment B.

a. Percent of place cells during criterion sessions in environment A and test sessions in

environment B. Light-colored bars represent averages within individual sessions, and

dark-colored bars represent overall averages in environments A and B. Overlaid bars

indicate standard deviation. On average, 9.3% ± 4.2% of cells were classified as place

cells. There was no significant difference in the percentage of place cells between

environments A and B (average in environment A: 8.1% ± 3.3%; average in environment

B: 11.1% ± 4.9%; two-tailed t-test, t(23) = -1.9, p > 0.05).

b. Example activity map of a specific single cell in environments A and B. Yellow indicates

the highest firing rates. This cell exhibited remapping between environments, showing

different place fields relative to external cues in the two contexts.

c. Distribution of distances between place field centers (determined by the highest calcium

event rate), comparing sessions A(n) and A(n-1) versus sessions A(n) and B(1). Place

field centers shifted significantly more when the animal was moved to environment B

compared to shifts between two consecutive sessions in environment A (Wilcoxon rank

sum test, $p = 0.002$; two-sided t-test, $t(1430) = -2.5$, $p = 0.01$; two-sample Kolmogorov-

Smirnov test, $p = 3.6 \times 10$). The brown shaded area represents the overlap between

distance histograms.

d. Distribution of the median distance between place field centers after shuffling center

locations 500 times. The actual median value for session A(n) to A(n-1) was smaller than

all shuffled medians ($p = 0$, dashed blue line), while the median for session A(n) to B(1)

was greater than or equal to 56% of shuffled medians ($p = 0.56$, dashed yellow line). The

brown shaded area represents the overlap between shuffled distributions.

e. Population vector correlation (PVC) based on calcium events for sessions A(n-1), A(n),

and B(1). A significant positive correlation was found between sessions A(n-1) and A(n)

for cells present in both sessions ($p = 0.0023$, $r = 0.11$). In contrast, there was no

significant correlation between sessions A(n) and B(1) for shared cells ($p > 0.05$, $r = $ -

0.04). These findings suggest that calcium event patterns are significantly similar

between sessions A(n-1) and A(n) but not between session A(n) and B(1). Dashed lines

represent lines of best fit. (a.u. = arbitrary units).

Fig. 3. A model trained in environment A can decode positions within environment A but not in environment B.

    a. A model trained on calcium trace and position data from session A(n), using cells present in both A(n) and A(n-1), predicted positions in environment A(n-1) with significantly greater accuracy than that of a model trained on shuffled position data. The model was run 500 times, and the accuracy of predictions was assessed. The model trained on actual data significantly outperformed the shuffled model across rats (double-sided t-tests: Rat 1: t(998) = -34.3, p = 5.0 × 10^{-171}; Rat 2: t(998) = -72.5, p = 0; Rat 3: t(998) = -1.96, p =

814	0.05; Rat 4: $t(998) = -53.7$, $p = 2.6 \times 10^{-299}$; Rat 5: $t(998) = -20.0$, $p = 4.1 \times 10^{-75}$). Error
815	bars represent standard error. A single asterisk (*) indicates $p \leq 0.05$, and three asterisks
816	(***) indicate $p < 10^{-74}$.
817	**b.** (Top row) Visualization of model performance for decoding animal position in
818	environment A(n). Left: The trained model demonstrated on the training data from
819	session A(n). Middle: The same model applied to held-out trace data (25%) from session
820	A(n). Right: The model applied to predict animal position in session A(n-1). (Bottom
821	row) The same models trained on shuffled position data. Shown here is the model for Rat
822	4, where the model trained on real data significantly outperformed model trained on
823	shuffled data for decoding position in session A(n-1) (500 simulations, $t(998) = -53.7$, $p =$
824	2.6×10^{-299}). For visualization purposes, distance from the corner of the environment is
825	plotted using normalized values in arbitrary units [a.u.].
826	**c.** A model trained on data from session A(n), using cells present in both A(n) and B(1),
827	was applied to environment B(1). The model's predictions were significantly less
828	accurate than those of a model trained on shuffled position data (double-sided t-tests: Rat
829	1: $t(998) = 84.1$, $p = 0$; Rat 2: $t(998) = 18.8$, $p = 7.4 \times 10^{-68}$; Rat 3: $t(998) = 74.7$, $p = 0$;
830	Rat 4: $t(998) = 13.1$, $p = 2.0 \times 10^{-36}$; Rat 5: $t(998) = 154.4$, $p = 0$). Error bars represent
831	standard error. Three asterisks (***) indicate $p < 10^{-35}$.
832	**d.** (Top row) Model trained on position and calcium trace data from session A(n), using
833	cells present in both A(n) and B(1). Left: The model demonstrated on the training data
834	from session A(n). Middle: The same model applied to held-out trace data (25%) from
835	session A(n). Right: The model applied to predict the animal position in session B(1).
836	(Bottom row) The same models trained on shuffled position data. Shown here is the
837	model for Rat 4, where the shuffled model performed significantly better than the model

838   trained on session A(n) for decoding position in session B(1) (500 simulations, $t(998) =$

839   13.1, $p = 2.0 \times 10^{-36}$). For visualization purposes, distance from the corner of the

840   environment is plotted using normalized values in arbitrary units [a.u.].

Fig 4

Figure 4. Identifying CS and US modulation in individual cells

a. Peristimulus time histograms (PSTHs) of six example cells responding to the CS, US,

and/or trace period. Bars represent calcium events summed across 50 trials. The

shaded regions indicate different stimulus periods: blue (CS), green (trace), and red

(US).

b. Mean calcium traces of eight example cells in response to the CS, US, and/or trace

period, averaged across 50 trials. Shaded regions are consistent with panel a (a.u. =

arbitrary units).

c. Scatter plots illustrating the significant positive correlation between spatial mutual

information (MI) and CSUS mutual information (MI). Data points represent binned

observations with error bars showing standard errors. The solid red line represents the

best fit from a linear regression analysis of the original unbinned dataset. Left: CSUS-

MI2 ($r^2 = 0.04$, $p = 2.1 \times 10^{-104}$). Right: CSUS-MI5 ($r^2 = 0.09$, $p = 1.7 \times 10^{-237}$).

d. Mosaic plots showing the odds ratios for cells modulated by spatial and/or CSUS

information. Cells were classified as significantly modulated if their mutual

information exceeded 95% of shuffled values. Left: Cells with significant spatial

modulation had a 1.35 times higher chance of being significantly modulated by

CSUS-MI2 compared to cells without spatial modulation (Fisher's exact test, $p =$

0.001). Right: Similar results for CSUS-MI5, where cells with significant spatial

modulation had a 1.27 times higher chance of being significantly modulated by

CSUS-MI5 (Fisher's exact test, $p = 0.002$).

e. Place field remapping during conditioning and non-conditioning periods. (Left) Heat

map comparing the spatial distances between place field centers during conditioning

(e.g., during CS/US trials) versus non-conditioning periods (e.g., intertrial intervals).

868   Color represents the absolute difference in distances between field centers: regions

869   with minimal color variation (blue) suggest similar place field centers between the

870   two conditions, while more yellow areas represent significant differences in distance

871   (Mantel statistic = 534.58, $p > 0.05$). (Right) Example of two cells showing spatial

872   calcium activity. Top: Heat maps showing calcium event rates during periods of

873   movement but not conditioning (e.g., intertrial intervals). Bottom: Calcium event

874   rates during CS/US conditioning periods. Dark blue signifies the lowest event rate,

875   and yellow represents the highest event rate. These results show that calcium events

876   during trials are not confined to a cell's place field.

877  **f.** Distribution of calcium events during CS (left) and US (right) periods in sessions

878   A(n) (blue) and B(1) (yellow). No significant difference was observed in

879   hippocampal firing during the CS (left) or US (right) periods between environments

880   A and B (CS: two-tailed t-test, $t(1174) = 0.68$, $p > 0.05$; US: two-tailed t-test, $t(1174)$

881   $= 1.20$, $p > 0.05$; KS tests, $p > 0.05$ for both comparisons).

Fig. 5. A model trained in environment A can decode CS and US periods in both

environment A and B at above chance levels, including fine-grained temporal decoding.

    a. A CEBRA model was trained using calcium imaging data and time-stamped CS/US

    periods from environment A, using only cells that were recorded in both environments A

    and B. The model was used to decode whether the animal was in a CS or US period in

    another session in environment A; all models successfully decoded these periods

    compared to shuffled data (all double-sided t-tests: Rat 1: $t(998) = 6.7$, $p = 2.5 \times 10^{-11}$;

    Rat 2: $t(998) = 16.1$, $p = 7.4 \times 10^{-52}$; Rat 3: $t(998) = 83.5$, $p = 0$; Rat 4: $t(998) = 80.1$, $p =$

0; Rat 5: t(998) = 61.0, p = 0). Bars indicate standard error; significance denoted as ***p

$< 10^{-10}$.

b. The same models from panel a were now applied to environment B; they significantly

outperformed chance level (all double-sided t-tests: Rat 1: t(998) = 10.4, p = 2.6×10^{-24};

Rat 2: t(998) = 2.7, p = 7.6×10^{-3}; Rat 3: t(998) = 75.3, p = 0; Rat 4: t(998) = 63.7, p = 0;

Rat 5: t(998) = 106.3, p = 0). Error bars represent standard error, *p < 10^{-3}, ***p < 10^{-10}.

c. (Top row) A CEBRA model trained on CS/US periods, divided into two time bins (data

from Rat 4). For the top first three graphs, the model was trained using position and

calcium trace data from session A(n), using cells present in both A(n) and A(n-1). First

graph: The trained model applied to training data from session A(n). Second graph: The

model applied to decode CS/US periods from held-out data (25%) from session A(n).

Third graph: The model applied to session A(n-1). For the top last three graphs, the

model was trained using position and calcium trace data from session A(n), using cells

present in both A(n) and B(1). Fourth graph: The trained model applied to training data

from session A(n). Fifth graph: The model applied to decode CS/US periods from held-

out data (25%) from session A(n). Sixth graph: The model applied to session B(1).

(Bottom row) The same model trained on shuffled data. The model trained on data from

session A(n) significantly outperformed the model trained on shuffled data in decoding

both session A(n-1) and session B(1) (double-sided t-tests, t(998) = 83.5, p = 0, and

t(998) = 75.3, p = 0, respectively).

d. A CEBRA model was trained on data from environment A to decode temporal order

within the CS, trace, and US periods split into five divisions, applied to an alternate

session in environment A (session A(n-1)). The model significantly outperformed chance

(all double-sided t-tests for accuracy: Rat 1: t(998) = 41.5, p = 3.3×10^{-220}; Rat 2: t(998)

$= 28.7$, $p = 7.8 \times 10^{-133}$; Rat 3: $t(998) = 122.6$, $p = 0$; Rat 4: $t(998) = 118.5$, $p = 0$; Rat 5:

$t(998) = 62.6$, $p = 0$). Bars indicate standard error; significance denoted as ***$p < 10^{-10}$.

e. The same five models from panel d were now applied to environment B and

outperformed models trained on shuffled data in decoding the temporal aspects of the

CS/US periods, indicating that fine-grained temporal encoding is stable across

environments (all double-sided t-tests for accuracy: Rat 1: $t(998) = 55.1$, $p = 4.9 \times 10^{-305}$;

Rat 2: $t(998) = 9.3$, $p = 1.1 \times 10^{-19}$; Rat 3: $t(998) = 71.4$, $p = 0$; Rat 4: $t(998) = 62.6$, $p =$

0; Rat 5: $t(998) = 106.2$, $p = 0$). Bars indicate standard error; significance denoted as

***$p < 10^{-10}$.

f. Same analysis as in panel c, but for a CEBRA model trained on CS/US periods split into

five time bins. Top: The five divisions of the conditioning period are shown. Data from

Rat 3 show that the model trained on session A(n) outperformed models trained on

shuffled data when decoding both session A(n-1) and session B(1) (double-sided t-tests,

$t(998) = 88.9$, $p = 0$, and $t(998) = 71.4$, $p = 0$, respectively).

g. Confusion matrices displaying CEBRA decoding of five CS/US time bins, as shown in

panel f. Top row: Models trained on data from session A(n). Bottom row: Models trained

on shuffled data. Darker colors along the diagonal indicate higher model accuracy.

h. Model accuracy for decoding sessions A(n-1) and B(1) using a model trained on data

from session A(n) with CSUS2 and CSUS5 divisions. For both CSUS2 and CSUS5, a

model trained in session A(n) decoded session B(1) with accuracy similar to that

achieved when decoding session A(n-1) (CSUS2: double-sided t-test, $t(8) = -0.13$, $p >$

0.05; CSUS5: $t(8) = 0.32$, $p > 0.05$). Bars represent standard error

Fig 6

	A(n-1)	A(n)	B(1)	B(2)	A(n-1) Shuffled	A(n) Shuffled	B(1) Shuffled	B(2) Shuffled
A(n)	96.1							
B(1)	98.9	96.7						
B(2)	80.5	92.3	95.5					
A(n-1) Shuffled	0.4	0.2	0.1	0.8				
A(n) Shuffled	2.1	0.4	0.5	1.7	1.3			
B(1) Shuffled	2.7	2.4	3.7	2.9	0.6	0.2		
B(2) Shuffled	1.3	1.1	0.8	0.5	0.3	0.9	0.3	

	A(n-1)	A(n)	B(1)	B(2)	A(n-1) Shuffled	A(n) Shuffled	B(1) Shuffled	B(2) Shuffled
A(n)	99.0							
B(1)	98.3	99.0						
B(2)	97.3	97.2	97.0					
A(n-1) Shuffled	0.5	0.5	0.6	0.5				
A(n) Shuffled	0.4	0.3	0.4	0.2	0.2			
B(1) Shuffled	2.5	3.0	2.6	3.1	0.9	1.0		
B(2) Shuffled	11.2	12.2	11.5	11.9	0.5	0.2	0.5	

Fig 6. High consistency in neural representations between environments A and B.

a.   Consistency scores for each rat, calculated with 2, 3, 5, 7, and 10 latents. Lighter bars

represent consistency percentage for actual data, while adjacent darker bars represent

consistency for shuffled data. The schematic below the x-axis illustrates the data sets

included in each comparison (for position and labelling of data sets see figure 6b).

For the CSUS2 division of conditioning periods (split into CS and US components),

all five rats show significantly higher consistency between environments A and B in

the actual data compared to shuffled data. This consistency remains significant with

up to 10 latents.

b. Example consistency measurements from Rat 5 for CSUS2 with 2, 3, 5, 7, and 10

latents.

c. Average consistency scores across all rats for CSUS2. The figure legend follows the

format of panel a. Bars represent standard error. Consistency scores for actual data

versus shuffled data were significantly different across all latent dimensions (all

double-sided t-tests: 2 latents, $p < 1 \times 10^{-5}$; 3 latents, $p < 1 \times 10^{-4}$; 5 latents, $p < 1 \times$

10^{-5}; 7 latents, $p < 1 \times 10^{-5}$; 10 latents, $p < 1 \times 10^{-4}$).

d.   Consistency scores for each rat, as in panel a, but for CSUS5, where the conditioning

period is divided into five time bins. All five rats show significantly higher

consistency between environments A and B in the actual data compared to shuffled

data; this is maintained across latents for up to 10.

e.   Example consistency measurements from Rat 3 for CSUS5 with 2, 3, 5, 7, and 10

latents.

f.   Average consistency scores across all rats for CSUS5. The figure legend matches that

of panel c. Bars represent standard error. Significant differences between actual and

shuffled data were observed for all latent dimensions (all double-sided t-tests: 2

latents, $p < 1 \times 10^{-12}$; 3 latents, $p < 1 \times 10^{-7}$; 5 latents, $p < 1 \times 10^{-6}$; 7 latents, $p < 1 \times$

10^{-5}; 10 latents, $p < 1 \times 10^{-5}$).

Figure 7. High degree of consistency between the conditioning representations across animals

a. Consistency across animals for CSUS2 with 2, 3, 5, 7, and 10 latents. The larger graph highlights the bracketed area from the graph with two latents.

b. Conditioning period is divided into 2 time bins (CSUS2): the similarity between models across animals is not significantly different from the similarity between models for an individual animal, regardless of the number of latents (2 latents: $t(188) = -0.45$, $p > 0.05$;

3 latents: $t(188) = -1.57$, $p > 0.05$; 5 latents: $t(188) = -0.40$, $p > 0.05$; 7 latents: $t(188) = -$

0.22, $p > 0.05$; 10 latents: $t(188) = 0.40$, $p > 0.05$). Error bars represent standard error.

c. Consistency across animals for CSUS5 with 2, 3, 5, 7, and 10 latents. The larger graph

highlights the bracketed area from the graph with two latents.

d. Conditioning period is divided into 5 time bins (CSUS5); the similarity across animal

models remains comparable to that for within-animal models, regardless of the number of

latents (2 latents: $t(188) = -0.79$, $p > 0.05$; 3 latents: $t(188) = 0.25$, $p > 0.05$; 5 latents:

$t(188) = -0.42$, $p > 0.05$; 7 latents: $t(188) = 0.70$, $p > 0.05$; 10 latents: $t(188) = 0.30$, $p >$

0.05). Error bars represent standard error.

Methods

LEAD CONTACT AND MATERIALS AVAILABILITY

Questions and requests for information should be directed to and will be fulfilled by the Lead Contact, Hannah Wirtshafter (hsw@northwestern.edu). This study did not generate new unique reagents. The data that support the findings of this study are available from the corresponding author.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

All procedures were performed within Northwestern Institutional Animal Care and Use Committee and NIH guidelines. Five male Long Evans rats (275–325 g) were sourced from Charles River Laboratories, injected with AAV9-GCaMP8m, implanted with a 2-mm GRIN lens, and trained and tested on eyeblink conditioning in two apparatuses (Fig. 1). Animals were individually housed in an animal facility with a 12/12 h light/dark cycle.

METHOD DETAILS

GCaMP7c injection, lens implantation, EMG implantation

GCaMP8 injection and lens implantation were completed as reported in Wirtshafter and Disterhoft, 2022 and Wirtshafter and Disterhoft, 2023[1,2]. Briefly, rats were anesthetized with isoflurane (induction 4%, maintenance 1-2%) and a craniotomy was performed at stereotaxic coordinates Bregma AP −4.00mm, ML 3.00mm. 0.06uL of GCaMP8m (obtained from AddGene, packaged AAV9 of pGP-AAV-syn-jGCaMP8m-WPRE, lot v175525, titer 1.3E+13 GC/mL) was injected over 12 minutes (approximate coordinates Bregma AP −4.00mm, ML 3mm, DV 2.95mm relative to skull); then the syringe was raised 0.2mm and an additional 0.6ul

of GCaMP7 was injected. We repeated this process once more and at slightly different

coordinates in the craniotomy hole, resulting in 4 total injections.

We then aspirated tissue from the craniotomy site using a vacuum pump and 25 gauge

needle. Tissue was aspirated up to and including the horizontal striations of the corpus collosum.

A 2mm GRIN lens (obtained from Go!Foton, CLH lens, 2.00mm diameter, 0.448 pitch, working

distance 0.30mm, 550nm wavelength) was then inserted into the craniotomy hole and cemented

in place using dental acrylic. Animals were given buprenorphine (0.05mg/kg) and 20mL saline,

taken off anesthesia, and allowed to recover in a clean cage placed upon a heat pad.

Six to eight weeks after surgery, animals were again anesthetized with isoflurane and

checked for GCaMP expression. If expression was seen, baseplates were attached using UV-

curing epoxy and dental acrylic. Electrode implantation to record obicularis oculi

electromyographic (EMG) activity occurred in the same surgery as baseplate attachment, as

described previously[3,4]. Briefly, a connector containing 5 wires was cemented on the front of the

animal's head: 4 wires were implanted directly above the eye in the surrounding muscle (2 for

recording, 2 for electrical stimulation). An additional wire was attached to a connector attached

to a ground screw located above the cerebellum; this screw was implanted during lens

implantation surgery.

1032

Behavioral environment and training

Two behavioral apparatuses were used in these experiments: Environment A was a

78.7cm x 50.8cm unscented rectangular enclosure with wire floor and walls and white lighting.

Environment B was a 50.1cm x 34.9cm scented (with two dabs of clove essential oil on opposite

walls) ovular enclosure with white solid floor and walls, and red lighting. Both environments

were located at the same spot in the room relative to external cues (see Figures 1b and S1).

A tether containing a plug to relay the EMG activity and to deliver a shock to the rat's eye

was attached to a the eyeblink connector on the rat's head. The miniscope was plugged into the

cemented baseplate. The miniscope and EMG cords were all attached to a commutator for ease

of animal movement.

The CS was a 250ms, 85dB free-field tone (5ms rise-fall time). The US was a 100ms

shock directed to the left eye. Shock amount varied per session per animal and was calibrated, if

needed, at the end of a training session for the next session's training. Shock level was deemed

appropriate when a shock was met with a firm shake of the animal's head.

The trace interval was 500ms and the intertrial interval (ITI) was randomized between

30s and 60s, with a 45s average. EMG signal output was amplified (5000×) and filtered (100 Hz

to 5 kHz), then digitized at 3 kHz and stored by computer.

A conditioned response (CR) was identified as an increase in integrated EMG activity

that exceeded the baseline mean amplitude by more than four standard deviations, sustained for a

minimum duration of 15ms. Baseline mean amplitude was calculated during the 500ms

preceding CS onset. Additionally, the response had to commence at least 50ms after the

conditioned stimulus (CS) onset and before the unconditioned stimulus (US) onset.

The animal's first exposure to each environment was a 38min exploration session, in

which the animal was able to freely move and explore the environment without any conditioning

(Figure 1a). Animals were then trained in one environment per session, with no more than one

session per day, and were considered to have learned the task after reaching criterion (70% CRs

in 50 trials) on three consecutive training sessions (termed 'criterion sessions') or when the

previous four training sessions averaged over 70% (in this instance, only the final three of those

sessions were considered 'criterion sessions'). Following the last session in environment A, the

animal was given an exploratory session in environment B. The session after that, the animal was

tested on eye blink conditioning in environment B, using the same parameters as used in

environment A.

1065

1066   **Calcium imaging**

1067   Calcium imaging was completed as reported in Wirtshafter and Disterhoft, 2022 and

1068   Wirtshafter and Disterhoft, 2023[1,2]. Briefly, calcium imaging was done using UCLA V4

1069   Miniscopes[5,6], assembled with two 3mm diameter, 6mm FL achromat lens used in the objective

1070   module and one 4mm diameter, 10mm FL achromat lens used in the emission module.

1071

1072   **QUANTIFICATION AND STATISTICAL ANALYSIS**

1073   Means are presented as mean+-standard deviation. All analysis code is available

1074   at https://github.com/hsw28/ca_imaging and https://github.com/hsw28/Hannahs-CEBRAs. Code

1075   to create specific figures is also available at the former github repository.

1076

1077   **Position and speed analysis**

1078   Position was sampled by an overhead camera at 30Hz. Position tracking was done post-

1079   recording using DeepLabCut[7]. Position was then converted from pixels to cm. Position was

1080   smoothed using a Gaussian filter with standard deviation of 2cm. Speed was calculated by taking

1081   the hypotenuse of the coordinates one before and after the time of interest.

1082

Video pre-processing and cell identification

Video pre-processing and cell identification were performed as reported in Wirtshafter

and Disterhoft, 2022 and Wirtshafter and Disterhoft, 2023[1,2]. In brief, videos were recorded with

Miniscope software at 15frames/second. Video processing was done using CIATAH software[8].

Videos were down sampled in space and normalized by subtracting the mean value of each

frame from the frame. Each frame was then normalized using a bandpass FFT filter (70-

100cycles/pixel) and motion corrected to a using TurboReg[9]. Videos were then converted to

relative florescence (dF/F_0); F_0 was the mean over the entire video.

Cells were automatically identified using CIATAH[8] using CNMF-E[10]. Images were

filtered with a gaussian kernel of width 2 pixels and neuron diameter was set at a pixel size of 8.

The threshold for merging neurons was set at a calcium trace correlation of 0.65; neurons were

merged if their distances were smaller than 4 pixels and they had highly correlated spatial shapes

(correlation>0.8) and small temporal correlations (correlation <0.4).

In vivo calcium imaging involves detecting changes in intracellular calcium levels, which

serve as proxies for neuronal activity. Calcium events refer to transient increases in calcium

concentration above a threshold level; these crossings putatively correspond to spikes in neuronal

firing. These events typically appear as peaks in the data and indicate an active response from the

neuron. Calcium traces are continuous recordings of calcium levels over time. Thus, calcium

events highlight specific neuronal activations, while calcium traces provide a full temporal

picture of these activations together with baseline activity.

All cells identified using CNMF-E were then scored as neurons or not by a human scorer.

Scoring was also done within CIATAH software in a Matlab GUI. Scoring was done while

visualizing and considering a calcium activity trace, average waveform, a montage of the

candidate cell's Ca2+ events, and a maximum projection of all cells on which the candidate cell

was highlighted. The relative fluorescence ($\Delta F/F_0$) local maxima of each identified cell were

considered calcium event times.

1109

Cell cross registration across sessions and within session

Validation and registration were completed as documented in Wirtshafter and Disterhoft[2].

Briefly, videos underwent five rounds of registration using Turboreg image rotation[9] with the

CIATAH software[8,11]. Background noise, axons, and dendrites were removed using an image

binarization threshold of 40% of the images' maximum value. Cells were matched across

sessions using a distance threshold of a maximum of five pixels, with a minimum 2-D correlation

coefficient of 0.5. Sessions were aligned to session A(n), the last session in environment A.

1117

Place cell identification and computing spatial mutual information

Place cells were identified using mutual information computed when the animals were

running at speeds greater than or equal to 4cm/s. MI was computed for all cells; there was no

calcium event rate criterion for included cells. To be considered significant, the computed mutual

information (MI) must be greater than 95% of MI scores computed 500 times from shuffled

positions[12]. To compute the MI for each cell, the training environments were divided into 2.5cm

x 2.5cm bins. The calcium event rate of each cell and the occupancy of the animal were found

for each bin. Rate and occupancy were smoothed with a Gaussian kernel with filter width of 3cm

and Sigma of 0.5cm. Mutual information was computed during periods of movement as

follows[2,12,13]:

1128
$$p = \frac{P_s}{P_o}$$

1129
$$M_s = \sum P_s$$

1130
$$M_o = \sum P_o$$

1131
$$MI = \sum p * \log_2\left(\frac{p}{M_s * M_o}\right)$$

1132   where:
1133 P_s = calcium event probability in each bin
1134 P_o = occupancy probability at each bin
1135

Mutual information using calcium traces was computed as above, except instead of P_s

being calcium event probability per bin, the value of P_s was the average value of calcium trace in

the bin.

We computed MI using both calcium events and calcium trace data. There was no

significant difference between the number of place cells detected using calcium event data and

calcium trace data, (paired t-test t(24)=1.01, p>0.05). Note that all place cell and place field

measurements are presented with conditioning periods included, as the animal was frequently

moving during conditioning periods. We also computed results while excluding conditioning

periods and found no significant differences.

1145

1146   **Computing CSUS mutual information**

1147   The computation of CSUS mutual information was very similarly to that for spatial

1148   mutual information. A 1.3 second period beginning at the start of the CS tone was either divided

1149   into 2 bins (CSUS-MI2) or 5 bins (CSUS-MI5) (Fig. 3c-3d). Mutual information was then

1150   computed using the following:

1151
$$p = \frac{P_s}{P_o}$$

1152
$$M_s = \sum P_s$$

1153
$$M_o = \sum P_o$$

1154
$$MI = \sum p * \log_2\left(\frac{p}{M_s * M_o}\right)$$

1155   Where:
1156 P_s = calcium event probability in each CSUS bin
1157 P_o = probability of individual CSUS occuring out of all CSUS bins
1158
1159

1160   Mutual information using calcium traces was computed as above, except that P_s did not

1161   represent the calcium event probability per bin, but the average value of calcium trace within the

1162   bin.

1163

Remapping quantification

The place cell center was defined as the occupancy-normalized location with maximum

number of calcium events while the animal was moving at 5cm/s or faster. Position was binned

into 2.5cm square bins. The place cell centers at environments A and B, as well as at

environment A across days and at environment B across days, were used to align each

environment across days, as well as to align environment A to environment B.

Population vector correlation was calculated between two environments using calcium

event data. Neurons present in both datasets (such as sessions A(n) and A(n-1), or A(n) and B(1))

were identified and their calcium event times were converted to rates using 0.75 second binning.

These firing rates were then normalized using z-score normalization across each neuron's activity

across time. The mean calcium event rate for each neuron in each environment was then

computed. The correlation among population vectors of firing rates in the two environments was

cpmputed:

1177

1178
$$population\ vector\ correlation = \frac{\sum_{i=1}^{N}\left(r_{\{i,A\}} - \overline{\{r\}}_A\right)\left(r_{\{i,B\}} - \overline{\{r\}}_B\right)}{\sqrt{\sum_{i=1}^{N}\left(r_{\{i,A\}} - \overline{\{r\}}_A\right)^2}\sqrt{\sum_{i=1}^{N}\left(r_{\{i,B\}} - \overline{\{r\}}_B\right)^2}}$$

1179
1180
1181   *Where:*
1182 $r_{\{i,A\}}$ and $r_{\{i,B\}}$ = firing rates of neuron i in environments A and B
1183 $\overline{\{r\}}_A$ and $\overline{\{r\}}_B$ = mean firing rates across neurons in environments A and B
1184   The factors in the denominator correspond to the standard deviation of the components of
1185   each population vector relative to their mean, computed in each environment.
1186

Use of CEBRA versus alternative methods

1188   We explored multiple different methods before settling on the use of CEBRA for this

1189 study. A short summary of each tested method can be found below:

1190

• **Principal Component Analysis (PCA)**[14]: Principal component analysis (PCA) is a

statistical method used to reduce the dimensionality of data while retaining as much

variability as possible. This linear technique identifies the axes (principal components) in

the dataset that maximize variance. The first principal component explains the most

variance, the second explains the second most, and so on. Principal components are

combinations of original features and may not always have clear or intuitive meanings. In

agreement with previous hippocampal data[15], PCA required upwards of 15-25

components to capture 95% of the variance of the data. In addition, across and within all

sessions and representations (spatial and task representations), the manifolds spanned by

the largest PCs remained highly similar, with small principal angles in pairwise

comparisons. This similarity in the orientation of the leading subspaces suggested that

PCA did not distinguish between spatial or behavioral components of the task (Figure

S5).

• **Independent Component Analysis (ICA)**[16]: Independent Component Analysis (ICA) is

a computational technique used to separate a multivariate signal into additive,

independent components. ICA operates under the assumption that observed data are

linear mixtures of underlying, independent sources. It aims to find a linear transformation

that maximizes the statistical independence of the estimated components. We found that

ICA embeddings were unstable throughout the length of the recordings, and also did not

clearly map onto behavioral states (Figure S6).

• **Isomap**[17]: Isomap is a manifold learning technique that seeks to capture the intrinsic

geometric structure of data. Isomap is useful when linear methods like PCA cannot

capture the intrinsic structure of the data, as it preserves the geodesic (curved) distances

in the reduced dimensionality space. Unlike linear methods such as PCA, Isomap can

capture nonlinear relationships in the data. Interestingly, using Isomap, only about 5

neural modes were required to achieve a residual variance of 5-10%. However, the

embedding shape did not relate to any discernable property of neural data or behavior

(Figure S7). Dimensionality reduction was achieved, but the resulting representations

were not interpretable (Figure S7).

•   **MIND**[15,18]: MIND is a decoding method designed for integrating multiple data modalities

to predict various features, particularly sensory and motor functions. MIND uses

recurrent neural networks whose hidden variables provide a memory mechanism for

remembering previous inputs; this approach is particularly apt for the analysis of time

series data such as neural recordings. While MIND was very robust at distinguishing the

different environments, it was not equipped to handle relatively short signals separated in

time, such as the conditioning trials separated by intertrial intervals. Our analyses using

MIND resulted in poor and unstable embeddings that could not be analyzed (Figure S8).

CEBRA[19] was chosen for this project for its ability to capture nonlinear relationships in the data

and to create stable embeddings over short and long time periods. Additionally, spatial

separations of components were well isolated and correlated well with observed behaviors.

1231

Use of CEBRA for position decoding

Optimal parameters for decoding the position of each animal from neural activity were

determined using an extensive grid search across learning rate, temperature, and number of

iterations (Figure S9). Models created to compare different sessions, such as a model trained on

data from session A(n) used to decode session B(1), were only trained on cells that occurred in

both sessions. Models were trained on spike traces of these cells, labeled with the animal's (X,Y)

position. In all cases, 75% of data was used to train the model while 25% of data was held out for

verification. All models were run 500 times. Optimal embeddings were determined based on the

minimum median absolute error between the predicted and true positions. The optimal

parameters for each rat are as follows:

	Rat 1	Rat 2	Rat 3	Rat 4	Rat 5
Model Architecture	'Offset10-model'				
Batch size	512				
Learning rate	$5.5*10^{-5}$	$6.625*10^{-4}$	$5.5*10^{-4}$	$1.0*10^{-3}$	$1.0*10^{-3}$
Temperature mode	'Auto'				
Minimum temperature	No minimum	1.5	0.95	$1.0*10^{-9}$	No minimum
Output dimensions (# of latents)	3				
Max iterations	25000	8000	26500	30000	18000
Distance	'Cosine'				
Conditional	'Time delta'				
Number of hidden units	32				
Time offsets	1				

1242

1243   The number of output dimensions was chosen based on the fewest number of dimensions under

1244   which all 5 models consistently outperformed shuffled data for both position and conditioning

1245   decoding (Figure S10-12).

1246

1247   **Use of CEBRA for conditioning decoding**

1248   As in position decoding, the optimal parameters for decoding conditioning were

determined for each animal using an extensive grid search across learning rate, temperature, and

number of iterations (Figure S11). Models created to compare different sessions of neural

activity, such as a model trained on data from session A(n) used to decode session B(1), were

only trained on cells that occurred in both sessions. Models were trained on spike traces of these

cells, with labels corresponding to the CSUS bin during which the signal occurred (either one out

of 2 bins or out of 5 bins, see Figures 3c-d). In all cases, 75% of data was used to train the model

while 25% of data was held out for verification. All models were run 500 times. Optimal

embeddings were determined based on the percent of correctly binned time points. The optimal

parameters for each rat are as follows:

1258

	Rat 1	Rat 2	Rat 3	Rat 4	Rat 5
Model Architecture	'Offset10-model'				
Batch size	512				
Learning rate	$3.5*10^{-3}$	$7.0*10^{-3}$	$3.5*10^{-3}$	$7.5*10^{-3}$	$9.5*10^{-3}$
Temperature mode	'Constant'	'Constant'	'Auto'	'Constant'	'Constant'
Minimum temperature	2.33	1.75	1.67	1.67	2.66
Output dimensions (# of latents)	3				
Max iterations	50000	7500	20000	18000	25000
Distance	'Euclidian'	'Cosine'	'Cosine'	'Euclidian'	'Cosine'
Conditional	'Time delta'				
Number of hidden units	32				
Time offsets	1				

1259

1260   The number of output dimension was chosen based on the fewest number of dimensions

1261   under which all 5 models consistently outperformed shuffled data for both position and

1262   conditioning decoding (Figure S10-12). The parameters listed above were used for decoding into

1263   2 or 5 bins, including the use of 3 output dimensions (# of latents).

1264   The accuracy of results was computed from the entries in the confusion matrix:

1265

1266 $$Accuracy = \frac{Sum\ of\ number\ of\ diagonal\ (correct)\ elements}{Sum\ of\ number\ of\ total\ elements}$$

1267

1268   Precision was calculated for each class i, $1 \leq i \leq n$, where n is the number of classes:
1269

1270
$$Precision_i = \frac{(TP_i)}{TP_i + FP_i}$$

1271   where:
1272 $TP = true\ positives$
1273 $FP = false\ positives$
1274
1275   The global precision is given by the average:

1276
$$Precision = \frac{1}{n} \sum_{i=1}^{n} Precision_i$$

1277
1278
1279   Recall, also known as sensitivity, was calculated for each class i, $1 \leq i \leq n$, where n is the

1280   number of classes:

1281

1282
$$Recall_i = \frac{(TP_i)}{TP_i + FN_i}$$

1283
1284   where:
1285 $TP = true\ positives$
1286 $FN = false\ negatives$
1287
1288   The global recall is given by the average:
1289

1290
$$Recall = \frac{1}{n} \sum_{i=1}^{n} Recall_i$$

1291
1292   The F1 score was calculated for each class i, $1 \leq i \leq n$, where n is the number of classes:
1293

1294
$$F1_i = 2 * \frac{Precision_i * Recall_i}{Precision_i + Recall_i}$$

1295
1296   The global F1 score is given by the average:
1297

1298
$$F1 = \frac{1}{n} \sum_{i=1}^{n} F1_i$$

1299
1300
1301   The area under the receiver operating characteristic (ROC) curve was calculated for each class i,

1302 $1 \leq i \leq n$, where n is the number of classes:

1303

$$AUC_i = roc\_auc\_score(y_{true-bin}[:,i], y_{pred-prob}[:,i])$$

1304

1305

1306   The global value is given by the average:

1307

$$ROC\ AUC = \frac{1}{n}\sum_{i=1}^{n} AUC_i$$

1308

1309

1310   where:

1311 $y_{true-bin} = binarized\ true\ labels$

1312 $y_{pred-prob} = predicted\ probabilities\ for\ each\ class$

1313

Model consistency

Model consistency was computed using a built-in CEBRA function which relies on the

function 'sklearn.metrics.consistency_score'[20]. The function compares the embeddings from

different models by calculating pairwise consistency scores. This comparison involves

measuring the similarity of the embeddings using statistical metrics; i.e. this metric calculates

how similar a model's labels are for similar instances in the data set.

To determine consistency between environments and across animals, the data was fit to

each model 20 times. The model with the lowest loss was selected and compared to other models

with the lowest loss. Models were created using each individual animal's optimal parameters

(see above).

1324

1325

1326 **Acknowledgements:** This work was supported by an NIA T32 (T32-AG020506/AG/NIA), an

1327 NIA R37 (R37-AG008796/AG/NIA), an NINDS R01 (R01 NS113804/NS/NINDS), and a K99

1328 award (K99 MH135062).

1329

1330 This research was supported in part through the computational resources and staff assistance

1331 provided by the Quest high performance computing facility at Northwestern University, which is

1332 jointly supported by the Office of the Provost, the Office for Research, and Northwestern

1333 University Information Technology.

1334

1335 We would like to thank all members of the Disterhoft lab, especially Mackenzie Kneisly. We

1336 would like to thank the following individuals for their assistance with CEBRA: Mackenzie

1337 Mathis and Steffen Schneider. And additional thank you to David Wirtshafter for his feedback.

1338

1339

1340 **Author contributions:** Investigation, H.S.W.; Formal Analysis, H.S.W.; Writing –Original

1341 Draft, H.S.W.; Writing – Review & Editing, H.S.W., S.A.S., J.F.D.; Funding: H.S.W, J.F.D.,

1342 Supervision, S.A.S., J.F.D.

1343

1344 **Declaration of interests:** The authors declare no competing interests.

1345

1353 **Supplementary figures**

1354

Figure S1

1355

Figure S1. Photos of testing chambers.

Photos of testing chambers. Top: Environment A, an unscented rectangular enclosure with wire

floor and walls, and white lighting. Bottom: Environment B, a scented ovular enclosure with

white solid floor and walls, and red lighting. Both environments were located at the same spot in

the room relative to external cues. The door to the chamber was closed during testing, to

accentuate the distinction between the white and red lighting.

1362

Figure S2. Learning curves for the five rats.

Substantial variability in the number of sessions required to learn the task; average number of sessions was 20 ± 4.2 (including the criterion sessions). The fastest learners (2 rats) reached criterion after 14 sessions, while the slowest rat required 24 sessions to reach criterion.

1367

1368 **Figure S3. Percent of place cells by session for each animal.**

1369 The percentage of cells classified as place cells for each session and each animal

1370

Figure S4. CSUS-MI2 and CSUS-MI5 differences between sessions.

    a. The trial period was divided into two segments: the CS and trace period (750 ms) and the

        US and post-US period (500 ms). Mutual information (MI) was calculated for cells based

        on these two periods and compared to shuffled data, where period IDs were shuffled 500

        times across all trials. Left: Using calcium event data, we found that $10.7\% \pm 4.9\%$ of

        cells contained significant CSUS information related to whether the animal was in a CS

        or US period. Right: Using calcium traces, $19.9\% \pm 8.2\%$ of cells contained significant

        information distinguishing the CS from the US period. No significant differences in

CSUS-MI were observed between environments A and B (double-sided t-tests, calcium

events: $t(23) = 0.48$, $p > 0.05$; calcium traces: $t(23) = -0.52$, $p > 0.05$).

b. The trial period was divided into five equal segments, each 250 ms. MI was calculated

for each cell based on these five periods. We compared the observed values to those

obtained after shuffling period IDs 500 times. Left: Using calcium event data, $15.5\% \pm$

7.8% of cells contained significant information distinguishing the five periods, compared

to $10.0\% \pm 7.8\%$ when using calcium trace data. No significant differences in these MI

metrics were found between environments A and B (double-sided t-tests, calcium events:

$t(23) = -0.32$, $p > 0.05$; calcium traces: $t(23) = -1.1$, $p > 0.05$).

c. Left: There was no significant difference in CSUS-MI2 values when comparing session

A(n) to session A(n-1) versus session A(n) to session B(1) (Wilcoxon rank sum test: $p >$

0.05; double-sided t-test: $t(1431) = 0.86$, $p > 0.05$). Right: A small but significant

difference was observed in CSUS-MI5 when comparing session A(n) to session A(n-1)

versus session A(n) to session B(1) (Wilcoxon rank sum test: $p = 0.049$; double-sided t-

test: $t(1431) = -2.2$, $p = 0.03$).

1394

Figure S5

1395

Figure S5. PCA computations for session A(n) and session B(1).

Only cells present in both sessions were used. Principal component analysis (PCA) revealed that

approximately 15-25 principal components (PCs) are needed to account for 95% of the variance

in the data. When using the complete cell population (not shown), more than 25 PCs are required

to achieve the same variance. Across and within all sessions and representations (spatial and

task), the principal angles between manifolds remain highly similar.

1402

Figure S6. ICA computations across different segments of a session.

Top: Independent component analysis (ICA) was for data over an entire session, using three

independent components (ICs). Blue dots represent non-trial times, while red dots represent trial

times. Middle: ICA computed over the last two-thirds of the same session shows variability in

1407   ICs across segments within a session. Bottom: ICA computed over the second half of the session

1408   shows additional variability in the results of the analysis depending on how the session is

1409   divided. These results indicate that independent components are highly variable over the course

1410   of one session and are sensitive to how the session is partitioned.

1411

1412

Figure S7. Isomap computations for session A(n) and session B(1).

Isomap computations suggest that approximately five neural modes are sufficient to achieve a

residual variance of 5-10%. However, the shape of the Isomap embedding does not correlate

with any discernable properties of neural activity or behavior, suggesting limited interpretability

of the embedding structure.

Figure S8

1418

Figure S8. MIND outputs for sessions A(n), B(1), and concatenated sessions.

Top row: MIND embeddings during movement, excluding trial periods, with color bars

representing frames. The temporal structure of the data is well captured, with clear separation

between A(n) and B(1). Bottom row: MIND embeddings during conditioning periods are highly

unstable. Small changes in parameters result in substantial shifts in the embedding structure,

transitioning from a linear structure (left) to an undefined, unstable cloud (middle and right).

1425

1426

Figure S9. Grid search over decoding parameters for position.

A grid search was performed over three parameters: minimum temperature, learning rate, and

number of iterations, for decoding position. Models were trained using cells from session A(n)

that also appeared in session A(n-1). The figure shows decoding accuracy for session A(n-1)

using the models trained on data from session A(n). Yellow areas indicate higher decoding

accuracy.

1433

Figure S10. Position decoding error as a function of the number of latents (Rat 5).

This figure shows the decoding error for position as the number of latents increases. Left panels:

model trained on data from A(n), tested on held out data from A(n). Upper left panel: As the

number of latents increases, the model's ability to decode a different session within the same

environment decreases. This effect is not consistent across all rats; see lower left panel. Right

panels: model trained on data from A(n), tested on A(n-1) (top) or B(1) (bottom). Performance is

particularly bad when the model is tested on a different environment (lower right panel).

1441

Figure S11. CSUS2 decoding accuracy with increasing number of latents (Rat 3).

Percent of incorrect decoding for the CSUS2 model as the number of latents increases. Left

panels: model trained on data from A(n), tested on held out data from A(n). As the number of

latents increases, the model's ability to decode a different session within the same environment

remains stable or slightly decreases. Right panels: model trained on data from A(n), tested on

A(n-1) (top) or B(1) (bottom). Performance remains stable for a different session in the same

environment (upper right panel) but deteriorates with increasing number of latents when the

model is tested in a different environment (lower right panel). Each model was run 100 times.

Figure S12

1450

Figure S12. CSUS5 decoding accuracy with increasing number of latents (Rat 5).

Same as Figure S11, but for CSUS5; the conditioning period has been divided into five time bins

instead of two. The percent of incorrect decoding is shown as the number of latents increases.

Left panels: model trained on data from A(n), tested on held out data from A(n). As the number

of latents increases, the model's ability to decode a different session within the same environment

remains stable or slightly increases. Right panels: model trained on data from A(n), tested on

A(n-1) (top) or B(1) (bottom). Performance remains stable for a different session in the same

environment (upper right panel) but deteriorates with increasing number of latents when the

model is tested in a different environment (lower right panel). Each model was run 100 times.

1460

1461

Figure S13. Grid search over decoding parameters for conditioning.

A grid search was performed over three parameters: minimum temperature, learning rate, and

number of iterations for conditioning decoding. Models were trained using cells from session

A(n) that also appeared in session B(1). The figure shows decoding accuracy for CSUS2 session

B(1) using the models trained on data from session A(n). Yellow areas indicate higher decoding

accuracy. For Rats 1 and 4, the 'Euclidean' distance with 'constant' temperature mode was used.

For Rats 2 and 5, 'cosine' distance with 'constant' temperature mode was used. For Rat 3, 'cosine'

distance with 'auto' temperature mode was used.

1470

1471

Methods References

Wirtshafter, H. S. & Disterhoft, J. F. Place cells are nonrandomly clustered by field location in CA1 hippocampus. *Hippocampus* **33**, 65-84 (2023). https://doi.org:10.1002/hipo.23489

Wirtshafter, H. S. & Disterhoft, J. F. In Vivo Multi-Day Calcium Imaging of CA1 Hippocampus in Freely Moving Rats Reveals a High Preponderance of Place Cells with Consistent Place Fields. *J Neurosci* (2022). https://doi.org:doi: 10.1523/JNEUROSCI.1750-21.2022

Weiss, C. & Thompson, R. F. The effects of age on eyeblink conditioning in the freely moving Fischer-344 rat. *Neurobiol Aging* **12**, 249-254 (1991). https://doi.org:10.1016/0197-4580(91)90105-s

Skelton, R. W. Bilateral cerebellar lesions disrupt conditioned eyelid responses in unrestrained rats. *Behav Neurosci* **102**, 586-590 (1988). https://doi.org:10.1037//0735-7044.102.4.586

Aharoni, D., Khakh, B. S., Silva, A. J. & Golshani, P. All the light that we can see: a new era in miniaturized microscopy. *Nat Methods* **16**, 11-13 (2019). https://doi.org:10.1038/s41592-018-0266-x

Silva, A. J. Miniaturized two-photon microscope: seeing clearer and deeper into the brain. *Light: Science & Applications* **6**, e17104-e17104 (2017).

Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nature neuroscience* **21**, 1281-1289 (2018).

Corder, G. *et al.* An amygdalar neural ensemble that encodes the unpleasantness of pain. *Science* **363**, 276-281 (2019). https://doi.org:10.1126/science.aap8586

Thevenaz, P., Ruttimann, U. E. & Unser, M. A pyramid approach to subpixel registration based on intensity. *IEEE Trans Image Process* **7**, 27-41 (1998). https://doi.org:10.1109/83.650848

Zhou, P. *et al.* Efficient and accurate extraction of in vivo calcium signals from microendoscopic video data. *eLife* **7** (2018). https://doi.org:10.7554/eLife.28728

CIAtah: a software package for analyzing one- and two-photon calcium imaging datasets. v. v1.0.0 (Zenodo, 2018).

Kinsky, N. R., Sullivan, D. W., Mau, W., Hasselmo, M. E. & Eichenbaum, H. B. Hippocampal Place Fields Maintain a Coherent and Flexible Map across Long Timescales. *Current biology : CB* **28**, 3578-3588 e3576 (2018). https://doi.org:10.1016/j.cub.2018.09.037

Olypher, A. V., Lansky, P., Muller, R. U. & Fenton, A. A. Quantifying location-specific information in the discharge of rat hippocampal place cells. *J Neurosci Methods* **127**, 123-135 (2003). https://doi.org:10.1016/s0165-0270(03)00123-7

Pearson, K. LIII. On lines and planes of closest fit to systems of points in space. *The London, Edinburgh, and Dublin philosophical magazine and journal of science* **2**, 559-572 (1901).

Nieh, E. H. *et al.* Geometry of abstract learned knowledge in the hippocampus. *Nature* **595**, 80-84 (2021).

1516   16   Ans, B., Hérault, J. & Jutten, C. Architectures neuromimétiques adaptatives: Détection
1517   de primitives. *Proceedings of Cognitiva* **85**, 593-597 (1985).
1518   17   Tenenbaum, J. B., Silva, V. d. & Langford, J. C. A global geometric framework for
1519   nonlinear dimensionality reduction. *science* **290**, 2319-2323 (2000).
1520   18   Low, R. J., Lewallen, S., Aronov, D., Nevers, R. & Tank, D. W. Probing variability in a
1521   cognitive map using manifold inference from neural dynamics. *BioRxiv*, 418939 (2018).
1522   19   Schneider, S., Lee, J. H. & Mathis, M. W. Learnable latent embeddings for joint
1523   behavioural and neural analysis. *Nature* **617**, 360-368 (2023).
1524   https://doi.org:10.1038/s41586-023-06031-6
1525   20   Pedregosa, F. *et al.* Scikit-learn: Machine learning in Python. *the Journal of machine*
1526   *Learning research* **12**, 2825-2830 (2011).
1527