Statistical method accounts for microscopic electric field distortions around neurons when simulating activation thresholds

Konstantin Weise[1,2*], Sergey N. Makaroff[3,4], Ole Numssen[2], Marom Bikson[5], Thomas R. Knösche[2,6]

[1]Leipzig University of Applied Sciences, Leipzig, Germany

[2]Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

[3]ECE Department, Math Department, Worcester Polytechnic Institute, Worcester MA USA

[4]Massachusetts General Hospital, Boston MA USA

[5]The City College of New York, NY USA

[6]Technical University of Ilmenau, Germany

*corresponding author (kweise@cbs.mpg.de)

Abstract

Introduction: Notwithstanding advances in computational models of neuromodulation, there are mismatches between simulated and experimental activation thresholds. Transcranial Magnetic Stimulation (TMS) of the primary motor cortex generates motor evoked potentials (MEPs). At the threshold of MEP generation, whole-head models predict macroscopic (at millimeter scale) electric fields (50-70 V/m) which are considerably below conventionally simulated cortical neuron thresholds (175-350 V/m).

Methods: We hypothesize that this apparent contradiction is in part a consequence of electrical field warping by brain microstructure. Classical neuronal models ignore the physical presence of neighboring neurons and microstructure and assume that the macroscopic field directly acts on the neurons. In previous work, we performed advanced numerical calculations considering realistic microscopic compartments (e.g., cells, blood vessels), resulting in locally inhomogeneous (micrometer scale) electric field and altered neuronal activation thresholds. Here we combine detailed neural threshold simulations under homogeneous field assumptions with microscopic field calculations, leveraging a novel statistical approach.

Results: We show that, provided brain-region specific microstructure metrics, a single statistically derived scaling factor between microscopic and macroscopic electric fields can be applied in predicting neuronal thresholds. For the cortical sample considered, the statistical methods match TMS experimental thresholds.

Conclusions: Our approach can be broadly applied to neuromodulation models, where fully coupled microstructure scale simulations may not be computationally tractable.

Keywords:

Multiscale brain modeling, brain stimulation, biophysical modeling, TMS

1. Introduction

Transcranial magnetic stimulation (TMS) of the primary motor cortex (M1) causes peripheral muscle activation, reflected by motor evoked potentials (MEP) recorded from surface electrodes over respective muscles. Such experiments are valuable for studying the motor system and its pathologies (e.g., Di Lazzaro & Ziemann, 2013), and underpin individual dosing of repetitive TMS (rTMS) therapies, such as for depression (Rossi et al., 2021). The amplitude of the MEP scales with the TMS device output and, more directly, with the electric field the relevant neurons are exposed to. Modeling the relation between stimulation intensity and cortical responses underpins explaining TMS and rTMS outcomes.

Numerical field modeling in conjunction with a non-linear regression approach has enabled localization of activated neuronal populations and the derivation of input-output (IO) curves that map TMS induced electric field strengths to the MEP amplitudes (Weise et al., 2020; Numssen et al., 2021; Weise et al., 2023a). With conventional biphasic pulses these sigmoidal IO dose responses have a half-maximum (50% of peak MEP response) at ~50-70 V/m (Numssen et al., 2021). However, explicit simulations of cortical L3/4 neurons predict higher thresholds of ~260 V/m for the same TMS waveform (Weise et al., 2023b) and 175-350 V/m for monophasic pulses (Aberra et al., 2020; Weise et al., 2023b). We hypothesize that this mismatch is a consequence of conventional numerical field models ignoring the presence of microscopic structures (cell membranes, blood vessels).

In calculation of electric fields produced during neuromodulation (TMS), classical models assume macroscopically (mm scale) homogenous tissue. At the microscopic (μm) scale, however, the conductivity is highly inhomogeneous given the low conductivity of cell membranes and vasculature. Recently, this effect has been investigated in detail by Qi et al. (2025) using a high-resolution boundary elements model of a sub-volume (250×140×90 μm) of the L2/3 P36 mouse primary visual cortex with detailed segmentation of microscopic compartments, taking into account neuronal and

glial membranes as well as blood vessels (Turner et al. 2022; MICrONs Consortium 2021), and

comprising ~0.5 billion facets in total.

Adapted from (Qi et al., 2025), Figure 1 shows how low conducting barriers cause charge

accumulation and Fig. 2 demonstrates how this changes the effective electric field at the cell

membranes. This electric field inhomogeneity may explain the apparent discrepancy between the

activation thresholds found in microscopic simulations to those obtained when relating

experimentally observed MEPs to macroscopic electric field simulations of TMS. The goal of this

work is to test this hypothesis.

To this end, we use the electric field simulations performed by Qi et al. (2025) to determine the

microscopic fields at the axons that correspond to a given macroscopic field. As prior simulations

(Aberra et al. (2020); Weise et al. (2023b)) have revealed that the axonal terminals have the lowest

threshold, we consider field differences between both approaches at axon terminals to estimate the

recruitment rate as a function of the macroscopic field. If the microscopic electric fields from Qi et

al. (2025) elicit action potentials at the axon terminals for macroscopic field strengths of 50-70 V/m

(Numssen et al., 2021), then microstructure-dependent electric field warping does indeed account

for the aforementioned activation threshold discrepancy seen in between conventional

(macroscopic) and explicit (microscopic) modeling.

Since realistic microscopic simulations are computationally expensive, it can be impractical to

directly replace the macroscopic field calculation methods with microscopic simulations in routine

analysis. For this reason, the second aim of this study is to provide an easy-to-compute method

that corrects the discrepancy between activation thresholds (or recruitment rates) based on

microscopic and macroscopic fields. Our method offers a principled way to create, for any

neuromodulation technology and any cortical tissue with available microanatomical representation,

a lookup table that maps macroscopic field strength and orientations to recruitment rates.

2. Materials and Methods

2.1 Modeling framework

The present multiscale modeling framework includes three distinct yet interconnected steps, each utilizing a different numerical method:

i. **Biophysical computations**: Activation thresholds for rodent neurons with neuron morphologies from Markram et al. (2015) are computed under external brain stimulation. These computations are performed using the 1D cable equation with appropriate channel parameters.

ii. **Novel statistical model**: Essentially random microscopic field (extracellular potential) fluctuations in the cortical gray matter are statistically incorporated into the standard multiscale brain stimulation pipeline (Aberra et al., 2020; Weise et al., 2023b), which previously accounted only for macroscopic deterministic (smooth) brain stimulation fields.

iii. **Representative set of microscopic field/potential fluctuations**: A realistic set of extracellular microscopic field/potential fluctuations is derived from Qi et al. (2025). This set is used to initialize the parameters of the proposed statistical model.

Below, we describe each modeling step in detail.

2.2 Threshold computations in morphologically realistic neuron models

In extension of an earlier study by Aberra et al. (2020), Weise et al. (2023b) computed the thresholds of different neuronal populations with respect to the electric field at the membrane. Detailed models of a large number of neuronal morphologies, taken from the Blue Brain Project (Markram et al. 2015), were used to account for the natural variability of neurons of different types. These types include layer 2 and 3 pyramidal cells (L2/3 PC), layer 4 small, nested, and large basket cells (L4 S/N/LBC), as well as layer 5 pyramidal cells (L5 PC) from mouse cortex. The neural morphologies were scaled up to human dimensions. In those simulations, we assumed a homogeneous or linearly changing field across the neuron, thus neglecting any effects of the presence of the neurons themselves and other structures. This essentially means treating

microscopic and macroscopic fields (see Introduction) as equal. The firing thresholds were

determined independently for each neuron by applying external electric fields with different angles

with respect to and different gradients along the somato-dendritic axis. Importantly, it turned out

that the initial generation of action potentials (i.e., the excitement of the respective neuron) almost

exclusively occurred at one of the axonal terminals. The excitation then spread over the entire

axonal arbor and activated all other synapses. We utilize these results in the present study.

2.3 Alteration of neuronal recruitment rate due to microscopic field perturbations

To allow drawing general conclusions regarding the relevance of differences between microscopic

and macroscopic electric fields on the apparent excitation threshold of neurons with respect to the

macroscopic field, the problem must be treated statistically. The ratio between microscopic and

macroscopic electric fields at an arbitrary location r on the (axonal) cell membrane $s_E(r) = \frac{E_{micro}(r)}{E_{macro}}$

can be considered a random variable with a probability density $p(s_E)$. Estimating this distribution

requires separate electric field simulations of (homogeneous) macroscopic and (inhomogeneous)

microscopic electric fields in a sample of neural tissue, which are described in the next section. This

provides the essential means to adapt the previously determined thresholds and recruitment rates

of Aberra et al. (2020) and Weise et al. (2023b) with respect to microscopic electric fields. The

corrected recruitment rate r_S of a *single* axon terminal is then given as a function of the external

macroscopic electric field E_{macro} by:

$$
133 \qquad r_s(E_{macro}) = \int\limits_{0}^{\infty} p(E_{thres}) \left(\int\limits_{\frac{E_{thres}}{E_{macro}}}^{\infty} p(s_E)\, ds_E \right) dE_{thres} \qquad (1)
$$

130   where E_{thres} and $p(E_{thres})$ are the threshold field strength and its probability density, respectively,

131   calculated over different samples of neurons (within a particular neuron type) with externally applied

132   electric fields from, e.g., Weise et al. (2023b). The inner integral $\int_{\frac{E_{thresh}}{E_{macro}}}^{\infty} p(s_E)\, ds_E$ describes the

probability that, at any arbitrary axonal terminal, the (relative) microscopic field $s_E = E_{micro}/E_{macro}$ is above a given (relative) threshold E_{thres}/E_{macro}. The outer integral then sums that value over all possible threshold values, weighted by their probabilities, thus yielding the final activation probability (recruitment rate) of axonal terminals.

The fact that excitation of *any* axonal terminal of a given neuron eventually excites the entire axonal arbor with all its terminals (see previous section) leads to a statistical problem that depends on the average number of terminals N per neuron in the respective cell population. Accordingly, the recruitment rate of a population of neurons $r_n(E_{macro})$ can be determined from the recruitment rate from a single axon segment $r_s(E_{macro})$ from eq. (1) as follows:

$$r_n(E_{macro}) = 1 - \left(1 - r_s(E_{macro})\right)^N \qquad (2)$$

2.4 Computing the microscopic extracellular electric field in cortical cube sample

To determine the statistics for the scaling factor between microscopic and macroscopic electric fields $p(s_E)$, we utilized extensive electromagnetic numerical simulations of a neural tissue sample performed with boundary element fast multipole method (BEM-FMM) (Makarov et al 2018, Noetscher et al., 2023). The details and results of these computations are reported elsewhere (Qi et al., 2025). In short, the simulations were based on a 250×140×90 μm cube of neural tissue from mouse primary visual cortex, obtained from electron microscopic images with a resolution of 3.6×3.6×40 nm (the "pinky" dataset of IARPA Phase I). The reconstruction comprises triangulated surfaces (membranes) of various cellular structures, including pyramidal and non-pyramidal neurons, astrocytes, microglia, oligodendrocytes and precursors, pericytes, vasculature, nuclei, mitochondria, etc. (Turner et al. 2022; MICrONs Consortium 2021). The surface resolution (or average computational mesh size) was set at 100 nm to account for microscopic closely spaced objects.

The computation of the extracellular electric field is based on the assumption that membranes are

non-conducting at the end of an initial polarization period, allowing to solve the extracellular field

problem as a decoupled problem, using Neumann boundary conditions at the outer surfaces of the

membranes (for details, see Noetscher et al., 2023; Makaroff et al., 2023; Qi et al., 2025). This

implies that intracellular fields are not computed. Intracellular space is considered isopotential in

the approximation of initial polarization, and intracellular structures, like mitochondria and nuclei,

thus have no influence.

For the numerical treatment of the problem, the BEM-FMM method (Makaroff et al. 2018; Makaroff

et al., 2023; Noetscher et al., 2023) was used. This method computes the electric charges deposited

(or induced) on membrane surfaces in response to an impressed electric field. These charges (more

precisely, the surface charge density) are responsible for the secondary electric field, which serves

as the source of microscopic electric field and extracellular potential perturbations. The secondary

extracellular electric field along its skeleton centerlines was forced to have the mean value zero to

assure that the total extracellular electric field for every neuron within the sample has exactly the

same mean value as in the homogeneous space.

According to classical quasi-static conduction theory (Wang et al., 2024; Jackson 1998, Balanis

2023, Stratton 2006), the resulting field perturbations are relatively strong (on the order of the

impressed field itself), but highly localized. These perturbations typically have a spatial extent

similar to the size of the perturber itself, such as the cross-section of a soma, dendrite, or axon.

However, perturbations of the same polarity can potentially form large clusters, for example, in

regions with several closely spaced somas. The size of such clusters may approach the axonal

space constant, thereby facilitating easier excitation or inhibition of action potentials.

The BEM-FMM approach was specifically adapted to a large neuronal ensemble with several

hundreds of closely spaced neurons and with the overall size of 0.5-1 billion facets, using a nested

iterative algorithm (see Qi et al., 2025). Its idea is to compute charge deposition for each neuron (a

membrane-based surface mesh with ca 1-2 million facets) separately, then update this charge

deposition by the effect of all other neurons, and finally repeat this process iteratively until a proper

convergence criterion is met. While the electromagnetic computational effort has been quite

extensive and took over half a year, its accuracy has been verified by excellent self-convergence.

Since the threshold computations by Weise et al. (2023b) and Aberra et al. (2020) (see Section 2.1)

are based on one-dimensional cable equations, we need the extracellular potential or the collinear

extracellular electric field at the centerlines of the neuronal processes. In the computations by Qi et

al. (2025), however, neurons are intrinsically modeled as three-dimensional objects. Therefore, for

performing the biophysical modeling, we extracted the average solution over the cross-sections of

the processes (dendrites, axons, soma) as described in Qi et al. (2025).

3. Results

3.2 Microscopic electric field variations

Figure 2 illustrates, for an example neuron, the impact of the presence of neurons and other

structures on the local collinear electric field at the membranes. In this particular case, we observe

an elevation of the field maximum by more than a factor of 2. For better visualization, the values

were limited to ±200 V/m, as extreme values can be up to a factor of 10. However, due to the

complex morphology and mutual influences, the locations of field increases and attenuations cannot

be assessed deterministically on a larger scale. This motivates a statistical approach to the problem,

as described in Section 2.2, in order to account for the field variability of the microscopic electric

field when calculating the neuronal thresholds and to correct the values determined by Aberra et al.

(2020) and Weise et al. (2023b).

3.2 Influence of microscopic electric field variations on neural excitability

In total, the activation functions of $\sim 1.4 \cdot 10^6$ axon segments were extracted from the cube sample.

The histogram and the probability density function are shown in Fig. 3. According to that distribution,

the median of the scaling factor is 1.06 and the probability that the electric field at an axon segment

is higher than macroscopically assumed is $p\left(\frac{E_{micro}}{E_{macro}} > 1\right) = 55.8\%$. The distribution also shows that

axon segments can be exposed to field strengths exceeding macroscopic electric field

approximations by a factor of 5 and more. However, also the opposite can be the case, as axon

segments may be exposed to very low field strengths, too.

The probability density function of the electric field scaling factor (Fig. 3), in conjunction with the

results from Weise et al. (2023b), allows for a realistic estimation of the neuronal recruitment rates

with respect to the macroscopic field, using eqs. (1) and (2). Fig. 4 shows an example of a corrected

recruitment rate taking into account microscopic electric field variations for the case of L2/3 PC from

Weise et al. (2023b), where it was assumed that the macroscopic electric field is homogeneous and

points along the somato-dendritic axis towards the soma. We assumed an average number of

terminals per neuron of N=35 (STD=13.7), informed by the L2/3 population used by Weise et al.

(2023b). The grey shaded lines are determined after sampling the number N of terminals per neuron

from a normal distribution with the given mean and standard deviation to illustrate the expected

variability of the recruitment curves. A comparison between the old and new recruitment curves

shows a considerable reduction of the activation threshold from about 225 V/m (dashed black line)

to 30-40 V/m half maximum (red line) when considering the effects of microscopic electric fields.

To enable a comparison of the results with experimental data, we also present the I/O curve of

motor evoked potentials (MEPs) of the first dorsal interosseous (FDI) of a representative subject

from Numssen et al. (2021) as a function of the macroscopic electric field calculated in that study

after successful motor mapping.

A comparison between the recruitment curves shows a clearly improved correspondence to the

field strength values observed in the experiment when considering microscopic electric field effects.

Note that it is straightforward to apply the correction of the recruitment rates to all neuronal

populations, i.e. L2/3 PC, L4 S/N/LBC, and L5 PC presented in Weise et al. (2023b) including

different electric field angles with respect to the cortical normal direction ($\theta = [0°, 180°]$) and linear

field gradients along that direction ($\Delta|\tilde{\mathbf{E}}| = [-20, 20] \, \%/mm$). As an example, Fig. 5 shows the

dependences of the uncorrected and corrected recruitment rates of L2/3 PC stimulated with

biphasic TMS pulses. The correction was applied to all recruitment rate interpolators from all

neuronal populations considered in Weise et al. (2023b) and can be downloaded from Weise et al.

(2025). Fig. 5 (c) and (d) show the recruitment rate curves for field angles $\theta = 0°$, $\theta = 90°$, and $\theta =$

$180°$ without and with microscopic electric field correction, respectively. It can be observed that the

field correction influences the shape of the recruitment rate curves. The directional sensitivity is still

present with comparable ratios. Comparing the required e-field values at a recruitment rate of 0.5

for example results in the case of uncorrected macroscopic fields $E(\theta = 90°) = 260 \, V/m$,

$E(\theta = 0°) = 227 \, V/m$, and $E(\theta = 180°) = 231 \, V/m$, which results in $\frac{E(\theta=0°)}{E(\theta=90°)} = 0.87$ and $\frac{E(\theta=180°)}{E(\theta=90°)} =$

0.89. With microscopic field correction, the electric fields are $E(\theta = 90°) = 43.3 \, V/m$, $E(\theta = 0°) =$

$37.7 \, V/m$, and $E(\theta = 180°) = 39.5 \, V/m$, resulting in $\frac{E(\theta=0°)}{E(\theta=90°)} = 0.87$ and $\frac{E(\theta=180°)}{E(\theta=90°)} = 0.91$.

4. Discussion

In recent years, TMS advances have been supported by macroscopic field stimulations tailored to

individual head and brain morphology (e.g., Makarov et al., 2020; Thielscher et al., 2015). These

models help explain and optimize TMS applications by quantifying cortical stimulation strength in

terms of a physical entity: the induced (macroscopic) electric field strength (Caulfield et al., 2021;

Numssen et al., 2024). However, using the macroscopic (voxel-based) electric field as a proxy for

neural excitation is a simplification, as neurons respond to the electric field in a non-linear fashion.

Significant efforts to account for this relationship have been made by detailed mechanistic modeling

of neurons at different spatial and temporal scales, from simulated single-cell responses to single

pulses (Aberra et al., 2020; Weise et al., 2023b) to network plasticity after repetitive TMS (rTMS;

Shirinpour et al., 2021). However, these approaches have not converged on the neuronal activation

threshold. Single biphasic pulse TMS of the motor cortex typically yields MEP thresholds of 60-100

V/m macroscopic electric fields (Numssen et al., 2021; Numssen, Kuhnke et al., 2024; Rosanova

et al., 2009; Caulfield et al., 2024), while EEG recordings detect changes in cortical activity for rTMS

at around 35 V/m (Zmeykina et al., 2020). This variability across methods may reflect different

noise levels, which affects detectability. In contrast, single-cell simulations of macroscopic electric

fields have yielded threshold estimates above 200 V/m for single monophasic pulses (Aberra et al.,

2020; Weise et al., 2023b).

In this study, we investigated local electric field perturbations when considering neural structures at

the microscopic level in the electric field simulations. This enabled us to bridge the gap between

activation thresholds observed in experiments with respect to macroscopically computed electric

fields and those predicted by detailed microscopic simulations of neurons. Our approach lowers

this threshold estimate to about 30…40 V/m (Fig. 4), which is in the order of EEG experiments (see

above). Also, the corrected half-maximum values around 50 V/m are much closer to the values

seen for experimental data using recorded MEPs together with macroscopic field calculations (e.g.,

60 V/m in Numssen, Kuhnke et al., 2024). Hence, the recruitment rates computed using the more

realistic (i.e., microscopic) electric fields provide a reasonable proxy for the activation thresholds of

hand muscles (and, thus, MEPs). This was not the case when using the macroscopically estimated

field strength as the field at the neuronal membranes (see Fig. 4).

Note, however, that our model describes the recruitment rate of neurons in the motor cortex, which

is not identical to muscle activation reflected by MEP, which involves further downstream

processing. Thus, including these processes (e.g., cortical dynamics, long range axonal

transmission, spinal dynamics, and muscle fiber activation function) is expected to yield an even

more accurate prediction of the MEP.

One key precondition to our approach is the ability to predict microscopic fields at very fine detail

by means of large-scale numerical computations. To this end, we utilized results from a method

using the BEM-FMM to model perturbations of an impressed electric field within a microscopically

realistic brain tissue sample, with many tightly spaced neuronal cells and other structures (Qi et al.,

2025). The obtained results (Fig. 1 and 2) demonstrate strong local field perturbations due to the

presence of membranes. The derived probability density function of the ratio between microscopic

and macroscopic electric fields allowed us to apply a statistical correction to the previously

determined recruitment rates by Weise et al. (2023b), who assumed equality between the

macroscopic field and the local field at the membranes. Critically, this approach entails that the time

consuming numerical field computations have to be performed only once (for a particular type of

tissue) and are then reused in the form of a statistical distribution to correct recruitment rates derived

from simple macroscopic field estimations. Note, however, that the probability density function of

the microscopic/macroscopic field ratio is potentially tissue dependent. Here, we applied an

available sample of mouse visual cortex to the activation threshold in human motor cortex. Although

we assume that the overall trend in microscopic field distortion would be similar in all cortical tissues,

we certainly lose some accuracy. Potentially, cell morphologies, cell alignment (anisotropy), cell

density, and other factors exercise an influence on the scaling factors. For the future, we envision

the construction of a library of different tissue types, for each of which a microscopic field simulation

is performed and a separate probability density function is derived. Such research would also shed

light on the question to what extent this relationship can be actually generalized and how much we

have to take into account local tissue properties.

Another limiting factor of the model refers to the neural density in the sample. In the model of Qi et

al. (2025), the neural density is about 30±5%, which is significantly less than previously reported

values of about 70% (Pilgrim et al., 1982; Lehmenkühler et al., 1993; Mota et al., 2014). This

discrepancy is due to the technical limitation that all neurons with soma outside the sample were

omitted and consequently their dendrites and axons are also missing in the segmentation leading

to a decrease in neural density. This is expected to have an influence on the electric field scaling

distribution. In order to reduce this influence, larger samples in the millimeter scale would be

necessary.

The simulations by Weise et al. (2023b) provide detailed insight into the statistics of the activation

thresholds for various neuronal populations, and how they depend on parameters of the external

electric field, such as the angle of incidence and field gradient. Already these simulations were

complex and computationally expensive, to a degree that impedes usage in whole-head models.

Here, we characterized the microscopic electric field considering the mutual influence of the

neurons in a realistic cubic sample of mouse visual cortex, based on simulations by Qi et al., 2025,

which were even more time-consuming. Theoretically, all simulations carried out by Weise et al.

(2023b) would also have to consider microscopic field effects. However, this would lead to an

exponential increase in the computing time and is currently far from being feasible. It would also

require a much larger tissue sample, because in the one currently utilized, large portions of the

neuronal arbors were cut out, rendering direct threshold simulations biased (see Qi et al., 2025).

The approach presented here elegantly decouples both problems and considers the determination

of the thresholds in the homogeneous field for different cell types and the deviation of microscopic

from macroscopic fields separately. Both approaches are then combined by statistically

incorporating the electric field deviations together with the recruitment curve in eqs. (1) and (2).

Currently, this is the computationally tractable approach to quantify the impact of microscopic

electric field perturbations on the firing thresholds and recruitment curves caused by TMS. It

represents a significant breakthrough by aligning neural modeling with experimental realities for the

first time. Apart from a better understanding of the TMS effect, this may open the door to more

systematic procedures for the design of effective stimulation protocols (Shaner et al., 2023).

A promising extension to the current approach of estimating the thresholds would be the use of

bidomain models (Czerwonky et al. 2023; Fellner et al. 2022) of the neurons in combination with

the microscopic field simulations, instead of the one-dimensional cable equation in conjunction with

a much larger 1 mm^3 MICrONS mouse brain sample (MICrONS Consortium 2021), which includes

considerably better developed axonal arbors and ~75,000 neurite cells. However, the necessary

computing power would be immense and is currently not yet available.

Another interesting question would be whether and to what extent the same effects would apply for

transcranial electric stimulation (TES). As both methods are based on externally caused electric

fields (Peterchev et al., 2012), one could speculate that also with TES the microscopic fields at the

cell membranes are larger than the macroscopically computed ones. However, the different ways

of impressing that field (magnetic induction vs. externally applied voltage) may give rise to

differences, which need to be explored in future computational research.

Data availability statement

The electric field data used in this study is published by Qi et al. (2025) via BossDB (https://bossdb.org/project/makaroff2024). It includes post-processed cell CAD models, microcapillary CAD models, post-processed neuron morphologies, extracellular electric fields and potential distributions. The derived electric field scaling distribution and recruitment rate operators are publicly available in a repository (Weise et al. 2025, https://osf.io/wu4sa/).

Acknowledgements

This study has received support from BMBF grant 01GQ2201 (KW, TRK), NIMH grant R01MH130490 and NIBIB grant R01EB035484 (SNM) and NINDS grant 1R01NS112996 (MB).

Credit Authorship Statement

KW, SNM, and TRK formulated the overarching research goals and aims. SNM computed and curated the electric field data. KW and TRK developed the threshold correction approach. KW and SNM prepared the visualizations. KW, SNM, and TRK acquired the financial support. SNM provided the computing resources for the computationally expensive electric field calculations. KW and TRK wrote the initial draft. SNM contributed to the methods and discussion section and critically reviewed the whole manuscript. ON and MB continued to the analysis and discussion, and critically reviewed the manuscript.

Conflicts of interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

Aberra, A. S., Wang, B., Grill, W. M., & Peterchev, A. V. (2020). Simulation of transcranial magnetic stimulation in head model with morphologically-realistic cortical neurons. Brain stimulation, 13(1), 175-189.

Balanis, Constantine A., Advanced Engineering Electromagnetics, Dec, 8, 2023, Print ISBN:9781394180011 |Online ISBN:9781394180042 |DOI:10.1002/9781394180042.

Caulfield, K. A., Li, X., & George, M. S. (2021). Four electric field modeling methods of dosing prefrontal transcranial magnetic stimulation (TMS): Introducing APEX MT dosimetry. Brain Stimul, 14(4), 1032–1034. https://doi .-org /-10 .-1016 /-j .-brs .-2021 .-06 .-012

Czerwonky DM, Aberra AS, Gomez LJ. (2023-12-16), "A Boundary Element Method of Bidomain Modeling for Predicting Cellular Responses to Electromagnetic Fields", BioRxiv: The Preprint Server for Biology, doi:10.1101/2023.12.15.571917.

Di Lazzaro, V., & Ziemann, U. (2013). The contribution of transcranial magnetic stimulation in the functional evaluation of microcircuits in human motor cortex. Frontiers in neural circuits, 7, 18.

Fellner A, Heshmat A, Werginz P, Rattay F. A finite element method framework to model extracellular neural stimulation. J Neural Eng. 2022 Apr 7;19(2). doi: 10.1088/1741-2552/ac6060.

Galanis, C., Neuhaus, L., Hananeia, N., Turi, Z., Jedlicka, P., & Vlachos, A. (2024). Axon morphology and intrinsic cellular properties determine repetitive transcranial magnetic stimulation threshold for plasticity. *Frontiers in Cellular Neuroscience*, *18*, 1374555.

Jackson, John David, Classical Electrodynamics, Aug, 1998, 832 pages, ISBN: 9780471309321

(Print) Published: 1998 by Wiley, NY

Lehmenkühler A, Sykov´a E, Svoboda J, Zilles K, Nicholson C. Extracellular space parameters in

the rat neocortex and subcortical white matter during postnatal development determined by

diffusion analysis. *Neuroscience* 1993 Jul;55(2): 339–51.

Makaroff, S. N., Qi, Z., Rachh, M., Wartman, W. A., Weise, K., Noetscher, G. M., ... &

Nummenmaa, A. R. (2023). A fast direct solver for surface-based whole-head modeling of

transcranial magnetic stimulation. Scientific reports, 13(1), 18657.

Makarov, S. N., Wartman, W. A., Daneshzand, M., Fujimoto, K., Raij, T., & Nummenmaa, A.

(2020). A software toolkit for TMS electric-field modeling with boundary element fast multipole

method: an efficient MATLAB implementation. *Journal of neural engineering*, *17*(4), 046023.

Makarov, S. N., Noetscher, G. M., Raij, T., & Nummenmaa, A. (2018). A quasi-static boundary

element approach with fast multipole acceleration for high-resolution bioelectromagnetic models.

IEEE Transactions on Biomedical Engineering, 65(12), 2675-2683.

Markram, H., Muller, E., Ramaswamy, S., Reimann, M. W., Abdellah, M., Sanchez, C. A., ... &

Schürmann, F. (2015). Reconstruction and simulation of neocortical microcircuitry. Cell, 163(2),

456-492.

414 MICrONs Consortium. Functional connectomics spanning multiple areas of mouse visual cortex.

415 bioRxiv originally published 2021.07.28.454025/updated 2023.04.19; doi:

416 https://www.biorxiv.org/content/10.1101/2021.07.28.454025v3.

418 MICrONS Consortium. Online: https://www.microns-explorer.org/cortical-mm3

420 Mota B, Herculano-Houzel S. All brains are made of this: a fundamental building block of brain

421 matter with matching neuronal and glial masses. *Front Neuroanat* 2014 Nov 12;8:127.

423 Noetscher, G. M., Tang, D., Nummenmaa, A. R., Bingham, C. S., McIntyre, C. C., & Makaroff, S.

424 N. (2023). Estimations of charge deposition onto convoluted axon surfaces within extracellular

425 electric fields. IEEE Transactions on Biomedical Engineering.

427 Numssen, O., Zier, A. L., Thielscher, A., Hartwigsen, G., Knösche, T. R., & Weise, K. (2021).

428 Efficient high-resolution TMS mapping of the human motor cortex by nonlinear regression.

429 NeuroImage, 245, 118654.

431 Numssen, O., Kuhnke, P., Weise, K., & Hartwigsen, G. (2024). Electric-field-based dosing for

432 TMS. *Imaging Neuroscience*, 2, 1-12.

434 Peterchev, A. V., Wagner, T. A., Miranda, P. C., Nitsche, M. A., Paulus, W., Lisanby, S. H., ... &

435 Bikson, M. (2012). Fundamentals of transcranial electric and magnetic stimulation dose: definition,

436 selection, and reporting practices. *Brain stimulation*, 5(4), 435-453.

438 Pilgrim C, Reisert I, Grab D. Volume densities and specific surfaces of neuronal and glial tissue

439 elements in the rat supraoptic nucleus. *J Comp Neurol* 1982 Nov 10;211 (4):427–31.

Qi, Z., Noetscher, G. M., Miles, A., Weise, K., Knösche, T. R., Cadman, C. R., ... & Makaroff, S. N. (2024). Electromagnetic Modeling within a Microscopically Realistic Brain–Implications for Brain Stimulation. bioRxiv, 2024-04.

Rosanova, M., Casali, A., Bellina, V., Resta, F., Mariotti, M., & Massimini, M. (2009). Natural frequencies of human corticothalamic circuits. J Neurosci, 29(24), 7679–7685.

Rossi, S., Antal, A., Bestmann, S., Bikson, M., Brewer, C., Brockmöller, J., ... & Hallett, M. (2021). Safety and recommendations for TMS use in healthy subjects and patient populations, with updates on training, ethical and regulatory issues: Expert Guidelines. Clinical Neurophysiology, 132(1), 269-306.

Shirinpour, S., Hananeia, N., Rosado, J., Tran, H., Galanis, C., Vlachos, A., ... & Opitz, A. (2021). Multi-scale modeling toolbox for single neuron and subcellular activity under Transcranial Magnetic Stimulation. *Brain stimulation*, *14*(6), 1470-1482.

Stratton, Julius A., Electromagnetic Theory, Wiley-IEEE Press, 2006, 640 p., DOI: 10.1002/9781119134640.

Thielscher, A., Antunes, A., & Saturnino, G. B. (2015). Field modeling for transcranial magnetic stimulation: A useful tool to understand the physiological effects of TMS? Annu Int Conf IEEE Eng Med Biol Soc, 2015, 222–225.

464 Turner, N. L., Macrina, T., Bae, J. A., Yang, R., Wilson, A. M., Schneider-Mizell, C., ... & Seung,

465 H. S. (2022). Reconstruction of neocortex: Organelles, compartments, cells, circuits, and activity.

466 Cell, 185(6), 1082-1100.

468 Wang, B., Peterchev, A. V., Gaugain, G., Ilmoniemi, R. J., Grill, W. M., Bikson, M., & Nikolayev,

469 D. (2024). Quasistatic approximation in neuromodulation. *Journal of neural engineering*, *21*(4),

470 041002.

472 Weise, K., Numssen, O., Thielscher, A., Hartwigsen, G., & Knösche, T. R. (2020). A novel

473 approach to localize cortical TMS effects. Neuroimage, 209, 116486.

475 Weise, K., Numssen, O., Kalloch, B., Zier, A. L., Thielscher, A., Haueisen, J., ... & Knösche, T. R.

476 (2023a). Precise motor mapping with transcranial magnetic stimulation. Nature protocols, 18(2),

477 293-318.

479 Weise, K., Worbs, T., Kalloch, B., Souza, V. H., Jaquier, A. T., Van Geit, W., ... & Knösche, T. R.

480 (2023b). Directional sensitivity of cortical neurons towards TMS-induced electric fields. Imaging

481 Neuroscience, 1, 1-22.

483 Weise, K. (2025, January 29). Statistical method accounts for microscopic electric field distortions

484 around neurons when simulating activation thresholds - Data and Code. Retrieved from

485 osf.io/wu4sa

487 Zmeykina, E., Mittner, M., Paulus, W., & Turi, Z. (2020). Weak rTMS--induced electric fields produce

488 neural entrainment in humans. Sci Rep, 10(1), 11994.

Figures

Fig. 1: Distribution of the extracellular electric field magnitude (V/m) inside the sample when a uniform impressed electric field of $E_x^{(p)} = 100\,\mathrm{V/m}$ is applied in the medial-lateral direction (from left to right in the (a)), showing how low conducting barriers (e.g., cell membranes) cause charge accumulation and associated field distortion in (b) and (c). The intracellular space (grey) is not included in the computations. Tissue segmentation and electric field computations are from Qi et al. (2025).

Fig. 2. The influence of the accumulation of charges on the membranes onto the collinear electric field at the centerlines of neuronal processes for an example neuron. Top: A uniform impressed electric field is applied along the *x*-axis (from dorsal to ventral) with $E_x^{(p)} = 100 \, \text{V/m}$. The influence of the charges is not considered and the impressed field is directly projected to the centerlines. Hence, the maximum achievable collinear field (for an optimally oriented neuronal segment) at the neurons $E_{max}^{(c)}$ is equal to the impressed field. Bottom: The same uniform impressed field is applied, but realistic neuronal (sub-)compartments are included into the voltage simulation and the impressed field is distorted by the field of the induced charges. In consequence, the maximum achievable collinear field $E_{max}^{(c)}$ can be larger than the impressed field $E_x^{(p)}$.

Fig. 3: Histogram and probability density $p(s_e)$ of the electric field scaling factor $\left(s_e = \frac{E_{micro}}{E_{macro}}\right)$ between

microscopic and macroscopic electric fields (black dashed line). The median of the scaling factor is $\bar{s}_e = 1.06$

(red dashed line).

Fig. 4: Uncorrected and corrected recruitment rates in comparison to experimental MEPs. Dashed black line:

original recruitment rates from Weise et al. (2023b) of L2/3 PC for $\theta = 0°$ and $\Delta|\tilde{\mathbf{E}}| = 0$ %/mm considering

homogeneous macroscopic electric fields and a biphasic TMS pulse, without taking into account microscopic

electric field effects. Red line: corrected recruitment rates determined from eqs. (1) and (2) using the

probability density $p(s_e)$ of the electric field scaling factor between microscopic and macroscopic electric fields

from Fig. 2, together with the recruitment rate determined using macroscopic electric fields from Weise et al.

(2023b) assuming an average number of axon terminals of N=35. Grey lines: Recruitment rate curves after

sampling the number of axon terminals N from a normal distribution with the given mean of N=35 and standard

deviation of 13.7. Colored dots: MEPs as function of the external macroscopic electric field determined

experimentally in Numssen et al. (2021) after motor mapping (different colors represent different subjects).

Fig. 5: Recruitment rates of L2/3 PC without (a) and with (b) microscopic field corrections stimulated by biphasic TMS pulses for different electric field angles. No e-field gradient ($\Delta|\tilde{\boldsymbol{E}}| = 0\,\%/\mathrm{mm}$) was assumed. (a) The neuronal recruitment rate from Weise et al. (2023b) did not consider microscopic electric field variations, yielding e-field thresholds of above 200 V/m. (b) Recruitment rate of a neuronal population taking electric field variations from Fig. 3 into account yields thresholds of below 50 V/m. (c) and (d) Recruitment rates for field angles of $\theta = 0°$, $\theta = 90°$, and $\theta = 180°$ without and with microscopic field correction, respectively.