Title: **Syntaxin11 Deficiency Inhibits CRAC Channel Priming To Suppress**
Cytotoxicity And Gene Expression In FHLH4 Patient T Lymphocytes.

Sritama Datta[1], Abhikarsh Gupta[1#], Kunal Mukesh Jagetiya[1#], Vikas Tiwari[2⊥], Megumi
Yamashita[3⊥], Sandra Ammann[4,5], Mohammad Shahrooei[6], Atharva Rahul Yande[1],
Ramanathan Sowdhamini[2], Adish Dani[1], Murali Prakriya[3], Monika Vig[1]*

Address and Affiliation: [1]Tata Institute of Fundamental Research, Hyderabad, India.
[2]National Centre for Biological Sciences, Bangalore, India. [3]Northwestern University,
Feinberg School of Medicine, Chicago, USA. [4]Institute for Immunodeficiency, Center for
Chronic Immunodeficiency, Medical center, University of Freiburg, Faculty of Medicine,
University of Freiburg, Freiburg, Germany. [5]Institute for Transfusion Medicine and Gene
Therapy, Medical Center, University of Freiburg, Faculty of Medicine, University of
Freiburg, Freiburg, Germany. [6]Department of Microbiology, Immunology and
Transplantation, Clinical and Diagnostic Immunology, KU Leuven, Leuven, Belgium.

[#, ⊥] These authors contributed equally.

***Correspondence to:** monika.vig@gmail.com

26 **Keywords:** Orai, Stim, SNAP, SNARE, CRAC, Syntaxin11, ion channels, SOCE, FHLH4,
27 autoimmunity, cytotoxicity, CTL, T lymphocytes

Abstract:

CRAC channels enable calcium entry from the extracellular space in response to a variety of stimuli and are crucial for gene expression and granule exocytosis in lymphocytes. Here we find that Syntaxin11, a Q-SNARE, associated with FHLH4 disease in human patients, directly binds Orai1, the pore forming subunit of CRAC channels. Syntaxin11 depletion strongly inhibited SOCE, CRAC currents, IL-2 expression and cytotoxicity in cell lines and FHLH4 patient T lymphocytes. Constitutively active H134 Orai1 mutant completely reconstituted calcium entry in Syntaxin11 depleted cells and the defects of granule exocytosis as well as gene expression could be bypassed by ionomycin induced calcium influx in FHLH4 T lymphocytes. Our data reveal a Syntaxin11 induced pre-activation state of Orai which is necessary for its subsequent coupling and gating by the endoplasmic reticulum resident Stim protein. We propose that ion channel regulation by specific SNAREs is a primary and conserved function which may have preceded their role in vesicle fusion.

Introduction:

Most eukaryotic cells have a limited amount of calcium sequestered inside intracellular stores, the largest of which are the endoplasmic reticulum (ER). Signaling from cell surface receptors induces the release of stored calcium which, in turn, activates store-operated calcium entry (SOCE) to replenish the stores and sustain signaling and other calcium dependent cellular processes [1]. Calcium release activated calcium (CRAC) channels play a major role in mediating SOCE [2]. In lymphocytes and mast cells, CRAC currents have been shown to be crucial for granule exocytosis as well as gene expression associated with effector functions [3] [4] [5]. Orai (CRACM) multimers form the pore of CRAC channels [6] and ER-resident Stim proteins sense store-depletion and subsequently trap and gate freely diffusing resting Orai in ER-PM junctions [7]. CRAC channels are therefore thought to be dependent on Stim proteins for any structural transitions that result in the activation of CRAC currents [8].

Genome-wide RNAi screens, which initially identified Stim and Orai [9-13], have yielded a wealth of information regarding additional players in SOCE. For instance, we have previously shown that alpha-soluble N-ethylmaleimide sensitive factor (NSF) attachment protein (α-SNAP), a well-known synaptic family adaptor protein that forms a part of the 20S SNARE super-complex[14, 15], is a crucial component of the CRAC channel supramolecular complex [12, 16]. α-SNAP independently bound Stim as well as Orai, with high affinity [17-19]. Single particle diffusion analysis showed that store-depletion induced arrest of Orai mobility was completely reversed in α-SNAP depleted cells, suggesting non-functional co-clustering with Stim proteins [17]. Step-photobleaching analysis of Orai1 further showed altered stoichiometry and ion selectivity of Orai1 multimers in α-SNAP depleted cells demonstrating that α-SNAP is required for the on-site assembly and ion selectivity of CRAC channels [17]. Therefore, analysis of the role of α-SNAP revealed that Stim:Orai coupling is necessary but not sufficient for SOCE and the molecular process likely involves multiple additional steps.

A key question related to the above described findings is whether and how do the

additional molecular players of the synaptic machinery regulate SOCE? Our previous

analyses have shown that alpha-SNAP functions independently of its usual binding

partner NSF-ATPase in the regulation of SOCE [16]. However, this question remains

incompletely addressed and becomes even more pertinent because specific SNAREs

appeared in two independent genome-wide RNAi screens previously conducted in

Drosophila cells to identify regulators of SOCE but were not characterized [11, 12].

Additionally, associations of SNAREs with pore-forming subunits of a variety of channels

have been reported previously, however, several of these interactions were reported to

be inhibitory [20] [21]. In others, SNAREs were initially thought to be required for the insertion

of pore subunits into the target membrane [22-24]. Several earlier reports documenting the

direct association of VGCCs with t-SNAREs have proposed that interactions with synaptic

machinery serve to localize the source of calcium influx close to the sites of secretory and

synaptic vesicle fusion. However, this hypothesis runs into following problems. First,

SNAREs also associate with the pore subunits of potassium (Kv2.1), chloride (CFTR) and

sodium channels, neither of which conduct calcium to facilitate membrane fusion [25-28].

Second, recent studies have identified several other adaptor proteins such as calcium-

calmodulin dependent serine kinase (CASK) and Mint-1, which directly bind Cav channels

and could serve the function of anchoring at desired locations [22]. Finally, synaptic proteins

should not have to bind close to the pore forming region, in order to perform the task of

anchoring the channels. Therefore, the exact role of these associations remains

unestablished, but a common theme that emerges is that none of them directly modulate

the surface expression of the candidate channels [16, 25, 26, 29, 30]. Furthermore, SNAREs are

highly expressed proteins in most cells [23], yet only three assembled SNARE complexes

are needed for the fast fusion exocytosis in chromaffin cells [31] and two synaptobrevin

molecules are sufficient for vesicle fusion in hippocampal neurons [32].

It is therefore necessary to envisage an alternate, and more fundamental, function

for the association of SNAREs with the ion channel pore subunits. We hypothesized that

additional synaptic proteins are involved in the direct regulation of SOCE. However,

because the repertoire of synaptic family proteins is much larger in mammalian cells, we

conducted a limited RNAi screen in HEK293 and Jurkat T cells, where we initially targeted

those synaptic family genes that are expressed in both cells. STX11 specifically regulated

SOCE, gene expression and cytolysis by directly binding to Orai and inducing a hitherto

unsuspected, early molecular transition which prepared Orai for subsequent gating by

Stim. Further, defective SOCE drove the granule exocytosis defects of STX11 deficient

FHLH4 patient T lymphocytes, which could be bypassed with ionomycin.

Results:

We have previously shown that α-SNAP, a ubiquitously expressed synaptic family protein, forms an integral component of the functional CRAC channel complex and binds Orai1 and Stim1 with high affinity [12, 16 17-19]. NSF-ATPase, a usual binding partner of α-SNAP, was not involved in regulating SOCE [16]. However, α-SNAP also binds the cis-SNARE complex and a Q-SNARE, Syntaxin 5 (STX5), showed inhibition of SOCE in one of the earlier genome-wide screen performed in *Drosophila* S2 cells [11]. Furthermore, STX1 has been previously shown to bind a variety of ion channels [22-24]. Therefore, to assess the potential role of Syntaxins in SOCE, we first knocked down STX5 (Supplementary Figure 1A) and STX1A (Supplementary Figure 1B) using five different sequences of shRNA targeting each gene in HEK293 cells and measured SOCE in response to Thapsigargin (TG). We did not see a significant defect in SOCE in either of the cases. However, several members of the synaptic family proteins are not ubiquitously expressed and many show redundancy in their function. Therefore, we searched online databases and initially only targeted synaptic family protein genes that are expressed in HEK293 and T cells (Supplementary Figure 1C-N). We found that Syntaxin11 knockdown showed strong inhibition of SOCE in a variety of cell lines (Supplementary Figure 1D) (Figure 1A-F). To establish specificity of the knockdown, we expressed STX11 in STX11 depleted cells, which largely restored SOCE (Figure 1G). In addition, we knocked down STX11 using an additional shRNA sequence, which also significantly inhibited SOCE in Jurkat T cells (Supplementary Figure 2). To assess the extent of STX11 mRNA depletion, we extracted total RNA from scramble (scr) and STX11 shRNA treated HEK293 cells and subjected it to quantitative PCR analysis using Taqman probes (Supplementary Figure 3). We observed nearly 70% depletion of STX11 mRNA. To determine whether ectopic expression of STX11 would enhance SOCE, we expressed STX11 in HEK293 (Figure 1H) and Jurkat T cells (Supplementary Figure 4) and measured SOCE. There was a significant increase in both the cases reinforcing a crucial role of STX11 in SOCE. To determine whether the increase in SOCE seen upon the expression of STX11 was due to CRAC channels, we depleted Orai1 expression before expressing STX11 in HEK293

cells. The STX11 mediated enhancement observed in SOCE was lost in Orai1 depleted
cells (Figure 1I).

Previous studies have shown that SOCE in T cells is mediated by CRAC channels
formed by Orai proteins. To test whether CRAC current (I_{CRAC}) is affected by knockdown
of STX11 in T cells, we performed whole cell patch clamp recordings on scramble (scr)
and STX11 shRNA treated Jurkat T cells (Figure 2A-C). Leak-subtracted I_{CRAC} was
recorded in 20 mM Ca^{2+} and a Na^+-based divalent cation free (DVF) solution. These
recordings showed that both the Ca^{2+} and DVF CRAC current is decreased more than
two-fold in the STX11 shRNA treated cells with the normalized currents shown in Figure
2C. The electrophysiological properties of the residual current in STX11 shRNA treated
cells was indistinguishable from I_{CRAC} in scramble shRNA treated cells in terms of
blockade of the Ca^{2+} current by La^{3+}, depotentiation of the DVF current over tens of
seconds, and fast inactivation of Ca^{2+} current. Moreover, the reversal potential of the
$Ca^{2+}$and DVF currents were similar between scramble and STX11 shRNA treated Jurkat
T cells, suggesting that STX11 depletion does not affect the calcium selectivity of CRAC
channels (Figure 2D). These results indicate that knockdown of STX11 significantly
decreases I_{CRAC} in Jurkat T cells.

CRAC channel mediated calcium influx is crucial for sustained calcium signaling
which, in turn, is required for the activation and nuclear translocation of specific
transcription factors such as nuclear factor of activated T (NFAT) [2] [33]. We stimulated
control and STX11 depleted Jurkat T cells with thapsigargin (TG) and phorbol myristate
acetate (PMA), prepared nuclear and cytosolic extracts and subjected them to Western
blot using anti-NFAT antibody (Figure 2E). The nuclear translocation of NFAT was
severely compromised in STX11 depleted T cells. To visualize NFAT translocation in
response to T cell receptor mediated stimulation, we incubated control and STX11
depleted Jurkat T cells with plate-coated anti-CD3 and soluble anti-CD28, fixed,
permeabilized and immunolabelled for NFAT. Representative images and quantification
of the nuclear NFAT fraction (Figure 2F, 2G) across multiple randomly chosen fields
shows that STX11 depleted cells are deficient in NFAT nuclear translocation upon anti-

CD3 stimulation. The defect of NFAT translocation also affects gene expression, as
assessed by IL-2 specific QPCR of control and STX11 depleted Jurkat cells upon anti-
CD3 stimulation (Figure 2H). Taken together, these data demonstrate an essential role
for STX11 in SOCE *via* CRAC channels, NFAT activation and gene expression.

STX11 is a Q-SNARE protein and most Q-SNAREs possess a C-terminal
transmembrane domain for membrane insertion [34]. However, STX11 is atypical in this
regard because it harbors a group of cysteine residues close to the C-terminus in place
of the transmembrane domain (Supplementary Figure 5). The cellular localization of
STX11 is not established [35] [36] [37]. This is primarily because using the available commercial
antibodies, native STX11 is undetectable in most cell lines and tagging STX11 on either
the N- or the C-terminus results in partial degradation and mis-localization of the protein
(data not shown) [35]. Therefore, to determine its localization in HEK293 cells, we inserted
an HA tag inside the N-terminal unstructured region of STX11, just before the beginning
of the Habc domain, expressed in HEK cells, permeabilized and labelled with anti-HA tag
antibody. STX11 with an internal HA tag localized in the plasma membrane (Figure 3A).
We next expressed untagged STX11 in cells co-expressing Orai1-YFP, permeabilized
and stained with anti-STX11 antibody. In majority of cells, STX11 co-localized with Orai1
in the plasma membrane (Figure 3B). Co-localization of STX11 and Orai1 suggested a
direct role in SOCE. To test this, we co-expressed STX11 with either Flag- or Myc-tagged
Orai1 in HEK293, prepared whole cell lysates and subjected them to co-
immunoprecipitation followed by Western Blot using either anti-Flag, anti-Myc or anti-
STX11 antibodies. We found that both Orai1 and STX11 could co-immunoprecipitate
each other under resting as well as store-depleted conditions (Figure 3C, 3D). To further
assess whether STX11 directly binds Orai1, we expressed and purified from *E. Coli*,
MBP-tagged N- and C-terminal cytosolic tails of Orai1 (Figure 3E). *In vitro* pull-down
assay performed by incubating Orai1 cytoplasmic domains with full length soluble His-
tagged STX11 showed that both the N-terminus as well as the C-terminus of Orai1 bound
STX11 albeit with different apparent affinities (Figure 3F). In both cases, the binding of
STX11 to Orai1 was similar or higher when compared to SNAP23, a known binding
partner of STX11. SNARE proteins typically utilize their SNARE domain for interacting

with other SNAREs [34]. To identify the domain of STX11 involved in the regulation of

SOCE, we expressed His-tagged H$_{abc}$ and SNARE domains of STX11 in *E. Coli*, purified

and assessed their binding to the MBP-tagged Orai1 tails *via* a similar pull-down assay.

We found that the H$_{abc}$ domain of STX11 showed significantly high binding to the C-

terminus of Orai1 but faint binding to the N-terminus was also detected (Figure 3G). The

SNARE domain did not show any binding to the Orai1 tails but showed faint binding to

SNAP23 (Supplementary Figure 6). We analyzed the ability of full length Orai1 and

STX11 to form a complex using AlphaFold3 (AF3), however, the scores were insignificant

(ipTM = 0.11 and pTM = 0.35). Therefore, we examined domain-domain interactions

STX11 (H$_{abc}$, SNARE) and Orai1 (N-terminus, C-terminus). The complex of STX11 H$_{abc}$

with Orai1 C-terminus resulted in a significantly high prediction score (Supplementary

Table 1). The AF3 predictions can vary with different seeds. Therefore, we executed AF3

predictions with different initial seeds to generate multiple models of STX11 H$_{abc}$ with

Orai1 C-terminus. The contact frequency of interface residues was calculated in different

models. Multiple residues of STX11 H$_{abc}$ domain and Orai1 C-terminus domain were

observed to have contact frequency of more than 0.8 (Supplementary Figure 7). The best

scoring model among all predicted models was considered for further analysis

(Supplementary Table 2).

Next, we performed all atom MD simulation in aqueous environment to assess the

interaction stability of the STX11 H$_{abc}$ and Orai1 C-terminus complex. The STX11 H$_{abc}$

remained conformationally stable throughout the simulation time as assessed through

RMSD (Supplementary Figure 8). The binding energy (Supplementary Figure 9) and the

interaction between the two subunits remained largely stable throughout the simulation

period (Movies 1-3). Major interactions observed include salt bridge between

Arg78_STX11 and Glu275_Orai1, Glu150_STX11 and Arg289_Orai1, Arg160_STX11

and Glu272_Orai1, H-bond between Asn147_STX11 and Arg289_Orai1 and

Gln164_STX11 and Glu272_Orai1. A potential cation-pi interaction between

Tyr146_STX11 and Arg289-Orai1 was also observed in one of the replicates

(Supplementary Figure 10).

The individual trajectories were concatenated and clustered to obtain a centroid
structure for this complex. There were 46 clusters, and the largest cluster had 52
members. The cluster representative of the largest cluster showed the elaborate protein-
protein interface, and the Orai1 C-terminus was found to be oriented in an anti-parallel
orientation to the STX11 H$_{abc}$ (Figure 3H, 3I) and we observed 4 salt bridge and 13 H-
bond interactions (Figure 3J). Overall, the docking and simulation analysis corroborated
the biochemical findings and demonstrated a direct role of STX11 in the regulation of
Orai1.

Upon store depletion Stim is known to localize to ER-PM junctional regions where
it traps and co-clusters with Orai1. To further understand the mechanism of action of
STX11 in SOCE, we first assessed whether the localization of Orai and Stim was normal
in control and STX11 depleted cells. We imaged HEK293 cells expressing CFP-Orai1,
Stim1-YFP under resting or store depleted conditions, comparing the levels of Orai1,
Stim1 in the PM or in ER-PM junctions (also termed puncta) across control or STX11
depleted cells. Although the distribution of resting and store-depleted Stim1-YFP
appeared normal (Figure 4A, 4B), CFP-Orai1 appeared partly diffuse in STX11 deficient
store-depleted cells (Figure 4B). Because Stim clusters Orai, we first quantified Stim1-
YFP intensity within ER-PM puncta of store-depleted HEK293 cells. We found no defect
in the ability of Stim1 to localize and cluster in the ER-PM junctions upon store-depletion
(Figure 4C). In line with these findings, measurement of the ER calcium content by
stimulating control and STX11 depleted cells with Ionomycin showed no significant
change (Supplementary Figure 11) and staining of ER and Golgi with organelle specific
markers also showed no abnormalities (Supplementary Figures 12A and 12B) suggesting
that the overall health and calcium content of ER and Golgi were not adversely affected
by STX11 depletion. However, quantification of Orai1 intensity revealed that the fraction
of Orai1 inside Stim:Orai puncta showed a consistent decrease (Figure 4D) while fraction
of Orai1 outside puncta was found to be higher (Figure 4E). To assess whether the level
of Orai1 expression in the plasma membrane was normal, we labelled a U2OS cell line
stably expressing Orai1 tagged with a Bungarotoxin binding site (BBS) in the second
extracellular loop and YFP at C-terminus (Orai-BBS-YFP) with Alexa-647 conjugated

bungarotoxin (BTX-A647) (Supplementary Figures 13) [16]. We found no change in the

expression levels of total Orai1 in the PM of STX11 depleted cells (Figure 4F). Large

fluorescent tags can sometimes cause steric hindrance in certain locations. Therefore,

we reversed the direction of fluorescent tags on Orai1 as well as Stim1 and repeated the

experiment described above. Expression of Orai1-CFP and YFP-Stim1 could overcome

the apparent defect in the entrapment of Orai by Stim in STX11 depleted HEK293 cells

(Figure 4G-J). However, SOCE was still significantly inhibited in Orai1-CFP and YFP-

Stim1 expressing HEK293 cells (Figure 4K). These data demonstrate that STX11 binding

to resting Orai1 has implications beyond mere co-entrapment of Stim1 and Orai1.

Collectively, these data suggest that STX11 regulates functional entrapment as well as

subsequent gating of Orai by Stim.

To further establish that STX11 depletion mediated suppression of SOCE

encompasses a defect in Orai gating, we expressed Orai1 tethered to two Stim-Orai

activating regions (SOAR) of Stim1 [38] and eGFP (Orai1-S-S-GFP) (Supplementary Figure

14) in HEK293 cells. Orai1-S-S-GFP has previously been shown to constitutively activate

Orai1 [39]. We found that in STX11-depleted, Orai1-S-S-GFP expressing cells, constitutive

calcium entry was significantly smaller in magnitude (Figure 5A-B). To determine whether

reduced calcium entry resulted from a defect in the ability of SOAR to independently bind

Orai1, we next expressed YFP-tagged soluble CRAC activation domain (CAD) of Stim1,

YFP-CAD, in Orai1-CFP expressing stable HEK293 cell line. YFP-CAD localizes to the

cytosol in HEK293 cells but in cells overexpressing Orai1, a significant majority of it

localizes to the plasma membrane due to its association with Orai1 [40]. We quantified the

amount of YFP-CAD localized to the PM as a fraction of total YFP-CAD expressed in

each cell and found no defect in its ability to localize to the plasma membrane in STX11

depleted Orai1-CFP expressing cells (Figure 5C-D). However, in line with our

observations with Orai1-S-S-GFP, constitutive calcium entry was significantly reduced

(Figure 5E-G). Finally, we expressed the constitutively active H134S mutant of Orai1 in

HEK293 cells and measured calcium influx. H134S Orai1 harbors an open pore and the

C-terminal cytosolic tails of Orai1 are unlatched, straightened and pointing towards the

cytosol [41, 42]. Remarkably, expression of H134S Orai1 completely restored constitutive

calcium influx in STX11 depleted cells (Figure 5H, 5I). Taken together, these data suggest

that by binding to the resting Orai1 cytosolic tails, STX11 facilitates a structural transition,

such as correct alignment of Orai1 C-termini, which prepares Orai for gating by Stim.

Therefore, our studies have unraveled a SNARE dependent priming step in the process

of Orai1 gating, which cannot be compensated by merely increasing the density of Stim

proteins bound to Orai.

Human patients with mutations in STX11 develop a rare but fatal autoimmune

disease known as familial hemophagocytic lymphohistiocytosis type 4 (FHLH4) [43]. The

primary cause of this life-threatening disease is a defect in the cytolytic activity of T

lymphocytes and NK cells, which renders the patients susceptible to recurrent infections.

Patients suffer from high fever and severe lymphopenia in early infancy and succumb to

the disease by adolescence unless given bone marrow transplants. Given that STX11

deficient T cells exhibit a strong defect in SOCE which is known to be crucial for

degranulation [3] [4] [5], we hypothesized that reduced SOCE is the primary cause of

cytotoxicity defects in FHL4 patient T cells. To test our hypothesis, we isolated PBMCs

from a 4 year old FHLH4 patient with a homozygous deletion frameshift mutation in the

STX11 coding sequence, c.752delA:p.Lys251fs. The patient presented with typical

symptoms of FHLH4 disease. Given the limited amount of sample available from the

patient, we periodically stimulated patient and healthy donor PBMCs with PHA and

cultured them in IL-2 to expand their numbers, prior to analysis *in vitro*. Isolation and

Sanger sequencing of the patient PBMC DNA showed deletion of a single Adenine at the

752nd position in the STX11 gene (Figure 6A), which would lead to frameshift as well as

elongation of the transcript, resulting in altered protein sequence following Lysine 251.

Supplementary Figure 15 shows the schematic of the wildtype and predicted mutant

STX11 proteins, with the estimated molecular weight of mutant STX11 predicted to be

39.5 KDa. To determine whether the elongated mutant STX11 protein was expressed,

whole cell lysates prepared from the FHLH4 patient and healthy donor PBMCs were

subjected to SDS-PAGE and Western Blot. While the wildtype STX11 band (~33 KDa)

was absent in the patient lysate, we observed a faint but distinct band running higher, at

around 37 KDa, in the FHLH4 sample (Figure 6B). Therefore, the observed and predicted

molecular weight of the FHLH4 mutant protein was higher but its expression was
significantly reduced compared to the wildtype STX11 likely due to protein instability and
degradation. In accordance with our findings in the STX11 depleted Jurkat T cells, SOCE
was found to be significantly defective in the FHLH4 patient T cells when compared to
healthy donor T cells (Figure 6C). Further, expression of wildtype STX11 in FHLH4 T cells
reversed the SOCE defect (Figure 6D), conclusively ruling out any additional
abnormalities in patient T cells and like in the case of Jurkat and HEK293 cells (Figure
1H and Supplementary Figure 4), expression of STX11 in wildtype PBMCs resulted in a
significant increase in SOCE (Figure 6E-F). We next performed granule release assay on
the *in vitro* cultured control and FHLH4 patient CD8 T cells in response to receptor (anti-
CD3+anti-CD28) mediated stimulation (Figure 6G, 6H). We observed a ~50% defect in
the FHLH4 patient CD8 T cell degranulation. To determine whether the reduced granule
release results from a direct defect in vesicle fusion or SOCE, we stimulated the cells with
Ionomycin+PMA (Figure 6G, 6H). Remarkably, cytolytic activity was fully restored in
FHLH4 patient T cells demonstrating that defective SOCE largely causes the cytolytic
defects in FHLH4 patient T cells. Further, we stimulated WT and FHLH4 PBMCs either
through the receptor (anti-CD3+anti-CD28) or using Ionomycin+PMA, extracted total
RNA and performed Q-PCR to assess Interleukin-2 (IL-2) expression. As seen previously
with Jurkat T cells (Figure 2), IL-2 expression was significantly defective in FHLH4 patient
T cells but could be largely restored with Ionomycin (Figure 6I). Taken together, these
data show that reduced SOCE primarily causes the cytotoxicity and gene expression
defects in FHLH4 patient T cells resulting in immune dysregulation. These defects likely
together initiate the pathogenesis of the complex FHLH4 disease in human patients.
393
Finally, Q-SNAREs, such as syntaxins, are often found in complex with other Q-
SNAREs such as SNAP23/25/29 in the target membranes. A number of previous studies
have implicated the complex of SNAP23/25 and STX1a in the regulation of a variety of
ion channels [26, 29]. The three SNAPs, SNAP23/SNAP25/SNAP29 are each capable of
contributing two SNARE domains to the *trans*- and *cis*-SNARE complex although they
themselves lack a transmembrane domain and are attached to the target membrane *via*
palmitoylation of specific residues. Depletion of SNAP23/SNAP25/SNAP29 did not show

a significant reduction in SOCE (Supplementary Figure 1E-1G). To determine whether SNAP23/SNAP25/SNAP29 might still form a part of the STX11:Orai complex to regulate SOCE, we co-expressed flag-tagged Orai with myc-tagged respective SNAPs and performed co-immunoprecipitations. We did not find any interaction between the three SNAPs and Orai1 (Supplementary Figure 16). Similar studies were performed by co-expressing YFP-Stim1 and the respective SNAPs. Again, no interaction was found between the three SNAPs and Stim1 (Supplementary Figure 17). These data show that, unlike in the case of *trans-* and *cis-*SNARE complexes, STX11 does not collaborate with SNAP23/SNAP25/SNAP29 in the regulation of SOCE. Figure 7 proposes a model to summarize the interaction of Orai1 with STX11 which is necessary for and precedes store-operated gating of Orai1 channels.

Discussion:

CRAC channels conduct a small but highly calcium specific current in response to store-depletion. According to the prevailing view, Orai proteins, which reside in the plasma membrane, depend on the ER resident store sensor Stims for the structural transitions leading up to their activation[7]. Unlike other ion channels, the process of activation of CRAC currents is exceptionally slow. For instance, following break-in or store-depletion, it takes, on average, ~5 minutes for Stims to cluster in the ER-PM junctions and for measurable CRAC currents to flow. It is believed that during this time, all the action happens at the level of Stim proteins, which undergo intramolecular transitions and slowly trap and gate freely diffusing Orai. This makes Orai proteins a completely passive player in SOCE.

We have shown that the cytosolic tails of Orai1 constitutively couple with STX11 in resting cells (Figure 3). Interestingly, Orai cytosolic tails have been shown to be crucial for gating but are missing from the structures of closed and open *Drosophila* Orai [41, 44]. The structure of closed Orai suggested that C-terminal tails of two adjacent Orai subunits, bend, pair with each other in an antiparallel fashion and sit closely apposed to the plasma membrane. However, in the structures of the constitutively active H134 mutant Orai, the C-terminal tails are found to orient away from the plasma membrane [41, 44]. Unlike wildtype Orai1, the H134S Orai1 mutant was insensitive to STX11 depletion. Therefore, it is reasonable to propose that STX11 binding to resting Orai1 facilitates a structural transition which enables Stim dependent gating of Orai in ER-PM junctions. Such molecular transition should logically precede the association of Orai with Stims because the STX11 and Stim1 interacting regions appear to be overlapping in Orai1. Of note, STX11 did not co-localize with Orai and Stim in ER-PM junctions (Supplementary Figure 18). In line with those findings, STX11 also did not appreciably co-immunoprecipiate with Stim1 (Supplementary Figure 19). Therefore, our data suggest that STX11 comes off Orai after having induced the proposed structural change. In a different scenario, change in either the structure of STX11 itself or antibody accessibility/ affinity could have precluded detection of STX11 in Stim:Orai clusters.

FHLH is a heterogeneous autoimmune disorder where patients suffer from
hyperinflammation and severe dysregulation of the immune system. Defects in the CTL
and NK cell cytotoxicity machinery underlie the development of this complex disease.
Several genes such as *STX11*, *UNC13D*, *Prf1*, *Rab27*, *Munc18-2* and sometimes *XIAP*
and *ITK* are broadly grouped together based on disease symptoms, although the age of
onset and severity varies across the spectrum [45] [46] [47]. Most mutations in the *STX11* coding
region result in a complete loss or a severe depletion of the protein levels [48]. The disease,
therefore, results from reduced STX11 protein in most FHLH4 patients. Of note, we saw
a ~50% defect in CTL degranulation and a ~70% defect in IL-2 expression even though
the patient exhibited all the typical symptoms of FHLH4 by 4 years of age. A partial defect
in degranulation is not surprising given that *in vitro* culture of FHLH patient CTLs in the
presence of IL-2 has been previously shown to overcome the cytolytic defect [49]. In those
studies, IL-2 dependent induction and compensation by STX3 was proposed to cause the
reversal of the granule release defect *in vitro* [49]. Therefore, it is likely that an analogous
mechanism partially rescued the degranulation defect in the *in vitro* cultured FHLH4
patient PBMCs in our study. Another possibility is that the residual STX11 mutant protein
retained function in FHLH4 T lymphocytes (Figure 6B). Nevertheless, in addition to the
cytolytic defects, for the first time, we have shown that STX11 deficient T cells harbor a
severe defect in SOCE, downstream signaling and gene expression. Therefore, STX11
deficiency affects T cell activation and FHLH4 disease progression at multiple levels. We
propose that the design and early administration of CRAC channel agonists could
potentially provide an alternative to bone marrow transplants for FHLH4 patients.

In summary, we have shown that binding to a Q-SNARE, STX11, provides an on-
site molecular switch necessary for a previously unsuspected crucial priming of the CRAC
channel pore, Orai. This, SNARE enabled, structural transition is necessary for the gating
of CRAC channels. We argue that ion channel regulation is a novel, direct and primary
role of specific Q-SNAREs, such as STX11. In accordance with this hypothesis, SNAREs
are now thought to have evolved from a common archaeal precursor found to be present
in the genomes of Asgard and members of *Legionella* [50]. Both organisms lack an
endomembrane system but do express ion channels and transporters.

Materials and Methods

Plasmid constructs and transfection

Orai1-Myc, Flag-Orai1, Stim1-Myc, Orai1-CFP, Orai1-YFP, CFP-Orai1, YFP-Orai1, YFP-Stim1, Orai-BBS-YFP, Orai1-S-S-EGFP, CFP-Stim1, Stim1-CFP, YFP-Stim1 and Stim1-YFP plasmids have been described previously [17] [19]. Stim1 CAD was sub-cloned in eYFPC1 vector after amplification from YFP-Stim1 plasmid to generate YFP-CAD. pLKO.1, psPAX2 and pMD2.G were purchased from Addgene. pLKO.1 cloned shRNA sequences targeting the genes of interest, were either purchased from Horizon Discovery, UK or designed and cloned in-house using the Broad Institute portal (https://portals.broadinstitute.org/gpp/public/gene/search). Full-length human STX11 was cloned from human cDNA prepared from HEK293 cells and subcloned into pcDNA3.1(+), pcDNA/4TO/Myc-HisA (Invitrogen, Grand Island, NY), pMSCV-IRES-mcherry (Addgene), pEF1alpha-IRES (Clontech) and pET28b vector [19] with an N-terminal 6XHis-tag. The fragments of STX11 were cloned in pET28b vector with an N-terminal 6xHis tag for expression in *E. Coli*. HA tag was inserted between H30 and G31 in Stx11 cloned in pMSCV-IRES-mCherry construct using PCR. Point mutants of Orai1 (H134S) were generated using site directed mutagenesis. The fragments of Orai1 (1-87, 1-47, 48-103, 256-301 and 272-292) were amplified from full length constructs and cloned into pMAL-c5X vector (New England Biolabs), in-frame with MBP protein coding sequence, as described previously [19]. SNAP23, SNAP25 and SNAP29 cDNA cloned in pCMV-Sport6.1 vectors were purchased from Dharmacon and subcloned into pcDNA4/TOMycHisA. SNAP23 was subcloned in pMAL-c5X vector in-frame with MBP. All plasmid DNA transfections in human cell lines and primary cells were done using Lipofectamine 2000 (Invitrogen)/ Lipofectamine 3000 (Invitrogen) or Amaxa nucleofection kit (Lonza, Basel, Switzerland) respectively, as per manufacturer's protocol.

Cell Lines

Lentiviral shRNA transduction experiments were performed in HEK 293, U2OS or Jurkat (ATCC, Manassas, VA) cell lines cultured in low glucose DMEM (Hyclone, Logan, UT) or RPMI (Hyclone) respectively, with 5-10% fetal bovine serum (Hyclone), 1X Penicillin

Streptomycin, and GlutaMax (Gibco, Grand Island, NY). Stable cell lines generated using

HEK293 and U2OS parent lines have been described previously [16]. HEK293-FT cell line

was cultured in high glucose DMEM with 10% FBS and 10mM HEPES, 1X Penicillin

Streptomycin, 1X GlutaMax and 1X non-essential amino acid and transfected with the

appropriate plasmids to generate viral supernatants. All cell lines were tested for

mycoplasma contamination twice every year, and found to be negative.

Antibodies and Reagents

Antibodies/Reagents	Source and Cat no	Dilution/ Conc.
Mouse anti-myc (9E10)	Abcam #ab32	1:1000
Mouse anti-myc (9E10)	Culture supernatant, in house	1:1000
Rabbit anti-SNAP23	SYSY#111203	1:1500
Mouse anti-SNAP25	SYSY#111011	1:1500
Rabbit anti-SNAP29	SYSY#111303	1:1500
Rabbit anti-GFP	Invitrogen#A11122	1:2500
Rabbit anti-flag	Sigma#F7425	1:1600
Mouse anti-flag (M2)	Sigma#F3165	1:1500
Mouse anti-6X His	Invitrogen #MA1-135	1:2000
Rabbit anti-Stx11	Sigma#SAB4301593	1:1000
Mouse anti-NFATc2 (4G6-G5)	Santacruz#7296	1:500
Goat anti-Lamin B (M-20)	Santacruz#6217	1:500
Rabbit monoclonal anti-HA	CST#3724S	1:1000
Alpaca VHH anti-rabbit AF488	Jackson #611-544-152	1:400 (1.375μg/ml)
DAPI	Invitrogen #D1306	2μg/ml
Rabbit anti-Stx11	SySy #110113	1.67μg/ml
Rabbit anti-Stx11	Invitrogen #PA5-50800	1:2000

Goat anti-rabbit A647	Invitrogen #A-21245	1:800
Rabbit anti-Stx11	Proteintech #13301-1-AP	0.3µg/ml
Donkey anti-rabbit A647	Jackson #711-605-152	1:800
HRP GAPDH	Sigma #G9295	1:40000
HRP Donkey anti-Rabbit	Jackson #711-035-152	1:50000
HRP Donkey anti-Mouse	Jackson #715-035-151	1:50000
Rabbit anti-NFAT1	Cell Signaling Technology	1:200
Ultra-LEAF Purified anti-human CD3	Biolegend #317326	5-10µg/ml
Ultra-LEAF Purified anti-human CD28	Biolegend #302934	2µg/ml
Ultra-LEAF Purified anti-human CD49d	Biolegend #304340	2µg/ml
PE anti-human CD107a (LAMP-1)	Biolegend #328608	1:50
APC anti-human CD8 antibody	Biolegend #344722	0.5µg/ml
Fura-2-AM	Invitrogen #F1221	1µM
Thapsigargin (TG)	Sigma #T9033-1MG	1µM
Polybrene (Hexadimethrine-bromide)	Sigma #107689-10G	8µg/ml
NE-PER kit	Thermo #78833	
Protein A/G Mag Sepharose xtra	Cytiva #28967056/66	
Protein A/G 4 Fast Flow Sepharose	Cytiva #17528001	
Protease Inhibitor Cocktail (Mammalian)	Sigma #539134	
Protease Inhibitor Cocktail (Bacterial)	Calbiochem #539132	
Talon Beads	Takara #635502	
Dextrin Sepharose Beads	Cytiva #28935597	
HEPES	Sigma #H3375-100G	
Lipofectamine 2000	Thermofisher #11668019	
Lipofectamine 3000	Thermofisher #L3000001	
Lemo21(DE3) *E.coli.* cells	NEB #C2528J	
IPTG	Himedia #MB072-10G Sigma #16758-10G	
Sarkosyl (N-lauroylsarcosine sodium	Sigma #L9150-50G	

salt)		
Lysozyme	Sigma #62970-5G-F	
DNaseI Recombinant RNase-free	Roche #04716728001	
PMA (Phorbol Myristate Acetate)	Sigma #P8139	20-50ng/ml
Ionomycin	Sigma #407952	$1\mu M$
Human IL-2 protein	Acro Biosystems #H5215	50ng/ml
Lectin from *Phaseolus vulgaris* (PHA)	Sigma #L8754	$2\mu g/ml$
Retronectin	Takara #T100B	$20\mu g/ml$
RNeasy Plus Mini kit	Qiagen#74134	
QIAshredder	Qiagen#79654	
Superscript IV	ThermoFisher#18091300	
Qubit ssDNA Assay kit	ThermoFisher# Q10212	
Human Stx11 TaqMan probe (FAM-MGB)	ThermoFisher# Hs00186823_m1	
Human Beta Actin TaqMan probe(FAM-MGB)	ThermoFisher# Hs01060665_g1	
Human IL2 Taqman probe (FAM-MGB)	ThermoFisher# Hs00174114_m1	
Paraformaldehyde	Electron Microscopy Sciences #157-8	
Bovine Serum Albumin	Jackson #001000173	
Luria Bertini Broth	Himedia #M1245	
HBSS	Sigma #H8264	
Terrific Broth	Sigma #T9179	
Alexa fluor 647 conjugated Alpha Bungarotoxin	Invitrogen #B35450	$1\mu g/ml$
HRP rabbit anti-GFP	Invitrogen #A10260	
HRP Goat anti-rabbit Fc	Jackson #111-035-008	1:50,000
Ampicillin	Sigma #A0166-5G	
30% Acrylamide:Bis Solution (29:1)	Biorad #1610156	

DTT (Dithiothreitol)	Biorad #1610611	
0.05% Trypsin-EDTA	Gibco #25300054	
100X Penicillin Streptomycin	Sigma #P4333-100ML	
Puromycin	Gibco #A11138-03	
Pluronic F-127 (20% solution in DMSO)	Invitrogen #P3000MP	0.02%

Lentiviral transductions

For the generation of lentiviral supernatants, shRNAs cloned in pLKO.1 were co-
transfected with psPAX2 packaging plasmid and pMD2.G envelope plasmid into HEK293-
FT cells using the calcium phosphate method of transfection. 48 and 72 hours post-
transfection, viral supernatants were collected, pooled and stored at -80°C till further use.
For transduction of HEK293 and Jurkat cells, 0.1 million cells were plated in 6- or 24-well
plates either the day before (HEK293) or the same day (Jurkat). Viral supernatants were
added to the cells along with 8ug/ml polybrene and cells were spun at 2500 RPM, 30°C
for 90 mins. 24 hours or 48 hours post-spinfection, Puromycin was added at a final
concentration of 1ug/ml to HEK293 and Jurkat cells, respectively. Cells were analyzed 3-
5 days post transduction.

Immunocytochemistry and confocal imaging

For Stx11 localization, resting or store-depleted (1μM Thapsigargin) WT HEK293 or
HEK293 cells stably expressing Orai1-YFP were either plated in 6 well plates and
spinfected with viral supernatants generated from pMSCV-Stx11(HA)-IRES-mCherry,
pMSCV-IRES-mCherry (empty vector control) or plated in carbon coated, glow
discharged 35mm glass bottom dishes (IBDI) and transfected with pEF1alpha-Stx11-
IRES-mCherry. 24 hr post-spinfection, cells were plated in 35mm glass bottom dishes.
For immunolabelling, cells were washed with Ringer's buffer, fixed with 4% PFA and
blocked with 3% BSA containing 0.1% NP40 for 1.5 hrs. Post-fixation and
permeabilization, cells were incubated either with anti-HA (CST) or rabbit anti-Stx11
(SySy/ Proteintech) primary antibodies O/N at 4°C, washed and incubated with either
Alpaca VHH anti-rabbit AF488, donkey anti-rabbit A647 or goat anti-rabbit A647

secondary antibodies for 1 hour at room temperature (RT). Cells were counter stained

with DAPI for 10 min at RT and imaged using Olympus FV3000 laser scanning

microscope. Images were acquired sequentially using the following parameters: DAPI

(DM405/488 dichroic,   405nm excitation, 430-470nm emission); anti-HA Alexa488

(DM405/488, 488nm excitation, 500-590nm emission); Stx11 (DM405/488/561/640,

640nm excitation, 650-750nm emission); Orai1-YFP (DM405/488/561/640, 488nm

excitation, 521-591nm emission); CFP-Stim1 (DM405-445/514/594, 405nm excitation,

448-510nm emission).

Quantification of Orai1 and Stim1 intensities inside/outside ER-PM puncta

HEK293 cells were plated in 6 well plates and transduced with scramble or STX11

shRNA. 48hr post transduction, cells were trypsinized and plated in carbon coated 35mm

dishes and co-transfected with CFP-Orai1 and Stim1-YFP. Images were acquired

sequentially for CFP and YFP using Olympus FV3000, using 405nm and 514nm lasers

respectively using a 60X objective. Cells were washed and incubated in Ringer's buffer

prior to imaging and the positions of CFP and YFP double positive cells were marked.

Resting Orai1 and Stim1 images were acquired first. Following this, cells were incubated

with Thapsigargin (1uM) plus EGTA (10mM) for ~8 mins before capturing the store-

depleted images. The imaging was done in live cell chamber to maintain the temperature

at 37C. For analysis, Stim1-YFP images were masked using Phansalkar local

thresholding in ImageJ software. Stim puncta were detected on the image mask using

"Analyze Particles" in ImageJ and used for determining the intensity values of CFP-Orai1

and Stim1-YFP inside Stim1:Orai1 co-clusters. Cell boundaries were manually drawn

using CFP-Orai1 images and total Orai1 intensities were obtained using these ROIs. The

Stim/ Orai intensity values obtained from the above analysis were plotted using Origin

software. In the experiment where the direction of fluorescent tags on Orai1 and Stim1

were flipped, Orai1-YFP or CFP-Stim1 expressing cells were transduced with scramble

or STX11 shRNA and plated as mentioned above. Images were acquired using a TIRF

microscope setup described before [14]. Images were captured in resting cells and positions

marked. Store-depleted images were acquired after incubation with 1uM Thapsigargin

(TG) and 10mM EGTA for ~6 mins. Analysis was done as mentioned above with Otsu

local thresholding in ImageJ software. Stim1 and Orai1 intensity (AU) and area (pixels)

values obtained from the analysis were plotted using Origin software.

Alpha-bungarotoxin binding assay

U2OS cell line stably expressing Orai1-BBS-YFP construct has been described

previously[14]. Briefly, Orai1-BBS-YFP expressing U2OS cells were transduced with

scramble or Stx11 shRNA. Cells were trypsinized 3-4 days post-transduction, centrifuged

and labelled on ice with AF647 conjugated alpha-bungarotoxin (1μg/ml) containing 2%

FBS in 1X HBSS for 30 mins. Following this, cells were washed with HBSS and fixed with

4% PFA (~20 min at RT). Samples were run using Cytoflex FACS analyzer (Beckman

Coulter) and data were analyzed using Flow Jo software.

Validation of Stx11 knockdown using QPCR

Scramble and Stx11 shRNA treated HEK293 cells were trypsinized, counted and equal

number of cells were used for lysis. The lysates were homogenized using QIA shredder

columns (Qiagen) and RNA isolation was done using RNeasy Plus Mini kit (Qiagen). Total

RNA isolated from both the groups was used for cDNA synthesis using oligo dT primers

and Superscript IV as per manufacturer's guidelines. The concentration of synthesized

cDNA was estimated using Qubit ssDNA assay kit and STX11 TaqMan probes were used

to perform the QPCR using LightCycler 96 (Roche). Beta-actin was used as

housekeeping control.

Single cell Ca^{2+} imaging

Cells were plated in carbon-coated glass bottom dishes (one day prior or 30 mins before

the assay for HEK293 and Jurkat T cells respectively) and loaded with 1uM Fura-2 AM

dye in HBSS (CaCl$_2$ 1.8mM, KCl 5.36mM, MgSO$_4$ 0.81mM, NaCl 136.89 mM, Na$_2$HPO4

0.335mM, D-Glucose 5.55mM) for 30 mins, at 37°C in the dark. After incubation, cells

were washed and incubated in Ringer's buffer (135mM NaCl, 5mM KCl, 1.8mM CaCl$_2$,

1mM MgCl$_2$, 5.6mM Glucose, 10mM HEPES) for an additional 10 mins, washed and

imaged in Ringer's buffer or Calcium-free Ringer's buffer (135mM NaCl, 5mM KCl, 1mM

MgCl$_2$, 5.6mM Glucose, 10mM HEPES at pH7.5), as indicated. Olympus IX-71 inverted

microscope equipped with a Lamda-LS illuminator (Sutter Instrument, Novato, CA), Fura-

2 (340/380) filter set (Chroma, Bellows Falls, VT), a 10X 0.3NA objective lens (Olympus,

UPLFLN, Japan), and a Photometrics Coolsnap HQ2 CCD camera was used to capture

images at a frequency of ~1 image pair every 2 or 4 seconds interval. Approximately 30-

50 cells were imaged per group in each experiment unless otherwise stated. Data were

acquired, analyzed and plotted using MetaFluor (Molecular Devices, Sunnyvale, CA),

Microsoft Excel, and Origin softwares.

For constitutive calcium influx assay, shRNA treated HEK293 cells were either

transfected or nucleofected with Orai1-S-S-EGFP or Orai1(H134S)-YFP and ~12-14 hour

post transfection (~4-5 hour post nucleofection), cells were loaded with Fura-2 AM,

washed and imaged in Ringer's buffer with 0mM calcium to acquire baselines and 2mM

calcium thereafter. To measure CAD induced constitutive calcium influx, Orai1-CFP

expressing stable HEK293 cells were nucleofected with YFP-CAD and analyzed ~4 hour

later as described above. To identify cells expressing Orai1 mutants and fusion proteins

in the plasma membrane, cells were imaged using a 20X 0.7NA water objective lens

(Olympus, UApoN340, Japan). Images were acquired at a frequency of ~1 image pair

every 10 seconds interval to avoid photobleaching and analyzed as described above.

Measurement of NFAT nuclear localization by Western Blot

Jurkat T cells were transduced with scramble or STX11 shRNA to knock-down STX11 as

described above. On day 4 post-transduction, cells were collected, spun down and

resuspended in plain RPMI media and rested for 1hr at 37°C. Following this, cells were

counted and divided into two equal groups. One group was resuspended in RPMI

(unstimulated) and the other in RPMI media containing 1uM TG + 10ng/ l PMA (Phorbol

Myristate Acetate) (stimulated) and incubated at 37°C for 30 mins. Following incubation,

cells were pelleted and the nuclear and cytosolic protein fractions were separated using

NE-PER kit according to the manufacturer's guidelines and subjected to SDS-PAGE and

Western Blot using the mouse anti-NFATc2 primary antibody (Santa Cruz) followed by

Donkey anti-mouse secondary antibody.

Estimation of nuclear translocation of NFAT in Jurkat T cells by imaging

Control and STX11 depleted Jurkat T cells were rested in plain RPMI for ~1 hour prior to
the assay. ~100,000 cells per group were plated on freshly carbon-coated coverslips for
40 minutes and stimulated for 1 hour at 37°C with 5 ug/ml anti-CD3 antibody, diluted in
plain RPMI. Following this, cells were washed and fixed using 4% PFA diluted in 1X PBS
for 20 minutes at RT, washed twice and incubated with 30 mM Glycine for 10 minutes at
RT. For permeabilization and blocking, cells were incubated with 0.1% Saponin, 3% BSA
diluted in 1X PBS for 1 hour at RT, washed and incubated with anti-NFAT primary
antibody (anti-NFAT1, CST) at 4°C, overnight. Following primary antibody application,
cells were washed with 1X PBS and incubated with anti-Rabbit AF647 secondary
antibody for 1 hour, washed and stained with DAPI for 5 minutes followed by additional
washes. Images were acquired in the FV3000 confocal microscope. DAPI was used to
identify the nuclear area. Nuclear *versus* whole cell (total) NFAT mean intensity ratio was
plotted across different groups.

Whole cell lysates (WCLs), Western Blot and Co-immunoprecipitation (Co-IP)
HEK293 cells transfected with the desired plasmids were lysed using buffer containing
50mM Tris-Cl (pH 8.0), 150mM NaCl, 1% NP-40, 1mM PMSF, and protease inhibitor
cocktail. The whole cell lysates were centrifuged at 21000g for 15 minutes and
supernatants were subjected to SDS-PAGE, and Western Blot. For
immunoprecipitations, lysates were divided into two equal parts. To the first tube, the
appropriate anti-mouse or anti-rabbit primary antibody was added and to the second,
same amount of the respective IgG control antibody was added. The antibody-lysate
mixtures were incubated overnight at 4°C. Following this, Protein A/G Mag Sepharose
beads were added and incubated with the antibody-lysate mixtures for 4 hours, washed
with the lysis buffer containing 0.1% NP-40, boiled with 1X Laemmli buffer + 120mM DTT
and subjected to SDS-PAGE and Western Blot analysis. Typically, 1/10th of the whole cell
lysate (WCL) was loaded in the input lane of the co-IP blots.

***E. coli* expression and *in-vitro* binding assays**

Full-length His$_6$-tagged Stx11 and truncation mutants were cloned in pET28b, expressed

in Lemo21 (DE3) *E. coli* cells and induced with 1mM IPTG (Isopropyl β- d-1-

thiogalactopyranoside) for 14-18 hours at 18°C. The cell pellets were lysed in buffer

containing 50mM Tris-Cl (pH 8.0), 150mM NaCl, 10% Glycerol, 1mM PMSF, 1%

Sarkosyl, 0.1mg/ml Lysozyme, protease inhibitor cocktail and sonicated on ice. DNase I

was added after sonication and the lysates were further incubated for ~60 minutes before

centrifugation at 21000g for 40 mins. The supernatants were subjected to SDS-PAGE to

confirm expression by Coomassie staining and subsequently used for pull-down assays.

MBP-tagged Orai1 constructs were expressed in Lemo21 (DE3) *E. coli* cells, induced with

0.3mM IPTG (Isopropyl β- d-1-thiogalactopyranoside) for 14-18 hours at 18°C. The cell

pellets were lysed in buffer containing 50mM Tris-Cl (pH 8.0), 150mM NaCl, 5% Glycerol,

1mM PMSF, 0.1mg/mL Lysozyme and protease inhibitor cocktail (1:200) and sonicated

on ice. DNase I was added after sonication and the lysates were further incubated for ~60

minutes before centrifugation at 21000g for 40 mins. The supernatants were collected

and subjected to SDS-PAGE to confirm expression by Coomassie staining and used for

pull-down assays. Following lysis, the His$_6$-tagged Stx11 proteins were purified using

Talon Beads and MBP-tagged Orai1 proteins using Dextrin Sepharose/Amylose resin

according to the manufacturer's guidelines. For *in-vitro* binding assays, lysates prepared

from cells expressing MBP or MBP-tagged Orai1 fragments were incubated with 25μL

Dextrin Sepharose Beads and incubated for 2 hours at 4°C. After incubation, beads were

washed thrice and ~50-125ng of purified His-tagged full length Stx11, His-tagged SNARE

or His-tagged H$_{abc}$ domains were diluted in buffer containing 50mM Tris-Cl (pH8), 150mM

NaCl, 5% Glycerol and 0.1% NP-40 and added to the beads. Following 1 hour of

incubation at 4°C, the beads were washed thrice, re-suspended and boiled in the binding

buffer containing 1X Laemmli and 120mM DTT. The eluate was subjected to SDS-PAGE

and Western blot. MBP and MBP-tagged proteins were detected with Ponceau staining,

Full length STX11 using Rabbit anti-STX11 primary antibody (Thermo) followed by

Donkey anti-rabbit HRP and His-tagged STX11 fragments were detected or mouse anti-

6XHis primary antibody (Invitrogen) followed by Donkey anti-mouse HRP secondary

antibody. 1/5th of the protein was loaded in the input lane for all *in vitro* binding assays.

QPCR to estimate gene expression in Jurkat T cells

Scramble (scr) and STX11 shRNA treated Jurkat T cells were rested for ~1 hour prior to the assay, stimulated with soluble anti-CD3 (5-10 μg/ml) for 3 hours and washed with HBSS. Total RNA was isolated using RNeasy Mini kit as per manufacturer's instructions. cDNA was synthesized using random hexamers and Invitrogen Superscript IV kit and quantified using Qubit ssDNA assay. Taqman probes for IL-2 and beta-actin housekeeping control were used for performing the QPCR in triplicates. Data analysis was done by calculating the double delta C_t values.

Isolation and culture of human PBMCs

All experiments with human PBMCs were approved by the Institutional Human Ethics Committee. Healthy human whole blood freshly collected in heparin or Citrate Phosphate Dextrose Adenine (CPDA) solution was subjected to density gradient centrifugation using Ficoll-Paque PLUS media. The buffy coat was separated, washed twice with HBSS and cultured with IL-2 (50 ng/ml). Cells were stimulated once per week with PHA (2 ug/ml) for 72 hours and rested in IL-2 for the remaining 72-96 hours. Unless specified, all the assays were performed following 24-48 hours of rest, post stimulation.

Isolation of genomic DNA and sequencing of FHLH4 patient mutation.

Genomic DNA was extracted from healthy donor and FHLH-4 patient PBMCs using Phenol-Chloroform-Isoamyl alcohol. Stx11 genomic region flanking the mutation was PCR amplified using primers: 5' Forward -CATGCACGACTACAACCAGGC and 3' Reverse -GGGACAGCAGAAGCAGCAGAGGG. The resulting PCR products were separated on 2% agarose gel, excised and extracted using the Macherey-Nagel Nucleospin columns and subjected to Sanger sequencing using the 5' Forward PCR primer to confirm the mutation.

Measurement of SOCE in human PBMCs

PBMCs in culture were washed, rested and allowed to adhere to freshly carbon-coated (IBDI) dishes for 1 hour in plain RPMI at 37°C. The cells were washed with HBSS and incubated with 1ml of 1μM Fura-2-AM diluted in Ringer's buffer for 40 minutes, washed

and incubated for an additional 10 minutes to allow de-esterification of the dye. The assay

was started with 1 ml of Calcium free Ringer's buffer in the imaging dish and images were

captured every 4 seconds. After capturing the baseline for ~60 seconds, stores were

depleted by the addition of 1 μM Thapsigargin. In other assays, 10 μg/ml of anti-CD3 and

5 μg/ml of the secondary antibody were used to cross-link the TCRs and thereby induce

store depletion. ~5 minutes post-store-depletion, Calcium Chloride ($CaCl_2$) was added

back to the dish at a final concentration of 2mM to estimate the magnitude of store

operated calcium entry.

Degranulation assay

PBMCs were cultured in RPMI containing 10% FBS and 50 ng/ml IL-2 and stimulated

with PHA (2 ug/ml) 48hrs before the assay. 24hrs prior to the assay, IL-2 was washed off

but PHA was re-added. On the day of the assay cells were washed twice to remove PHA

and any growth factors. To measure degranulation, cells were either left unstimulated,

stimulated with a combination of anti-CD3 (10 μg/ml plate-coated), anti-CD28 (2 μg/ml

soluble) and anti-CD49d (2 μg/ml soluble) or a combination of Ionomycin (1 μM) and PMA

(50 ng/ml) for 3.5 hours. CD107a-PE antibody (1:50 dilution) was added to each of the

three groups at the start of stimulation. Following stimulations, cells were transferred to

ice, washed with cold HBSS containing 2% FBS and incubated with anti-CD8 APC for 20

minutes, washed twice with cold HBSS and analyzed using Cytoflex (Beckman Coulter)

flow cytometer.

QPCR to estimate gene expression in human PBMCs

PBMCs were taken off IL-2 48 hours prior to the assay. On the day of the assay, cells

were washed twice, left unstimulated and either stimulated with anti-CD3 (10 μg/ml, plate-

coated), anti-CD28 (2 μg/ml) and anti-CD49d (2 μg/ml) or with Ionomycin (1 μM) and PMA

(50 ng/ml) for 6 hours. To end the stimulation, cells were washed with cold HBSS, pelleted

and used for RNA extraction. Total RNA was isolated using Qiagen RNeasy mini Kit and

cDNA was synthesized using random hexamers and Superscript IV (Invitrogen). cDNA

was quantified using Qubit and subjected to QPCR analysis using Taqman probes for IL-

2 and beta-actin housekeeping gene control in triplicates. Analysis was performed by
calculating the double delta C_t values.

Transduction of human PBMCs with pMSCV-STX11

For viral transduction of PBMCs, polystyrene non-TC treated 24-well plates were coated
with retronection (20 μg/ml) overnight at 4°C, blocked with 2% BSA for 30 minutes at
room temperature and washed twice with HBSS. The retroviral supernatants were added
onto the coated wells and the plates were spun at 1800g for 2 hours at 30°C. Following
spin, the wells were washed with the blocking solution. PBMCs cultured in RPMI
containing 10% FBS were stimulated with PHA (2 μg/ml) and IL-2 (50 ng/ml) for 48 hours
prior to transduction. Stimulated PBMCs were transferred to coated plates at a density of
0.25 million cells/well centrifuged at 400g for 40 minutes at 30°C. The cells were analyzed
48-72 hours post transduction.

Stx11-Orai1 complex prediction

Full length STX11 and Orai1 and their two domains, (H_{abc} 41-167 and SNARE 183-
267) and (N-terminus 1-87 and C-terminus 256-301), respectively were used to generate
the complex of STX11 and Orai1 using AlphaFold3 (AF3) in all combinations. The
resultant models were assessed using ipTM and pTM scores. The best scoring
combination (STX11- H_{abc} and Orai1-C-terminus) was further considered to generate
more models by changing the seed values. The best model in terms of highest ipTM and
pTM values was considered for further analysis. Custom script was used to analyze the
contact frequency of interacting residues of Stx11-H_{abc} and Orai1-C-terminus complex
across all predicted models.

Molecular dynamics (MD) simulation

The STX11- H_{abc}-Orai1-C-terminus complex was subjected to all-atom MD
simulation. Initially, the complex was prepared using protein-preparation module of
Schrodinger which involves H-bond network optimization and restrained minimization of

the initial structure. The prepared structure was solvated using TIP3P water model in an
orthorhombic box and neutralized by adding counter ions. OPLS4 force field was used,
and simulation system was generated by specifying 150mM salt (NaCl). The solvated
system was subjected to default relaxation protocol of Desmond followed by production
run for 500 ns at 300K and 1 atm pressure in NPT ensemble. The default relaxation
protocol includes several short simulation steps. (1) Brownian dynamics simulation for
100 ps at 10 K temperature in NVT with restraints on solute heavy atoms (2) Simulation
in NVT ensemble at 10 K for 12 ps with restraints on solute heavy atoms (3) 12 ps
simulation in NPT ensemble at 10 K with restraints on solute heavy atoms (4) Simulation
in NPT ensemble for 12 ps with restraints on solute heavy atoms (5) Simulation in NPT
ensemble for 24 ps without restraints. Three independent runs were executed with
different initial seeds.

Analysis of MD simulation

The MD runs were analyzed for the stability of the complex and interactions
between the subunits. The RMSD was calculated using simulation interaction diagram
(SID) module of Schrodinger. The first frame was used as reference frame for RMSD
calculation. Script "analyze_trajectory_ppi.py" was used to calculate the interactions
between the STX11 and Orai1. The binding energy (ΔG) at each nanosecond across
simulation was calculated using "thermal_mmgbsa.py" script of Schrodinger.

The representative structure was generated through trajectory clustering. The last
200 ns frames from each run were combined and clustered using scripts "trj_merge.py"
and "trajectory cluster" of Schrodinger. The cluster representative of the largest cluster
was used to analyze the protein-protein interactions. PDB Sum was used to calculate the
interactions in the cluster representative. Figures were generated using Pymol and MD
movies using Maestro.

Figure Legends:-

Figure 1: STX11 is required for SOCE.

(**A-F**) RNAi mediated depletion of STX11 reduces SOCE. Measurement of thapsigargin

(TG) induced SOCE in various cell lines after shRNA mediated depletion of STX11. The

traces in panels **A,C,E** show representative average single cell Fura-2 calcium imaging

assays. Bars in panels **B,D,F** show mean % SOCE $\pm$ SE from three independent

experiments each where mean SOCE from scramble (scr) shRNA treated group in each

experiment was set at 100% and the relative response of STX11 shRNA treated groups

was calculated respectively. *$P<0.05$; **$P<0.01$; ***$P<0.001$ using two-tailed Student's *t*

test. (**G**) Representative Fura-2 calcium imaging assay showing reconstitution of SOCE

in STX11 depleted Jurkat T cells by ectopic expression of STX11. Black (scr shRNA), red

(STX11 shRNA), green (STX11 shRNA with STX11 expression). N=2 (**H**) Ectopic

expression of STX11 enhances SOCE. Representative Fura-2 calcium imaging assay

showing SOCE in HEK293 cells expressing STX11 (red) or empty vector (EV) (black).

(N=3) (**I**) STX11 mediated enhancement of SOCE is dependent on Orai1. A

representative Fura 2 calcium imaging assay showing measurement of thapsigargin (TG)

induced SOCE in HEK293 cells where Orai1 expression was depleted using shRNA and

STX11 was over-expressed. (N=2)

Figure 2: STX11 depletion suppresses I$_{CRAC}$, downstream signaling and gene

expression in Jurkat T cells. I$_{CRAC}$ was recorded from Jurkat T-cells in the whole-cell

recording configuration in 20 mM extracellular Ca^{2+} Ringer's solution. I$_{CRAC}$ was induced

by passive depletion of intracellular Ca^{2+} stores by dialyzing 8 mM BAPTA into the cell via

the patch-pipette. (**A**) Representative current at -100 mV in Jurkat T cell transfected with

scr shRNA construct. The current is blocked by extracellular La^{3+} (100 µM) and replacing

the 20 mM Ca^{2+} Ringer's solution with a divalent free solution (DVF) evokes a large Na+

current which depotentiates over tens of seconds. The current-voltage (I-V) relationship

of the Ca^{2+} and DVF currents are shown on the right. (**B**) I$_{CRAC}$ from a Jurkat T cell

transfected with STX11 shRNA. Both Ca^{2+} and Na^+ current amplitudes are reduced

relative to control cells. The I-V relationships (*right plots)* show no change in ion

selectivity. (**C-D**) Summary of the current amplitudes of Ca^{2+} and Na^{+} currents and

current reversal potentials in scr and STX11 knockdown cells. (**E&F**) Estimation of

nuclear translocation of NFAT. (**E**) Western Blot showing nuclear translocation of NFAT

in Jurkat T cells treated with scr or STX11 shRNA for 4 days and stimulated with PMA+TG

for 30min prior to the preparation of nuclear and cytoplasmic extracts. (N=3) (**F**)

Representative confocal images of Jurkat T cells treated with scr or STX11 shRNA for 4

days and stimulated with 10ug/ml anti-CD3 for 1 hour. Following stimulation, cells were

fixed, permeabilized and stained using anti-NFAT primary antibody, followed by donkey

anti-rabbit AF647 secondary antibody, and counter-stained with DAPI to mark the nuclei.

(N=2) (**G**) Box and whisker plot showing percent nuclear NFAT in Jurkat T cells quantified

from 40-50 cells populating 10 randomly chosen fields per group in (**F**). Boundaries of the

box plots represent 25^{th} and 75^{th} percentile values, horizontal line represents mean, white

circle represents median and whiskers denote the outliers. (**H**) Quantitative PCR to

assess IL-2 transcription in anti-CD3 stimulated Jurkat T cells. Jurkat T cells were treated

with scr or STX11 shRNA for 4 days and stimulated with 5ug/ml anti-CD3 for 3 hours.

Total RNA was extracted from cells and subjected to QPCR analysis using Taqman

probes for IL-2 and beta-actin. The bars show relative IL-2 mRNA expression levels with

the scr shRNA treated group set at 100%. Shown here are mean $\pm$ SE. (N=3) **(F-H)**

*P<0.05; **P<0.01; ***P<0.001 using two-tailed Student's t test.

Figure 3: STX11 directly binds resting Orai1 in the plasma membrane. (**A**) STX11

localizes in the plasma membrane. Representative confocal images of HEK293 cells

expressing internal-HA-tagged STX11, stained using anti-HA antibody followed by alpaca

VHH anti-rabbit secondary nanobody. Scale bar 15um. (N=3) (**B**) Co-localization of

STX11 and Orai1 in the plasma membrane. Representative confocal images of HEK293

cells expressing Orai1-YFP and STX11, stained using anti-STX11 antibody followed by

donkey anti-rabbit secondary antibody. Scale bar 10um. (N=3). (**C&D**) Co-

immunoprecipitation of STX11 with Orai1. Whole cell lysates of resting and store-depleted

HEK293 cells expressing either Flag-Orai1 and STX11 (**C**) or STX11 and Orai1-Myc-His

(**D**) were subjected to immunoprecipitation and Western Blot using anti-Myc, anti-Flag or

anti-STX11 antibodies, as indicated. (N=3) **(E)** Schematic showing key domains of STX11

and Orai1 used for *in vitro* pulldown assays. (**F**) Pull-down assay showing *in vitro* binding of His-tagged full length STX11 to MBP-tagged cytosolic domains of Orai1. MBP-tagged Orai1 fragments, expressed in *E. Coli* and immobilized on the amylose resin, were incubated with purified His-tagged STX11 protein. Post incubation, beads were washed, boiled and subjected to Western Blot analysis using anti-STX11 antibody. (Top panel) Ponceau S staining showing the input of MBP alone or MBP-tagged fragments. (Bottom panel) Western Blot using anti-STX11 antibody. (N=3) (**G**) Pull-down assay showing *in vitro* binding of His-tagged H_{abc} domain of STX11 to MBP-tagged Orai1 N- and C-termini performed as described above. (Top panel) Ponceau S staining showing the input of MBP alone or MBP-tagged Orai1 cytosolic tails. (Bottom panel) Western Blot using anti-His antibody. (N=3). (**H-I**) Representative structure of STX11 H_{abc} and Orai1 C-terminus complex after MD simulation (**H**) Sphere representation of STX11 H_{abc} (green) and Orai1 C-terminus (cyan) complex. The N-termini are highlighted in blue, and C-termini are highlighted in red (**I**) Cartoon representation of STX11-H_{abc} (green) and Orai1 C-terminus (cyan) complex highlighting the CA of terminal residues as spheres (**J**) Protein-protein interactions between the STX11 H_{abc} and the Orai1 C-terminus.

Figure 4: STX11 depletion compromises the functional assembly of Orai1 with Stim1 in ER-PM junctions. (**A-B**) Representative confocal images of resting (**A**) and store-depleted (**B**) scr and STX11 shRNA treated HEK293 cells expressing N-terminal CFP tagged Orai1 (CFP-Orai1) and C-terminal YFP tagged Stim1 (Stim1-YFP). (**C-H**) Box and whisker plots showing (**C**) Quantification of Stim1-YFP intensities inside Stim:Orai clusters of scr and STX11 shRNA treated HEK293 cells, ~8 min post store-depletion. (**D-E**) Quantification of CFP-Orai1 intensities inside (**D**) and outside (**E**) Stim1-YFP clusters of scr and STX11 shRNA treated cells, ~8 minute post store-depletion. N=3. Boundaries of the box plots represent 25^{th} and 75^{th} percentile values, horizontal line represents mean, white circle median and whiskers denote the outliers. (**F**) Quantification of total Orai1 levels in the plasma membrane of STX11 depleted cells. U2OS cells stably expressing Orai1-BBS-YFP were transduced with scr (black) or STX11 (red) shRNA, stimulated with 1uM TG, incubated with alpha-bungarotoxin alexa 647 (BTX-A647) and washed. BTX binding to surface Orai1 was measured using FACS, where binding to

wildtype HEK293 cells was used as control. (N=3). (**G-J**) Quantification of C-term tagged Orai1 (Orai1-YFP) and N-term tagged Stim1 (CFP-Stim1) inside puncta in control and STX11 depleted cells, ~6 minute post store-depletion. (**G-H**) Quantification of mean intensities of Orai1-YFP (**G**) and CFP-Stim1 (**H**) inside puncta. n=8. Quantification of mean area of Orai1-YFP (**I**) and CFP-Stim1 (**J**) puncta. The continuous and dotted lines represent mean and median respectively. *P<0.05, **P<0.01 and ***P<0.001 using two-tailed Student's *t* test. (**K**) Representative Fura-2 calcium imaging assay to measure Thapsigargin induced SOCE in Orai1-YFP and CFP-Stim1 expressing HEK293 cells treated with scr or STX11 shRNA.

Figure 5: STX11 binding to resting Orai1 allows successful gating by Stim1. (A) Representative Fura-2 calcium imaging assay to measure constitutive calcium influx in Orai1-S-S-GFP expressing control or STX11 depleted HEK293 cells. Cells were imaged in Ringer's buffer containing 0mM followed by 2mM extracellular Ca^{2+}. (**B**) Box and whisker plot showing quantification of constitutive calcium influx across experiments as shown in (**A**). N=3. (**C**) Representative images of YFP-CAD localization in Orai1-CFP expressing, control and STX11 depleted HEK293 cells. (**D**) Quantification of YFP-CAD in the plasma membrane of Orai1-CFP expressing HEK293 cells represented as % of total YFP-CAD in respective cells. (**E**) Representative Fura-2 calcium imaging assay to measure constitutive calcium influx in Orai1-CFP and YFP-CAD expressing control or STX11 depleted HEK293 cells. (**F-G**) Quantification of constitutive calcium influx in Orai1-CFP and YFP-CAD expressing cells described in **E,** at 0mM (**F**) and 2mM (**G**) extracellular Ca^{2+}. N=3. (**H**) Representative Fura-2 calcium imaging assay to measure constitutive calcium influx in Orai1-H134S mutant expressing control and STX11 depleted HEK293 cells as described in (**A**). (**I**) Quantification of constitutive calcium influx across experiments as shown in (**H**). N=3. Boundaries of the box plots in **B,D,F,G&I** represent 25th and 75th percentile values, horizontal line represents mean, white circle median and whiskers denote the outliers. *P<0.05, **P<0.01 and ***P<0.001 in (C-D,F-G,I) using two-tailed Student's *t* test.

Figure 6: Ionomycin rescues the cytotoxicity and gene expression defects in STX11 deficient FHLH4 patient T lymphocytes. (**A**) Sanger sequencing of the patient DNA showing deletion of a single of Adenine at the 752nd position and the resulting frameshift. (**B**) Western Blot of whole cell lysates prepared from healthy human control and FHLH4 patient PBMCs showing the relative molecular weight and abundance of the wildtype and mutant STX11 bands. (**C**) Fura-2 calcium imaging assay measuring anti-CD3 induced SOCE in wildtype and FHLH4 PBMCs. (**D**) Fura-2 calcium imaging assay measuring SOCE in wildtype and FHLH4 PBMCs expressing either empty vector (EV) or wildtype human STX11, respectively. (**E**) Representative Fura-2 calcium imaging assay measuring SOCE in wildtype PBMCs with or without ectopic expression of wildtype STX11. (**F**) Bar plot showing quantification of SOCE across multiple experiments in (**E**). N=3. (**G**) Granule release assay performed on the *in vitro* cultured healthy human control and FHLH4 patient CD8 T cells. PBMCs isolated from control and FHLH4 patient blood were stimulated with (2ug/ml) PHA for 48 hours. On the day of assay, cells were washed and either left unstimulated or restimulated with plate coated anti-CD3 + soluble anti-CD28 or PMA + Ionomycin in the presence of CD107a-PE antibody, stained with anti-CD8 antibody and analyzed. (N=2) (**H**) Quantification of the granule release assays as shown in (**G**). (**I**) Analysis of IL-2 expression in control and FHLH4 T cells. *In vitro* cultured healthy human control and FHLH4 patient T cells were rested and stimulated with either plate coated anti-CD3 + soluble anti-CD28 or PMA + Ionomycin for 6 hours. Total RNA was extracted and subjected to quantitative PCR analysis using Taqman probes for IL-2 and beta-actin in triplicates. *$P<0.05$; **$P<0.01$; ***$P<0.001$ using two-tailed Student's *t*-test.

Figure 7: Hypothetical model showing Orai1 bound to STX11 for its priming. Blue ribbons depict Orai1 in plasma membrane (PM). Green and Orange ribbons represent H$_{abc}$ and SNARE domains of STX11, respectively. Also shown are the C-terminal cysteines of STX11 which facilitate membrane attachment of the protein.

Movies 1-3: Three independent molecular dynamic simulations of Orai1 C-terminus interaction with STX11 H$_{abc}$ domain. Due to BioRxiv file size limit, the

990   movies can be found at the following link:

991   https://www.dropbox.com/scl/fo/3j07d3sua4g58w3yfmyv7/AAKjvIiniA-

992   z994hxV9DNHo?rlkey=e7yj5wyukvwg5seg1awkuctc4&st=z6vp57dr&dl=0

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

References:

1. Clapham, D.E. Calcium signaling. *Cell* **131**, 1047-1058 (2007).
2. Vig, M. & Kinet, J.P. The long and arduous road to CRAC. *Cell Calcium* **42**, 157-162 (2007).
3. Vig, M. *et al.* Defective mast cell effector functions in mice lacking the CRACM1 pore subunit of store-operated calcium release-activated calcium channels. *Nat Immunol* **9**, 89-96 (2008).
4. Vig, M. & Kinet, J.P. Calcium signaling in immune cells. *Nat Immunol* **10**, 21-27 (2009).
5. Maul-Pavicic, A. *et al.* ORAI1-mediated calcium influx is required for human cytotoxic lymphocyte degranulation and target cell lysis. *Proc Natl Acad Sci U S A* **108**, 3324-3329 (2011).
6. Vig, M. *et al.* CRACM1 multimers form the ion-selective pore of the CRAC channel. *Curr Biol* **16**, 2073-2079 (2006).
7. Prakriya, M. & Lewis, R.S. Store-Operated Calcium Channels. *Physiol Rev* **95**, 1383-1436 (2015).
8. Soboloff, J., Rothberg, B.S., Madesh, M. & Gill, D.L. STIM proteins: dynamic calcium signal transducers. *Nat Rev Mol Cell Biol* **13**, 549-565 (2012).
9. Roos, J. *et al.* STIM1, an essential and conserved component of store-operated Ca2+ channel function. *J Cell Biol* **169**, 435-445 (2005).
10. Liou, J. *et al.* STIM is a Ca2+ sensor essential for Ca2+-store-depletion-triggered Ca2+ influx. *Curr Biol* **15**, 1235-1241 (2005).
11. Zhang, S.L. *et al.* Genome-wide RNAi screen of Ca(2+) influx identifies genes that regulate Ca(2+) release-activated Ca(2+) channel activity. *Proc Natl Acad Sci U S A* **103**, 9357-9362 (2006).
12. Vig, M. *et al.* CRACM1 is a plasma membrane protein essential for store-operated Ca2+ entry. *Science* **312**, 1220-1223 (2006).
13. Feske, S. *et al.* A mutation in Orai1 causes immune deficiency by abrogating CRAC channel function. *Nature* **441**, 179-185 (2006).
14. Yoon, T.Y. & Munson, M. SNARE complex assembly and disassembly. *Curr Biol* **28**, R397-R401 (2018).
15. Zhao, M. *et al.* Mechanistic insights into the recycling machine of the SNARE complex. *Nature* **518**, 61-67 (2015).
16. Miao, Y. *et al.* An essential and NSF independent role for alpha-SNAP in store-operated calcium entry. *Elife* **2**, e00802 (2013).
17. Li, P., Miao, Y., Dani, A. & Vig, M. alpha-SNAP regulates dynamic, on-site assembly and calcium selectivity of Orai1 channels. *Mol Biol Cell* **27**, 2542-2553 (2016).
18. Miao, Y., Bhushan, J., Dani, A. & Vig, M. Na+ influx via Orai1 inhibits intracellular ATP induced mTORC2 signaling to disrupt CD4 T cell gene expression and differentiation. *Elife* **6** (2017).
19. Ramanagoudr-Bhojappa, R., Miao, Y. & Vig, M. High affinity associations with alpha-SNAP enable calcium entry via Orai1 channels. *PLoS One* **16**, e0258670 (2021).

1066 20. Lu, Q. *et al.* Syntaxin 1A supports voltage-dependent inhibition of alpha1B Ca2+
1067   channels by Gbetagamma in chick sensory neurons. *J Neurosci* **21**, 2949-2957
1068   (2001).
1069 21. Bezprozvanny, I., Scheller, R.H. & Tsien, R.W. Functional impact of syntaxin on
1070   gating of N-type and Q-type calcium channels. *Nature* **378**, 623-626 (1995).
1071 22. Weiss, N. & Zamponi, G.W. Regulation of voltage-gated calcium channels by
1072   synaptic proteins. *Adv Exp Med Biol* **740**, 759-775 (2012).
1073 23. Pozzi, D., Corradini, I. & Matteoli, M. The Control of Neuronal Calcium
1074   Homeostasis by SNAP-25 and its Impact on Neurotransmitter Release.
1075   *Neuroscience* **420**, 72-78 (2019).
1076 24. Catterall, W.A. & Few, A.P. Calcium channel regulation and presynaptic
1077   plasticity. *Neuron* **59**, 882-901 (2008).
1078 25. He, Y. *et al.* Modulation of Kv2.1 channel gating and TEA sensitivity by distinct
1079   domains of SNAP-25. *Biochem J* **396**, 363-369 (2006).
1080 26. Cormet-Boyaka, E. *et al.* CFTR chloride channels are regulated by a SNAP-
1081   23/syntaxin 1A complex. *Proc Natl Acad Sci U S A* **99**, 12477-12482 (2002).
1082 27. Grimm, C. *et al.* High susceptibility to fatty liver disease in two-pore channel 2-
1083   deficient mice. *Nat Commun* **5**, 4699 (2014).
1084 28. Lin-Moshier, Y. *et al.* The Two-pore channel (TPC) interactome unmasks
1085   isoform-specific roles for TPCs in endolysosomal morphology and cell
1086   pigmentation. *Proc Natl Acad Sci U S A* **111**, 13087-13092 (2014).
1087 29. Toft-Bertelsen, T.L., Ziomkiewicz, I., Houy, S., Pinheiro, P.S. & Sorensen, J.B.
1088   Regulation of Ca2+ channels by SNAP-25 via recruitment of syntaxin-1 from
1089   plasma membrane clusters. *Mol Biol Cell* **27**, 3329-3341 (2016).
1090 30. Nieto-Rostro, M., Ramgoolam, K., Pratt, W.S., Kulik, A. & Dolphin, A.C. Ablation
1091   of alpha2delta-1 inhibits cell-surface trafficking of endogenous N-type calcium
1092   channels in the pain pathway in vivo. *Proc Natl Acad Sci U S A* **115**, E12043-
1093   E12052 (2018).
1094 31. Mohrmann, R., de Wit, H., Verhage, M., Neher, E. & Sorensen, J.B. Fast vesicle
1095   fusion in living cells requires at least three SNARE complexes. *Science* **330**, 502-
1096   505 (2010).
1097 32. Sinha, R., Ahmed, S., Jahn, R. & Klingauf, J. Two synaptobrevin molecules are
1098   sufficient for vesicle fusion in central nervous system synapses. *Proc Natl Acad
1099   Sci U S A* **108**, 14318-14323 (2011).
1100 33. Vig, M. & Kinet, J.P. Calcium signaling in immune cells. *Nature immunology* **10**,
1101   21-27 (2009).
1102 34. Jahn, R. & Scheller, R.H. SNAREs--engines for membrane fusion. *Nat Rev Mol
1103   Cell Biol* **7**, 631-643 (2006).
1104 35. Tang, B.L., Low, D.Y. & Hong, W. Syntaxin 11: a member of the syntaxin family
1105   without a carboxyl terminal transmembrane domain. *Biochem Biophys Res
1106   Commun* **245**, 627-632 (1998).
1107 36. Dabrazhynetskaya, A. *et al.* Syntaxin 11 marks a distinct intracellular
1108   compartment recruited to the immunological synapse of NK cells to colocalize
1109   with cytotoxic granules. *J Cell Mol Med* **16**, 129-141 (2012).
1110 37. Zhang, G. *et al.* The vesicular transporter STX11 governs ATGL-mediated
1111   hepatic lipolysis and lipophagy. *iScience* **25**, 104085 (2022).

38. Yuan, J.P. *et al.* SOAR and the polybasic STIM1 domains gate and regulate Orai
channels. *Nature cell biology* **11**, 337-343 (2009).
39. Li, Z. *et al.* Graded activation of CRAC channel by binding of different numbers of
STIM1 to Orai1 subunits. *Cell Res* **21**, 305-315 (2011).
40. Park, C.Y. *et al.* STIM1 clusters and activates CRAC channels via direct binding
of a cytosolic domain to Orai1. *Cell* **136**, 876-890 (2009).
41. Hou, X., Burstein, S.R. & Long, S.B. Structures reveal opening of the store-
operated calcium channel Orai. *Elife* **7** (2018).
42. Yeung, P.S. *et al.* Mapping the functional anatomy of Orai1 transmembrane
domains for CRAC channel gating. *Proc Natl Acad Sci U S A* **115**, E5193-E5202
(2018).
43. Zhang, K. *et al.* Familial Hemophagocytic Lymphohistiocytosis, in
GeneReviews((R)). (eds. M.P. Adam *et al.*) (Seattle (WA); 1993).
44. Hou, X., Pedi, L., Diver, M.M. & Long, S.B. Crystal structure of the calcium
release-activated calcium channel Orai. *Science* **338**, 1308-1313 (2012).
45. Jessen, B. *et al.* Graded defects in cytotoxicity determine severity of
hemophagocytic lymphohistiocytosis in humans and mice. *Front Immunol* **4**, 448
(2013).
46. Chinn, I.K. *et al.* Genetic and mechanistic diversity in pediatric hemophagocytic
lymphohistiocytosis. *Blood* **132**, 89-100 (2018).
47. Zhizhuo, H. *et al.* Screening the PRF1, UNC13D, STX11, SH2D1A, XIAP, and
ITK gene mutations in Chinese children with Epstein-Barr virus-associated
hemophagocytic lymphohistiocytosis. *Pediatr Blood Cancer* **58**, 410-414 (2012).
48. Hellewell, A.L., Foresti, O., Gover, N., Porter, M.Y. & Hewitt, E.W. Analysis of
familial hemophagocytic lymphohistiocytosis type 4 (FHL-4) mutant proteins
reveals that S-acylation is required for the function of syntaxin 11 in natural killer
cells. *PLoS One* **9**, e98900 (2014).
49. Hackmann, Y. *et al.* Syntaxin binding mechanism and disease-causing mutations
in Munc18-2. *Proc Natl Acad Sci U S A* **110**, E4482-4491 (2013).
50. Neveu, E., Khalifeh, D., Salamin, N. & Fasshauer, D. Prototypic SNARE Proteins
Are Encoded in the Genomes of Heimdallarchaeota, Potentially Bridging the Gap
between the Prokaryotes and Eukaryotes. *Curr Biol* **30**, 2468-2480 e2465 (2020).

1144

1145

1146

1147

Acknowledgements: We thank Vivien Beziat, Yenan Bryceson, Heinrich Schlums, Jelve

Nejati Zendegani, Stephan Ehl, and Jasmin Mann for help with human PBMC handling,

isolation, storage and shipping. Anand Vaidya for advice on STX11 purification. Jyoti

Rohilla for technical support. This work was supported in part by Department of Atomic

Energy, Government of India under Project Identification No. RTI 4007 and NIH-NIAID

grant AI108636.

Figure 1

Figure 1: STX11 is required for SOCE.

(A-F) RNAi mediated depletion of STX11 reduces SOCE. Measurement of thapsigargin (TG) induced SOCE in various cell lines after shRNA mediated depletion of STX11. The traces in panels A,C,E show representative average single cell Fura-2 calcium imaging assays. Bars in panels B,D,F show mean % SOCE   SE from three independent experiments each where mean SOCE from scramble (scr) shRNA treated group in each experiment was set at 100% and the relative response of STX11 shRNA treated groups was calculated respectively. *P<0.05; **P<0.01; ***P<0.001 using two-tailed Student's t test. (G) Representative Fura-2 calcium imaging assay showing reconstitution of SOCE in STX11 depleted Jurkat T cells by ectopic expression of STX11. Black (scr shRNA), red (STX11 shRNA), green (STX11 shRNA with STX11 expression). N=2 (H) Ectopic expression of STX11 enhances SOCE. Representative Fura-2 calcium imaging assay showing SOCE in HEK293 cells expressing STX11 (red) or empty vector (EV) (black). (N=3) (I) STX11 mediated enhancement of SOCE is dependent on Orai1. A representative Fura 2 calcium imaging assay showing measurement of thapsigargin (TG) induced SOCE in HEK293 cells where Orai1 expression was depleted using shRNA and STX11 was over-expressed. (N=2)

Figure 2

Figure 2: STX11 depletion suppresses ICRAC, downstream signaling and gene expression in Jurkat T cells. ICRAC was recorded from Jurkat T-cells in the whole-cell recording configuration in 20 mM extracellular Ca2+ Ringer's solution. ICRAC was induced by passive depletion of intracellular Ca2+ stores by dialyzing 8 mM BAPTA into the cell via the patch-pipette. (A) Representative current at -100 mV in Jurkat T cell transfected with scr shRNA construct. The current is blocked by extracellular La3+ (100 μM) and replacing the 20 mM Ca2+ Ringer's solution with a divalent free solution (DVF) evokes a large Na+ current which depotentiates over tens of seconds. The current-voltage (I-V) relationship of the Ca2+ and DVF currents are shown on the right. (B) ICRAC from a Jurkat T cell transfected with STX11 shRNA. Both Ca2+ and Na+ current amplitudes are reduced relative to control cells. The I-V relationships (right plots) show no change in ion selectivity. (C-D) Summary of the current amplitudes of Ca2+ and Na+ currents and current reversal potentials in scr and STX11 knockdown cells. (E&F) Estimation of nuclear translocation of NFAT. (E) Western Blot showing nuclear translocation of NFAT in Jurkat T cells treated with scr or STX11 shRNA for 4 days and stimulated with PMA+TG for 30min prior to the preparation of nuclear and cytoplasmic extracts. (N=3) (F) Representative confocal images of Jurkat T cells treated with scr or STX11 shRNA for 4 days and stimulated with 10ug/ml anti-CD3 for 1 hour. Following stimulation, cells were fixed, permeabilized and stained using anti-NFAT primary antibody, followed by donkey anti-rabbit AF647 secondary antibody, and counter-stained with DAPI to mark the nuclei. (N=2) (G) Box and whisker plot showing percent nuclear NFAT in Jurkat T cells quantified from 40-50 cells populating 10 randomly chosen fields per group in (F). Boundaries of the box plots represent 25th and 75th percentile values, horizontal line represents mean, white circle represents median and whiskers denote the outliers. (H) Quantitative PCR to assess IL-2 transcription in anti-CD3 stimulated Jurkat T cells. Jurkat T cells were treated with scr or STX11 shRNA for 4 days and stimulated with 5ug/ml anti-CD3 for 3 hours. Total RNA was extracted from cells and subjected to QPCR analysis using Taqman probes for IL-2 and beta-actin. The bars show relative IL-2 mRNA expression levels with the scr shRNA treated group set at 100%. Shown here are mean SE. (N=3) (F-H) *P<0.05; **P<0.01; ***P<0.001 using two-tailed Student's t test.

Figure 3

Figure 3: STX11 directly binds resting Orai1 in the plasma membrane. (A) STX11 localizes in the plasma membrane. Representative confocal images of HEK293 cells expressing internal-HA-tagged STX11, stained using anti-HA antibody followed by alpaca VHH anti-rabbit secondary nanobody. Scale bar 15um. (N=3) (B) Co-localization of STX11 and Orai1 in the plasma membrane. Representative confocal images of HEK293 cells expressing Orai1-YFP and STX11, stained using anti-STX11 antibody followed by donkey anti-rabbit secondary antibody. Scale bar 10um. (N=3). (C&D) Co-immunoprecipitation of STX11 with Orai1. Whole cell lysates of resting and store-depleted HEK293 cells expressing either Flag-Orai1 and STX11 (C) or STX11 and Orai1-Myc-His (D) were subjected to immunoprecipitation and Western Blot using anti-Myc, anti-Flag or anti-STX11 antibodies, as indicated. (N=3) (E) Schematic showing key domains of STX11 and Orai1 used for in vitro pulldown assays. (F) Pull-down assay showing in vitro binding of His-tagged full length STX11 to MBP-tagged cytosolic domains of Orai1. MBP-tagged Orai1 fragments, expressed in E. Coli and immobilized on the amylose resin, were incubated with purified His-tagged STX11 protein. Post incubation, beads were washed, boiled and subjected to Western Blot analysis using anti-STX11 antibody. (Top panel) Ponceau S staining showing the input of MBP alone or MBP-tagged fragments. (Bottom panel) Western Blot using anti-STX11 antibody. (N=3) (G) Pull-down assay showing in vitro binding of His-tagged Habc domain of STX11 to MBP-tagged Orai1 N- and C-termini performed as described above. (Top panel) Ponceau S staining showing the input of MBP alone or MBP-tagged Orai1 cytosolic tails. (Bottom panel) Western Blot using anti-His antibody. (N=3). (H-I) Representative structure of STX11 Habc and Orai1 C-terminus complex after MD simulation (H) Sphere representation of STX11 Habc (green) and Orai1 C-terminus (cyan) complex. The N-termini are highlighted in blue, and C-termini are highlighted in red (I) Cartoon representation of STX-11-Habc (green) and Orai1 C-terminus (cyan) complex highlighting the CA of terminal residues as spheres (J) Protein-protein interactions between the STX11 Habc and the Orai1 C-terminus.

Figure 4

Figure 4: STX11 depletion compromises the functional assembly of Orai1 with Stim1 in ER-PM junctions.
(A-B) Representative confocal images of resting (A) and store-depleted (B) scr and STX11 shRNA treated HEK293 cells expressing N-terminal CFP tagged Orai1 (CFP-Orai1) and C-terminal YFP tagged Stim1 (Stim1-YFP). (C-H) Box and whisker plots showing (C) Quantification of Stim1-YFP intensities inside Stim:Orai clusters of scr and STX11 shRNA treated HEK293 cells, ~8 min post store-depletion. (D-E) Quantification of CFP-Orai1 intensities inside (D) and outside (E) Stim1-YFP clusters of scr and STX11 shRNA treated cells, ~8 minute post store-depletion. N=3. Boundaries of the box plots represent 25th and 75th percentile values, horizontal line represents mean, white circle median and whiskers denote the outliers. (F) Quantification of total Orai1 levels in the plasma membrane of STX11 depleted cells. U2OS cells stably expressing Orai1-BBS-YFP were transduced with scr (black) or STX11 (red) shRNA, stimulated with 1uM TG, incubated with alpha-bungarotoxin alexa 647 (BTX-A647) and washed. BTX binding to surface Orai1 was measured using FACS, where binding to wildtype HEK293 cells was used as control. (N=3). (G-J) Quantification of C-term tagged Orai1 (Orai1-YFP) and N-term tagged Stim1 (CFP-Stim1) inside puncta in control and STX11 depleted cells, ~6 minute post store-depletion. (G-H) Quantification of mean intensities of Orai1-YFP (G) and CFP-Stim1 (H) inside puncta. n=8. Quantification of mean area of Orai1-YFP (I) and CFP-Stim1 (J) puncta. The continuous and dotted lines represent mean and median respectively. *P<0.05, **P<0.01 and ***P<0.001 using two-tailed Student's t test. (K) Representative Fura-2 calcium imaging assay to measure Thapsigargin induced SOCE in Orai1-YFP and CFP-Stim1 expressing HEK293 cells treated with scr or STX11 shRNA.

Figure 5

Figure 5: STX11 binding to resting Orai1 allows successful gating by Stim1. (A) Representative Fura-2 calcium imaging assay to measure constitutive calcium influx in Orai1-S-S-GFP expressing control or STX11 depleted HEK293 cells. Cells were imaged in Ringer's buffer containing 0mM followed by 2mM extracellular Ca^{2+}. (B) Box and whisker plot showing quantification of constitutive calcium influx across experiments as shown in (A). N=3. (C) Repre­sentative images of YFP-CAD localization in Orai1-CFP expressing, control and STX11 depleted HEK293 cells. (D) Quantification of YFP-CAD in the plasma membrane of Orai1-CFP expressing HEK293 cells represented as % of total YFP-CAD in respective cells. (E) Representative Fura-2 calcium imaging assay to measure constitutive calcium influx in Orai1-CFP and YFP-CAD expressing control or STX11 depleted HEK293 cells. (F-G) Quantification of constitutive calcium influx in Orai1-CFP and YFP-CAD expressing cells described in E, at 0mM (F) and 2mM (G) extracellular Ca^{2+}. N=3. (H) Representative Fura-2 calcium imaging assay to measure constitutive calcium influx in Orai1-H134S mutant expressing control and STX11 depleted HEK293 cells as described in (A). (I) Quantification of constitutive calcium influx across experiments as shown in (H). N=3. Boundaries of the box plots in B,D,F,G&I repre­sent 25th and 75th percentile values, horizontal line represents mean, white circle median and whiskers denote the outliers. *$P<0.05$, **$P<0.01$ and ***$P<0.001$ in (C-D,F-G,I) using two-tailed Student's t test.

Figure 6

Figure 6: Ionomycin rescues the cytotoxicity and gene expression defects in STX11 deficient FHLH4 patient T lymphocytes. (A) Sanger sequencing of the patient DNA showing deletion of a single of Adenine at the 752nd position and the resulting frameshift. (B) Western Blot of whole cell lysates prepared from healthy human control and FHLH4 patient PBMCs showing the relative molecular weight and abundance of the wildtype and mutant STX11 bands. (C) Fura-2 calcium imaging assay measuring anti-CD3 induced SOCE in wildtype and FHLH4 PBMCs. (D) Fura-2 calcium imaging assay measuring SOCE in wildtype and FHLH4 PBMCs expressing either empty vector (EV) or wildtype human STX11, respectively. (E) Representative Fura-2 calcium imaging assay measuring SOCE in wildtype PBMCs with or without ectopic expression of wildtype STX11. (F) Bar plot showing quantification of SOCE across multiple experiments in (E). N=3. (G) Granule release assay performed on the in vitro cultured healthy human control and FHLH4 patient CD8 T cells. PBMCs isolated from control and FHLH4 patient blood were stimulated with (2ug/ml) PHA for 48 hours. On the day of assay, cells were washed and either left unstimulated or restimulated with plate coated anti-CD3 + soluble anti-CD28 or PMA + Ionomycin in the presence of CD107a-PE antibody, stained with anti-CD8 antibody and analyzed. (N=2) (H) Quantification of the granule release assays as shown in (G). (I) Analysis of IL-2 expression in control and FHLH4 T cells. In vitro cultured healthy human control and FHLH4 patient T cells were rested and stimulated with either plate coated anti-CD3 + soluble anti-CD28 or PMA + Ionomycin for 6 hours. Total RNA was extracted and subjected to quantitative PCR analysis using Taqman probes for IL-2 and beta-actin in triplicates. *P<0.05; **P<0.01; ***P<0.001 using two-tailed Student's t-test.

Figure 7

Figure 7: Hypothetical model showing Orai1 bound to STX11 for its priming. Blue ribbons depict Orai1 in plasma membrane (PM). Green and Orange ribbons represent Habc and SNARE domains of STX11, respectively. Also shown are the C-terminal cysteines of STX11 which facilitate membrane attachment of the protein.