Syntaxin11 Deficiency Inhibits CRAC Channel Priming to Suppress Cytotoxicity and Gene Expression in FHLH4 Patient T Lymphocytes.

Sritama Datta[1,#], Abhikarsh Gupta[1,#], Kunal Mukesh Jagetiya[1], Atharva Rahul Yande[1], Vikas Tiwari[2,⊥], Megumi Yamashita[3,⊥], Resmi Bera[1], Sandra Ammann[4,5], Mohammad Shahrooei[6], Ramanathan Sowdhamini[2], Murali Prakriya[3], Adish Dani[1], Monika Vig[1]*

Address and Affiliation: [1]Tata Institute of Fundamental Research, Hyderabad, India. [2]National Centre for Biological Sciences, Bangalore, India. [3]Northwestern University, Feinberg School of Medicine, Chicago, USA. [4]Institute for Immunodeficiency, University of Freiburg, Germany. [5]Institute for Transfusion Medicine and Gene Therapy, University of Freiburg, Germany. [6]Department of Microbiology, Immunology and Transplantation, Clinical and Diagnostic Immunology, KU Leuven, Leuven, Belgium.

[#,⊥] These authors contributed equally.

*able**Correspondence to:** monika.vig@gmail.com

Keywords: Orai, FHLH4, CRAC, Syntaxin11, SOCE, Stim, SNARE, ion channels, autoimmunity, cytotoxicity, CTL, T lymphocytes

Abstract:

Mutations in Syntaxin11, a Q-SNARE, result in a fatal immune disorder known as familial hemophagocytic lymphohistiosis 4 (FHLH4) in human patients. A key diagnostic feature of FHLH4 is defective T and natural killer cell cytotoxicity. Here we show that Syntaxin11 directly binds and regulates Orai1, the pore forming subunit of calcium release activated calcium (CRAC) channels. CRAC channels enable store-operated calcium entry (SOCE) from the extracellular space and are crucial for granule exocytosis and nuclear factor of activated T cells (NFAT) dependent gene expression in activated lymphocytes. Syntaxin11 depletion strongly inhibited SOCE, CRAC currents, NFAT activation, interleukin-2 gene expression and degranulation in FHLH4 patient T lymphocytes and cell lines. Remarkably, defects of granule exocytosis as well as interleukin-2 expression could be reversed by ionomycin induced calcium influx in patient T lymphocytes and a constitutively active, H134S, mutant of Orai1 rescued calcium entry in Syntaxin11 depleted cells. Further mechanistic analyses showed that Syntaxin11 primes Orai1 subunits for correct on-site assembly which is a prerequisite for functional entrapment and gating of CRAC channels by the endoplasmic reticulum resident Stim1. We propose that priming of ion channel pore subunits is a primary and conserved function of specific SNAREs which likely preceded their role in vesicle fusion.

Introduction:

Familial hemophagocytic lymphohistiocytosis 4 (FHLH4) is a life-threatening
immune disease caused due to mutations of STX11 [1]. The underlying cause of FHLH
is reduced cytolytic activity of T and NK cells [2], which renders patients susceptible to
recurrent infections. Patients suffer from high fever, severe lymphopenia in early infancy
and succumb to disease by adolescence, unless given bone marrow transplants. STX11
is highly expressed in almost all cells of the immune system and, being a Q-SNARE, is,
by default, associated with exocytosis of vesicles. Intriguingly, FHLH4 patients also
manifest cytokine storm, which cannot be reconciled with a presumed general defect in
membrane trafficking and vesicle fusion.

Most eukaryotic cells have a limited amount of calcium sequestered inside
intracellular stores, the largest of which is the endoplasmic reticulum (ER). Signaling from
cell surface receptors induces the release of stored calcium which, in turn, activates store-
operated calcium entry (SOCE) to replenish stores, sustain signaling and drive other
calcium dependent cellular processes [3]. Calcium release activated calcium (CRAC)
channels are the major conductors of SOCE [4]. In lymphocytes and mast cells, CRAC
currents have been shown to be crucial for cytotoxicity, degranulation as well as gene
expression [5, 6]. Working in concert with a number of accessory proteins, Orai (CRACM)
multimers conduct SOCE upon functional clustering with Stim proteins in ER-PM
junctions [7-9]. Yet, the prevailing thought attributes all structural transitions that result in
the activation of CRAC currents to Stim proteins [10].

Genome-wide RNAi screens, which initially identified Orai (CRACM), have also
yielded a wealth of information regarding additional crucial players in SOCE [8, 11, 12].
For instance, we have previously shown that alpha-soluble N-ethylmaleimide sensitive
factor (NSF) attachment protein (α-SNAP), a well-known synaptic family adaptor protein
that forms a part of the 20S SNARE super-complex [13], is a crucial component of the
CRAC channel supramolecular complex [8, 9]. α-SNAP independently binds Stim as well

as Orai and is required for the on-site assembly and calcium selectivity of CRAC channels

[14-16]. α-SNAP depletion in cells revealed that Stim:Orai coupling is necessary but not

sufficient for SOCE, a process that requires distinct time-resolved molecular steps *in vivo*.

In addition to α-SNAP, some SNAREs also came up as candidates in two genome

wide RNAi screens for regulators of SOCE but were not characterized [8, 12].

Furthermore, specific SNAREs have been previously found to form inhibitory associations

with some ion channels, but their exact role was not defined [17]. Therefore, we

hypothesized a more fundamental function for the association of SNAREs with ion

channel pore subunits and set out to test this in the context of SOCE. Using a targeted

RNAi screen, biochemical and cell biological analyses, we have found a direct role of

STX11 in the correct, on-site, assembly of CRAC channel pore subunit, Orai1, for

subsequent gating by Stim1. Analyses of T cells derived from a symptomatic FHLH4

patient showed that the previously reported [2] as well as newly identified defects in

degranulation and interleukin-2 gene expression predominantly stem from suppressed

SOCE. Based on the findings presented here, we demonstrate a novel SNARE

dependent priming step for the CRAC channel pore in the molecular sequence of SOCE

and suggest preventive pharmacological interventions for FHLH4 patients.

Results:

RNAi dependent ablation of a Q-SNARE, Syntaxin 5 (STX5), has earlier shown inhibition of SOCE in a genome-wide screen performed in *Drosophila* S2 cells [12] and STX1A has been previously shown to bind a variety of ion channels [17]. Therefore, we first knocked down STX5 (Supplementary Figure 1A) and STX1A (Supplementary Figure 1B) using different sequences of shRNA targeting each gene in HEK293 cells and measured SOCE in response to Thapsigargin (TG). In both cases, we did not see a significant defect in SOCE. Since several proteins involved in synaptic function are not ubiquitously expressed and many show redundancy in function, we searched online databases to target SNAREs that are expressed in HEK293 and/or T cells (Supplementary Figure 1C-H). We found that Syntaxin11 knockdown showed strong and consistent inhibition of SOCE by day 4 (Supplementary Figure 2A-B) (Figure 1A-B). Quantitative PCR (qPCR) analysis of total RNA extracted from scramble (scr) and STX11 shRNA treated cells showed nearly 70% depletion of STX11 mRNA (Supplementary Figure 2C) and ectopic expression of STX11 in STX11 depleted Jurkat cells rescued SOCE (Figure 1C&D). The efficiency of STX11 depletion was assessed for each experiment reported in this study. We restricted our analysis to day 3 to 4 post RNAi treatment as longer incubations with RNAi or CRISPR knockout of STX11 affected cell viability.

Given that STX11 is highly expressed in most immune cells and is the only known SNARE that, when mutated, causes defects in T and NK cell cytotoxicity in humans and mice [2], we hypothesized that reduced SOCE contributes to the cytotoxicity defects of FHLH4 patient T cells. To test this hypothesis, we isolated PBMCs from an FHLH4 patient with a homozygous deletion frameshift mutation in the STX11 coding sequence, c.752delA:p.Lys251fs. The patient presented with all symptoms of FHLH4 disease by 4 years of age. Given the limited sample amount, we expanded PBMC numbers by periodically stimulating patient and healthy donor cells with PHA and IL-2, prior to analysis. Isolation and Sanger sequencing of DNA from patient PBMCs showed deletion of a single Adenine at position 752 in the STX11 gene which lead to frameshift following Lysine 251 (Figure 1E). The absence of any contaminating wildtype peaks showed that

the mutation was present in all blood cells. Figure 1F shows schematic of the predicted mutant STX11 protein with an estimated molecular weight of ~39.5 KDa compared to ~33 KDa of wildtype STX11. Frameshift mutation resulted in the disruption of SNARE domain, ablation of the terminal cysteines and elongation of the resulting transcript. We subjected whole cell lysates prepared from the FHLH4 patient and healthy donor (HD) PBMCs to SDS-PAGE and western blot and found that while the wildtype STX11 band was absent, a faint but distinct band running at ~37 KDa was present in the FHLH4 patient sample lysate (Figure 1G and Supplementary Figure 2D). Therefore, the predicted and observed molecular weight of the FHLH4 mutant protein was higher but its levels appeared significantly reduced compared to STX11 from HD. This is likely due to protein instability and degradation because ectopic expression of the FHLH4 mutant in HEK293 cells showed at least two clear bands (Figure 1H and Supplementary Figure 2E), although reduced reactivity to anti-STX11 antibody is also a possibility. In accordance with our findings in the STX11 depleted cell lines (Figure 1A-B and Supplementary Figure 2A-B), SOCE was found to be significantly defective in FHLH4 T cells when compared to HD (Figure 1I). Further, expression of wildtype (Figure 1J), but not FHLH4 mutant STX11 (Figure 1K-L), reversed defective SOCE in STX11 deficient patient (Figure 1J) and Jurkat T cells (Figure 1K-L), respectively. These data suggest that FHLH4 mutant STX11 was functionally compromised in addition to being unstable and no additional unknown mutations resulted in reduced SOCE.

Overexpression of STX1A has been earlier shown to inhibit voltage gated calcium channels (VGCCs) [18, 19]. Since STX11 depletion inhibited SOCE, we next sought to determine the effect of STX11 overexpression. We ectopically expressed STX11 in healthy donor PBMCs (Figure 2A-B), HEK293 and Jurkat T cells (Supplementary Figure 3A-B) and measured SOCE. In each of these cases, there was an increase in SOCE, reinforcing a positive role of STX11 in SOCE although ectopic expression of FHLH4 STX11 mutant failed to augment SOCE (Supplementary Figure 2F). To determine whether the inhibition in SOCE (Figure 1) and its augmentation (Figure 2A-B) resulted from changes in the activity of CRAC channels, we depleted Orai1 from STX11 over-expressing HEK293 cells, which resulted in complete loss of enhancement observed in

SOCE (Figure 2C&D). To quantify the level of Orai1 expression in plasma membrane, we

labelled a previously characterized U2OS cell line stably expressing Orai1 tagged with a

bungarotoxin binding site (BBS) and YFP (Orai-BBS-YFP) with Alexa-647 conjugated

alpha-bungarotoxin (BTX-A647) (Supplementary Figures 3C) [9]. There was no change

in the expression of total Orai1 in the PM of STX11 over-expressing (Supplementary

Figure 3D) or STX11 depleted cells (Supplementary Figure 3E), even though SOCE was

significantly suppressed in these U2OS cells (Supplementary Figures 3F-G). The ER-

targeted CEPIA expressing control and STX11 depleted cells showed comparable ER

calcium content (Supplementary Figure 3H). Labelling STX11 depleted cells with ER,

Golgi organelle specific markers also showed no abnormalities (Supplementary Figures

3I&J) suggesting that the overall health, function and calcium content of Golgi and ER

were not adversely affected by STX11 depletion.

Previous studies have shown that SOCE in T cells is mediated by CRAC channels

formed by Orai proteins. To test whether CRAC current (I_{CRAC}) was affected by

knockdown of STX11 in T cells, we performed whole cell patch clamp recordings on

scramble and STX11 shRNA treated Jurkat T cells (Figure 2E-G). Leak-subtracted I_{CRAC}

was recorded in 20 mM Ca^{2+} and a Na^+-based divalent cation free (DVF) solution. These

recordings showed that both the Ca^{2+} and DVF CRAC currents were significantly

decreased in the STX11 shRNA treated cells with the normalized currents shown in

Figure 2E-G. The electrophysiological properties of the residual current in STX11 shRNA

treated cells was indistinguishable from I_{CRAC} in scramble shRNA treated cells in terms of

blockade of the Ca^{2+} current by La^{3+} and depotentiation of the DVF current over tens of

seconds. Moreover, the reversal potential of the Ca^{2+} and DVF currents were similar

between scramble and STX11 shRNA treated Jurkat T cells, suggesting that STX11

depletion does not affect the calcium selectivity of CRAC channels (Figure 2H). Together,

these results demonstrate that STX11 works by specifically modulating I_{CRAC} in T cells.

CRAC channel mediated sustained calcium influx is crucial for the activation of a

key calcium dependent transcription factor known as nuclear factor of activated T cells

(NFAT) [4, 20]. We stimulated control and STX11 depleted Jurkat T cells with

Thapsigargin (TG)+PMA, prepared nuclear, cytosolic extracts and subjected them to
western blot using anti-NFAT1 antibody (Figure 3A). Nuclear translocation of NFAT was
severely compromised in STX11 depleted T cells. In addition, immunofluorescence
imaging and quantification of total *versus* nuclear NFAT from anti-CD3 stimulated Jurkat
cells showed a defect in nuclear translocation of NFAT1 in STX11 depleted cells (Figure
3B-C). To determine whether NFAT induced gene expression was affected [4, 20], we
quantified IL-2 mRNA by qPCR and secreted IL-2 levels from culture supernatants of anti-
CD3 stimulated Jurkat T cells. STX11 depleted cells showed a significant decrease in IL-
2 gene expression as well as secretion (Figure 3D, 3E). We stimulated HD and FHLH4
patient T cells either through T cell receptor using anti-CD3 or with phorbol myristate
acetate (PMA) + Ionomycin and assessed IL-2 gene expression. Importantly, induction of
IL-2 expression was significantly defective in FHLH4 patient T cells but could be largely
restored with Ionomycin (Figure 3F). We next performed granule release assay on the *in*
vitro cultured HD and FHLH4 patient CD8 T cells in response to anti-CD3 mediated
stimulation and observed a significant defect in the FHLH4 patient CD8 T cell
degranulation (Figure 3G, 3H). To determine whether the reduced granule release results
from a direct defect in vesicle fusion or SOCE, we stimulated cells with Ionomycin along
with (PMA). Remarkably, CD8 T cell degranulation was fully restored in FHLH4 T cells
suggesting that defective SOCE primarily causes cytolytic defects in FHLH4 T cells. Of
note, the patient exhibited all symptoms of FHLH by 4 years of age and we observed
strong suppression of SOCE (Figure 1I), IL-2 gene expression and secretion (Figure 3D-
F). However, *in vitro* culture of FHLH4 patient CD8 T cells with IL-2 has been previously
shown to partially or fully restore cytolytic defects by other groups [21, 22]. Therefore,
considering previous reports, a relatively lower (50%) defect in CD8 T cells degranulation
was not surprising [21, 22]. Collectively, these data suggest that defects in T cell
degranulation as well as interleukin-2 gene expression originate from suppressed SOCE
and significantly contribute to the immune dysregulation of FHLH4 patients.

Unlike most Q-SNAREs, STX11 lacks a transmembrane domain but harbors a
stretch of cysteines close to the C-terminus (Supplementary Figure 4A), which have been
proposed to be essential for its membrane localization [23]. The cellular localization of

STX11 is not established primarily because, unlike primary T cells, native STX11 is undetectable in Jurkat and HEK293 cell lines using available commercial antibodies and tagging STX11 on either N- or C-terminus result in partial degradation (Supplementary Figure 4B) and complete mis-localization of the protein (Supplementary Figure 4C). Therefore, we inserted HA tag inside the N-terminal unstructured region of STX11, expressed in HEK cells, permeabilized and labelled with anti-HA tag antibody. The HA tagged human STX11 localized in the plasma membrane (Supplementary Figure 4D). We next expressed untagged human STX11 in cells co-expressing Orai1-YFP, permeabilized and stained with anti-STX11 antibody. In most cells, STX11 co-localized with Orai1 in plasma membrane (Figure 4A). To test whether STX11 and Orai1 formed a complex, we co-expressed STX11 with either Flag- or Myc-tagged Orai1 in HEK293, prepared whole cell lysates (WCLs) and subjected them to co-immunoprecipitation (co-IP) followed by western blot using either anti-Flag, anti-Myc or anti-STX11 antibodies. Both Orai1 and STX11 could co-IP each other (Figure 4B-C) (Supplementary Figure 5A-B). To further assess whether STX11 directly bound Orai1, we expressed and purified from *E. coli*, MBP-tagged N- and C-terminal cytosolic tails of Orai1 (Figure 4D). *In vitro* pull-down assay performed by incubating MBP-tagged Orai1 cytoplasmic tails with full length soluble His-tagged STX11 showed that the C-terminus of Orai1 directly bound STX11 (Figure 4E) (Supplementary Figure 5C). SNAREs typically utilize their SNARE domain for associating with other SNAREs [24]. To identify the domain of STX11 involved in the regulation of SOCE, we expressed His-tagged H_{abc} and SNARE domains of STX11 in *E. coli*, purified and assessed their binding to the MBP-tagged Orai1 tails using pull-down assay. We found that the H_{abc} domain of STX11 bound to the C-terminus of Orai1 (Figure 4F) (Supplementary Figure 5D). The SNARE domain did not show any binding to Orai1 and only a faint binding to SNAP23 (Supplementary Figure 5E). We next used AlphaFold3 (AF3) to obtain deeper insights into the domain-domain interactions of STX11 and Orai1. Of all the possible combinations, the complex of STX11 H_{abc} and Orai1 C-terminus resulted in a significantly high prediction score (Supplementary Table 1). We, therefore, executed AF3 predictions with different initial seeds to generate multiple models of STX11 H_{abc} with Orai1 C-terminus. We calculated the contact frequency of interface residues in different models and found that multiple residues showed frequency greater than 0.8

(Supplementary Figure 6). For further analysis, we considered the predicted model with

the best score (Supplementary Table 2). Next, we performed all-atom MD simulation in

aqueous environment (Movies 1-3) to assess the conformation, interaction stability of the

STX11 H_{abc} and Orai1 C-terminus complex, which was stable throughout the simulation

time as assessed through RMSD (Supplementary Figure 7), and binding free energy (ΔG)

(Supplementary Figure 8A) (Figure 4G). The individual trajectories were concatenated

and clustered to obtain a centroid structure for this complex. There were 46 clusters, and

the largest cluster, with 52 members, showed an elaborate protein-protein interface where

STX11 H_{abc} was found to be oriented in an anti-parallel orientation to Orai1 C-terminus

(Figure 4H).

Because STX11 binds the C-terminus of Orai1 and the region largely overlaps with

Stim1 binding domain, we next sought to determine whether STX11 co-localizes with

Orai:Stim puncta. We over-expressed untagged STX11 in a Orai1-YFP and CFP-Stim1

expressing stable HEK293 cell line, store-depleted with TG and stained using anti-STX11

antibody. We did not observe STX11 in Orai:Stim puncta (Figure 4I). Accordingly, STX11

did not co-IP with Stim1 (Supplementary Figure 9A). These data suggest that STX11

comes off Orai1 C-terminus prior to Stim1 dependent clustering of Orai1. Since co-IP

studies suggested sustained interaction between STX11 with Orai1 (Figure 4B-C), we

quantified the fraction of Orai1 in Stim:Orai puncta w.r.t total Orai1 in PM and found that

nearly 60% Orai1 remains outside of Stim1 puncta in unmanipulated cells, even 8 minutes

post TG addition (Supplementary Figure 9B). We next performed all-atom MD simulations

on Orai1 C-terminus (256-301) and the Orai activation domain of human Stim1 (312-387),

previously studied in an NMR study [25] (Supplementary Figure 9C). We used AF3 to

generate the complex of Stim1 and Orai1. The single copy as well as two copies of Stim1

with monomeric Orai1 resulted in low prediction scores (Supplementary Table 3-4).

Nonetheless, ΔG values between the protomers of Stim1 and Orai1 (Figure 4J)

(Supplementary Figure 9D) were comparable to that of STX11 and Orai1 (Figure 4G).

We next used two copies each of Stim1 and Orai1 to generate the complex. The best

scoring model among multiple runs of AF3 (Supplementary Table 5), was considered for

the MD simulation. Three independent MD simulations were run for the Stim1:Orai1

tetramer (Movies 4-6). The Stim1:Orai1 complex showed initial large conformational
change but stabilized thereafter (RMSD, Supplementary Figure 10A). The ΔG values for
dimers of Stim1 and Orai1 remained stable throughout the simulation period and were
found to be significantly more negative compared to STX11:Orai complex and Stim:Orai
protomers (Supplementary Figure 10B) (Figure 4K). All three simulation trajectories were
concatenated and clustered to obtain a centroid structure for this complex
(Supplementary Figure 10C). There were 47 clusters, and the largest cluster had 63
structures. Collectively, these data suggest that STX11 directly binds Orai1 with affinities
similar to that of Stim1:Orai1 protomers. However, presence of higher density of Stim1 in
ER-PM junctions of store-depleted cells could, in theory, outcompete STX11:Orai1
interactions by progressively allowing formation of Stim1:Orai1 dimers/ oligomers.
322
The major interactions predicted between STX11 and Orai1 from all-atom MD
simulations are highlighted in (Supplementary Figure 8B-C). To assess the significance
of predicted interactions, we performed site-directed mutagenesis of two key residues of
STX11 predicted to be involved in binding to Orai1 (Figure 5A). Ectopic expression of
untagged N147A_E150A STX11 showed correct localization (Figure 5A) as well as stable
expression (Figure 5B) (Supplementary Figure 11A) but failed to rescue SOCE in STX11
depleted Jurkat T cells (Figure 5C-D). We next expressed His-tagged wildtype and
N147A_E150A mutant STX11 in *E. coli.*, purified (Figure 5E) and assessed their ability to
bind MBP-tagged Orai1 C-terminal tail in an *in vitro* pull-down assay (Figure 5F)
(Supplementary Figure 11B). Compared to wildtype STX11, N147A_E150A mutant
STX11 showed reduced binding to Orai1 C-terminus. Further, we introduced four
mutations, R289A_E272A_E275A_E278A, in the Orai1 C-terminus (Figure 5G), tagged
with YFP and expressed in HEK293 cells. The Orai1 mutant localized to PM (Figure 5G)
but failed to rescue SOCE in Orai1 deficient cells (Figure 5H-I). We expressed and
purified MBP-tagged R289A_E272A_E275A_E278A mutant Orai1 C-terminus from *E.*
coli and assessed its ability to bind purified wildtype as well as N147A_E150A mutant
STX11 in a pull-down assay (Figure 5J) (Supplementary Figure 11C). Binding of Orai1
mutant to both was significantly compromised. Because Stim1 binds an overlapping
region of Orai1 C-terminus, we next tested whether mutation of Orai1 residues affected

Stim1 binding. We co-expressed YFP-tagged soluble CRAC activation domain (CAD) of
Stim1 (YFP-CAD) (Figure 5K) with either CFP-tagged wildtype or mutant Orai1 in
HEK293 cell line. YFP-CAD localizes to the cytosol in HEK293 cells but in cells
overexpressing Orai1, a significant fraction of it localizes to plasma membrane due to its
association with Orai1 [10]. We quantified relative YFP-CAD levels in the plasma
membrane of both groups (Figure 5K-L) but found no difference in PM recruitment of CAD
suggesting that the inability of R289A_E272A_E275A_E278A mutant Orai1 to support
SOCE does not result from defective binding to Stim1. Collectively, these data
demonstrate a direct role of STX11 in the binding and regulation of Orai1.

Upon store depletion, Stim1 localizes to ER-PM junctional regions where it traps
and co-clusters with Orai1. To further understand the mechanism, we first imaged
HEK293 cells expressing CFP-Orai1, Stim1-YFP under resting conditions and then
quantified the levels of Orai1 and Stim1 in ER-PM junctions (puncta) of store-depleted
control and STX11 depleted cells. The distribution of Orai1 and Stim1 was normal in
resting cells (Figure 6A). However, in store-depleted cells, clustering of CFP-Orai1 was
visibly defective in STX11 RNAi treated cells (Figure 6B), even though the intensities
(Figure 6C) and area of Stim1-YFP puncta were normal (Figure 6D). Quantification of
Orai1 intensities confirmed a significant defect in entrapment (Figure 6E) with relatively
higher fraction of Orai1 present outside Stim1 puncta in STX11 depleted cells (Figure 6F).
To determine whether reduced Orai1 clustering resulted from structural defects within
ER-PM junctions, we reversed the direction of fluorescent tags on Orai1 as well as Stim1
and repeated the experiment in Figure 6A-F. Interestingly, the defect in Stim1 mediated
entrapment of Orai could be rescued (Supplementary Figure 12A-B) (Figure 6G-J) but
SOCE remained significantly inhibited even when Orai and Stim were co-overexpressed
(Figure 6K-L). Taken together, these data show that STX11 binding to resting Orai1 has
implications beyond mere co-entrapment of Orai1 with Stim1 and STX11 depletion does
not alter the structure or proximity of junctional ER to the PM. Importantly, STX11 was
found to be limiting in cells co-overexpressing Orai and Stim that have been proposed to
be sufficient for amplifying CRAC currents in earlier studies using patch clamp [26] [27]
[28].

Defects in the entrapment of Orai1 suggest absence of a necessary molecular

transition within Orai1 which precedes its interaction with Stim1. Using step

photobleaching, we have previously shown that under resting conditions, Orai1 exists as

a dimer but forms higher order oligomers, on-site, upon store-depletion [14] [29]. To

directly test whether SNAREs contributed to the on-site assembly of CRAC channel pore

subunit, we co-expressed Orai1-CFP along with Orai1-YFP in control or STX11 depleted

HEK293 cells and measured Orai:Orai FRET. Remarkably, we found significantly higher

Orai:Orai FRET in STX11 depleted cells even in the absence of Stim1 or store-depletion

(Figure 7A). We then sought to determine whether store-depletion could induce a similar

transition in Orai1 in absence of Stim1 in store-depleted scramble control cells. Indeed,

we found Orai:Orai FRET to be significantly higher post store-depletion when compared

to resting state control cells (Figure 7B). In STX11 depleted cells, however, store-

depletion did not show any further increase in FRET over and above what was already

observed in resting cells.

Increasing the density of Stim1 has been earlier proposed to bypass the

requirements for accessory factors in SOCE. To determine whether increasing the levels

of Stim1 could compensate for STX11 depletion, we expressed Orai1 tethered to two

Stim-Orai activating regions (SOAR) of Stim1 and GFP (Orai1-S-S-GFP) (Supplementary

Figure 13A-B) in HEK293 cells. Orai1-S-S-GFP has been previously shown to

constitutively activate Orai1 [30]. In STX11-depleted Orai1-S-S-GFP expressing cells, the

magnitude of calcium entry was significantly smaller (Supplementary Figure 13C-D). To

determine whether reduced calcium entry resulted from a defect in the ability of SOAR to

bind Orai1, we expressed YFP-CAD (Supplementary Figure 13E), in control or STX11

depleted Orai1-CFP expressing HEK293 cell line. Quantification of PM localized YFP-

CAD as a fraction of total CAD showed no defect in STX11 depleted cells (Supplementary

Figure 13E-F). Yet, as seen with Orai1-S-S-GFP, constitutive calcium entry was

significantly reduced (Figure 7C-E). We next expressed the constitutively active H134S

mutant of Orai1 (Supplementary Figure 13G) as it harbors an open pore and the C-

terminal cytosolic tails of Orai1 are unlatched, pointing towards the cytosol [31].

Remarkably, expression of H134S Orai1 completely restored constitutive calcium influx

in STX11 depleted cells (Figure 7F-G). To further determine whether binding with STX11

induces straightening of Orai1 tails or molecular transitions within transmembrane

helices, we expressed in STX11 depleted cells the constitutively active ANSGA (261-265)

mutant of Orai1 (Supplementary Figure 13H), which harbors 4 consecutive mutations in

the Orai1 C-terminus resulting in the constitutive activation of CRAC currents along with

straightening of the C-terminal tail [32]. Intriguingly, constitutive calcium entry in STX11

deficient, Orai1 ANSGA mutant expressing cells was significantly suppressed (Figure 7H-

I). Taken together, these data show that STX11 induces a molecular switch in the pore-

forming subunit of CRAC channels, which precedes and primes Orai1 for correct

assembly and successful gating by Stim1. Such SNARE dependent priming of Orai1 is

crucial and cannot be compensated by merely increasing the density of Stim proteins

bound to Orai or by introducing mutations that alter the orientation of Orai1 tails *in vivo*.

A hypothetical model depicting unprimed and primed Orai1 is shown in Figure 7J.

Q-SNAREs are often found in complex with other SNAREs such as SNAP23/25/29

in target membranes. Some previous studies have implicated the complex of SNAP23/25

and STX1A in the regulation of specific ion channels [33, 34]. The three SNAPs,

SNAP23/SNAP25/SNAP29 are each capable of contributing two SNARE domains to the

trans- and *cis*-SNARE complex. RNAi mediated depletion of SNAP23/SNAP25/SNAP29

did not show a significant reduction in SOCE (Supplementary Figure 14A-C). To

determine whether SNAP23/SNAP25/SNAP29 might still form a part of the STX11:Orai

complex to regulate SOCE, we co-expressed myc-tagged Orai1 with untagged SNAPs

and performed respective co-IPs. We did not find any interaction between either of the

three SNAPs and Orai1 (Supplementary Figure 14D). Similar studies were performed by

co-expressing YFP-Stim1 and the three myc-tagged SNAPs. Again, no interaction was

found between any of the SNAPs and Stim1 (Supplementary Figure 14E). These data

show that unlike in the case of vesicle fusion, STX11 does not collaborate with

SNAP23/SNAP25/SNAP29 in the regulation of SOCE.

FHLH is a heterogeneous immune disorder where diverse genes such as *STX11*,

UNC13D, *Prf1*, *Munc18-2*, *MAGT1, XIAP, LYST, AP3B1, RAB27A*, *ITK*, amongst several

others, are broadly grouped together based on disease symptoms, although the age of
onset and severity varies across the spectrum [35, 36]. Congenital immunodeficiency
involving defects in the CTL and NK cell cytotoxicity appear to be the common factors
underlying the development of FHLH, although the individual pathways involved are
highly diverse. Using shRNA, we screened 6 additional genes, *Perforin1*, *RAB27A*,
UNC13D and *Munc18-2*, *Synaptotagmin11* and *13*, that have been either associated with
vesicle fusion or routinely grouped together with STX11 in diagnosing FHLH
(Supplementary Figure 15). SOCE was not affected by depletion of most genes. Future
studies will evaluate the role of others.

444
445
446   **Discussion:**
447

CRAC channels conduct a small but highly calcium specific current in response to
the depletion of intracellular calcium stores. According to the prevailing view, preformed
Orai hexamers reside in the PM and depend on the ER resident store sensor Stim for all
structural transitions leading up to their activation [10]. Unlike other ion channels, the
process of activation of CRAC currents is exceptionally slow. For instance, following
break-in or store-depletion, on average, it takes ~5 minutes for Stims to cluster in the ER-
PM junctions and for measurable CRAC currents to flow from Orai. During this time, Stim
proteins undergo intramolecular transitions and slowly trap and gate freely diffusing Orai.
It is therefore believed that no molecular transitions take place within the pore forming
subunit, Orai, until its clustering by Stim in the ER-PM junctions.

In STX11 depleted cells, despite normal distribution of resting and store-depleted
Stim1, the clustering of CFP-Orai1, which is thought to be entirely dependent on
associations with Stim1, was visibly defective (Figure 6B-F). Given that Orai:Orai FRET
was defective in STX11 deficient cells even in the absence of Stim1 and store-depletion,
we have shown that Orai1 undergoes, a hitherto unrecognized, SNARE dependent early
molecular transition which precedes clustering and gating by Stim1. Given our earlier
findings of variable Orai1 stoichiometry in resting *versus* store-depleted cells [14], the

most likely explanation for ~25% increase over basal Orai:Orai FRET upon STX11

depletion or store-depletion is oligomerization of Orai1 dimers into higher order oligomers.

A mere 5% increase in absolute NFRET values, for instance, can indicate up to 10-fold

increase in interaction.

Stim1 binds both the N- as well as the C-terminus of Orai1 but their individual

relevance in the trapping *versus* gating of Orai1 remains unclear [37]. Switching the large

fluorescent tag from the Orai1 N- to the C-terminus could rescue its clustering but not

gating by Stim1, essentially reinforcing the idea that STX11 induced molecular shifts in

Orai1 do not merely correct orientations of cytosolic tails for trapping. Orai cytosolic tails

have been shown to be crucial for CRAC channel gating but are missing from the

structures of *Drosophila* Orai [31]. A previous NMR structure suggested that C-terminal

tails of two adjacent Orai1 subunits, bend, pair with each other in an antiparallel fashion

and sit closely apposed to PM [25]. However, in structures of constitutively active H134

mutant Orai, the C-terminal tails were found to orient away from plasma membrane [31].

We have shown that the ANSGA mutant of Orai1 [32], but not H134S Orai1, was sensitive

to STX11 depletion. These data agree with FRET and suggest that STX11 facilitated

molecular transitions occur within the transmembrane regions of Orai1 rather than

changing the orientations of Orai1 tails. The changes in resting cell Orai:Orai FRET

showed that such transition precedes the association and clustering of Orai with Stims.

These data align with our findings of an overlapping region of Orai1 bound by STX11 and

Stim1. Because tagging of STX11 on either end with fluorescent tags results in mis

localization and degradation, our study could not demonstrate the real-time removal of

STX11 from the Orai1- C-terminus to allow for Stim1 binding in live cells. It is also unclear

whether binding or removal of STX11 induces the correct transition in Orai1 assembly.

Further, it remains to be determined whether STX11 removal is actively induced by an

additional chaperone or passive competition due to formation of relatively higher affinity

dimeric/oligomeric associations between Orai1:Stim1 in ER-PM junctions (Figure 4K).

Future studies will be able to systematically evaluate each of these possibilities.

We have shown that the C-terminal cytosolic tail of Orai1 directly couples with

STX11 in resting cells (Figure 4). Although binding appeared reasonably strong in co-

IPs, STX11:Orai1 interaction appeared relatively weak in pull-downs (Figure 4). This is

likely due, in part, to the instability of purified His-STX11 in solution which necessitated

shorter co-incubation times during pull-down assays. Instability could arise from the

absence of a co-chaperone, such as one of the known binding partners of the synaptic

family or membranes to which STX11 attaches in lives cells *via* its C-terminal cysteines

[23]. Together, these factors could be crucial for the stabilization of purified STX11 *in vitro*.

Previous whole cell patch clamp analysis of HEK293 cells co-overexpressing Orai1

and Stim1 has shown up to 50-100-fold amplification of CRAC currents whereas Fura-2

based live imaging of the same cells routinely shows only 2-fold amplification [26] [27]

[28]. The difference in the fold change between these two readouts is often used to

extrapolate and conclude that all other accessory factors that have since been identified

are either not limiting or unnecessary for native CRAC currents. A major problem with this

interpretation is that even though amplification of CRAC currents was shown, none of the

previous patch clamp studies established whether the higher currents resulted from a

greater number of active channels or unchecked conductance per channel by performing

single channel recordings. The fact that recording of I_{CRAC} in the whole cell configuration

involves dialysis of cells with high concentrations of BAPTA, a fast calcium chelator, the

calcium dependent inactivation of CRAC channels, a key mechanism of regulation of

I_{CRAC} amplitude in live cells is disrupted. Furthermore, BAPTA mediated chelation of Ca^{2+}

ions and voltage clamping at hyperpolarizing potentials to record amplified CRAC

currents prevents the progressive depolarization of cells due to influx of positively charged

ions which would have otherwise dampened the driving force for CRAC currents in intact

cells. Given these limitations, Fura-2 assays offer a more physiological reflection of SOCE

amplitudes in live cells as they allow for these regulatory mechanisms, which are

necessary for the regulation of native currents to be still operational in live cells. While

patch clamp remains a direct method of verification of CRAC currents, exercising logic

and caution in interpreting amplitudes of I_{CRAC} is imperative in order for the field to move

forward. Using Fura-2 imaging assays in STX11 depleted cells, we have conclusively

established that STX11 is necessary as well as limiting not only in live and unperturbed

HEK, U2OS, Jurkat and primary T cells but also HEK293 cells co-overexpressing Orai1

and Stim1 (Figure 6K-L).

Most human mutations in the *STX11* coding region result in the complete loss or

severe depletion of STX11 protein levels [23]. The FHLH4 disease, therefore, results from

reduced STX11 expression in most reported patients. Targeted mutagenesis of the

STX11:Orai1 interface lining residues based on the AlphaFold 3 model, however, didn't

destabilize   or   mis-localize   either   N147A_E150A   STX11   or

R289A_E272A_E275A_E278A Orai1 and yet both mutants showed reduced binding to

their respective wildtype counterparts. Importantly, the R289A_E272A_E275A_E278A

mutant of Orai1 retained the ability to bind Stim1.

In addition to the established cytolytic defects in FHLH4 T cells, for the first time,

we have shown that STX11 deficient and FHLH4 T cells harbor a severe defect in SOCE,

NFAT signaling, IL-2 cytokine expression and secretion. Therefore, our data strongly

suggest that T cell calcium signaling defects significantly contribute to FHLH4 disease

progression, independent of exocytotic defects. We propose that design and early

administration of CRAC channel agonists could potentially provide an alternative to

immunosuppressants and bone marrow transplants for FHLH4 patients.

In summary, we have shown that binding to a Q-SNARE, STX11, provides an on-

site molecular switch in the CRAC channel pore, Orai1, necessary for its correct

assembly, functional coupling and gating by Stim1. We suggest that ion channel priming

is a novel, direct and primary role of specific SNAREs. In accordance with this hypothesis,

early SNARE-like proteins have been found in bacteria, which lack an endomembrane

system but express a variety of ion channels [38-40]. SNAREs are now thought to have

evolved from a common archaeal precursor found in the genomes of Asgard and

members of *Legionella* [41].

Materials and Methods

Plasmid constructs and transfection

Stim1 CAD was sub-cloned in eYFPC1 vector after amplification from YFP-Stim1 plasmid to generate YFP-CAD. pLKO.1, psPAX2 and pMD2.G were purchased from Addgene. pLKO.1 cloned shRNA sequences targeting the genes of interest, were either purchased from Horizon Discovery, UK or designed and cloned in-house using the Broad Institute portal (https://portals.broadinstitute.org/gpp/public/gene/search). Full-length human STX11 was cloned from human cDNA prepared from HEK293 cells and subcloned into pcDNA3.1(+), pcDNA/4TO/Myc-HisA (Invitrogen, Grand Island, NY), pMSCV-IRES-mcherry (Addgene), pEF1alpha-IRES (Clontech) and pET28b vector [16] with an N-terminal 6XHis-tag. The fragments of STX11 were cloned in pET28b vector with an N-terminal 6xHis tag for expression in *E. coli*. HA tag was inserted between H30 and G31 in STX11 cloned in pMSCV-IRES-mCherry construct using PCR. H134S and ANSGA mutants of Orai1 were generated using site directed mutagenesis and PCR, respectively. The fragments of Orai1 (1-87, 1-47, 48-103, 256-301 and 272-292) were amplified from full length constructs and cloned into pMAL-c5X vector (New England Biolabs), in-frame with MBP protein coding sequence, as described previously [16]. SNAP23, SNAP25 and SNAP29 cDNA cloned in pCMV-Sport6.1 vectors were purchased from Dharmacon and subcloned into pcDNA4/TO/Myc-HisA. SNAP23 was subcloned in pMAL-c5X vector in-frame with MBP. All plasmid DNA transfections in human cell lines and primary cells were done using Lipofectamine 2000 (Invitrogen)/ Lipofectamine 3000 (Invitrogen) or Amaxa nucleofection kit (Lonza, Basel, Switzerland) respectively, as per manufacturer's protocol.

Cell Lines

Lentiviral shRNA transduction experiments were performed in HEK293, U2OS or Jurkat (ATCC, Manassas, VA) cell lines cultured in low glucose DMEM (Hyclone, Logan, UT) or RPMI (Hyclone) respectively, with 5-10% fetal bovine serum (Hyclone), 1X Penicillin Streptomycin, and GlutaMax (Gibco, Grand Island, NY). Stable cell lines generated using HEK293 and U2OS parent lines have been described previously [9]. HEK293-FT cell line was cultured in high glucose DMEM with 10% FBS and 10mM HEPES, 1X Penicillin

589 Streptomycin, 1X GlutaMax and 1X non-essential amino acid and transfected with the

590 appropriate plasmids to generate viral supernatants. All cell lines were tested for

591 mycoplasma contamination twice every year and found to be negative.

593 **Antibodies and Reagents**

Antibodies/Reagents	Source and Cat no	Dilution/ Conc.
Mouse anti-myc (9E10)	Abcam #ab32	1:1000
Mouse anti-myc (9E10)	Culture supernatant, in house	1:1000
Rabbit anti-SNAP23	SYSY#111203	1:1500
Mouse anti-SNAP25	SYSY#111011	1:1500
Rabbit anti-SNAP29	SYSY#111303	1:1500
Rabbit anti-GFP	Invitrogen#A11122	1:2500
Rabbit anti-flag	Sigma#F7425	1:1600
Mouse anti-flag (M2)	Sigma#F3165	1:1500
Mouse anti-6X His	Invitrogen #MA1-135	1:2000
Rabbit anti-Stx11	Sigma#SAB4301593	1:1000
Mouse anti-NFATc2 (4G6-G5)	Santacruz#7296	1:500
Rabbit anti-NFAT1	Cell Signaling Technology #4389	1:200
Goat anti-Lamin B (M-20)	Santacruz#6217	1:500
Rabbit monoclonal anti-HA	CST#3724S	1:1000
Alpaca VHH anti-rabbit AF488	Jackson #611-544-152	1:400 (1.375μg/ml)
DAPI	Invitrogen #D1306	2μg/ml
Rabbit anti-Stx11	SySy #110113	1.67μg/ml
Rabbit anti-Stx11	Invitrogen #PA5-50800	1:2000
Goat anti-rabbit A647	Invitrogen #A-21245	1:800
Rabbit anti-Stx11	Proteintech #13301-1-AP	0.3μg/ml
Donkey anti-rabbit A647	Jackson #711-605-152	1:800

HRP GAPDH	Sigma #G9295	1:40000
HRP Donkey anti-Rabbit	Jackson #711-035-152	1:50000
HRP Donkey anti-Mouse	Jackson #715-035-151	1:50000
Purified Human IL-2	BD #555051	1 μg/ml
Purified Human IL-2 Biotin	BD #555040	1 μg/ml
Streptavidin HRP antibody	Jackson #016-030-084	0.1 μg/ml
Ultra-LEAF Purified anti-human CD3	Biolegend #317326	5-10μg/ml
Ultra-LEAF Purified anti-human CD28	Biolegend #302934	2μg/ml
Ultra-LEAF Purified anti-human CD49d	Biolegend #304340	2μg/ml
PE anti-human CD107a (LAMP-1)	Biolegend #328608	1:50
APC anti-human CD8 antibody	Biolegend #344722	0.5μg/ml
Fura-2-AM	Invitrogen #F1221	1μM
Thapsigargin (TG)	Sigma #T9033-1MG	1μM
Polybrene (Hexadimethrine-bromide)	Sigma #107689-10G	8μg/ml
NE-PER kit	Thermo #78833	
Protein A/G Mag Sepharose xtra	Cytiva #28967056/66	
Protein A/G 4 Fast Flow Sepharose	Cytiva #17528001	
Protease Inhibitor Cocktail (Mammalian)	Sigma #539134	
Protease Inhibitor Cocktail (Bacterial)	Calbiochem #539132	
Talon Beads	Takara #635502	
Dextrin Sepharose Beads	Cytiva #28935597	

HEPES	Sigma #H3375-100G	
Lipofectamine 2000	Thermofisher #11668019	
Lipofectamine 3000	Thermofisher #L3000001	
Lemo21(DE3) *E.coli.* cells	NEB #C2528J	
IPTG	Himedia#MB072-10G Sigma #16758-10G	
Sarkosyl (N-lauroylsarcosine sodium salt)	Sigma #L9150-50G	
Lysozyme	Sigma #62970-5G-F	
DNaseI Recombinant RNase-free	Roche #04716728001	
PMA (Phorbol Myristate Acetate)	Sigma #P8139	20-50ng/ml
Ionomycin	Sigma #407952	1μM
Human IL-2 protein	Acro Biosystems #H5215	50ng/ml
Lectin from *Phaseolus vulgaris* (PHA)	Sigma #L8754	2μg/ml
Retronectin	Takara #T100B	20μg/ml
RNeasy Plus Mini kit	Qiagen#74134	
QIAshredder	Qiagen#79654	
Superscript IV	ThermoFisher#18091300	
Qubit ssDNA Assay kit	ThermoFisher# Q10212	
Human STX11 TaqMan probe (FAM-MGB)	ThermoFisher# Hs00186823_m1	
Human Beta Actin TaqMan probe(FAM-MGB)	ThermoFisher# Hs01060665_g1	
Human IL2 Taqman probe (FAM-MGB)	ThermoFisher# Hs00174114_m1	
Paraformaldehyde	Electron Microscopy Sciences #157-8	

Bovine Serum Albumin	Jackson #001000173	
Luria Bertini Broth	Himedia #M1245	
HBSS	Sigma #H8264	
Terrific Broth	Sigma #T9179	
Alexa fluor 647 conjugated Alpha Bungarotoxin	Invitrogen #B35450	1μg/ml
HRP rabbit anti-GFP	Invitrogen #A10260	
HRP Goat anti-rabbit Fc	Jackson #111-035-008	1:50,000
Ampicillin	Sigma #A0166-5G	100ug/ml
Kanamycin	Sigma #K1876-1G	50ug/ml
Chloramphenicol	Sigma #C 0378-5G	100ug/ml
30% Acrylamide:Bis Solution (29:1)	Biorad #1610156	
DTT (Dithiothreitol)	Biorad #1610611	
0.05% Trypsin-EDTA	Gibco #25300054	
100X Penicillin Streptomycin	Sigma #P4333-100ML	
Puromycin	Gibco #A11138-03	
Pluronic F-127 (20% solution in DMSO)	Invitrogen #P3000MP	0.02%

Lentiviral transductions

For the generation of lentiviral supernatants, shRNAs cloned in pLKO.1 were co-transfected with psPAX2 packaging and pMD2.G envelope plasmids into HEK293-FT cells using the calcium phosphate method of transfection. 48- and 72-hours post-transfection, viral supernatants were collected, pooled and stored at -80°C till further use. For transduction of HEK293 and Jurkat cells, 0.1 million cells were plated in 6- or 24-well plates either the day before (HEK293) or the same day (Jurkat). Viral supernatants were added to the cells along with 8ug/ml polybrene and cells were spun at 2500 RPM, 30°C for 90 mins. 24 hours or 48 hours post-spinfection, Puromycin was added at a final

concentration of 1ug/ml to HEK293 and Jurkat cells, respectively. Cells were analyzed 3-

5 days post transduction.

Immunocytochemistry and confocal imaging

For STX11 localization, resting or store-depleted (1μM Thapsigargin) WT HEK293 or

HEK293 cells stably expressing Orai1-YFP were either plated in 6 well plates and

spinfected with viral supernatants generated from pMSCV-STX11(HA)-IRES-mCherry,

pMSCV-IRES-mCherry (empty vector control) or plated in carbon coated, glow

discharged 35mm glass bottom dishes (IBDI) and transfected with pEF1alpha-STX11-

IRES-mCherry. 24 hr post-spinfection, cells were plated in 35mm glass bottom dishes.

For immunolabelling, cells were washed with Ringer's buffer, fixed with 4% PFA and

blocked with 3% BSA containing 0.1% NP40 for 1.5 hrs. Post-fixation and

permeabilization, cells were incubated either with anti-HA (CST) or rabbit anti-Stx11

(SySy/ Proteintech) primary antibodies O/N at 4°C, washed and incubated with either

Alpaca VHH anti-rabbit AF488, donkey anti-rabbit A647 or goat anti-rabbit A647

secondary antibodies for 1 hour at room temperature (RT). Cells were counter stained

with DAPI for 10 min at RT and imaged using Olympus FV3000 laser scanning confocal

microscope. Images were acquired sequentially using the following parameters: DAPI

(DM405/488 dichroic, 405nm excitation, 430-470nm emission); anti-HA Alexa488

(DM405/488, 488nm excitation, 500-590nm emission); STX11 (DM405/488/561/640,

640nm excitation, 650-750nm emission); Orai1-YFP (DM405/488/561/640, 488nm

excitation, 521-591nm emission); CFP-Stim1 (DM405-445/514/594, 405nm excitation,

448-510nm emission). For STX11-N147A-E150A mutant, HEK293 cells were spinfected

with viral supernatants and imaged as described above where WGA (Wheat germ

agglutinin) conjugated with Alexa 488 was used to mark plasma membrane.

Quantification of Orai1 and Stim1 intensities inside/outside ER-PM puncta

HEK293 cells stably expressing Orai1-YFP with CFP-Stim1 or CFP-Orai1 with Stim1-YFP

were plated in 6 well plates and transduced with scramble or STX11 shRNA. 48hr post

transduction, cells were trypsinized and plated in carbon coated 35mm dishes. Images

were acquired using a TIRF microscope setup described before [14]. Briefly, resting cells

were imaged first and positions were marked. Store-depleted images were acquired after

incubation with 1 μM Thapsigargin (TG) and 10mM EGTA for ~6 mins. Analysis was done

using Otsu local thresholding in ImageJ software. Stim1 and Orai1 intensity (AU) and area

μm^2 values obtained from the analysis were plotted using Origin software.

Orai:Orai FRET

Scramble and STX11 shRNA transduced HEK293 cells were plated on freshly carbon

coated, glow discharged 35mm glass bottom dishes (IBDI) and transfected with 0.5 μg

each of Orai1-CFP and Orai1-YPF plasmid DNA using Lipofectamine 3000. 16 hours post

transfection, co-transfected cells were imaged using PlanApoN 60XO/1.42 NA

objective with Olympus FV3000 laser scanning microscope. Orai1-CFP, Orai1-YFP and

FRET images were acquired with the following excitation laser and emission bandwidth

parameters using DM405-445/514 dichroic mirror: CFP Ex-405nm Em-460-630nm; YFP

Ex-514nm Em- 530-630nm; FRET Ex-405nm, Em-530-630nm. ROIs representing the

plasma membrane rim were drawn around each cell, followed by local thresholding using

Bernsen's algorithm in Fiji software to generate an image mask which was applied to

CFP, YFP, FRET images to extract mean intensity values. The background subtracted

mean intensities of FRET (I_{FRET}), CFP (I_{Donor}) and YFP($I_{acceptor}$) images were measured

for each cell. Cells singly expressing Orai1-CFP or Orai1-YFP only were used as CFP

donor or YFP acceptor channel bleed-through (BT) controls. The BT was calculated using

the following formula:

$BT_{Donor} = \dfrac{I_{FRET}}{I_{Donor}}$ when the Orai1-CFP donor alone is expressed,

$BT_{acceptor} = \dfrac{I_{FRET}}{I_{acceptor}}$ when the Orai1-YFP acceptor alone is expressed.

FRET and NFRET percentage was calculated by subtracting the contributions of CFP

and YFP bleed through from the FRET signal using the following:

663
$$FRET = I_{FRET} - (BT_{Donor} * I_{Donor}) - (BT_{Acceptor} * I_{Acceptor})$$

$$NFRET\% = \left(\frac{FRET}{\sqrt{(I_{Acceptor} * I_{Donor})}} \right) * 100$$

Alpha-bungarotoxin binding assay

U2OS cell line stably expressing Orai1-BBS-YFP construct has been described previously[14]. Briefly, Orai1-BBS-YFP expressing U2OS cells were either transduced with scramble shRNA/ STX11 shRNA/ pMSCV-IRES-mcherry(EV) or pMSCV-STX11-IRES-mcherry. Transduced cells were trypsinized either 3-4 days post-transduction or 48 hour post-transduction (in the case of pMSCV), centrifuged and labelled on ice with AF647 conjugated alpha-bungarotoxin (1µg/ml) containing 2% FBS in 1X HBSS for 30 mins. Following this, cells were washed with HBSS and fixed with 4% PFA (~20 min at RT). Samples were run using Cytoflex FACS analyzer (Beckman Coulter) and data were analyzed using Flow Jo software.

Validation of STX11 knockdown using qPCR

Scramble and STX11 shRNA treated HEK293 cells were trypsinized, counted and equal number of cells were used for lysis. The lysates were homogenized using QIA shredder columns (Qiagen) and RNA isolation was done using RNeasy Plus Mini kit (Qiagen). Total RNA isolated from both the groups was used for cDNA synthesis using oligo dT primers and Superscript IV as per manufacturer's guidelines. The concentration of synthesized cDNA was estimated using Qubit ssDNA assay kit and STX11 TaqMan probes were used to perform the qPCR using LightCycler 96 (Roche). Beta-actin was used as housekeeping control.

Single cell Ca^{2+} imaging

Cells were plated in carbon-coated glass bottom dishes (one day prior or 30 mins before the assay for HEK293 and Jurkat T cells respectively) and loaded with 1µM Fura-2 AM dye in HBSS (CaCl$_2$ 1.8mM, KCl 5.36mM, MgSO$_4$ 0.81mM, NaCl 136.89 mM, Na$_2$HPO4 0.335mM, D-Glucose 5.55mM) for 30 mins, at 37°C in the dark. After incubation, cells were washed and incubated in Ringer's buffer (135mM NaCl, 5mM KCl, 1.8mM CaCl$_2$,

1mM $MgCl_2$, 5.6mM Glucose, 10mM HEPES) for an additional 10 mins, washed and

imaged in Ringer's buffer or Calcium-free Ringer's buffer (135mM NaCl, 5mM KCl, 1mM

$MgCl_2$, 5.6mM Glucose, 10mM HEPES at pH7.5), as indicated. Olympus IX-71 inverted

microscope equipped with a Lamda-LS illuminator (Sutter Instrument, Novato, CA), Fura-

2 (340/380) filter set (Chroma, Bellows Falls, VT), a 10X 0.3NA objective lens (Olympus,

UPLFLN, Japan), and a Photometrics Coolsnap HQ2 CCD camera was used to capture

images at a frequency of ~1 image pair every 2 or 4 seconds interval. Data were acquired,

analyzed and plotted using MetaFluor (Molecular Devices, Sunnyvale, CA), Microsoft

Excel, and Origin softwares. Approximately 30-40 cells were imaged per group in each

experiment unless otherwise stated. Mean SOCE ± SE for each experiment was

calculated from three independent experiments where the maximum value of the

scramble shRNA group (amongst the three experiments) was set at 100% and the relative

responses of other scr and STX11 shRNA groups were normalized w.r.t it.

For constitutive calcium influx assay, shRNA treated HEK293 cells were either

transfected or nucleofected with Orai1-S-S-EGFP, Orai1(H134S)-YFP or Orai1(ANSGA)-

YFP and ~12-14 hour post transfection (~4-5 hour post nucleofection), cells were loaded

with Fura-2 AM, washed and imaged in Ringer's buffer with 0 mM calcium to acquire

baselines and 2 mM calcium thereafter. To measure CAD induced constitutive calcium

influx, Orai1-CFP expressing stable HEK293 cells were nucleofected with YFP-CAD and

analyzed ~4 hour later as described above. To identify cells expressing Orai1 mutants

and fusion proteins in the plasma membrane, cells were imaged using a 20X 0.7NA water

objective lens (Olympus, UApoN340, Japan). Images were acquired at a frequency of ~1

image pair every 10 seconds interval to avoid photobleaching and analyzed as described

above.

Measurement of NFAT nuclear localization by Western blot

Jurkat T cells were transduced with scramble or STX11 shRNA to knock-down STX11 as

described above. On day 4 post-transduction, cells were collected, spun down and

resuspended in plain RPMI media and rested for 1hr at 37°C. Following this, cells were

counted and divided into two equal groups. One group was resuspended in RPMI

(unstimulated) and the other in RPMI media containing 1 μM TG + 10 ng/ml PMA (Phorbol

Myristate Acetate) (stimulated) and incubated at 37°C for 30 mins. Following incubation,

cells were pelleted and the nuclear and cytosolic protein fractions were separated using

NE-PER kit according to the manufacturer's guidelines and subjected to SDS-PAGE and

western blot using the mouse anti-NFATc2 primary antibody (Santa Cruz) followed by

Donkey anti-mouse secondary antibody.

Estimation of nuclear translocation of NFAT by immunolabeling

Control and STX11 depleted Jurkat T cells were rested in plain RPMI for ~1 hour prior to

the assay. ~100,000 cells per group were plated on freshly carbon-coated coverslips for

40 minutes and stimulated for 1 hour at 37°C with 5 µg/ml anti-CD3 antibody, diluted in

plain RPMI. Following this, cells were washed and fixed using 4% PFA diluted in 1X PBS

for 20 minutes at RT, washed twice and incubated with 30 mM Glycine for 10 minutes at

RT. For permeabilization and blocking, cells were incubated with 0.1% Saponin, 3% BSA

diluted in 1X PBS for 1 hour at RT, washed and incubated with anti-NFAT primary

antibody (anti-NFAT1, CST) at 4°C, overnight. Following primary antibody application,

cells were washed with 1X PBS and incubated with anti-Rabbit AF647 secondary

antibody for 1 hour, washed and stained with DAPI for 5 minutes followed by additional

washes. Images were acquired in the FV3000 confocal microscope using a 100X

objective lens. DAPI was used to identify the nuclear area. Nuclear *versus* whole cell

(total) NFAT mean intensity ratio was plotted across different groups.

Whole cell lysates (WCLs), Western blot and Co-immunoprecipitation (Co-IP)

HEK293 cells transfected with the desired plasmids were lysed using buffer containing

50 mM Tris-Cl (pH 8.0), 150 mM NaCl, 1% NP-40, 1 mM PMSF, and protease inhibitor

cocktail. The whole cell lysates were centrifuged at 21000 g for 15 minutes and

supernatants were subjected to SDS-PAGE, and western blot. For immunoprecipitations,

lysates were divided into two equal parts. To the first tube, the appropriate anti-mouse or

anti-rabbit primary antibody was added and to the second, same amount of the respective

IgG control antibody was added. The antibody-lysate mixtures were incubated overnight

at 4°C. Following this, Protein A/G Mag Sepharose beads were added and incubated with

the antibody-lysate mixtures for 4 hours, washed with the lysis buffer containing 0.1% NP-
40, boiled with 1X Laemmli buffer + 120 mM DTT and subjected to SDS-PAGE and
western blot analysis. Typically, 1/10ᵗʰ of the whole cell lysate (WCL) was loaded in the
input lane of the co-IP blots.
760
***E. coli* expression and *in-vitro* binding assays**
Full-length His_6-tagged Stx11 and truncation mutants were cloned in pET28b, expressed
in Lemo21 (DE3) *E. coli* cells and induced with 1 mM IPTG (Isopropyl β- d-1-
thiogalactopyranoside) for 14-18 hours at 18°C. The cell pellets were lysed in buffer
containing 50 mM Tris-Cl (pH 8.0), 150 mM NaCl, 10% Glycerol, 1 mM PMSF, 1%
Sarkosyl, 0.1 mg/ml Lysozyme, protease inhibitor cocktail and sonicated on ice. DNase I
was added after sonication and the lysates were further incubated for ~60 minutes before
centrifugation at 21000 g for 40 mins. The supernatants were subjected to SDS-PAGE to
confirm expression by Coomassie staining and subsequently used for pull-down assays.
MBP-tagged Orai1 constructs were expressed in Lemo21 (DE3) *E. coli* cells, induced with
0.3mM IPTG (Isopropyl β- d-1-thiogalactopyranoside) for 14-18 hours at 18°C. The cell
pellets were lysed in buffer containing 50 mM Tris-Cl (pH 8.0), 150 mM NaCl, 5% Glycerol,
1 mM PMSF, 0.1 mg/mL Lysozyme and protease inhibitor cocktail (1:200) and sonicated
on ice. DNase I was added after sonication and the lysates were further incubated for ~60
minutes before centrifugation at 21000 g for 40 mins. The supernatants were collected
and subjected to SDS-PAGE to confirm expression by Coomassie staining and used for
pull-down assays. Following lysis, the His_6-tagged Stx11 proteins were purified using
Talon Beads and MBP-tagged Orai1 proteins using Dextrin Sepharose/Amylose resin
according to the manufacturer's guidelines. For *in-vitro* binding assays, lysates prepared
from cells expressing MBP or MBP-tagged Orai1 fragments were incubated with 25 μL
Dextrin Sepharose Beads and incubated for 2 hours at 4°C. After incubation, beads were
washed thrice and ~50-125 ng of purified His-tagged full length Stx11, His-tagged SNARE
or His-tagged H_{abc} domains were diluted in buffer containing 50mM Tris-Cl (pH8), 150
mM NaCl, 5% Glycerol and 0.1% NP-40 and added to the beads. Following 1 hour of
incubation at 4°C, the beads were washed thrice, re-suspended and boiled in the binding
buffer containing 1X Laemmli and 120 mM DTT. The eluate was subjected to SDS-PAGE

and Western blot. MBP and MBP-tagged proteins were detected with Ponceau staining,

Full length STX11 using Rabbit anti-STX11 primary antibody (Thermo) followed by

Donkey anti-rabbit HRP and His-tagged STX11 fragments were detected or mouse anti-

6XHis primary antibody (Invitrogen) followed by Donkey anti-mouse HRP secondary

antibody.

qPCR to estimate gene expression in Jurkat T cells

Scramble (scr) and STX11 shRNA treated Jurkat T cells were rested for ~1 hour prior to

the assay, stimulated with soluble anti-CD3 (5-10 μg/ml) for 3 hours and washed with

HBSS. Total RNA was isolated using RNeasy Mini kit as per manufacturer's instructions.

cDNA was synthesized using random hexamers and Invitrogen Superscript IV kit and

quantified using Qubit ssDNA assay. Taqman probes for IL-2 and beta-actin

housekeeping control were used for performing the qPCR in triplicates. Data analysis was

done by calculating the double delta C_t values.

EIA to measure IL-2 level in cell supernatant of Jurkat T cells

Scramble (scr) and STX11 shRNA treated Jurkat T cells were stimulated with plate-

coated anti-CD3 (10 μg/ml) and the culture supernatants were collected at 12 and 24 hour

post-stimulation. EIA plates were coated with purified mouse anti-human IL-2 capture

antibody (1 μg/ml) overnight at 4°C, washed with 0.05% PBST and blocked with 2% BSA

for 2 hours. After blocking, the plates were incubated with the culture supernatants for 2

hours at RT, washed and incubated with the detection antibody (Biotin mouse anti-human

IL-2, 1 μg/ml) for 2 hours. After washing, Streptavidin HRP (0.1 μg/ml) followed by TMB

reagent were added. 250 mM HCl was added to stop the reaction and absorbance was

measured at 450 nm.

Isolation and culture of human PBMCs

Whole blood freshly collected in Heparin or Citrate Phosphate Dextrose Adenine (CPDA)

solution was subjected to density gradient centrifugation using Ficoll-Paque PLUS media.

The buffy coat was separated, washed twice with HBSS and cultured with IL-2 (50 ng/ml).

Cells were stimulated once per week with PHA (2 μg/ml) for 72 hours and rested in IL-2

for the remaining 72-96 hours. Unless specified, all the assays were performed following

24-48 hours of rest, post stimulation.

Isolation of genomic DNA and sequencing

Genomic DNA was extracted from PBMCs using Phenol-Chloroform-Isoamyl alcohol.

STX11 genomic region flanking the mutation was PCR amplified using primers: 5'

Forward   -CATGCACGACTACAACCAGGC   and   3'   Reverse   -

GGGACAGCAGAAGCAGCAGAGGG. The resulting PCR products were separated on

2% agarose gel, excised and extracted using the Macherey-Nagel Nucleospin columns

and subjected to Sanger sequencing using the 5' Forward PCR primer to confirm the

mutation.

Measurement of SOCE in human PBMCs

PBMCs in culture were washed, rested and allowed to adhere to freshly carbon-coated

(IBDI) dishes for 1 hour in plain RPMI at 37°C. The cells were washed with HBSS and

incubated with 1 ml of 1 μM Fura-2-AM diluted in Ringer's buffer for 40 minutes, washed

and incubated for an additional 10 minutes to allow de-esterification of the dye. The assay

was started with 1 ml of Calcium free Ringer's buffer in the imaging dish and images were

captured every 4 seconds. After capturing the baseline for ~60 seconds, stores were

depleted by the addition of 1 μM Thapsigargin. In other assays, 10 μg/ml of anti-CD3 and

5 μg/ml of the secondary antibody were used to cross-link the TCRs and thereby induce

store depletion. ~5 minutes post-store-depletion, Calcium Chloride ($CaCl_2$) was added

back to the dish at a final concentration of 2 mM to estimate the magnitude of store

operated calcium entry.

Degranulation assay

PBMCs were cultured in RPMI containing 10% FBS and 50 ng/ml IL-2 and stimulated

with PHA (2 μg /ml) 48hrs before the assay. 24hrs prior to the assay, IL-2 was washed

off but PHA was re-added. On the day of the assay cells were washed twice to remove

PHA and any growth factors. To measure degranulation, cells were either left

unstimulated, stimulated with a combination of anti-CD3 (10 μg/ml plate-coated), anti-

CD28 (2 μg/ml soluble) and anti-CD49d (2 μg/ml soluble) or a combination of Ionomycin

(1 μM) and PMA (50 ng/ml) for 3.5 hours. CD107a-PE antibody (1:50 dilution) was added

to each of the three groups at the start of stimulation. Following stimulations, cells were

transferred to ice, washed with cold HBSS containing 2% FBS and incubated with anti-

CD8 APC for 20 minutes, washed twice with cold HBSS and analyzed using Cytoflex

(Beckman Coulter) flow cytometer.

qPCR to estimate gene expression in human PBMCs

PBMCs were taken off IL-2 48 hours prior to the assay. On the day of the assay, cells

were washed twice, left unstimulated and either stimulated with anti-CD3 (10 μg/ml, plate-

coated), anti-CD28 (2 μg/ml) and anti-CD49d (2 μg/ml) or with Ionomycin (1 μM) and PMA

(50 ng/ml) for 6 hours. To end the stimulation, cells were washed with cold HBSS, pelleted

and used for RNA extraction. Total RNA was isolated using Qiagen RNeasy mini kit and

cDNA was synthesized using random hexamers and Superscript IV (Invitrogen). cDNA

was quantified using Qubit and subjected to QPCR analysis using Taqman probes for IL-

2 and beta-actin housekeeping gene control in triplicates. Analysis was performed by

calculating the double delta C_t values.

Transduction of human PBMCs with pMSCV-STX11

For viral transduction of PBMCs, polystyrene non-TC treated 24-well plates were coated

with retronection (20 μg/ml) overnight at 4°C, blocked with 2% BSA for 30 minutes at

room temperature and washed twice with HBSS. The retroviral supernatants were added

onto the coated wells and the plates were spun at 1800g for 2 hours at 30°C. Following

spin, the wells were washed with the blocking solution. PBMCs cultured in RPMI

containing 10% FBS were stimulated with PHA (2 μg/ml) and IL-2 (50 ng/ml) for 48 hours

prior to transduction. Stimulated PBMCs were transferred to coated plates at a density of

0.25 million cells/well centrifuged at 400g for 40 minutes at 30°C. The cells were analyzed

48-72 hours post transduction.

Patch clamp measurements

Patch clamp recordings were performed by using an Axopatch 200B amplifier (Axon
Instruments, Foster City, CA) interfaced to an ITC-18 input/output board (Instrutech, Port
Washington, NY) and an iMac G5 computer. Currents were filtered at 1 kHz with a four-
pole Bessel filter and sampled at 5 kHz. Recording electrodes were pulled from 100-µl
pipettes (VWR), coated with Sylgard, and fire-polished to a final resistance of 2 to 5 MΩ.
Stimulation and data acquisition and analysis were performed by using in-house routines
developed by RS Lewis (Stanford University) on the Igor Pro platform (Wavemetrics, Lake
Oswego, OR). All data were corrected for the liquid junction potential of the pipette
solution relative to Ringer's in the bath (-10 mV) and for leak currents collected in 20 mM
extracellular free calcium plus 10µM La^{3+}. The holding potential was +30 mV. The
standard voltage stimulus consisted of a 100-ms step to –100 mV followed by a 100-ms
ramp from –100 to +100 mV applied at 1 s intervals. I_{CRAC} was typically activated by
passive depletion of ER Ca^{2+} stores by intracellular dialysis of 8 mM BAPTA.

Solutions and chemicals
The standard extracellular Ringer solution contained 130 mM NaCl, 4.5 mM KCl, 20 mM
$CaCl_2$, 10 mM tetraethylammoniumchloride (TEA-Cl), 10 mM D -glucose and 5 mM
HEPES (pH 7.4 with NaOH). The standard divalent-free (DVF) Ringer solutions contained
(in mM) 150 NaCl, 10 HEDTA, 1 EDTA, and 10 HEPES (pH 7.4). The internal solution
contained: 135 mM Cs aspartate, 8 mM $MgCl_2$, 8 mM Cs-BAPTA and 10 mM HEPES (pH
7.2 with CsOH).

Data analysis
Analysis of current amplitudes was typically performed by measuring the peak currents
during the -100 mV pulse. Averaged results are presented as the mean value standard
error of the mean (SEM). Reversal potentials were measured from the average of several
leak-subtracted sweeps (4–6) in each cell. For datasets with two groups, statistical
analysis was performed with two- tailed t test to compare between control and test
conditions with a confidence level of 95%, and results with $p<0.05$ were considered
statistically significant.

STX11:Orai1 and Stim1:Orai complex prediction

Full length STX11 and its two domains, (H_{abc} 41-167 and SNARE 183-267) and Orai1 (N-terminus 1-87 and C-terminus 256-301) were used to generate the complex of STX11 and Orai1 using AlphaFold3 (AF3) in all combinations. For Stim1, residue 312- 387 were used along with Orai C-terminus 256-301. The resultant models were assessed using ipTM and pTM scores. The best scoring combinations were further considered to generate more models by changing the seed values. The best model in terms of highest ipTM and pTM values was considered for further analysis. Custom script was used to analyze the contact frequency of interacting residues of STX11-H_{abc} and Orai1 C-terminus complex as well as Stim1 and Orai1 C-terminus, across all predicted models.

Molecular dynamics (MD) simulation

The STX11 H_{abc}:Orai1 C-terminus and Stim1(312-387):Orai1 C-terminus complexes were subjected to all-atom MD simulations. Initially, the complexes was prepared using protein-preparation module of Schrodinger which involves H-bond network optimization and restrained minimization of the initial structure. The prepared structures were solvated using TIP3P water model in an orthorhombic box and neutralized by adding counter ions. OPLS4 force field was used, and simulation system was generated by specifying 150 mM salt (NaCl). The solvated system was subjected to default relaxation protocol of Desmond followed by production run for 500 ns at 300K and 1 atm pressure in NPT ensemble. The default relaxation protocol includes several short simulation steps. (1) Brownian dynamics simulation for 100 ps at 10 K temperature in NVT with restraints on solute heavy atoms (2) Simulation in NVT ensemble at 10 K for 12 ps with restraints on solute heavy atoms (3) 12 ps simulation in NPT ensemble at 10 K with restraints on solute heavy atoms (4) Simulation in NPT ensemble for 12 ps with restraints on solute heavy atoms (5) Simulation in NPT ensemble for 24 ps without restraints. Three independent runs were executed with different initial seeds for both structures.

Analysis of MD simulation

The MD runs were analyzed for the stability of the complex and interactions between the subunits. The RMSD was calculated using simulation interaction diagram (SID) module

of Schrodinger. The first frame was used as reference frame for RMSD calculation. Script

"analyze_trajectory_ppi.py" was used to calculate the interactions between STX11:Orai1

and Stim1:Orai1. The binding energy (ΔG) at each nanosecond across simulation was

calculated using "thermal_mmgbsa.py" script of Schrodinger. The representative

structure was generated through trajectory clustering. The last 200 ns frames from each

run were combined and clustered using scripts "trj_merge.py" and "trajectory cluster" of

Schrodinger. The cluster representative of the largest cluster was used to analyze the

protein-protein interactions. PDB Sum was used to calculate the interactions in the cluster

representative. Figures were generated using Pymol and MD movies using Maestro.

Statistical Analysis

Statistical analysis was done using unpaired two-tailed Student's *t* test where only two

groups were being compared and ANOVA where more than two groups were present.

Normality of distribution was checked in Origin. Mann Whitney U Test was used for

datasets with non-normal (or skewed) distribution. P values for all significance tests were

*P<0.05; **P<0.01; ***P<0.001.

Acknowledgements: We thank Vivien Beziat, Yenan Bryceson, Heinrich Schlums, Jelve

Nejati Zendegani, Stephan Ehl, and Jasmin Mann for help with human PBMC handling,

isolation, storage and shipping. Anand Vaidya for advice on STX11 purification. Jyoti

Rohilla for technical support. Bio Render for the generation of models and IBS 2.0 for the

generation of schematics shown in this manuscript. This work was supported in part by

Department of Atomic Energy, Government of India under Project Identification

No. RTI 4007 and NIH-NIAID grant AI108636.

References:

1.	Zhang, K., et al., *Familial Hemophagocytic Lymphohistiocytosis*, in *GeneReviews((R))*, M.P. Adam, et al., Editors. 1993: Seattle (WA).
2.	D'Orlando, O., et al., *Syntaxin 11 is required for NK and CD8(+) T-cell cytotoxicity and neutrophil degranulation.* Eur J Immunol, 2013. **43**(1): p. 194-208.
3.	Clapham, D.E., *Calcium signaling.* Cell, 2007. **131**(6): p. 1047-58.
4.	Vig, M. and J.P. Kinet, *The long and arduous road to CRAC.* Cell Calcium, 2007. **42**(2): p. 157-62.
5.	Vig, M., et al., *Defective mast cell effector functions in mice lacking the CRACM1 pore subunit of store-operated calcium release-activated calcium channels.* Nat Immunol, 2008. **9**(1): p. 89-96.
6.	Maul-Pavicic, A., et al., *ORAI1-mediated calcium influx is required for human cytotoxic lymphocyte degranulation and target cell lysis.* Proc Natl Acad Sci U S A, 2011. **108**(8): p. 3324-9.
7.	Vig, M., et al., *CRACM1 multimers form the ion-selective pore of the CRAC channel.* Curr Biol, 2006. **16**(20): p. 2073-9.
8.	Vig, M., et al., *CRACM1 is a plasma membrane protein essential for store-operated Ca2+ entry.* Science, 2006. **312**(5777): p. 1220-3.
9.	Miao, Y., et al., *An essential and NSF independent role for alpha-SNAP in store-operated calcium entry.* Elife, 2013. **2**: p. e00802.
10.	Prakriya, M. and R.S. Lewis, *Store-Operated Calcium Channels.* Physiol Rev, 2015. **95**(4): p. 1383-436.
11.	Feske, S., et al., *A mutation in Orai1 causes immune deficiency by abrogating CRAC channel function.* Nature, 2006. **441**(7090): p. 179-85.
12.	Zhang, S.L., et al., *Genome-wide RNAi screen of Ca(2+) influx identifies genes that regulate Ca(2+) release-activated Ca(2+) channel activity.* Proc Natl Acad Sci U S A, 2006. **103**(24): p. 9357-62.
13.	Yoon, T.Y. and M. Munson, *SNARE complex assembly and disassembly.* Curr Biol, 2018. **28**(8): p. R397-R401.
14.	Li, P., et al., *alpha-SNAP regulates dynamic, on-site assembly and calcium selectivity of Orai1 channels.* Mol Biol Cell, 2016. **27**(16): p. 2542-53.
15.	Miao, Y., et al., *Na+ influx via Orai1 inhibits intracellular ATP induced mTORC2 signaling to disrupt CD4 T cell gene expression and differentiation.* Elife, 2017. **6**.
16.	Ramanagoudr-Bhojappa, R., Y. Miao, and M. Vig, *High affinity associations with alpha-SNAP enable calcium entry via Orai1 channels.* PLoS One, 2021. **16**(10): p. e0258670.
17.	Jarvis, S.E. and G.W. Zamponi, *Masters or slaves? Vesicle release machinery and the regulation of presynaptic calcium channels.* Cell Calcium, 2005. **37**(5): p. 483-8.
18.	Lu, Q., et al., *Syntaxin 1A supports voltage-dependent inhibition of alpha1B Ca2+ channels by Gbetagamma in chick sensory neurons.* J Neurosci, 2001. **21**(9): p. 2949-57.
19.	Bezprozvanny, I., R.H. Scheller, and R.W. Tsien, *Functional impact of syntaxin on gating of N-type and Q-type calcium channels.* Nature, 1995. **378**(6557): p. 623-6.

1014   20.   Vig, M. and J.P. Kinet, *Calcium signaling in immune cells.* Nature immunology, 2009.
1015   **10**(1): p. 21-7.
1016   21.   Bryceson, Y.T., et al., *Defective cytotoxic lymphocyte degranulation in syntaxin-11*
1017   *deficient familial hemophagocytic lymphohistiocytosis 4 (FHL4) patients.* Blood, 2007.
1018   **110**(6): p. 1906-15.
1019   22.   Hackmann, Y., et al., *Syntaxin binding mechanism and disease-causing mutations in*
1020   *Munc18-2.* Proc Natl Acad Sci U S A, 2013. **110**(47): p. E4482-91.
1021   23.   Hellewell, A.L., et al., *Analysis of familial hemophagocytic lymphohistiocytosis type 4*
1022   *(FHL-4) mutant proteins reveals that S-acylation is required for the function of syntaxin*
1023   *11 in natural killer cells.* PLoS One, 2014. **9**(6): p. e98900.
1024   24.   Jahn, R. and R.H. Scheller, *SNAREs--engines for membrane fusion.* Nat Rev Mol Cell Biol,
1025   2006. **7**(9): p. 631-43.
1026   25.   Stathopulos, P.B., et al., *STIM1/Orai1 coiled-coil interplay in the regulation of store-*
1027   *operated calcium entry.* Nat Commun, 2013. **4**: p. 2963.
1028   26.   Peinelt, C., et al., *Amplification of CRAC current by STIM1 and CRACM1 (Orai1).* Nat Cell
1029   Biol, 2006. **8**(7): p. 771-3.
1030   27.   Mercer, J.C., et al., *Large store-operated calcium selective currents due to co-expression*
1031   *of Orai1 or Orai2 with the intracellular calcium sensor, Stim1.* J Biol Chem, 2006.
1032   **281**(34): p. 24979-90.
1033   28.   Soboloff, J., et al., *Orai1 and STIM reconstitute store-operated calcium channel function.*
1034   J Biol Chem, 2006. **281**(30): p. 20661-20665.
1035   29.   Penna, A., et al., *The CRAC channel consists of a tetramer formed by Stim-induced*
1036   *dimerization of Orai dimers.* Nature, 2008. **456**(7218): p. 116-20.
1037   30.   Li, Z., et al., *Graded activation of CRAC channel by binding of different numbers of STIM1*
1038   *to Orai1 subunits.* Cell Res, 2011. **21**(2): p. 305-15.
1039   31.   Hou, X., S.R. Burstein, and S.B. Long, *Structures reveal opening of the store-operated*
1040   *calcium channel Orai.* Elife, 2018. **7**.
1041   32.   Zhou, Y., et al., *The STIM1-binding site nexus remotely controls Orai1 channel gating.*
1042   Nat Commun, 2016. **7**: p. 13725.
1043   33.   Cormet-Boyaka, E., et al., *CFTR chloride channels are regulated by a SNAP-23/syntaxin*
1044   *1A complex.* Proc Natl Acad Sci U S A, 2002. **99**(19): p. 12477-82.
1045   34.   Toft-Bertelsen, T.L., et al., *Regulation of Ca2+ channels by SNAP-25 via recruitment of*
1046   *syntaxin-1 from plasma membrane clusters.* Mol Biol Cell, 2016. **27**(21): p. 3329-3341.
1047   35.   Jessen, B., et al., *Graded defects in cytotoxicity determine severity of hemophagocytic*
1048   *lymphohistiocytosis in humans and mice.* Front Immunol, 2013. **4**: p. 448.
1049   36.   Chinn, I.K., et al., *Genetic and mechanistic diversity in pediatric hemophagocytic*
1050   *lymphohistiocytosis.* Blood, 2018. **132**(1): p. 89-100.
1051   37.   Park, C.Y., et al., *STIM1 clusters and activates CRAC channels via direct binding of a*
1052   *cytosolic domain to Orai1.* Cell, 2009. **136**(5): p. 876-90.
1053   38.   Delevoye, C., et al., *Conservation of the biochemical properties of IncA from Chlamydia*
1054   *trachomatis and Chlamydia caviae: oligomerization of IncA mediates interaction*
1055   *between facing membranes.* J Biol Chem, 2004. **279**(45): p. 46896-906.
1056   39.   Paumet, F., et al., *Intracellular bacteria encode inhibitory SNARE-like proteins.* PLoS One,
1057   2009. **4**(10): p. e7375.

1058   40.   Barocchi, M.A., V. Masignani, and R. Rappuoli, *Opinion: Cell entry machines: a common*
1059   *theme in nature?* Nat Rev Microbiol, 2005. **3**(4): p. 349-58.
1060   41.   Neveu, E., et al., *Prototypic SNARE Proteins Are Encoded in the Genomes of*
1061   *Heimdallarchaeota, Potentially Bridging the Gap between the Prokaryotes and*
1062   *Eukaryotes.* Curr Biol, 2020. **30**(13): p. 2468-2480 e5.

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

1089

1090

1091

1092 **Figure Legends:**

1093

Figure 1: STX11 is required for SOCE. Measurement of SOCE in STX11 depleted

Jurkat and FHLH4 patient T cells. (**A-B**) Representative Fura-2 calcium imaging assay

(**A**) and quantification of independent repeats (**B**) measuring thapsigargin (TG) induced

SOCE in Jurkat T cells treated with scr (black), STX11#1 (red) or STX11#2 (blue)

shRNAs. (n=30-40 cells/group, N=3). (**C-D**) Representative Fura-2 calcium imaging

assay (**C**) and quantification of repeats (**D**) showing reconstitution of SOCE in STX11

depleted Jurkat T cells by ectopic expression of STX11. scr shRNA with empty vector

(EV) expression (black), STX11 shRNA with EV expression (red), STX11 shRNA with

STX11 expression (green). (**E**) Sanger sequencing of FHLH4 patient DNA showing

deletion of a single Adenine at the 752nd position and the resulting frameshift. (**F**)

Schematic showing the predicted domain distribution of the wildtype *versus* FHLH4

mutant STX11 protein. (**G**) Western blot on WCLs of healthy donor (HD) and FHLH4

patient PBMCs showing the relative molecular weight and abundance of the wildtype and

mutant STX11 bands. (**H**) Western blot on WCLs of HEK293 cells over-expressing

wildtype or FHLH4 mutant STX11 and empty vector. (**I**) Fura-2 calcium imaging assay

measuring anti-CD3 induced SOCE in HD (black) and FHLH4 (red) T cells. (n=30-40

cells/group). (**J**) Fura-2 calcium imaging assay measuring SOCE in HD expressing empty

vector (EV) (black) and FHLH4 T cells expressing human STX11 (red), respectively.

(n=30-40 cells/group). (**K-L**) Representative Fura-2 calcium imaging assay (**K**) and

quantification of repeats (**L**) measuring SOCE in Jurkat T cells treated either with scr

shRNA expressing empty vector (EV) (black), STX11 shRNA, expressing EV (red),

STX11 shRNA, expressing wildtype STX11 (green), STX11 shRNA, expressing FHLH4

mutant STX11 (blue).

Figure 2. STX11 regulates I$_{CRAC}$. Effect of over-expression of STX11 on SOCE and

measurement of I$_{CRAC}$. (**A-B**) Representative Fura-2 calcium imaging assay (**A**) and

quantification of repeats (**B**) measuring SOCE in HD PBMCs, expressing empty vector

(EV) (black) or wildtype STX11 (red). (**C-D**) Dependence of STX11 mediated

enhancement of SOCE on Orai1. A representative Fura-2 calcium imaging assay (**C**) and

quantification of repeats (**D**) showing measurement of TG induced SOCE in HEK293 cells

where Orai1 was depleted and STX11 was over-expressed. Scr shRNA + empty vector

(black), Orai1 shRNA + empty vector (red), Orai1 shRNA + STX11 (blue). N=3. (**E-H**)

Measurement of I_{CRAC} in STX11 depleted Jurkat T-cells in the whole-cell recording

configuration in 20 mM extracellular Ca^{2+} Ringer's solution. I_{CRAC} was induced by passive

depletion of intracellular Ca^{2+} stores by dialyzing 8 mM BAPTA into the cell via the patch-

pipette. (**E**) Representative current at -100 mV in Jurkat T cells treated with scr shRNA

construct. The current is blocked by extracellular La^{3+} (10 µM) and replacing the 20 mM

Ca^{2+} Ringer's solution with a divalent free solution (DVF) evokes a large Na^+ current which

depotentiates over tens of seconds. The current-voltage (I-V) relationship of the Ca^{2+} and

DVF currents are shown on the right. (**F**) I_{CRAC} from a Jurkat T cell treated with STX11

shRNA. Both Ca^{2+} and Na^+ current amplitudes are reduced relative to control cells. The

I-V relationships (*right plots)* show no change in ion selectivity. (**G-H**) Summary of the

current amplitudes of Ca^{2+} and Na^+ currents and current reversal potentials in scr and

STX11 knockdown cells.

Figure 3. STX11 regulates NFAT activation, IL-2 gene expression and degranulation

in FHLH4 patient T cells. (**A-C**) Estimation of nuclear translocation of NFAT. (**A**) Western

blot showing nuclear translocation of NFAT in Jurkat T cells treated with scr or STX11

shRNA and stimulated with PMA+TG. N=3. (**B**) Representative confocal images of control

and STX11 depleted Jurkat T cells stimulated with anti-CD3, immunolabelled with anti-

NFAT1 antibody and counter-stained with DAPI. (N=3) (**C**) Box and whisker plot showing

quantification of nuclear NFAT from cells populating 10 randomly chosen fields per group

in (**B**). (**D**) Quantification of IL-2 mRNA in anti-CD3 stimulated Jurkat T cells treated with

scr or STX11 shRNA using qPCR. The bars show mean $\pm$ SE of relative IL-2 mRNA. N=3.

(E) Quantification of IL-2 EIA on supernatants of scr (black) or STX11 (red) shRNA treated

Jurkat T cells stimulated with anti-CD3. (**F**) qPCR analysis of IL-2 mRNA expression in

various stimulated HD (black) and FHLH4 T cells (red). (**G-H**) Granule release assay

performed on HD (black) and FHLH4 CD8 T (red) cells (**G**) and its quantification (**H**).

1150

Figure 4: STX11 directly binds resting Orai1 in plasma membrane. Intracellular localization of STX11 and its association with Orai1. (**A**) Representative confocal images of HEK293 cells expressing Orai1-YFP with either STX11 or empty vector (Vector), stained using anti-STX11 antibody. N=3. (**B-C**) Co-IP to assess STX11 binding to Orai1. Whole cell lysates of resting and store-depleted HEK293 cells co-expressing either Flag-Orai1 and STX11 (**B**) or Orai1-Myc and STX11 (**C**) were subjected to IP and western blot using anti-Myc, anti-Flag or anti-STX11 antibodies, as indicated. (N=3) (**D**) Schematic showing key domains of STX11 and Orai1 used for *in vitro* pull-down assays. (**E**) Pull-down assay showing *in vitro* binding of His-tagged STX11 to Orai1. (Top) Ponceau S staining showing the input of MBP alone or MBP-tagged Orai1 fragments. (Bottom) Western blot using anti-STX11 antibody. Input:1/5th of the protein. N=4. (**F**) Pull-down assay showing *in vitro* binding of His-tagged H$_{abc}$ domain of STX11 to Orai1 fragments. (Top) Ponceau S staining showing the input of MBP alone or MBP-tagged Orai1 fragments. Input:1/5th of the protein. (Bottom) Western blot using anti-His antibody. N=3. (**G**) Binding free energy distribution of STX11-H$_{abc}$ and Orai1 C-terminus interactions. (**H**) Sphere and cartoon representation of the structure of STX11 H$_{abc}$ (green) and Orai1 C-terminus (cyan) complex after MD simulation. The N-termini are highlighted in blue, and C-termini in red. (**I**) Representative confocal images showing localization of STX11 with respect to Orai1:Stim1 puncta in store-depleted cells. HEK293 cells expressing Orai1-YFP, CFP-Stim1 were transfected with STX11 and stained using anti-STX11 antibody. (N=3) (**J-K**) Binding free energy distribution for the interactions between the protomers (**J**) and dimers (**K**) of Stim1(312-387) and Orai1 C-terminus interactions. (N=3)

Figure 5. Residues involved in Orai1:STX11 complex formation are crucial for SOCE. Analysis of STX11 and Orai1 mutants. (**A**) Schematic showing STX11 mutations (Top). Expression of wildtype STX11 or STX11 N147A_E150A mutant in HEK293 followed by immunolabeling (**A**, bottom) and Western blot (**B**) using anti-STX11 antibody. (**C-D**) Representative Fura-2 calcium imaging assay (**C**) and quantification of repeats (**D**) measuring rescue of SOCE in Jurkat cells treated either with scr shRNA (black) or STX11 shRNA expressing empty vector (red), wildtype STX11 (green) or N147A_E150A mutant STX11 (blue). (**E**) Coomassie blue stained SDS-PAGE showing expression and

purification of His-tagged wildtype or N147A_E150A mutant STX11 from *E. coli.* (**F**) Pull-

down assay measuring their binding to MBP-tagged Orai1 C-terminus. (Top) Ponceau S

staining showing the input of MBP-tagged fragments. (Bottom) Western blot using anti-

STX11 antibody. N=4 (**G**) Schematic showing Orai1 mutations (Top). Confocal images of

HEK293 cells expressing YFP-tagged wildtype or R289A_E272A_E275A_E278A mutant

Orai1 (Bottom). (**H-I**) Representative Fura-2 calcium imaging assay (**H**) and quantification

of repeats (**I**) measuring rescue of SOCE in HEK293 cells treated either with scr (black)

or Orai1 shRNA expressing empty vector (red), wildtype Orai1 (green) or

R289A_E272A_E275A_E278A mutant Orai1 (blue). (**J**) Pull-down assay measuring the

binding of MBP-tagged wildtype or R289A_E272A_E275A_E278A mutant Orai1 C-

terminus to wildtype or N147A_E150A mutant STX11. (Top) Ponceau S staining showing

the input of MBP-tagged Orai1 fragments. (Bottom) Western blot using anti-STX11

antibody. N=4. (**K-L**) Representative confocal images (**K**) and quantification (**L**) of PM

localized YFP-CAD in CFP-tagged wildtype or R289A_E272A_E275A_E278A mutant

Orai1 expressing HEK293 cells.

1197

Figure 6: Functional clustering of Orai1 is compromised in STX11 depleted cells.

Quantification of Orai1 and Stim1 intensities in the ER-PM puncta of store-depleted cells.

(**A-B**) Representative confocal images of resting (**A**) and store-depleted (**B**) shRNA

treated HEK293 cells expressing N-terminal CFP tagged Orai1 (CFP-Orai1) and C-

terminal YFP tagged Stim1 (Stim1-YFP). (**C-F**) Mean Stim1-YFP intensity (**C**) and area

(**D**) of Stim1 clusters. Mean CFP-Orai1 intensities inside (**E**) and outside (**F**) Stim1-YFP

clusters of shRNA treated cells, post store-depletion. (**G-J**) Quantification of Orai1 and

Stim1 intensities in the ER-PM puncta of C-terminal YFP-tagged Orai1 (Orai1-YFP) and

N-terminal CFP-tagged Stim1 (CFP-Stim1). (**G-H**) Mean CFP-Stim1 intensity (**G**) and

area (**H**) of Stim1 clusters. Mean Orai1-YFP intensity (**I**) and area (**J**) of Orai1-YFP

clusters in control and STX11 depleted cells. (**K-L**) Representative Fura-2 calcium

imaging assay (**K**) and quantification of repeats (**L**) measuring TG induced SOCE in

Orai1-YFP and CFP-Stim1 expressing HEK293 cells treated with scr (black) or STX11

(red) shRNA.

1212

Figure 7: STX11 primes Orai1 for successful gating by Stim1. Orai:Orai FRET. **(A)**
FRET analysis of control and STX11 shRNA treated HEK293 cells co-expressing Orai1-
CFP and Orai1-YFP under resting conditions. (**B**) FRET analysis of cells described in (**A**)
before and after store-depletion. (**C-E**) Fura-2 calcium imaging assay (**C**) and
quantification of repeats (**D-E**) to measure constitutive calcium influx in Orai1-CFP and
YFP-CAD expressing, control (black) or STX11 depleted (red) HEK293 cells at 0 mM (**D**)
and 2 mM (**E**) extracellular Ca^{2+}. (**F-G**) Fura-2 calcium imaging assay (**F**) and
quantification of repeats (**G**) to measure constitutive calcium influx in Orai1-H134S mutant
expressing control (black) and STX11 depleted (red) HEK293 cells. (**H-I**) Fura-2 calcium
imaging assay (**H**) and quantification of repeats (**I**) to measure constitutive calcium influx
in Orai1-ANSGA mutant expressing control (black) and STX11 depleted (red) HEK293
cells. (**J**) Proposed model depicting STX11 unbound (unprimed) and bound Orai1
undergoing priming and correct assembly before Stim1 mediated gating. For simplicity,
interactions for only two subunits in the front have been highlighted in primed Orai1.
1227

Supplementary Movies 1-3: Molecular dynamic simulations of Orai1 C-terminus
interaction with the STX11_H$_{ABC}$ domain.

Supplementary Movies 4-6: Molecular dynamic simulations of Orai1 C-terminus
interaction with Stim1 (312-387).
Due to BioRxiv file size limit, the movies can be found at the following link:
https://www.dropbox.com/scl/fo/3j07d3sua4g58w3yfmyv7/AAKjvIiniA-
z994hxV9DNHo?rlkey=e7yj5wyukvwg5seg1awkuctc4&st=z6vp57dr&dl=0
1235

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

STX11 sufficient:

Resting Orai1 — Store depletion → Primed Orai1 — Stim1 dependent entrapment → Active Orai1

Orai1
STX11

Plasma membrane
ER-PM junction
Stim1
Endoplasmic reticulum

STX11 deficient:

Unprimed Orai1

Plasma membrane