Syntaxin11 Deficiency Inhibits CRAC Channel Priming to Suppress Cytotoxicity and Gene Expression in T Lymphocytes.

Sritama Datta[1#], Abhikarsh Gupta[1#], Kunal Mukesh Jagetiya[1], Resmi Bera[1], Vikas Tiwari[2⊥], Atharva Rahul Yande[1], Megumi Yamashita[3⊥], Abdul Rishad, Vishal Malik[1], Sreejith Raran-Kurussi[1], Sandra Ammann[4,5], Mohammad Shahrooei[6], Kalyaneswar Mandal[1], Ramanathan Sowdhamini[2], Murali Prakriya[3], Adish Dani[1], Monika Vig[1]*

Address and Affiliation: [1]Tata Institute of Fundamental Research, Hyderabad, India. [2]National Centre for Biological Sciences, Bangalore, India. [3]Northwestern University, Feinberg School of Medicine, Chicago, USA. [4]Institute for Immunodeficiency, University of Freiburg, Germany. [5]Institute for Transfusion Medicine and Gene Therapy, University of Freiburg, Germany. [6]Department of Microbiology, Immunology and Transplantation, Clinical and Diagnostic Immunology, KU Leuven, Leuven, Belgium.

[#,⊥] These authors contributed equally.

***Correspondence to:** monika.vig@gmail.com

Keywords: Orai, FHLH4, CRAC, Syntaxin11, SOCE, calcium, Stim, SNARE, ion channels, immunodeficiency, cytotoxicity, CTL, CD8 T lymphocytes

Abstract:

Mutations in Syntaxin11, a Q-SNARE, result in a fatal immune disorder known as familial hemophagocytic lymphohistiocytosis 4 (FHLH4) in human patients. A key diagnostic feature of FHLH4 is defective T and natural killer (NK) cell cytotoxicity. Here we show that Syntaxin11 directly binds and regulates Orai1, the pore forming subunit of calcium release activated calcium (CRAC) channels. CRAC channels enable store-operated calcium entry (SOCE) from the extracellular space and are crucial for granule exocytosis and nuclear factor of activated T cells (NFAT) dependent gene expression in activated lymphocytes. Syntaxin11 depletion strongly inhibited SOCE, CRAC currents, NFAT activation, interleukin-2 gene expression and degranulation in FHLH4 patient T lymphocytes and cell lines. Remarkably, defects of granule exocytosis as well as interleukin-2 expression could be reversed by ionomycin in patient T lymphocytes and a constitutively active, H134S, mutant of Orai1 rescued calcium entry in Syntaxin11 depleted cells. Further analyses showed that Syntaxin11 primes Orai1 for optimal on-site multimeric assembly which was Stim independent but required for gating. Priming of ion channel pore subunits is, therefore, a primary function of specific SNAREs which may have preceded their role in vesicle fusion.

Introduction:

Familial hemophagocytic lymphohistiocytosis 4 (FHLH4) is a life-threatening immune disease caused due to mutations of STX11 [1]. The underlying cause of FHLH is reduced cytolytic activity of T and NK cells [2], which renders patients susceptible to recurrent infections. Patients suffer from high fever, severe lymphopenia in early infancy and succumb to disease by adolescence, unless given bone marrow transplants. STX11 is highly expressed in almost all cells of the immune system and, being a Q-SNARE, is, by default, associated with exocytosis of vesicles. Intriguingly, FHLH4 patients also manifest cytokine storm, which cannot be reconciled with a presumed general defect in membrane trafficking and vesicle fusion.

Most eukaryotic cells have a limited amount of calcium sequestered inside intracellular stores, the largest of which is the endoplasmic reticulum (ER). Signaling from cell surface receptors induces the release of stored calcium which, in turn, activates store-operated calcium entry (SOCE) to replenish stores, sustain signaling and drive other calcium dependent cellular processes [3]. Calcium release activated calcium (CRAC) channels are the major conductors of SOCE [4]. In lymphocytes and mast cells, CRAC currents have been shown to be crucial for cytotoxicity, degranulation as well as gene expression [5, 6]. Working in concert with a number of accessory proteins, Orai (CRACM) multimers conduct SOCE upon functional clustering with Stim proteins in ER-PM junctions [7-9]. Yet, the prevailing thought attributes all structural transitions that result in the activation of CRAC currents to Stim proteins [10].

Genome-wide RNAi screens, which initially identified Orai (CRACM), have also yielded a wealth of information regarding additional crucial players in SOCE [8, 11, 12]. For instance, we have previously shown that alpha-soluble N-ethylmaleimide sensitive factor (NSF) attachment protein (α-SNAP), a well-known synaptic family adaptor protein that forms a part of the 20S SNARE super-complex [13], is a crucial component of the CRAC channel supramolecular complex [8, 9]. α-SNAP independently binds Stim as well as Orai and is required for the correct assembly and calcium selectivity of CRAC channels

[14-16]. α-SNAP depletion in cells revealed that Stim:Orai coupling is necessary but not

sufficient for SOCE, a process that requires distinct time-resolved molecular steps *in vivo*.

In addition to α-SNAP, SNAREs also came up as candidates in two genome wide

RNAi screens for regulators of SOCE but were not characterized [8, 12]. Furthermore,

specific SNAREs have been previously found to form inhibitory associations with specific

ion channels, but their exact role was not defined [17]. Therefore, we hypothesized a

more fundamental function for the association of SNAREs with ion channel pore subunits

and set out to test this in the context of SOCE. Using a targeted RNAi screen, biochemical

and biophysical analyses, we have found a direct role of STX11 in facilitating assembly

of CRAC channel pore subunit, Orai1, into optimal multimers. Assembly of Orai1 was

independent of, and preceded, trapping and gating by Stim1. Analyses of T cells derived

from symptomatic STX11 deficient, FHLH4 patient showed that the previously reported

[2], as well as newly identified, defects in degranulation and interleukin-2 gene

expression predominantly stem from suppressed SOCE. Based on the findings presented

here, we demonstrate a novel SNARE dependent priming step for CRAC channel pore in

the molecular sequence of SOCE and suggest preventive pharmacological interventions

for FHLH4 patients.

Results:

RNAi dependent ablation of a Q-SNARE, Syntaxin 5 (STX5), has earlier shown inhibition of SOCE in a genome-wide screen performed in *Drosophila* S2 cells [12] and STX1A has been previously shown to bind a variety of ion channels [17]. Therefore, we first knocked down STX5 (Supplementary Figure 1A) and STX1A (Supplementary Figure 1B) using different sequences of shRNA targeting each gene in HEK293 cells and measured SOCE in response to Thapsigargin (TG). In both cases, we did not see a significant defect in SOCE. Since several proteins involved in synaptic function are not ubiquitously expressed and many show redundancy in function, we searched online databases to target SNAREs that are expressed in HEK293 and/or T cells (Supplementary Figure 1C-H). We found that Syntaxin11 knockdown showed strong and consistent inhibition of SOCE by day 3 to 4 (Supplementary Figure 2A-B) (Figure 1A-B). Quantitative PCR (qPCR) analysis of total RNA extracted from scramble (scr) and STX11 shRNA treated cells showed nearly 70% depletion of STX11 mRNA (Supplementary Figure 2C) and ectopic expression of STX11 in STX11 depleted Jurkat cells rescued SOCE (Figure 1C,D). The efficiency of STX11 depletion was assessed for each experiment reported in this study. We restricted our analysis to day 3 to 4, post RNAi treatment, as longer incubations with RNAi or CRISPR knockout of STX11 affected cell viability.

Given that STX11 is highly expressed in most immune cells and is the only known SNARE that, when mutated, causes defects in T and NK cell cytotoxicity in humans and mice [2], we hypothesized that reduced SOCE contributes to the cytotoxicity defects of FHLH4 patient T cells. To test this hypothesis, we isolated PBMCs from an FHLH4 patient with a homozygous deletion frameshift mutation in the STX11 coding sequence, c.752delA:p.Lys251fs. The patient presented with all symptoms of FHLH4 disease by 4 years of age. Given the limited sample amount, we expanded PBMC numbers by periodically stimulating patient and healthy donor cells with PHA and IL-2, prior to analysis. Isolation and Sanger sequencing of DNA from patient PBMCs showed deletion of a single Adenine at position 752 in the STX11 gene which lead to frameshift following Lysine 251 (Figure 1E). The absence of any contaminating wildtype peaks showed that

the mutation was present in all blood cells. Figure 1F shows schematic of the predicted

mutant STX11 protein with an estimated molecular weight of ~39.5 KDa compared to ~33

KDa of wildtype STX11. The frameshift mutation resulted in disruption of the SNARE

domain, ablation of the terminal cysteines and elongation of the resulting transcript. We

subjected whole cell lysates prepared from the FHLH4 patient and healthy donor (HD)

PBMCs to SDS-PAGE and western blot and found that while the wildtype STX11 band

was absent, a faint but distinct band running at ~37 KDa was present in the FHLH4 patient

sample lysate (Figure 1G). Therefore, the predicted and observed molecular weight of

the FHLH4 mutant protein was higher but its levels appeared significantly reduced

compared to STX11 from HD. This is likely due to protein instability and degradation

because ectopic expression of the FHLH4 STX11 mutant in HEK293 cells showed at least

two clear bands (Figure 1H), although reduced reactivity to anti-STX11 antibody is also a

possibility. In accordance with our findings in the STX11 depleted cell lines (Figure 1A-B

and Supplementary Figure 2A-B), SOCE was found to be significantly defective in FHLH4

T cells when compared to HD (Figure 1I). Further, expression of wildtype (Figure 1J), but

not FHLH4 mutant STX11 (Figure 1K-L), reversed defective SOCE in STX11 deficient

patient (Figure 1J) and Jurkat T cells (Figure 1K-L), respectively. These data suggest that

FHLH4 mutant STX11 was functionally compromised in addition to being unstable and

no additional unknown mutations resulted in reduced SOCE.

To determine whether the inhibition of SOCE in Figure 1 resulted from changes in

the activity of CRAC channels or their expression level, we quantified the level of Orai1

expression in plasma membrane. To do this, we labelled a previously characterized

U2OS cell line stably expressing Orai1 tagged with a bungarotoxin binding site (BBS) in

the extracellular loop and YFP in the C-terminus (Orai-BBS-YFP) with Alexa-647

conjugated alpha-bungarotoxin (BTX-A647) (Supplementary Figures 3A) [9]. There was

no change in the expression of total Orai1 in the PM of STX11 depleted U2OS cells

(Supplementary Figure 3B), even though SOCE was significantly suppressed

(Supplementary Figures 3C-D). The ER-targeted CEPIA expressing control and STX11

depleted cells showed comparable ER calcium content (Supplementary Figure 3E).

Labelling STX11 depleted cells with ER, Golgi organelle specific markers also showed no

abnormalities (Supplementary Figures 3F-G) suggesting that the overall health, function

and calcium content of Golgi and ER were not adversely affected by STX11 depletion.

Previous studies have shown that SOCE in T cells is mediated by CRAC channels

formed by Orai proteins. To test whether CRAC current (I_{CRAC}) was affected by

knockdown of STX11 in T cells, we performed whole cell patch clamp recordings on

scramble and STX11 shRNA treated Jurkat T cells (Figure 2A-B). Leak-subtracted I_{CRAC}

was recorded in 20 mM Ca^{2+} and a Na^+-based divalent cation free (DVF) solution. These

recordings showed that both the Ca^{2+} and DVF CRAC currents were significantly

decreased in the STX11 shRNA treated cells with the normalized currents shown in

Figure 2C. The electrophysiological properties of the residual current in STX11 shRNA

treated cells was indistinguishable from I_{CRAC} in scramble shRNA treated cells in terms of

blockade of the Ca^{2+} current by La^{3+} and depotentiation of the DVF current over tens of

seconds. Moreover, the reversal potential of the Ca^{2+} and DVF currents were similar

between scramble and STX11 shRNA treated Jurkat T cells, suggesting that STX11

depletion does not affect the calcium selectivity of CRAC channels (Figure 2D). Together,

these results demonstrate that STX11 works by specifically modulating I_{CRAC} in T cells.

CRAC channel mediated sustained calcium influx is crucial for the activation of a

key calcium dependent transcription factor known as nuclear factor of activated T cells

(NFAT) [4, 18]. We stimulated control and STX11 depleted Jurkat T cells with

Thapsigargin (TG)+PMA, prepared nuclear, cytosolic extracts and subjected them to

western blot using anti-NFAT1 antibody (Figure 3A). Nuclear translocation of NFAT was

severely compromised in STX11 depleted T cells. In addition, immunofluorescence

imaging and quantification of total *versus* nuclear NFAT from anti-CD3 stimulated Jurkat

cells showed a defect in nuclear translocation of NFAT1 in STX11 depleted cells (Figure

3B-C). To determine whether NFAT induced gene expression was affected [4, 18], we

quantified IL-2 mRNA by qPCR and secreted IL-2 levels from culture supernatants of anti-

CD3 stimulated Jurkat T cells. STX11 depleted cells showed a significant decrease in IL-

2 gene expression as well as secretion (Figure 3D, 3E). We stimulated HD and FHLH4

patient T cells either through T cell receptor using anti-CD3 or with phorbol myristate

acetate (PMA) + Ionomycin and assessed IL-2 gene expression. Importantly, induction of

IL-2 expression was significantly defective in FHLH4 patient T cells but could be largely

restored with Ionomycin (Figure 3F). We next performed granule release assay on the *in*

vitro cultured HD and FHLH4 patient CD8 T cells in response to anti-CD3 mediated

stimulation and observed a significant defect in the FHLH4 patient CD8 T cell

degranulation (Figure 3G, 3H). To determine whether the reduced granule release results

from a direct defect in vesicle fusion or SOCE, we stimulated cells with Ionomycin along

with (PMA). Remarkably, CD8 T cell degranulation was fully restored in FHLH4 T cells

suggesting that defective SOCE primarily causes cytolytic defects in FHLH4 T cells. Of

note, the patient exhibited all symptoms of FHLH4 disease and we observed strong

suppression of SOCE (Figure 1I), IL-2 gene expression and secretion (Figure 3D-F).

However, *in vitro* culture of FHLH4 patient CD8 T cells with IL-2 has been previously

shown to partially or fully restore cytolytic defects by other groups [19, 20]. Therefore,

considering previous reports, a relatively lower (50%) defect in CD8 T cells degranulation

was not surprising [19, 20]. Collectively, these data suggest that defects in T cell

degranulation as well as interleukin-2 gene expression originate from suppressed SOCE

and significantly contribute to the immune dysregulation of FHLH4 patients.

Unlike most Q-SNAREs, STX11 lacks a transmembrane domain but harbors a

stretch of cysteines proposed to be essential for its membrane localization close to the C-

terminus (Supplementary Figure 4A) [21]. The cellular localization of STX11 is not

established because, unlike primary T cells, native STX11 is undetectable in Jurkat and

HEK293 cell lines using available commercial antibodies. Tagging STX11 on either N- or

C-terminus results in partial degradation (Supplementary Figure 4B) and complete mis-

localization of the protein (Supplementary Figure 4C). Therefore, we inserted an HA tag

inside the N-terminal unstructured region of human STX11. The internal HA tagged

STX11 as well as un-tagged STX11 localized in the plasma membrane of most HEK 293

cells (Figure 4A) and co-localized with Orai1-YFP in cells co-expressing Orai1 (Figure

4B). To test whether STX11 and Orai1 formed a complex, we co-expressed STX11 with

either Flag- or Myc-tagged Orai1 in HEK293, prepared whole cell lysates (WCLs) and

subjected them to co-immunoprecipitation (co-IP) followed by western blot using either

anti-Flag, anti-Myc or anti-STX11 antibodies. Both Orai1 and STX11 could co-IP each other (Figure 4C-D). To further assess whether STX11 directly bound Orai1, we expressed and purified from *E. coli*, MBP-tagged N- and C-terminal cytosolic tails of Orai1 (Figure 4E). *In vitro* pull-down assay performed by incubating MBP-tagged Orai1 cytoplasmic tails with full length soluble His-tagged STX11 showed that the C-terminus of Orai1 directly bound STX11 (Figure 4F). SNAREs typically utilize their SNARE domain for associating with other SNAREs [22]. To identify the domain of STX11 involved in the regulation of SOCE, we expressed His-tagged H_{abc} and SNARE domains of STX11 in *E. coli*, purified and assessed their binding to the MBP-tagged Orai1 tails using pull-down assay. We found that the H_{abc} domain of STX11 bound to the C-terminus of Orai1 (Figure 4G). The SNARE domain of STX11 did not show any binding to Orai1 and only a faint binding to SNAP23 (Supplementary Figure 4D). We next used AlphaFold3 (AF3) to obtain deeper insights into the domain interactions of STX11 and Orai1. Of all the possible combinations, the complex of STX11 H_{abc} and Orai1 C-terminus resulted in a significantly high prediction score (Supplementary Table 1). We, therefore, executed AF3 predictions with different initial seeds to generate multiple models of STX11 H_{abc} with Orai1 C-terminus. We calculated the contact frequency of interface residues in different models and found that multiple residues showed frequency greater than 0.8 (Supplementary Figure 5A). For further analysis, we considered the predicted model with the best score (Supplementary Table 2). Next, we performed all-atom MD simulation in aqueous environment (Movies 1-3) to assess the conformation, interaction stability of the STX11 H_{abc} and Orai1 C-terminus complex, which was stable throughout the simulation time suggested by RMSD (Supplementary Figure 5B), and binding free energy (ΔG) values (Supplementary Figure 6A) (Figure 4H). The individual trajectories were concatenated and clustered to obtain a centroid structure for this complex. There were 46 clusters, and the largest cluster, with 52 members, showed an elaborate protein-protein interface where STX11 H_{abc} was found to be oriented in an anti-parallel orientation to the Orai1 C-terminus (Figure 4I).

Because STX11 binds the C-terminus of Orai1 and the region largely overlaps with the Stim1 binding domain, we next sought to determine whether STX11 clusters with Orai:Stim puncta. We over-expressed HA-tagged STX11 in a Orai1-YFP and CFP-Stim1

expressing stable HEK293 cell line, store-depleted with TG, and stained using anti-HA

antibody. We did not observe co-clustering of STX11 in Orai:Stim puncta even 8 minutes

post store-depletion (Figure 4J-K). Accordingly, STX11 did not co-IP with Stim1 in store-

depleted cells (Supplementary Figure 7A). These data suggest that STX11 comes off

Orai1 C-terminus prior to Stim1 dependent clustering of Orai1. Since co-IP studies

suggested sustained interaction between STX11 with Orai1 (Figure 4B-C), we quantified

the fraction of Orai1 that resides inside Stim:Orai puncta w.r.t total Orai1 in PM and found

that post TG addition, nearly 60% of Orai1 remains outside of Stim1 puncta even in

unmanipulated cells (Supplementary Figure 7B). We used AF3 to generate the complex

of Stim-Orai activating region (SOAR) domain of Stim1 and Orai1 (Supplementary Figure

8A). Given that SOAR is a dimer under resting state, we used two copies each of SOAR

and Orai1 [23] [24]. The prediction scores for their complex are shown in Supplementary

Table 3. The best scoring model among multiple runs of AF3 was considered for MD

simulations on the complex of Orai1 C-terminus (256-301) and SOAR. The ΔG values for

the complex are shown in (Supplementary Figure 8B-C). Three independent MD

simulations were run for the complex (Movies 4-6) which showed an initial conformational

change which stabilized thereafter (Supplementary Figure 9A). All three simulation

trajectories were concatenated and clustered to obtain a centroid structure for this

complex (Supplementary Figure 9B). There were 43 clusters, and the largest cluster had

77 structures. Collectively, these data suggest that STX11 directly binds Orai1 with an

affinity that is physiological but relatively poorer when compared to the binding of SOAR

dimers to Orai1. Therefore, presence of high density of SOAR dimers in ER-PM junctions

of store-depleted cells could potentially outcompete STX11:Orai1 interactions by

progressively allowing formation of SOAR:Orai1 oligomers. It is also possible that

another, yet to be characterized, molecular step and/or protein actively catalyzes the

removal of STX11 from Orai1 prior to its entrapment in ER-PM puncta. Future studies will

be able to systematically assess these possibilities.

The major interactions predicted between STX11 and Orai1 from all-atom MD

simulations are highlighted in Supplementary Figure 6B-C. To assess the significance of

these interactions, we performed site-directed mutagenesis of two key residues of STX11

predicted to be involved in binding to Orai1 (Figure 5A). Ectopic expression of untagged

N147A_E150A STX11 showed correct localization (Figure 5A) as well as stable

expression (Figure 5B) but failed to rescue SOCE in STX11 depleted Jurkat T cells

(Figure 5C-D). We next expressed His-tagged wildtype and N147A_E150A mutant

STX11 in *E. coli.*, purified (Figure 5E) and assessed their ability to bind MBP-tagged Orai1

C-terminal tail in an *in vitro* pull-down assay (Figure 5F). Compared to wildtype STX11,

N147A_E150A mutant STX11 showed reduced binding to Orai1 C-terminus. Further, we

introduced four mutations, R289A_E272A_E275A_E278A, in the Orai1 C-terminus

(Figure 5G), tagged with YFP and expressed in HEK293 cells. The Orai1 mutant localized

to PM (Figure 5G) but failed to rescue SOCE in Orai1 deficient cells (Figure 5H-I). We

expressed and purified MBP-tagged R289A_E272A_E275A_E278A mutant Orai1 C-

terminus from *E. coli* and assessed its ability to bind purified wildtype as well as

N147A_E150A mutant STX11 in a pull-down assay (Figure 5J). Binding of Orai1 mutant

to both was significantly compromised. Because Stim1 binds an overlapping region of the

Orai1 C-terminus, we next tested whether mutation of Orai1 residues affected Stim1

binding. We co-expressed YFP-tagged soluble CRAC activation domain (CAD) of Stim1

(YFP-CAD) (Figure 5K) with either CFP-tagged wildtype or mutant Orai1 in HEK293 cell

line. YFP-CAD localizes to the cytosol in HEK293 cells but in cells overexpressing Orai1,

a significant fraction of it localizes to plasma membrane due to its association with Orai1

[10]. We quantified relative YFP-CAD levels in the plasma membrane of both groups

(Figure 5K-L) but found no difference in the PM recruitment of CAD suggesting that the

inability of R289A_E272A_E275A_E278A mutant Orai1 to support SOCE does not result

from defective binding to CAD/SOAR. Collectively, these data demonstrate a direct role

of STX11 in the binding and regulation of Orai1.

Upon store depletion, Stim1 localizes to ER-PM junctional regions where it traps

and co-clusters with Orai1. To further understand the mechanism, we first imaged

HEK293 cells expressing CFP-Orai1, Stim1-YFP under resting conditions and then

quantified the levels of Orai1 and Stim1 in ER-PM junctions (puncta) of store-depleted

control and STX11 depleted cells. The distribution of Orai1 and Stim1 was normal in

resting cells (Figure 6A). However, in store-depleted cells, clustering of CFP-Orai1 was

visibly defective in STX11 RNAi treated cells (Figure 6B), even though the intensities

(Figure 6C) and area of Stim1-YFP puncta were normal (Figure 6D). Quantification of

Orai1 intensities confirmed a defect in entrapment (Figure 6E) with relatively higher
fraction of Orai1 present outside Stim1 puncta in STX11 depleted cells (Figure 6F). To
determine whether reduced clustering of Orai1 resulted from structural defects within ER-
PM junctions, we reversed the direction of fluorescent tags on Orai1 as well as Stim1 and
repeated the experiment in Figure 6A-F. Interestingly, the defect in Stim1 mediated
entrapment of Orai1 could be rescued (Supplementary Figure 10A-B) (Figure 6G-J) but
SOCE remained significantly inhibited even when Orai1 and Stim1 were co-
overexpressed (Figure 6K-L). Taken together, these data show that STX11 binding to
resting Orai1 has implications beyond mere co-entrapment of Orai1 with Stim1. STX11
depletion did not alter the structure or proximity of junctional ER to the PM and yet
reduced SOCE even in cells co-overexpressing Orai and Stim that have been proposed
to be sufficient for amplifying CRAC currents in earlier studies [25] [26] [27].

Defects in the functional entrapment of Orai1 suggest the absence of a necessary
molecular transition within Orai1 which precedes its interaction and gating by Stim1. To
test whether STX11 induced structural changes within CRAC channel pore subunit,
Orai1, we co-expressed Orai1-CFP along with Orai1-YFP in scramble control or STX11
depleted HEK293 cells and measured Orai:Orai FRET. We found significantly higher
Orai1:Orai1 FRET in STX11 depleted resting cells even in the absence of ectopically
expressed Stim1. (Figure 7A). We have previously shown that store-depletion induces
on-site assembly of Orai1 dimers into oligomers [14]. Notably, store-depletion induced a
similar level of increase in Orai1:Orai1 FRET in scramble control cells. To confirm these
findings, we performed acceptor photobleaching analysis on Orai1-CFP (donor), Orai1-
YFP (acceptor) co-expressing HEK293 resting cells where we had stably knocked down
the expression of endogenous Orai1 and Stim1. Although the trends were largely similar,
the differences in Orai1:Orai1 FRET were much more significant in these cells (Figure
7B-C). These data demonstrate a previously unsuspected, novel, molecular transition in
Orai1 necessary for CRAC channel activation, which we define as 'priming'. Further,
these data show that priming precedes the interaction of Orai1 with Stim1 in ER-PM
junctions. To determine whether 'priming' simply induced a structural change or facilitated
correct assembly of Orai1 multimers, we expressed Flag-Orai1 in control or STX11

depleted resting cells, mechanically lysed to isolate membranes, crosslinked Orai1
oligomers using BS3 and resolved using SDS-PAGE (Figure 7D) (Supplementary Figure
11A). We found a greater percentage of higher order oligomers in STX11 depleted cells
suggesting stoichiometric changes (Figure 7E) (Supplementary Figure 11A).

Increasing the density of Stim1 has been earlier proposed to bypass the
requirements for accessory factors in SOCE. To determine whether increasing the levels
of Stim1 could compensate for STX11 depletion, we expressed Orai1 tethered to two
Stim-Orai activating region (SOAR) domains of Stim1 and GFP (Orai1-S-S-GFP)
(Supplementary Figure 11B-C) in HEK293 cells. Orai1-S-S-GFP has been previously
shown to constitutively activate Orai1 [23]. In STX11-depleted Orai1-S-S-GFP expressing
cells, the magnitude of calcium entry was significantly smaller (Supplementary Figure
11D-E). To determine whether reduced calcium entry resulted from a defect in the ability
of SOAR to bind Orai1, we expressed YFP-CAD, in control or STX11 depleted Orai1-CFP
expressing HEK293 cell line (Figure 7F). Quantification of PM localized YFP-CAD as a
fraction of total CAD showed no defect in STX11 depleted cells (Figure 7F). Yet, as seen
with Orai1-S-S-GFP, constitutive calcium entry was significantly reduced (Figure 7G-H).
We next expressed a constitutively active H134S mutant of Orai1 (Supplementary Figure
11F) as it harbors an open pore and the C-terminal cytosolic tails of Orai1 are unlatched,
pointing towards the cytosol [28]. Remarkably, expression of H134S Orai1 completely
restored constitutive calcium influx in STX11 depleted cells (Figure 7I-J). To further
establish whether binding with STX11 induces straightening of Orai1 tails or molecular
transitions within transmembrane helices, we expressed in STX11 depleted cells the
constitutively active ANSGA (261-265) mutant of Orai1 (Supplementary Figure 11G),
which harbors 4 consecutive mutations in the Orai1 C-terminus resulting in the
straightening of the C-terminal tail along with constitutive activation of CRAC currents
[29]. The constitutive calcium entry in STX11 deficient Orai1 ANSGA mutant expressing
cells was significantly suppressed (Figure 7K-L). Taken together, these data show that
STX11 induces a molecular switch in the pore-forming subunit, Orai1, which precedes
and primes it for optimal on-site assembly into multimers and subsequent gating by Stim1.
Such SNARE dependent priming of Orai1 is crucial and cannot be compensated by

merely increasing the density of Stim proteins bound to Orai or by introducing mutations
that alter the orientation of Orai1 tails *in vivo*. The proposed models depicting unprimed
and primed Orai1 (Figure 8A) and its role in T cell function (Figure 8B) are shown.

Q-SNAREs are often found in complex with other SNAREs such as SNAP23/25/29
in target membranes. Some previous studies have implicated the complex of SNAP23/25
and STX1A in the regulation of specific ion channels [30, 31]. The three SNAPs,
SNAP23/SNAP25/SNAP29 are each capable of contributing two SNARE domains to the
trans- and *cis*-SNARE complex. RNAi mediated depletion of SNAP23/SNAP25/SNAP29
did not show a significant reduction in SOCE of HEK 293 cells (Supplementary Figure
12A-C). To determine whether SNAP23/SNAP25/SNAP29 might still form a part of the
STX11:Orai complex to regulate SOCE, we co-expressed myc-tagged Orai1 with
untagged SNAPs and performed respective co-IPs. We did not find any interaction
between either of the three SNAPs and Orai1 (Supplementary Figure 12D). Similar
studies were performed by co-expressing YFP-Stim1 and the three myc-tagged SNAPs.
Again, no interaction was found between any of the SNAPs and Stim1 (Supplementary
Figure 12E). These data show that, unlike in the case of vesicle fusion, STX11 does not
collaborate with SNAP23/SNAP25/SNAP29 in the regulation of SOCE.

FHLH is a heterogeneous immune disorder where diverse genes such as *Prf1*,
STX11, *UNC13D*, *Munc18-2*, *MAGT1, XIAP, LYST, AP3B1, RAB27A*, *ITK*, amongst
several others, are broadly grouped together based on disease symptoms, although the
age of onset and severity varies across the spectrum [32, 33]. Congenital
immunodeficiency involving defects in the CTL and NK cell cytotoxicity appear to be the
common factors underlying the development of FHLH, although the individual pathways
involved are highly diverse. Using shRNA, we screened 6 additional genes, *Perforin1*,
RAB27A, *UNC13D* and *Munc18-2*, *Synaptotagmin11* and *13*, that have been either
associated with vesicle fusion or routinely grouped together with STX11 in diagnosing
FHLH (Supplementary Figure 13). SOCE was not affected by depletion of most genes.
Future studies will evaluate the role of others.

Discussion:

CRAC channels conduct a small, highly specific calcium current in response to the depletion of intracellular calcium stores. According to the prevailing view, Orai multimers reside in the PM and depend on the ER resident store sensor, Stim, for all structural transitions leading up to their activation [10]. Unlike other ion channels, the process of activation of CRAC currents is exceptionally slow. For instance, following break-in or store-depletion, on average, it takes ~5 minutes for Stims to cluster in the ER-PM junctions and for measurable CRAC currents to flow from Orai1. During this time, Stims undergo intramolecular transitions and slowly trap and gate freely diffusing Orai. It is, therefore, believed that no molecular transitions take place within the pore forming subunit, Orai, until its entrapment by Stim in the ER-PM junctions.

We have shown that in STX11 deficient cells, the basal Orai:Orai FRET is significantly higher even in the absence of Stim1 or store-depletion. These data establish that Orai1 undergoes a, hitherto unrecognized, SNARE dependent molecular transition which precedes trapping and gating by Stim1. We have earlier shown that store-depletion induces dynamic on-site assembly of Orai1 dimers into oligomers [14]. In STX11 depleted resting cells, the ~25-50% increase over basal Orai:Orai FRET largely results from a change in the oligomerization of Orai1 because the proportion of BS3 crosslinked Orai1 oligomers was also found to be shifted towards higher molecular weight. It is likely, though, that a change in the overall conformation of Orai1 protomers facilitated the change in store-dependent Orai assembly and contributed to the overall change in Orai:Orai FRET.

The change in Orai:Orai FRET induced by STX11 depletion not only persisted but increased in Orai1, Stim1 double deficient cells. Furthermore, STX11 bound a region of Orai1 C-terminus that has been previously shown to bind Stim1. Therefore, our data suggest that STX11 induced molecular transitions in Orai are independent of and precede its association and clustering by Stims. Because tagging of STX11 with fluorescent tags on either end results in mis localization and degradation, our study could not demonstrate

the real-time removal of STX11 from the Orai1- C-terminus to allow for Stim1 binding in

live cells. It is also unclear whether binding or removal of STX11 induced the proposed

molecular transition defined as 'priming' of Orai1. Further, it remains to be determined

whether STX11 removal is actively induced by an additional chaperone or passive

displacement from the formation of relatively higher avidity oligomeric associations

between Orai1:Stim1 in ER-PM junctions. Future studies will be able to systematically

evaluate each of these possibilities.

We have shown that the C-terminal cytosolic tail of Orai1 directly couples with

STX11 in resting cells.   Although binding appeared reasonably strong in co-IPs,

STX11:Orai1 interaction appeared somewhat weak in pull-downs. This is likely due, in

part, to the instability of purified His-STX11 in solution which necessitated shorter co-

incubation times during pull-down assays. Instability could arise from the absence of a

stabilizing co-chaperone, such as one of the known binding partners of SNAREs [34] [35]

or membranes to which STX11 attaches in lives cells *via* its C-terminal cysteines [21].

Together, these factors appear crucial for the stabilization of purified STX11 *in vitro*.

Stim1 binds the N- as well as the C-terminus of Orai1 but their individual relevance

in the trapping *versus* gating of Orai1 remains unclear [36]. Though crucial for trapping

and gating, Orai1 tails were missing from early structures of *Drosophila* Orai [28]. A

previous NMR structure suggested that C-terminal tails of two adjacent Orai1 subunits,

bend, pair with each other in an antiparallel fashion and sit closely apposed to PM [37].

However, in recent structures of constitutively active H134 mutant Orai, the C-terminal

tails were found to orient away from the membrane [28]. In STX11 depleted cells,

switching the CFP-tag from the Orai1 N- to the C-terminus could rescue its clustering but

not gating by Stim1. Furthermore, STX11 depletion inhibited the constitutively active

ANSGA mutant of Orai1 [29], where the tails of Orai1 are proposed to be constitutively

unlatched. These data essentially reinforce our conclusions that STX11 induced

molecular shifts encompass Orai1 transmembranes.

Using Fura-2 imaging assays in STX11 depleted cells, we have conclusively
established that STX11 is necessary as well as limiting not only in live and unperturbed
HEK, U2OS, Jurkat and primary T cells but also HEK293 cells co-overexpressing Orai1
and Stim1. Previous whole cell patch clamp analysis of HEK293 cells co-overexpressing
Orai1 and Stim1 showed up to 50-100-fold amplification of CRAC currents whereas Fura-
2 based imaging of live cells routinely shows only 2-fold amplification [25] [26] [27]. The
patch clamp studies performed on Stim, Orai over-expressing cells are often extrapolated
to conclude that all other accessory factors that have since been identified are
unnecessary for native CRAC currents. A major problem with this interpretation is that
even though amplification of CRAC currents was shown, none of the previous patch
clamp studies established whether the higher currents resulted from a greater number of
active channels or unchecked conductance per channel by performing single channel
recordings. The fact that recording of I_{CRAC} in the whole cell configuration involves use of
20mM extracellular calcium and dialysis of cells with high concentrations of BAPTA, a
fast calcium chelator, the calcium dependent inactivation of CRAC channels, a key
mechanism of regulation of I_{CRAC} magnitude in live cells is disrupted. Furthermore,
hyperpolarizing potentials and voltage clamping used to record CRAC currents prevents
the progressive depolarization of cells due to influx of cations which would otherwise
dampen the driving force for CRAC currents in intact cells. Given these limitations, Fura-
2 assays offer a more realistic and physiological reflection of SOCE amplitudes in live
cells as they allow for these regulatory mechanisms, which are necessary for the
regulation of native currents to be still operational. While patch clamp remains a direct
method of verification of CRAC currents, exercising logic in extrapolation of amplitudes
of I_{CRAC} using patch clamp is imperative for the field to move forward.

Most human mutations in the *STX11* coding region result in the complete loss or
severe depletion of STX11 protein levels [21]. The FHLH4 disease, therefore, results from
reduced STX11 expression in most reported patients. Targeted mutagenesis of the
STX11:Orai1 interface lining residues based on the AlphaFold 3 model, however, didn't
destabilize   or   mis-localize   either   N147A_E150A   STX11   or
R289A_E272A_E275A_E278A Orai1 and yet both mutants showed reduced binding to

their respective wildtype counterparts. Importantly, the R289A_E272A_E275A_E278A
mutant of Orai1 retained the ability to bind Stim1.

In addition to the established cytolytic defects in FHLH4 T cells, for the first time,
we have shown that STX11 deficient, FHLH4 T cells harbor a severe defect in store-
operated calcium signaling and IL-2 gene expression. Our data strongly suggest that
SOCE defects significantly contribute to FHLH4 disease progression *via*, as well as
independent of, exocytotic/ degranulation defects. We propose that design and early
administration of CRAC channel agonists could potentially provide an alternative to
immunosuppressants and bone marrow transplants for FHLH4 patients.

In summary, we have shown that binding to a Q-SNARE, STX11, provides an on-
site molecular switch in the CRAC channel pore, Orai1, necessary for its optimal
assembly and function. We suggest that ion channel priming is a novel, direct and primary
role of specific SNAREs. In accordance with this hypothesis, early SNARE-like proteins
have been found in bacteria, which lack an endomembrane system but express a variety
of ion channels [38-40]. SNAREs are now thought to have evolved from a common
archaeal precursor found in the genomes of Asgard and members of *Legionella* [41].

Materials and Methods

Plasmid constructs and transfection

Stim1 CAD was sub-cloned in eYFPC1 vector after amplification from YFP-Stim1 plasmid to generate YFP-CAD. pLKO.1, psPAX2 and pMD2.G were purchased from Addgene. pLKO.1 cloned shRNA sequences targeting the genes of interest, were either purchased from Horizon Discovery, UK or designed and cloned in-house using the Broad Institute portal (https://portals.broadinstitute.org/gpp/public/gene/search). Full-length human STX11 was cloned from human cDNA prepared from HEK293 cells and subcloned into pcDNA3.1(+), pcDNA/4TO/Myc-HisA (Invitrogen, Grand Island, NY), pMSCV-IRES-mcherry (Addgene), pEF1alpha-IRES (Clontech) and pET28b vector [16] with an N-terminal 6XHis-tag. The fragments of STX11 were cloned in pET28b vector with an N-terminal 6xHis tag for expression in *E. coli*. HA tag was inserted between H30 and G31 in STX11 cloned in pMSCV-IRES-mCherry construct using PCR. H134S and ANSGA mutants of Orai1 were generated using site directed mutagenesis and PCR, respectively. The fragments of Orai1 (1-87, 1-47, 48-103, 256-301 and 272-292) were amplified from full length constructs and cloned into pMAL-c5X vector (New England Biolabs), in-frame with MBP protein coding sequence, as described previously [16]. SNAP23, SNAP25 and SNAP29 cDNA cloned in pCMV-Sport6.1 vectors were purchased from Dharmacon and subcloned into pcDNA4/TO/Myc-HisA. SNAP23 was subcloned in pMAL-c5X vector in-frame with MBP. All plasmid DNA transfections in human cell lines and primary cells were done using Lipofectamine 2000 (Invitrogen)/ Lipofectamine 3000 (Invitrogen) or Amaxa nucleofection kit (Lonza, Basel, Switzerland) respectively, as per manufacturer's protocol.

Cell Lines

Lentiviral shRNA transduction experiments were performed in HEK293, U2OS or Jurkat (ATCC, Manassas, VA) cell lines cultured in low glucose DMEM (Hyclone, Logan, UT) or RPMI (Hyclone) respectively, with 5-10% fetal bovine serum (Hyclone), 1X Penicillin Streptomycin, and GlutaMax (Gibco, Grand Island, NY). Stable cell lines generated using HEK293 and U2OS parent lines have been described previously [9]. HEK293-FT cell line was cultured in high glucose DMEM with 10% FBS and 10mM HEPES, 1X Penicillin

589   Streptomycin, 1X GlutaMax and 1X non-essential amino acid and transfected with the

590   appropriate plasmids to generate viral supernatants. All cell lines were tested for

591   mycoplasma contamination twice every year and found to be negative.

593   **Antibodies and Reagents**

Antibodies/Reagents	Source and Cat no	Dilution/ Conc.
Mouse anti-myc (9E10)	Abcam #ab32	1:1000
Mouse anti-myc (9E10)	Culture supernatant, in house	1:1000
Rabbit anti-SNAP23	SYSY#111203	1:1500
Mouse anti-SNAP25	SYSY#111011	1:1500
Rabbit anti-SNAP29	SYSY#111303	1:1500
Rabbit anti-GFP	Invitrogen#A11122	1:2500
Rabbit anti-flag	Sigma#F7425	1:1600
Mouse anti-flag (M2)	Sigma#F3165	1:1500
Mouse anti-6X His	Invitrogen #MA1-135	1:2000
Rabbit anti-Stx11	Sigma#SAB4301593	1:1000
Mouse anti-NFATc2 (4G6-G5)	Santacruz#7296	1:500
Rabbit anti-NFAT1	Cell Signaling Technology #4389	1:200
Goat anti-Lamin B (M-20)	Santacruz#6217	1:500
Rabbit monoclonal anti-HA	CST#3724S	1:1000
Alpaca VHH anti-rabbit AF488	Jackson #611-544-152	1:400 (1.375μg/ml)
DAPI	Invitrogen #D1306	2μg/ml
Rabbit anti-Stx11	SySy #110113	1.67μg/ml
Rabbit anti-Stx11	Invitrogen #PA5-50800	1:2000
Goat anti-rabbit A647	Invitrogen #A-21245	1:800
Rabbit anti-Stx11	Proteintech #13301-1-AP	0.3μg/ml
Donkey anti-rabbit A647	Jackson #711-605-152	1:800

HRP GAPDH	Sigma #G9295	1:40000
HRP Donkey anti-Rabbit	Jackson #711-035-152	1:50000
HRP Donkey anti-Mouse	Jackson #715-035-151	1:50000
Purified Human IL-2	BD #555051	1 μg/ml
Purified Human IL-2 Biotin	BD #555040	1 μg/ml
Streptavidin HRP antibody	Jackson #016-030-084	0.1 μg/ml
Ultra-LEAF Purified anti-human CD3	Biolegend #317326	5-10μg/ml
Ultra-LEAF Purified anti-human CD28	Biolegend #302934	2μg/ml
Ultra-LEAF Purified anti-human CD49d	Biolegend #304340	2μg/ml
PE anti-human CD107a (LAMP-1)	Biolegend #328608	1:50
APC anti-human CD8 antibody	Biolegend #344722	0.5μg/ml
Fura-2-AM	Invitrogen #F1221	1μM
Thapsigargin (TG)	Sigma #T9033-1MG	1μM
Polybrene (Hexadimethrine-bromide)	Sigma #107689-10G	8μg/ml
NE-PER kit	Thermo #78833	
Protein A/G Mag Sepharose xtra	Cytiva #28967056/66	
Protein A/G 4 Fast Flow Sepharose	Cytiva #17528001	
Protease Inhibitor Cocktail (Mammalian)	Sigma #539134	
Protease Inhibitor Cocktail (Bacterial)	Calbiochem #539132	
Talon Beads	Takara #635502	
Dextrin Sepharose Beads	Cytiva #28935597	

HEPES	Sigma #H3375-100G	
Lipofectamine 2000	Thermofisher #11668019	
Lipofectamine 3000	Thermofisher #L3000001	
Lemo21(DE3) *E.coli.* cells	NEB #C2528J	
IPTG	Himedia#MB072-10G Sigma #16758-10G	
Sarkosyl (N-lauroylsarcosine sodium salt)	Sigma #L9150-50G	
Lysozyme	Sigma #62970-5G-F	
DNaseI Recombinant RNase-free	Roche #04716728001	
PMA (Phorbol Myristate Acetate)	Sigma #P8139	20-50ng/ml
Ionomycin	Sigma #407952	$1\mu M$
Human IL-2 protein	Acro Biosystems #H5215	50ng/ml
Lectin from *Phaseolus vulgaris* (PHA)	Sigma #L8754	$2\mu g/ml$
Retronectin	Takara #T100B	$20\mu g/ml$
RNeasy Plus Mini kit	Qiagen#74134	
QIAshredder	Qiagen#79654	
Superscript IV	ThermoFisher#18091300	
Qubit ssDNA Assay kit	ThermoFisher# Q10212	
Human STX11 TaqMan probe (FAM-MGB)	ThermoFisher# Hs00186823_m1	
Human Beta Actin TaqMan probe(FAM-MGB)	ThermoFisher# Hs01060665_g1	
Human IL2 Taqman probe (FAM-MGB)	ThermoFisher# Hs00174114_m1	
Paraformaldehyde	Electron Microscopy Sciences #157-8	

Bovine Serum Albumin	Jackson #001000173	
Luria Bertini Broth	Himedia #M1245	
HBSS	Sigma #H8264	
Terrific Broth	Sigma #T9179	
Alexa fluor 647 conjugated Alpha Bungarotoxin	Invitrogen #B35450	1μg/ml
HRP rabbit anti-GFP	Invitrogen #A10260	
HRP Goat anti-rabbit Fc	Jackson #111-035-008	1:50,000
Ampicillin	Sigma #A0166-5G	100ug/ml
Kanamycin	Sigma #K1876-1G	50ug/ml
Chloramphenicol	Sigma #C 0378-5G	100ug/ml
30% Acrylamide:Bis Solution (29:1)	Biorad #1610156	
DTT (Dithiothreitol)	Biorad #1610611	
0.05% Trypsin-EDTA	Gibco #25300054	
100X Penicillin Streptomycin	Sigma #P4333-100ML	
Puromycin	Gibco #A11138-03	
Pluronic F-127 (20% solution in DMSO)	Invitrogen #P3000MP	0.02%

Lentiviral transductions

For the generation of lentiviral supernatants, shRNAs cloned in pLKO.1 were co-transfected with psPAX2 packaging and pMD2.G envelope plasmids into HEK293-FT cells using the calcium phosphate method of transfection. 48- and 72-hours post-transfection, viral supernatants were collected, pooled and stored at -80°C till further use. For transduction of HEK293 and Jurkat cells, 0.1 million cells were plated in 6- or 24-well plates either the day before (HEK293) or the same day (Jurkat). Viral supernatants were added to the cells along with 8ug/ml polybrene and cells were spun at 2500 RPM, 30°C for 90 mins. 24 hours or 48 hours post-spinfection, Puromycin was added at a final

605 concentration of 1ug/ml to HEK293 and Jurkat cells, respectively. Cells were analyzed 3-

606 5 days post transduction.

Immunocytochemistry and confocal imaging

For STX11 localization, resting or store-depleted (1μM Thapsigargin) WT HEK293 or

HEK293 cells stably expressing Orai1-YFP were either plated in 6 well plates and

spinfected with viral supernatants generated from pMSCV-STX11(HA)-IRES-mCherry,

pMSCV-IRES-mCherry (empty vector control) or plated in carbon coated, glow

discharged 35mm glass bottom dishes (IBDI) and transfected with pEF1alpha-STX11-

IRES-mCherry. 24 hr post-spinfection, cells were plated in 35mm glass bottom dishes.

For immunolabelling, cells were washed with Ringer's buffer, fixed with 4% PFA and

blocked with 3% BSA containing 0.1% NP40 for 1.5 hrs. Post-fixation and

permeabilization, cells were incubated either with anti-HA (CST) or rabbit anti-Stx11

(SySy/ Proteintech) primary antibodies O/N at 4°C, washed and incubated with either

Alpaca VHH anti-rabbit AF488, donkey anti-rabbit A647 or goat anti-rabbit A647

secondary antibodies for 1 hour at room temperature (RT). Cells were counter stained

with DAPI for 10 min at RT and imaged using Olympus FV3000 laser scanning confocal

microscope. Images were acquired sequentially using the following parameters: DAPI

(DM405/488 dichroic, 405nm excitation, 430-470nm emission); anti-HA Alexa488

(DM405/488, 488nm excitation, 500-590nm emission); STX11 (DM405/488/561/640,

640nm excitation, 650-750nm emission); Orai1-YFP (DM405/488/561/640, 488nm

excitation, 521-591nm emission); CFP-Stim1 (DM405-445/514/594, 405nm excitation,

448-510nm emission). For STX11-N147A-E150A mutant, HEK293 cells were spinfected

with viral supernatants and imaged as described above where WGA (Wheat germ

agglutinin) conjugated with Alexa 488 was used to mark plasma membrane.

Quantification of Orai1 and Stim1 intensities inside/outside ER-PM puncta

HEK293 cells stably expressing Orai1-YFP with CFP-Stim1 or CFP-Orai1 with Stim1-YFP

were plated in 6 well plates and transduced with scramble or STX11 shRNA. 48hr post

transduction, cells were trypsinized and plated in carbon coated 35mm dishes. Images

were acquired using a TIRF microscope setup described before [14]. Briefly, resting cells

were imaged first and positions were marked. Store-depleted images were acquired after
incubation with 1 µM Thapsigargin (TG) and 10mM EGTA for ~6 mins. Analysis was done
using Otsu local thresholding in ImageJ software. Stim1 and Orai1 intensity (AU) and area
µm^2 values obtained from the analysis were plotted using Origin software.
640

Orai:Orai FRET
Scramble and STX11 shRNA transduced HEK293 cells were plated on freshly carbon
coated, glow discharged 35mm glass bottom dishes (IBDI) and transfected with 0.5 µg
each of Orai1-CFP and Orai1-YPF plasmid DNA using Lipofectamine 3000. 16 hours post
transfection, co-transfected cells were imaged using PlanApoN 60XO/1.42 NA
objective with Olympus FV3000 laser scanning microscope. Orai1-CFP, Orai1-YFP and
FRET images were acquired with the following excitation laser and emission bandwidth
parameters using DM405-445/514 dichroic mirror: CFP Ex-405nm Em-460-630nm; YFP
Ex-514nm Em- 530-630nm; FRET Ex-405nm, Em-530-630nm. ROIs representing the
plasma membrane rim were drawn around each cell, followed by local thresholding using
Bernsen's algorithm in Fiji software to generate an image mask which was applied to
CFP, YFP, FRET images to extract mean intensity values. The background subtracted
mean intensities of FRET (I_{FRET}), CFP (I_{Donor}) and YFP($I_{acceptor}$) images were measured
for each cell. Cells singly expressing Orai1-CFP or Orai1-YFP only were used as CFP
donor or YFP acceptor channel bleed-through (BT) controls. The BT was calculated using
the following formula:

$BT_{Donor} = \frac{I_{FRET}}{I_{Donor}}$ when the Orai1-CFP donor alone is expressed,

$BT_{acceptor} = \frac{I_{FRET}}{I_{acceptor}}$ when the Orai1-YFP acceptor alone is expressed.

FRET and NFRET percentage was calculated by subtracting the contributions of CFP
and YFP bleed through from the FRET signal using the following:

$$FRET = I_{FRET} - (BT_{Donor} * I_{Donor}) - (BT_{Acceptor} * I_{Acceptor})$$

$$NFRET\% = \left(\frac{FRET}{\sqrt{\left(I_{Acceptor} * I_{Donor} \right)}} \right) * 100$$

Acceptor Photobleaching

Scramble and STX11 shRNA transduced, Orai1, Stim1 double knockdown HEK293 cells

where were plated on Poly-D-Lysine coated glass bottom dishes (Cellvis) and transfected

with 0.5 ug DNA each of Orai1-CFP and Orai1-YPF using Lipofectamine 2000. 16 hours

post transfection, co-transfected cells were imaged using PlanApoN 60XO/1.42 NA

objective with Olympus FV3000 laser scanning microscope. Images were acquired with

the following excitation laser and emission bandwidth parameters using DM405-445/514

dichroic mirror: CFP Ex-405nm Em-460-508nm; YFP Ex-514nm Em- 530-630nm, with

4% offset. The pre bleach images of Orai1-CFP (the donor) and Orai1-YFP (the acceptor)

were captured, sequentially. This was followed by bleaching of the acceptor, Orai1-YFP,

using 100% 514 laser power for 2 mins and then acquiring the donor and acceptor images

post bleach. ROIs were drawn around the cell periphery using the Orai1-CFP image.

Local thresholding was performed using Bernsen's algorithm to generate binary masks,

which were then applied to the CFP images to measure mean intensities. The background

subtracted mean intensities of CFP images were measured for each cell and the FRET

efficiencies were calculated as follows: E (%) = (1 -I_{before}/I_{after}) *100. I_{before} and I_{after} denotes

the average fluorescence intensity of the donor in the plasma membrane before and after

acceptor photobleaching.

Alpha-bungarotoxin binding assay

U2OS cell line stably expressing Orai1-BBS-YFP construct has been described

previously[14]. Briefly, Orai1-BBS-YFP expressing U2OS cells were either transduced with

scramble shRNA/ STX11 shRNA/ pMSCV-IRES-mcherry (EV) or pMSCV-STX11-IRES-

mcherry. Transduced cells were trypsinized either 3-4 days post-transduction or 48 hour

post-transduction (in the case of pMSCV), centrifuged and labelled on ice with AF647

conjugated alpha-bungarotoxin (1μg/ml) containing 2% FBS in 1X HBSS for 30 mins.

Following this, cells were washed with HBSS and fixed with 4% PFA (~20 min at RT).

Samples were run using Cytoflex FACS analyzer (Beckman Coulter) and data were

analyzed using Flow Jo software.

Validation of STX11 knockdown using qPCR

Scramble and STX11 shRNA treated HEK293 cells were trypsinized, counted and equal

number of cells were used for lysis. The lysates were homogenized using QIA shredder

columns (Qiagen) and RNA isolation was done using RNeasy Plus Mini kit (Qiagen). Total

RNA isolated from both the groups was used for cDNA synthesis using oligo dT primers

and Superscript IV as per manufacturer's guidelines. The concentration of synthesized

cDNA was estimated using Qubit ssDNA assay kit and STX11 TaqMan probes were used

to perform the qPCR using LightCycler 96 (Roche). Beta-actin was used as housekeeping

control.

Single cell Ca^{2+} imaging

Cells were plated in carbon-coated glass bottom dishes (one day prior or 30 mins before

the assay for HEK293 and Jurkat T cells respectively) and loaded with 1μM Fura-2 AM

dye in HBSS (CaCl$_2$ 1.8mM, KCl 5.36mM, MgSO$_4$ 0.81mM, NaCl 136.89 mM, Na$_2$HPO4

0.335mM, D-Glucose 5.55mM) for 30 mins, at 37°C in the dark. After incubation, cells

were washed and incubated in Ringer's buffer (135mM NaCl, 5mM KCl, 1.8mM CaCl$_2$,

1mM MgCl$_2$, 5.6mM Glucose, 10mM HEPES) for an additional 10 mins, washed and

imaged in Ringer's buffer or Calcium-free Ringer's buffer (135mM NaCl, 5mM KCl, 1mM

MgCl$_2$, 5.6mM Glucose, 10mM HEPES at pH7.5), as indicated. Olympus IX-71 inverted

microscope equipped with a Lamda-LS illuminator (Sutter Instrument, Novato, CA), Fura-

2 (340/380) filter set (Chroma, Bellows Falls, VT), a 10X 0.3NA objective lens (Olympus,

UPLFLN, Japan), and a Photometrics Coolsnap HQ2 CCD camera was used to capture

images at a frequency of ~1 image pair every 2 or 4 seconds interval. Data were acquired,

analyzed and plotted using MetaFluor (Molecular Devices, Sunnyvale, CA), Microsoft

Excel, and Origin softwares. Approximately 30-40 cells were imaged per group in each

experiment unless otherwise stated. Mean SOCE ± SE for each experiment was

calculated from three independent experiments where the maximum value of the

scramble shRNA group (amongst the three experiments) was set at 100% and the relative

responses of other scr and STX11 shRNA groups were normalized w.r.t it.

For constitutive calcium influx assay, shRNA treated HEK293 cells were either

transfected or nucleofected with Orai1-S-S-EGFP, Orai1(H134S)-YFP or Orai1(ANSGA)-

YFP and ~12-14 hour post transfection (~4-5 hour post nucleofection), cells were loaded

with Fura-2 AM, washed and imaged in Ringer's buffer with 0 mM calcium to acquire

baselines and 2 mM calcium thereafter. To measure CAD induced constitutive calcium

influx, Orai1-CFP expressing stable HEK293 cells were nucleofected with YFP-CAD and

analyzed ~4 hour later as described above. To identify cells expressing Orai1 mutants

and fusion proteins in the plasma membrane, cells were imaged using a 20X 0.7NA water

objective lens (Olympus, UApoN340, Japan). Images were acquired at a frequency of ~1

image pair every 10 seconds interval to avoid photobleaching and analyzed as described

above.

Measurement of NFAT nuclear localization by Western blot

Jurkat T cells were transduced with scramble or STX11 shRNA to knock-down STX11 as

described above. On day 4 post-transduction, cells were collected, spun down and

resuspended in plain RPMI media and rested for 1hr at 37°C. Following this, cells were

counted and divided into two equal groups. One group was resuspended in RPMI

(unstimulated) and the other in RPMI media containing 1 μM TG + 10 ng/ml PMA (Phorbol

Myristate Acetate) (stimulated) and incubated at 37°C for 30 mins. Following incubation,

cells were pelleted and the nuclear and cytosolic protein fractions were separated using

NE-PER kit according to the manufacturer's guidelines and subjected to SDS-PAGE and

western blot using the mouse anti-NFATc2 primary antibody (Santa Cruz) followed by

Donkey anti-mouse secondary antibody.

Estimation of nuclear translocation of NFAT by immunolabeling

Control and STX11 depleted Jurkat T cells were rested in plain RPMI for ~1 hour prior to

the assay. ~100,000 cells per group were plated on freshly carbon-coated coverslips for

40 minutes and stimulated for 1 hour at 37°C with 5 μg/ml anti-CD3 antibody, diluted in

plain RPMI. Following this, cells were washed and fixed using 4% PFA diluted in 1X PBS

for 20 minutes at RT, washed twice and incubated with 30 mM Glycine for 10 minutes at

RT. For permeabilization and blocking, cells were incubated with 0.1% Saponin, 3% BSA

diluted in 1X PBS for 1 hour at RT, washed and incubated with anti-NFAT primary

antibody (anti-NFAT1, CST) at 4°C, overnight. Following primary antibody application,

cells were washed with 1X PBS and incubated with anti-Rabbit AF647 secondary

antibody for 1 hour, washed and stained with DAPI for 5 minutes followed by additional

washes. Images were acquired in the FV3000 confocal microscope using a 100X

objective lens. DAPI was used to identify the nuclear area. Nuclear *versus* whole cell

(total) NFAT mean intensity ratio was plotted across different groups.

Whole cell lysates (WCLs), Western blot and Co-immunoprecipitation (Co-IP)

HEK293 cells transfected with the desired plasmids were lysed using buffer containing

50 mM Tris-Cl (pH 8.0), 150 mM NaCl, 1% NP-40, 1 mM PMSF, and protease inhibitor

cocktail. The whole cell lysates were centrifuged at 21000 g for 15 minutes and

supernatants were subjected to SDS-PAGE, and western blot. For immunoprecipitations,

lysates were divided into two equal parts. To the first tube, the appropriate anti-mouse or

anti-rabbit primary antibody was added and to the second, same amount of the respective

IgG control antibody was added. The antibody-lysate mixtures were incubated overnight

at 4°C. Following this, Protein A/G Mag Sepharose beads were added and incubated with

the antibody-lysate mixtures for 4 hours, washed with the lysis buffer containing 0.1% NP-

40, boiled with 1X Laemmli buffer + 120 mM DTT and subjected to SDS-PAGE and

western blot analysis. Typically, $1/10_{th}$ of the whole cell lysate (WCL) was loaded in the

input lane of the co-IP blots. The full-length blots pertaining to each figure have been

deposited in Zenodo https://doi.org/10.5281/zenodo.19045452.

E. coli **expression and** *in-vitro* **binding assays**

Full-length His_6-tagged Stx11 and truncation mutants were cloned in pET28b, expressed

in Lemo21 (DE3) *E. coli* cells and induced with 1 mM IPTG (Isopropyl β- d-1-

thiogalactopyranoside) for 14-18 hours at 18°C. The cell pellets were lysed in buffer

containing 50 mM Tris-Cl (pH 8.0), 150 mM NaCl, 10% Glycerol, 1 mM PMSF, 1%

Sarkosyl, 0.1 mg/ml Lysozyme, protease inhibitor cocktail and sonicated on ice. DNase I

was added after sonication and the lysates were further incubated for ~60 minutes before

centrifugation at 21000 g for 40 mins. The supernatants were subjected to SDS-PAGE to

confirm expression by Coomassie staining and subsequently used for pull-down assays.

MBP-tagged Orai1 constructs were expressed in Lemo21 (DE3) *E. coli* cells, induced with

0.3mM IPTG (Isopropyl β- d-1-thiogalactopyranoside) for 14-18 hours at 18°C. The cell

pellets were lysed in buffer containing 50 mM Tris-Cl (pH 8.0), 150 mM NaCl, 5% Glycerol,

1 mM PMSF, 0.1 mg/mL Lysozyme and protease inhibitor cocktail (1:200) and sonicated

on ice. DNase I was added after sonication and the lysates were further incubated for ~60

minutes before centrifugation at 21000 g for 40 mins. The supernatants were collected

and subjected to SDS-PAGE to confirm expression by Coomassie staining and used for

pull-down assays. Following lysis, the His$_6$-tagged Stx11 proteins were purified using

Talon Beads and MBP-tagged Orai1 proteins using Dextrin Sepharose/Amylose resin

according to the manufacturer's guidelines. For *in-vitro* binding assays, lysates prepared

from cells expressing MBP or MBP-tagged Orai1 fragments were incubated with 25 μL

Dextrin Sepharose Beads and incubated for 2 hours at 4°C. After incubation, beads were

washed thrice and ~50-125 ng of purified His-tagged full length Stx11, His-tagged SNARE

or His-tagged H$_{abc}$ domains were diluted in buffer containing 50mM Tris-Cl (pH8), 150

mM NaCl, 5% Glycerol and 0.1% NP-40 and added to the beads. Following 1 hour of

incubation at 4°C, the beads were washed thrice, re-suspended and boiled in the binding

buffer containing 1X Laemmli and 120 mM DTT. The eluate was subjected to SDS-PAGE

and Western blot. MBP and MBP-tagged proteins were detected with Ponceau staining,

Full length STX11 using Rabbit anti-STX11 primary antibody (Thermo) followed by

Donkey anti-rabbit HRP and His-tagged STX11 fragments were detected or mouse anti-

6XHis primary antibody (Invitrogen) followed by Donkey anti-mouse HRP secondary

antibody.

BS3 Crosslinking of Flag-Orai Oligomers:

Scramble and STX11 shRNA transduced HEK 293 cells were transfected with Flag-Orai1,

harvested, washed, pelleted and subjected to one cycle of snap freeze-thaw. The pellet

was resuspended in ice cold lysis buffer (1X HBSS, 1mM PMSF and protease inhibitor

cocktail), sonicated for 30 secs (Branson 2510) and mechanically homogenized using

dounce homogenizer. The homogenized supernatant was spun at 1500 x g for 10 mins

at 4° C, to pellet the cell debris. The supernatant was ultracentrifuged at 105,000 x g for

1 hour at 4° C (Optima MAX-XP ultracentrifuge, Beckman Coulter). The membrane

fraction pellet was resuspended in lysis buffer and re-homogenized using dounce

homogenizer. The protein concentration was estimated using BCA assay and BS3 cross-

linker was added to the resuspended membranes, at a final concentration of 1mM, and

incubated for 30 mins at room temperature. BS3 was quenched using 50mM Tris-HCl pH

7.5 for 15 mins at room temperature, followed by addition of 1X Laemmli and 100 mM

DTT and incubation at 37° C for 10 mins. The samples were resolved using 4-15%

gradient SDS-PAGE and subjected to Western blot using anti-Flag antibody.

Quantification of Flag-Orai oligomers using Western blot:

Equal size ROIs were drawn over corresponding protein bands of same size, using

ImageJ. The total intensity of each band was measured by calculating the area under the

intensity curve and its relative proportion was calculated by dividing the raw intensity of

each band by the sum of intensities of all the bands in the respective lanes.

qPCR to estimate gene expression in Jurkat T cells

Scramble (scr) and STX11 shRNA treated Jurkat T cells were rested for ~1 hour prior to

the assay, stimulated with soluble anti-CD3 (5-10 μg/ml) for 3 hours and washed with

HBSS. Total RNA was isolated using RNeasy Mini kit as per manufacturer's instructions.

cDNA was synthesized using random hexamers and Invitrogen Superscript IV kit and

quantified using Qubit ssDNA assay. Taqman probes for IL-2 and beta-actin

housekeeping control were used for performing the qPCR in triplicates. Data analysis was

done by calculating the double delta C_t values.

EIA to measure IL-2 levels in supernatants of Jurkat T cells

Scramble (scr) and STX11 shRNA treated Jurkat T cells were stimulated with plate-

coated anti-CD3 (10 μg/ml) and the culture supernatants were collected at 12 and 24 hour

post-stimulation. EIA plates were coated with purified mouse anti-human IL-2 capture

antibody (1 μg/ml) overnight at 4°C, washed with 0.05% PBST and blocked with 2% BSA

for 2 hours. After blocking, the plates were incubated with the culture supernatants for 2

hours at RT, washed and incubated with the detection antibody (Biotin mouse anti-human

IL-2, 1 μg/ml) for 2 hours. After washing, Streptavidin HRP (0.1 μg/ml) followed by TMB

reagent were added. 250 mM HCl was added to stop the reaction and absorbance was

measured at 450 nm.

Isolation and culture of human PBMCs

Whole blood freshly collected in Heparin or Citrate Phosphate Dextrose Adenine (CPDA)

solution was subjected to density gradient centrifugation using Ficoll-Paque PLUS media.

The buffy coat was separated, washed twice with HBSS and cultured with IL-2 (50 ng/ml).

Cells were stimulated once per week with PHA (2 μg/ml) for 72 hours and rested in IL-2

for the remaining 72-96 hours. Unless specified, all the assays were performed following

24-48 hours of rest, post stimulation.

Isolation of genomic DNA and sequencing

Genomic DNA was extracted from PBMCs using Phenol-Chloroform-Isoamyl alcohol.

STX11 genomic region flanking the mutation was PCR amplified using primers: 5'

Forward   -CATGCACGACTACAACCAGGC   and   3'   Reverse   -

GGGACAGCAGAAGCAGCAGAGGG. The resulting PCR products were separated on

2% agarose gel, excised and extracted using the Macherey-Nagel Nucleospin columns

and subjected to Sanger sequencing using the 5' Forward PCR primer to confirm the

mutation.

Measurement of SOCE in human PBMCs

PBMCs in culture were washed, rested and allowed to adhere to freshly carbon-coated

(IBDI) dishes for 1 hour in plain RPMI at 37°C. The cells were washed with HBSS and

incubated with 1 ml of 1 μM Fura-2-AM diluted in Ringer's buffer for 40 minutes, washed

and incubated for an additional 10 minutes to allow de-esterification of the dye. The assay

was started with 1 ml of Calcium free Ringer's buffer in the imaging dish and images were

captured every 4 seconds. After capturing the baseline for ~60 seconds, stores were

depleted by the addition of 1 μM Thapsigargin. In other assays, 10 μg/ml of anti-CD3 and

5 μg/ml of the secondary antibody were used to cross-link the TCRs and thereby induce

store depletion. ~5 minutes post-store-depletion, Calcium Chloride ($CaCl_2$) was added

back to the dish at a final concentration of 2 mM to estimate the magnitude of store

operated calcium entry.

Degranulation assay

PBMCs were cultured in RPMI containing 10% FBS and 50 ng/ml IL-2 and stimulated

with PHA (2 μg /ml) 48hrs before the assay. 24hrs prior to the assay, IL-2 was washed

off but PHA was re-added. On the day of the assay cells were washed twice to remove

PHA and any growth factors. To measure degranulation, cells were either left

unstimulated, stimulated with a combination of anti-CD3 (10 μg/ml plate-coated), anti-

CD28 (2 μg/ml soluble) and anti-CD49d (2 μg/ml soluble) or a combination of Ionomycin

(1 μM) and PMA (50 ng/ml) for 3.5 hours. CD107a-PE antibody (1:50 dilution) was added

to each of the three groups at the start of stimulation. Following stimulations, cells were

transferred to ice, washed with cold HBSS containing 2% FBS and incubated with anti-

CD8 APC for 20 minutes, washed twice with cold HBSS and analyzed using Cytoflex

(Beckman Coulter) flow cytometer.

qPCR to estimate gene expression in human PBMCs

PBMCs were taken off IL-2 48 hours prior to the assay. On the day of the assay, cells

were washed twice, left unstimulated and either stimulated with anti-CD3 (10 μg/ml, plate-

coated), anti-CD28 (2 μg/ml) and anti-CD49d (2 μg/ml) or with Ionomycin (1 μM) and PMA

(50 ng/ml) for 6 hours. To end the stimulation, cells were washed with cold HBSS, pelleted

and used for RNA extraction. Total RNA was isolated using Qiagen RNeasy mini kit and

cDNA was synthesized using random hexamers and Superscript IV (Invitrogen). cDNA

was quantified using Qubit and subjected to QPCR analysis using Taqman probes for IL-

2 and beta-actin housekeeping gene control in triplicates. Analysis was performed by

calculating the double delta C_t values.

Transduction of human PBMCs with pMSCV-STX11

For viral transduction of PBMCs, polystyrene non-TC treated 24-well plates were coated

with retronection (20 μg/ml) overnight at 4°C, blocked with 2% BSA for 30 minutes at

room temperature and washed twice with HBSS. The retroviral supernatants were added

onto the coated wells and the plates were spun at 1800g for 2 hours at 30°C. Following

spin, the wells were washed with the blocking solution. PBMCs cultured in RPMI

containing 10% FBS were stimulated with PHA (2 μg/ml) and IL-2 (50 ng/ml) for 48 hours

prior to transduction. Stimulated PBMCs were transferred to coated plates at a density of

0.25 million cells/well centrifuged at 400g for 40 minutes at 30°C. The cells were analyzed

48-72 hours post transduction.

Patch clamp measurements

Patch clamp recordings were performed by using an Axopatch 200B amplifier (Axon

Instruments, Foster City, CA) interfaced to an ITC-18 input/output board (Instrutech, Port

Washington, NY) and an iMac G5 computer. Currents were filtered at 1 kHz with a four-

pole Bessel filter and sampled at 5 kHz. Recording electrodes were pulled from 100-μl

pipettes (VWR), coated with Sylgard, and fire-polished to a final resistance of 2 to 5 MΩ.

Stimulation and data acquisition and analysis were performed by using in-house routines

developed by RS Lewis (Stanford University) on the Igor Pro platform (Wavemetrics, Lake

Oswego, OR). All data were corrected for the liquid junction potential of the pipette

solution relative to Ringer's in the bath (-10 mV) and for leak currents collected in 20 mM

extracellular free calcium plus 10μM La^{3+}. The holding potential was +30 mV. The

standard voltage stimulus consisted of a 100-ms step to −100 mV followed by a 100-ms

ramp from −100 to +100 mV applied at 1 s intervals. I_{CRAC} was typically activated by

passive depletion of ER Ca^{2+} stores by intracellular dialysis of 8 mM BAPTA.

Solutions and chemicals

The standard extracellular Ringer solution contained 130 mM NaCl, 4.5 mM KCl, 20 mM

CaCl$_2$, 10 mM tetraethylammoniumchloride (TEA-Cl), 10 mM D -glucose and 5 mM

HEPES (pH 7.4 with NaOH). The standard divalent-free (DVF) Ringer solutions contained

(in mM) 150 NaCl, 10 HEDTA, 1 EDTA, and 10 HEPES (pH 7.4). The internal solution

contained: 135 mM Cs aspartate, 8 mM MgCl$_2$, 8 mM Cs-BAPTA and 10 mM HEPES (pH

7.2 with CsOH).

Data analysis

Analysis of current amplitudes was typically performed by measuring the peak currents

during the -100 mV pulse. Averaged results are presented as the mean value standard

error of the mean (SEM). Reversal potentials were measured from the average of several

leak-subtracted sweeps (4–6) in each cell. For datasets with two groups, statistical

analysis was performed with two- tailed t test to compare between control and test

conditions with a confidence level of 95%, and results with $p<0.05$ were considered

statistically significant.

STX11:Orai1 and SOAR:Orai complex prediction

Full length STX11 and its two domains, (H$_{abc}$ 41-167 and SNARE 183-267) and Orai1 (N-

terminus 1-87 and C-terminus 256-301) were used to generate the complex of STX11

and Orai1 using AlphaFold3 (AF3) in all combinations. For Stim1, SOAR, residue 344-

444 were used along with Orai C-terminus 256-301. The resultant models were assessed

using ipTM and pTM scores. The best scoring combinations were further considered to

generate more models by changing the seed values. The best model in terms of highest

ipTM and pTM values was considered for further analysis. Custom script was used to

analyze the contact frequency of interacting residues of STX11-H$_{abc}$ and Orai1 C-terminus

complex as well as SOAR and Orai1 C-terminus, across all predicted models.

Molecular dynamics (MD) simulation

The STX11 H$_{abc}$:Orai1 C-terminus and SOAR:Orai1 C-terminus complexes were

subjected to all-atom MD simulations. Initially, the complexes were prepared using

protein-preparation module of Schrodinger which involves H-bond network optimization

and restrained minimization of the initial structure. The prepared structures were solvated

using TIP3P water model in an orthorhombic box and neutralized by adding counter ions.

OPLS4 force field was used, and simulation system was generated by specifying 150 mM

salt (NaCl). The solvated system was subjected to default relaxation protocol of Desmond

followed by production run for 500 ns at 300K and 1 atm pressure in NPT ensemble. The

default relaxation protocol includes several short simulation steps. (1) Brownian dynamics

simulation for 100 ps at 10 K temperature in NVT with restraints on solute heavy atoms

(2) Simulation in NVT ensemble at 10 K for 12 ps with restraints on solute heavy atoms

(3) 12 ps simulation in NPT ensemble at 10 K with restraints on solute heavy atoms (4)

Simulation in NPT ensemble for 12 ps with restraints on solute heavy atoms (5) Simulation

in NPT ensemble for 24 ps without restraints. Three independent runs were executed with

different initial seeds for both structures.

Analysis of MD simulation

The MD runs were analyzed for the stability of the complex and interactions between the

subunits. The RMSD was calculated using simulation interaction diagram (SID) module

of Schrodinger. The first frame was used as reference frame for RMSD calculation. Script

"analyze_trajectory_ppi.py" was used to calculate the interactions between STX11:Orai1

and Stim1:Orai1. The binding energy (ΔG) at each nanosecond across simulation was

calculated using "thermal_mmgbsa.py" script of Schrodinger. The representative

structure was generated through trajectory clustering. The last 200 ns frames from each

run were combined and clustered using scripts "trj_merge.py" and "trajectory cluster" of

Schrodinger. The cluster representative of the largest cluster was used to analyze the

protein-protein interactions. PDB Sum was used to calculate the interactions in the cluster

representative. Figures were generated using Pymol and MD movies using Maestro.

Statistical Analysis

Statistical analysis was done using unpaired two-tailed Student's *t* test where only two

groups were being compared and ANOVA where more than two groups were present.

Normality of distribution was checked in Origin. Mann Whitney U Test was used for

datasets with non-normal (or skewed) distribution. P values for all significance tests were

*P<0.05; **P<0.01; ***P<0.001.

1000

1001

Acknowledgements: We thank Vivien Beziat, Yenan Bryceson, Heinrich Schlums, Jelve

Nejati Zendegani, Stephan Ehl, and Jasmin Mann for help with human PBMC handling,

isolation, storage and shipping. Anand Vaidya for advice on STX11 purification. Jyoti

Rohilla for technical support. Bio Render for the generation of models and IBS 2.0 for the

generation of schematics shown in this manuscript. This work was supported in part by

Department of Atomic Energy, Government of India under Project Identification

No. RTI 4007 and NIH-NIAID grant AI108636.

1009

1010

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022

1023

1024

1025

1026

1027

1028

References:

1. Zhang, K., et al., *Familial Hemophagocytic Lymphohistiocytosis*, in *GeneReviews((R))*, M.P. Adam, et al., Editors. 1993: Seattle (WA).
2. D'Orlando, O., et al., *Syntaxin 11 is required for NK and CD8(+) T-cell cytotoxicity and neutrophil degranulation.* Eur J Immunol, 2013. **43**(1): p. 194-208.
3. Clapham, D.E., *Calcium signaling.* Cell, 2007. **131**(6): p. 1047-58.
4. Vig, M. and J.P. Kinet, *The long and arduous road to CRAC.* Cell Calcium, 2007. **42**(2): p. 157-62.
5. Vig, M., et al., *Defective mast cell effector functions in mice lacking the CRACM1 pore subunit of store-operated calcium release-activated calcium channels.* Nat Immunol, 2008. **9**(1): p. 89-96.
6. Maul-Pavicic, A., et al., *ORAI1-mediated calcium influx is required for human cytotoxic lymphocyte degranulation and target cell lysis.* Proc Natl Acad Sci U S A, 2011. **108**(8): p. 3324-9.
7. Vig, M., et al., *CRACM1 multimers form the ion-selective pore of the CRAC channel.* Curr Biol, 2006. **16**(20): p. 2073-9.
8. Vig, M., et al., *CRACM1 is a plasma membrane protein essential for store-operated Ca2+ entry.* Science, 2006. **312**(5777): p. 1220-3.
9. Miao, Y., et al., *An essential and NSF independent role for alpha-SNAP in store-operated calcium entry.* Elife, 2013. **2**: p. e00802.
10. Prakriya, M. and R.S. Lewis, *Store-Operated Calcium Channels.* Physiol Rev, 2015. **95**(4): p. 1383-436.
11. Feske, S., et al., *A mutation in Orai1 causes immune deficiency by abrogating CRAC channel function.* Nature, 2006. **441**(7090): p. 179-85.
12. Zhang, S.L., et al., *Genome-wide RNAi screen of Ca(2+) influx identifies genes that regulate Ca(2+) release-activated Ca(2+) channel activity.* Proc Natl Acad Sci U S A, 2006. **103**(24): p. 9357-62.
13. Yoon, T.Y. and M. Munson, *SNARE complex assembly and disassembly.* Curr Biol, 2018. **28**(8): p. R397-R401.
14. Li, P., et al., *alpha-SNAP regulates dynamic, on-site assembly and calcium selectivity of Orai1 channels.* Mol Biol Cell, 2016. **27**(16): p. 2542-53.
15. Miao, Y., et al., *Na+ influx via Orai1 inhibits intracellular ATP induced mTORC2 signaling to disrupt CD4 T cell gene expression and differentiation.* Elife, 2017. **6**.
16. Ramanagoudr-Bhojappa, R., Y. Miao, and M. Vig, *High affinity associations with alpha-SNAP enable calcium entry via Orai1 channels.* PLoS One, 2021. **16**(10): p. e0258670.
17. Jarvis, S.E. and G.W. Zamponi, *Masters or slaves? Vesicle release machinery and the regulation of presynaptic calcium channels.* Cell Calcium, 2005. **37**(5): p. 483-8.
18. Vig, M. and J.P. Kinet, *Calcium signaling in immune cells.* Nature immunology, 2009. **10**(1): p. 21-7.
19. Bryceson, Y.T., et al., *Defective cytotoxic lymphocyte degranulation in syntaxin-11 deficient familial hemophagocytic lymphohistiocytosis 4 (FHL4) patients.* Blood, 2007. **110**(6): p. 1906-15.

1072   20.   Hackmann, Y., et al., *Syntaxin binding mechanism and disease-causing mutations in*
1073   *Munc18-2.* Proc Natl Acad Sci U S A, 2013. **110**(47): p. E4482-91.
1074   21.   Hellewell, A.L., et al., *Analysis of familial hemophagocytic lymphohistiocytosis type 4*
1075   *(FHL-4) mutant proteins reveals that S-acylation is required for the function of syntaxin*
1076   *11 in natural killer cells.* PLoS One, 2014. **9**(6): p. e98900.
1077   22.   Jahn, R. and R.H. Scheller, *SNAREs--engines for membrane fusion.* Nat Rev Mol Cell Biol,
1078   2006. **7**(9): p. 631-43.
1079   23.   Li, Z., et al., *Graded activation of CRAC channel by binding of different numbers of STIM1*
1080   *to Orai1 subunits.* Cell Res, 2011. **21**(2): p. 305-15.
1081   24.   McNally, B.A., et al., *Gated regulation of CRAC channel ion selectivity by STIM1.* Nature,
1082   2012. **482**(7384): p. 241-5.
1083   25.   Peinelt, C., et al., *Amplification of CRAC current by STIM1 and CRACM1 (Orai1).* Nat Cell
1084   Biol, 2006. **8**(7): p. 771-3.
1085   26.   Mercer, J.C., et al., *Large store-operated calcium selective currents due to co-expression*
1086   *of Orai1 or Orai2 with the intracellular calcium sensor, Stim1.* J Biol Chem, 2006.
1087   **281**(34): p. 24979-90.
1088   27.   Soboloff, J., et al., *Orai1 and STIM reconstitute store-operated calcium channel function.*
1089   J Biol Chem, 2006. **281**(30): p. 20661-20665.
1090   28.   Hou, X., S.R. Burstein, and S.B. Long, *Structures reveal opening of the store-operated*
1091   *calcium channel Orai.* Elife, 2018. **7**.
1092   29.   Zhou, Y., et al., *The STIM1-binding site nexus remotely controls Orai1 channel gating.*
1093   Nat Commun, 2016. **7**: p. 13725.
1094   30.   Cormet-Boyaka, E., et al., *CFTR chloride channels are regulated by a SNAP-23/syntaxin*
1095   *1A complex.* Proc Natl Acad Sci U S A, 2002. **99**(19): p. 12477-82.
1096   31.   Toft-Bertelsen, T.L., et al., *Regulation of Ca2+ channels by SNAP-25 via recruitment of*
1097   *syntaxin-1 from plasma membrane clusters.* Mol Biol Cell, 2016. **27**(21): p. 3329-3341.
1098   32.   Jessen, B., et al., *Graded defects in cytotoxicity determine severity of hemophagocytic*
1099   *lymphohistiocytosis in humans and mice.* Front Immunol, 2013. **4**: p. 448.
1100   33.   Chinn, I.K., et al., *Genetic and mechanistic diversity in pediatric hemophagocytic*
1101   *lymphohistiocytosis.* Blood, 2018. **132**(1): p. 89-100.
1102   34.   Dulubova, I., et al., *Munc18-1 binds directly to the neuronal SNARE complex.* Proc Natl
1103   Acad Sci U S A, 2007. **104**(8): p. 2697-702.
1104   35.   Lee, S., et al., *Munc18-1 induces conformational changes of syntaxin-1 in multiple*
1105   *intermediates for SNARE assembly.* Sci Rep, 2020. **10**(1): p. 11623.
1106   36.   Park, C.Y., et al., *STIM1 clusters and activates CRAC channels via direct binding of a*
1107   *cytosolic domain to Orai1.* Cell, 2009. **136**(5): p. 876-90.
1108   37.   Stathopulos, P.B., et al., *STIM1/Orai1 coiled-coil interplay in the regulation of store-*
1109   *operated calcium entry.* Nat Commun, 2013. **4**: p. 2963.
1110   38.   Delevoye, C., et al., *Conservation of the biochemical properties of IncA from Chlamydia*
1111   *trachomatis and Chlamydia caviae: oligomerization of IncA mediates interaction*
1112   *between facing membranes.* J Biol Chem, 2004. **279**(45): p. 46896-906.
1113   39.   Paumet, F., et al., *Intracellular bacteria encode inhibitory SNARE-like proteins.* PLoS One,
1114   2009. **4**(10): p. e7375.

1115   40.   Barocchi, M.A., V. Masignani, and R. Rappuoli, *Opinion: Cell entry machines: a common*
1116   *theme in nature?* Nat Rev Microbiol, 2005. **3**(4): p. 349-58.
1117   41.   Neveu, E., et al., *Prototypic SNARE Proteins Are Encoded in the Genomes of*
1118   *Heimdallarchaeota, Potentially Bridging the Gap between the Prokaryotes and*
1119   *Eukaryotes.* Curr Biol, 2020. **30**(13): p. 2468-2480 e5.

1120

1121

1122

1123

1124

1125

1126

1127

1128

1129

1130

1131

1132

1133

1134

1135

1136

1137

1138

1139

1140

1141

1142

1143

1144

1145

1146

1147

1148 **Figure Legends:**

1149

Figure 1: STX11 is required for SOCE. Measurement of SOCE in STX11 depleted

Jurkat and FHLH4 patient T cells. (**A-B**) Representative Fura-2 calcium imaging assay

(**A**) and quantification of independent repeats (**B**) measuring thapsigargin (TG) induced

SOCE in Jurkat T cells treated with scr (black), STX11#1 (red) or STX11#2 (blue)

shRNAs. (n=30-40 cells/group, N=3). (**C-D**) Representative Fura-2 calcium imaging

assay (**C**) and quantification of repeats (**D**) showing reconstitution of SOCE in STX11

depleted Jurkat T cells by ectopic expression of STX11. scr shRNA with empty vector

(EV) expression (black), STX11 shRNA with EV expression (red), STX11 shRNA with

STX11 expression (green). (**E**) Sanger sequencing of FHLH4 patient DNA showing

deletion of a single Adenine at the 752^{nd} position and the resulting frameshift. (**F**)

Schematic showing the predicted domain distribution of the wildtype *versus* FHLH4

mutant STX11 protein. (**G**) Western blot on WCLs of healthy donor (HD) and FHLH4

patient PBMCs showing the relative molecular weight and abundance of the wildtype and

mutant STX11 bands. (**H**) Western blot on WCLs of HEK293 cells over-expressing

wildtype or FHLH4 mutant STX11 and empty vector. (**I**) Fura-2 calcium imaging assay

measuring anti-CD3 induced SOCE in HD (black) and FHLH4 (red) T cells. (n=30-40

cells/group). (**J**) Fura-2 calcium imaging assay measuring SOCE in HD expressing empty

vector (EV) (black) and FHLH4 T cells expressing human STX11 (red), respectively.

(n=30-40 cells/group). (**K-L**) Representative Fura-2 calcium imaging assay (**K**) and

quantification of repeats (**L**) measuring SOCE in Jurkat T cells treated either with scr

shRNA expressing empty vector (EV) (black), STX11 shRNA, expressing EV (red),

STX11 shRNA, expressing wildtype STX11 (green), STX11 shRNA, expressing FHLH4

mutant STX11 (blue).

Figure 2. STX11 regulates I$_{CRAC}$. (**A-D**) Measurement of I$_{CRAC}$ in STX11 depleted Jurkat

T-cells in the whole-cell recording configuration in 20 mM extracellular Ca^{2+} Ringer's

solution. I$_{CRAC}$ was induced by passive depletion of intracellular Ca^{2+} stores by dialyzing

8 mM BAPTA into the cell via the patch-pipette. (**A**) Representative current at -100 mV

in Jurkat T cells treated with scr shRNA construct. The current is blocked by extracellular

La^{3+} (10 µM) and replacing the 20 mM Ca^{2+} Ringer's solution with a divalent free solution

(DVF) evokes a large Na^+ current which depotentiates over tens of seconds. The current-

voltage (I-V) relationship of the Ca^{2+} and DVF currents are shown on the

right. (**B**) I_{CRAC} from a Jurkat T cell treated with STX11 shRNA. Both Ca^{2+} and Na^+ current

amplitudes are reduced relative to control cells. The I-V relationships (*right plots)* show

no change in ion selectivity. (**C-D**) Summary of the current amplitudes of Ca^{2+} and

Na^+ currents and current reversal potentials in scr and STX11 knockdown cells.

Figure 3. STX11 regulates NFAT activation, IL-2 gene expression and degranulation

in FHLH4 patient T cells. (**A-C**) Estimation of nuclear translocation of NFAT. (**A**) Western

blot showing nuclear translocation of NFAT in Jurkat T cells treated with scr or STX11

shRNA and stimulated with PMA+TG. N=3. (**B**) Representative confocal images of control

and STX11 depleted Jurkat T cells stimulated with anti-CD3, immunolabelled with anti-

NFAT1 antibody and counter-stained with DAPI. (N=3) (**C**) Box and whisker plot showing

quantification of nuclear NFAT from cells populating 10 randomly chosen fields per group

in (**B**). (**D**) Quantification of IL-2 mRNA in anti-CD3 stimulated Jurkat T cells treated with

scr or STX11 shRNA using qPCR. The bars show mean $\pm$ SE of relative IL-2 mRNA. N=3.

(E) Quantification of IL-2 EIA on supernatants of scr (black) or STX11 (red) shRNA treated

Jurkat T cells stimulated with anti-CD3. (**F**) qPCR analysis of IL-2 mRNA expression in

various stimulated HD (black) and FHLH4 T cells (red). (**G-H**) Granule release assay

performed on HD (black) and FHLH4 CD8 T (red) cells (**G**) and its quantification (**H**).

1198

Figure 4: STX11 directly binds resting Orai1 in plasma membrane. Intracellular

localization of STX11 and its association with Orai1. (**A**) Representative confocal images

of HEK293 cells transfected with either HA-tagged STX11 or untagged STX11 and

stained using anti-HA antibody or anti-STX11 antibody respectively. N=3 (**B**) Localization

of HA-tagged STX11 with respect to Orai1-YFP in the basal as well as nuclear plane of

HEK293 cells. N=3 (**C-D**) Co-IP to assess STX11 binding to Orai1. Whole cell lysates of

resting and store-depleted HEK293 cells co-expressing either Flag-Orai1 and STX11 (**C**)

or Orai1-Myc and STX11 (**D**) were subjected to IP and Western blot using anti-Myc, anti-

Flag or anti-STX11 antibodies, as indicated. (N=3) (**E**) Schematic showing key domains

of STX11 and Orai1 used for *in vitro* pull-down assays. (**F**) Pull-down assay showing *in*

vitro binding of His-tagged STX11 to Orai1. (Top) Ponceau S staining showing the input

of MBP alone or MBP-tagged Orai1 fragments. (Bottom) Western blot using anti-STX11

antibody. Input:1/5th of the protein. N=4. (**G**) Pull-down assay showing *in vitro* binding of

His-tagged H_{abc} domain of STX11 to Orai1 fragments. (Top) Ponceau S staining showing

the input of MBP alone or MBP-tagged Orai1 fragments. Input:1/5th of the protein.

(Bottom) Western blot using anti-His antibody. N=3. (**H**) Binding free energy distribution

of STX11-H_{abc} and Orai1 C-terminus interactions. (**I**) Sphere and cartoon representation

of the structure of STX11 H_{abc} (green) and Orai1 C-terminus (cyan) complex after MD

simulation. The N-termini are highlighted in blue, and C-termini in red. (**J**) Representative

confocal images showing localization of STX11 with respect to Orai1:Stim1 puncta in

store-depleted cells. HEK293 cells expressing Orai1-YFP, CFP-Stim1 were transfected

with HA-tagged STX11, store-depleted and stained using anti-HA antibody. (N=3) (**K**)

Quantification of the localization of STX11 with respect to Stim1:Orai1 puncta in store-

depleted cells.

1223

Figure 5. Residues involved in Orai1:STX11 complex formation are crucial for

SOCE. Analysis of STX11 and Orai1 mutants. (**A**) Schematic showing STX11 mutations

(Top). Expression of wildtype STX11 or STX11 N147A_E150A mutant in HEK293

followed by immunolabeling (**A**, bottom) and Western blot (**B**) using anti-STX11 antibody.

(**C-D**) Representative Fura-2 calcium imaging assay (**C**) and quantification of repeats (**D**)

measuring rescue of SOCE in Jurkat cells treated either with scr shRNA (black) or STX11

shRNA expressing empty vector (red), wildtype STX11 (green) or N147A_E150A mutant

STX11 (blue). (**E**) Coomassie blue stained SDS-PAGE showing expression and

purification of His-tagged wildtype or N147A_E150A mutant STX11 from *E. coli.* (**F**) Pull-

down assay measuring their binding to MBP-tagged Orai1 C-terminus. (Top) Ponceau S

staining showing the input of MBP-tagged fragments. (Bottom) Western blot using anti-

STX11 antibody. N=4 (**G**) Schematic showing Orai1 mutations (Top). Confocal images of

HEK293 cells expressing YFP-tagged wildtype or R289A_E272A_E275A_E278A mutant

Orai1 (Bottom). **(H-I)** Representative Fura-2 calcium imaging assay (**H**) and quantification

of repeats (**I**) measuring rescue of SOCE in HEK293 cells treated either with scr (black)

or Orai1 shRNA expressing empty vector (red), wildtype Orai1 (green) or

R289A_E272A_E275A_E278A mutant Orai1 (blue). (**J**) Pull-down assay measuring the

binding of MBP-tagged wildtype or R289A_E272A_E275A_E278A mutant Orai1 C-

terminus to wildtype or N147A_E150A mutant STX11. (Top) Ponceau S staining showing

the input of MBP-tagged Orai1 fragments. (Bottom) Western blot using anti-STX11

antibody. N=4. (**K-L**) Representative confocal images (**K**) and quantification (**L**) of PM

localized YFP-CAD in CFP-tagged wildtype or R289A_E272A_E275A_E278A mutant

Orai1 expressing HEK293 cells.

1247

Figure 6: Orai1 clustering is dysfunctional in STX11 depleted cells. Quantification of

Orai1 and Stim1 intensities in the ER-PM puncta of store-depleted cells. (**A-B**)

Representative confocal images of resting (**A**) and store-depleted (**B**) shRNA treated

HEK293 cells expressing N-terminal CFP tagged Orai1 (CFP-Orai1) and C-terminal YFP

tagged Stim1 (Stim1-YFP). (**C-F**) Mean Stim1-YFP intensity (**C**) and area (**D**) of Stim1

clusters. Mean CFP-Orai1 intensities inside (**E**) and outside (**F**) Stim1-YFP clusters of

shRNA treated cells, post store-depletion. (**G-J**) Quantification of Orai1 and Stim1

intensities in the ER-PM puncta of C-terminal YFP-tagged Orai1 (Orai1-YFP) and N-

terminal CFP-tagged Stim1 (CFP-Stim1). (**G-H**) Mean CFP-Stim1 intensity (**G**) and area

(**H**) of Stim1 clusters. Mean Orai1-YFP intensity (**I**) and area (**J**) of Orai1-YFP clusters in

control and STX11 depleted cells. (**K-L**) Representative Fura-2 calcium imaging assay

(**K**) and quantification of repeats (**L**) measuring TG induced SOCE in Orai1-YFP and

CFP-Stim1 expressing HEK293 cells treated with scr (black) or STX11 (red) shRNA.

1261

Figure 7: STX11 primes Orai1 for optimal on-site assembly. Oligomerization of Orai1.

(A) N-FRET analysis of scramble control and STX11 shRNA treated HEK293 cells co-

expressing Orai1-CFP and Orai1-YFP before and after store-depletion. **(B&C)**

Representative images (**B**) and quantification (**C**) of FRET analysis using acceptor

photobleaching of control and STX11 shRNA treated Orai1 Stim1 double knockdown

resting HEK293A cells co-expressing Orai1-CFP (donor) and Orai1-YFP (acceptor). **(D-**

E) Representative western blot (**D**) and quantification of band intensities from repeats (**E**)

showing distribution of BS3 crosslinked Flag-Orai1 oligomeric bands in control and

STX11 depleted HEK293 cells. N=3. (**F-H**) Fura-2 calcium imaging assay (**F**) and

quantification of repeats (**G-H**) to measure constitutive calcium influx in Orai1-CFP and

YFP-CAD expressing, control (black) or STX11 depleted (red) HEK293 cells at 0 mM (**G**)

and 2 mM (**H**) extracellular Ca^{2+}. (**I-J**) Fura-2 calcium imaging assay (**I**) and quantification

of repeats (**J**) to measure constitutive calcium influx in Orai1-H134S mutant expressing

control (black) and STX11 depleted (red) HEK293 cells. (**K-L**) Fura-2 calcium imaging

assay (**K**) and quantification of repeats (**L**) to measure constitutive calcium influx in Orai1-

ANSGA mutant expressing control (black) and STX11 depleted (red) HEK293 cells.

1278

1279 **Figure 8: Models depicting STX11 mediated priming of Orai1 and its role in T cell**

1280 **signaling and function.** (**A**) STX11 unbound (unprimed) and bound (primed) Orai1 in

1281 resting and store-depleted cells. For simplicity, STX11 interaction with only two subunits

1282 of Orai1 has been shown. (**B**) Regulation of T cell SOCE, NFAT activation and effector

1283 function by STX11.

1284

1285

1286 Due to BioRxiv file size limit, the Supplementary Movies can be found at the following

1287 link: https://www.dropbox.com/scl/fo/3j07d3sua4g58w3yfmyv7/AAKjvIiniA-

1288 z994hxV9DNHo?rlkey=e7yj5wyukvwg5seg1awkuctc4&st=z6vp57dr&dl=0

1289

1290 **Supplementary Movies 1-3: Molecular dynamic simulations of Orai1 C-terminus**

1291 **interaction with the STX11_H_{ABC} domain.**

1292 **Supplementary Movies 4-6: Molecular dynamic simulations of Orai1 C-terminus**

1293 **interaction with Stim1 SOAR domain.**

1294

1295

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

A **STX11 sufficient:**

STX11 deficient:

B

Supplementary Figure 1

Supplementary Figure 1. (A-H) Targeted RNAi screen for SNAREs involved in SOCE. Fura-2 calcium imaging assays measuring thapsigargin (TG) induced SOCE in HEK293 and/or Jurkat cell lines treated with scramble (scr) shRNA or different shRNA sequences against the gene of interest for 3-4 days. Shown here are the average traces from 30-40 cells per group.

Supplementary Figure 2

Supplementary figure 2 (A-B) Representative Fura-2 calcium imaging assay **(A)** and its quantification from repeats **(B)** measuring thapsigargin (TG) induced SOCE in HEK293 cells treated with scr or STX11 shRNA. Bars show relative SOCE ± SE from three independent experiments in **(A)** where SOCE from scramble shRNA treated group in one experiment was set at 100%. **(C)** Representative quantitative PCR to assess the efficiency of knockdown in shRNA treated cells from **(B)**. Total RNA extracted from cells treated with STX11 shRNA was subjected to qPCR analysis using Taqman probes for STX11. Data were normalized to Beta-actin housekeeping control.

Supplementary Figure 3

Supplementary Figure 3. (A) Schematic showing the design of recombinant Orai1-BBS-YFP fusion protein. Shown here is the placement of the bungarotoxin binding site (BBS) inside the second extracellular loop and YFP tag in the C-terminus of Orai1 as well as binding of BTX-A647 to BBS. **(B)** Quantification of total Orai1 levels in the plasma membrane of STX11 depleted cells. U2OS cells stably expressing Orai1-BBS-YFP were transduced with scr (black) or STX11 (red) shRNA, stimulated with 1uM TG, incubated with alpha-bungarotoxin alexa 647 (BTX-A647) and washed. BTX binding to surface Orai1 was measured using flow-cytometer, where its binding to wildtype U2OS cells were used as control. (N=3). **(C-D)** Measurement of thapsigargin (TG) induced SOCE in U2OS cells treated with STX11 RNAi in **(B)**. **(C)** Representative average single cell Fura-2 calcium imaging assay. **(D)** Bars showing relative mean % SOCE ± SE from three independent experiments in **(C)**. **(E)** Measurement of ER calcium content in Scr and STX11 shRNA treated HEK293 cells using ER-localized CEPIA. Cells were incubated in Ringer's buffer containing 0 mM calcium followed by 1 mM EGTA and stimulated with 1 uM ionomycin. (N=3).**(F-G)** Immunolabelling of HEK293 cells to assess ER **(F)** and Golgi **(G)** health. Scr and STX11 shRNA treated HEK293 cells were fixed, permeabilized and stained with anti-KDEL antibody **(F)** and anti-GM130 antibody **(G)** to assess ER and Golgi morphology respectively.

Supplementary Figure 4

Supplementary Figure 4. (A) Schematic showing a comparison of the key domains in STX11 and STX1A. **(B)** Western blot of whole cell lysates prepared from HEK293 cells expressing YFP-STX11 and STX11-YFP. Note the mismatch in molecular weight of the N- versus C-terminally tagged STX11 and the degradation products (arrows). N=3. **(C)** Confocal images showing localization of ectopically expressed mCherry-STX11 or STX-11-mCherry in HEK293 cells. N=3. **(D)** Pull-down assay showing *in vitro* binding of His-tagged SNARE domain of STX11 to MBP-tagged Orai1 N- and C-termini. (Top panel) Ponceau S staining showing the input of MBP alone or MBP-tagged Orai1 cytosolic tails. (Bottom panel) Western blot using anti-His antibody.

Supplementary Figure 5

Supplementary Figure 5: (A) Frequency of interface residues of STX11-H$_{abc}$ and Orai1 C-terminus across AF3 predicted models. Residues with more than 50% frequency are shown. **(B)** RMSD of STX11 H$_{abc}$ and Orai1 C-terminus upon superposition on the 0th ns frame. The STX11-H$_{abc}$ RMSD indicates the backbone RMSD upon superposition of all frames on the STX11-H$_{abc}$ backbone of 0th ns. The Orai1 C-terminus RMSD suggests the stability of Orai1 with respect to STX11 and is calculated for the Orai1 C-terminus after superposition of all frames on the STX11-H$_{abc}$ backbone of 0th ns.

Supplementary Figure 6

Supplementary Figure 6: (A) Binding energy (ΔG) of STX11-H$_{abc}$ and Orai1 C-terminus complex across the simulation time for all three replicates. **(B)** Interactions between STX11-H$_{abc}$ and Orai1 C-terminus across the simulation time in three replicates. The interactions that persist for more than 30% of simulation time are shown and highlighted in different colors. H-bond has been categorized as sidechain-sidechain (ss), sidechain-backbone (sb) and backbone-sidechain (bs). **(C)** Residues predicted to be involved in interaction between STX11 H$_{abc}$ and the Orai1 C-terminus identified from all-atom MD simulations.

Supplementary Figure 7

Supplementary Figure 7. (A) Co-immunoprecipitation (co-IP) to test the association of Stim1 with STX11. HEK293 cells were co-transfected with Stim1-Myc-His and STX11, store-depleted, lysed and subjected to co-IP followed by Western blot as indicated. (N=2) **(B)** Quantification of fraction of Orai1 inside and outside Stim1:Orai1 clusters in store-depleted HEK293 cells expressing CFP-Orai1 and Stim1-YFP.

Supplementary Figure 8

Supplementary Figure 8: (A) Schematic of human Stim1 and Orai1 showing the domains used for MD simulation. TM:Transmembrane, P/S: Proline/Serine rich and CC: represent Coiled coil domains. **(B)** The binding free energy distribution of the three runs for the interactions between the SOAR dimer and Orai1 C termini. **(C)** Binding energy (Δ G) of SOAR dimer and Orai1 C-termini complex across the simulation time for all three runs.

Supplementary Figure 9

A

Supplementary Figure 9. (A) RMSD of SOAR dimer and Orai1 C-termini. **(B)** Cartoon representation of the structure of Stim1-SOAR(344-444) dimer in complex with Orai1 C-termini.

Supplementary Figure 10

Supplementary Figure 10. (A-B) Orai1-YFP and CFP-Stim1 clusters in control and STX11 depleted cells. Representative images of Scr and STX11 shRNA treated resting **(A)** and store-depleted **(B)** HEK293 cells expressing C-terminal YFP-tagged Orai1 (Orai1-YFP) and N-terminal CFP-tagged Stim1 (CFP-Stim1). Scale bar 10µm.

Supplementary Figure 11

Supplementary Figure 11. (A) Western blot of BS3 cross-linked Flag-Orai1 acquired at a higher exposure showing the monomeric Flag-Orai1 band (labelled as A) **(B)** Schematic representation of SOAR domain in full length Stim1. **(C)** Cartoon showing the design of the Orai1-SOAR-SOAR-EGFP (Orai1-S-S-GFP) construct. **(D)** Representative Fura-2 calcium imaging assay to measure constitutive calcium influx in Orai1-S-S-GFP expressing control (black) or STX11 depleted (red) HEK293 cells. Cells were imaged in Ringer's buffer containing 0 mM followed by 2 mM extracellular Ca^{2+}. **(E)** Quantification of constitutive calcium influx across experiments as shown in **(D)**. N=3. **(F-G)** Schematic representations of H134S mutation **(F)** and ANSGA mutation **(G)** in full length Orai1.

Supplementary Figure 12

Supplementary Figure 12. SOCE in RNAi treated Jurkat cells to assess the role of SNAPs. **(A-C)** Fura-2 calcium imaging assay measuring SOCE in cells treated with scramble (scr) shRNA or different shRNA sequences against SNAP23 **(A)** SNAP25 **(B)** SNAP29 **(C)**. Average traces from 30-40 cells per group are shown. **(D)** Western blot images of co-IPs to test the association of Orai1 with SNAP23/ SNAP25/ SNAP29. HEK293 cells were co-transfected with Orai1-Myc and untagged SNAP23/ SNAP25/ SNAP29, store-depleted, lysed and subjected to co-IP followed by western blot as indicated. (N=2) **(E)** Western blot images of co-IP to test the association of Stim1 with SNAP23/ SNAP25/ SNAP29. HEK293 cells were co-transfected with YFP-Stim1 and Myc-tagged SNAP23/ SNAP25/ SNAP29, store-depleted, lysed and subjected to co-IP followed by western blot as indicated. (N=2).

Supplementary figure 13

Supplementary Figure 13. RNAi screen for genes involved in primary HLH and/or vesicle fusion for their role in SOCE. **(A-F)** Fura-2 calcium imaging assays measuring thapsigargin (TG) induced SOCE in HEK293 or Jurkat cells treated with scramble (scr) shRNA or different shRNA sequences against the gene of interest for 3-4 days. Shown here are the average traces from 30-40 cells per group.

Supplementary Table 1: AF3 prediction for STX11:Orai1 complex formation

Proteins/Domains	ipTM	pTM
Full-length Orai1 - Full-length STX11	0.11	0.35
STX11-Habc_41-167 and Orai1_N-term_1-87	0.27	0.59
STX11-Habc_41-167 and Orai1_C-term_256-301	**0.53**	**0.79**
STX11-SNARE_183-267 and Orai1_N-term_1-87	0.08	0.34
STX11-SNARE_183-267 and Orai1_C-term_256-301	0.18	0.45
STX11-Habc_41-167 and full-length Orai1	0.24	0.38
STX11-SNARE_183-267 and full-length Orai1	0.08	0.37
Orai1_N-term_1-87 and full-length STX11	0.15	0.5
Orai1_C-term_256-301 and full-length STX11	0.16	0.54

Supplementary Table 2: AF3 predictions for STX11-H_{abc} and Orai1 C-terminus with different seeds

Seed values for STX11-H$_{abc}$ and Orai1 C-terminus	ipTM	pTM
111	**0.69**	**0.85**
236	0.51	0.78
174	0.5	0.77
802	0.49	0.76
455	0.56	0.78

Supplementary Table 3: AF3 predictions for SOAR and Orai1 in 2:2 ratio

2 copies of SOAR and 2 copies of Orai1_C-term_256-301	ipTM	pTM
Seed_101	0.73	0.78
Seed_102	0.73	0.78
Seed_103	0.73	0.78
Seed_104	0.73	0.78
Seed_105	**0.74**	**0.79**