Three-dimensional memory of nuclear organization through cell cycles

Shin Fujishiro[*1] and Masaki Sasai[**1,2]

[1)]*Fukui Institute for Fundamental Chemistry, Kyoto University, Kyoto 606-8103, Japan*
[2)]*Department of Complex Systems Science, Nagoya University, Nagoya 464-8601, Japan*

(*sinfu@nagoya-u.jp)

(**masakisasai@nagoya-u.jp)

(Dated: 25 October 2024)

The genome in the cell nucleus is organized by a dynamic process influenced by structural memory from mitosis. In this study, we develop a model of human genome dynamics through cell cycles by extending the previously developed whole-genome model to cover the mitotic phase. With this extension, we focus on the role of mitotic and cell cycle memory in genome organization. The simulation progresses from mitosis to interphase and the subsequent mitosis, leading to successive cell cycles. During mitosis, our model describes microtubule dynamics, showing how forces orchestrate the assembly of chromosomes into a rosette ring structure at metaphase. The model explains how the positioning of chromosomes depends on their size in metaphase. The memory of the metaphase configuration persists through mitosis and into interphase in dimensions perpendicular to the cell division axis, effectively guiding the distribution of chromosome territories over multiple cell cycles. At the onset of each G1 phase, phase separation of active and inactive chromatin domains occurs, leading to A/B compartmentalization. Our cycling simulations show that the compartments are unaffected by structural memory from previous cycles and are consistently established in each cell cycle. The genome model developed in this study highlights the interplay between chromosome dynamics and structural memory across cell cycles, providing a new perspective for the analyses of cellular processes.

I. INTRODUCTION

In human cells, chains of chromosomes are confined within a nucleus with a radius of typically $5\,\mu$m in the G1 phase. The nucleus is far from being in a well-stirred equilibrium as the microscope observation showed that each chromosome moves less than a few μm during interphase in the human cells.[1] Although chromatin exhibits rapid, liquid-like movements on a scale of $\lesssim 100\,$nm,[2–4] it shows slow, glass-like movements on a μm scale.[4,5] Therefore, an intriguing and important issue is the nonequilibrium memory effects in the slow process of chromatin structural organization.

An example suggesting the memory effect is how chromosomes form territories in mammalian nuclei.[6,7] In mammalian cells, chromatin domains, or topologically associating domains (TADs), with a sequence length of 10^5–$10^6\,$bp[8] and a diameter of 200–400 nm,[4,9] are considered as the structural and functional units of chromosomes.[10] One way to represent a chromosome is as a chain connecting these domains. For instance, if a chromosome of length $10^8\,$bp exhibited a Gaussian-like behavior, its radial distribution would extend to $R_{\text{ideal}} \sim (10^8/10^5)^{1/2} \times 200\,$nm $\approx 6\,\mu$m, reaching the nuclear radius. If multiple chains, which tend to spread over the entire nucleus, were in thermal equilibrium within cells, the nucleus would resemble a polymer melt,[11] with multiple distributions of chains significantly overlapping. However, observations with fluorescence microscopy have shown that chromosome chains in mammalian cells are not mixed but instead form individual territories.[12–14] One possible explanation for the territory formation is that the interphase chromosomes retain the memory of the mitotic phase, in which individual chromosomes are separated. When entering the G1 phase, condensed chromosomes in the mitotic phase expand due to rapid, liquid-

FIG. 1. Cells go through a cycle of events as they proliferate. An important question is how the genome structure's memory is retained or erased through cell cycles.

like movements during decondensation, but the large-scale movements of chromosomes are slower than the expansion. Therefore, the system reaches a nonequilibrium stationary G1 state before the chains are mixed with each other.

The hypothesis that the memory of the mitotic phase influences the interphase genome architecture is also supported by observations on other organisms such as fungi or mosquitoes.[15] In these organisms, the formation of chromosome territories is less obvious, and instead, telomeres or centromeres are clustered, indicating genome architectures with Rabl-like features. Moreover, in these organisms, the subunits of condensin II, a factor determining chromosome structure during mitosis, are not expressed,[15] suggesting that the difference in the mitotic configuration determines the difference in the genome architecture in interphase.

Fujishiro and Sasai analyzed the memory effects in the genome architecture by developing a polymer model of the

human genome.[6,16] In this model, chromosomes in the mitotic phase were represented as chains of beads, with each bead representing a 10-Mb chromatin region. The simulation was started from a random metaphase (RMP) configuration based on the assumption that the memory of the configuration in metaphase would be erased by the large motion of chromosomes during anaphase (Fig. 1). Starting from the RMP configuration, the anaphase process was simulated by moving the kinetochore of each chromosome toward a spindle pole. After the chromosomes were assembled near the spindle pole, the nuclear envelope was generated to wrap the genome in telophase. Upon exiting telophase, the chromosome model was fine-grained by replacing the 10-Mb resolution chains with chains connecting 100-kb chromatin regions. Using this 100-kb resolution model, the decondensation of chromosomes was simulated at the entry to the G1 phase. In the G1 phase, the nucleus reached a steady state, with simulated chromosomes forming territories. The simulated size of each territory quantitatively reproduced the experimental observation,[17] and the simulated radial position of each territory within the nucleus explained the experimental data.[18] The simulation also revealed a memory persistence within each chromosome. The dragging force toward the spindle pole during anaphase caused chromosomes to form V shapes, with the memory of this configuration persisting as a relatively large frequency of contacts between the p- and q-arms of chromosomes, which were necessary to explain the observed Hi-C contact map in interphase.[8] In this way, the memory persisting through the decondensation from the mitotic phase to the interphase explains the territory formation and the shape of each chromosome in the territory.

In the previous version of the Fujishiro and Sasai model,[6,16] the RMP approximation was used to represent the starting configuration of the simulation in the mitotic phase. However, it is uncertain whether the metaphase memory is erased during anaphase, so a more precise structure should replace this approximation to examine the memory effect during mitosis. Here, to address this issue, we develop a model to simulate the entire cell cycle of human cells (Fig. 1). The simulation proceeds in the following steps: (i) We generate a prophase genome configuration at a 10-Mb resolution using the simulated interphase genome configuration at a 100-kb resolution. (ii) During the prometaphase, we assume microtubule fibers connect a pair of spindle poles and daughter chromosomes. The daughter chromosomes are moved through the microtubule dynamics to form a rosette distribution,[12,19] and aligned on the spindle equator through chromosome congression, giving a metaphase configuration. (iii) Then, we focus on a half spindle to simulate the movement of a sister genome toward one of the spindle poles during anaphase. (iv) Similarly to the previous version of the model, the configuration of assembled chromosomes is relaxed in telophase, and (v) chromosomes are fine-grained to a 100-kb resolution to simulate decondensation toward the interphase to complete the cell cycle.

Mitosis is a highly dynamic process that attracted significant interest from both experimental and theoretical studies.[20–25] There have been extensive discussions on various aspects of the spindle formation and dynamics, including the elongation of microtubules to reach the chromosome kinetochores,[26–28] the determination of spindle shape and size,[29] and the regulation of chromosome oscillations within a spindle.[30–33] The present study focuses on how chromosomes are moved and configured during mitosis,[12,19,34,35] rather than the molecular basis of spindle structure and dynamics. For this purpose, we adopt a phenomenological model of microtubule dynamics. With this model, we analyze how chromosomes self-organize to form configurations during mitosis and examine how genome structural memory is retained or erased during a single cell cycle and through multiple cell cycles. It is expected that the model developed in this study will help address questions on fluctuations in the position of each chromosome within the nucleus, and how errors or perturbations during the mitotic phase affect chromosome structure and dynamics during interphase.

In the cell cycle simulation, it is reasonable to assume repulsive interactions between chromatin regions to explain the chromosome expansion at the beginning of the G1 phase. In addition, as the genome enters the G1 phase, transcription becomes active, and we need to distinguish between regions with high and low gene activity. We classify 100-kb chromatin regions into type-A (gene-active), type-B (gene-inactive or gene-poor), and type-u (intermediate) regions. Because domains in type-B regions are more tightly compacted than those in type-A regions,[36,37] there should be stronger repulsive interactions between type-B regions than between type-A regions. In the Fujishiro and Sasai model,[6,16] the heterogeneous repulsive interactions between 100-kb chromatin regions induce phase separation of chromatin, causing type-A regions to assemble into compartment A and type-B regions into compartment B, and spontaneously forming the lamina associated domains (LADs)[38,39] and the nucleolus associated domains (NADs).[40] This model explains the Hi-C and other biochemical data with high precision without using the global Hi-C data as an input to the simulation.[6] As compartments A and B spread across boundaries of chromosome territories,[8] it is crucial to analyze whether the compartments are enhanced or fluctuate through multiple cell cycles.

Thus, in this study, we develop a model of the human genome to examine its essential structural characteristics along the cell cycle. With this model, we analyze the effects of the persistence of structural memory through iterated cycles of human cells.

II. 3D MODEL OF MITOSIS

We first describe a model of the three-dimensional (3D) genome conformation during mitosis. By integrating it with a previously developed model of the interphase genome,[6,16] we can simulate the 3D genome dynamics throughout the entire cell cycle. In our model, we represent chromosomes as chains of volume-excluding beads. During the mitotic phase, the observed Hi-C contact maps showed that chromosomes consist of condensed regions of approximately 10 Mb, and at scales above this, the contact frequencies showed similar patterns to

those of linear chains.[41,42] Therefore, we use each bead to represent a 10-Mb segment in chromosomes during mitosis.

A. 3D dynamics of mitotic chromosomes

The motion of each 10-Mb bead was simulated by numerically integrating the overdamped Langevin equation,

$$\frac{d\mathbf{r}_i}{dt} = -\mu \frac{\partial U}{\partial \mathbf{r}_i} + \sqrt{2\mu k_{\mathrm{B}} T}\, \boldsymbol{\xi}_i, \tag{1}$$

where $\mathbf{r}_i$ is the position of the ith bead with $1 \leq i \leq 2N$, and N is the number of beads in each daughter genome. μ is the mobility of a bead, U is the force-field potential, k_{B} is the Boltzmann constant, and T is the effective temperature in the simulation. $\boldsymbol{\xi}_i$ is a Gaussian white noise vector, satisfying $\langle \xi_{i,\alpha}(t)\xi_{j,\beta}(s)\rangle = \delta_{i,j}\delta_{\alpha,\beta}\delta(t-s)$ with $1 \leq i, j \leq 2N$ and $\alpha, \beta = x, y, z$, where $\delta(\cdot)$ is the Dirac delta and $\delta_{.,.}$ is the Kronecker delta. The effective temperature T should reflect the movements in degrees of freedom that are not considered in the present coarse-grained model. These movements may vary depending on the amount of chemical reactions produced in the system. Changing in reactions for the motor protein activity or transcriptional activation may increase the effective temperature. Therefore, by writing a unit temperature as T^*, we used $T = T^*$ in prometaphase and $T = 4T^*$ in anaphase and telophase. As explained later in this paper, increasing the effective temperature in anaphase and telophase was necessary to explain how the Rabl-like feature of the clustered centromeres formed in anaphase is lost in telophase. Since the mobility μ is uniform for all degrees of freedom in Eq. 1, choosing the value of μ is equivalent to determining the unit of time. Here, for the simulation convenience, we chose μ as $\mu = 1\,\mu\mathrm{m}^2/(\mathrm{hr} \cdot k_{\mathrm{B}} T^*) \approx 0.016\,\mu\mathrm{m}^2/(\mathrm{min} \cdot k_{\mathrm{B}} T^*)$.[6] The Langevin equation Eq. 1 was discretized with a time step of $\delta t = 10^{-4}\,\mathrm{min}$ in simulations.

The force-field potential U consists of the linear-chain component U_{chain}, the sister chromatid cohesion component U_{sister}, the packing potential U_{pack}, the polar ejection force (PEF) component U_{pe}, and the kinetochore-microtubule force (KMF) component U_{km}:

$$U = U_{\mathrm{chain}} + U_{\mathrm{sister}} + U_{\mathrm{pack}} + U_{\mathrm{pe}} + U_{\mathrm{km}}. \tag{2}$$

In Eq. 2, $U_{\mathrm{sister}} = 0$ in anaphase and telophase, and $U_{\mathrm{pack}} \neq 0$ only in telophase. We explain U_{pack} later in Eqs. 20 and 21 in section III, and U_{pe} and U_{km} in subsections, *Polar ejection force* and *Kinetochore-microtubule force*, of the current section II.

The linear-chain component, U_{chain}, is composed of volume-excluding repulsion, bond spring, and bending-cost potentials:

$$U_{\mathrm{chain}} = \sum_{i=1}^{2N}\sum_{j=i+1}^{2N} u_{\mathrm{rep}}(\mathbf{r}_{i,j}) + \sum_{k=1}^{4n}\sum_{i=s_k}^{e_k-1} u_{\mathrm{bond}}(\mathbf{r}_{i,i+1})$$
$$+ \sum_{k=1}^{4n}\sum_{i=s_k+1}^{e_k-1} u_{\mathrm{bend}}(\mathbf{r}_{i-1,i}, \mathbf{r}_{i,i+1}), \tag{3}$$

where the vector $\mathbf{r}_{i,j} = \mathbf{r}_i - \mathbf{r}_j$ is the displacement vector between the ith and jth beads, k is the index to label chromosomes, and $n = 23$. The kth chromosome consists of beads numbered as $s_k \leq i \leq e_k$. The functional forms of potentials in Eq. 3 are:

$$u_{\mathrm{rep}}(\mathbf{r}) = \begin{cases} \varepsilon_{\mathrm{rep}}\left(1 - \left(\frac{r}{\sigma}\right)^2\right)^3 & \text{for } r \leq \sigma, \\ 0 & \text{for } r > \sigma, \end{cases} \tag{4}$$

$$u_{\mathrm{bond}}(\mathbf{r}) = \begin{cases} 0 & \text{for } r \leq \sigma, \\ \frac{1}{2}K(r-\sigma)^2 & \text{for } r > \sigma, \end{cases} \tag{5}$$

$$u_{\mathrm{bend}}(\mathbf{r}, \mathbf{s}) = \varepsilon_{\mathrm{bend}}\frac{\mathbf{r} \cdot \mathbf{s}}{rs}. \tag{6}$$

We set the values of repulsive energy to prevent the chains from crossing each other as $\varepsilon_{\mathrm{rep}} = 100k_{\mathrm{B}}T^*$, where $k_{\mathrm{B}}T^*$ is a unit energy in simulations. The bead size was set to $\sigma = 0.3\,\mu\mathrm{m}$ to make the length of mitotic chromosomes approximately $3\,\mu\mathrm{m}$, and the bending energy and the spring constant were set to allow chromosomes for taking a rod-like shape as $\varepsilon_{\mathrm{bend}} = 1k_{\mathrm{B}}T^*$ and $K = 10000k_{\mathrm{B}}T^*/\mu\mathrm{m}^2$.

The association of sister chromatids in prometaphase and metaphase is maintained through additional bonds:

$$U_{\mathrm{sister}} = \sum_{(k,l)\in\mathsf{Sis}} u_{\mathrm{bond}}(r_{k,l}), \tag{7}$$

where Sis is a set of pairs of kinetochore beads on sister chromatids. Here, a kinetochore bead is the chromosomal bead that covers the centromere of the chromosome. We assume that cohesin, which bundles sister chromatids, disappears after metaphase and set $U_{\mathrm{sister}} = 0$ in anaphase.

In addition to these bead-spring forces described by U_{chain} and U_{sister} in Eq. 2, mitotic chromosomes are subject to the forces in the spindle (Fig. 2a). In the following subsections, we explain these forces, PEF and KMF, and the corresponding potential components, U_{pe} and U_{km}.

B. Polar ejection force

Polar ejection force (PEF) is the force exerted by the action of chromokinesin residing on the arms of mitotic chromosomes. An ambient microtubule binds a chromokinesin molecule, whose motor activity drives the chromosome towards the plus-end of the microtubule, or the direction away from the spindle pole, generating PEF on the chromosome.[43,44] When we regard a spindle pole as a point source from which microtubules grow, the density of microtubules around a chromosome should decay with the inverse-square law. Hence, when a chromosome is distance R away from a spindle pole, the binding rate of a chromokinesin molecule to a microtubule is proportional to $1/R^2$. In this microtubule field model, the binding probability p_{ck} follows the master equation,

$$\frac{dp_{\mathrm{ck}}}{dt} = \frac{\Sigma}{R^2}a_{\mathrm{ck}}(1-p_{\mathrm{ck}}) - k_{\mathrm{ck}}p_{\mathrm{ck}}, \tag{8}$$

FIG. 2. (a) Forces acting on mitotic chromosomes. Microtubules (MTs) grow from a spindle pole (SP), attach to the kinetochore (black band) on sister chromatids (Ch), and generate kinetochore-microtubule force (KMF) on a chromosome. Other non-attached MTs exert polar ejection force (PEF) on chromosomal arms through the binding of chromokinesin molecules; the reactive cross-section is shown as a surface Σ. (b) Potentials for fields of KMF (red dashed), PEF (blue dashed), and their summation (black), represented as functions of distance R from a spindle pole. (c) Two-dimensional potential surface for the field of the summed force of KMF and PEF generated by a pair of spindle poles. The inset is a magnification, showing that the potential has double minima due to the combination of the KMF acting inward and the PEF acting outward.

where a_{ck} is the binding rate, k_{ck} is the unbinding rate, and Σ is the reactive cross-section for chromokinesin molecules on a chromosomal segment and microtubules. Assuming that the binding and unbinding are much faster than chromosomal dynamics in prometaphase and metaphase, the stationary solution,

$$p_{ck} = \frac{1}{1+(R/b)^2}, \quad b^2 = \frac{a_{ck}\Sigma}{k_{ck}}, \qquad (9)$$

can be used to describe the force field. Here, b^2 is the effective cross-section. If chromokinesin molecules bound on a chromosomal segment generate force f_{pe} when binding to a microtubule, the PEF, F_{pe}, acting on the chromosomal segment is

$$F_{pe} = p_{ck}f_{pe} = \frac{f_{pe}}{1+(R/b)^2}. \qquad (10)$$

The potential for this force is

$$u_{pe}(R) = -\int F_{pe}\,dR = u_{0,pe} - f_{pe}b\arctan\frac{R}{b}. \qquad (11)$$

Setting the constant of integration $u_{0,pe} = f_{pe}b\pi/2$, and noting that $\arctan(x) = \pi/2 - \arctan(1/x)$, we obtain the PEF potential,

$$u_{pe}(R) = f_{pe}b\arctan\frac{b}{R}. \qquad (12)$$

In the prometaphase simulation, all beads on chromosomal chains are subject to the PEF potential from each spindle pole:

$$U_{pe} = \sum_{i=1}^{4N}\sum_{s\in SP} u_{pe}(R_{i,s}). \qquad (13)$$

Here, $SP = \{1,2\}$ is the set of spindle poles, and $R_{i,s} = |\mathbf{r}_i - \mathbf{c}_s|$ is the distance between the ith bead and the sth spindle pole. $\mathbf{c}_s$ is the position vector of the sth spindle pole.

We assume that PEF acts during prometaphase to metaphase and diminishes in anaphase. In Eq. 12, we set

$f_{pe} = 200k_BT^*/\mu m \sim 1\,pN$ based on the experimental measurements during prometaphase,[45] and $f_{pe} = 0$ in anaphase. We used $b^2 = 0.8\,\mu m^2$, so that the chromosomes show reasonable shapes and the arrangement on the metaphase plate.

C. Kinetochore-microtubule force

A bundle of microtubules, known as k-fiber, binds to the kinetochore of a chromosome through the Ndc80 complex.[46] The polymerization and depolymerization of microtubules result in pushing and pulling of the kinetochore from/to a spindle pole, respectively. The depolymerization rate of a microtubule is proportional to its length.[47–49] Therefore, if the rates of elongation and shortening are a_m and k_m, respectively, the length of a microtubule bundle, ℓ, follows the equation,

$$\frac{d\ell}{dt} = a_m - k_m\ell. \qquad (14)$$

Although the altering polymerization and depolymerization oscillate the length of microtubules,[50] we average out the oscillations in our model to more focus on the global motion of chromosomes. Because a chromosome is bound to the tip of a k-fiber, its position x along the k-fiber is $x = \ell$; hence, the motion of the chromosome obeys

$$\frac{dx}{dt} = \frac{d\ell}{dt} = a_m - k_mx = -k_m\left(x - \frac{a_m}{k_m}\right). \qquad (15)$$

This can be written as an overdamped motion under an effective spring potential,

$$\frac{dx}{dt} = -\mu_c u'_{km}(x), \quad u_{km}(x) = \frac{1}{2}\frac{k_m}{\mu_c}(x-\ell^*)^2, \qquad (16)$$

where ℓ^* is the stationary length of the k-fiber, $\ell^* = a_m/k_m$, and μ_c is the effective mobility of a chromosome. When a chromosome is represented as a linear chain of L beads of mobility μ, the effective mobility should approximately be $\mu_c = \mu/L$. Therefore, we define the kinetochore-microtubule

force (KMF) potential for the kinetochore of a chromosome of length L as

$$u_{\text{km}}(R;L) = \frac{1}{2}K_L(R-\ell^*)^2, \quad K_L = \frac{k_{\text{m}}L}{\mu}. \quad (17)$$

We should note that $u_{\text{km}}(R;L)$ is not an elastic potential under the passive thermal fluctuation but represents the active microtubule dynamics in Eq. 14, which led to the dependence of $u_{\text{km}}(R;L)$ on L in Eq. 17 through the constraint of $x = \ell$ in Eq. 15.

Then, we have

$$U_{\text{km}} = \sum_{k=1}^{4n} u_{\text{km}}(R_{i(k),s(k)}), \quad (18)$$

where $i(k)$ is the index of the kinetochore bead of the kth chromosome and $s(k)$ is the spindle pole associated with the kth chromosome. Again, $R_{i,s} = |\mathbf{r}_i - \mathbf{c}_s|$ is the distance between the ith bead and the sth spindle pole at position $\mathbf{c}_s$. By construction, the kth and the $k+2n$th chromosomes constitute sister chromatids. We neglect the cases of erroneous connections and define $s(k) = 1$ for $k \leq 2n$ and $s(k) = 2$ for $k > 2n$ to ensure the proper association of sister chromatids with a pair of spindle poles.

In simulations, we used different values of ℓ^* in Eq. 17 in different mitotic phases to represent the transition from metaphase to anaphase. However, for simplicity, we used a constant value of the microtubule depolymerization rate as $k_{\text{m}} = 0.8\,\text{min}^{-1}$ from prometaphase through anaphase. We chose this value, so that the poleward motion in the early anaphase finishes within 5 min.

The combined potential of forces acting on a chromosome of L beads can be approximated by considering a virtual particle, which is the distance R away from the spindle pole. This particle is subjected to the force, whose potential is $u_{\text{spindle}}(R;L) = Lu_{\text{pe}}(R) + u_{\text{km}}(R;L) = Lu_{\text{spindle}}(R;1)$. In Fig. 2b, we show $u_{\text{spindle}}(R;1)$ as a function of R. $u_{\text{spindle}}(R;1)$ has a minimum at $R \approx \ell^*$, showing the combination of PEF and KMF yields a stable state. Fig. 2c is a two-dimensional plot of $u_{\text{spindle}}(R_1;1) + u_{\text{spindle}}(R_2;1)$, where R_1 and R_2 are distances between the particle and spindle poles 1 and 2, respectively. $u_{\text{spindle}}(R_1;1) + u_{\text{spindle}}(R_2;1)$ shows the tendency of the particle to reside at the equator between two spindle poles. We should note that u_{spindle} has two minima of about $2\,\mu\text{m}$ apart on the relatively flat potential surface along the equator due to the balance between the KMF acting inward and the PEF acting outward. The flat potential on the equator prevents chromatids from collapsing toward the spindle center and instead distributes them in a rosette pattern on the metaphase plate. In this way, the potential $U_{\text{pe}} + U_{\text{km}}$ in Eq. 2 should generate the stable metaphase configuration of sister chromatids. Then, with the depletion of PEF and the loss of cohesion as $U_{\text{pe}} = U_{\text{sister}} = 0$ in anaphase, sister genomes are separated and dragged toward two poles. In the next section, we show how this scenario was realized in simulations.

III. SIMULATION OF THE ENTIRE CELL CYCLE

To represent the high-resolution chromatin organization during interphase, such as the organization of compartments, we assumed each bead in the interphase represents a 100-kb chromatin region. The 100-kb resolution chains in the G1 phase were coarse-grained and duplicated to generate the 10-Mb resolution chains at prophase. Simulation of the prometaphase process was then performed using the 10-Mb resolution chains to generate the metaphase genome. After the metaphase, the dynamics of one of the two sister genomes were followed in anaphase and telophase. When entering the G1 phase, the 10-Mb resolution chains were fine-grained to the 100-kb resolution chains. See Appendix A and Appendix B for more details on the procedures of coarse-graining and fine-graining.

As the starting configuration at the 0th round of the cell cycle, the RMP configuration was used (See Appendix C). From this configuration, through the anaphase and telophase simulations, the G1-phase genome at the 1st cycle was generated. Then, the prophase configuration at the cth cycle was generated from the G1-phase configuration at the cth cycle, and the next $c+1$st G1-phase configuration was generated through the simulated mitosis. Although we used human lymphoblastoid cells (GM12878) as an example in this simulation, the developed simulation scheme is also applicable to other human cells.

A. Prophase

The prophase genome configuration was generated in the simulation as the initial configuration to start the prometaphase dynamics. Chromosomes are duplicated during the transition from the G1 phase to the S phase, and compacted from the G2 phase to prophase to form the prophase configuration. Although each chromosome shows detailed structural changes during these phases, these changes should be erased through the drastic movement in chromosome condensation and congression during mitosis. Therefore, we generated the prophase conformation directly from the G1-phase conformation by neglecting those detailed changes produced in the S and G2 phases and focusing on the chromosome territory distribution.

We first coarse-grained each 10Mb chromatin segment of the G1-phase genome into a single bead, as explained in Appendix A. Then, the resulting coarse-grained structures were duplicated in the neighborhood of each chain as

$$\mathbf{r}_{N+i} = \mathbf{r}_i + \mathbf{\Delta}_k, \quad s_k \leq i \leq e_k, \quad (19)$$

where s_k and e_k are the start and end numbers of beads in the kth chain, and $\mathbf{\Delta}_k$ is a vector of length $\sigma = 0.3\,\mu\text{m}$, whose direction was randomly chosen for each chain k. This duplication produces a coarse-grained genome $\{\mathbf{r}_i\}_{i=1}^{2N}$ that approximates the changes via S and G2 phases. Starting from the prophase configuration thus generated (Fig. 3a), we performed the prometaphase simulation.

FIG. 3. Snapshots of the 3D genome structure during a whole cell-cycle simulation: (a) Prometaphase onset, displaying $4n$ chromosomes. (b-c) Metaphase configuration at the end of prometaphase, shown from different angles. (d) Positional autocorrelation of sister kinetochores in the prometaphase simulation. (e) Anaphase starts with the $2n$ half of the metaphase structure and (f) allows chromosomes to be dragged towards a spindle pole, (g) resulting in a Rabl-like configuration. (h) Telophase, where chromosomes are relaxed in a spherical container. (i) Positional autocorrelation of centromeres in the anaphase simulation. (j) Two-dimensional projection of chromosomal beads with color-coded centromeres at the beginning of telophase, showing clustered centromeres. The x axis indicates the equatorial position and the y axis is taken along the spindle axis. (k) The centromere clustering is relaxed at the end of telophase. (l) Cross-section of the nucleus at the beginning of the G1 phase and (m) at the end of the G1 phase in an expanded nucleus.

B. Prometaphase to metaphase

During the simulation from prometaphase to metaphase, we placed a pair of spindle poles at $c_1 = (0\,\mu\text{m}, 6\,\mu\text{m}, 0\,\mu\text{m})$ and $c_2 = (0\,\mu\text{m}, -6\,\mu\text{m}, 0\,\mu\text{m})$. As the center of the nucleus in the parental cell was at $(0, 0, 0)$ and its radius was several μm, c_1 and c_2 are at symmetric positions just near the surface of the nucleus in the parental cell. The stationary length of a k-fiber was set to $\ell^* = 6\,\mu$m, so that the length of k-fibers matches the distance between a spindle pole and the metaphase plate.

The KMF drags kinetochores, allowing centromeres to gather toward the inner region near the spindle axis. Meanwhile, the PEF pushes the chromosomal arms outwards. The combined interactions of KMF and PEF show two minima on the two-dimensional surface, as illustrated in Fig. 2c. These minima form a ring of approximately $2\,\mu$m in diameter in 3D space. Consequently, the KMF and PEF work together to assemble chromosomes toward the ring, leading to "chromosome congression."

Larger chromosomes are subjected to the stronger KMF and PEF. Therefore, when we followed the chromosome behaviors in the two-dimensional directions on the plane perpendicular to the spindle axis, we found that larger chromosomes tend to arrive at the ring at an earlier time than smaller chromosomes. Smaller chromosomes gathered later to find a room in the inner region near the ring. This alignment created a "chromosome rosette" on the spindle-equator plane (the metaphase plate), as shown in Figs. 3b and 3c. This simulated process of chromosome positioning resembles the observed process in mouse oocyte cells,[35] in which smaller chromosomes gather in the inner region of the ring after the other chromosomes assemble around the ring. Since KMF acts only on kinetochores and PEF acts on whole chromosomes, kinetochores gather toward the center of the rosette ring, while chromosomal arms point outwards (Fig. 3c).

The simulation for prometaphse to metaphase was run for 3×10^5 steps, or 30 minutes, which was long enough for all chromosomes to align on a metaphase plate. Once aligned,

volume exclusion and congestion prevent the chromosomes from repositioning on the metaphase plate. As a result, the memory of the chromosomal positioning in the parental nucleus is partially retained on the metaphase plane. This partial retention of memory through chromosome congression is discussed quantitatively in the first subsection of the next section, *Memory retention and erasure during mitosis*.

C. Anaphase

In anaphase, sister chromatids dissociate, the PEF is turned off,[51] and $4n$ chromosomes are dragged towards one of two spindle poles, which segregates $4n$ chromosomes into two sets of $2n$ chromosomes. Therefore, in the anaphase simulation, we changed the total potential to $U = U_{chain} + U_{km}$ and halved the system by choosing one set of $2n$ chromosomes associated with the first spindle pole, i.e., choosing the beads $\{r_i\}_{i=1}^{N}$. The spindle pole was moved to $c_1 = (0\,\mu m, 7.5\,\mu m, 0\,\mu m)$, which is near the surface of the daughter nucleus to be formed in the telophase. The stationary k-fiber length for the U_{km} was set to $\ell^* = 1.5\,\mu m$, which allows the chromosomes to be dragged toward $6\,\mu m$ away from the previous metaphase plate. See Figs. 3e-3g for the snapshots during anaphase. In simulations, the quantities were calculated using the coordinates translated by $(0, \mu m, -6\,\mu m, 0\,\mu m)$ to place the system center around the coordinate origin. The effective temperature representing the noise strength was raised to $T = 4T^*$.

After U_{sister} and the PEF were turned off (Fig. 3e), centromeres arrived at the region about ℓ^* apart from the spindle pole in several minutes (Fig. 3f). Then, centromeres fluctuated around the spindle pole, which relaxed chromosomes' configuration, allowing telomeres to form a Rabl-like distribution. In this configuration, centromeres were distributed around a sphere with a radius ℓ^* around the spindle pole. Because smaller chromosomes are more diffusive, centromeres of small chromosomes tend to diffuse along the sphere surface and reside in the forward positions along the spindle axis, leading to a bias in the chromosome distribution (Fig. 3g). This diffusion was necessary in the present model to avoid centromeres from clustering in telophase and interphase, so that the anaphase simulation was run long enough for 3×10^5 steps, or 30 minutes.

D. Telophase

At the onset of telophase, microtubules dissociate from kinetochores, and the nuclear membrane starts to form. Hence, the kinetochore-microtubule component U_{km} was turned off and instead a pre-nuclear packing component U_{pack} was turned on: $U = U_{chain} + U_{pack}$, where

$$U_{pack} = \sum_{i=1}^{N} u_{pack}(r_i) \qquad (20)$$

and u_{pack} represents a harmonic potential constraining the beads into a packing sphere

$$u_{pack}(r) = \begin{cases} 0 & \text{for } r \leq R_{pack}, \\ \frac{1}{2}K_{pack}\left(r - R_{pack}\right)^2 & \text{for } r > R_{pack}, \end{cases} \qquad (21)$$

The packing radius was set to $R_{pack} = 1.5\,\mu m$, so that the volume fraction of the beads is approximately $N\sigma^3/(2R_{pack})^3 \approx 0.6$, indicating a loose random packing. The spring constant was set to $K_{pack} = 50k_B T^*/\mu m^2$; this value is large enough to prevent chains from leaving the sphere, but much smaller than K in $u_{bond}(r)$, allowing chains to fluctuate near the packing surface. See Fig. 3h for the snapshot of the telophase structure. The effective temperature of $T = 4T^*$ allowed chromosomes to move inside the crowded packing sphere, which resolved the Rabl-like cluster of centromeres (Fig. 3j) formed during the anaphase, distributing centromeres homogeneously inside the packing sphere (Fig. 3k). The telophase simulation was run for 5×10^5 steps, or 50 minutes, to achieve sufficient movement. Here, the bias of chromosome distribution generated in the late stage of anaphase, in which small chromosomes are positioned in the advanced direction along the spindle axis (the y-axis in Fig. 3j), was left at the beginning of telophase. This bias was relaxed during telophase, but some bias remained at the end of telophase (Fig. 3k), which affected the distribution of chromosomes in the G1 phase as discussed in the subsection, *Inheritance and imprinting of radial distributions through cell cycles*, in the next section.

E. G1 phase

The spherical surface that packed the genome in telophase becomes the nuclear envelope. Inside this sphere, we fine-grained the chromosome chains to 100-kb resolution chains. See Appendix B for fine-graining. At the entry into the G1 phase, structural constraints due to condensins disappear, and chromosomes begin to decondensate, leading to genome expansion. In order to describe this expansion, it is natural to assume the repulsive interactions between 100-kb chromatin regions.

In addition, we need to consider the heterogeneity in repulsive interactions because the transcription activity is heterogeneous in the genome, which should affect the physical properties of each 100-kb region. Using the NCI analysis as explained in Appendix D, we distinguished the gene-active type-A regions, the gene-inactive or gene-poor type-B regions, and the intermediate type-u regions in the GM12878 genome. The previous 1-kb resolution model,[6] which takes account of the nucleosome-nucleosome attractive interactions and the cohesin activity within domains, showed that the repulsive interactions dominate at the 100-kb resolution when chromatin is packed in high density as in the nucleus, while both attractive and repulsive interactions coexist at the 1-kb level. The similar dominance of the repulsive interactions in the coarse-grained models of polymer solutions has been discussed.[52] The 100-kb type-A regions calculated with this 1-kb resolution model are more extended than type-B regions,

leading to the milder repulsive interactions between type-A regions, U_{AA}, and the harder repulsive interactions between type-B regions, U_{BB}. We defined the ABu interactions as $U_{\alpha\beta} = (U_{\alpha\alpha} + U_{\beta\beta})/2$ with α, β =A, B, or u. See the previous publication[6] for more details.

We performed the simulation of the G1 phase with the 100-kb resolution model using the thus defined interaction potentials, $U_{\alpha\beta}$. With these heterogeneous repulsive interactions, the genome expanded as cells entered the G1 phase (Fig. 3l). Through this expansion, the genome showed phase separation: type-A regions assembled into compartment A, type-B regions into compartment B, and type-u regions resided at the interface of compartments A and B. The LADs and NADs were spontaneously formed with this phase separation. Fig. 3m shows the genome structure after reaching the stationary state in the G1 phase.

We should note that the mean-field criterion of enegetically driven phase separation, the Flory-Huggins parameter of $\chi = U_{AB} - (U_{AA} + U_{BB})/2$, is $\chi = 0$ by definition in the present simulation, indicating that the phase separation is not driven energetically but entropically; the gathering of type-A regions allows their overlap due to the milder repulsion between them, leading to the system's larger entropy and resulting in the phase separation. The G1 phase simulation was stopped at 7 hr after the onset of the G1 phase as in the previous study.[6] Then, the phase-separated genome configuration was used for generating the prophase configuration in the next round of the cell cycle.

IV. MEMORY RETENTION THROUGH CELL CYCLES

To explore how the structural memory of the genome is retained or erased through cell cycles, we computed 30 independent trajectories of five consecutive cell cycles. The correlation functions, distributions, and contact maps described in this section are the averages over these 30 trajectories.

A. Memory retention and erasure during mitosis

During mitosis, chromosomes undergo significant movements, leading to the loss of some structural features of the interphase genome. However, certain essential components are retained through mitosis. We investigate these memory effects by calculating the autocorrelation between the positions of sister kinetochores at different time points in prometaphase and the autocorrelation between the positions of centromeres at different time points in anaphase. The positions' autocorrelation is anisotropic due to the chromosomes being pushed or pulled anisotropically from/to the spindle poles during mitosis. Therefore, we calculated two types of autocorrelation functions (ACFs): the axial ACF, which describes the autocorrelation of positions projected on the spindle axis, and the in-plane ACF, which describes the autocorrelation of positions projected on the plane perpendicular to the spindle axis. For the definitions of the axial ACF and the in-plane ACF, please refer to Appendix E. The ACFs in prometaphase to metaphase

are shown in Fig. 3d, and the in-plane ACF in anaphase is shown in Fig. 3i.

In Fig. 3d, the axial ACF was calculated from prometaphase to metaphase. The significant decay of the axial ACF during prometaphase indicates the intense structural mixing and the memory loss along the axis direction. Then, at metaphase, all chromosomes align in a thin layer perpendicular to the axis, resulting in the almost complete loss of memory along the spindle axis (Appendix E).

The in-plane ACF remains close to 1 in both Fig. 3d and Fig. 3i, indicating that the arrangement of chromosomes projected on the plane perpendicular to the spindle axis remains similar from the beginning of prometaphase (prophase) to the end of anaphase (beginning of telophase). The slow kinetics in the plane directions are caused by the crowded alignment of chromosomes. Thus, the genome conformation in the interphase is partially retained from prophase to the end of anaphase in the two-dimensional directions perpendicular to the spindle axis, but the conformation is intensely mixed along the spindle axis and reset at metaphase. This anisotropy in the memory is consistent with the observations from microscopy[53,54].

The axial structural order was newly generated in anaphase. Centromeres were dragged toward the spindle pole. Starting from metaphase (Fig. 3e), the large chromosomes at the periphery of the ring approached the spindle pole earlier than the small chromosomes located in the inner region of the ring at the early stage of anaphase (Fig. 3f). Then, after all chromosomes reached the sphere of approximate radius ℓ^* from the spindle pole, small chromosomes diffused more easily on the sphere surface than long chromosomes and appeared in the advanced direction along the spindle axis (Fig. 3g). This configuration became a starting configuration (Fig. 3j) of the subsequent relaxation in telophase.

During telophase, the retained memory in anaphase was partially erased due to the structural relaxation (Fig. 3h), but this process was incomplete and was cut short by the start of chromosome decondensation towards the G1 phase. Consequently, the structural features emphasized during mitosis persisted after telophase; large chromosomes located outside and small chromosomes inside on the plane perpendicular to the spindle axis, and small chromosomes tended to reside ahead along the spindle axis (Fig. 3k). These structural features were inherited by the G1-phase configuration. In the next subsection, we examine how these structural features change across multiple cell cycles.

B. Inheritance and imprinting of radial distributions through cell cycles

In telophase, microtubules detach from kinetochores, which allows chromosomes to relax from a Rabl-like configuration. However, the system can not equilibrate within a short duration of telophase. Therefore, a significant amount of positional memory from the previous interphase is carried over to the next interphase through telophase. Illustrative effects are shown in Fig.4a. The excess densities of chromosome 1 (chr1)

FIG. 4. Shifts in radial distributions of chromosomes through cell cycles and phases. (a) Radial distributions of chr1 and chr16 in the first G1 phase after RMP initialization (upper panel) and the second G1 phase (lower panel), shown as differences from the whole-genome reference density. (b) Growth of the radial distributions of chr1 (upper panel) and chr16 (lower panel) through cell cycles and phases. Tc, Ic, and Mc are telophase, interphase, and metaphase in the cth cycle, respectively. Excess density is plotted as areas in the histogram at each phase, with positive parts shown as solid areas and negative parts shown as hollow areas. (c) Average radial positions of kinetochores on the metaphase plate in each cell cycle. Cell cycle history is shown as dimmed solid lines, and the positioning at the 5th cycle is shown as dots. Radial positions of dots are color-coded in the range 1.7-2.0 μm, and the lower and the upper bounds of the range are shown as concentric circles in broken lines in the plot. The positioning at the 4th cycle (M4) is also shown in (d). (e) Average radial positions of centromeres in the G1 phase in each cell cycle. The projected radial positions projected on the last metaphase plate (left panel) and the three-dimensional radial positions (right panel) are shown. The positioning at the 5th cycle (I5), which follows after M4, is shown as dots in each panel.

and chromosome 16 (chr16) at the G1 phase in the first cycle (upper panel) and in the second cycle (lower panel) are plotted as functions of the normalized radial distance ($R/R_{\max}$) from the nuclear center. Here, as chromosomes distribute on average with the radial density of $\overline{\rho}(R) \sim R^2$, the excess density defined by $\rho(R) - \overline{\rho}(R)$ shows the characteristic distribution of each chromosome. See Appendix F for the precise definitions of $\rho(R)$ and $\overline{\rho}(R)$. In the first G1 phase after RMP configuration, chr1 and chr16 both had a higher probability of being located at the nuclear periphery (upper panel). As discussed earlier, smaller chromosomes tend to settle in the inner region of the metaphase plate. This memory in metaphase is inherited through telophase by the next G1 phase. As a result, after a single cell cycle, the radial distribution of chr16 became skewed towards the interior of the nucleus in the G1 phase, whereas chr1 remained near the nuclear periphery (lower panel).

Consecutive changes in the radial distributions over five cell cycles are tracked in Fig. 4b. The distribution of chr1 was con-

sistently biased towards the nuclear periphery throughout cell cycles. Meanwhile, the distribution of chr16 shifted in the second telophase (T2), remained interior in the second and third cycles, suddenly spread in the fourth telophase (T4), and then migrated back to the interior in the fifth telophase (T5). The inward migration of chr16 was initiated as a weak excess density of $R/R_{\max} < 0.5$ in the first and fourth metaphase (M1 and M4) and was amplified in the subsequent telophase (through anaphase).

The diagram in Fig. 4c shows the average radial positions of the midpoints of sister kinetochores on the metaphase plate. The kinetochores tend to align around the ring-shaped potential minimum defined by KMF and PEF. Longer chromosomes, such as chr1-5, tended to be biased outwards, while shorter ones, such as chr16-22, were biased inwards. Notably, medium-sized chromosomes chr9 and chr10 were located in the interior of the plane, and the smaller chr18, despite having a similar size to chr17 and chr19, was positioned as an outer-distributed chromosome. It is important to note

that in prometaphase, chromosomes are characterized solely by the lengths of the p- and q-arms. The longer the arms, the stronger the PEF and KMF, leading to the faster congression towards the metaphase plate. Once chromosomes reach the metaphase plate, they compete to fit their arms within the two-dimensional plate to reach the ring-shaped potential minimum. Additionally, the outward-directed PEF acts on the p- and q-arms of each chromosome, with the magnitude approximately proportional to the arms' length, which results in imbalanced tension on the kinetochores of the chromosomes. These factors collectively determine the characteristic positioning of chromosomes on the metaphase plate, which was consistently reproduced, if not completely, in every cycle (Figs. 4c and 4d). Consequently, from prometaphase to metaphase, the intrinsic radial distribution bias on chromosomes is imprinted over the previous distribution.

The daughter cells in the subsequent G1 phase inherit the characteristic radial positioning of chromosomes from metaphase. This inheritance is evident in the similarity between the metaphase radial distribution in the fourth cycle (Fig. 4d) and the in-plane radial distribution in the subsequent G1 phase starting the fifth cycle (left panel in Fig. 4e). In the G1 phase, the in-plane radial distribution is defined by the distribution projected on the same plane as the metaphase plate, even though the plate itself no longer exists. The full 3D radial distribution in the G1 phase (right panel in Fig. 4e) results from both the in-plane distribution and the out-plane distribution in the axial direction. The 3D radial distribution in the G1 phase for chr8-10 and chr19-22 differed significantly from their in-plane radial distribution, indicating the influence of the out-plane axial distribution. This axial distribution originates from the bias found in telophase (Fig. 3k), which arises from the chromosome distribution around a spindle pole at the late stage of anaphase. Therefore, the chromosomal assembly towards a spindle pole in anaphase also leaves an imprint on the 3D radial distribution, which is evident in the subsequent G1 phase.

Interestingly, the simulated in-plane distribution of autosomes in the G1 phase (left panel of Fig. 4e) appears similar to the microscopically observed radial distribution in lymphoblast cells,[18] with the exception of the two small chromosomes chr20 and chr21. However, the simulated 3D radial distribution in the G1 phase (right panel of Fig. 4e) does not match the microscopically observed distribution,[18] indicating that the simulation was placing too much emphasis on the axial distribution. This discrepancy could be due to always using the same axis direction in different cell cycles in simulations, whereas in natural cells, the axis direction can vary due to nuclear deformation and centrosome location change. While we fixed the axis direction in simulations to distinguish the in-plane and axial imprinting on the structural memory during mitosis, the fluctuation of the axis should be considered when comparing the simulated distribution with the observed one.

The simulated in-plane distribution agrees with the microscopically observed distribution for most chromosomes. Here, the discrepancy in reproducing the positioning of chr20 is likely due to the short arm length of chr20. We need to use a model with a finer resolution than the current 10-Mb res-

olution in order to accurately assess the PEF acting on the short arms. Additionally, the discrepancy in the positioning of chr21 may be attributed to the inaccurate evaluation of the rDNA length in the simulation. We were unable to compare the positioning of the X chromosomes because the microscopic observations were made on male lymphoblast cells,[18] while the simulations were based on female lymphoblastoid cells.

In this section, we have seen that the radial distribution of chromosomes in the interphase nucleus is partly carried over from one cell cycle to the next, with imprinting added to the distribution during mitosis. After exiting the interphase, cells progress through prophase, prometaphase, and metaphase. Through this process, the bias in the radial distribution shown in Fig. 4c is imprinted on memory from the previous interphase configuration. During metaphase, the chromosome distribution along the axis is reset as chromosomes are aligned on a two-dimensional metaphase plate. In anaphase, a new bias in the axial distribution is generated as chromosomes move toward a spindle pole. The resulting configuration relaxes during telophase and serves as the starting point for the next interphase. In the following sections, we explore how the growth of the radial distribution impacts genome-wide contacts between chromosomes and compartments in the interphase genome.

C. Contacts between chromosomes organized through cell cycles

The larger chromosomes show persistent positioning in their radial distribution due to the memory effects through cell cycles, and the smaller chromosomes tend to reside in the inner region in the in-plane distribution. We can highlight such size-dependent behaviors by monitoring the contacts between chromosomes in interphase. See Appendix G for the inter-chromosome contacts examined in this subsection.

Fig. 5a shows the inter-chromosome contact enrichment in the G1 phase in each cell cycle. Contacts between small chromosomes were more frequent than others, showing the tendency of small chromosomes positioned in a correlated way in the nucleus. The tendency was also manifested in the less frequent contacts between large and small chromosomes. Fig. 5a shows that the contrast in the contact enrichment became larger through multiple cell cycles, indicating the memory effects on the chromosome positioning in interphase. Fig. 5b summarizes this tendency, showing the gradual separation of large and small chromosomes through multiple cycles. The first cycle behaves differently because of the RMP approximation used in the preceding metaphase.

D. Compartments are consistently established in each cell cycle

The present simulations of cell cycles showed that memory of the metaphase-plate structure affects the distribution of chromosome territories in the next interphase, and mem-

FIG. 5. Inter-chromosomal contacts through cell cycles. (a) The inter-chromosomal $\log_2$ contact enrichment in each cell cycle. (b) The $\log_2$ contact enrichment within/between groups of large (L, chr1-5) and small (S, chr16-22 excluding chr18) chromosomes. (c) The genome-wide compartment PC1 signal along chr16 in each cell cycle. (d) Distribution of genome-wide compartment PC1 signals in each cell cycle.

ory of the interphase distribution is retained as the persistent in-plane distribution of chromosomes in the next metaphase plate. Through this partial memory persistence, distributions of chromosomes are gradually organized through multiple cell cycles though they bear large fluctuations. In the G1 phase, phase separation of type-A and type-B chromosomal segments results in the spatial compartmentalization of the genome (Fig. 3m). These compartments spread across different chromosomes; therefore, an intriguing question is whether the compartment distribution is partially inherited or further enhanced through cell cycles.

To investigate this issue, we have calculated the 1st principal component vector (PC1), which is an index distinguishing compartments, known as the compartment signal. To eliminate the contribution of possibly robust intra-chromosomal compartmentalization, we masked out the intra-chromosomal contacts when deriving PC1. See Appendix G on the derivation of the compartment signal, PC1.

Surprisingly, the genome-wide compartment signal was nearly invariant across cell cycles despite the repositioning of chromosomes (Fig. 5c and 5d). After the nucleus reaches the stationary state in the G1 phase, the genome shows only limited structural changes with small-scale fluctuations, and compartments established at the early stage of the G1 phase are not altered through the G1 phase. Therefore, the present result suggests that phase separation occurring in the early stage of each G1 phase is responsible for compartmentalization, and the inherited memory does not affect compartmentalization. This robustness is realized when each chromosome deforms and adjusts its local 1 to 10-Mb scale structures both inside each chromosome and at the interface between different chromosomes, even in the case the chromosome territory distribution largely fluctuates and differs from cell to cell.

The present multi-cell-cycle simulations showed that compartments are robust without being much affected by the

memory of the mitotic structures. The robust compartmentalization should be responsible for the stable regulation of transcription and replication, and this is in sharp contrast to the flexibility of the chromosome territory distribution.

V. DISCUSSION

We have developed a model of the 3D genome structure during mitosis. In this model, the active dynamics of microtubules cause forces that push chromosome arms away from the spindle pole (PEF) and drag chromosomes along k-fibre (KMF). Our simulation of mitosis demonstrated that the balance between PEF and KMF leads to the assembly of chromosomes into a rosette ring structure on the metaphase plate. In this structure, larger chromosomes tend to be located on the periphery, while smaller chromosomes tend to reside in the inner region of the ring. This structure reflects the memory of the previous interphase distribution of chromosome territory, as the structural memory in the plane perpendicular to the spindle axis persists from prophase to anaphase. This memory is further enhanced or modified by the competition among chromosomes and fluctuations during their assembly toward the metaphase plate. The distribution of chromosomes along the spindle axis becomes mixed in metaphase, losing its memory, but the memory perpendicular to the axis is partially retained through telophase and inherited by the subsequent interphase genome.

By integrating the mitosis model with the previously developed interphase genome model, we conducted simulations of consecutive cell cycles. As a result of memory retention and erasure during mitosis, the distribution of chromosome territories fluctuates in each cell cycle but consistently shows a tendency for small chromosomes to reside in the inner region and large chromosomes at the periphery in the in-

plane distribution. During interphase, contacts between small chromosomes are strengthened, while contacts between small and large chromosomes are weakened, as small chromosomes tend to cluster in the nucleus across multiple cycles. However, compartments remain robust against fluctuations in chromosome positioning occurring in every cell cycle. This demonstrates that the phase separation at the beginning of the G1 phase adjusts structures at the 1 to 10 Mb scale, generating compartments and regulating chromatin functions.

There are several important directions to improve the current simulation model. First, the movement of cells during interphase should impact the structure inside the nucleus. Rotations and deformations of the nucleus should cause long-wavelength chromatin movement inside the nucleus, as observed in microscopic measurements.[55] In the current model, chromosomes do not move much and memory is retained during interphase. However, the chromatin fluctuations induced by the movement of the nucleus may alter the memory. It is also interesting to explore how the fluctuation in the shape of the nucleus is influenced by the distribution of chromosomes inside the nucleus.

During cell division, the movement of cellular components can cause changes in the orientation of the spindle axis by altering the positions of centrosomes.[56] While the structural characteristics along the spindle axis are mixed and reset during mitosis, the memory of structures in the plane perpendicular to the axis is partially retained. As a result, the angular distribution in this plane is influenced by the orientation of the spindle. The increased contrast of the contact pattern depicted in Figs. 5a and 5b should be reviewed to determine whether the contrast remains consistent even when the axis direction fluctuates significantly across cell cycles.

During prometaphase and metaphase, mitotic chromosomes oscillate due to the intermittent switching of microtubules between elongation and shortening modes.[30,57,58] These oscillations were not considered in the present study, and a static potential field was used as the basis for the simulation. However, it is possible that the oscillations could alleviate the congestion of chromosomes on the metaphase plate, leading to a larger repositioning of chromosomes in metaphase compared to the present simulation (see Fig. 3d). It would be interesting to investigate whether such repositioning contributes to a size-dependent distribution of chromosomes or results in a more homogeneous distribution.

Finally, the model should be thoroughly tested using a different set of parameters and a larger number of trajectories. Although the current model has its limitations, it opens up new research opportunities. It is particularly important to investigate whether the positioning of the centrosomes and the change in spindle axis direction are regulated during the differentiation process to control chromosome distribution in daughter cells.[59,60] Mouse rod cells, which have an inverted nuclear structure with heterochromatin located in the inner region of the nucleus,[61,62] should be examined to understand nuclear generation throughout cell cycles. The combined efforts of simulating mitosis and the interphase chromosomes should provide new insights into the cellular processes.

ACKNOWLEDGMENTS

This work was supported by JSPS-KAKENHI Grants 22H00406 and 24H00061.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study were derived by using the computational codes, which are available in a repository, https://github.com/snsinfu/3d-genome-cycle

CONFLICT OF INTEREST STATEMENT

The authors have no conflicts to disclose.

Appendix A: Coarse-graining

We write the 3D position vector of the mth bead in the 100-kb resolution genome as $\boldsymbol{r}'_m$ with $1 \leq m \leq N'$, where N' is the total number of 100-kb beads in the genome. We write the 3D position vector of the ith bead in the 10-Mb resolution model of the mitotic genome as $\boldsymbol{r}_i$ with $1 \leq i \leq 2N$, where N is the number of 10-Mb beads in each daughter genome.

Given the G1 phase genome structure $\{\boldsymbol{r}'_m\}_{m=1}^{N'}$ and the ranges of chromosomal beads $s'_k \leq m \leq e'_k$ for each chain $k = 1, \ldots, 2n$ ($n = 23$), the coarse-grained coordinates $\{\boldsymbol{r}_i\}_{i=1}^{N}$ are defined as the centroids of corresponding regions in the original structure:

$$\boldsymbol{r}_{s_k+i} = \frac{1}{M} \sum_{m=0}^{M-1} \boldsymbol{r}'_{s'_k+iM+m}, \quad 0 \leq i < \frac{L'_k}{M}. \tag{A1}$$

Here, $L'_k = e'_k - s'_k + 1$ is the number of beads constituting the kth chain in the 100-kb resolution structure, and $M = 10\text{Mb}/100\text{kb} = 100$ is the coarsening factor. The coarse ranges of chromosomal beads $s_k \leq i \leq e_k$ are defined as $s_1 = 1$, $e_k + 1 = s_{k+1}$, and $e_{2n} = N$.

Appendix B: Fine-graining

We use cubic spline functions to interpolate the coarse-grained structures. $\boldsymbol{r}_i = (x_i, y_i, z_i)$ with $s_k \leq i \leq e_k$ is the 3D position vector of the ith bead on the chain k. Writing the 3D curve of the 10-Mb resolution chain k as $\tilde{\boldsymbol{r}}^k = (\tilde{x}^k, \tilde{y}^k, \tilde{z}^k)$ with $\tilde{x}^k = (x_{s_k}, \ldots, x_i, \ldots, x_{e_k})$, $\tilde{y}^k = (y_{s_k}, \ldots, y_i, \ldots, y_{e_k})$, and $\tilde{z}^k = (z_{s_k}, \ldots, z_i, \ldots, z_{e_k})$, we trace the curve $\tilde{\boldsymbol{r}}^k$ with a vector of cubic spline functions $(f_x^k(u), f_y^k(u), f_z^k(u))$. Here, u is the curve parameter, satisfying $(f_x^k(u_i), f_y^k(u_i), f_z^k(u_i)) = (x_i, y_i, z_i)$ with

$$u_i = \frac{i - s_k + 0.5}{e_k - s_k + 1} \qquad (s_k \leq i \leq e_k). \tag{B1}$$

Then, we obtain the interpolated coordinates $r'_m = (x'_m, y'_m, z'_m) = (f^k_x(u_m), f^k_y(u_m), f^k_z(u_m))$ of fine-grained beads, using

$$u_m = \frac{m - s'_k + 0.5}{e'_k - s'_k + 1} \qquad (s'_k \le m \le e'_k). \qquad \text{(B2)}$$

r'_m is used as the initial configuration of the 100-kb resolution model at the entry into the G1 phase.

Appendix C: Random metaphase approximation

At the 0th round of cell cycle, the configuration with the random metaphase (RMP) approximation was used as the initial configuration of the simulation. In the RMP configuration, each chromosome chain is a straight rod having a random direction. The RMP coordinate of the ith bead ($s_k \le i \le e_k$) of the kth chain is

$$r_i = g_k + \left(i - \frac{s_k + e_k}{2}\right) \sigma w_k, \qquad \text{(C1)}$$

where $\sigma = 0.3$ μm and g_k is a random 3D point normally distributed with the standard deviation of $1\,\mu$m. w_k is a unit vector having a random direction.

Appendix D: Type-ABu annotation with the NCI classification of chromatin regions

As cells enter the G1 phase, transcription becomes active, and the genome starts to show heterogeneous features as mixtures of gene-active type-A regions and gene-inactive or gene-poor type-B regions. To simulate the G1 genome, we need to distinguish these different types of chromatin regions. *Neiboring region Contact Index* (NCI) is a simple method for classifying chromatin regions. This index estimates the relative contact frequencies within a 50-100 kb scale. These local contact frequencies should come from the formation of functional complexes, such as the enhancer-promoter complex and the replication complex, reflecting the functional activity of those local regions. Using the 1-kb resolution contact matrix, m_{kl}, obtained from the Hi-C measurements,[8] we define the contact counts between the ith 50-kb region and the jth 50-kb region as

$$C_{i,j} = \sum_{k \in i\text{th region}} \sum_{l \in j\text{th region}} m_{kl}. \qquad \text{(D1)}$$

Then, the NCI at a 50-kb resolution is defined as

$$W_{50\text{kb}}(i) = \frac{1}{2}\left[\frac{C_{i,i+1}}{\sqrt{C_{i,i}C_{i+1,i+1}}} + \frac{C_{i,i-1}}{\sqrt{C_{i,i}C_{i-1,i-1}}}\right]. \qquad \text{(D2)}$$

We should note that when the correction for the bias in the Hi-C measurements, $b(i)$, is defined at the ith site, these bias-correction factors are canceled between denominator and numerator in $W_{50\text{kb}}(i)$. In other words, the NCI already accounts for this bias correction. We smoothed $W_{50\text{kb}}(i)$ over a 500-kb range around the kth 100-kb region and obtained the smoothed NCI, $w(k)$. Using the Z score of $w(k)$, $Z_w(k)$, we classified each 100-kb region into type A for $Z_w \ge 0.3$, type B for $Z_w \le -0.3$, and type u for $-0.3 < Z_w < 0.3$. For more detailed explanations, please refer to the previous publications.[6,16]

We performed the simulation of the G1 phase with this ABu annotation of 100-kb regions. In this annotation, we used the Hi-C contact map near the diagonal part, but we did not use further information from the Hi-C contact map in the rest of the simulation. The NCI provides a convenient method to annotate chromatin regions, but instead, it is possible to use the other method based on the ChIp-seq data of histone modifications or other biochemical data for the ABu annotation of 100-kb regions.

Appendix E: Autocorrelation functions

Autocorrelation functions (ACFs) of positions of centromeres in anaphase or sister kinetochores in prometaphase are calculated to quantify the memory of chromosome positioning. The degree of memory erasure is anisotropic because of the directed forces from mitotic spindle. Therefore, ACFs are calculated for the direction along the spindle axis, giving *axial ACF*, and for the directions perpendicular to the spindle axis, giving *in-plane ACF*.

Consider a trajectory $\{q_k(t)\}^{2n}_{k=1}$. $q_k = r_{i(k)}$ in anaphase, where $i(k)$ is the bead covering the centromere of the kth chromosome. For prometaphase structures, we use

$$q_k = \frac{r_{i(k)} + r_{i(k)+N}}{2}, \qquad \text{(E1)}$$

where $i(k)$ is the index of the kinetochore bead of the kth chromosome. The in-plane ACF, $C_{\text{ip}}(t)$, is defined by using configuration vectors projected on the plane perpendicular (the xz plane in the present study) to the spindle axis (the y axis),

$$Q_{\text{ip}}(t) = \big(q_{1,\text{x}}(t) - \overline{q}_{\text{x}}(t),\ q_{1,\text{z}}(t) - \overline{q}_{\text{z}}(t),$$
$$\dots,\ q_{2n,\text{x}}(t) - \overline{q}_{\text{x}}(t),\ q_{2n,\text{z}}(t) - \overline{q}_{\text{z}}(t)\big), \qquad \text{(E2)}$$

where, $\overline{q}(t) = 1/(2n)\sum^{2n}_{k=1} q_k(t)$ is the whole-genome centroid at time t. Then, the in-plane ACF is calculated as

$$C_{\text{ip}}(t) = \left\langle \frac{Q_{\text{ip}}(t) \cdot Q_{\text{ip}}(0)}{|Q_{\text{ip}}(t)|\,|Q_{\text{ip}}(0)|} \right\rangle, \qquad \text{(E3)}$$

where, the average $\langle \cdot \rangle$ is taken over the population of independent simulations. The axial ACF, $C_{\text{ax}}(t)$, is defined in the same way, except for the configuration vector being projected on the spindle axis as

$$C_{\text{ax}}(t) = \left\langle \frac{Q_{\text{ax}}(t) \cdot Q_{\text{ax}}(0)}{|Q_{\text{ax}}(t)|\,|Q_{\text{ax}}(0)|} \right\rangle, \qquad \text{(E4)}$$

with

$$Q_{\text{ax}}(t) = \big(q_{1,\text{y}}(t) - \overline{q}_{\text{y}}(t),\ \dots,\ q_{2n,\text{y}}(t) - \overline{q}_{\text{y}}(t)\big). \qquad \text{(E5)}$$

The ACFs, $C_{ax}(t)$ and $C_{ip}(t)$, are evaluated to 1 at $t = 0$ and decay with time as the memory of centromere positioning is erased.

We should note that the ACF is defined by a correlation between the normalized positional vectors. When the vector has a small length, a small normalization factor in the denominator enlarges the ACF value. This indeed happens for the axial ACF at metaphase, where all kinetochores align in a thin layer near the metaphase plate. Therefore, the finite axial ACF value of about 0.5 at metaphase, shown in Fig. 3d, does not imply memory retention. Because the alignment of centromeres in a thin layer at the starting time point does not define a meaningful axial ACF in anaphase, we only show the in-plane ACF in Fig. 3i.

Appendix F: Analysis of radial distributions

The radial density $\rho_{\phi,k}(R)$ of each chromosome k is estimated from the simulation snapshots at telophase $\phi = T_c$, the G1 phase $\phi = I_c$, and metaphase $\phi = M_c$ in cell cycles $c = 1, 2, 3, 4, 5$.

First, a set of radial positions of the kth chromosome $R_{\phi,k}$ at phase ϕ are defined from the snapshots. The definition of $R_{\phi,k}$ is explained in the last part of this Appendix subsection (See Eqs. F5, F6, F7 and F8). Then, the maximum $R_{\phi,\max} = \max_k R_{\phi,k}$ is identified. The radial density is estimated from a histogram binned at $\zeta_m = R_{\phi,\max} m/B$ ($m = 0, 1, ..., B$) as

$$\rho_{\phi,k}(\zeta_m) = \frac{\left| \{ R_k \in R_{\phi,k} \mid \zeta_m \le R_k < \zeta_{m+1} \} \right|}{|R_{\phi,k}| R_{\phi,\max}/B}. \tag{F1}$$

In addition, the total density $\overline{\rho}_\phi(R)$ is estimated as a reference, from the total sample $R_\phi = \cup_{k=1}^{2n} R_{\phi,k}$,

$$\overline{\rho}_\phi(\zeta_m) = \frac{\left| \{ R \in R_\phi \mid \zeta_m \le R < \zeta_{m+1} \} \right|}{|R_\phi| R_{\phi,\max}/B}. \tag{F2}$$

Then, the excess density is defined for each chromosome as the deviation of $\rho_{\phi,k}(R)$ from the total density,

$$\Delta\rho_{\phi,k}(R) = \rho_{\phi,k}(R) - \overline{\rho}_\phi(R). \tag{F3}$$

Finally, the excess densities for homologous chromosomes are merged as

$$\Delta\rho'_{\phi,k}(R) = \frac{\Delta\rho_{\phi,k}(R) + \Delta\rho_{\phi,k+n}(R)}{2}, \tag{F4}$$

which was used for plotting Figs. 4a-4e.

The radial positions of chromosomes are sampled at telophase and interphase as the collection of the radial positions of all beads comprising each chromosomal chain,

$$R_{T_c,k} = \{ r_i(t) \mid s_k \le i \le e_k, t \in T_T \}, \tag{F5}$$

$$R_{I_c,k} = \{ r'_i(t) \mid s'_k \le i \le e'_k, t \in T_I \}, \tag{F6}$$

$$R_{ipI_c,k} = \{ (r'^2_{i,x}(t) + r'^2_{i,z}(t))^{1/2} \mid s'_k \le i \le e'_k, t \in T_I \}, \tag{F7}$$

in each cell cycle $c = 1, 2, 3, 4, 5$. $R_{ipI,k}$ is used to compute in-plane radial distribution in Fig. 4e. The sampled time points are the final 10 snapshots $T_T = \{49.1 \,\text{min}, \ldots, 50.0 \,\text{min}\}$ in telophase and $T_I = \{6.1 \,\text{hr}, \ldots, 7 \,\text{hr}\}$ in interphase. In metaphase, kinetochores settle in an inner region of a metaphase plate and chromosomal arms point outwards from there, forming a rosette structure. This ordered configuration produces misleading densities if whole chromosome chains are considered. Hence, we analyzed the distributions of kinetochores of sister chromatids (Eq. E1) in metaphase,

$$R_{M_c,k} = \{ q_k(t) \mid t \in T_M \}. \tag{F8}$$

The sampled time points are the final 10 snapshots $T_M = \{29.1 \,\text{min}, \ldots, 30.0 \,\text{min}\}$ in prometaphase.

Appendix G: Analysis of contacts and compartment signals

In each cell cycle, the genome-wide contact matrix at a 100-kb resolution was calculated as described,[6] using the contact distance of $0.24 \,\mu\text{m}$ and G1-phase structures from $t = 4 \,\text{hr}$ to $7 \,\text{hr}$. The contact matrix was balanced with the `balance_cooler` function in the Cooler package[63] and denoted as $C_{(c)}$, where $c = 1, 2, 3, 4, 5$ designates the cell cycle.

In the present study, we are interested in the effect of chromosomal repositioning on genome-wide compartmentalization in cycling cells. Therefore, to eliminate the contribution of possibly robust intra-chromosomal compartmentalization, we mask out the intra-chromosomal contacts when deriving the compartment signal. Specifically, we define the trans-observed-over-expected (TO/E) matrix as

$$X_{(c),ij} = \begin{cases} \dfrac{C_{(c),ij}}{\overline{C}_{(c)}}, & \text{if } (i,j) \in TC, \\ 1, & \text{otherwise}, \end{cases} \tag{G1}$$

where $TC = \{ (i,j) \mid s'_k \le i \le e'_k, \, s'_l \le j \le e'_l, \, 1 \le k, l \le n \}$ is the set of all inter-chromosomal pairs of bins, and $\overline{C}_{(c)}$ is the average inter-chromosomal contact frequency

$$\overline{C}_{(c)} = \frac{1}{|TC|} \sum_{(i,j) \in TC} C_{(c),ij}. \tag{G2}$$

The TO/E matrix then is standardized by the row-wise average and standard deviation,

$$\hat{X}_{(c),ij} = \frac{X_{(c),ij} - mX_{(c),\bullet j}}{sX_{(c),\bullet j}}, \tag{G3}$$

where

$$mX_{(c),\bullet j} = \langle X_{(c),i'j} \rangle_{i'} \tag{G4}$$

$$sX_{(c),\bullet j} = \langle (X_{(c),i'j} - mX_{(c),\bullet j})^2 \rangle_{i'}^{1/2}. \tag{G5}$$

Then, singular value decomposition (SVD) is applied on the standardized matrix $\hat{X}_{(c)}$, and the first singular value $\sigma_{(c),1}$ and associated singular vectors $\boldsymbol{u}_{(c)}, \boldsymbol{v}_{(c)}$ are obtained,

$$\hat{X}_{(c)} = \underbrace{\sigma_{(c),1} \boldsymbol{u}_{(c)} \boldsymbol{v}^\mathsf{T}_{(c)} + \cdots}_{\text{SVD}}. \tag{G6}$$

Finally, the first left singular vector $\boldsymbol{u}_{(c)}$ is normalized to its genome-wide maximum, giving the genome-wide compartment PC1 signal for cell cycle c,

$$\text{PC1}_{(c),i} = \frac{u_{(c),i}}{\max_{i'} u_{(c),i'}}, \tag{G7}$$

which was used to plot Figs 5c and 5d. In Fig. 5a, the log-transformed TO/E matrix for each cycle was plotted at the chromosome-level resolution. For each pair of chromosome (k,l), the chromosome-level O/E value $\tilde{X}_{(c),kl}$ is the average value

$$\tilde{X}_{(c),kl} = \frac{1}{|I(k)|\,|I(l)|} \sum_{i \in I(k)} \sum_{j \in I(l)} X_{(c),ij} \tag{G8}$$

where $I(k) = \{i \mid s'_k \le i \le e'_k\}$ is the set of indices of 100-kb bins covering chromosome k.

REFERENCES

[1] J. Walter, L. Schermelleh, M. Cremer, S. Tashiro, and T. Cremer, "Chromosome order in HeLa cells changes during mitosis and early G1, but is stably maintained during subsequent interphase stages," J Cell Biol **160**, 685–697 (2003).

[2] K. Maeshima, S. Ide, K. Hibino, and M. Sasai, "Liquid-like behavior of chromatin," Curr Opin Genet Dev **37**, 36–45 (2016).

[3] S. S. Ashwin, T. Nozaki, K. Maeshima, and M. Sasai, "Organization of fast and slow chromatin revealed by single-nucleosome dynamics," Proc Natl Acad Sci U S A **116**, 19939–19944 (2019).

[4] T. Nozaki, S. Shinkai, S. Ide, K. Higashi, S. Tamura, M. A. Shimazoe, M. Nakagawa, Y. Suzuki, Y. Okada, M. Sasai, S. Onami, K. Kurokawa, S. Iida, and K. Maeshima, "Condensed but liquid-like domain organization of active chromatin regions in living human cells," Sci Adv **9**, eadf1488 (2023).

[5] H. Strickfaden, T. O. Tolsma, A. Sharma, D. A. Underhill, J. C. Hansen, and M. J. Hendzel, "Condensed chromatin behaves like a solid on the mesoscale in vitro and in living cells," Cell **183**, 1772–1784.e13 (2020).

[6] S. Fujishiro and M. Sasai, "Generation of dynamic three-dimensional genome structure through phase separation of chromatin," Proc Natl Acad Sci U S A **119**, e2109838119 (2022).

[7] I. Solovei and L. Mirny, "Spandrels of the cell nucleus," Curr Opin Cell Biol **90**, 102421 (2024).

[8] S. S. Rao, M. H. Huntley, N. C. Durand, E. K. Stamenova, I. D. Bochkov, J. T. Robinson, A. L. Sanborn, I. Machol, A. D. Omer, E. S. Lander, and E. L. Aiden, "A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping," Cell **159**, 1665–1680 (2014).

[9] T. Nozaki, R. Imai, M. Tanbo, R. Nagashima, S. Tamura, T. Tani, Y. Joti, M. Tomita, K. Hibino, M. T. Kanemaki, K. S. Wendt, Y. Okada, T. Nagai, and K. Maeshima, "Dynamic organization of chromatin domains revealed by super-resolution live-cell imaging," Mol Cell **67**, 282–293 e7 (2017).

[10] T. Misteli, "The self-organizing genome: Principles of genome architecture and function," Cell **183**, 28–45 (2020).

[11] P. G. de Gennes, *Scaling Concepts in Polymer Physics* (Cornell University Press, 1979).

[12] A. Bolzer, G. Kreth, I. Solovei, D. Koehler, K. Saracoglu, C. Fauth, S. Müller, R. Eils, C. Cremer, M. R. Speicher, and T. Cremer, "Three-dimensional maps of all chromosomes in human male fibroblast nuclei and prometaphase rosettes," PLoS Biol **3**, e157 (2005).

[13] T. Cremer and M. Cremer, "Chromosome territories," Cold Spring Harb Perspect Biol **2**, a003889 (2010).

[14] T. Cremer, M. Cremer, and C. Cremer, "The 4D nucleome: Genome compartmentalization in an evolutionary context," Biochemistry (Mosc) **83**, 313–325 (2018).

[15] C. Hoencamp, O. Dudchenko, A. M. O. Elbatsh, S. Brahmachari, J. A. Raaijmakers, T. van Schaik, A. S. Cacciatore, et al., "3D genomics across the tree of life reveals condensin II as a determinant of architecture type," Science **372**, 984–989 (2021).

[16] S. Fujishiro and M. Sasai, "Three-dimensional simulation of whole-genome structuring through the transition from anaphase to interphase," in *Computational Methods for 3D Genome Analysis*, Methods in Molecular Biology, edited by R. Nakato (Springer Nature, 2024) pp. 293–308.

[17] J. H. Su, P. Zheng, S. S. Kinrot, B. Bintu, and X. Zhuang, "Genome-scale imaging of the 3D organization and transcriptional activity of chromatin," Cell **182**, 1641–1659.e26 (2020).

[18] S. Boyle, S. Gilchrist, J. M. Bridger, N. L. Mahy, J. A. Ellis, and W. A. Bickmore, "The spatial organization of human chromosomes within the nuclei of normal and emerin-mutant cells," Human Mol Genet **10**, 211–219 (2001).

[19] R. Nagele, T. Freeman, L. McMorrow, and H. Y. Lee, "Precise spatial positioning of chromosomes during prometaphase: evidence for chromosomal order," Science **270**, 1831–1835 (1995).

[20] A. Mogilner, R. Wollman, G. Civelekoglu-Scholey, and J. Scholey, "Modeling mitosis," Trends Cell Biol **16**, 88–96 (2006).

[21] J. R. McIntosh, M. I. Molodtsov, and F. I. Ataullakhanov, "Biophysics of mitosis," Q Rev Biophys **45**, 147–207 (2012).

[22] G. Civelekoglu-Scholey and D. Cimini, "Modelling chromosome dynamics in mitosis: a historical perspective on models of metaphase and anaphase in eukaryotic cells," Interface Focus **4**, 20130073 (2014).

[23] S. Reber and A. A. Hyman, "Emergent properties of the metaphase spindle," Cold Spring Harb Perspect Biol **7**, a015784 (2015).

[24] D. Oriola, D. J. Needleman, and J. Brugués, "The physics of the metaphase spindle," Annu Rev Biophys **47**, 655–673 (2018).

[25] V. A. Valdez, L. Neahring, S. Petry, and S. Dumont, "Mechanisms underlying spindle assembly and robustness," Nat Rev Mol Cell Biol **24**, 523–542 (2023).

[26] R. Paul, R. Wollman, W. T. Silkworth, I. K. Nardi, D. Cimini, and A. Mogilner, "Computer simulations predict that chromosome movements and rotations accelerate mitotic spindle assembly without compromising accuracy," Proc Natl Acad Sci U S A **106**, 15708–15713 (2009).

[27] C. Edelmaier, A. R. Lamson, Z. R. Gergely, S. Ansari, R. Blackwell, J. R. McIntosh, M. A. Glaser, and M. D. Betterton, "Mechanisms of chromosome biorientation and bipolar spindle assembly analyzed by computational modeling," Elife **9** (2020), 10.7554/eLife.48787.

[28] M. A. Krivov, F. I. Ataullakhanov, and P. S. Ivanov, "Computer simulation of merotelic kinetochore-microtubule attachments: corona size is more important than other cell parameters," Chromosome Res **29**, 327–349 (2021).

[29] D. Oriola, F. Jülicher, and J. Brugués, "Active forces shape the metaphase spindle through a mechanical instability," Proc Natl Acad Sci U S A **117**, 16154–16159 (2020).

[30] R. V. Skibbens, V. P. Skeen, and E. D. Salmon, "Directional instability of kinetochore motility during chromosome congression and segregation in mitotic newt lung cells: a push-pull mechanism," J Cell Biol **122**, 859–875 (1993).

[31] S. W. Grill, K. Kruse, and F. Jülicher, "Theory of mitotic spindle oscillations," Phys Rev Lett **94**, 108104 (2005).

[32] O. Campàs and P. Sens, "Chromosome oscillations in mitosis," Phys Rev Lett **97**, 128102 (2006).

[33] J. Häfner, M. I. Mayr, M. M. Möckel, and T. U. Mayer, "Pre-anaphase chromosome oscillations are regulated by the antagonistic activities of Cdk1 and PP1 on Kif18A," Nat Commun **5**, 4397 (2014).

[34] F. A. Habermann, M. Cremer, J. Walter, G. Kreth, J. von Hase, K. Bauer, J. Wienberg, C. Cremer, T. Cremer, and I. Solovei, "Arrangements of macro- and microchromosomes in chicken cells," Chromosome Res **9**, 569–584 (2001).

[35] O. Takenouchi, Y. Sakakibara, and T. S. Kitajima, "Live chromosome identifying and tracking reveals size-based spatial pathway of meiotic errors in oocytes," Science **385**, eadn5529 (2024).

[36] A. N. Boettiger, B. Bintu, J. R. Moffitt, S. Wang, B. J. Beliveau, G. Fudenberg, M. Imakaev, L. A. Mirny, C. T. Wu, and X. Zhuang, "Super-resolution imaging reveals distinct chromatin folding for different epigenetic states," Nature **529**, 418–422 (2016).

[37] Y. T. Hsiao, I. H. Liao, B. K. Wu, H. C. Chu, and C. L. Hsieh, "Probing chromatin condensation dynamics in live cells using interferometric scat-

tering correlation spectroscopy," Commun Biol **7**, 763 (2024).

[38]M. Sadaie, R. Salama, T. Carroll, K. Tomimatsu, T. Chandra, A. R. Young, M. Narita, P. A. Pérez-Mancera, D. C. Bennett, H. Chong, H. Kimura, and M. Narita, "Redistribution of the Lamin B1 genomic binding profile affects rearrangement of heterochromatic domains and SAHF formation during senescence," Genes Dev **27**, 1800–8 (2013).

[39]J. Kind, L. Pagie, H. Ortabozkoyun, S. Boyle, S. S. de Vries, H. Janssen, M. Amendola, L. D. Nolen, W. A. Bickmore, and B. van Steensel, "Single-cell dynamics of genome-nuclear lamina interactions," Cell **153**, 178–92 (2013).

[40]S. Dillinger, T. Straub, and A. Németh, "Nucleolus association of chromosomal domains is largely maintained in cellular senescence despite massive nuclear reorganisation," PLoS One **12**, e0178821 (2017).

[41]N. Naumova, M. Imakaev, G. Fudenberg, Y. Zhan, B. R. Lajoie, L. A. Mirny, and J. Dekker, "Organization of the mitotic chromosome," Science **342**, 948–953 (2013).

[42]J. H. Gibcus, K. Samejima, A. Goloborodko, I. Samejima, N. Naumova, J. Nuebler, M. T. Kanemaki, L. Xie, J. R. Paulson, W. C. Earnshaw, L. A. Mirny, and J. Dekker, "A pathway for mitotic chromosome formation," Science **359**, eaao6135 (2018).

[43]S. Cane, A. A. Ye, S. J. Luks-Morgan, and T. J. Maresca, "Elevated polar ejection forces stabilize kinetochore-microtubule attachments," J Cell Biol **200**, 203–18 (2013).

[44]H. Maiato, A. M. Gomes, F. Sousa, and M. Barisic, "Mechanisms of chromosome congression during mitosis," Biology (Basel) **6**, 13 (2017).

[45]G. J. Brouhard and A. J. Hunt, "Microtubule movements on the arms of mitotic chromosomes: Polar ejection forces quantified *in vitro*," Proc Natl Acad Sci U S A **102**, 13903–13908 (2005).

[46]J. K. Monda and I. M. Cheeseman, "The kinetochore–microtubule interface at a glance," J Cell Sci **131**, jcs214577 (2018).

[47]V. Varga, J. Helenius, K. Tanaka, A. A. Hyman, T. U. Tanaka, and J. Howard, "Yeast kinesin-8 depolymerizes microtubules in a length-dependent manner," Nat Cell Biol **8**, 957–962 (2006).

[48]V. Varga, C. Leduc, V. Bormuth, S. Diez, and J. Howard, "Kinesin-8 motors act cooperatively to mediate length-dependent microtubule depolymerization," Cell **138**, 1174–1183 (2009).

[49]M. I. Mayr, S. Hümmer, J. Bormann, T. Grüner, S. Adio, G. Woehlke, and T. U. Mayer, "The human kinesin Kif18A is a motile microtubule depolymerase essential for chromosome congression," Curr Biol **17**, 488–498 (2007).

[50]S. L. Kline-Smith and C. E. Walczak, "Mitotic spindle assembly and chromosome segregation," Mol Cell **15**, 317–327 (2004).

[51]S. Soeda, K. Yamada-Nomoto, and M. Ohsugi, "The microtubule-binding and coiled-coil domains of Kid are required to turn off the polar ejection force at anaphase," J Cell Sci **129**, 3609–3619 (2016).

[52]V. Krakoviack, J. P. Hansen, and A. A. Louis, "Influence of solvent quality on effective pair potentials between polymers in solution," Phys Rev E **67**, 041801 (2003).

[53]D. Gerlich, J. Beaudouin, B. Kalbfuss, N. Daigle, R. Eils, and J. Ellenberg, "Global chromosome positions are transmitted through mitosis in mammalian cells," Cell **112**, 751–764 (2003).

[54]H. Strickfaden, A. Zunhammer, S. van Koningsbruggen, D. Köhler, and T. Cremer, "4D chromatin dynamics in cycling cells: Theodor Boveri's hypotheses revisited," Nucleus **1**, 284–297 (2010).

[55]A. Zidovska, D. A. Weitz, and T. J. Mitchison, "Micron-scale coherence in interphase chromatin dynamics," Proc Natl Acad Sci U S A **110**, 15555–15560 (2013).

[56]V. Nunes, M. Dantas, D. Castro, E. Vitiello, I. Wang, N. Carpi, M. Balland, M. Piel, P. Aguiar, H. Maiato, and J. G. Ferreira, "Centrosome-nuclear axis repositioning drives the assembly of a bipolar spindle scaffold to ensure mitotic fidelity," Mol Biol Cell **31**, 1675–1690 (2020).

[57]A. Desai and T. J. Mitchison, "Microtubule polymerization dynamics," Annu Rev Cell Dev Biol **13**, 83–117 (1997).

[58]L. Laan, J. Husson, E. L. Munteanu, J. W. Kerssemakers, and M. Dogterom, "Force-generation and dynamic instability of microtubule bundles," Proc Natl Acad Sci U S A **105**, 8920–8925 (2008).

[59]J. A. Kaltschmidt, C. M. Davidson, N. H. Brown, and A. H. Brand, "Rotation and asymmetry of the mitotic spindle direct asymmetric cell division in the developing central nervous system," Nat Cell Biol **2**, 7–12 (2000).

[60]N. Tang and W. F. Marshall, "Centrosome positioning in vertebrate development," J Cell Sci **125**, 4951–4961 (2012).

[61]I. Solovei, M. Kreysing, C. Lanctot, S. Kosem, L. Peichl, T. Cremer, J. Guck, and B. Joffe, "Nuclear architecture of rod photoreceptor cells adapts to vision in mammalian evolution," Cell **137**, 356–368 (2009).

[62]I. Solovei, A. S. Wang, K. Thanisch, C. S. Schmidt, S. Krebs, M. Zwerger, T. V. Cohen, D. Devys, R. Foisner, L. Peichl, H. Herrmann, H. Blum, D. Engelkamp, C. L. Stewart, H. Leonhardt, and B. Joffe, "Lbr and lamin a/c sequentially tether peripheral heterochromatin and inversely regulate differentiation," Cell **152**, 584–598 (2013).

[63]N. Abdennur and L. A. Mirny, "Cooler: scalable storage for Hi-C data and other genomically labeled arrays," Bioinformatics **36**, 311–316 (2020).