Social Experience Shapes Fighting Strategies in *Drosophila*

Can Gao[1*], Mingze Ma[1*], Jie Chen[1], Xiaoxiao Ji[1], Qionglin Peng[1#], Yufeng Pan[1,2#]

Affiliations

[1]The Key Laboratory of Developmental Genes and Human Disease, Jiangsu Key Laboratory of Brain Science and Medicine, School of Life Science and Technology, Southeast University, Nanjing, 210096, China.

[2]Co-innovation Center of Neuroregeneration, Nantong University, Nantong, 226019, China.

[*]These authors contributed equally to this study.

[#]Correspondence to: pany@seu.edu.cn (Y.P.) or pengqionglin@seu.edu.cn (Q.P.)

Abstract

Social isolation generally increases aggression but decreases mating competition, resulting in an intricate and ambiguous relationship between social experience, aggression, and reproductive success. In male *Drosophila*, aggression is often characterized by lunging, a frequent and comparatively low-intensity combat behavior. Here, we provide a behavioral paradigm for studying a less frequent but more vigorous fighting form known as tussling. While social enrichment decreases lunging, aligning with past observations, it heightens the more forceful tussling behavior. These two forms of aggression rely on different olfactory receptor neurons, specifically Or67d for lunging and Or47b for tussling. We further identify three pairs of central pC1 neurons that specifically promote tussling. Moreover, shifting from lunging to tussling in socially enriched males is accompanied with better territory control and mating success. Our findings identify distinct sensory and central neurons for two fighting forms and suggest that social experience may shape fighting strategies to optimize reproductive success.

Keywords: Fighting strategies, social experience, aggression, *Drosophila melanogaster*, *Or47b* ORNs, pC1 neurons, reproductive competition

Introduction

Competition for resources such as territory, food, and mates is crucial for animal survival and reproduction (Bergman et al., 2010; Hoffmann and Cacoyianni, 1990; Stockley and Campbell, 2013; Zwarts et al., 2012). Among diverse competitive strategies, physical combat is a prevalent form of fighting to repel rivals and regulated by both genetic factors and social environment (Chen, 2022; Zwarts et al., 2012). Understanding how social experience influences combat behaviors in animals to maximize their survival and reproduction remains a fundamentally important yet inadequately understood question.

In the fruit fly, *Drosophila melanogaster*, inter-male aggression includes relative low-intensity fights such as lunging, in which one fly rears up on its hind legs and snaps down onto its opponent, as well as relative high-intensity fights like tussling, where two males tumble over each other (Certel and Kravitz, 2012; Chen et al., 2002; Hoopfer, 2016; Palavicino-Maggio and Sengupta, 2022). A recent work showed that lunges are insufficient to establish dominance but mainly serves to maintain social status, while tussling frequence is correlated with dominance establishment (Simon and Heberlein, 2020). It is noteworthy that the majority of studies focus on the low-intensity lunging behavior, owing to its higher frequency and ease of observation. Substantial studies have identified regulatory genes and the neural circuit underlying the lunging-based aggressive behaviors (Hoopfer, 2016; Kravitz and Fernandez, 2015). In particular, the male-specific volatile pheromone 11-*cis*-vaccenyl acetate (cVA) acutely promotes lunging behavior via its olfactory receptor Or67d (Wang and Anderson, 2010). Interestingly, chronic cVA exposure in socially-enriched males reduces the level of lunging via another olfactory receptor Or65a (Liu et al., 2011). Consistently, it has been found that social isolation increases lunging frequency (Palavicino-Maggio and Sengupta, 2022; Yadav et al., 2024) by altering the expression of genes such as *Cyp6a20* (Wang et al., 2008), *Drosulfakinin* (*Dsk*) (Agrawal et al., 2020), and *Tachykinin* (*Tk*) (Asahina et al., 2014). Social isolation also increases male aggression in mice (Matsumoto et al., 2005; Zelikowsky et al., 2018) and even in humans (Ferguson et al., 2005). In mice, the TK homologue Tachykinin 2 (Tac2) responds to social isolation and promotes inter-male aggressive behaviors (Zelikowsky et al., 2018). These findings contribute to the consensus that social isolation escalates aggression, whereas social enrichment mitigates it in animal models.

Meanwhile, social enrichment increases male mating success by enhancing the sensitivity of *Or47b*-expressing olfactory receptor neurons (ORNs) to fly pheromones common to both sexes (Lin et al., 2016; Sethi et al., 2019). Artificially activating *Or47b* ORNs also improves mating advantage in a competitive assay (Dweck et al., 2015). Group housing and age-dependent juvenile hormone (JH) jointly increase the expression of the male-specific proteins from *fruitless* (*fru*[M]), which is a master gene for male courtship behaviors (Peng et al., 2021; Sato et al., 2019; Yamamoto and

Koganezawa, 2013), in the *Or47b*-expressing ORNs (Hueston et al., 2016; Sethi et al.,
2019; Zhao et al., 2020). Notably, courtship levels in single-housed (SH) wild-type
males are equal to (Dankert et al., 2009; Pan and Baker, 2014), in some cases, even
stronger than those in group-housed (GH) males (Dankert et al., 2009). In fact, group-
housing decreases excitability of the courtship-promoting P1 neurons (Inagaki et al.,
2014), which express both terminal genes in the sex determination hierarchy, *fru* and
doublesex (*dsx*). These findings raise a question on how social enrichment, which
suppresses lunging-based aggression and does not increase courtship intensity, would
facilitate mating competition.
Here, we provide a behavioral paradigm for studying inter-male tussling in
Drosophila and discover that this infrequent yet intense form of combat is enhanced
by social enrichment, while the low-intensity lunging is suppressed by social
enrichment. This social enhancement of tussling requires olfactory receptor Or47b,
but not Or67d or Or65a. We further identify three pairs of central *dsx*-expressing pC1
neurons that specifically promote tussling. Moreover, we demonstrate that socially
enriched males can overcome mating disadvantages associated with aging.
92
93 **Results**
94
A Paradigm for High-intensity Aggression in *Drosophila*
While studying the lunging-based aggression in *Drosophila* using a high-throughput
aggression assay based on previously aggression paradigms (Certel and Kravitz, 2012;
Wu et al., 2020; Zhou et al., 2008), we occasionally observed the high-intensity
boxing and tussling behavior in male flies as previously reported (Chen et al., 2002;
Nilsen et al., 2004), which are difficult to distinguish and hereafter simply referred to
as tussling. We reasoned that the intensity, instead of the frequency, could be decisive
for a combat (Simon and Heberlein, 2020). Thus, although tussling is a less frequent
behavior, it could serve a vital function.
To induce a higher level of tussling for further manipulation, we first introduced
a fixed live virgin female in the same behavioral chamber for assaying lunging. The
female fly was immobilized in the middle of the food, with its head embedded in the
food and its body exposed to enhance male competition (Figure 1A). We found that
males that were group-housed for 7 days showed a significantly higher frequency of
tussling behavior and reduced latency to tussle in the presence of an embedded virgin
female (Figure 1B). We next compared tussling in male flies of different ages. We
observed a significantly higher proportion of 14-day-old males raised in groups that
showed tussling behavior during the 2-hour observation period, compared to the 7-
day-old males (Figure 1B). Furthermore, 2-day-old males never displayed tussling
behavior. In contrast, 14-day-old males showed a lower level of lunging behavior than
7-day-old males did (Figure 1-figure supplement 1). Previous studies have shown that,
in addition to mating resources, food quality is another crucial factor affecting
aggressive behaviors (Lim et al., 2014). Therefore, we next assayed tussling behavior

in 14-day-old males in the presence of an embedded virgin female with different
concentrations of sucrose or yeast in the food during the behavioral test. We found
that these males displayed similar levels of tussling behavior with different
concentrations of sucrose in the food (Figure 1C). On the contrary, male tussling
intensity increased significantly with higher yeast concentrations such that more than
70% of males displayed this high-intensity fighting behavior (Figures 1D, see Figure
1-video supplement 1). Taken together, these results provide an efficient behavioral
paradigm for the high-intensity tussling behavior and indicate that older males display
more tussling but less lunging than younger ones.
127
Social Enrichment Inhibits Low-intensity Lunging but Enhances High-intensity
Tussling
Previous studies have shown that social enrichment inhibits aggressive behaviors,
specifically the low-intensity lunging, in male flies (Yadav et al., 2024). Therefore,
we wondered whether social enrichment might also inhibit the high-intensity tussling
behavior in males. To answer this question, we compared both fighting behaviors in
SH and GH 14-day-old males (Figure 2A). Lunging and tussling events from each
group of representative samples were exhibited in raster plots for comparison (Figure
2B). We scored lunging behavior mainly in the first 10-min of test, due to its
relatively higher occurrence, and tussling behavior during the whole 2-hour
observational period since males take much longer time to initiate tussling (Figure
1D). Interestingly, we found that while SH males performed more frequent lunging
behavior in 10 min than GH males did, which is consistent with previous findings,
they rarely showed tussling behavior. In contrast, GH males displayed a large number
of tussling events within 2 hours (Figure 2B). Statistical analysis of tussling and
lunging behaviors showed that GH males exhibited reduced tussling latency and
enhanced tussling frequency, in addition to increased lunging latency and decreased
lunging frequency, compared to SH males (Figures 2C-F). We also quantified the
duration of each tussling and lunging events, and found that the average duration of
individual lunging events was less than 0.2 second, while a tussling event lasted from
seconds to minutes (Figure 2G). This further suggests that tussling is a more intense
form of aggressive behavior compared to lunging. These results indicate that social
enrichment inhibits low-intensity lunging but promotes high-intensity tussling in male
flies.
152
Or47b ORNs Mediate Social Experience-induced Tussling Behavior
To further investigate the sensory basis underlying the opposite regulation of lunging
and tussling behaviors by social experience, we tried to examine the role of cVA-
sensitive neurons, as well as other fru^M-expressing ORNs, in mediating intermale
aggression. Two cVA sensory neurons, *Or67d* and *Or65a* ORNs, play crucial roles in
enhancing the lunging-based aggression in SH males, in which *Or67d* ORNs mediate
acute cVA stimulation and promote male aggression (Wang and Anderson, 2010),

while *Or65a* ORNs mediate chronic cVA stimulation and inhibit male aggression in
GH males (Liu et al., 2011). In addition, the *fru^M*-expressing *Or47b* ORNs are
sensitive to a class of pheromones common to both sexes, such as palmitoleic acid
(PA) (Lin et al., 2016), and their sensitivity is enhanced under group rearing
conditions (Sethi et al., 2019). *Ir84a* ORNs, which are also *fru^M*-positive, are involved
in the perception of mixed food odors and promote male courtship behavior (Grosjean
et al., 2011). We silenced *Or67d*, *Or65a*, *Or47b* or *Ir84a* ORNs by expressing the
inwardly rectifying potassium channel Kir2.1 (Baines et al., 2001) and assayed
lunging and tussling behaviors in 14-day-old GH males. We found that silencing
Or47b ORNs, but not *Or67d*, *Or65a* or *Ir84a* ORNs, significantly decreased male
tussling, compared to the control (Figure 3A, B, see Table S1 for detailed genotypes).
In contrast, silencing *Or67d* ORNs, but not other ORNs, significantly reduced male
lunging (Figure 3C, D). These loss-of-function results suggest that *Or47b* ORNs
mediate male tussling behavior, whereas *Or67d* ORNs regulate male lunging behavior.
To elucidate the potential role of the Or47b receptor in mediating tussling
behavior, we knocked down its expression in *Or47b* ORNs using RNA interference
(RNAi). Our results showed a significant decrease in male tussling behavior when the
Or47b receptor expression was reduced (Figure 3E, F), indicating that the
functionality of *Or47b* ORNs in tussling is dependent on the Or47b receptor. As
previous studies suggested that the male-specific *fru^M* sustains *Or47b* expression in an
activity-dependent manner (Sethi et al., 2019), we next expressed microRNAs
targeting *fru^M* (*UAS-fruMi*) (Meissner et al., 2016) in *Or47b* ORNs and assayed male
tussling. The efficiency of *Or47b* and *fru^M* RNAi knockdown was both validated by
quantitative PCR (Figure 3-figure supplement 1). We observed a slight increase of
tussling latency in *fru^M* knocked-down males, although not significantly different
from the control (Figure 3E, F), suggesting that Or47b, rather than Fru^M, plays a more
crucial role in mediating male tussling.
To further confirm that the effect of social enrichment on tussling depends on the
activity of *Or47b* ORNs, we artificially activated *Or47b* ORNs by expressing the
bacterially derived sodium channel (NaChBac) (Ren et al., 2001) in 14-day-old SH
males. We found that these SH males with enhanced activity of *Or47b* ORNs showed
a significant increase in tussling behavior compared to control SH males. Interestingly,
the level of tussling in these SH males with activated *Or47b* ORNs was
indistinguishable from that in GH males (Figure 3G, H). These gain-of-function
results indicate that social enrichment acts through *Or47b* ORNs to promote male
tussling. Using the trans-Tango technique that labels downstream neurons through
ligand-receptor mediated signaling (Talay et al., 2017), we found substantially more
downstream signals of *Or47b* ORNs in GH males than those in SH males (Figure 3-
figure supplement 2), suggesting the involvement of *Or47b* ORNs under social
enrichment.
200
Distinct Central Brain Neurons for Lunging and Tussling

We next set out to identify the central neurons involved in regulating the high-
intensity tussling. We found that silencing the mushroom body neurons by expressing
an inwardly rectifying potassium channel Kir2.1 (Baines et al., 2001) driven by
R19B03-GAL4 did not affect male tussling, suggesting that the experience-dependent
tussling behavior is not dependent on the classical learning and memory center.
Surprisingly, inactivation of the previously-identified aggression-promoting P1[a] and
TK neurons did not affect tussling behavior (Figure 4A, B). These results imply that
the central circuit for tussling behavior may be distinct from the previously reported
aggression circuits. As *dsx*-expressing pC1 neurons, also termed P1 if co-expressing
fru[M], are crucial for sexual and aggressive behaviors in both males and females, we
also silenced two subsets of pC1 neurons (pC1[SS1] and pC1[SS2]), which were sexually
dimorphic, and their functions were previously characterized in females but not in
males (Wang et al., 2021). We found that inactivation of pC1[SS2] but not pC1[SS1]
neurons significantly reduced the tussling behavior in male flies (Figure 4A, B). The
cell bodies of pC1[SS2] and P1[a] neurons are both located in the center of posterior brain
and project to overlapping but distinct parts of the lateral protocerebral complex
(Figure 4C-E). These results suggest that pC1[SS2] neurons may be key central neurons
regulating tussling behavior.

To further compare the function of P1[a] and pC1[SS2] neurons in regulating male
behaviors, we performed optogenetic activation experiments on these neurons
expressing *CsChrimson*. Compared to control male pairs that did not show any
significant change upon red light stimulation (Figure 4F-H), optogenetic activation of
P1[a] neurons induced acute courtship behavior as indicated by unilateral wing
extension (UWE), and upon red light removal, there was a significant increase in
lunging but not tussling (Figure 4I-K), consistent with previous findings. In contrast,
optogenetic activation of pC1[SS2] neurons induced strong and acute tussling behavior,
with more than 50% of the time during red light in tussling, and the tussling behavior
ceased immediately upon removal of red light (Figure 4L, see Figure 4-video
supplement 1). Activation of pC1[SS2] neurons did not significantly affect either male
lunging or courtship (Figure 4M, N). These results indicate that pC1[SS2] neurons
acutely promote the high-intensity tussling behavior.

Previous studies have revealed a crucial role of P1[a] neurons in promoting a
persistent internal state that enhances male aggression (Hoopfer et al., 2015). To test
whether P1[a] neurons can promote both lunging and tussling behaviors, we further
performed thermal activation experiments. We found that activation of P1[a] neurons
via dTrpA1 at 30°C induced male tussling, just like activation of pC1[SS2] neurons did
(Figure 4-figure supplement 1). As thermal activation but not optogenetic activation
of P1[a] neurons induced tussling, we reasoned that vision might be a key factor for this
discrepancy, since all optogenetic experiments were performed in dark or red light.
Thus, we re-performed these thermal activation experiments in dark and found that
activation of pC1[SS2] but not P1[a] neurons induced tussling, although the level of
tussling by pC1[SS2]-activated males was lower compared to that in light, suggesting a

crucial role of vision in male tussling. These optogenetic and themogenetic activation experiments suggest that P1^a neurons can promote both lunging and tussling behaviors, but their function on tussling requires visual stimulation, while pC1^{SS2} neurons play a more direct role in acutely promoting male tussling.

We next used *retro*-Tango (Sorkaç et al., 2023) and *trans*-Tango (Talay et al., 2017) techniques to explore the potential upstream and downstream circuits of P1^a and pC1^{SS2} neurons, and observed substential differences in both conditions (Figure 4-figure supplement 2), suggesting largely parallel inputs and outputs for P1^a and pC1^{SS2} neurons.

Dsx^M Regulates the Development and Function of pC1^{SS2} Neurons

To further compare the differences between P1^a neurons and pC1^{SS2} neurons, we investigated whether P1^a and pC1^{SS2} neurons express *fru* and/or *dsx* using Fru^M and Dsx^M antibodies (Chen et al., 2021; Peng et al., 2022). We found that most P1^a neurons were *fru*-positive and *dsx*-positive, while pC1^{SS2} neurons were *dsx*-positive but *fru*-negative (Figure 5A-5D). These results suggest that *dsx* instead of *fru* may play crucial roles in the development and function of pC1^{SS2} neurons.

To investigate the function of *dsx* in pC1^{SS2} neurons, we knocked down *dsx* via RNAi and compared the morphological and functional changes of these neurons. We found that *dsx^M* knockdown in pC1^{SS2} neurons slightly increased the number of cells from 2-3 pairs to 3-4 pairs (Figures 5E-5G), and significantly lengthened the axonal projections ventrally (Figure 5H). We reasoned that these developmental abnormality may effect the tussling-promoting functions of pC1^{SS2} neurons, and tried to activate these neurons with or without knocking down *dsx*. We found that males with wild-type *dsx* function displayed intensive tussling behavior for approximately 50% and 70% of the total activation time, respectively, during the first and second activation periods, with the response returning to baseline levels after the removal of red light (Figure 5I, 5K). However, males with *dsx* knocked down in pC1^{SS2} neurons showed no tussling behavior during the first activation session, and few tussling behavior during the second activation session (Figure 5J, 5K). Together, these results reveal the crucial role of the *dsx* gene in the development and tussling-promoting function of pC1^{SS2} neurons.

Territory and Mating Advantage in Tussling-favored Experienced Males

As our results showed that GH males and SH males were more likely to perform the low-frequency, high-intensity tussling and the high-frequency, low intensity lunging, respectively, we wondered which fighting strategy might be more practical in the competition for territory and mating resources.

To at least partially address this question, we first designed an experimental paradigm for territory competition (Figure 6A). A circular territory, which is provided with food and a fixed virgin female and separated by a wall with a small door open to the outer area, was competed by two tester males, one of which was marked on their

wings. We manually analyzed the first 10 winning events, each was defined as one male successfully displacing the other from the inner resource area. The winning index was calculated as the proportional difference in the number of wins between the marked male and unmarked male. In control groups such as two GH males of the same age, we observed some 'winner takes all' phenomenon (one male won nearly all of the 10 encounters), but overall, the winning index was close to zero due to the randomness of marking (see methods). We further found that while GH 7-day-old males (G7) won slightly, but not significantly, more than SH 7-day-old males (S7) (Figure 6B), older G14 males had a significant advantage over the S14 males (Figure 6C and Figure 6-figure supplement 1, Table S2, see Figure 6-video supplement 1). Additionally, we observed that when G14 males were paired with S14 males, the proportion of males performing tussling behavior was not significantly different from that between two G14 males. However, the duration of each tussling event in G14 and S14 male pairs was significantly shorter than that between two G14 males (Figure 6-figure supplement 2).

Next, we conducted further investigations into the role of social experience and the two types of fighting behaviors in mating competition (Figure 6D). Two males and one virgin female were loaded into the behavioral chamber at the same time and recorded for courtship behaviors. A winner was defined as the male that first copulated with the female. We used the copulation advance index, which was calculated as the proportional difference between marked males and unmarked males in winning copulation (see methods). We found that G7 males had no significant advantage over S7 males (Figure 6E); however, G14 males displayed a significantly increased mating advantage over S14 males (Figure 6F, see Figure 6-video supplement 2). Together, these results suggest that the tussling-favored GH males are more competitive in territory control and mating compared to the lunging-favored SH males, supporting a correlation between tussling and reproductive competetion.

The above results showed that GH males were more competitive compare to SH males, especially in older males, suggesting that social experience may play a potential role in altering age-related mating competition. To test this hypothesis, we first tested mating competition between males of different ages under the same rearing conditions. The results showed that younger males, whether reared in group or isolation, had a greater mating advantage compared to older males (G7 > G14, S7 > S14, G14 > G21, S14 > S21) (Figure 6G, Table S2). These results indicate that there is generally a disadvantage in mating competition associated with aging. We next investigated whether the mating advantage induced by social experience could compensate for the mating disadvantage caused by aging. We tested pairs of males with different ages and rearing conditions (G7 vs. S14, S7 vs. G14, G14 vs. S21, S14 vs. G21) for mating competition. We found that while 14-day-old males were less competitive than 7-day-old males regardless of the rearing condition, indicative of the aging-related disadvantage, 21-day-old GH males were significantly more competitive than 14-day-old SH males (G21 > S14) (Figure 6H).

To further investigate whether the territorial control and mating competition
advantage in GH males are related to their high tussling behavior, we paired males
with low tussling behavior (*Or47b-GAL4/UAS-Kir2.1* and *pC1^{SS2}/UAS-Kir2.1*) with
control males for territorial control and mating competition tests. We found that males
with inactivated Or47b or pC1^{SS2} neurons were incompetitive for both territorial
control and mating competition compared to control males (Figure 6-figure
supplement 3), further supporting a correlation between the occurrence of tussling
behavior and the acquisition of territorial and mating competition advantages.
Taken together, these results indicate that long-term social enrichment in older
males increases their territorial and mating competition, which can overcome their
mating disadvantage associated with aging (Figure 7).
339
340   **Discussion**
341
Regulation of physiology and behavior by social experience is an important feature
across animal species. Here, we found that social experience shapes the fighting
strategies of male flies, in which GH males tend to reduce the low-intensity lunging
and increase the high-intensity tussling. We further showed that the two forms of
fighting are mediated by distinct sensory and central neurons. The complex
implications of these findings are discussed below.
Previous studies have found that social isolation generally enhances aggression
but decreases mating competition in animal models (Palavicino-Maggio and Sengupta,
2022; Sethi et al., 2019; Yadav et al., 2024), presenting a paradox between social
experience, aggression and reproductive success. Our results provide an explanation
for this paradox by dissecting aggressive behaviors into the well-studied, frequently
observed lunging behavior and the less frequent but high-intensity tussling behavior.
We improved the behavioral paradigm for studying tussling behavior and revealed
that social experience oppositely regulates lunging and tussling. It is important to note
that the frequency of lunging decreases while the frequency of tussling increases from
7-day-old to 14-day-old males, suggesting a shift in fighting strategy with age. Our
results further suggest that social experience may enhance male mating competition
by shifting their fighting strategy to the low-frequency, high-intensity tussling, which
may be more effective in combat, although direct evidence is still lacking. These
results are consistent with a previous study showing a correlation between the
establishment of male dominance and tussling behavior, but not lunging (Simon and
Heberlein, 2020). Interestingly, we found that the mating advantage gained through
social enrichment can even offset the mating disadvantage associated with aging.
Note that a previous study found a mating advantage in 7-day-old males compared to
2-day-old males (Lin et al., 2016), which may be due to different levels of sexual
maturity, but not necessatily reflect the aging effect.
Our results revealed distinct neural circuits underlying the two forms of fighting,
spanning from primary sensory neurons to central regulatory neurons. Specifically,

silencing the cVA-sensing Or67d neurons reduced male lunging but not tussling,
whereas inactivating Or47b neurons impaired male tussling but not lunging. Notably,
Or47b neurons respond to pheromones common to both sexes, and the sensitivity of
these sensory neurons is enhanced by social enrichment (Lin et al., 2016; Peng et al.,
2021; Sethi et al., 2019). Unlike the Or67d-dependent lunging, which is primarily
male-induced and socially inhibited, the Or47b-dependent tussling is promoted by
social enrichment with either male or female flies. Recently, it has been found that
female flies increase aggression towards mating pairs or mated females, depending on
Or47b and Or67d, respectively (Gaspar et al., 2022; Wan et al., 2023). These results
suggest a common function of Or47b and Or67d neurons in both sexes for regulating
aggression. However, whether these neurons mediate different fighting patterns in
females and how they respond to social experience in females remain unknown.
According to the FlyWire connectome database (Dorkenwald et al., 2024; Schlegel et
al., 2024), connecting Or47b and Or67d ORNs to pC1 neurons in females requires at
least two intermediate neurons, suggesting a high degree of parallele processings from
ORNs to pC1s.

Regarding central regulatory neurons, we identified a novel class of pC1 neurons,
$pC1^{SS2}$, that specifically promote male tussling. In male flies, there are ~60 pairs of
dsx-expressing pC1 neurons, including the well-characterized P1[a] neurons that also
express fru^M (Asahina, 2018; Jiang and Pan, 2022); however, there are only three
pairs of $pC1^{SS2}$ neurons that specifically express *dsx* but not fru^M, and are crucial for
the high-intensity tussling in males. The three pairs of $pC1^{SS2}$ neurons may have
distinct function from P1[a] neurons in several ways. (1) P1[a] neurons promotes both
male courtship and aggression (Hoopfer et al., 2015; Zhao et al., 2024), whereas
$pC1^{SS2}$ neurons specifically promote aggression but not courtship. A recent study also
identified $pC1^{SS2}$ neurons as aggression-promoting, but their roles in different fighting
modes are not depicted (Hindmarsh Sten et al., 2025). That tussling-promoting $pC1^{SS2}$
neurons specifically express *dsx* but not fru^M is also consistent with a previous finding
that *dsx*-positive pC1 neurons, rather than fru^M-positive pC1 neurons, are crucial for
male aggression (Koganezawa et al., 2016). Moreover, the anatomical features of
$pC1^{SS2}$ neurons are highly similar to the male-specific aggression-promoting (MAP)
neurons identified previously (Chiu et al., 2021). (2) P1[a] neurons promotes both
lunging and tussling, although their function on tussling requires additional visual
stimulation. In contrast, $pC1^{SS2}$ neurons specifically promote tussling, and the
presence of light enhances but is not absolutely necessary for tussling. These results
align with previous findings that visual stimuli amplify P1 activation and related
behaviors (Hindmarsh Sten et al., 2021; Kohatsu and Yamamoto, 2015; Pan et al.,
2012), and visual stimulation is important for male aggression (Duistermars et al.,
2018; Ramin et al., 2014; Schretter et al., 2020). (3) Male lunging is induced after P1[a]
activation; whereas male tussling is induced during $pC1^{SS2}$ activation. These results
suggest that P1[a] neurons may not directly regulate aggression but instead induce a
persistent state that promote aggression in general, while $pC1^{SS2}$ neurons may play a

more direct role in mediating tussling. Future studies are needed to further dissect the
seemingly parallel sensory and central neuronal pathways involved in lunging and
tussling, and to explore how these pathways interact during social experience and
aging to optimize reproductive success.
416
Limitations of the Study
417
418
We manually analyzed aggressive behaviors and strived to ensure objectivity in
behavioral analysis; however, precisely tracking and quantifying specific occurrences
of lunging and tussling events for each fly over extended time periods remained
methodologically challenging. Furthermore, tussling events often exhibit prolonged
durations, which may also include other forms of aggression such as holding. While
our results indicate an association between tussling behavior and reproductive
competition, a direct link between them is lacking. Future investigations are needed to
establish a causal relationship between fighting strategies and reproductive success.
427
Materials and Methods
428
429
Fly Stocks
430
Flies were maintained at 22 or 25 °C in a 12 h:12 h light: dark cycle. *Canton-S* flies
were used as the wild-type strain and obtained from Janelia Research Campus (Pan
and Baker, 2014). *Or67d-GAL4* (BDSC_9998), *Or65a-GAL4* (BDSC_9993), *Or47b-*
GAL4 (BDSC_9984), *IR84a-GAL4* (BDSC_41734), *TK-GAL4* (BDSC_51795),
R19B03-GAL4 (BDSC_49830), *UAS-dTrpA1* (BDSC_26263), *UAS-Kir2.1*
(BDSC_6595), *UAS-CsChrimson* (BDSC_82181), *UAS-myrGFP* (BDSC_32198),
retro-Tango (BDSC_99661) and *trans-Tango* (BDSC_77124) flies were obtained
from Bloomington *Drosophila* Stock Center (BDSC). *UAS-NaChBac* (Ren et al.,
2001)*, fru^{GAL4}* (Stockinger et al., 2005)*, P1^a-spGAL4* (Zhang et al., 2018), *pC1^{SS1}-*
spGAL4 (Wang et al., 2021)*, pC1^{SS2}-spGAL4* (Wang et al., 2021), *UAS-dsx^M-RNAi*
(Peng et al., 2022) and *UAS-fruMi* (Meissner et al., 2016) were used as previously
described. *UAS-Or47b RNAi* (THU_2599) was obtained from TsingHua Fly Center.
Detailed genotypes of flies were listed in Table S1.
444
Male aggression assay
445
For the aggression assays presented in Figure 1, the prepared food was evenly spread
into the center of behavioral chambers. There was no female in the traditional
paradigm (Figure 1A, left) (Wu et al., 2020; Zhou et al., 2008), while a 3-5 day old
virgin female was gently fixed in the center of the food with its abdomen outside in
the modified paradigm (Figure 1A, right). We used a fixed female to restrict its
position in the center of food. Then two male testers were placed into the upper
chambers separated from the food (with or without a fixed female) by a film barrier.
All testers were loaded by cold anesthesia. After a 30-minute adaptation, the film was

gently removed to allow the two males to fall into the behavioral chamber, and the aggressive behavior was recorded for 2 hours. The aggression assays in Figure 1-figure supplement 1 was performed by using the paradigm without female, while other data were acquired using the modified paradigm with a fixed female unless otherwise stated. In some experiments, both single-housed and group-housed (30-40 males per vial) males were assayed for comparison.

We manually analyzed lunging and tussling behaviors in this study. Lunging is characterized by a male raising its forelegs and quickly striking the opponent, and each lunge typically lasts less than 0.2 seconds through detailed slow-motion analysis. Tussling is characterized by both males using their forelegs and bodies to tumble over each other, and this behavior may last from seconds to minutes. Tussling is often mixed with boxing, in which both flies rear up and strike the opponent with forelegs. Since boxing is often transient and difficult to distinguish from tussling, we referred to the mixed boxing and tussling behavior simply as tussling. As we manually analyze tussling for 2 hours for each pair of males, it is possible that we may miss some tussling events, especially those quick ones.

For the quantification of lunge latency and frequency in Figures 2E, 2F, 3C, 3D, and Figure 1- figures supplement 1A, 1B, we manually analyzed the videos and defined the lunge latency as the time point at which the first lunging event occurred between the two male flies after the start of the recording. We then counted the number of lunge events occurring within 10 minutes after the first lunge. For the quantification of tussle latency and frequency in Figures 1B, 1C, 1D, 2C, 2D, 3A, 3B, 3E, 3F, 3G, 3H, 4A and 4B, we manually analyzed the videos and defined the tussle latency as the time point at which the first tussling event occurred between the two male flies after the start of the recording. We defined the tussle frequency (No. of tussles in 2 hr) as the total number of tussling events that occurred within two hours from the beginning of the video.

The food recipes for the male aggression assay were listed below. A mixture of 2.5% yeast, 2.5% sucrose and 1% agar dissolved in apple juice (Huiyuan Apple juice concentrate) was used for Figure 1B; a mixture of 2.5% yeast, different concentrations of sucrose (0, 2.5%, 5% or 10%) and 1% agar dissolved in apple juice was used for Figure 1C; a mixture of different concentrations of yeast (0, 2.5%, 5% or 10%), 2.5% sucrose and 1% agar dissolved in apple juice was used for Figure 1D; A mixture of 10% yeast and 1% agar dissolved in apple juice was used for other experiments.

dTrpA1-mediated activation experiments

For thermogenetic activation assay showed in Figure 4-figure supplement 1, two 5-7 days old males were loaded into the circular chamber (d=15mm) under cold anesthesia and separated with a transparent film. Before the recording start, the chamber was transferred to the recording platform at a specific temperature (22°C or 30°C) for 30 minutes adaptation. After that, the film was removed, and the behavior

495 was recorded for 1 hour. The 10-minute video after film extraction was used for
496 behavioral analysis.
497
CsChrimson-mediated optogenetic activation experiments
For optogenetic activation assay showed in Figure 4 and 5, males were raised at 25°C
on standard fly food, collected after eclosion and housed in the dark for 5-7 days on
food containing 0.4 mM all *trans*-Retinal (MFCD00001550, Sigma-Aldrich, St. Louis,
MO). The specific light stimulation protocols are showed in the schematic (Figure 4F).
In brief, Two 5-7 days old males were loaded into the circular chamber (d=10mm)
under freeze anesthesia and separated with a transparent film in the dark. Then the
chamber was moved on the plane light source (IR channel: 865nm, red-light channel:
630nm) for 15 mins adaptation. Before video recording, the IR light channel was
switched on and adjusted to the appropriate intensty for recording. The film was
removed before the recording start, with the red-light channel switched on (constant
red light, 0.02 mW/mm^2) during the 2nd and 5th minutes of the experiment. The
statistics for the tussling, lunging, and unilateral wing extension (UWE) are as follows.
For tussling time and UWE time, we used LifesongY software (website:
https://sourceforge.net/projects/lifesongy/), which is written by Dr. Chao Guo based
on LifesongX (Bernstein et al., 1992), to calculate the percentage of the total time
these behaviors occurred per minute. Specifically, the "tussling time (%)" refers to the
percentage of time during the observation period in which tussling behavior occurred,
relative to the total observation time. Similarly, "UWE time (%)" represents the
percentage of time during the observation period in which UWE occurred. For lunges,
we counted the number of lunging occurrences per minute.
519
Territorial competition experiment
For the territorial competition experiment in Figures 6A-6C, Figure 6-figure
supplement 3A, the behavioral setup was produced using a 3D printer (Bambu Lab
A1), as illustrated in the schematic, with a transparent cover plate placed on the top.
The preparation of food and female were the same as the above aggression paradigm.
The female was fixed in the center of the food and two male testers were placed
separately in the two small chambers on both sides. Before initiating the experiment,
the two barriers were closed. After a 30-minute adaptation, the barriers were opened
to allow the males access to the central large chamber, and were closed once the
males went inside, and the territorial behavior was recorded for 2 hours. The first 10
encounters occurring after 1 hour of recording were used for statistical analysis. The
winning event was defined as a male driving the other male out of the food area. To
discriminate between tester males, about half of them were marked on one of their
wings using a red maker pen 24 hours before the experiment. For expreiments using
males with different ages and/or housing conditions, we calculated the winning index
as (num. of wins by males [upper panel] – num. of wins by males [lower panel])/10
encounters * 100%. For control expreiments using males of the same age and housing

condition, we calculated the winning index as (num. of wins by unmarked males –
num. of wins by marked males)/10 encounters * 100%, which is roughly zero due to
the randomness of marking.
540
Mating competition experiment
For the mating competition experiment shown in Figure 6D-6H, Figure 6-figure
supplement 3B, the wings of males were marked 24 hours prior to the experiment the
same as described above. During the experiment, a 3-5 days old virgin female was
placed into the chamber under freeze anesthesia and covered with a transparent film.
Then the unmarked males were loaded into the chamber in the same manner and also
covered with a transparent film. Finally, the marked males were loaded. After 30
minutes adaptation, two films were removed to allow the three flies to interact, and
the behavior was recorded for 1 hour. The chambers without successful copulation
during the 1-hour period were excluded from further analysis. For each condition, we
used 120 efficient chambers with 10 random chambers collectively as one sample. For
expreiments using males with different ages and/or housing conditions, the copulation
advance index was defined as (num. of male winners [upper panel] – num. of male
winners [lower panel])/10 chambers * 100%. For control expreiments using males of
the same age and housing condition, the copulation advance index was calculated as
(num. of winners by unmarked males – num. of winners by marked males)/10
chambers * 100%, which is roughly zero due to the randomness of marking. The
winner was defined as the male that first copulated with the female.
559
Quantitative Real-time PCR
Adult fly samples were frozen in liquid nitrogen and decapitated by vortex. The heads
were then separated from the bodies using metal sieves. Each sample, consisting of 40
frozen heads, was used for total RNA preparation using TRIzol reagent (15596026,
Thermo Fisher Scientific, Waltham, MA). We purified total RNA using a DNA-
freeTM Kit (AM1906, Thermo Fisher Scientific, Waltham, MA) and performed
reverse transcription using SuperScriptTM IV (18091050, Thermo Fisher Scientific,
Waltham, MA) to obtain cDNA used for templates. Quantitative PCR was performed
on the Roche LightCycler® 96 Real-Time PCR machine using AceQ qPCR SYBR
Green Master Mix (Q121-02, Vazyme, Nanjing). Transcript levels were analyzed by
the $2^{-\Delta CT}$ method using *actin* as control. Each sample was run in triplicate. Each
experiment was repeated three times using independent sets of genetic crosses.
Primers used for RT-qPCR quantification were:
actin forward: 5'-GTCGCGATTTAACCGACTACCTGA-3'
actin reverse: 5'-TCTTGCTT CGAGATCCACATCTGC-3'
fru-P1 forward: 5'-GTGTGCGTACGTTTGAGTGT-3'
fru-P1 reverse: 5'-TAATCCTGTGACGTCGCCAT-3'
Or47b forward: 5'-CAAATCTCAGCCTTCTGCGG-3'
Or47b reverse: 5'-GATACTGGCACAGCAAACTCA-3'

Tissue Dissection, Staining, and Imaging

Brains were dissected in Schneider's insect medium (S2) and fixed in 4% paraformaldehyde in 0.5% Triton X-100 and 0.5% bovine serum albumin in phosphate-buffered saline (PAT) for 30 min at room temperature. After 4 × 10-min washes, tissues were blocked in 3% normal goat serum (NGS) for 60 min, then incubated in primary antibodies diluted in 3% NGS for 4 h at room temperature and 1 day at 4°C, then washed in PAT, and incubated in secondary antibodies diluted in 3% NGS for 4 h at room temperature and 1 day at 4°C. The tissues were then washed thoroughly in PAT and mounted for imaging. Primary antibodies used in this study include mouse anti-HA (BioLegend, 901501, 1:200), rabbit anti-GFP (Invitrogen, A11122, 1:1000), monoclonal mouse anti-GFP (Sigma-Aldrich, G6539, 1:500), mouse anti-Bruchpilot (Developmental Studies Hybridoma Bank, nc82, 1:50), rabbit anti-Dsx[M] (1:500) and mouse anti-Fru[M] (1:2000). Secondary antibodies used include donkey anti-mouse IgG conjugated to Alexa 488 (Invitrogen, A21206, 1:500), donkey anti-mouse IgG conjugated to Alexa 555 (Invitrogen, A-31570, 1:500), donkey anti-rabbit IgG conjugated to Alexa 488 (Invitrogen, A-21206, 1:500) and donkey anti-rabbit IgG conjugated to Alexa 555 (Invitrogen, A-31572, 1:500). Samples were imaged on Zeiss 900 confocal microscopes using ZEN and processed with Fiji software.

Trans-synaptic labeling by *trans*-Tango and *retro*-Tango

The *trans*-Tango or *retro*-Tango ligand with GFP was expressed under the GAL4 driver. Trans-synaptic signals were marked by a membrane-associated form of tdTomato (QUAS-mtdTomato-3xHA). The anti-HA antibody (BioLegend, 901501, 1:200) was applied to characterize the postsynaptic or presynaptic signal. For Figure 3-figure supplement 2, male testers were kept at 18 °C until eclosion and subsequently single- or group-housed at the same temperature for two weeks before dissection and imaging. For Figure 4-figure supplement 2, flies were kept at 18 °C for *trans*-Tango and 25 °C for *retro*-Tango experiments.

Brain image registration

The protocol for standard brain registration is described previously (Zhou et al., 2014). Confocal images for brains containing P1[a] neurons and pC1[SS2] neurons were registered onto a standard reference brain (Zhou et al., 2014), using a Fiji graphical user interface as described previously (Ostrovsky et al., 2013).

Statistics

Experimental flies and genetic controls were tested under the same conditions, and data were collected from at least two independent experiments. Statistical analysis was performed using GraphPad Prism 9 (https://www.graphpad.com/scientific-software/prism/) as indicated in each figure legend. Data were first verified for normal

distribution by the D'Agostino–Pearson normality test. For normally-distributed data, Student's *t* test was used for pairwise comparisons, and one-way ANOVA was used for comparisons among multiple groups, followed by Tukey's multiple comparisons. For data not normally-distributed, the Mann-Whitney test was used for pairwise comparisons, and the Kruskal–Wallis test was used for comparison among multiple groups, followed by Dunn's multiple comparisons. For group matched data that were normally-distributed, the Paired *t* test signed rank test was used for pairwise comparisons, and RM one-way ANOVA was used for comparison among multiple groups, followed by RM one-way ANOVA multiple comparisons. For group matched data that were not normally-distributed, the Wilcoxon matched-pairs signed rank test was used for pairwise comparisons, and the Friedman test was used for comparison among multiple groups, followed by Dunn's multiple comparisons. For qPCR experiments, the average relative expression of three independent experiments was analyzed using Mann-Whitney test. For one group compared with zero, we used one sample *t* test. The Chi-square test was used to compare male tussling percentage between two groups.

Data availability

All data generated or analyzed during this study are included in the article and its supporting information files. Fly stocks and reagents used in this study are available from the corresponding author upon reasonable request.

Author contributions

C. Gao, Q. Peng and Y. Pan conceived the study; C. Gao and M. Ma performed all experiments and analyzed the data with help from J. Chen and X. Ji; Q. Peng and Y. Pan supervised the study; C. Gao and Y. Pan wrote the manuscript with input from all authors.

Competing interests

The authors declare no competing interests.

Acknowledgments

We thank the Bloomington *Drosophila* Stock Center, Tsinghua Fly Center, and Dr. Kaiyu Wang and Fei Wang for fly stocks. This work was supported by grants from National Key R&D Program of China (2021YFA1101300), the National Natural Science Foundation of China (32371067 to Y.P.), Shenzhen Medical Academy of Research and Translation (B2402005 to Y.P.), the Natural Science Foundation from Jiangsu Province (BK20231418 to Q.P.), and the Fundamental Research Funds for the Central Universities (2242023R40054 to Q.P.).

References

Agrawal, P., Kao, D., Chung, P. and Looger, L.L. (2020). The neuropeptide Drosulfakinin

regulates social isolation-induced aggression in Drosophila. Journal of Experimental Biology

223, https://doi.org/10.1242/jeb.207407.

Asahina, K. (2018). Sex differences in Drosophila behavior: qualitative and quantitative

dimorphism. Current Opinion in Physiology 6, 35-45.

https://doi.org/10.1016/j.cophys.2018.04.004.

Asahina, K., Watanabe, K., Duistermars, B.J., Hoopfer, E., González, C.R., Eyjólfsdóttir, E.A.,

Perona, P. and Anderson, D.J. (2014). Tachykinin-expressing neurons control male-specific

aggressive arousal in Drosophila. Cell 156, 221-235.

https://doi.org/10.1016/j.cell.2013.11.045.

Baines, R.A., Uhler, J.P., Thompson, A., Sweeney, S.T. and Bate, M. (2001). Altered

electrical properties in Drosophila neurons developing without synaptic transmission. The

Journal of Neuroscience 21, 1523-1531. https://doi.org/10.1523/JNEUROSCI.21-05-

01523.2001.

Bergman, M., Olofsson, M. and Wiklund, C. (2010). Contest outcome in a territorial butterfly:

the role of motivation. Proceedings of the Royal Society B: Biological Sciences 277, 3027-

3033. https://doi.org/10.1098/rspb.2010.0646.

Bernstein, A.S., Neumann, E.K. and Hall, J.C. (1992). Temporal analysis of tone pulses within

the courtship songs of two sibling Drosophila species, their interspecific hybrid, and

behavioral mutants of D. melanogaster (Diptera: Drosophilidae). Journal of Insect Behavior 5,

15-36. https://doi.org/10.1007/BF01049155.

Certel, S.J. and Kravitz, E.A. (2012). Scoring and analyzing aggression in Drosophila. Cold

Spring Harbor Protocols 2012, 319-325. https://doi.org/10.1101/pdb.prot068130.

Chen, J., Jin, S., Chen, D., Cao, J., Ji, X., Peng, Q. and Pan, Y. (2021). fruitless tunes

functional flexibility of courtship circuitry during development. eLife 10, e59224.

https://doi.org/10.7554/eLife.59224.

Chen, M., And Sokolowski, M. B. (2022). How Social Experience and Environment Impacts

Behavioural Plasticity in Drosophila. Fly 16, 68-84.

https://doi.org/10.1080/19336934.2021.1989248.

Chen, S., Lee, A.Y., Bowens, N.M., Huber, R. and Kravitz, E.A. (2002). Fighting fruit flies: a

model system for the study of aggression. Proceedings of the National Academy of Sciences

of the United States of America 99, 5664-5668. https://doi.org/10.1073/pnas.082102599.

Chiu, H., Hoopfer, E.D., Coughlan, M.L., Pavlou, H.J., Goodwin, S.F. and Anderson, D.J.

(2021). A circuit logic for sexually shared and dimorphic aggressive behaviors in Drosophila.

Cell 184, 507-520.e516. https://doi.org/10.1016/j.cell.2020.11.048.

Dankert, H., Wang, L., Hoopfer, E.D., Anderson, D.J. and Perona, P. (2009). Automated

monitoring and analysis of social behavior in Drosophila. Nature Methods 6, 297-303.

https://doi.org/10.1038/nmeth.1310.

Dorkenwald, S., Matsliah, A., Sterling, A.R., Schlegel, P., Yu, S.C., Mckellar, C.E., Lin, A.,

Costa, M., Eichler, K., Yin, Y., et al. (2024). Neuronal wiring diagram of an adult brain. Nature

634, 124-138. https://doi.org/10.1038/s41586-024-07558-y.

Duistermars, B.J., Pfeiffer, B.D., Hoopfer, E.D. and Anderson, D.J. (2018). A Brain Module for

Scalable Control of Complex, Multi-motor Threat Displays. Neuron 100, 1474-1490.e1474.

https://doi.org/10.1016/j.neuron.2018.10.027.

Dweck, H.K., Ebrahim, S.A., Thoma, M., Mohamed, A.A., Keesey, I.W., Trona, F., Lavista-

Llanos, S., Svatos, A., Sachse, S., Knaden, M., et al. (2015). Pheromones mediating

copulation and attraction in Drosophila. Proceedings of the National Academy of Sciences of

the United States of America 112, E2829-2835. https://doi.org/10.1073/pnas.1504527112.

Ferguson, C.J., Averill, P.M., Rhoades, H., Rocha, D., Gruber, N.P. and Gummattira, P.

(2005). Social isolation, impulsivity and depression as predictors of aggression in a

psychiatric inpatient population. Psychiatric Quarterly 76, 123-137.

https://doi.org/10.1007/s11089-005-2335-1.

Gaspar, M., Dias, S. and Vasconcelos, M.L. (2022). Mating pair drives aggressive behavior in

female Drosophila. Current Biology 32, 4734-4742 e4734.

https://doi.org/10.1016/j.cub.2022.09.009.

Grosjean, Y., Rytz, R., Farine, J.P., Abuin, L., Cortot, J., Jefferis, G.S. and Benton, R. (2011).

An olfactory receptor for food-derived odours promotes male courtship in Drosophila. Nature

478, 236-240. https://doi.org/10.1038/nature10428.

Hindmarsh Sten, T., Li, R., Otopalik, A. and Ruta, V. (2021). Sexual arousal gates visual

processing during Drosophila courtship. Nature 595, 549-553. https://doi.org/10.1038/s41586-

021-03714-w.

Hindmarsh Sten, T., Li, R., Hollunder, F., Eleazer, S. and Ruta, V. (2025). Male-male

interactions shape mate selection in Drosophila. Cell 0, S0092867425000376.

https://doi.org/10.1016/j.cell.2025.01.008.

Hoffmann, A.A. and Cacoyianni, Z. (1990). Territoriality in Drosophila melanogaster as a

conditional strategy. Animal Behaviour 40, 526-537. https://doi.org/10.1016/S0003-

3472(05)80533-0.

Hoopfer, E.D. (2016). Neural control of aggression in Drosophila. Current Opinion in

Neurobiology 38, 109-118. https://doi.org/10.1016/j.conb.2016.04.007.

Hoopfer, E.D., Jung, Y., Inagaki, H.K., Rubin, G.M. and Anderson, D.J. (2015). P1

interneurons promote a persistent internal state that enhances inter-male aggression in

Drosophila. eLife 4, e11346. https://doi.org/10.7554/eLife.11346.

Hueston, C.E., Olsen, D., Li, Q., Okuwa, S., Peng, B., Wu, J. and Volkan, P.C. (2016).

Chromatin Modulatory Proteins and Olfactory Receptor Signaling in the Refinement and

Maintenance of Fruitless Expression in Olfactory Receptor Neurons. PLoS Biology 14,

e1002443. https://doi.org/10.1371/journal.pbio.1002443.

Inagaki, H.K., Jung, Y., Hoopfer, E.D., Wong, A.M., Mishra, N., Lin, J.Y., Tsien, R.Y. and

Anderson, D.J. (2014). Optogenetic control of Drosophila using a red-shifted

channelrhodopsin reveals experience-dependent influences on courtship. Nat Methods 11,

325-332. https://doi.org/10.1038/nmeth.2765.

Jiang, X. and Pan, Y. (2022). Neural Control of Action Selection Among Innate Behaviors.

Neuroscience Bulletin 38, 1541-1558. https://doi.org/10.1007/s12264-022-00886-x.

Koganezawa, M., Kimura, K.-I. and Yamamoto, D. (2016). The Neural Circuitry that Functions

as a Switch for Courtship versus Aggression in Drosophila Males. Current Biology 26, 1395-

1403. https://doi.org/10.1016/j.cub.2016.04.017.

Kohatsu, S. and Yamamoto, D. (2015). Visually induced initiation of Drosophila innate

courtship-like following pursuit is mediated by central excitatory state. Nature

Communications 6, 6457. https://doi.org/10.1038/ncomms7457.

Kravitz, E.A. and Fernandez, M.P. (2015). Aggression in Drosophila. Behavioral

Neuroscience 129, 549-563. https://doi.org/10.1037/bne0000089.

Lim, R.S., Eyjólfsdóttir, E., Shin, E., Perona, P. and Anderson, D.J. (2014). How food controls

aggression in Drosophila. PLoS One 9, e105626.

https://doi.org/10.1371/journal.pone.0105626.

Lin, H.H., Cao, D.S., Sethi, S., Zeng, Z., Chin, J.S.R., Chakraborty, T.S., Shepherd, A.K.,

Nguyen, C.A., Yew, J.Y., Su, C.Y., et al. (2016). Hormonal Modulation of Pheromone

Detection Enhances Male Courtship Success. Neuron 90, 1272-1285.

https://doi.org/10.1016/j.neuron.2016.05.004.

Liu, W., Liang, X., Gong, J., Yang, Z., Zhang, Y.H., Zhang, J.X. and Rao, Y. (2011). Social

regulation of aggression by pheromonal activation of Or65a olfactory neurons in Drosophila.

Nature Neuroscience 14, 896-902. https://doi.org/10.1038/nn.2836.

Matsumoto, K., Pinna, G., Puia, G., Guidotti, A. and Costa, E. (2005). Social isolation stress-

induced aggression in mice: a model to study the pharmacology of neurosteroidogenesis.

Stress 8, 85-93. https://doi.org/10.1080/10253890500159022.

Meissner, G.W., Luo, S.D., Dias, B.G., Texada, M.J. and Baker, B.S. (2016). Sex-specific

regulation of Lgr3 in Drosophila neurons. Proceedings of the National Academy of Sciences

of the United States of America 113, E1256-1265. https://doi.org/10.1073/pnas.1600241113.

Nilsen, S.P., Chan, Y.-B., Huber, R. and Kravitz, E.A. (2004). Gender-selective patterns of

aggressive behavior in Drosophila melanogaster. Proceedings of the National Academy of

Sciences of the United States of America 101, 12342-12347.

https://doi.org/10.1073/pnas.0404693101.

Ostrovsky, A., Cachero, S. and Jefferis, G. (2013). Clonal analysis of olfaction in Drosophila:

image registration. Cold Spring Harbor Protocols 2013, 347-349.

https://doi.org/10.1101/pdb.prot071738.

Palavicino-Maggio, C.B. and Sengupta, S. (2022). The neuromodulatory basis of aggression:

Lessons from the humble fruit fly. Frontiers in Behavioral Neuroscience 16,

https://doi.org/10.3389/fnbeh.2022.836666.

Pan, Y. and Baker, B.S. (2014). Genetic identification and separation of innate and

experience-dependent courtship behaviors in Drosophila. Cell 156, 236-248.

https://doi.org/10.1016/j.cell.2013.11.041.

Pan, Y., Meissner, G.W. and Baker, B.S. (2012). Joint control of Drosophila male courtship

behavior by motion cues and activation of male-specific P1 neurons. Proceedings of the

National Academy of Sciences of the United States of America 109, 10065-10070.

https://doi.org/10.1073/pnas.1207107109.

Peng, Q., Chen, J. and Pan, Y. (2021). From fruitless to sex: On the generation and

diversification of an innate behavior. Genes Brain Behavior 20, e12772.

https://doi.org/10.1111/gbb.12772.

Peng, Q., Chen, J., Su, X., Wang, R., Han, C. and Pan, Y. (2022). The sex determination

gene doublesex is required during adulthood to maintain sexual orientation. Journal of

Genetics and Genomics 49, 165-168. https://doi.org/10.1016/j.jgg.2021.08.006.

Ramin, M., Domocos, C., Slawaska-Eng, D. and Rao, Y. (2014). Aggression and social

experience: genetic analysis of visual circuit activity in the control of aggressiveness in

Drosophila. Molecular Brain 7, 55. https://doi.org/10.1186/s13041-014-0055-0.

Ren, D., Navarro, B., Xu, H., Yue, L., Shi, Q. and Clapham, D.E. (2001). A prokaryotic

voltage-gated sodium channel. Science 294, 2372-2375.

https://doi.org/10.1126/science.1065635.

Sato, K., Goto, J. and Yamamoto, D. (2019). Sex Mysteries of the Fly Courtship Master

Regulator Fruitless. Frontiers in Behavioral Neuroscience 13, 245.

https://doi.org/10.3389/fnbeh.2019.00245.

Schlegel, P., Yin, Y., Bates, A.S., Dorkenwald, S., Eichler, K., Brooks, P., Han, D.S., Gkantia,

M., Dos Santos, M., Munnelly, E.J., et al. (2024). Whole-brain annotation and multi-

connectome cell typing of Drosophila. Nature 634, 139-152. https://doi.org/10.1038/s41586-

024-07686-5.

Schretter, C.E., Aso, Y., Robie, A.A., Dreher, M., Dolan, M.J., Chen, N., Ito, M., Yang, T.,

Parekh, R., Branson, K.M., et al. (2020). Cell types and neuronal circuitry underlying female

aggression in Drosophila. Elife 9, https://doi.org/10.7554/eLife.58942.

Sethi, S., Lin, H.H., Shepherd, A.K., Volkan, P.C., Su, C.Y. and Wang, J.W. (2019). Social

Context Enhances Hormonal Modulation of Pheromone Detection in Drosophila. Current

Biology 29, 3887-3898.e3884. https://doi.org/10.1016/j.cub.2019.09.045.

Simon, J.C. and Heberlein, U. (2020). Social hierarchy is established and maintained with

distinct acts of aggression in male Drosophila☐melanogaster. Journal of Experimental Biology

223, jeb232439. https://doi.org/10.1242/jeb.232439.

Sorkaç, A., Mo☐neanu, R.A., Crown, A.M., Savaş, D., Okoro, A.M., Memiş, E., Talay, M. and

Barnea, G. (2023). retro-Tango enables versatile retrograde circuit tracing in Drosophila. eLife

12, e85041. https://doi.org/10.7554/eLife.85041.

Stockinger, P., Kvitsiani, D., Rotkopf, S., Tirián, L. and Dickson, B.J. (2005). Neural circuitry

that   governs   Drosophila   male   courtship   behavior.   Cell   121,   795-807.

https://doi.org/10.1016/j.cell.2005.04.026.

Stockley, P. and Campbell, A. (2013). Female competition and aggression: interdisciplinary

perspectives. Philosophical Transactions of the Royal Society B: Biological Sciences 368,

20130073. https://doi.org/10.1098/rstb.2013.0073.

Talay, M., Richman, E.B., Snell, N.J., Hartmann, G.G., Fisher, J.D., Sorkaç, A., Santoyo, J.F.,

Chou-Freed, C., Nair, N., Johnson, M., et al. (2017). Transsynaptic Mapping of Second-Order

Taste   Neurons   in   Flies   by   trans-Tango.   Neuron   96,   783-795.e784.

https://doi.org/10.1016/j.neuron.2017.10.011.

Wan, X., Shen, P., Shi, K., Li, J., Wu, F. and Zhou, C. (2023). A Neural Circuit Controlling

Virgin Female Aggression Induced by Mating-related Cues in Drosophila. Neuroscience

Bulletin 39, 1396-1410. https://doi.org/10.1007/s12264-023-01050-9.

Wang, K., Wang, F., Forknall, N., Yang, T., Patrick, C., Parekh, R. and Dickson, B.J. (2021).

Neural circuit mechanisms of sexual receptivity in Drosophila females. Nature 589, 577-581.

https://doi.org/10.1038/s41586-020-2972-7.

Wang, L. and Anderson, D.J. (2010). Identification of an aggression-promoting pheromone

and   its   receptor   neurons   in   Drosophila.   Nature   463,   227-231.

https://doi.org/10.1038/nature08678.

Wang, L., Dankert, H., Perona, P. and Anderson, D.J. (2008). A common genetic target for

environmental and heritable influences on aggressiveness in Drosophila. Proceedings of the

National   Academy   of   Sciences   of   the   United   States   of   America   105,   5657-5663.

https://doi.org/10.1073/pnas.0801327105.

Wu, F., Deng, B., Xiao, N., Wang, T., Li, Y., Wang, R., Shi, K., Luo, D.-G., Rao, Y. and Zhou,

C. (2020). A neuropeptide regulates fighting behavior in Drosophila melanogaster. eLife 9,

e54229. https://doi.org/10.7554/eLife.54229.

Yadav, R.S.P., Ansari, F., Bera, N., Kent, C. and Agrawal, P. (2024). Lessons from lonely flies:

Molecular   and   neuronal   mechanisms   underlying   social   isolation.   Neuroscience   and

Biobehavioral Reviews 156, 105504. https://doi.org/10.1016/j.neubiorev.2023.105504.

Yamamoto, D. and Koganezawa, M. (2013). Genes and circuits of courtship behaviour in

Drosophila   males.   Nature   Reviews   Neuroscience   14,   681-692.

https://doi.org/10.1038/nrn3567.

Zelikowsky, M., Hui, M., Karigo, T., Choe, A., Yang, B., Blanco, M.R., Beadle, K., Gradinaru,

V., Deverman, B.E. and Anderson, D.J. (2018). The neuropeptide Tac2 controls a distributed

brain   state   induced   by   chronic   social   isolation   stress.   Cell   173,   1265-1279.

https://doi.org/10.1016/j.cell.2018.03.037.

Zhang, W., Guo, C., Chen, D., Peng, Q. and Pan, Y. (2018). Hierarchical Control of

Drosophila Sleep, Courtship, and Feeding Behaviors by Male-Specific P1 Neurons.

Neuroscience Bulletin 34, 1105-1110. https://doi.org/10.1007/s12264-018-0281-z.

Zhao, H., Jiang, X., Ma, M., Xing, L., Ji, X. and Pan, Y. (2024). A neural pathway for social

modulation of spontaneous locomotor activity (SoMo-SLA) in Drosophila. Proceedings of the

National Academy of Sciences of the United States of America 121, e2314393121.

https://doi.org/10.1073/pnas.2314393121.

Zhao, S., Deanhardt, B., Barlow, G.T., Schleske, P.G., Rossi, A.M. and Volkan, P.C. (2020).

Chromatin-based reprogramming of a courtship regulator by concurrent pheromone

perception and hormone signaling. Science Advances 6, eaba6913.

https://doi.org/10.1126/sciadv.aba6913.

Zhou, C., Rao, Y. and Rao, Y. (2008). A subset of octopaminergic neurons are important for

Drosophila aggression. Nature Neuroscience 11, 1059-1067. https://doi.org/10.1038/nn.2164.

Zhou, C., Pan, Y., Robinett, C.C., Meissner, G.W. and Baker, B.S. (2014). Central brain

neurons expressing doublesex regulate female receptivity in Drosophila. Neuron 83, 149-163.

https://doi.org/10.1016/j.neuron.2014.05.038.

Zwarts, L., Versteven, M. and Callaerts, P. (2012). Genetics and neurobiology of aggression

in Drosophila. Fly 6, 35-48. https://doi.org/10.4161/fly.19249.

871
872

Figures and figure legends

Figure 1. A paradigm for tussling behavior in male *Drosophila*.

(A) Diagrams of two fighting paradigms for *Drosophila* tussling behavior. The left diagram shows a traditional fighting paradigm with no female (NF) in the food area and a modified fighting paradigm with a fixed female (FF) in the center of the food area. Before the experiment began, two male flies were placed on the upper layer, separated from the food by a film. After a 30-minute adaptation, the film was removed to start recording. The right diagram shows a pair of tussling male *Drosophila*. All tester males in this figure were group housed.

(B) The influence of age and female factors on male tussling behavior. From top to
bottom, the tussle latency, the number of tussles, and the proportion of tussles
occurring are shown respectively. n=30, 30, 30, 46, 46, 46 from left to right. Bars
sharing the same letter are not significantly different according to Kruskal-Wallis test
(tussle latency, number of tussles) and Chi-square test (proportion of tussles
occurring).
(C) Sucrose concentration of food have no effect on male tussling behavior. n=30 for
each group, n.s., not significant, Kruskal-Wallis test (tussle latency, number of
tussles), and Chi-square test (proportion of tussles occurring).
(D) The influence of yeast concentration on male tussling behavior. n=30 for each
group, n.s., not significant, *P < 0.05, ***P < 0.001, ****P < 0.0001, Kruskal-Wallis
test (tussle latency, number of tussles), and Chi-square test (proportion of tussles
occurring). Error bars correspond to SEM.
897

898
Figure 1-figure supplement 1. Aging suppresses lunging behavior in male
***Drosophila*.**
(A) The lunge latency in 7-day-old single-housed (S7), 14-day-old single-housed
(S14), 7-day-old group-housed (G7) and 14-day-old group-housed (G14) males, n=12
for each group, n.s., not significant, *P < 0.05, Mann-Whitney test.
(B) The number of lunges in S7, S14, G7 and G14 males within 10 minutes, n=12 for
each group, *P < 0.05, ****P < 0.0001, Mann-Whitney test. Error bars correspond to
SEM.
907
908　**Figure 1-video supplement 1.** Tussling behavior between two 14-day-old group-
909　housed (G14) wild-type males.
910

911
Figure 2. Social enrichment inhibits lunging behavior but promotes tussling
behavior in males.
(A) Schematic diagram of *Drosophila* rearing conditions and experimental procedure.
(B) Representative raster plots illustrating lunge events in the first 10 minutes and
tussle events over 2 hours in single-housed (SH) and group-housed (GH) males.
(C, D) Latency (C) and number (D) of tussling behavior in SH and GH males, n=27,
30 for SH and GH males, respectively, ****P < 0.0001, Mann-Whitney test for both.
(E, F) Latency (E) and number (F) of lunging behavior in SH and GH males, n=27, 30
for SH and GH males, respectively, ****P < 0.0001, Mann-Whitney test for both.
Error bars correspond to SEM.

926
Figure 3. Or47b neurons are required for male tussling but not lunging.

(A, B) The effect of inactivating different sensory neurons on latency (A) and number
(B) of tussling behavior in 14-day-old group-housed males (G14). n=30 for each
group, n.s., not significant, **P < 0.01, Kruskal-Wallis test.
(C, D) The effect of inactivating different sensory neurons on the latency (C) and
number (D) of lunging behavior in G14 males. n=30 for each group, n.s., not
significant, **P < 0.01, ***P < 0.001, Kruskal-Wallis test.
(E, F) The effect of knocking down *Or47b* or *fru^M* expression in *Or47b* ORNs on the
latency (E) and number (F) of tussling behavior in G14 males, respectively. n=30 for
each group, n.s., not significant, ****P < 0.0001, Kruskal-Wallis test.
(G, H) The effect of activating *Or47b* ORNs on latency (G) and number (H) of
tussling behavior under conditions of GH or SH for 14 days, n=60, 60, 60, 49
respectively, n.s., not significant, ****P < 0.0001, Kruskal-Wallis test. Error bars
correspond to SEM.
941

943 **Figure 3-figure supplement 1. Validation of the *Or47b* and *fru^M* RNAi efficiency**.
944 Validation of *UAS-Or47b-RNAi* (A) and *UAS-fruMi* (B) efficiency by quantitative
945 qPCR. n=9 for each of three replicates, ****P < 0.0001, ***P < 0.001, Mann-
946 Whitney test. Error bars correspond to SEM.
947

Figure 3-figure supplement 2. Group housing increases downstream signals of
***Or47b* ORNs.**
(A) Projection of *Or47b* ORNs (green) and their downstream neurons (red) in GH and
SH conditions. Scale bar, 100μm.
(B) The fluorescence intensity of neurons downstream of *Or47b* ORNs is stronger
under GH conditions compared to SH conditions. Both groups of male testers were 14
days old, n=10 and 8, respectively, ****$P < 0.0001$, Mann-Whitney test. Error bars
correspond to SEM.
957

Figure 4. A subset of pC1 neurons specifically promote male tussling behavior.

(A, B) The effect of inactivating candidate interneurons on latency (A) and number (B) of male tussling behavior, n=30 for each group, bars sharing the same letter are not significantly different according to Kruskal-Wallis test.

(C-E) The expression pattern of *P1ᵃ-spGAL4* (C) and *pC1^{SS2}-spGAL4* (D) in male
brain, and (E) the registration of P1ᵃ (magenta) and pC1^{SS2} (green) neurons in a
standard brain. The GFP expression indicates P1ᵃ (magenta) and pC1^{SS2} (green)
neurons in the male brain counterstained with nc82 (blue). White dashed boxes in (C-
E) were zoomed in and shown below each panel. Scale bars, 100μm.

(F-H) Statistics on tussling (F), lunging (G) and unilateral wing extension (UWE, H)
by *UAS-CsChrimson/+* males. Red light activation with a light intensity of 0.02
mW/mm^2 for 1 minute was applied at the 2nd and 5th minutes of this and subsequent
experiments. n=32 for each group, n.s., not significant, Friedman test.

(I-K) Statistics on tussling (I), lunging (J) and UWE (K) in *P1ᵃ>UAS-CsChrimson*
males. n=32 for each group, n.s., not significant, ****P < 0.0001, Friedman test.

(L-N) Statistics on tussling (L), lunging (M) and UWE (N) in *pC1^{SS2}>UAS-*
CsChrimson males. n=32 for each group, n.s., not significant, ***P < 0.001, ****P <
0.0001, Friedman test.

Figure 4-figure supplement 1. light-dependent initiation of male tussling
behavior.

(A) Schematic diagram of the thermal activation experiment.

(B) The percentage of tussling occurrence (left) and total tussling time (right) within
10 mins. All tester males were 5 to 7-day-old group-housed males. n=18 for each
group, bars sharing the same letter are not significantly different according to
Kruskal-Wallis test based on the data of total tussling time.

987
988 **Figure 4-figure supplement 2.** Immunostaining images of potential upstream signals
989 (*retro*-Tango) and downstream signals (*trans*-Tango) of P1^a and pC1^{SS2} neurons.
990 Scale bars, 100μm.

991
992 **Figure 4-video supplement 1.** Optogenetic activation of pC1^{SS2} neurons acutely
993 induces tussling in 5 to 7-day-old group-housed males (*pC1^{SS2}>UAS-CsChrimson*).

Figure 5. Dsx^M regulates the development and tussling-promoting function of pC1^{SS2} neurons.

(A, B) The co-staining of P1^a or pC1^{SS2} neurons with the Fru^M antibody (A), and the quantification of *fru*-positive neurons in P1^a and pC1^{SS2} populations (B). n=5 for each group. Scale bars, 50μm.

(C, D) The co-staining of P1^a or pC1^{SS2} neurons with the Dsx^M antibody, and The quantification of *dsx*-positive neurons in P1^a and pC1^{SS2} populations (D), n=9 for each group. Scale bars, 50μm.

(E-K) Knocking down *dsx* in pC1^{SS2} neurons impact their development and tussling-promoting function. (E, F) Immunostaining of pC1^{SS2} neurons in control (E) and *dsx* knock-down (F) males. Magenta represents nc82 signals, and green represents GFP signals. The images within the yellow dashed box and white dashed box are magnified 2× and 4×, respectively, and displayed on the right. Scale bars, 100μm. (G) Quantification of the number of pC1^{SS2} neurons in control and *dsx* knock-down males.

1017

1018

Figure 6. Social enrichment enhances male territorial control and mating competition.

(A) Schematic diagram of the paradigm for testing territorial control in males.

(B) The territorial control test of 7-day-old GH (G7) and SH (S7) males. G7a and G7b stand for marked G7 male and unmarked G7 male, respectively. n=12 for each group, n.s., not significant, one sample t test.

(C) The territorial control test of 14-day-old GH (G14) and SH (S14) males. n=12 for
each group, n.s., not significant, *P < 0.05, one sample *t* test.
(D) Schematic diagram of the paradigm for testing male mating advantage.
(E) The mating competition test between G7 and S7 males. n=12 for each group, n.s.,
not significant, one sample *t* test.
(F) The mating competition test between G14 and S14 males. n=12 for each group,
n.s., not significant, *P < 0.05, one sample *t* test.
(G and H) The mating competition test between young males and old males, n=12 for
each group, n.s., not significant, *P < 0.05, ****P < 0.0001, one sample *t* test for
comparison within group (red labeling), Mann-Whitney test for comparison between
groups (black labeling). Error bars correspond to SEM.
1036

1037

1038 **Figure 6-figure supplement 1. Winning events of representative samples for**
1039 **territorial competition.**
1040 The blue and red bars represent the wins for S14 males and G14 males, respectively.
1041 An example formula is provided for calculating the winning index in the dashed box.
1042

1043

Figure 6-figure supplement 2. The occurrence and duration of tussling behavior in male flies under different pairing conditions.

(A) Proportion of male pairs performing tussling within 2 hours under three different pairing conditions in the territorial control paradigm. n=12 for each group, bars sharing the same letter are not significantly different according to Chi-square test.

(B) Duration of each tussling event in males under the G14-G14 and G14-S14 pairing conditions. n=15 events from 8-12 samples for each group, ****$P < 0.0001$, Mann-Whitney test. Error bars correspond to SEM.

1052

(a indicates unmarked, b indicates marked)

1053

Figure 6-figure supplement 3. Silencing Or47b or pC1^{SS2} neurons compromises male territorial control and mating competition.

(A, B) Effects of Or47b neurons inactivation and pC1^{SS2} neurons inactivation on territorial control (A) and mating competition (B) in male flies. n=12 for each group, n.s., not significant, ***$P < 0.001$, ****$P < 0.0001$, one sample *t* test.

Figure 6-video supplement 1. In a territory competition paradigm, a 14-day-old group-housed (G14, marked) wild-type male consistently outcompetes a 14-day-old single-housed (S14, unmarked) wild-type male.

Figure 6-video supplement 2. In a mating competition paradigm, a 14-day-old group-housed (G14, unmarked) wild-type male outperformed a 14-day-old single-housed (S14, marked) wild-type male and successfully copulated with the female.

1067

1068
Figure 7. Summary model of fighting strategies and reproductive success in male
Drosophila.

Age and social enrichment inhibit the low-intensity, high-frequency lunging while promoting the low-frequency, high-intensity tussling. This shift in fighting strategies among experienced, aged males is accompanied with better territorial control and mating competition, even offsetting aging-related mating disadvantages. Lunging and tussling behaviors are regulated by distinct sensory and central neurons.

1076

1077   **Table S1. Detailed information for fly stocks.**

1078

Figures	Genotypes	Housing conditions	Sample sizes
Figure 1			
B	*Canton-S*	GH-2d, GH-7d, GH-14d	n=30, 30, 30, 46, 46, 46
C	*Canton-S*	GH-14d	n=30 for each group
D	*Canton-S*	GH-14d	n=30 for each group
Figure 2			
B	*Canton-S*	SH-14d, GH-14d	
C, D, E, F	*Canton-S*	SH-14d, GH-14d	n=27 for SH, n=30 for GH
G	*Canton-S*	SH-14d, GH-14d	n=30 events from 10 samples for each group
Figure 3			
A, B, C, D	*w+;;UAS-Kir2.1/+*	GH-14d	30
	w+;;Or67d-GAL4/UAS-Kir2.1	GH-14d	30
	w+;Or65a-GAL4/+;UAS-Kir2.1/+	GH-14d	30
	w+;;Or47b-GAL4/UAS-Kir2.1	GH-14d	30
	w+;;Ir84a-GAL4/UAS-Kir2.1	GH-14d	30
E, F	*w+;;Or47b-GAL4/+*	GH-14d	30
	w+;;Or47b-GAL4/UAS-Or47b-RNAi	GH-14d	30
	w+;;Or47b-GAL4/UAS-fruMi	GH-14d	30
G, H	*w+;;Or47b-GAL4/+*	GH-14d	60
	w+;;Or47b-GAL4/UAS-NaChBac	GH-14d	60
	w+;;Or47b-GAL4/+	SH-14d	60
	w+;;Or47b-GAL4/UAS-NaChBac	SH-14d	49
Figure 4			
A, B	*w+;;UAS-Kir2.1/+*	GH-14d	30
	w+;;R19B03-GAL4/UAS-Kir2.1	GH-14d	30
	w+;R15A01-AD/+;R71G01-DBD/UAS-Kir2.1 ($P1^a$ > UAS-Kir2.1)	GH-14d	30
	w+;;TK-GAL4/UAS-Kir2.1 (GH-14d)	GH-14d	30
	w+;VT002064-AD/+;VT008469DBD/UAS-Kir2.1 ($pC1^{SS1}$ > UAS-Kir2.1)	GH-14d	30
	w+;;VT002064-AD,dsxDBD/UAS-Kir2.1 ($pC1^{SS2}$ > UAS-Kir2.1)	GH-14d	30
C	*w-;R15A01-AD/UAS-myrGFP; R71G01-DBD/+ ($P1^a$ > UAS-GFP)*	GH-5-7d	

D	*w-;UAS-myrGFP/+;VT002064-AD,dsxDBD/+ (pC1^{SS2} > UAS-GFP)*	GH-5-7d	
F, G, H	*w-;;UAS-CsChrimson/+*	GH-5-7d	32
I, J, K	*w-;R15A01-AD/+;R71G01-DBD/ UAS-CsChrimson (P1^a > UAS-CsChrimson)*	GH-5-7d	32
L, M, N	*w-;;VT002064-AD,dsxDBD/UAS-CsChrimson (pC1^{SS2} > UAS-CsChrimson)*	GH-5-7d	32
Figure 5			
A, B, C, D	*w-;R15A01-AD/UAS-myrGFP; R71G01-DBD/+ (P1^a > UAS-GFP)*	GH-5-7d	n=5 for each group (B)
	w-;UAS-myrGFP/+;VT002064-p65AD,dsxDBD/+ (pC1^{SS2} > UAS-GFP)	GH-5-7d	n=9 for each group (D)
E	*w-;UAS-CsChrimson/+;VT002064-AD, dsxDBD/+ (pC1^{SS2} > UAS-CsChrimson)*	GH-5-7d	
F	*w-;UAS-CsChrimson/+; VT002064-AD, dsxDBD/UAS-dsx^M-RNAi (pC1^{SS2} > UAS-CsChrimson & UAS-dsx^M-RNAi)*	GH-5-7d	
G	*w-;UAS-CsChrimson/+;VT002064-AD, dsxDBD/+ (pC1^{SS2} > UAS-CsChrimson)*	GH-5-7d	10
	w-;UAS-CsChrimson/+;VT002064-AD, dsxDBD/UAS-dsx^M-RNAi (pC1^{SS2} > UAS-CsChrimson & UAS-dsx^M-RNAi)	GH-5-7d	9
H	*w-;UAS-CsChrimson/+;VT002064-AD, dsxDBD/+ (pC1^{SS2} > UAS-CsChrimson)*	GH-5-7d	10
	w-;UAS-CSChrimson/+;VT002064-AD, dsxDBD/UAS-dsx^M-RNAi (pC1^{SS2} > UAS-CsChrimson & UAS-dsx^M-RNAi)	GH-5-7d	10
I	*w-;UAS-CsChrimson/+;VT002064-AD, dsxDBD/+ (pC1^{SS2} > UAS-CsChrimson)*	GH-5-7d	12
J	*w-;UAS-CsChrimson/+;VT002064-AD, dsxDBD/UAS-dsx^M-RNAi (pC1^{SS2} > UAS-CsChrimson & UAS-dsx^M-RNAi)*	GH-5-7d	12
K	*w-;UAS-CsChrimson/+;VT002064-AD, dsxDBD/+ (pC1^{SS2} > UAS-CsChrimson)*	GH-5-7d	12
	w-;UAS-CsChrimson/+;VT002064-AD, dsxDBD/UAS-dsx^M-RNAi (pC1^{SS2} > UAS-CsChrimson & UAS-	GH-5-7d	12

	dsx^M-RNAi)		
	w-;UAS-CsChrimson/+;VT002064-AD, dsxDBD/+ (pC1^{SS2} > UAS-CsChrimson)	GH-5-7d	12
	w-;UAS-CsChrimson/+;VT002064-AD, dsxDBD/UAS-dsx^M-RNAi (pC1^{SS2} > UAS-CsChrimson & UAS-dsx^M-RNAi)	GH-5-7d	12
Figure 6			
B	*Canton-S*	GH-7d	12
	Canton-S	SH-7d	12
	Canton-S	GH-7d, SH-7d	12
C	*Canton-S*	GH-14d	12
	Canton-S	SH-14d	12
	Canton-S	GH-14d, SH-14d	12
E	*Canton-S*	GH-7d	12
	Canton-S	SH-7d	12
	Canton-S	GH-7d, SH-7d	12
F	*Canton-S*	GH-14d	12
	Canton-S	SH-14d	12
	Canton-S	GH-14d, SH-14d	12
G	*Canton-S*	GH-7d, GH-14d	12
	Canton-S	SH-7d, SH-14d	12
	Canton-S	GH-14d, GH-21d	12
	Canton-S	SH-14d, SH-21d	12
H	*Canton-S*	GH-7d, SH-14d	12
	Canton-S	SH-7d, GH-14d	12
	Canton-S	GH-14d, SH-21d	12
	Canton-S	SH-14d, GH-21d	12
Figure 1-figure supplement 1			
A, B	*Canton-S*	SH-7d	12
	Canton-S	SH-14d	12
	Canton-S	GH-7d	12
	Canton-S	GH-14d	12
Figure 3-figure supplement 1			
A	*w+;;Or47b-GAL4/+*	GH-5-7d	9
	w+;;Or47b-GAL4/UAS-Or47b-RNAi	GH-5-7d	9

B	$w+;;fru^{GAL4}/+$	GH-5-7d	9
	$w+;;fru^{GAL4}/UAS$-$fruMi$	GH-5-7d	9
Figure 3-figure supplement 2			
A	*UAS-myrGFP,QUAS-mtdtomato*(*3xHA*)*;trans-Tango/+;Or47b-GAL4/+*	GH-14d, 18°C	
	UAS-myrGFP,QUAS-mtdtomato(*3xHA*)*;trans-Tango/+;Or47b-GAL4/+*	SH-14d, 18°C	
B	*UAS-myrGFP,QUAS-mtdtomato*(*3xHA*)*;trans-Tango/+;Or47b-GAL4/+*	GH-14d, 18°C	10
	UAS-myrGFP,QUAS-mtdtomato(*3xHA*)*;trans-Tango/+;Or47b-GAL4/+*	SH-14d, 18°C	8
Figure 4-figure supplement 1			
B	*w+;UAS-dTrpA1/+* (light, 22°C)	GH-5-7d	18
	w+;R15A01-AD/UAS-dTrpA1; R71G01-DBD/+ ($P1^a > UAS$-$dTrpA1$) (light, 22°C)	GH-5-7d	18
	w-;UAS-dTrpA1/+; VT002064-AD, dsxDBD/+ ($pC1^{SS2} > UAS$-$dTrpA1$) (light, 22°C)	GH-5-7d	18
	w+;UAS-dTrpA1/+ (dark, 22°C)	GH-5-7d	18
	w+;R15A01-AD/UAS-dTrpA1;R71G01-DBD/+ ($P1^a > UAS$-$dTrpA1$) (dark, 22°C)	GH-5-7d	18
	w-;UAS-dTrpA1/+;VT002064-AD, dsxDBD/+ ($pC1^{SS2} > UAS$-$dTrpA1$) (dark, 22°C)	GH-5-7d	18
	w+;UAS-dTrpA1/+ (light, 30°C)	GH-5-7d	18
	w+;R15A01-AD/UAS-dTrpA1;R71G01-DBD/+ ($P1^a > UAS$-$dTrpA1$) (light, 30°C)	GH-5-7d	18
	w-;UAS-dTrpA1/+;VT002064-AD, dsxDBD/+ ($pC1^{SS2} > UAS$-$dTrpA1$) (light, 30°C)	GH-5-7d	18
	w+;UAS-dTrpA1/+ (dark, 30°C)	GH-5-7d	18
	w+;R15A01-AD/UAS-dTrpA1;R71G01-DBD/+ ($P1^a > UAS$-$dTrpA1$) (dark, 30°C)	GH-5-7d	18
	w-;UAS-dTrpA1/+;VT002064-AD, dsxDBD/+ ($pC1^{SS2} > UAS$-$dTrpA1$) (dark, 30°C)	GH-5-7d	18
Figure 4-figure supplement 2			
retro- Tango	*QUAS-mtdTomato*(*3xHA*)*;retro-Tango/R15A01-AD;UAS-EGFP/R71G01-DBD* ($P1^a > retro$-$Tango$)	GH-14-21d	

	QUAS-mtdTomato(3xHA);retro-Tango/+;UAS-EGFP/VT002064AD,dsxDBD ($pC1^{SS2} > retro\text{-}Tango$)	GH-14-21d	
GFP	*w-;R15A01-AD/+;R71G01-DBD/UAS-myrGFP* ($P1^a > GFP$)	GH-5-7d	
	w-;;VT002064AD,dsxDBD/UAS-myrGFP ($pC1^{SS2} > GFP$)	GH-5-7d	
*trans-*Tango	*UAS-myrGFP,QUAS-mtdtomato(3xHA);trans-Tango/R15A01-AD;R71G01-DBD/+* ($P1^a > trans\text{-}Tango$)	GH-14-21d	
	UAS-myrGFP,QUAS-mtdtomato(3xHA);trans-Tango/+;VT002064AD,dsxDBD/+ ($pC1^{SS2} > trans\text{-}Tango$)	GH-14-21d	
Figure 6-figure supplement 1			
	Canton-S	GH-14d, SH-14d	12
Figure 6-figure supplement 2			
A	*Canton-S*	GH-14d	12
	Canton-S	SH-14d	12
	Canton-S	GH-14d, SH-14d	12
B	*Canton-S*	GH-14d	n=15 events from 8-12 samples for each group
	Canton-S	GH-14d, SH-14d	n=15 events from 8-12 samples for each group
Figure 6-figure supplement 3			
A	*w+;;UAS-Kir2.1/+*	GH-14d	12
	w+;;Or47b-GAL4/UAS-Kir2.1	GH-14d	12
	w+;;Or47b-GAL4/UAS-Kir2.1 w+;;UAS-Kir2.1/+	GH-14d	12
	w+;;VT002064AD,dsxDBD/ UAS-Kir2.1($pC1^{SS2} > UAS\text{-}Kir2.1$)	GH-14d	12
	w+;;VT002064AD, dsxDBD/ UAS-Kir2.1($pC1^{SS2} > UAS\text{-}Kir2.1$) *w+;;UAS-Kir2.1/+*	GH-14d	12
B	*w+;;UAS-Kir2.1/+*	GH-14d	12
	w+;;Or47b-GAL4/UAS-Kir2.1	GH-14d	12
	w+;;Or47b-GAL4/UAS-Kir2.1 w+;;UAS-Kir2.1/+	GH-14d	12
	w+;;VT002064AD,dsxDBD/ UAS-Kir2.1($pC1^{SS2} > UAS\text{-}Kir2.1$)	GH-14d	12
	w+;;VT002064AD, dsxDBD/ UAS-	GH-14d	12

	Kir2.1(pC1^{SS2} > UAS-Kir2.1) *w+;;UAS-Kir2.1/+*			

1079

1080

1081   **Table S2. Detailed comparison for territorial control and mating competition.**

	pairs	winning index	P value	conclusion
Territorial control assay				
Fig. 6B	G7a vs G7b	-10±30.15	0.275	/
	S7a vs S7b	8.33±43.87	0.524	/
	G7 vs S7	21.66±50.06	0.1619	/
Fig. 6C	G14a vs G14b	8.33±49.33	0.5702	/
	S14a vs S14b	-3.33±59.60	0.8499	/
	G14 vsS14	48.33±65.76	0.0272	G14>S14

	pairs	copulation advance index	P value	conclusion
Mating competition assay				
Fig. 6E	G7a vs G7b	3.33±22.29	0.6147	/
	S7a vs S7b	3.33±14.35	0.4382	/
	G7 vs S7	-5±17.32	0.3388	/
Fig. 6F	G14a vs G14b	0±19.07	>0.9999	/
	S14a vs S14b	0±17.06	>0.9999	/
	G14 vs S14	26.67±32.29	0.0155	G14>S14
Fig. 6G	G7 vs G14	46.67±24.62	<0.0001	G7>G14
	S7 vs S14	43.33±25.35	<0.0001	S7>S14
	G14 vs G21	58.33±27.58	<0.0001	G14>G21
	S14 vs S21	51.67±24.80	<0.0001	S14>S21
Fig. 6H	G7 vs S14	45±24.31	<0.0001	G7>S14
	S7 vs G14	18.33±21.67	0.0137	S7>G14
	G14 vs S21	41.67±21.67	<0.0001	G14>S21
	S14 vs G21	-25±28.44	0.0112	S14<G21
		S21<S14<G21<G14		