Title: Experimental evolution of *Drosophila* without its microbiome

Authors: Akshay Malwade[1], Anish Koner[1], B. Vibishan[1,2], Chinmay Gadkari[1], Sarthak

Saini[1], Vinayak Khodake[1], and Sutirth Dey[1*]

Affiliations:

[1]Indian Institute of Science Education and Research (IISER) Pune, Dr. Homi Bhabha

Road, Pashan, Pune, Maharashtra, 411008, India.

[2]Present address: Department of Bioengineering, Indian Institute of Science (IISC),

Bengaluru, India

***Name and address of the corresponding author:**

Sutirth Dey

Professor, Department of Biology

Indian Institute of Science Education and Research, Pune

Dr. Homi Bhabha Road, Pune, Maharashtra, India 411 008 Tel: +91-20-25908054

Email: s.dey@iiserpune.ac.in

Running title: Experimental evolution of *Drosophila* without its microbiome

Keywords: experimental evolution, *Drosophila melanogaster*, gut microbiome, RNA-Seq,

phenotypic plasticity, evolutionary addiction

Author contributions: AM and SD formulated the study. AM, AK, BV, CG, SS, and VK

carried out the assays. AM and SD did the data analysis. AM and SD wrote the manuscript

with inputs from the other authors.

Acknowledgments: We acknowledge the help of Krishna Chaitanya Sattaru, Navanath

Kadam, Saee Patil, Shreya Mishra, Sohum Ranade, Viraj Gaonkar, and Yashvita

Subramanian in data collection.

Data availability: All the data relevant to this study will be made publicly available in the

Dryad Digital Repository upon acceptance.

Funding: AM was supported by the Council for Scientific and Industrial Research (CSIR)

Senior Research Fellowship. AK was supported by the DST-INSPIRE fellowship. BV was

supported by a fellowship from IISER Pune. CG was supported by the UGC Junior

Research Fellowship.

Conflicts of interest: The authors do not have any conflict of interest.

Abstract

The ability of the gut microbiome to contribute to diverse host phenotypes and, in turn,

alter host fitness, raises the possibility that the presence of such microbes can influence

the host's evolution. However, empirical studies on the evolutionary aspects of host-

microbe relationships are rare. Here, we used experimental evolution on laboratory

populations of *D. melanogaster* to understand how a host can evolve without its

microbiome. For this, we compared replicate *Drosophila* populations reared without the

microbes (labeled as MBL_{1-4}, for "Microbiota-less") for 54 host generations with their

corresponding controls (labeled as MB_{1-4}), whose microbiome was reconstituted every

generation. This comparison was done in two common-garden assay environments: with

microbes and microbe-free. We found that contrary to what we expected, only modest

adaptations were seen in the MBL populations. When looking at the phenotypic plasticity

of MB-MBLs across with-microbe and microbe-free environments, we saw that MBLs had

lesser plasticity than MBs. This might hint at MBLs reducing their dependence on

microbes. We performed RNA-Seq on MBL_1 and MB_1 populations to see if there were

changes in gene expression patterns. We found that a cluster of anti-microbial peptides

(AMPs) was up-regulated, and another cluster of heat shock proteins (HSPs) was down-

regulated in MBL_1 compared to MB_1. These results show that while the host's homeostasis

is perturbed by microbiome manipulation on short timescales, the host-microbiome

integration seems more labile on longer timescales, with hosts showing potential for

adaptation without the microbes.

Introduction

Host-associated microbes are shown to modulate host biology (Cryan and Dinan 2012; McFall-Ngai et al. 2013; Sommer and Bäckhed 2013; Lynch and Hsiao 2019; McFall-Ngai 2024). These symbiont microbes can enhance the host's nutritional capacity (Douglas 1998; Ridley et al. 2012; Sun et al. 2022), offer protection from pathogens through colonization resistance (Caballero-Flores et al. 2023; Spragge et al. 2023), and shape the host's physiology (Sommer and Bäckhed 2013; Grenier and Leulier 2020), life history (Shin et al. 2011; Gould et al. 2018; Storelli et al. 2018; Lee et al. 2019; Gnainsky et al. 2021; Arias-Rojas and Iatsenko 2022; Suyama et al. 2023), behavior (Cryan and Dinan 2012; Vuong et al. 2017; Wong et al. 2017; Schretter et al. 2018), and immunity (Chervonsky 2013; Lee et al. 2013; Belkaid and Hand 2014; Selosse et al. 2014; Ansaldo et al. 2021; Jordan and Clarke 2024). In hosts where the impact of this symbiont-mediated modulation is substantial, microbes can alter the host's fitness (Rosenberg and Zilber-Rosenberg 2016; Ma et al. 2023). The ability of symbionts to contribute to diverse host phenotypes and, in turn, influence host fitness raises the possibility that the presence of such microbes can influence the host's evolution (Gilbert et al. 2015; Macke et al. 2017; Kolodny et al. 2020; Henry et al. 2021).

As microbiome acquisition can happen on faster timescales than the host's generation time, the microbiome can confer rapid adaptation to the hosts (Zilber-Rosenberg and Rosenberg 2008; Lau and Lennon 2012; Alberdi et al. 2016; Voolstra and Ziegler 2020). These adaptations can begin with the hosts expanding their niche with the help of microbes (Kopac and Klassen 2016; Borges 2017; Kolodny and Schulenburg 2020; Hoang et al. 2021; Koide 2023). For example, a study showed that stinkbugs can gain insecticide resistance within a single generation by acquiring the bacterial strain of *Burkholderia* that can degrade the insecticide fenitrothion (Kikuchi et al. 2012). This study also showed that along with the insecticidal resistance, the stinkbugs gained a fitness advantage which could help this symbiosis to spread in the population.

Theoretical studies have shown that under certain conditions (strong vertical selection and short generation time), hosts can get a fitness advantage from symbiont microbes, even if the microbes pay costs for such an advantage (Vliet and Doebeli 2019). It has been argued that the microbiome can alter the mean and the variance of the host's phenotypic distribution, which in turn can enable the latter to explore a novel fitness landscape (Henry

et al. 2021). These authors also suggest that the variance contributed by microbes can

alter the heritability of the traits, thus affecting the host's response to selection.

Another way in which the microbiota can affect the host's evolutionary potential is through

phenotypic plasticity. It is known that symbiotic microbes can be an important source of

phenotypic plasticity for their host (Alberdi et al. 2016), which can serve as an initial step in

the host's adaptation to novel environments (Hoang et al. 2021).

In light of the above discussion, it comes as no surprise that the potential impacts of

microbiota on the evolution of their host has become a major topic of investigation in the

field (Macke et al. 2017; Kolodny et al. 2020; Henry et al. 2021). However, there are more

pragmatic anthropocentric reasons for understanding the long-term impact of microbiota

on host evolution. In modern human societies, the overuse of antibiotics and a hygienic

lifestyle, along with other factors, could lead to the disappearance of beneficial microbes

(Noverr and Huffnagle 2004; Blaser and Falkow 2009). The recent COVID-19 pandemic-

induced lifestyle changes, such as the overuse of soaps and sanitizers, can add to this

selection pressure on certain symbiotic microbes. It has been hypothesized that such loss

of ancestral microbes (e.g., *H. Pyroli*) can increase the risk factors for conditions such as

gastroesophageal reflux disease (GERD) and childhood asthma (Blaser 2008; Blaser and

Falkow 2009). Even the microbiota of many naturally occurring species might have been

altered due to increased antibiotic load in the environment (Baquero et al. 2019; Larsson

and Flach 2022) and microenvironmental alterations brought by climate change (Williams

et al. 2023). Therefore, understanding the evolutionary implications of altering or removing

the microbiota on the hosts has emerged as a key question in evolutionary biology (Macke

et al. 2017; Henry et al. 2021).

Here, we used experimental evolution using *D. melanogaster* as a model system to

understand how a host can evolve in the absence of its microbiome. For this, we kept four

large (~2400 individuals) outbred populations of *D. melanogaster* microbe-free using

bleach (labeled 'MBL' populations) for more than 50 generations. This involved bleaching

D. melanogaster eggs every generation and rearing them in sterile food in a sterile

environment. Each of these four selected populations had an ancestry-matched control

(labeled 'MB' populations) where the microbiome was reconstituted just after the bleach

treatment from a common source pool. For this reconstitution, microbes were sampled

from the environment of the flies via fecal microbiome transfer. We compared the selected

MBL and control MB populations in two assay environments: (a) in the presence of

microbes (labeled "with-microbes" environment) and (b) in the absence of microbes

(labeled "microbe-free" environment). Previous observations from the lab showed that the

symbiotic microbiome associated with our outbred baseline lab population affects diverse

host traits and host fitness (Malwade et al., manuscript under prep). This observation

underscored the important role that the microbiome plays in maintaining the host's

homeostasis. Based on these results and the available literature, we hypothesized that the

host would face substantial selection pressure in the absence of microbes. Therefore, we

expected a rapid evolution of the host to maintain the homeostasis seen in the presence of

microbes in our ancestral baseline populations.

We assayed these populations twice during selection: generations 17-20 and generations

54-57. We found that even after 54 generations without microbes, contrary to

expectations, there were only modest adaptations in MBL populations. The percent

survival from egg to adulthood was higher for MBLs when tested under the microbe-free

assay environment. We also found that in the with-microbe assay environment, MBL

males had lower desiccation resistance. The increase in survival might be an adaptation,

and the reduction of male desiccation resistance might be a cost paid by MBLs. We also

saw that the effect sizes over evolutionary differences between MB-MBLs shifted from

"small to medium" effects in Gen 17-20 towards "medium to large" effects in Gen 54-57.

When looking at the phenotypic plasticity of MB-MBLs across with-microbe and microbe-

free environments, we saw that MBLs had lesser plasticity than MBs. This might hint at

MBLs reducing their dependence on microbes. To see if there are changes in gene

expression patterns, we performed RNA-Seq on one population each of MB and MBL. We

found that a cluster of anti-microbial peptides (AMPs) was up-regulated, and another

cluster of heat shock proteins (HSPs) was down-regulated in MBL as compared to MBs.

These RNA-Seq results also hinted that the peritrophic matrix could be a site of interest for

carrying out further investigations related to these populations.

───

Methods

Details of the experimental evolution lines

The experimental evolution lines were derived from the four ancestral baseline populations named NDB_{1-4} (at their generation 21 in the lab). The details about the NDB_{1-4} are provided in Supplementary Text S1.1. Each ancestral NDB line gave rise to two populations: MBL ("Microbiome-Less," selected population that is without the microbiome) and MB (corresponding ancestry-matched control population with microbiome) (Supplementary Figure S1). For example, NDB_1 gave rise to MBL_1 and MB_1, NDB_2 gave rise to MBL_2 and MB_2, and so on. Thus, populations with the same numerical index (e.g., index 2 for MBL_2 and MB_2) share a common ancestry and were always assayed together and treated as blocks in statistical analysis. This design means a total of four replicate selected populations (MBL_{1-4}) were compared against four replicate control populations (MB_{1-4}) (Supplementary Figure S1). Each population in this setup is maintained at a population size of ~2400 in the plexiglass cages with the added implementation of a selection protocol and selection conditions described in the next section. NDB_1 served as the common source of the microbiome for all the MBs (for the rationale of this step, see microbiome reconstitution section).

Experimental evolution protocol employed each generation

MBLs were handled aseptically inside a Level II biosafety cabinet (make: Microfilt, India). All things that came in contact with MBLs for their maintenance were either autoclaved, filtered through a 0.2-micron filter, or surface-sterilized with 70% ethanol.

Every generation, on day 12 after egg deposition, each experimental evolution population of MBLs and MBs (total 8 populations) was provided with autoclaved yeast in the form of moist paste to boost their egg output. This paste also contained a few drops of acetic acid so that the flies are attracted to yeast. A fresh food plate with an exposed vertical food surface was provided for 12-16 hrs to these populations at the end of day 13 to collect eggs for the next generation. On day 14, these eggs were collected using a moist paintbrush and transferred to cell strainers (TCP026, HiMedia) for egg dechorionation (Supplementary Figure S2A). The cell strainer was then placed into one of the wells of a 6-well plate. These eggs were washed with 2% bleach for 2 min to remove the chorion layer and any associated microbes (Supplementary Figure S2B). The cell strainer with

dechorionated eggs was transferred to another well of the 6-well plate for another quick

wash with 2% bleach to kill any residual microbes. Next, the cell strainer with eggs was

transferred back to back to two fresh wells of the 6-well plate for two successive washes

with autoclaved dH$_2$O to remove bleach. To create MBLs, ~400 eggs were transferred to

translucent, plastic *Drosophila* milk bottles (Laxbro Inc., Catalog No. FLBT-20) that had

~50-60ml autoclaved banana-jaggery media at the bottom (Supplementary Figure S2C).

To make MBs, the native microbiome of NDB$_1$ was reconstituted (see next section) on the

dechorionated eggs before they were transferred to the food bottles (Figure 4.2C).

The presence of microbes in MBs and their absence in MBLs is shown by PCR using 16S

rRNA universal bacterial primers in Supplementary Figure S5 as described in

Supplementary Text S3.1 and S3.2.

The procedure of microbiome reconstitution

We chose passive microbiome reconstitution to create flies with microbes for long-term

experimental evolution. Supplementary Figure S3 illustrates the reconstitution protocol in

detail. The microbiome was reconstituted from the ancestral NDB$_1$ population for all the

MBs (i.e., MB$_{1-4}$). This ensured that there were no differences across the four MB

populations in terms of the microbiota that they were harboring. Several different sources

of the microbiome (say, every MB$_i$ getting their microbiome from their ancestral NDB$_i$)

would mean there is a greater chance of differential microbial association.

We describe the microbiome reconstitution protocol in detail in Supplementary Figure S3.

The source population for the microbiome reconstitution is NDB$_1$ unless stated otherwise.

A total of eight fly vials with non-autoclaved normal banana-jaggery food were set to keep

the flies for 24 hrs. These flies were collected on day 13 after egg collection. In each vial,

50 flies with an equal male-to-female ratio were transferred using mild CO$_2$ anesthesia.

After 24 hrs, flies were discarded, and 1 ml of autoclaved water was added to each vial to

extract their microbiome. Vials were swirled with water till no visible fecal spots were seen

on the vial walls. This method of sampling of microbes from the frass of flies housed in

vials has been used earlier in the literature (Chandler et al. 2022). The microbe-free eggs,

after the bleach treatment, were washed with autoclaved water twice and then dipped

twice into this microbiome slurry to seed the source microbiome on the eggs. These eggs

were then gently transferred to *Drosophila* vials or bottles with autoclaved food to generate

flies with reconstituted microbiomes derived from NDB_1.

Common-garden assay environments for the phenotypic assessment

We explain the assay environments using the MB_1-MBL_1 block (Figure 1). However, the

same is true for the rest of the three blocks of MB-MBL as well. MB_1 received microbiome

from NDB1 every generation. MBL_1 is maintained without any microbes. If we directly

compare the phenotypes of MB_1-MBL_1 in their experimental evolution conditions (i.e., MB

with microbes and MBL without microbes) (Figure 1A), any observed differences between

MB/MBL lines can be due to their evolutionary history (with/without microbes) or the assay

environment (with/without microbes) or both. Therefore, we needed to provide a common

assay condition for the phenotypic comparison of MB_1-MBL_1. We performed a phenotypic

comparison between MB_1-MBL_1 in the native environment of MB_1, which is with-microbes

environment, and also in the native environment of MBL_1, which is microbe-free (Figure

1B). For the with-microbe environment, MB_1 as well as MBL_1 were maintained in this

common-garden environment in the presence of microbes for a generation. F1 flies with-

microbes on which we performed the assays were raised as described in Supplementary

Text S2 and Supplementary Figure S4. For a microbe-free environment, MB_1 and MBL_1

were maintained in this common-garden environment in the absence of microbes for a

generation. Microbe-free F1 flies on which we performed the assays were raised as

described in Supplementary text Figure S4. Matching of environments via common-garden

rearing for a generation minimizes the non-genetic effects, giving us the evolutionary

effects in which we are primarily interested.

The detailed validation of these assay conditions is reported in Supplementary Text S5.1-

S5.2.

Figure 1. Illustration of environments in which experimental evolution was carried out and the environments in which phenotyoic assays are carried out.

RNA isolation, RNA-Seq, and analysis

A subset of flies from the phenotypic assays (at generation 54-57) were flash-frozen in

liquid nitrogen and stored at -80^{0}C. These flies from MB$_1$ and MBL$_1$ populations were sent

to a commercial vendor (miBiome Therapeutics LLP, Mumbai, India) for RNA-Seq, starting

with total RNA isolation. RNA-Seq was done only on female flies.

RNA isolation

Total RNA was extracted using miRNeasy Mini kit (Qiagen, Catalog No. 1038703)

following the manufacturer's protocol. About 30 females were homogenized and lysed in

Trizol reagent, followed by chloroform extraction. The aqueous phase was mixed with 1.5x

chilled ethanol, and the mix was loaded on the columns provided in the kit. RNA was

eluted in 30 μl RNase-free water. The RNA samples were quantitated on a Qubit

fluorometer. Appropriate dilutions were loaded on a high-sensitivity RNA screen tape to

determine the RNA integrity number (RIN).

Library preparation and sequencing

1.5 μg of RNA per sample was depleted of fly rRNA using QIAseq FastSelect - rRNA Fly

Kit (Qiagen, Catalog No. 333262). The NEBNext Ultra II directional RNA library prep kit

(Illumina, Catalog No. E7760L) was used to construct double-stranded cDNA libraries from

the rRNA-depleted RNA. The cleaned libraries were quantitated on a Qubit fluorometer,

and appropriate dilutions were loaded on high-sensitivity D1000 screen tape to determine

the library size.

These libraries were sequenced Illumina NGS platform with 2x150bp chemistry to

generate 10 million paired-end reads (3 Gb data per sample).

RNA-seq analysis

The quality of the reads was assessed using *FastQC v0.11.9*. These reads were passed

through *fastp* to remove adaptors and low-quality bases. The filtered reads were mapped

against the *Drosophila melanogaster* BDGP6.46 top-level chromosome

(GCA_000001215.4) retrieved from ENSEMBL using *HISAT2*. The program *featureCounts*

(mode = fragment) was used to count mapped reads. The count table from *featureCounts*

was given as input to DESeq2 to find differentially expressed genes in the populations.

Transcripts with counts less than ten were removed from all samples. The false discovery

rate (FDR) cut-off was at 0.05, and the log fold change (LFC) threshold was at 1.5.

Details of the phenotypic assays

For all the following assays, eggs were processed for dechorionation using bleach as per

the protocol illustrated in Supplementary Figure S2. These dechorionated eggs were

transferred to the *Drosophila* vials with $\sim$ 6 ml of food at a specified density (30 eggs per

vial unless stated otherwise). All the assays were carried out in autoclaved banana-jaggery

food.

We assayed the following phenotypes in MB_{1-4} vs. MBL_{1-4}.

Percentage survival to adulthood

For this assay, the eggs were collected via synchronized egg collection within 2 hr window.

These eggs were then put into food vials after being counted precisely under a microscope

using a paintbrush to the specified egg density (here, 30 eggs per vial). In this procedure,

eggs were transferred over a thin strip of solidified 1.3% non-nutritive agar with a moist

paintbrush. These eggs were spread over the agar strip such that a single egg could be

counted and moved into a stack. This egg stack was cut away along with the agar layer

using a scalpel that was pre-cleaned with cotton soaked in 70% ethanol. This piece of agar

with exactly 30 eggs was transferred to the food vials with autoclaved banana-jaggery

food. Males and females were sex separated under mild CO_2 anesthesia upon eclosion,

and their numbers were noted. The eclosion of the flies was scored at regular intervals of 2

hours. A total of ten vial replicates were used per treatment for this assay.

Dry body weight

On day 12 after egg deposition, the adult flies were sorted by sex in batches of 20

individuals per sex into 1.5ml micro-centrifuge tubes using CO_2 anesthesia. These flies

were then flash-frozen using liquid nitrogen and, if required, stored at -80^0C till weighing.

The flies were then dried at 60^0C for 72 hrs and weighed on an analytical weighing

balance (ME104, Mettler Toledo, least count = 0.1mg). First, the total weight of (micro-

centrifuge tube + flies) was recorded. Next, flies were removed from the tubes with a dry

paintbrush, and the weight of the empty micro-centrifuge tube was recorded. The

difference between these two gave us the body weight of 20 flies. This number was

divided by 20 to get the average dry body weight per fly. When flies were removed, we

counted these flies to double-check if every micro-centrifuge tube had 20 flies. If not, then

the total weight was divided by that count. A total of ten micro-centrifuge tubes were

weighed per treatment per sex. Thus, 200 flies per treatment per sex were sampled.

Egg-to-adult developmental time

This assay was conducted similarly to the percent survival to adulthood assay, with the

only difference being that eclosion time was noted for emerging flies in the given vial for

male and female flies separately. Males and females were sex separated under mild CO_2

anesthesia upon eclosion. The eclosion of the flies was scored at regular intervals of 2

hours in the case of assays in generations 17-20 and at regular intervals of 6 hours in the

case of assays in generations 54-57.

Female fecundity

On day 12 after egg deposition, flies were transferred to fresh food vials. On day 13, a

male-female pair was introduced into an aerated 50 ml falcon tube containing a food cup

using aspirators, thus completely avoiding CO_2 anesthesia. The food cup provided a

surface for egg-laying. The falcon tubes were kept for 12 hrs in an incubator maintained at

25^0C. After 12 hours, the total number of eggs laid was counted under a microscope. A

total of 50 replicates per treatment were set for this assay.

Desiccation resistance

Ten adult flies of the given sex were introduced into an empty plastic vial (Height: 3.5

inches x Diameter: 1 inch). The flies had no access to food or water. The vials were

plugged with fresh cotton plugs. These vials were checked every two hours for any deaths,

till al the flies were dead. The datasheet kept track of flies alive (total flies to start with -

count of total dead flies) at a given time point. The average time to death due to

desiccation was calculated for the vial. A total of ten such vial replicates were set per

treatment per sex.

Locomotor activity

The flies' locomotor activity was measured using the *Drosophila* Activity Monitoring (DAM)

system (Trikinetics Inc). Using an aspirator, a single fly was introduced into a glass DAM

tube (Length: 6.5 cm, Diameter: 4 mm). The locomotor activity for a given fly is the

average number of times the fly crosses the infrared beam running through the middle of

the glass DAM tube per hour, recorded over six hours. Adult flies (day 12 after egg

deposition) were used for this assay. A total of about 32 flies were used per treatment per

sex. The dead flies during the six hours were excluded from the analysis. The first 15

minutes of activity were excluded from the analysis as this was taken as the time taken for

flies to get acclimatized to the glass DAM tubes.

Mating latency and mating time

The mating assay was performed on virgin males and females. After the eclosion (typically

around day eight after egg deposition), male and female flies were separated under mild

CO_2 anesthesia into vials with fly food. The interval of the fly collection was six hours,

starting with the very first eclosion, to ensure that the eclosed flies were virgins. Males and

females were held separately at the density of ten flies per vial till the mating experiment

on day 12 after egg deposition.

On the day of the experiment, a female was introduced into a plastic *Drosophila* vial using

an aspirator, followed by the introduction of a male. The floor of the vials was covered with

solid 1.3% non-nutritive agar to avoid desiccation of the flies. The time at which the male

fly was introduced (T1), the time at which mating started (T2), and the time at which

mating ended (T3) were noted by live manual observation. The mating latency was

calculated as (T2-T1), and the mating time was calculated as (T3-T2). A total of 40

replicates were used for this assay.

Statistical analysis

The selected populations (MBL_{1-4}) were compared with their corresponding controls (MB_{1-4})

in two types of environments (with and without microbiota). The analysis for these two

environments was done separately. In each environment, four block means of MB_{1-4} were

compared against four block means of MBL_{1-4} using a paired t-test performed in GraphPad

Prism (v10.2.1).

For interpreting effect sizes using Cohen's d, the following criteria were followed: d > 0.8

(high); 0.8 > d > 0.5 (medium); d < 0.5 (low) (Cohen 2013). The graphs are plotted in

GraphPad Prism (v10.2.1). The figures drawn using BioRender have that specification in

their legend description.

In the Results section, along with the trends that are statistically significant (i.e., P < 0.05),

we have also highlighted trends that are not statistically significant (i.e., P > 0.05) but have

come close to statistical significance (P ~ 0.1) <u>with large effect sizes</u> (using Cohen's d).

This is highlighted in blue in Supplementary Tables S5-S8, which provide detailed

information about the statistical comparisons for the phenotypic assays.

Results

First phenotypic assessment of experimental evolution lines at host generations 17-
20

We assayed the selected and control populations twice, first in generations 17-20 and then

in generations 54-57 (The experimental design details of these two assessments are given

in detail in Supplementary Text S4.1 and S4.2, respectively). The results for generations

17-20 are provided in Supplementary Text S6. At this point, we observed that only one trait

(mating time in the with-microbe environment) showed a difference, and the rest of the

traits did not change for the flies kept without the microbiome for 17-20 generations

(Supplementary Tables S5 and S6).

Second phenotypic assessment of experimental evolution lines at host generations
54-57

Supplementary Tables S7 and S8 provide detailed information on the statistical

comparison between MB_{1-4} and MBL_{1-4} for the with-microbe assay environment and

microbe-free assay environment, respectively. Comparing these tables with the

corresponding ones for generation 17 (i.e., Supplementary Tables S5 and S6) shows that

there were a larger number of cases in the later generation where the effect sizes were

medium or large. We interpret this as a signature of evolutionary divergence.

Male traits in host generations 54-57

There was no difference between the body weight of the male MBL_{1-4} and MB_{1-4} flies in the

with-microbe assay environment (Paired t-test, $t_3 = 0.32$, $P = 0.7723$, Figure 2A-left). In the

microbe-free environment, the body weight of the male MBL_{1-4} was more than their

corresponding controls MB_{1-4} with a large effect size even though the P-value was

marginally greater than 0.05 (Paired t-test, $t_3 = 2.89$, $P = 0.0628$, Cohen's d = 1.74 (large),

Figure 2A-right).

The egg-to-adult development time was largely unaffected by the multi-generation

absence of microbes, as seen in both assay environments (Supplementary Tables S7 and

S8, Figure 2B).

MBL_{1-4} had lower desiccation resistance than the MB_{1-4} in the with-microbe environment

with a large effect size, although the P-value was marginally greater than 0.05 (Paired t-

test, t_3 = 3.16, P = 0.0510, Cohen's d = 1.62, Figure 2C-left). No difference was observed

in the microbe-free environment (Paired t-test, t_3 = 0.40, P = 0. 7164, Figure 2C-right).

In both environments, MB_{1-4} and MBL_{1-4} showed similar levels of locomotor activity

(Supplementary Tables S7 and S8, Figure 2D).

Figure 2. Male traits assessed in Gen 54-57 in with-microbe and microbe-free environments. (A) Body weight. (B) Egg-to-adult development time. (C) Desiccation resistance. (D) Locomotor activity using DAM system. In each assay environment, four filled circles are means for each of the four control populations MB_{1-4} and, similarly, four empty circles are means for each of the four selected populations MBL_{1-4}. The black horizontal line represent the grand mean over four population means and the error bars represent the SEM. For each host phenotype, the grand mean over MB_{1-4} was compared against the grand mean over MBL_{1-4} using a paired t-test.

Female traits in host generations 54-57

The body weights of the MB_{1-4} and MBL_{1-4} females were similar in the with-microbes

regime (Paired t-test, $t_3 = 0.82$, $P = 0.4726$, Figure 3A-left) but showed a marginal increase

for MBLs in the no-microbe regime which was not statistically significant, although with a

large effect size (Paired t-test, $t_3 = 2.34$, $P = 0.1015$, Cohen's d = 1.35, Figure 3A-right).

Female egg-to-adult development time was similar for MB-MBLs after multiple generations

without the microbes when assayed in both assay environments (Supplementary Tables

S7 and S8, Figure 3B).

MBL_{1-4} females had lower desiccation resistance in the with-microbe regime, which was

not statistically significant (Paired t-test, $t_3 = 2.45$, $P = 0.0916$, Cohen's d = 0.89 (large),

Figure 3C-left). MBs and MBLs had similar desiccation resistance in the no-microbe

regime (Paired t-test, $t_3 = 0.70$, $P = 0.5365$, Figure 3C-right).

MB_{1-4} and MBL_{1-4} females had similar locomotor activity levels in the with-microbes

environment (Paired t-test, $t_3 = 0.10$, $P = 0.9260$, Figure 3D-left). In the microbe-free

environment, there was a trend of MBL_{1-4} females being more active than the MB_{1-4} that

was not statistically significant (Paired t-test, $t_3 = 2.23$, $P = 0.1122$, Cohen's d = 0.89

(large), Figure 3C-left).

Figure 3. Female traits assessed in Gen 54-57 in with-microbe and microbe-free environments. (A) Body weight. (B) Egg-to-adult development time. (C) Desiccation resistance. (D) Locomotor activity using DAM system. In each assay environment, four filled circles are means for each of the four control populations MB_{1-4} and, similarly, four empty circles are means for each of the four selected populations MBL_{1-4}. The black horizontal line represent the grand mean over four population means and the error bars represent the SEM. For each host phenotype, the grand mean over MB_{1-4} was compared against the grand mean over MBL_{1-4} using a paired t-test.

Pair traits in host generations 54-57

There was a trend of lower survival for MBL_{1-4} than MB_{1-4} in the with-microbe regime, but it

is not statistically significant (Paired t-test, $t_3 = 2.31$, $P = 0.1044$, Cohen's d = 1.81 (large),

Figure 4A-left). In the microbe-free environment, MBL_{1-4} had better survival than MB_{1-4}

(Paired t-test, $t_3 = 3.72$, $P = 0.0388$, Cohen's d = 0.83 (large), Figure 4A-right).

There was a trend of MBL_{1-4} laying more eggs than the MB_{1-4} in the with-microbe regime,

but this difference was not statistically significant (Paired t-test, $t_3 = 1.75$, $P = 0.1791$,

Cohen's d = 1.67 (large), Figure 4B-left). In the microbe-free environment, MBL_{1-4} had a

trend of slightly higher fecundity, but it was not statistically significant either (Paired t-test,

$t_3 = 1.20$, $P = 0.3161$, Cohen's d = 0.58 (medium), Figure 4B-right).

MBL_{1-4} did not differ from MB_{1-4} in the latency to start the mating, as assessed in both with-

microbe and no-microbe regimes (Supplementary Table S7 and S8, Figure 4C).

There was no difference in MB_{1-4} and MBL_{1-4} mating time in the with-microbe regime

(Paired t-test, $t_3 = 0.15$, $P = 0.8928$, Figure 4D-left). In the microbe-free environment, there

is a trend of shorter mating times for MBL_{1-4} than MB_{1-4}, but this is not statistically

significant (Paired t-test, $t_3 = 2.35$, $P = 0.1007$, Cohen's d = 1.39 (large), Figure 4D-right).

Figure 4. Pair traits assessed in Gen 54-57 in with-microbe and microbe-free environments. (A) Survival to adulthood. (B) Female fecundity. (C) Mating latency. (D) Mating time. In each assay environment, four filled circles are means for each of the four control populations MB$_{1-4}$ and similarly, four empty circles are means for each of the four selected populations MBL$_{1-4}$. The black horizontal lines represent the grand means over four population means and the error bars represent the SEM. For each host phenotype, the grand mean over MB$_{1-4}$ was compared against the grand mean over MBL$_{1-4}$ using a paired t-test.

MBLs have reduced phenotypic plasticity than the MBs across with-microbe and
microbe-free environments (Gen 54-57 assays)

Figures 2 to 4 suggested that, compared to the MBs, the trait values of the MBLs changed
relatively less across the two assay environments (i.e., with microbes and microbe-free). In
other words, the reaction norms for the various traits across these two environments are
expected to be flatter for the MBLs than for the MBs. To quantify this observation, we used

effect size (Cohen's d) as a proxy for the measure of phenotypic plasticity shown by MB

vs. MBL populations *across two environments*. (Note: These across-environment effect

sizes are different than those provided in Supplementary Tables S5-S8). Supplementary

Text S4.1 and S4.2 discusses the experimental design used to calculate these plastic

effect changes and why such analysis could not be done in Gen 17-20.

We found that the MBLs, on average, had smaller effect sizes than MBs (Paired t-test, t_{11}

= 2.14, P = 0.0554, Cohen's d = 0.49 (small), Figure 5A), possibly hinting at lower

phenotypic plasticity than the MBs across two assay environments. Another way of looking

at the trait's reaction norm could be through the percentage change that the trait has

undergone across environments. The more the percent change, the more the phenotypic

plasticity. We found that for the traits that we assessed, MBLs had a lower percentage of

change in their trait values across two environments than MBs (Paired t-test, t_{11} = 2.99, P

= 0.0123, Cohen's d = 0.39 (small), Figure 5B).

Figure 5. (A) Effect sizes and (B) percentage changes as a proxy for plastic changes across two environments. Means are indicated by black horizontal lines. The error bars represent SEM. The data is analyzed using a paired t-test. For percent change, Wilcoxon matched-pairs signed rank test (a non-parametric test) is also significant (W = -64.0, P = 0.0093).

The big picture across both assessments: divergence in MB-MBLs with time?

In the previous section, we used effect sizes on phenotype differences across assay environments for MBs as well as MBLs as a proxy *for the phenotypic plasticity* exhibited by these two populations. In this section, we look at how effect sizes have changed *for the evolutionary difference between MB-MBLs over time* from the first phenotypic assessment in generations 17-20 to the second assessment in generations 54-57 (Figure 6). These effect sizes are provided in Supplementary Tables S5-S8.

The effects sizes (using Cohen's d) for the phenotypic comparison between MB-MBLs in generations 17-20 were mostly small, with some effects being medium or large (Figure 6-left). In generations 54-57, it looks like the distribution of effect sizes is shifted in the direction of large effects (Figure 6-right). This pattern is seen in both the assay environments: with microbes (circles with a coral fill) and microbe-free (circles without any fill). This might hint that the MBLs are still diverging from MBs even after 54-57 generations of experimental evolution.

Figure 6. Changes in effect sizes (Cohen's d) over all traits with time. Means are indicated by black horizontal lines. The error bars represent SEM. Coral circles indicate with-microbes environment and empty circles indicate the microbe-free environment.

RNA-Seq result – Gen 54-57

After investigating the effects of long-term microbiome absence using broad phenotypic

assays, we used RNA-Seq to determine if specific transcripts are differentially expressed

in MBL_1 vs. MB_1 as a direct consequence of their different evolutionary history. This

comparison was done in two assay environments: (a) microbe-free and (b) with-microbe.

The protocol for RNA-seq is described in detail in the method section.

In the microbe-free regime, where both selected and control populations were assayed

without their microbiome, only a very few transcripts showed differential expression when

analyzed using DESeq2 (Supplementary Text S7.1).

In the with-microbe regime, anti-microbial peptide (AMP) expression is up-regulated in

MBL_1 compared to their control MB_1 (Figure 4.20). The full list of up-regulated genes in

MBL_1 is given in Supplementary Table S9. We also find that a cluster of heat shock

proteins (HSPs) is down-regulated along with several other genes in MBL_1. The full list of

down-regulated genes in MBL_1 is given in Supplementary Table S10.

Among all the differentially expressed genes, only the significant genes with Log_2(Fold

Change) > 1.5 and adjusted P-value < 0.05 are highlighted in Figure 7 and given in

Supplementary Tables S9 and S10.

Gene ontology analysis of differentially expressed genes is presented in Supplementary

Text S7.3.

Figure 7. Differentially expressed genes in MBL₁ vs MB₁ in the with-microbes environment using DESeq2. The red dots represents the up-regulated genes, while blue dots represent down-regulated genes in MBL₁. The protocol for the RNA-Seq is described in the method section.

Discussion

MBL$_{1-4}$ show few changes even after 54 generations without microbes

In previous observations from the lab (Malwade et al., manuscript under prep), we saw that almost all the traits we investigated were affected by the removal of the microbiome. This led us to infer that microbiota is crucial for the realization of a large number of traits in *Drosophila melanogaster*, an observation that corroborates the results of countless studies from the literature (Broderick and Lemaitre 2012; Erkosar et al. 2013; Douglas 2018b; Lesperance and Broderick 2020; Ludington and Ja 2020). This led to the hypothesis that the removal of microbiota is likely to be extremely stressful for the flies, as many traits would be affected at once. Hence, we expected the microbiota-less environment to exert very strong selection pressure on the MBL1-4 flies, leading to changes in multiple life-history traits, maybe including some trade-offs, within relatively few generations. However, the results proved to be otherwise.

In the first set of phenotypic assays done in host generations 17-20 (Supplementary Text S6), we saw that the mating time of MBLs was shorter than the MBs (Supplementary Figure S13D, with-microbes environment) and female body weight was slightly elevated for MBLs (Supplementary Figure S12A), microbe-free environment). Apart from these two changes, we did not see any evidence of evolutionary divergence between MB$_{1-4}$-MBL$_{1-4}$ in either of the two assay environments.

In the second set of phenotypic assays done in host generations 54-57 (Figures 2 to 4), we saw that two traits showed significant change. In the microbe-free environment, we found that MBL$_{1-4}$ had better egg-to-adult survival than MB$_{1-4}$ (Figure 4A). In the with-microbes environment, MBL$_{1-4}$ males had lower desiccation resistance (Figure 2C). It is possible that this lower desiccation resistance might be a cost incurred by MBLs in this environment for improving their egg-to-adult survival in the selection environment (i.e., microbe-free environment).

When we looked at the effect sizes over all the traits assessed in Gen 17-20 vs. Gen 54-57 (Figure 6), we saw that the effect size distribution shifted from mostly "small and medium" effects to "small, medium, and large" effects. This suggests that even after 54 generations, the selected flies are still diverging from the control flies.

We assayed the life-history traits both in the presence and the absence of microbes. Thus, any improvement the fitness of the MBLs (compared to the MBs) in the microbe-free

environment can be construed as the effect of adaptation in the selection environment. On

the other hand, any reduction in the fitness of the MBLs (again compared to the MBs) in

either assay environment would signify a cost of adaptation under microbe-free conditions.

Our results showed that there were very few adaptations or trade-offs.

There are several potential ways to interpret these observations, some of which may not

be mutually exclusive. One possibility is that the selection pressure on the MBLs was so

strong that they responded extremely fast, and therefore, by the time we assayed them

first (i.e., by 17-20 generation), they had already responded maximally to the selection

pressure and were back to the MB levels across all traits. However, if this were to be the

case, in the microbe-free assay environments, we would have expected the MBLs to have

generally greater fitness than the MBs. This is because the microbe-free assay

environment is the "selection" environment for the MBLs but a novel environment for the

MBs. Since the MBLs did not do better than the MBs in this assay environment, therefore

we conclude that the former had not adapted too well in the selection environment.

Another possibility is that even though the selection pressure is strong, the flies did not

possess sufficient genetic variability to respond to the selection pressure. This might have

been a valid concern if, like many studies on *Drosophila*-microbiota systems, the

experiments were performed on highly inbred or iso-female strains (e.g., *Canton-S,*

Oregon-R, and *w^{1118}*). However, our fly populations are outbred and were reinvigorated

through genetic mixing of four outbred populations only one year prior to the initiation of

this study (Supplementary Text S1.3). Our regular observation of these flies indicates that

they are vigorous in the cage (in terms of activity) and have high fecundity and egg-to-

adult survivorship (A Malwade, personal observations). Therefore, we do not have any

reason to believe that these flies did not have sufficient genetic variations. More

importantly, the fact that egg-to-adult survivorship did change over time indicates that the

populations were indeed evolving, which would not have been possible in the absence of

genetic variation.

A third possibility is that while some divergence has indeed happened, they have not been

picked up by our assays. For example, some previous studies have shown that the role of

microbiota in affecting host physiology becomes prominent only in nutritionally unbalanced

diets (Shin et al. 2011; Storelli et al. 2011). It is entirely possible that we will see more

differences between the MBs and the MBLs if we assay them on a different diet. However,

previous observations from the lab (Malwade et al., manuscript under prep) have shown

that removing the microbiota leads to changes in almost every trait, even under the

standard food medium used in our lab. Therefore, it is unlikely that the lack of differences

between MBs and MBLs is due to the food on which they were assayed or selected.

This brings us to the fourth possibility that although the removal of microbiota can lead to

large-scale changes in a number of life-history and behavioral traits in a single generation,

this does not translate into strong selection pressure on an evolutionary timescale. Our

results would make sense when we consider the possibility that if the selection pressure

on the host to evolve in the absence of microbes is low, then the evolutionary response

can be slow and mild, as we observed in this study. We hypothesize that the results of

long-term experimental evolution can be explained by the so-called "evolutionary addiction

hypothesis" (Douglas 2018a; Moran et al. 2019; Hammer 2023). This hypothesis suggests

that when the microbiome is perturbed or culled entirely, hosts experience "withdrawal

symptoms" as the host and microbes have lived together for an extended period (Hammer

2023). Stated differently, the host can take care of itself in the absence of the microbes, it

is just that there is an inertia, borne out of very-long coexistence, towards the same. Thus,

when we remove the microbiome for the first time, the host goes in an altered state of

physiology as the host had evolved some dependence on the microbes that were always

present many generations prior to the manipulation. But, the host still had the capacity to

perform its function without the microbiome. Once several generations passed without the

microbes, the host adjusted itself to the absence of a microbiome without any major

evolutionary changes. This idea of evolutionary addiction might explain why this

experimental evolution happened under low selection pressure and why we haven't seen

major evolutionary changes in our experimental evolution so far.

MBL$_{1-4}$ might be reducing their dependence on the microbes as inferred by the

overall lesser phenotypic plasticity than the MBs

For the phenotypic assays done in generations 54-57, the differential response of MBs and

MBLs in two different environments (with microbes vs. microbe-free) represents the

phenotypic plasticity of these two populations. We wanted to find out if losing microbial

partners for multiple generations has made the MBLs more independent of them. If MBLs

have indeed started becoming less dependent on their microbial partners, then we might

see that, across the two environments, MBLs might be phenotypically less plastic, as the

presence or absence of microbes would affect MBLs less than the MBs.

If there is only one trait under consideration, the phenotypic plasticity of two populations

can be compared using the slopes of the reaction norms for each population across the

environments: the greater the slope, the more the phenotypic plasticity. What we were

looking at was phenotypic plasticity *across multiple traits*. To calculate average phenotypic

plasticity across different traits, one needs a dimensionless (or a normalized) measure.

This is needed because slopes for different traits (e.g., mg of bodyweight, no. eggs laid,

percent survival) cannot be added as they have different units. We used two

normalized/dimensionless proxies of phenotypic plasticity that fit the above criteria and

then compared the mean phenotypic plasticity for the MB-MBL populations. These proxies

were (a) effect size using Cohen's d (Figure 5A) and (b) percent change (Figure 5B). The

results suggest that MBLs had lower phenotypic plasticity across with-microbes and

microbe-free environments. This observation, in turn, might hint that MBLs might be less

dependent on the microbes.

MBL$_1$ and MB$_1$ have at least three sets of differentially expressed gene clusters

To see if MBLs have started showing differences in gene expression compared to MBs,

we performed an RNA-seq on one block of the populations, namely MBL$_1$ and MB$_1$.

A cluster of heat shock proteins (HSPs) is downregulated in MBL$_1$

Heat shock proteins are the proteins that are involved when the organism experiences

stress (Feder and Hofmann 1999). Even though, as their name suggests, they play an

important part primarily in the heat shock response (Lindquist 1986), some of these

proteins can be expressed in response to various other stresses such as cold, crowding,

anoxia, desiccation, etc., where these proteins can act as chaperones (King and MacRae

2015). We find that several HSPs, such as HSP23, HSP26, HSP67, and HSP68, and

subunits from HSPs, such as HSP67 and HSP70, are down-regulated in flies that were

maintained without the microbiome for multiple generations.

RNA-Seq results on MBL$_1$ agree with the desiccation resistance data over all four replicate
MBLs

Lower expression of HSPs in MBL$_1$ might hint at a muted stress response of the MBLs due
to the lower availability of HSPs to counter the stress. This correlation of stress resistance
with the expression of HSPs is well documented (Feder and Hofmann 1999). When we
looked at the desiccation resistance of MBL$_{1-4}$ vs. MB$_{1-4}$ in the with-microbes regime, we
found that MBL$_{1-4}$ indeed had a lower desiccation resistance than MB$_{1-4}$ for males (Figure
2C). For females, it comes close to statistical significance as well (Figure 3C). Both these
effects are large. While we haven't established a causal link in this trend of lower
desiccation resistance, this observation correlates the lower expression of HSPs to a
phenotypic readout of lower desiccation resistance.

Looking at this desiccation-HSP expression correlation, one can speculate that the thermal
stress tolerance of MBL$_{1-4}$ will be lower than that of MB$_{1-4}$ due to lower expression and,
hence, lower levels of HSPs. But that need not be necessarily true. A study in land snails
has shown that the desiccation-sensitive species maintain more HSPs than the
desiccation-resistant species (Mizrahi et al. 2010). If the same logic holds true for thermal
stress tolerance, then the lower expression of HSP might hint at a better ability of MBLs to
resist heat stress.

Antimicrobial peptides (AMPs) upregulated in MBL$_1$

Antimicrobial peptides (AMPs) are small cationic molecules that interact with bacterial
membranes to damage and kill the bacteria (Brogden 2005). They are widespread across
the tree of life, with animals and plants harboring various types of AMPs to deal with
pathogens (Mookherjee et al. 2020). In *Drosophila* melanogaster, there are seven types of
AMPs whose functions are known, and a detailed discussion on them is available in
(Lemaitre and Hoffmann 2007; Hanson and Lemaitre 2020). In addition to their classical
role of defense against pathogens, these AMPs are also shown to control the host
microbiome (Franzenburg et al. 2013; Mergaert 2018; Marra et al. 2021; Maritan et al.
2024).

A study that looked at the gene expression changes in the *Drosophila* gut using
transcriptomics when the host is colonized with the microbes vs. it is kept without microbes
found that the presence of a microbiome leads to upregulated expression of several

immune response genes, including AMPs such as Attacin, Drosomycin and stress

response genes such as HSP23, HSP67Bb, and HSP70Bc (Broderick et al. 2014). While

their observation of the up-regulation of the immune response matches our results in

MBL_1, the HSP response is down-regulated in MBL_1, in contrast to the observation by

Broderick et al., which was done over the host's lifetime (Broderick et al. 2014). The up-

regulation of the immune system upon microbiome colonization is also reported in other

subsequent single-generational studies that are done over the host's lifetime (Elya et al.

2016; Chandler et al. 2022).

These gene expression studies discussed above on *Drosophila* within a host's single

generation show that the presence of a microbiome leads to AMP up-regulation compared

to when a microbiome is absent. In our case, both MBL_1 and MB_1 were given the same

native microbiome for a single generation after 54 generations of microbiome absence for

MBLs. As both MB and MBL received the same native microbiome from a common pool

(shown in Supplementary Text S5.1-S5.2), their different evolutionary history might be the

main driving force in the up-regulation of AMPs, not the differential presence of

microbiome.

A hygiene hypothesis-like phenomenon can explain the overactive humoral immune

response in MBLs

MBLs have been kept away from any microbes for more than 50 generations. Several

studies have shown that microbes are indispensable for the proper calibration and

modulation of the immune system, especially early in life (Mazmanian et al. 2005; Abt et

al. 2012; Selosse et al. 2014; Gensollen et al. 2016; Knoop et al. 2017; Henrick et al.

2021; Hoang and King 2022; Metcalf et al. 2022; Donald and Finlay 2023). This calibration

includes proper sorting of self vs. non-self (Lee 2008; Lhocine et al. 2008). There is a

possibility that MBLs might have lost the delicate balance that maintains this calibration

and, as a result, are over-expressing AMPs compared to MBs to actively regulate the

microbiome.

In humans, this phenomenon is often discussed in the context of the hygiene hypothesis

(or a refined version of the same, called the old friend hypothesis) (Wills-Karp et al. 2001;

Bach 2018; Blackwell 2022). This hypothesis states that exposure to certain

microorganisms, particularly in early life, can reduce the frequency of atopic diseases and

immune-mediated disorders in organisms (Stiemsma et al. 2015). It is possible that the

observed increased levels of AMPs signify immune overactivation via immune

miscalibration over many generations. It is already known that *Drosophila* can suffer from

autoimmune disorders (Mortimer et al. 2021). Therefore, if our conjecture about the

hygiene hypothesis is true then one can predict that the MBLs would be suffering from

higher instances of auto-immune disorders, something that we would like to investigate in

detail in a future study.

Apart from the HSPs and AMPs, another smaller set of chitin-binding genes also seem to

be differentially expressed in MBL_1. We have discussed this in Supplementary Text S8.

Conclusion

Contrary to our expectation, removing microbes for the extended period of 54 host

generations did not result in prominent evolutionary changes in the host. This might be due

to weak evolutionary pressure on the host and can be explained using the "evolutionary

addiction hypothesis." This might also mean that, in the case of outbred *Drosophila* as the

host, the "host + microbe" unit (as a whole) may not be well-integrated on evolutionary

timescales as proposed by the hologenome theory of evolution. This has implications for

how host-microbe interactions are viewed and modeled. While in a given host, certain

traits can be more affected by the microbes (e.g., egg-to-adult survival in our case), certain

other traits may not show this functional integration with microbes (e.g., developmental

time in our case) and, hence, remain unaffected after microbiome removal. So, instead of

focusing on the entire "holobiont" as an independent entity, we think looking at the

functional integration between host and microbes trait-by-trait might be of greater value to

appreciate the microbiome's role in host evolution.

———————————————————————————————————————

References

Abt, M. C., L. C. Osborne, L. A. Monticelli, T. A. Doering, T. Alenghat, G. F. Sonnenberg, M. A. Paley, M. Antenus, K. L. Williams, J. Erikson, E. J. Wherry, and D. Artis. 2012. Commensal Bacteria Calibrate the Activation Threshold of Innate Antiviral Immunity. Immunity 37:158–170.

Alberdi, A., O. Aizpurua, K. Bohmann, M. L. Zepeda-Mendoza, and M. T. P. Gilbert. 2016. Do Vertebrate Gut Metagenomes Confer Rapid Ecological Adaptation? Trends Ecol. Evol. 31:689–699.

Ansaldo, E., T. K. Farley, and Y. Belkaid. 2021. Control of Immunity by the Microbiota. Annu. Rev. Immunol. 39:449–479.

Arias-Rojas, A., and I. Iatsenko. 2022. The Role of Microbiota in Drosophila melanogaster Aging. Front. Aging 3.

Bach, J.-F. 2018. The hygiene hypothesis in autoimmunity: the role of pathogens and commensals. Nat. Rev. Immunol. 18:105–120. Nature Publishing Group.

Baquero, F., T. M. Coque, J.-L. Martínez, S. Aracil-Gisbert, and V. F. Lanza. 2019. Gene Transmission in the One Health Microbiosphere and the Channels of Antimicrobial Resistance. Front. Microbiol. 10. Frontiers.

Belkaid, Y., and T. W. Hand. 2014. Role of the Microbiota in Immunity and Inflammation. Cell 157:121–141.

Blackwell, A. D. 2022. The Ecoimmunology of Health and Disease: The Hygiene Hypothesis and Plasticity in Human Immune Function. Annu. Rev. Anthropol. 51:401–418.

Blaser, M. J. 2008. Disappearing Microbiota: Helicobacter pylori Protection against Esophageal Adenocarcinoma. Cancer Prev. Res. (Phila. Pa.) 1:308–311.

Blaser, M. J., and S. Falkow. 2009. What are the consequences of the disappearing human microbiota? Nat. Rev. Microbiol. 7:887–894. Nature Publishing Group.

Borges, R. M. 2017. Co-niche construction between hosts and symbionts: ideas and evidence. J. Genet. 96:483–489.

Broderick, N. A., N. Buchon, and B. Lemaitre. 2014. Microbiota-Induced Changes in Drosophila melanogaster Host Gene Expression and Gut Morphology. mBio 5.

Broderick, N. A., and B. Lemaitre. 2012. Gut-associated microbes of Drosophila melanogaster. Gut Microbes 3:307–321.

Brogden, K. A. 2005. Antimicrobial peptides: pore formers or metabolic inhibitors in bacteria? Nat. Rev. Microbiol. 3:238–250. Nature Publishing Group.

Caballero-Flores, G., J. M. Pickard, and G. Núñez. 2023. Microbiota-mediated colonization resistance: mechanisms and regulation. Nat. Rev. Microbiol. 21:347–360. Nature Publishing Group.

Chandler, J. A., L. V. Innocent, D. J. Martinez, I. L. Huang, J. L. Yang, M. B. Eisen, and W. B.
Ludington. 2022. Microbiome-by-ethanol interactions impact Drosophila melanogaster
fitness, physiology, and behavior. iScience 25. Elsevier.

Chervonsky, A. V. 2013. Microbiota and Autoimmunity. Cold Spring Harb. Perspect. Biol.
5:a007294.

Cohen, J. 2013. Statistical Power Analysis for the Behavioral Sciences. Routledge.

Cryan, J. F., and T. G. Dinan. 2012. Mind-altering microorganisms: the impact of the gut microbiota
on brain and behaviour. Nat. Rev. Neurosci. 13:701–712. Nature Publishing Group.

Donald, K., and B. B. Finlay. 2023. Early-life interactions between the microbiota and immune
system: impact on immune system development and atopic disease. Nat. Rev. Immunol. 1–
14. Nature Publishing Group.

Douglas, A. E. 2018a. Fundamentals of Microbiome Science: How Microbes Shape Animal
Biology. Princeton University Press.

Douglas, A. E. 1998. Nutritional Interactions in Insect-Microbial Symbioses: Aphids and Their
Symbiotic Bacteria Buchnera. Annu. Rev. Entomol. 43:17–37. Annual Reviews.

Douglas, A. E. 2018b. The Drosophila model for microbiome research. Lab Anim. 47:157–164.

Elya, C., V. Zhang, W. B. Ludington, and M. B. Eisen. 2016. Stable Host Gene Expression in the
Gut of Adult Drosophila melanogaster with Different Bacterial Mono-Associations. PLOS
ONE 11:e0167357. Public Library of Science.

Erkosar, B., G. Storelli, A. Defaye, and F. Leulier. 2013. Host-Intestinal Microbiota Mutualism:
"Learning on the Fly." Cell Host Microbe 13:8–14.

Feder, M. E., and G. E. Hofmann. 1999. HEAT-SHOCK PROTEINS, MOLECULAR
CHAPERONES, AND THE STRESS RESPONSE: Evolutionary and Ecological Physiology.
Annu. Rev. Physiol. 61:243–282.

Franzenburg, S., J. Walter, S. Künzel, J. Wang, J. F. Baines, T. C. G. Bosch, and S. Fraune. 2013.
Distinct antimicrobial peptide expression determines host species-specific bacterial
associations. Proc. Natl. Acad. Sci. 110:E3730–E3738. Proceedings of the National
Academy of Sciences.

Gensollen, T., S. S. Iyer, D. L. Kasper, and R. S. Blumberg. 2016. How colonization by microbiota
in early life shapes the immune system. Science 352:539–544. American Association for
the Advancement of Science.

Gilbert, S. F., T. C. G. Bosch, and C. Ledón-Rettig. 2015. Eco-Evo-Devo: developmental symbiosis
and developmental plasticity as evolutionary agents. Nat. Rev. Genet. 16:611–622. Nature
Publishing Group.

Gnainsky, Y., N. Zfanya, M. Elgart, E. Omri, A. Brandis, T. Mehlman, M. Itkin, S. Malitsky, J.
Adamski, and Y. Soen. 2021. Systemic Regulation of Host Energy and Oogenesis by
Microbiome-Derived Mitochondrial Coenzymes. Cell Rep. 34:108583.

Gould, A. L., V. Zhang, L. Lamberti, E. W. Jones, B. Obadia, N. Korasidis, A. Gavryushkin, J. M.
Carlson, N. Beerenwinkel, and W. B. Ludington. 2018. Microbiome interactions shape host
fitness. Proc. Natl. Acad. Sci. 115:E11951–E11960. Proceedings of the National Academy
of Sciences.

Grenier, T., and F. Leulier. 2020. How commensal microbes shape the physiology of Drosophila
melanogaster. Curr. Opin. Insect Sci., doi: 10.1016/j.cois.2020.08.002.

Hammer, T. J. 2023. Why do hosts malfunction without microbes? Missing benefits versus
evolutionary addiction. Trends Microbiol. 0. Elsevier.

Hanson, M. A., and B. Lemaitre. 2020. New insights on *Drosophila* antimicrobial peptide function in
host defense and beyond. Curr. Opin. Immunol. 62:22–30.

Henrick, B. M., L. Rodriguez, T. Lakshmikanth, C. Pou, E. Henckel, A. Arzoomand, A. Olin, J.
Wang, J. Mikes, Z. Tan, Y. Chen, A. M. Ehrlich, A. K. Bernhardsson, C. H. Mugabo, Y.
Ambrosiani, A. Gustafsson, S. Chew, H. K. Brown, J. Prambs, K. Bohlin, R. D. Mitchell, M.
A. Underwood, J. T. Smilowitz, J. B. German, S. A. Frese, and P. Brodin. 2021.
Bifidobacteria-mediated immune system imprinting early in life. Cell 184:3884-3898.e11.

Henry, L. P., M. Bruijning, S. K. G. Forsberg, and J. F. Ayroles. 2021. The microbiome extends
host evolutionary potential. Nat. Commun. 12:5141.

Hoang, K. L., N. M. Gerardo, and L. T. Morran. 2021. Association with a novel protective microbe
facilitates host adaptation to a stressful environment. Evol. Lett. 5:118–129.

Hoang, K. L., and K. C. King. 2022. Symbiont-mediated immune priming in animals through an
evolutionary lens. Microbiology 168:001181. Microbiology Society,.

Jordan, C. K. I., and T. B. Clarke. 2024. How does the microbiota control systemic innate
immunity? Trends Immunol. 0. Elsevier.

Kikuchi, Y., M. Hayatsu, T. Hosokawa, A. Nagayama, K. Tago, and T. Fukatsu. 2012. Symbiont-
mediated insecticide resistance. Proc. Natl. Acad. Sci. 109:8618–8622. Proceedings of the
National Academy of Sciences.

King, A. M., and T. H. MacRae. 2015. Insect Heat Shock Proteins During Stress and Diapause.
Annu. Rev. Entomol. 60:59–75.

Knoop, K. A., J. K. Gustafsson, K. G. McDonald, D. H. Kulkarni, P. E. Coughlin, S. McCrate, D.
Kim, C.-S. Hsieh, S. P. Hogan, C. O. Elson, P. I. Tarr, and R. D. Newberry. 2017. Microbial
antigen encounter during a preweaning interval is critical for tolerance to gut bacteria. Sci.
Immunol. 2:eaao1314. American Association for the Advancement of Science.

Koide, R. T. 2023. On Holobionts, Holospecies, and Holoniches: the Role of Microbial Symbioses
in Ecology and Evolution. Microb. Ecol. 85:1143–1149.

Kolodny, O., B. J. Callahan, and A. E. Douglas. 2020. The role of the microbiome in host evolution.
Philos. Trans. R. Soc. B Biol. Sci. 375:20190588.

Kolodny, O., and H. Schulenburg. 2020. Microbiome-mediated plasticity directs host evolution
865   along several distinct time scales. Philos. Trans. R. Soc. B Biol. Sci. 375:20190589.

866   Kopac, S. M., and J. L. Klassen. 2016. Can They Make It on Their Own? Hosts, Microbes, and the
867   Holobiont Niche. Front. Microbiol. 7. Frontiers.

868   Larsson, D. G. J., and C.-F. Flach. 2022. Antibiotic resistance in the environment. Nat. Rev.
869   Microbiol. 20:257–269.

870   Lau, J. A., and J. T. Lennon. 2012. Rapid responses of soil microorganisms improve plant fitness
871   in novel environments. Proc. Natl. Acad. Sci. 109:14058–14062. Proceedings of the
872   National Academy of Sciences.

873   Lee, H.-Y., S.-H. Lee, J.-H. Lee, W.-J. Lee, and K.-J. Min. 2019. The role of commensal microbes
874   in the lifespan of Drosophila melanogaster. Aging 11:4611–4640.

875   Lee, K.-A., S.-H. Kim, E.-K. Kim, E.-M. Ha, H. You, B. Kim, M.-J. Kim, Y. Kwon, J.-H. Ryu, and W.-
876   J. Lee. 2013. Bacterial-Derived Uracil as a Modulator of Mucosal Immunity and Gut-
877   Microbe Homeostasis in Drosophila. Cell 153:797–811.

878   Lee, W.-J. 2008. How Do Flies Tolerate Microorganisms in the Gut? Cell Host Microbe 4:91–93.

879   Lemaitre, B., and J. Hoffmann. 2007. The Host Defense of Drosophila melanogaster. Annu. Rev.
880   Immunol. 25:697–743.

881   Lesperance, D. N., and N. A. Broderick. 2020. Microbiomes as modulators of Drosophila
882   melanogaster homeostasis and disease. Curr. Opin. Insect Sci. 39:84–90.

883   Lhocine, N., P. S. Ribeiro, N. Buchon, A. Wepf, R. Wilson, T. Tenev, B. Lemaitre, M. Gstaiger, P.
884   Meier, and F. Leulier. 2008. PIMS Modulates Immune Tolerance by Negatively Regulating
885   *Drosophila* Innate Immune Signaling. Cell Host Microbe 4:147–158.

886   Lindquist, S. 1986. THE HEAT-SHOCK RESPONSE. Annu. Rev. Biochem. 55:1151–1191. Annual
887   Reviews.

888   Ludington, W. B., and W. W. Ja. 2020. Drosophila as a model for the gut microbiome. PLOS
889   Pathog. 16:e1008398. Public Library of Science.

890   Lynch, J. B., and E. Y. Hsiao. 2019. Microbiomes as sources of emergent host phenotypes.
891   Science 365:1405–1409. American Association for the Advancement of Science.

892   Ma, Y., J. He, M. Sieber, J. von Frieling, I. Bruchhaus, J. F. Baines, U. Bickmeyer, and T. Roeder.
893   2023. The microbiome of the marine flatworm Macrostomum lignano provides fitness
894   advantages and exhibits circadian rhythmicity. Commun. Biol. 6:1–12. Nature Publishing
895   Group.

896   Macke, E., A. Tasiemski, F. Massol, M. Callens, and E. Decaestecker. 2017. Life history and eco-
897   evolutionary dynamics in light of the gut microbiota. Oikos 126:508–531.

898   Maritan, E., A. Quagliariello, E. Frago, T. Patarnello, and M. E. Martino. 2024. The role of animal
899   hosts in shaping gut microbiome variation. Philos. Trans. R. Soc. B Biol. Sci.
900   379:20230071. Royal Society.

Marra, A., M. A. Hanson, S. Kondo, B. Erkosar, and B. Lemaitre. 2021. Drosophila Antimicrobial
Peptides and Lysozymes Regulate Gut Microbiota Composition and Abundance. mBio
12:e00824-21. American Society for Microbiology.

Mazmanian, S. K., C. H. Liu, A. O. Tzianabos, and D. L. Kasper. 2005. An Immunomodulatory
Molecule of Symbiotic Bacteria Directs Maturation of the Host Immune System. Cell
122:107–118.

McFall-Ngai, M. 2024. Symbiosis takes a front and center role in biology. PLOS Biol. 22:e3002571.
Public Library of Science.

McFall-Ngai, M., M. G. Hadfield, T. C. G. Bosch, H. V. Carey, T. Domazet-Lošo, A. E. Douglas, N.
Dubilier, G. Eberl, T. Fukami, S. F. Gilbert, U. Hentschel, N. King, S. Kjelleberg, A. H. Knoll,
N. Kremer, S. K. Mazmanian, J. L. Metcalf, K. Nealson, N. E. Pierce, J. F. Rawls, A. Reid,
E. G. Ruby, M. Rumpho, J. G. Sanders, D. Tautz, and J. J. Wernegreen. 2013. Animals in a
bacterial world, a new imperative for the life sciences. Proc. Natl. Acad. Sci. 110:3229–
3236.

Mergaert, P. 2018. Role of antimicrobial peptides in controlling symbiotic bacterial populations.
Nat. Prod. Rep. 35:336–356. The Royal Society of Chemistry.

Metcalf, C. J. E., B. Tepekule, M. Bruijning, and B. Koskella. 2022. Hosts, microbiomes, and the
evolution of critical windows. Evol. Lett. 6:412–425.

Mizrahi, T., J. Heller, S. Goldenberg, and Z. Arad. 2010. Heat shock proteins and resistance to
desiccation in congeneric land snails. Cell Stress Chaperones 15:351–363.

Mookherjee, N., M. A. Anderson, H. P. Haagsman, and D. J. Davidson. 2020. Antimicrobial host
defence peptides: functions and clinical potential. Nat. Rev. Drug Discov. 19:311–332.
Nature Publishing Group.

Moran, N. A., H. Ochman, and T. J. Hammer. 2019. Evolutionary and Ecological Consequences of
Gut Microbial Communities. Annu. Rev. Ecol. Evol. Syst. 50:451–475.

Mortimer, N. T., M. L. Fischer, A. L. Waring, P. KR, B. Z. Kacsoh, S. E. Brantley, E. S. Keebaugh,
J. Hill, C. Lark, J. Martin, P. Bains, J. Lee, A. D. Vrailas-Mortimer, and T. A. Schlenke.
2021. Extracellular matrix protein N-glycosylation mediates immune self-tolerance in
Drosophila melanogaster. Proc. Natl. Acad. Sci. 118:e2017460118. Proceedings of the
National Academy of Sciences.

Noverr, M. C., and G. B. Huffnagle. 2004. Does the microbiota regulate immune responses outside
the gut? Trends Microbiol. 12:562–568.

Ridley, E. V., A. C.-N. Wong, S. Westmiller, and A. E. Douglas. 2012. Impact of the Resident
Microbiota on the Nutritional Phenotype of Drosophila melanogaster. PLoS ONE 7:e36765.

Rosenberg, E., and I. Zilber-Rosenberg. 2016. Microbes Drive Evolution of Animals and Plants: the
Hologenome Concept. mBio 7:10.1128/mbio.01395-15. American Society for Microbiology.

Schretter, C. E., J. Vielmetter, I. Bartos, Z. Marka, S. Marka, S. Argade, and S. K. Mazmanian.
2018. A gut microbial factor modulates locomotor behaviour in Drosophila. Nature 563:402–
406.
Selosse, M.-A., A. Bessis, and M. J. Pozo. 2014. Microbial priming of plant and animal immunity:
symbionts as developmental signals. Trends Microbiol. 22:607–613.
Shin, S. C., S.-H. Kim, H. You, B. Kim, A. C. Kim, K.-A. Lee, J.-H. Yoon, J.-H. Ryu, and W.-J. Lee.
2011. Drosophila Microbiome Modulates Host Developmental and Metabolic Homeostasis
via Insulin Signaling. Science 334:670–674.
Sommer, F., and F. Bäckhed. 2013. The gut microbiota — masters of host development and
physiology. Nat. Rev. Microbiol. 11:227–238.
Spragge, F., E. Bakkeren, M. T. Jahn, E. B. N. Araujo, C. F. Pearson, X. Wang, L. Pankhurst, O.
Cunrath, and K. R. Foster. 2023. Microbiome diversity protects against pathogens by
nutrient blocking. Science 382:eadj3502. American Association for the Advancement of
Science.
Stiemsma, L. T., L. A. Reynolds, S. E. Turvey, and B. B. Finlay. 2015. The hygiene hypothesis:
current perspectives and future therapies. ImmunoTargets Ther. 4:143–157. Dove Press.
Storelli, G., A. Defaye, B. Erkosar, P. Hols, J. Royet, and F. Leulier. 2011. Lactobacillus plantarum
Promotes Drosophila Systemic Growth by Modulating Hormonal Signals through TOR-
Dependent Nutrient Sensing. Cell Metab. 14:403–414.
Storelli, G., M. Strigini, T. Grenier, L. Bozonnet, M. Schwarzer, C. Daniel, R. Matos, and F. Leulier.
2018. Drosophila Perpetuates Nutritional Mutualism by Promoting the Fitness of Its
Intestinal Symbiont Lactobacillus plantarum. Cell Metab. 27:362-377.e8.
Sun, Q., N. M. Vega, B. Cervantes, C. P. Mancuso, N. Mao, M. N. Taylor, J. J. Collins, A. S. Khalil,
J. Gore, and T. K. Lu. 2022. Enhancing nutritional niche and host defenses by modifying
the gut microbiome. Mol. Syst. Biol. 18:e9933. John Wiley & Sons, Ltd.
Suyama, R., N. Cetraro, J. Y. Yew, and T. Kai. 2023. Microbes control Drosophila germline stem
cell increase and egg maturation through hormonal pathways. Commun. Biol. 6:1–18.
Nature Publishing Group.
Vliet, S. van, and M. Doebeli. 2019. The role of multilevel selection in host microbiome evolution.
Proc. Natl. Acad. Sci. 116:20591–20597. National Academy of Sciences.
Voolstra, C. R., and M. Ziegler. 2020. Adapting with Microbial Help: Microbiome Flexibility
Facilitates Rapid Responses to Environmental Change. BioEssays 42:2000004.
Vuong, H. E., J. M. Yano, T. C. Fung, and E. Y. Hsiao. 2017. The Microbiome and Host Behavior.
Annu. Rev. Neurosci. 40:21–49.
Williams, C. E., C. L. Williams, and M. L. Logan. 2023. Climate change is not just global warming:
Multidimensional impacts on animal gut microbiota. Microb. Biotechnol. 16:1736–1744.

Wills-Karp, M., J. Santeliz, and C. L. Karp. 2001. The germless theory of allergic disease: revisiting
the hygiene hypothesis. Nat. Rev. Immunol. 1:69–75. Nature Publishing Group.
Wong, A. C.-N., Q.-P. Wang, J. Morimoto, A. M. Senior, M. Lihoreau, G. G. Neely, S. J. Simpson,
and F. Ponton. 2017. Gut Microbiota Modifies Olfactory-Guided Microbial Preferences and
Foraging Decisions in Drosophila. Curr. Biol. 27:2397-2404.e4.
Zilber-Rosenberg, I., and E. Rosenberg. 2008. Role of microorganisms in the evolution of animals
and plants: the hologenome theory of evolution. FEMS Microbiol. Rev. 32:723–735.

980
981

982 **Supplementary information**
983
984 **Part 1 – Figures (S1 to S3) – Total 3**
985 **Part 2 – Text (S1 to S8) – Total 8**
986
987

Figure S1. Details of the ancestral and experimental evolution populations.

988
989
990

Figure S2. Egg dechorionation protocol using bleach to make MBL$_{1-4}$ and MB$_{1-4}$ populations. Created in BioRender.com.

992
993

Figure S3. Detailed protocol of microbiome reconstitution from NDB₁. Created in BioRender.com.

Text S1 Details of the baseline populations and amplicon sequencing

S1.1 *Drosophila* NDB stocks and rearing

Wolbachia-free *Drosophila* melanogaster used in the single-generation study were derived from four large, outbred laboratory populations named New Dey Baselines (NDB$_{1-4}$). These fly lines were maintained on the standard banana-jaggery food with a 14-day discrete generation cycle at 250C under constant light conditions with a breeding population of ~2400 individuals. NDBs trace their ancestry to the DB populations (see Supplementary Text S1.3 for the details), which in turn trace back to the wild-caught IV lines (Ives 1970; Rose 1984) via the JB$_{1-4}$ populations (Sheeba et al. 1998).

S1.2 Maintenance regime of baseline *Drosophila* populations (NDB$_{1-4}$)

In the lab, adult *Drosophila* of four baseline populations (NDB$_{1-4}$) are maintained in plexiglass cages (25 cm x 21 cm x 16 cm) at a large population size of ~2400 per population. On day 12 (after egg deposition), the fly population is provided with a moist yeast paste on top of the banana-jaggery food to boost egg output. This yeast paste also contains a few drops of glacial acetic acid to attract the flies to the paste. At the end of day 13, a food plate, with vertical edges of food exposed for oviposition, is provided for 12-16 hrs. On day 14, deposited eggs are collected by cutting thin food strips from the exposed surface, under a binocular stereo microscope. The size of the strips is adjusted under the microscope using a clean scalpel such that each strip contains about 400 eggs. These strips are then transferred to plastic and translucent *Drosophila* milk bottles containing ~50-60ml of banana-jaggery food at the bottom. This marks the new generation's 0th day after egg deposition. Adult flies arising from six such bottles are transferred to fresh plexiglass cages on the 12th day (post egg collection). This gives rise to a new generation with a population size of ~2400 individuals, which do not coexist with the adults of their parental generation at any stage.

S1.3 *Drosophila* DB stocks and rearing

NDB$_{1-4}$ were derived from their most recent ancestors in our lab, Dey Baselines (DB$_{1-4}$) (Sah et al. 2013). These four DBs were maintained in our lab for 220 generations and were used as baselines for this period. We pooled an equal proportion of eggs from each DB population and created four independent NDBs. We maintained NDBs separately for ~10 generations without any assays for them to stabilize after this intermixing. The maintenance of DBs was similar to NDBs except for two differences: (a) DBs were maintained in Drosophila 8-dram vials at an egg density of ~ 60 per vial with ~ 6ml of food as opposed to NDBs that are reared in milk bottles (b) DBs were yeasted on 18th day, provided with a cut plate on 20th day, and their eggs were collected for next generation on day 21. Thus, DBs followed a 21-day discrete generation cycle as opposed to a 14-day discrete generation cycle followed by NDBs.

S1.4 Population-level 16S rRNA amplicon sequencing

To know the relative microbiota composition in *Drosophila* populations, DNA was extracted as described in section S3.2.2. DNA was run on 1% agarose gel to see the extraction quality. After the integrity check on agarose gel, DNA was sent for 16S rRNA V3-V4 region amplicon sequencing to a commercial vendor (1st BASE Laboratories, Malaysia). Sample quality was again confirmed by the vendor using agarose gel, spectrophotometer, and fluorometric method. All samples passed DNA quality check and were subjected to amplicon PCR followed by a quality check. These samples were then processed to prepare amplicon libraries using 2-step PCR according to Illumina's 16S metagenomic library preparation guidelines (Link to the guidelines -

1050   https://sapac.support.illumina.com/downloads/16s_metagenomic_sequencing_library_preparation.
1051   html). These libraries were processed using Illumina's MiSeq platform using 2x300bp chemistry.

1052

S1.5 Data analysis of amplicon sequencing

Sequence adapters and low-quality reads were removed from the paired-end reads before the first
2,00,000 raw reads were extracted using BBTools. Then, the forward and reverse reads were
merged using QIIME. DADA2 pipeline (https://benjjneb.github.io/dada2/) was used to remove low-
quality regions and chimeric errors. The resulting data was in the form of amplicon sequence
variants (ASVs). The taxonomic classification was done using scikit-learn (https://scikit-
learn.org/stable/) and the Naive Bayes classifier against the SILVA database (release 132). This
part of the analysis was performed by the commercial vendor (1st BASE Laboratories, Malaysia).

1061

1062

Text S2 - Note on the use of F1 flies for the assays

In our flies, we removed microbes via dechorionation at the egg stage. Even if we have taken care of the confounding effect of bleach by using proper control, we wanted to keep the effect of the bleach on flies to a bare minimum (ideally none). To that end, we employed an additional measure of using F1 flies for the assays, as explained below (Supplementary Figure S4).

In this approach, to generate flies without the microbes, we followed a protocol employed by several other studies where F1 flies (progeny of the flies that experienced the bleach at the egg stage) were used in the assays (Pais et al. 2018; Selkrig et al. 2018; Lee et al. 2019). This ensured that the F1 assay flies of both types, microbiome-reconstituted (M+) and microbe-free (M-), did not experience bleach in their lifetime (their respective parent population did). To generate F1 M- assay flies, F1 eggs were collected from the flies without microbes (i.e., F0 M- flies, Supplementary Figure S4) and were raised in aseptic conditions, ensuring the absence of microbes. To generate control F1 M+ assay flies, F1 eggs were collected without any manipulation from the flies whose microbiome was reconstituted (i.e., F0 M+ flies, Supplementary Figure S4).

In our conventional/untreated baseline lab populations, about 86 percent of the flies survive till adulthood. With the approach of using the F1 flies we described, control F1 assay flies whose microbiome was reconstituted (i.e., M+ flies) showed just an additional ~5 percent mortality, showing that the overall mortality is kept to a minimum.

Figure S4. Illustration of the full experimental design to study microbiomes' contribution to host phenotypes. To create two types of flies (with the minimal effect of bleach), flies with microbes (M+) and without them (M-), we combined two procedures. First, we chose microbiome-reconstituted flies as a control instead of using untreated flies. The F0 flies of both (M+ and M-) types were generated using bleach at the egg stage. This protocol is illustrated in detail in Figure S2. Second, as discussed in Supplementary Text S2, assays were conducted on progeny of these F0 flies (i.e., F1 flies) that have not experienced bleach in their lifetime to avoid any recency effect of bleach. The eggs collected for F1 experimental flies either had the microbes (M+) or lacked the microbes (M-) depending on the state of their parent's microbiome. Created in BioRender.com.

1082

1083 **Text S3.1 - Assessing the microbiome status of MBs and MBLs during experimental**
1084 **evolution**
1085
1086 We plated adult flies from MBL_{1-4} every generation to see if the microbiome is indeed absent in
1087 these populations (data not shown here). Except for generations and blocks specified in
1088 Supplementary Table S1, in all other generations, MBLs did not show the presence of a
1089 microbiome. We sometimes did observe ~1-2 colonies (per 10 flies plated) in some replicates.
1090
1091 **Table S1. Generations in which MBLs showed transient contamination**
1092

Generation	Affected MBL blocks
37	Block 1 and 3
38	Block 1
39	Block 1

1093
1094 MB_{1-4} were plated every 2-3 generations and CFUs were counted to make sure they received the
1095 reconstituted microbiome (data not shown here).
1096
In generation 40 of the selection line, we performed PCR with 16S rRNA universal primers
(Supplementary Text S3.2.2) on DNA extracted from adult MB_{1-4} and MBL_{1-4} females (day 15 after
egg collection). In Supplementary Figure S5, we see that MB_{1-4} show the presence of the
reconstituted microbiome, and the microbiome is absent from MBL_{1-4}. This also confirms that, by
generation 40, the contamination that we saw in MBLs in previous generations was completely
gone.
1103

Figure S5. 16S rRNA PCR to check the status of microbiome in MB_{1-4} and MBL_{1-4} .

1104
1105
1106
1107

Text S3.2 Fly-associated bacterial load analysis

S3.2.1. Culture-dependent method

Adult flies (day 12 after egg deposition, unless stated otherwise) were sorted by sex under mild
CO_2 anesthesia. Five male and five female flies under anesthesia were surface sterilized with 70%
ethanol in 1 ml microcentrifuge tubes. These flies were then homogenized in 200 μl of autoclaved
dH_2O and plated on the nutrient agar (HiMedia) or the de Man, Rogosa, and Sharpe (MRS)
medium (HiMedia) and incubated for 72 hrs and 48 hrs, respectively, at 25^0C under aerobic
conditions. Dilutions with 10-100 colonies were counted, and the counts were multiplied by the
overall dilution factor to estimate colony-forming units (CFUs) per fly.

S3.2.2. Culture-independent method

To test the bacterial load using PCR, total DNA was extracted from eight to ten adult flies using the
QIAamp DNA Mini kit (Qiagen, Catalogue No. 51304) as per the manufacturer's instructions. We
modified three steps of the manufacturer's protocol in line with the protocol (Jehrke et al. 2018).
First, we used mechanical homogenization (with a homemade handheld motorized pestle mixer) to
lyse the tissue in 1.5ml Eppendorf tubes instead of just Proteinase-K treatment. Second, 20 μl of
20 mg/ml RNase A (HiMedia, Catalogue No. DS0003-5ML) was added to each sample after tissue
lysis. Third, the sample was incubated with Buffer AL at 70^0C for 30 min instead of 10 min. We
avoided any vortex step after tissue lysis to minimize the shearing of the DNA. The quality of total
DNA was checked on the agarose gel. This DNA, composed of the fly and microbial genomes, was
used as a template for PCR using 16S bacterial universal primers with a product size of about
1.5kb. The details of the primers are given below in Supplementary Table S2. The primers were
ordered from Sigma-Aldrich, India. The PCR reactions were set up using GoTaq® Flexi DNA
Polymerase (Promega, Catalogue No. M829) with 40 ng gDNA input. The initial denaturation was
at 94^0C for 2 min, followed by 30 cycles of (denaturation at 94^0C for 30s, annealing at $T_m = 60^0C$
for 1 min, extension at 72^0C for 1 min), with final extension 72^0C for 10 min. The PCR products
were visualized on 1% agarose gel with 1kb plus DNA ladder (NEB, Catalogue No. N3200S) as the
molecular weight marker.

Table S2. 16S bacterial universal primers used for PCR

Primer	Sequence	Label	Reference
Forward	AGAGTTTGATCCTGGCTCAG	fD1	(Weisburg et al. 1991)
Reverse	ACGGCTACCTTGTTACGACTT	rP2	(Weisburg et al. 1991)

1136
1137

Text S4.1 – First phenotypic assessment of experimental evolution lines at host generations 17-20

After 17 generations of selection, we looked at various host traits to see if the experimental evolution without the microbes has led to any divergence in MBLs vis-à-vis the MBs. This assessment was done in two assay environments: (1) with microbes and (2) microbe-free.

In this design, logistically, it was not possible to assay all four blocks of MB and MBL together. Hence, we assayed them in sets of two blocks, namely 1 - 2 and 3 - 4. Supplementary Table S3 gives details of the generations in which the assays took place. MB and MBL populations in each block were always assayed together in the same generation to know the effect of the selection procedure (e.g., MB_1 compared with MBL_1). For all assays done in this set, flies for separate environments (i.e., with microbes and microbe-free) were generated in different host generations (Supplementary Table S3). As two different environments were not assayed in the same host generation, we do not discuss the effect of the environment on these populations. We only discuss the effect of selection in this first set of assays.

Table S3. Details of the assay environments and host populations for Gen 17-20 assays

No.	Assay environment	Block	Host generation
1	With-microbes	3 and 4	17
		1 and 2	18
2	Microbe-free	1 and 2	19
		3 and 4	20

The validation of the microbiome states of flies raised in with-microbes and microbe-free assay environments is reported in Supplementary section S5.1.

Text S4.2 - Second phenotypic assessment of experimental evolution lines at host generations 54-57

In the first assessment of MBL_{1-4} vs. MB_{1-4} that happened in generations 17 to 20 (Supplementary Text S4.1), we noticed that plastic effects across two assay environments (with-microbes and microbes-free) on both of these populations were more significant than the effects of the selection (Figures S11-S13). (By plastic effects, we mean differences seen in a phenotype when the environment changes without a change in host genotype). However, as with-microbes and microbe-free flies were generated in separate host generations (see Supplementary Table S3), we couldn't compare these plastic effects quantitatively with this design. We amended our experimental design so that these plastic effects could be compared directly by conducting the assays in the two environments simultaneously. Supplementary Table S4 shows the details of our experimental plan for the set of assays conducted in generations 54 to 57. This amendment meant, due to logistical reasons, we could only assess one block at a time in the same host generation (in contrast to the Gen 17-20 assays where two blocks were assessed in the same host generation).

This experimental design allows us to make two kinds of comparisons: (1) the effect of experimental evolution in a common-garden assay environment obtained by directly comparing MBs vs. MBLs (Main text, Figures 2 to 4) and (2) the plastic effect of the presence vs. absence of microbes on both MB and MBL populations (Main text, Figure 5)

Table S4. Details of the assay environments and host populations for Gen 54-57 assays

No.	Common-garden environment	Block	Host generation
1	With-microbes	1	54
		2	55*
		3	56
		4	57
2	Microbe-free	1	54
		2	55*
		3	56
		4	57

*For block 2, body weight and female fecundity were assessed in Gen 59

Text S5.1 - Validation of microbiome status in the flies raised in two different assay
environments
S5.1.1 By plating fly homogenates
In the with-microbe environment, on both MRS and NA media, we see the presence of microbes (
Supplementary Figure S6, Set-B and Set-C). Set-A is the autoclaved water that is used to
homogenize the flies. The serial dilutions with ~10-100 colonies were counted to get the CFUs/fly.
While Supplementary Figure S6A shows the representative picture from block 1, CFUs/fly for all
four blocks of MBs and MBLs are compared using paired t-test on block means in Figure S7.
In the microbe-free environment, both MB_1 (Supplementary Figure S6B) and MBL_1 (Supplementary
Figure S6C) show the absence of the microbiome. While block 1 plates do not have any microbes,
other blocks show a few colonies in some replicates (data not shown here).
1226

Figure S6. MRS and NA plates showing presence of microbes in with-microbe
environment and their absence in microbe-free environment. (A) MB_1 and MBL_1 in with-
microbe environment (n = 4). The spots show different serial dilutions used to calculate CFUs/fly.

These CFU/fly for all the replicate populations are shown in Supplementary Figure S7. (B) MB_1 microbe-free environment (n = 3). (C) MBL_1 microbe-free environment (n = 3).

1227
1228
1229   **S5.1.2 MBs and MBLs received similar microbial loads in the with-microbe environment**
1230   **(Gen 54-57 assays)**
1231

Figure S7. Bacterial load in the MB_{1-4}-MBL_{1-4} flies (with-microbe environment) as determined by plating fly homogenates. CFUs/fly on (A) MRS-agar (B) NA-agar. The bacterial load in MBs vs. MBLs is compared using paired t-test on block means. Block means are indicated by black filled circles. The error bar represents SEM. We see that the process of microbiome reconstitution leads to MBs and MBLs carrying similar bacterial loads.

1232
1233
1234   **S5.1.3 PCR showing the presence of microbiome in the with-microbiome environment and**
1235   **depletion of the microbiome in microbe-free environment**
1236
To validate the status of the microbiome in both environments with a culture-independent method,
we extracted DNA from the four types of populations (Main text, Figure 1B). We performed PCR
with 16S rRNA universal primers (details in Supplementary Text S3.2). For PCR, a subset of flies
used for the assay on day 12 were flash-frozen in liquid nitrogen and kept at -80^0C for further use.
For the with-microbe environment, 12 female flies from only MB_1 and MBL_1 (block 1) were used for
the DNA extraction. For the microbe-free environment, total DNA was extracted from 12 female
flies (3 flies per block, four such blocks). The PCR was done with 40ng of input DNA with 30
cycles.
1245
In the with-microbe environment, MB_1 and MBL_1 showed the presence of microbes as expected
(Supplementary Figure S8). In the microbe-free environment, while two replicates have no band,
one replicate each from pooled MB_{1-4} and MBL_{1-4} contains a faint band. This faint band might be
indicative of a few colonies that show up on NA and MRS-agar plates, even after aseptic handling
(data not shown). This signal might get picked up due to saturating PCR conditions. As we are not
bleaching assay flies or using any antibiotics to avoid their toxic effects, we are essentially
maintaining flies without any chemicals for about two consecutive generations, i.e., for ~30 days
(Supplementary Figure S4). So, there might be a tradeoff between the efficiency of keeping all the
microbes away without any chemicals and how long that efficiency can be sustained without any
chemical agent. Another possibility is that the signal is from very few non-culturable bacteria.
1256
1257   These results match the observations in the Supplementary Text S5.1.1 obtained after plating flies
1258   on NA and MRS. Together, the evidence showed that both MB-MBL populations in a with-microbe
1259   environment harbor microbiome and in microbe-free environments do not have microbes with
1260   them.
1261

1262

Figure S8. PCR on MB$_1$ and MBL$_{1-4}$ to confirm the status of microbiome.

1263
1264
1265 **S5.2 Validation of microbiome composition of flies raised in with-microbes assay**
1266 **environment**
1267
1268 **S5.2.1 MB-MBL generation 17-20 assays**
1269
In the previous section, through plating fly homogenates and PCR, we showed that flies raised in
the with-microbes environment indeed have the microbes with them. However, the PCR only
indicates that microbes are present - it does not tell us the composition of the reconstituted
microbiome. In Supplementary Figure S7, we have established that the number of microbes
received by MB and MBL populations is similar. In this section, we test if the reconstituted
microbiome composition is similar for MB-MBLs, and this, in turn, matches the source microbiome
composition of NDB$_1$.
1277
To validate that the composition of the reconstituted microbiome of all four blocks (MB$_{1-4}$ and MBL$_{1-}$
$_4$) reflects the native NDB$_1$ composition, we performed amplicon sequencing on the V3-V4 region of
the 16S rRNA gene (Supplementary Figure S9, method details in Supplementary Text S1.4). Three
biological replicates were processed per population, giving us 24 samples (8 populations: 4 MBs +
4 MBLs, 3 replicates each). The native source pool of NDB$_1$ (at Gen 56) was also sequenced.
Thus, Supplementary Figure S9 shows a total of 27 samples (24 from MB$_{1-4}$-MBL$_{1-4}$ and 3 from
NDB$_1$).
1285
1286 The MB$_{1-4}$-MBL$_{1-4}$ populations also mirror this broad microbiome composition of NDB$_1$ with the
1287 dominance of *Acetobacter* spp. but with small fluctuations across MB-MBLs in *Acetobacter* and
1288 *Lactobacillus* fractions (Supplementary Figure S9). (In 2020, the old genus *Lactobacillus* was
1289 reclassified into 25 new different genera. *Lactiplantibacilus* and *Levilactobacilus* both belong to the
1290 old composite *Lactobacillus* genus.)
1291

Figure S9. Validation of microbiome composition after reconstitution in Gen 17-20 assays. The result of population-level 16S rRNA amplicon sequencing across four replicate populations of MB$_{1-4}$ and four replicate populations of MBL$_{1-4}$. Native microbiome composition of source population NDB$_1$ is also given for the reference (last three columns). Most abundant 20 taxa are shown for each population. For each population, three independent replicates where processed (labelled "A", "B", "C"). Each replicate consisted of pooled DNA extracted from surface sterilized full body homogenates of 8-10 females. The sequencing method and sequencing data analysis are described in Supplementary Text S1.4 and S1.5 respectively.

1292
1293
1294 **S5.2.2 MB-MBL generation 54-57 assays**
1295
1296 In generation 54-57 assays, we sequenced microbiomes from MB$_1$ and MBL$_1$ that were raised in
1297 the with-microbes environments (Supplementary Figure S10). We see that MB$_1$ and MBL$_1$ have
1298 received microbiomes that are very close in composition. *Acetobacter* spp. dominates the
1299 microbiomes of both populations.
1300
1301 Together, these sequencing results for generations 17-20 assay and for generations 54-57 assay
1302 show that microbiome composition received by MBs and MBLs is broadly similar to each other and
1303 to the source microbiome of NDB$_1$ (Supplementary Figure S9-last three lanes).
1304

Figure S10. Validation of microbiome composition after reconstitution in Gen 54-57 assays. Most abundant 100 taxa are shown for each population. For each population, three independent replicates where processed (labelled "A", "B", "C"). Each replicate consisted of pooled DNA extracted from surface sterilized full body homogenates of 8-10 females. The sequencing method and sequencing data analysis are described in Supplementary Text S1.4 and S1.5 respectively.

1305
1306
1307
1308

1309 **Text S6 - First phenotypic assessment of experimental evolution lines at host generations**
1310 **17-20**
1311

1312 **S6.1 Male traits in host generations 17-20**
1313

1314 None of the four phenotypes tested for males – adult body weight, egg-to-adult development time,
1315 desiccation resistance, and locomotor activity- showed an effect of selection as assessed by the
1316 paired t-test over block means (Supplementary Figure S11A-D). Supplementary Tables S5 and S6
1317 provide details of the statistical comparison for the with-microbe assay environment and microbe-
1318 free assay environment, respectively.
1319

Figure S11. Male traits assessed in Gen 17-20 in with-microbe and microbe-free environments. (A) Body weight. (B) Egg-to-adult development time. (C) Desiccation resistance. (D) Locomotor activity using DAM system. In each assay environment, four filled circles are means for each of the four control populations MB_{1-4} and, similarly, four empty circles are means for each of the four selected populations MBL_{1-4}. The black horizontal lines represent the grand means over four population means and the error bars represent the SEM. For each host phenotype, the grand mean over MB_{1-4} was compared against the grand mean over MBL_{1-4} using a paired t-test. DAM data for block 4 in with-microbes environment is not included here in the analysis.

1328 **S6.2 Female traits in host generations 17-20**
1329
In the case of females, egg-to-adult development time, desiccation resistance, and locomotor
activity didn't differ between MBs and MBLs (Supplementary Figure S12A-D) (Supplementary
Tables S5 and S6). The only trait in females that showed some difference between the MBs and
the MBLs was female body weight in the microbe-free environment (Paired t-test, $t_3 = 3.05$, P =
0.0554, Cohen's d = 0.81 (large), Supplementary Figure S12A). Here, although the p-value is
slightly larger than our pre-determined level of statistical significance (i.e., 0.05), we still interpret
this trait as the effect size was found to be large.
1337

Figure S12. Female traits assessed in Gen 17-20 in with-microbe and microbe-free environments. (A) Body weight. (B) Egg-to-adult development time. (C) Desiccation resistance. (D) Locomotor activity using DAM system. In each assay environment, four filled circles are means for each of the four control populations MB$_{1-4}$ and similarly, four empty circles are means for each of the four selected populations MBL$_{1-4}$. The black horizontal lines represent the grand means over four population means and the error bars represent the SEM. For each host phenotype, the grand mean over MB$_{1-4}$ was compared against the grand mean over MBL$_{1-4}$ using a paired t-test. DAM data for block 4 in with-microbes environment is not included here in the analysis.

1338

1339 **S6.3 Pair traits in host generations 17-20**
1340

1341 In traits where both males and females were involved in the assays - percent survival to adulthood,
1342 female fecundity over 12 hrs (assayed in the presence of a male), and mating latency - did not
1343 show any difference (Supplementary Figure S13A-C, Supplementary Tables S5 and S6). The
1344 mating time for MBL_{1-4} was lower than MB_{1-4} in the with-microbes environment (Paired t-test, t_3 =
1345 3.67, P = 0.0349, Cohen's d = 2.35 (large), Supplementary Figure S13D).
1346

**Figure S13. Pair traits assessed in Gen 17-20 in with-microbe and microbe-free
environments.** (A) Survival to adulthood. (B) Female fecundity. (C) Mating latency. (D) Mating
time. In each assay environment, four filled circles are means for each of the four control
populations MB_{1-4} and similarly, four empty circles are means for each of the four selected
populations MBL_{1-4}. The black horizontal lines represent the grand means over four population
means and the error bars represent the SEM. For each host phenotype, the grand mean over
MB_{1-4} was compared against the grand mean over MBL_{1-4} using a paired t-test.

1347
1348
1349

1350 **Table S5.** Summary of all assays in the <u>with-microbes environment (Gen 17-20).</u> T refers to the
1351 grand mean of the MBLs, while C refers to the grand mean of the MBs.

1352

	Assay	Test statistic	P-value	% change (T - C)/C	Cohen's d	Effect size*	Sample size$
1	Dry body weight (Male)	$t_3 = 0.03$	0.9785	-0.3	0.02	Small	10
2	Dry body weight (Female)	$t_3 = 0.23$	0.8356	-3.1	0.18	Small	10
3	Development time (Males)	$t_3 = 0.85$	0.4560	-1.1	0.41	Small	10
4	Development time (Females)	$t_3 = 0.77$	0.4976	-0.9	0.37	Small	10
5	Survival to adulthood	$t_3 = 0.21$	0.8465	0.4	0.10	Small	10
6	Fecundity (Females)	$t_3 = 1.96$	0.1413	-20.1	0.75	Medium	50
7	Desiccation resistance (Males)	$t_3 = 1.18$	0.3236	-13.7	0.54	Medium	10
8	Desiccation resistance (Females)	$t_3 = 1.17$	0.3272	-12.9	0.66	Medium	10
9	Locomotor activity (Males)	$t_3 = 0.82$	0.4969	-3.7	0.30	Small	25 to 32
10	Locomotor activity (Females)	$t_3 = 0.60$	0.6088	3.9	0.32	Small	31 to 32
11	Mating Latency	$t_3 = 0.96$	0.4092	10.4	0.37	Small	30
12	Mating time	$t_3 = 3.67$	0.0349	-7.3	2.35	Large	30

1353
1354 **Table S6.** Summary of all assays in the <u>microbe-free environment (Gen 17-20).</u> T refers to the
1355 grand mean of the MBLs, while C refers to the grand mean of the MBs.

1356

	Assay	Test statistic	P-value	% change (T - C)/C	Cohen's d	Effect size*	Sample size$
1	Dry body weight (Male)	$t_3 = 1.17$	0.3252	2.1	0.38	Small	10
2	Dry body weight (Female)	$t_3 = 3.05$	0.0554	4.8	0.81	Large	10
3	Development time (Males)	$t_3 = 0.80$	0.4833	0.4	0.28	Small	10
4	Development time (Females)	$t_3 = 1.64$	0.2000	0.8	0.82	Large	10
5	Survival to adulthood	$t_3 = 0.96$	0.4086	1.8	0.73	Medium	9 to 10
6	Fecundity (Females)	$t_3 = 0.73$	0.5185	-15.9	0.39	Small	50
7	Desiccation resistance (Males)	$t_3 = 0.10$	0.9246	0.2	0.02	Small	10
8	Desiccation resistance (Females)	$t_3 = 1.32$	0.2797	-4.1	0.39	Small	10
9	Locomotor activity (Males)	$t_3 = 0.83$	0.4670	-3.3	0.25	Small	26 to 32
10	Locomotor activity (Females)	$t_3 = 0.53$	0.6336	-4.8	0.22	Small	30 to 32
11	Mating Latency	$t_3 = 0.10$	0.9239	1.4	0.09	Small	30
12	Mating time	$t_3 = 0.30$	0.7845	-0.5	0.14	Small	30

1357

*Interpretation for effect sizes using Cohen's d: d > 0.8 (Large); 0.8 > d > 0.5 (Medium); < 0.5 (Small)

$Note: See methods for detailed information about sample sizes. For example, the sample size of ten per treatment for body weight indicates ten micro-centrifuge tubes with 20 flies each. Also, this is a sample size per block, i.e., each circle in the plots is an average over this sample size.

1366 **Table S7.** Summary of all assays in the <u>with-microbes environment (Gen 54-57).</u> T refers to the
1367 grand mean of the MBLs while C refers to the grand mean of the MBs.
1368

	Assay	Test statistic	P-value	% change (T - C)/C	Cohen's d	Effect size*	Sample size$
1	Dry body weight (Male)	$t_3 = 0.32$	0.7723	-0.6	0.07	Small	10
2	Dry body weight (Female)	$t_3 = 0.82$	0.4726	-1.8	0.17	Small	10
3	Development time (Males)	$t_3 = 2.62$	0.0788	0.4	0.17	Small	9 to 10
4	Development time (Females)	$t_3 = 0.91$	0.4299	0.2	0.06	Small	10
5	Survival to adulthood	$t_3 = 2.31$	0.1044	-2.7	1.81	Large	9 to 10
6	Fecundity (Females)	$t_3 = 1.75$	0.1791	26.0	1.67	Large	50
7	Desiccation resistance (Males)	$t_3 = 3.16$	0.0510	-18.2	1.62	Large	10
8	Desiccation resistance (Females)	$t_3 = 2.45$	0.0916	-6.1	0.89	Large	10
9	Locomotor activity (Males)	$t_3 = 0.50$	0.6498	6.4	0.37	Small	30 to 32
10	Locomotor activity (Females)	$t_3 = 0.10$	0.9260	1.1	0.07	Small	31 to 32
11	Mating Latency	$t_3 = 0.72$	0.5211	11.1	0.44	Small	34 to 40
12	Mating time	$t_3 = 0.15$	0.8928	-0.4	0.04	Small	34 to 40

1369
1370 **Table S8.** Summary of all assays in the <u>microbe-free environment (Gen 54-57)</u>. T refers to the
1371 grand mean of the MBLs while C refers to the grand mean of the MBs.
1372
1373

	Assay	Test statistic	P-value	% change (T - C)/C	Cohen's d	Effect size*	Sample size$
1	Dry body weight (Male)	$t_3 = 2.89$	0.0628	8.5	1.74	Large	10
2	Dry body weight (Female)	$t_3 = 2.34$	0.1015	7.4	1.35	Large	10
3	Development time (Males)	$t_3 = 1.27$	0.2952	-0.9	0.42	Small	10
4	Development time (Females)	$t_3 = 0.25$	0.8161	0.1	0.05	Small	10
5	Survival to adulthood	$t_3 = 3.72$	0.0338	6.0	0.83	Large	10
6	Fecundity (Females)	$t_3 = 1.20$	0.3161	15.2	0.58	Medium	50
7	Desiccation resistance (Males)	$t_3 = 0.40$	0.7164	1.3	0.11	Small	10
8	Desiccation resistance (Females)	$t_3 = 0.70$	0.5365	-3.6	0.32	Small	10
9	Locomotor activity (Males)	$t_3 = 0.01$	0.9913	-0.1	0.01	Small	27 to 32
10	Locomotor activity (Females)	$t_3 = 2.23$	0.1122	18.6	0.99	Large	32
11	Mating Latency	$t_3 = 0.06$	0.9526	-1.0	0.05	Small	32 to 38
12	Mating time	$t_3 = 2.35$	0.1007	-5.4	1.39	Large	32 to 38

1374

*Interpretation for effect sizes using Cohen's d: d > 0.8 (Large); 0.8 > d > 0.5 (Medium); < 0.5
(Small)
$Note: See methods for detailed information about sample sizes. For example, the sample size of
ten per treatment for body weight indicates ten micro-centrifuge tubes with 20 flies each. Also, this
is a sample size per block, i.e., each circle in the plots is an average over this sample size.
____
1381
1382

1383 **Text S7 – RNA-Seq result details**

1384

1385 **S7.1 No major changes in the gene expression profile of MB_1 vs MBL_1 in the microbe-free**
1386 **environment**

1387

In the microbe-free regime, where both selected and control populations were assayed without
their microbiome, only a few transcripts showed differential expression when analyzed using
DESeq2 (Supplementary Figure S14). Only the significant genes with Log_2(Fold Change) > 1.5 and
adjusted P-value < 0.05 are highlighted in Figure S14. Except for nAChRbeta2, which is a nicotinic
acetylcholine receptor subunit, and CG13617, which codes for cilium assembly protein DZIP1 (zinc
finger transcription factor), the functions of the other genes (CG14625, CG12693) are not known in
Drosophila melanogaster.

1395

**Figure S14. Differentially expressed genes in MBL_1 vs MB_1 in the no-microbe environment
using DESeq2.** The red dots represents the up-regulated genes, while blue dots represent
down-regulated genes in MBL_1. The protocol for the RNA-Seq is described in the method section
of the main text.

1396

1397 **S7.2 List of differentially regulated genes in with-microbe assay environment**

1398

1399 **Table S9. List of <u>up-regulated</u> genes in MBL$_1$ in RNA-Seq (with-microbes environment)**

No.	Gene ID	Log$_2$(FC)	Adjusted P-value	Symbol	Chr	Description
1	FBgn0036948	2.59	5.29E-46	CG7298	3L	-
2	FBgn0041579	2.74	8.56E-32	AttC	2R	Attacin-C
3	FBgn0010388	3.43	4.99E-31	Dro	2R	Drosocin
4	FBgn0034407	2.47	1.91E-10	DptB	2R	Diptericin B
5	FBgn0286723	2.63	2.44E-10	hll	2L	heimdall
6	FBgn0014865	2.28	1.46E-07	Mtk	2R	Metchnikowin
7	FBgn0004240	3.11	1.55E-05	DptA	2R	Diptericin A
8	FBgn0052284	2.28	0.0004	CG32284	3L	-
9	FBgn0034957	3.78	0.0039	Rsph4a	2R	Radial spoke head protein 4a
10	FBgn0034712	1.76	0.0122	Alp8	2R	Alkaline phosphatase 8
11	FBgn0037323	2.37	0.0227	CG2663	3R	-
12	FBgn0012042	1.64	0.0490	AttA	2R	Attacin-A

1400
1401
1402
1403 **Table S10. List of <u>down-regulated</u> genes in MBL$_1$ in RNA-Seq (with-microbes environment)**
1404

No.	Gene ID	Log$_2$(FC)	Adjusted P-value	Symbol	Chr	Description
1	FBgn0033978	-1.53	3.30E-19	Cyp6a23	2R	Cyp6a23
2	FBgn0001225	-3.94	1.75E-13	Hsp26	3L	Heat shock protein 26
3	FBgn0001229	-2.03	1.43E-09	Hsp67Bc	3L	Heat shock gene 67Bc
4	FBgn0013279	-2.37	7.09E-07	Hsp70Bc	3R	Heat-shock-protein-70Bc
5	FBgn0001224	-2.74	5.69E-06	Hsp23	3L	Heat shock protein 23
6	FBgn0039151	-1.84	6.07E-06	CG13607	3R	-
7	FBgn0013276	-2.48	1.55E-05	Hsp70Ab	3R	Heat-shock-protein-70Ab
8	FBgn0051354	-2.28	1.55E-05	Hsp70Bbb	3R	Hsp70Bbb
9	FBti0059721	-4.34	2.33E-05	NA	3R	-
10	FBgn0001230	-3.18	2.48E-05	Hsp68	3R	Heat shock protein 68
11	FBti0019359	-2.89	4.19E-05	NA	3R	-
12	FBgn0013278	-2.50	0.0003	Hsp70Bb	3R	Heat-shock-protein-70Bb
13	FBgn0030769	-1.75	0.0005	CG13012	X	-
14	FBgn0039152	-1.60	0.0005	Root	3R	Rootletin
15	FBgn0036363	-6.29	0.0006	CG10140	3L	-
16	FBgn0013275	-2.24	0.0009	Hsp70Aa	3R	Heat-shock-protein-70Aa
17	FBgn0013277	-2.41	0.0023	Hsp70Ba	3R	Heat-shock-protein-70Ba
18	FBgn0052495	-1.61	0.0078	Gss2	X	Glutathione synthetase 2
19	FBgn0039307	-2.28	0.0079	CR13656	3R	-
20	FBti0020330	-2.39	0.0227	NA	3R	-
21	RR43222_transposable_element	-2.87	0.0300	NA	X	-
22	FBti0059720	-2.94	0.0306	NA	3R	-
23	FBti0019362	-4.05	0.0393	NA	3R	-
24	FBgn0034784	-2.69	0.0476	CG9826	2R	-

1405   **S7.3 GO analysis of differentially regulated genes in the with-microbe assay environment**
1406
1407   To see the biological processes these genes are involved in, we used the ShinyGo tool (version
1408   0.80) (http://bioinformatics.sdstate.edu/go/) for gene ontology enrichment analysis with 'GO
1409   biological process' pathway database. We found that the upregulated genes belong to the pathway
1410   involved in the fly's humoral immune response (Figure 4.21), while down-regulated genes belong
1411   to the stress response pathway (Figure 4.22).
1412

Figure S15. Gene ontology (GO) enrichment analysis for 12 Up-regulated genes.

1413

Figure S16. Gene ontology (GO) enrichment analysis for 24 down-regulated genes.

1414

1415 **Text S8 - Another set of differently expressed genes hints at the peritrophic matrix**
1416 **(PM) as a potential region of interest in the MBLs**
1417
Along with HSPs and AMPs, another set of genes that have an altered expression are chitin-
interacting genes. CG7298 and CG32284 are up-regulated chitin-binding enzymes in MBL1, and
down-regulated CG10140 is a probable chitinase. As the RNA-seq is done on the full body of flies,
it is difficult to predict what this means for the fly. But, the annotation of CG7298 in FlyBase offers a
clue. CG7298 is expressed in the proventriculus, which is part of the fly digestive system. A recent
study supports the notion that proventriculus is the site where gut bacteria stably interact with flies,
possibly through interactions with N-acetyl glucosamine, which is a building block of chitin (Dodge
et al. 2023). Moreover, Drosophila melanogaster CG7298 has an ortholog called peritrophin-48 in
related Drosophila species, which is a component of the peritrophic matrix. The peritrophic matrix
is a chitinous layer that separates food bolus from the gut epithelia. This is also implicated as a
physical barrier that keeps gut bacteria away from the gut tissue (Kuraishi et al. 2011; Charroux
and Royet 2012; Rodgers et al. 2017; Erlandson et al. 2019). Taking this all together, we think this
is the region where chitin-interacting proteins are carrying out various processes that might be of
interest. However, the limited available information about these hits prevents us from drawing any
further conclusions at this point in time. In the future, we plan to confirm all the RNA-seq hits and
do additional experiments that can give us more insights.
1434
In a recent study, Kathedar et al. showed that the nicotinic acetylcholine receptor (nAchR)
regulates gut barrier function in Drosophila through the remodeling of the peritrophic matrix
(Katheder et al. 2023). We have seen that in our no-microbe assay regime, nAchRbeta2 gene
expression showed down-regulation in MBL1. This raises the possibility that the processes of
interest happening at the proventriculus that involve the peritrophic matrix might be mediated
through nAch signaling.
1441
1442

References for supplementary information

Charroux, B., and J. Royet. 2012. Gut-microbiota interactions in non-mammals: What can we learn from Drosophila? Semin. Immunol. 24:17–24.

Dodge, R., E. W. Jones, H. Zhu, B. Obadia, D. J. Martinez, C. Wang, A. Aranda-Díaz, K. Aumiller, Z. Liu, M. Voltolini, E. L. Brodie, K. C. Huang, J. M. Carlson, D. A. Sivak, A. C. Spradling, and W. B. Ludington. 2023. A symbiotic physical niche in Drosophila melanogaster regulates stable association of a multi-species gut microbiota. Nat. Commun. 14:1557. Nature Publishing Group.

Erlandson, M. A., U. Toprak, and D. D. Hegedus. 2019. Role of the peritrophic matrix in insect-pathogen interactions. J. Insect Physiol. 117:103894.

Ives, P. T. 1970. FURTHER GENETIC STUDIES OF THE SOUTH AMHERST POPULATION OF DROSOPHILA MELANOGASTER12. Evolution 24:507–518.

Jehrke, L., F. A. Stewart, A. Droste, and M. Beller. 2018. The impact of genome variation and diet on the metabolic phenotype and microbiome composition of Drosophila melanogaster. Sci. Rep. 8.

Katheder, N. S., K. C. Browder, D. Chang, A. De Maziere, P. Kujala, S. van Dijk, J. Klumperman, T.-C. Lu, H. Li, Z. Lai, D. Sangaraju, and H. Jasper. 2023. Nicotinic acetylcholine receptor signaling maintains epithelial barrier integrity. eLife 12:e86381. eLife Sciences Publications, Ltd.

Kuraishi, T., O. Binggeli, O. Opota, N. Buchon, and B. Lemaitre. 2011. Genetic evidence for a protective role of the peritrophic matrix against intestinal bacterial infection in Drosophila melanogaster. Proc. Natl. Acad. Sci. 108:15966–15971. Proceedings of the National Academy of Sciences.

Lee, H.-Y., S.-H. Lee, J.-H. Lee, W.-J. Lee, and K.-J. Min. 2019. The role of commensal microbes in the lifespan of Drosophila melanogaster. Aging 11:4611–4640.

Pais, I. S., R. S. Valente, M. Sporniak, and L. Teixeira. 2018. Drosophila melanogaster establishes a species-specific mutualistic interaction with stable gut-colonizing bacteria. PLOS Biol. 16:e2005710.

Rodgers, F. H., M. Gendrin, C. A. S. Wyer, and G. K. Christophides. 2017. Microbiota-induced peritrophic matrix regulates midgut homeostasis and prevents systemic infection of malaria vector mosquitoes. PLoS Pathog. 13:e1006391.

Rose, M. R. 1984. Laboratory Evolution of Postponed Senescence in Drosophila melanogaster. Evolution 38:1004–1010. [Society for the Study of Evolution, Wiley].

Sah, P., J. Paul Salve, and S. Dey. 2013. Stabilizing biological populations and metapopulations through Adaptive Limiter Control. J. Theor. Biol. 320:113–123.

Selkrig, J., F. Mohammad, S. H. Ng, J. Y. Chua, T. Tumkaya, J. Ho, Y. N. Chiang, D. Rieger, S. Pettersson, C. Helfrich-Förster, J. Y. Yew, and A. Claridge-Chang. 2018. The Drosophila microbiome has a limited influence on sleep, activity, and courtship behaviors. Sci. Rep. 8.

Sheeba, V., N. A. A. Madhyastha, and A. Joshi. 1998. Oviposition preference for novel versus normal food resources in laboratory populations ofDrosophila melanogaster. J. Biosci. 23:93–100.

Weisburg, W. G., S. M. Barns, D. A. Pelletier, and D. J. Lane. 1991. 16S ribosomal DNA amplification for phylogenetic study. J. Bacteriol. 173:697–703. American Society for Microbiology.