Conserved gene content and unique phylogenetic history characterize the 'bloopergene' underlying *Triturus'* balanced lethal system

Manon de Visser[1,2]*, James France[1,2], Olga Paulouskaya[1,2], Thomas Brown[3,4], Michael Fahrbach[5], Chris van der Ploeg[6,7], Ben Wielstra[1,2]

[1] Institute of Biology Leiden, Faculty of Science, Leiden University, Leiden, NL

[2] Understanding Evolution Group, Naturalis Biodiversity Center, Leiden, NL

[3] Max Planck Institute of Molecular Cell Biology and Genetics, Dresden, DE

[4] Leibniz Institute for Zoo and Wildlife Research, Berlin, DE

[5] Niedernhaller Str. 8/ 2, 74653 Criesbach, DE

[6] Dynamics and Control Group, Department of Mechanical Engineering, Eindhoven University of Technology, Eindhoven, NL

[7] Integrated Vehicle Safety Group, Netherlands Organisation for Applied Scientific Research, Helmond, NL

* Corresponding author

Abstract

In a balanced lethal system, half of the reproductive output always succumbs. A clear example is observed in *Triturus* newts, which contain two distinct versions of chromosome 1. Developmental arrest occurs in *Triturus* embryos with two copies of the same chromosome 1 version, but not in those that possess two different versions, suggesting that each version carries something uniquely vital. We use a target capture approach to investigate over 7,000 nuclear DNA markers across the genus *Triturus*. We sequence the same markers for all the main lineages of the Salamandridae family (to which *Triturus* belongs) to explore the evolutionary history of chromosome 1 versus other chromosomes. We discover dozens of genes that are completely missing from either one or the other version of chromosome 1. We also show that

the gene content of 1A and 1B is remarkably similar across all *Triturus* species, as both chromosome versions have a unique, but consistent set of deleted genes. This suggests that the balanced lethal system evolved rapidly and reached its current form before the radiation of *Triturus*. The tree topology of *Triturus'* chromosome 1 distinctly differs from the rest of the genome. We attribute this to pervasive, ancient hybridization between the ancestor of *Triturus* and other modern European newt lineages. Our findings accentuate the intriguingly complex nature of *Triturus'* chromosome 1 – the 'bloopergene' driving the evolutionarily enigmatic balanced lethal system.

Introduction

A balanced lethal system is an extreme hereditary disease that, in diploid organisms, causes the loss of fifty percent of offspring every generation (Wielstra, 2020, Schwander et al., 2014). A balanced lethal system is characterized by the presence of two versions of a specific chromosome that do not recombine and that each harbor one or more unique, recessive lethal alleles (Wielstra, 2020, Dawson, 1967). As a consequence, individuals heterokaryotypic for this chromosome (i.e., individuals that inherited both versions), possess all genes required for survival and are thus viable (Berdan et al., 2022b, Wielstra, 2020). However, given the laws of Mendelian inheritance, half the offspring in each generation comprise individuals that are homokaryotypic for the chromosome, meaning they possess either one or the other version twice (Mendel, 1865, Dobzhansky and Pavlovsky, 1955). Such individuals miss certain crucial genes, causing them to be inviable. It seems an evolutionary paradox that, despite the huge loss of fitness incurred, balanced lethal systems have evolved repeatedly across the tree of life (Dobzhansky and Pavlovsky, 1955, Dawson, 1967, James, 1965, Steiner, 1956, Lin, 1979, Thien, 1969).

Given that the two chromosome versions involved in a balanced lethal system do not recombine, they should be considered supergenes: physically linked set of genes that are inherited together (Thompson and Jiggins, 2014, Schwander et al., 2014, Hill and Robertson, 1966). Different versions of a supergene generally encode complex, distinct phenotypes that are maintained by balancing selection when each offers a unique advantage (Black and Shuker, 2019, Llaurens et al., 2017). However, suppressed recombination in supergenes – often caused by structural variations such as chromosomal inversions (Thompson and Jiggins, 2014) – also comes with drawbacks. Because purifying selection is weakened under reduced recombination,

supergenes tend to accumulate deleterious alleles and recessive lethality may develop in homokaryotypes over time (Wielstra, 2020, Berdan et al., 2022b). For instance, this has been proposed to have caused the evolution of the recessive lethal mutations embedded in the supergenes found in fire ants (Kay et al., 2022, Wang et al., 2013). Moreover, supergene-bound recessive lethal genes may originate suddenly, due to a disruptive inversion breakpoint. For example, such a breakpoint mutation affects the inversion-based supergene that underlies the complex male mating strategy polymorphism in ruffs and is lethal in homozygous state (Kupper et al., 2016, Hill et al., 2023). Thus, a supergene might instantly qualify as, or gradually turn into, what we call a 'bloopergene': a genetic construction that can boost the rapid evolution of adaptive traits, but likewise is an evolutionary trap because of maladaptive consequences. Following this rationale, a balanced lethal system should be considered the most extreme case of a bloopergene.

The best-known example of a balanced lethal system concerns 'chromosome 1 syndrome' in the salamander genus *Triturus*: the crested and marbled newts (Macgregor and Horner, 1980a, Wallace, 1994, Wielstra, 2020). Over two centuries ago, Rusconi already observed that half of *Triturus* embryos perish while still inside the egg (Rusconi, 1821). From karyotyping studies in the 1980s it became clear that the fundamental cause of this die-off is a balanced lethal system (Macgregor and Horner, 1980b, Horner and Macgregor, 1985). *Triturus* newts possess two versions of chromosome 1, dubbed 1A and 1B, that lack chiasmata and do not recombine along most of the long arm (Macgregor and Horner, 1980b). Only heterokaryotypic (1A1B) individuals survive the balanced lethal system, while homokaryotypic individuals (1A1A and 1B1B) experience developmental arrest (D'Amen et al., 2006, Wallace, 1994, Sessions et al., 1988a, Sims et al., 1984).

All *Triturus* species share chromosome 1 syndrome, indicating that this balanced lethal system originated in their common ancestor and has, remarkably, persisted for at least 24 million years (Wielstra, 2020, Steinfartz et al., 2007). On the one hand, a block-like deletion structure is known to characterize the 1A- and 1B-specific regions in a *Triturus* linkage map, suggesting a rapid origin due to a chromosomal re-arrangement – a 'cytogenetic accident' (France et al., in prep, Sessions et al., 1988a). On the other hand, different *Triturus* species exhibit different Giemsa C-banding patterns in both 1A and 1B, suggesting that 1A and 1B have accumulated differences in genomic content between species after the balanced lethal system became fixed (Callan and Lloyd, 1960, Sims et al., 1984). Thus, determining the degree of any genetic divergence across *Triturus* is required to reconstruct the ancestral constitution

of chromosome 1 and help us better understand the evolutionary origin of the balanced lethal system.

Supergenes are known to arise occasionally through introgressive hybridization (Jay et al., 2018, Stolle et al., 2022) and such a scenario could theoretically kickstart a balanced lethal system by bringing together two distinct versions of a chromosome, each 'pre-loaded' with private deleterious mutations, in a single population (Berdan et al., 2022a). Salamandridae, the salamander family to which *Triturus* belongs, is known to have a complex evolutionary history and bears an extensive signal of introgressive hybridization (Rancilhac et al., 2021b, Veith et al., 2018, Kieren et al., 2018). While the 'cytogenetic accident' hypothesis would predict that 1A and 1B originated in a single ancestral *Triturus* population (France et al., in prep), the possibility that 1A and 1B were brought together into a single population (Berdan et al., 2022b) should not be dismissed out of hand. To test these hypotheses we can compare the evolutionary history of *Triturus'* chromosome 1A and 1B to that of the other chromosomes.

Here, we conduct a genome-wide investigation using target capture data of 148 samples from 16 genera within the Salamandridae family (covering 26 recognized species in total), with the aim of unraveling the evolution of *Triturus'* peculiar chromosome 1. Firstly, we identify presence/absence variation associated with the balanced lethal system in *Triturus* by comparing 1A- and 1B-linked markers to explore how the gene content varies across the different *Triturus* species. Secondly, we investigate the evolutionary history of *Triturus'* chromosome 1 versus that of the rest of the *Triturus* genome by conducting phylogenomic analyses including all main Salamandridae lineages. We interpret our results in the light of existing and new theories regarding the natural origin of balanced lethal systems.

Results

Consistent deletions on 1A and 1B across *Triturus* newts

To test the hypothesis that the 1A- and 1B-specific regions of different *Triturus* species vary in genomic content, we use target enrichment by sequence capture to aim to obtain 7,139 nuclear DNA markers (Wielstra et al., 2019, De Visser et al., in prep-a) from 1A1A, 1A1B and 1B1B embryos belonging to ten *Triturus* species (n=3 for each genotype, for each species, as listed in Table S1). In healthy (1A1B) embryos, 6,785 genes are recovered. We explore which of these genes are missing in either 1A1A embryos (and therefore potentially absent from chromosome 1A) or 1B1B embryos (and therefore potentially absent from chromosome 1B)

for each *Triturus* species. To strengthen our confidence that genes are truly absent from 1A or 1B we determine their position on linkage maps of *Triturus* and the sister genus *Lissotriton* (the smooth newts; France et al., in prep), the chromosome-scale assembly of the distantly related *Pleurodeles* (the sharp-ribbed newt; Brown et al., 2023), and an Oxford Nanopore sequencing-based *Triturus* assembly, scaffolded against the *Pleurodeles* genome, given the knowledge that synteny in newts is high (France et al., in prep). We find that 29 genes are consistently absent *Triturus*-wide in diseased 1A1A embryos (but present in 1A1B and 1B1B embryos; Fig.1 and Table S2) and another 35 genes are consistently absent across *Triturus* in diseased 1B1B embryos (but present in 1A1B and 1A1A embryos; Fig.1 and Table S2).

The 1A and 1B chromosomes are remarkably similar in the different *Triturus* species in terms of gene content: variation in gene-deletions is low between species, and there is no apparent phylogenetic signal for the few gene-deletions that do deviate between species (Fig. 1). This indicates that deletions mainly happened before the radiation of *Triturus* and that only a few genes were lost afterwards. Also, the gene order in the chromosome-scale genome assembly of *Pleurodeles waltl* (Brown et al., 2023), suggests that chromosomes 1A and 1B are characterized by unique, big deletions of DNA stretches that contain dozens of genes and together span ~600 Mbp (in *Pleurodeles*). The genes deleted from 1A- and 1B generally form two consecutive clusters, followed by a minor 'mixed' block consisting of both presence/absence variation markers, as well as markers with distinct copies on A and B (Fig.1 and Table S2). This small 'mixed' block suggests the presence of a relatively small, evolutionary stratum present on both 1A and 1B, adjacent to the 1A-block in the estimated, ancestral constitution of chromosome 1 (although the pattern is notably minor).

Given the conserved synteny across different, distantly related newt genera (France et al., in prep), and given our observation of little variation in 1A- and 1B-linked markers across the entire genus *Triturus*, we can now further explore the evolutionary history of chromosomes 1A and 1B compared to the rest of the *Triturus genome.*

Discordant evolutionary history for *Triturus'* chromosome 1

To test the hypothesis that chromosomes 1A and 1B originated in different evolutionary lineages of newt and were later brought together in a single population through introgressive hybridization, we sequence our c. 7k nuclear DNA markers for all main evolutionary lineages in the Salamandridae family (Table S1). We then build phylogenetic trees for markers from all twelve linkage groups (presumed to correspond to the twelve chromosomes) of *Triturus*

separately (France et al., in prep), and we also build trees for only the recombining section of chromosome 1 (i.e. predominantly the short arm), the 1A-linked markers, and the 1B-linked markers, to explore whether either the 1A- or the 1B-linked markers display an aberrant evolutionary history.

We observe phylogenetic discordance in the 'modern European newts', a group of newts composed of four main lineages: the genera *Triturus*, *Lissotriton* and *Calotriton*, as well as the 'NIO clade', which comprises the genera *Neurergus*, *Ichthyosaura* and *Ommatotriton* (Rancilhac et al., 2021a). The topology for the entirety of chromosome 1, rather than only the 1A-specific or 1B-specific regions, deviates from that of the rest of the genome (Fig. 2 and Figs. S1-S18). Phylogenies for linkage groups 2 through 12 suggest that *Lissotriton* is the sister lineage of *Triturus* and that together these two genera comprise the sister lineage of the NIO clade. On the other hand, both the recombining section of chromosome 1, as well as the separate 1A- and 1B-linked markers, suggest that the phylogenetic positions of the NIO clade and *Lissotriton* are switched, with the NIO clade recovered as the sister lineage of *Triturus* (although for 1B-linked markers the support for *Triturus* + the NIO clade is relatively low). The 1B-linked markers show further deviation still: not *Lissotriton*, but *Calotriton* (more distantly related for all the other linkage groups) is suggested to be sister to the *Triturus* + the NIO clade, albeit with very low support (Fig. 2). Lastly, molecular dating suggests that the radiation of all newt clades involved (thus *Triturus*, *Lissotriton*, the NIO clade, and *Calotriton*, together known as the 'modern European newts') occurred in the distant past and in a brief time window (c. 38-42 million years ago; Figs. S18-S34). Given this topological discordance, we investigate the genomic pattern of introgression in more detail.

A complex history of pervasive introgression in modern European newts

The discordant phylogenetic position of *Triturus* within the modern European newts, based on chromosome 1 versus the rest of the genome, implicates a history of introgression. We test for deviations from a strictly bifurcating evolutionary history in order to identify potential key introgression events, again for markers positioned on the different *Triturus* linkage groups, including 1A, 1B and the recombining section of chromosome 1. Our analyses show that a signal of introgression involving the NIO clade affects not only *Triturus'* chromosome 1, but most other linkage groups as well (Fig. 3). To a lesser extent, introgression involving *Calotriton* is also implicated, which is in line with the observation that – based on mtDNA – this taxon is the sister lineage of *Triturus* (Rancilhac et al., 2021a). Overall, the dominant signal of

introgressive hybridization suggests introgression from the ancestor of the NIO clade into the ancestor of the *Triturus* lineage (Fig. 4).

Discussion

The chromosome underlying the balanced lethal system in *Triturus* has a remarkable evolutionary history. Gene-content-wise, little has changed in 1A and 1B since the balanced lethal system became established in the *Triturus* ancestor. This is surprising, because natural selection would be expected to be less efficient at preventing the further shedding of genes since the system became fixed, as homokaryotypes from that point on would have been destined to die anyway (Berdan et al., 2022b, Wielstra, 2020, Berdan et al., 2019).

We see that the 1A- and 1B-specific regions together span ~600 Mbp in the genome of *Pleurodeles* (Brown et al., 2023), which is considerably smaller than would be expected based on the physical size inferred in karyotyping studies of *Triturus*, estimated to be at least 1.2 Gb (Sims et al., 1984), which is over a third of the size of the human genome. The variation in Giemsa C-banding patterns that is observed between different lineages of the genus presumably reflects the independent accumulation of non-coding DNA (Callan and Lloyd, 1960, Sims et al., 1984). This is not unexpected given that salamander genomes are dominated by transposable elements, which are expected to accumulate in non-recombining regions (Sessions, 2008, Kent et al., 2017, McCartney-Melstad et al., 2016). However, even though the non-recombining long arms of chromosome 1 – comprising 1A and 1B – are physically large, the gene density is evidently low.

The near-identical gene content of 1A and 1B across the genus *Triturus* (in combination with the lack of a phylogenetic signal for the minor variation between species that is observed), supports that the 'bloopergene' driving the *Triturus* balanced lethal system originated *before* the adaptive radiation of *Triturus* occurred. This fits the hypothesis that chromosomes 1A and 1B evolved instantaneously due to a 'cytogenetic accident' (France et al., in prep, Sessions et al., 1988a). But does that preclude that introgressive hybridization played a role in the origin of the balanced lethal system?

Our phylogenomic approach reveals a divergent topology for chromosome 1 compared to the rest of the *Triturus* genome. The entire *Triturus* genome bears the signature of introgression derived from the NIO clade. However, introgressive hybridization seems to have affected chromosome 1 in particular, to the extent that a sister relationship between *Triturus*

and NIO clade is supported in our phylogenetic analysis (rather than *Lissotriton* being the sister lineage of *Triturus* as for the rest of the genome). Yet, the scenario that chromosome 1 reflects the actual evolutionary relationships and was relatively resistant to introgression from *Lissotriton* compared to the rest of the genome can perhaps not be dismissed out of hand. This is because including more molecular markers does not strictly lead to a more accurate species tree (Mirarab et al., 2024, Zhang et al., 2021, Stankowski et al., 2024).

Our findings do not fit the hypothesis that introgression brought together supergenes that evolved in different ancestral populations (Berdan et al., 2022b). The confounding topology was observed not only for 1A-specific regions or for 1B-specific regions of chromosome 1, but for both of them – as well as for the rest of chromosome 1. This further supports that the balanced lethal system arose strictly within the ancestral *Triturus* population – in line with the hypothesis that chromosomes 1A and 1B originated in a cytogenetic accident (France et al., in prep, Sessions et al., 1988a). Still, the aberrant evolution of chromosome 1 in *Triturus* evidently did involve introgressive hybridization.

In conclusion, we reconstruct the ancestral constitution of *Triturus'* chromosome 1 and provide new insights into the evolutionary mechanisms that twisted the *Triturus* genome in a distant past. The bloopergene that drives the balanced lethal system evolved over 24 million years ago within the ancestor of *Triturus*, against a backdrop of pervasive, introgressive hybridization, involving the other lineages that gave rise to the modern European newts.

Methods

Sampling scheme and phenotypic + genotypic classification

To investigate the intergeneric variation of *Triturus'* chromosome 1A and 1B, we obtain embryos from captive populations of all *Triturus* species listed in Table S1 (Wielstra and Arntzen, 2016). We include both the Balkan and Italian clade of *T. carnifex*, which are genetically distinct and presumably represent distinct species (Wielstra et al., 2021). We do not include *T. rudolfi* (Arntzen, 2024), closely related to *T. pygmaeus*, as it was only described after our sampling. Individuals were (pure)bred for an unknown number of generations in captivity, but the original founders of all breeding lines are known to derive from localities away from any (potential) hybrid zones (Wielstra et al., 2014), ensuring interspecific admixture is limited. Husbandry and breeding practices are described in (Fahrbach and Gerlach, 2018).

The two types of homokaryotypic *Triturus* embryos (1A1A and 1B1B) express different phenotypes upon developmental arrest: the so-called "slim-tailed" (1A1A) embryos appear

morphologically similar to healthy heterokaryotypic (1A1B) embryos, whereas the so-called "fat-tailed" embryos (1B1B) show more severe malformations (Sessions et al., 1988b, Vucic et al., 2024, Sims et al., 1984). During the sampling of *Triturus* embryos, embryos are thus classified as being healthy when they developed beyond the critical late tailbud stage, or as being diseased (in one, or the other, phenotypic group) when they experienced developmental arrest during this critical stage ("slim tailed" 1A1A or "fat tailed" 1B1B, Sessions et al., 1988a, Sims et al., 1984). In addition, we confirm this embryonic classification by a more standardized method: genotyping samples using a 1A-linked marker (PLEKHM1), a 1B-linked marker (NAGLU), and a control marker from elsewhere in the genome (CDK) with multiplex PCR (as described in Meilink et al., in prep). In total, we include 90 *Triturus* embryo samples: three individuals per 1A1B, 1A1A, and 1B1B genotype, for ten species/populations.

For building a *Triturus* species tree, we use previously published data of 27 adult *Triturus* newts (Wielstra et al., 2019, Kazilas et al., 2024) and add three *T. carnifex* samples (Wielstra et al., 2021) to account for presumed species-level divergence within this taxon (Table S1). Furthermore, to determine the phylogenetic relationships of 1A and 1B compared to the rest of the *Triturus* genome, we include other newts from the Salamandridae family in our study: we use three samples for closely related lineage (the modern European newts, comprising of the genera *Lissotriton, Ommatotriton, Ichthyosaura, Neurergus* and *Calotriton*) and at least one sample as representative for more distantly related lineages (De Visser et al., in prep-a, Rancilhac et al., 2021a). This means that, next to the *Triturus* samples, we use 28 Salamandridae samples from 15 different genera (Table S1).

Laboratory procedures and pre-processing of sequence data

To obtain genomic data we use "NewtCap": a target enrichment by sequence capture workflow (De Visser et al., in prep-a). In brief, we apply a standard salt-based method for extracting DNA using the Promega Wizard^{TM} Genomic DNA Purification kit (Promega, Madison, WI, USA), followed by library preparation using the NEBNext Ultra^{TM} II FS DNA Library Preparation Kit for Illumina (New England Biolabs, Ipswich, MA, USA). Then, we perform target enrichment using a MyBaits-II kit using sequence-specific, *Triturus*-based RNA probes (Arbor Bioscience, Ann Arbor, MI, USA, product Ref: # 170210-32, for details see; Wielstra et al., 2019, De Visser et al., in prep-a) and Illumina sequencing of the enriched targets (outsourced to Baseclear B.V., Leiden, the Netherlands). An upstream bioinformatics pipeline is used to

check the quality of the reads and to perform clean-up and mapping upstream (De Visser et al., in prep-a). Downstream analyses are described below.

Oxford Nanopore sequencing

To gather as much information about the (likely) genomic position of markers within the DNA of *Triturus*, we perform whole genome sequencing, followed by *de novo* assembly. Liver tissue from a *T. ivanbureshi* individual is flash frozen in liquid nitrogen and stored at -80 °C prior to whole genome sequencing. High molecular weight DNA extraction and DNA sequencing using Nanopore is performed by Future Genomics Technologies. DNA is prepared using the SQK-LSK110 ligation library kit. FLO-MIN106 and FLO-PRO002 (R9.4.1) flow cells are used for sequencing on MinION and PromethION platforms. For basecalling, Guppy v.5.0.17 (Wick et al., 2019) is used with the high-accuracy model (Guppy is available to ONT customers via https://community.nanoporetech.com/). From 14 flowcells, we obtain 91,494,016 reads, with an N50 ranging from 14,7 kb to 27,2 kb per flow cell. Adaptor sequences from the raw reads are trimmed using Porechop (https://github.com/rrwick/Porechop).

Genome assembly, scaffolding and alignment

We assemble the genome of *Triturus de novo* using the Oxford Nanopore Technology reads and scaffold the rather fragmentary output using the *Pleurodeles waltl* genome (56) as a reference to get a chromosome-level assembly. Although *P. waltl* does not suffer from a balanced lethal system, it is the most closely related genus with a published reference genome available and synteny is high (France et al., in prep), so it allows us to estimate the order of genes in *Triturus'* chromosome 1.

We run Shasta v.0.10.0 (Shafin et al., 2020) with arguments: "--Reads.minReadLength 5000" and "--config Nanopore-May2022", using reads > 5 kb (https://github.com/paoloshasta/shasta), followed by removal of any retained haplotigs using purge-dups v.1.2.6 (Guan et al., 2020) to produce a draft *Triturus* genome assembly of 21 Gb in size (https://github.com/dfguan/purge_dups). The assembly contains ~65k contigs, with an N50 of 1.27 Mb. For scaffolding, we run RagTag (Alonge et al., 2022) with the *P. waltl* genome as input reference (https://github.com/malonge/RagTag). This eventually scaffolds ~17.6Gb of the 21Gb contigs assembly into the 12 separate chromosomes. We then use BLAST+ v.2.2.31 (Camacho et al., 2009) to align the genes from the target capture set against our scaffolded

genome assembly and *P. waltl* reference genome with default parameters. Only the genes that have one hit in each of the genomes are considered for further analysis.

Presence/absence variation analysis with PAV-spotter

We use "PAV-spotter" (De Visser et al., in prep-b, https://github.com/Wielstra-Lab/PAVspotter) to search for presence/absence variation in the target capture data of three diseased fat tailed (1B1B), three diseased slim tailed (1A1A), and three healthy/control (1A1B) embryos, for each *Triturus* species/population independently. PAV-spotter uses BAM file information to calculate signal-cross-correlation values that are based on the resemblance of the distribution of reads mapped against reference sequences for a given target. We use the following rationale: an absence of reads for a target across 1B1B samples is indicative of a 1A-linked marker in case reads are mapped for the same target across 1A1A and 1A1B samples, and likewise an absence of reads for a certain target across 1A1A samples denotes a 1B-linked marker when reads are mapped for the same target across 1B1B and 1A1B samples.

We prepare the read depth data for PAV-spotter and we apply the tool in the same way as described in (De Visser et al., in prep-b), using default settings. We use three samples per phenotypic category to correct for any stochasticity and variation in capture-rates between samples. To identify chromosome 1-linked PAV, we used a 80% dissimilarity threshold (see the PAV-spotter manuscript for more details). We confirm the PAV patterns by visually inspecting the read content in the BAM files for targets that are identified as either 1A- or 1B-linked markers using IGV (Integrative Genomics Viewer; Robinson et al., 2011) and we again do this in the same way as described in (De Visser et al., in prep-b).

Reconstructing the ancestral constitution of chromosome 1

After obtaining the list of candidates for 1A-linked and 1B-linked markers, we cross-compare the results for the different *Triturus* species. To determine if targets were true positives, we cross-check the data with 1) a *Triturus* linkage map (France et al., in prep), 2) a *Lissotriton* linkage map (France et al., in prep), 3) the *Pleurodeles* genome assembly (Brown et al., 2023), and 4) the *Triturus* RagTag assemblies/scaffolds (which are based on the *Pleurodeles* genome). In case targets do not show up in expected regions within these comparative genomic datasets, they are to be excluded due to the higher likelihood of them being false positives (approximately 70% of putative 1A- and 1B-linked markers concerns singletons, meaning they

are only discovered in one species. These are also frequently positioned outside the relevant region based on our cross-checking, and are therefore considered 'accidental' hits). As we are aware that synteny across newt genera is high, but not absolute, we follow a rationale in the order of cross-comparisons that is based on the relatedness to *Triturus*, where data of more closely related genera overrule the data of more distantly related genera.

If a potential 1A- or 1B-linked target is found in the region of the *Triturus* linkage map that corresponds to the region of interest on chromosome 1 (between ~0–50 cM on linkage group 1), we consider it a true positive. Next, for any remaining target that is not present in the *Triturus* linkage map, we consider it a true positive if it is present on the relevant region of the *Lissotriton* linkage map (between ~50–100 cM on linkage group 4). For any remaining target at this point, we perform a cross-check with the best blast hits to the less closely related genus *Pleurodeles*. In case the target is recovered with a best blast hit on *Pleurodeles'* scaffold 6 between ~0–500 Mb (which corresponds to the regions of interest in *Triturus* and *Lissotriton*), we include it. Finally, if we have so far still no knowledge on any genomic position, we cross-check against our *Triturus* RagTag assembly. In case we observe that a target is on scaffold 6 of the RagTag assembly between ~0–500 Mb (which corresponds to the regions of interest in *Triturus* and *Lissotriton*), we include it. Any remaining targets are excluded.

Building the *Triturus* phylogeny

To help visualize potential phylogenetic signal in the Presence/Absence Variation matrix, we build a *Triturus* phylogeny. We exclude the entirety of chromosome 1 by restricting the input for the analysis to the 3,728 targets that were placed on linkage groups 2-12 in the *Triturus* linkage map (France et al., in prep). To extract only high quality SNPs, we apply quality filtering by removing heterozygote excess, as well as by; extracting SNPs and discarding INDELs, applying stringent quality filtering options, and removing sites with more than 50% missing data (as described in De Visser et al., in prep-a). Next, we remove invariant sites using an ascertainment bias correction Python script (https://github.com/btmartin721/raxml_ascbias) and convert the filtered VCF to PHYLIP format using the vcf2phylip script, v.2.8 (Ortiz, 2019) to serve as input for RAxML. Then, we run RAxML v.8.2.12 with 100 rapid bootstrap replicates under the ASC_GTRGAMMA model and Lewis ascertainment correction to perform a concatenation analysis and obtain the best-scoring maximum likelihood tree (Stamatakis, 2014), following the methods of (De Visser et al., in prep-a, https://github.com/Wielstra-Lab/NewtCap_bioinformatics). In the end, this phylogeny (Fig. 1 and Fig. S35) was based on in total 19,827 SNPs. We visualized the tree using FigTree v.1.4.4

(http://tree.bio.ed.ac.uk/software/figtree/) and enhanced the coloured version in Figure 1 using iTOL v.6 (Letunic and Bork, 2024).

Estimating Hardy-Weinberg Equilibrium deviations

To search for genes that are potentially still present on both chromosome 1A and 1B across (most of the) *Triturus* species, we search for targets that deviate from Hardy-Weinberg Equilibrium and display excessive heterozygosity explicitly in 1A1B embryos (and not in 1A1A or 1B1B embryos). We create three multi-sample VCF files for *Triturus*; one with the 30 1A1B embryos, one with the 30 1A1A embryos, and one with the 30 1B1B embryos. Then, we use BCFtools v.1.18 (Danecek et al., 2021, Li, 2011) to estimate Hardy-Weinberg and related statistics for every site in each of the VCF file. To take into account potential gene losses, or stochasticity in the data, we allow for a maximum of three individuals to have a missing genotype at any particular site per each VCF file, and we filter as such using the *--max-missing* function of VCFtools (Danecek et al., 2011).

From the data we extract the sites for which heterozygote excess was discovered in the 1A1B set (with a p-value below 0.05), but not in the 1A1A nor in the 1B1B set (i.e., the p-value was above 0.05 in both). As before (see the "The Presence/Absence Variation matrix" section), we check whether the discovered targets fall in the (vicinity of the) heteromorphic *Triturus* chromosome 1 region. Finally, we again perform a visual inspection in IGV (Robinson et al., 2011) to determine if heterozygous SNPs are restricted to 1A1B embryos. Based on a p-value of 0.05, we discover a total of only six targets that show significant heterozygote excess in explicitly 1A1B embryos, pass our IGV inspection, and appear in or near the region(s) of interest (Table S1).

Phylogenetic analyses of Salamandridae

To investigate if the topology of chromosome 1 differs from that of the rest of the genome, we build a species tree of Salamandridae following the same phylogenetic approaches as described above. We perform separate RAxML analyses based on SNPs from chromosome-level sets of targets mapped to different *Triturus* linkage groups: "Chr1" (all 390 targets on *Triturus* linkage group 1, including 1A- and 1B-linked markers), "Chr1*" (a subset of 329 recombining targets on *Triturus'* linkage group 1, thus excluding 1A- or 1B-linked markers), "1A" (the subset of 28 markers that was 1A-linked for all *Triturus* species analyzed), "1B" (the subset of 35

markers that was 1B-linked for all *Triturus* species analyzed), and "Chr2-12" (all 3,728 targets on *Triturus* linkage groups 2-12, see Table S3). We time-calibrate the best-scoring ML trees discovered by RAxML using TreePL (Smith and O'Meara, 2012) using to strict calibration points: 67.49 MYA, for the split between the "New World Newts" and "Old World Newts" and 24 MYA for the most recent common ancestor of *Triturus* (Steinfartz et al., 2007). The trees are visualized in FigTree v.1.4.4 and shown in Figs. S1 – S34. (http://tree.bio.ed.ac.uk/software/figtree/).

Calculating D- and f-statistics

To test the hypothesis that chromosomes 1A and 1B originated in different evolutionary newt lineages and later ended up in a single population through introgressive hybridization, we calculate Patterson's D, also known as 'ABBA-BABA statistic' (Patterson et al., 2012) and related statistics such as the admixture fraction f (also known as the 'f4-ratio) using Dsuite (Malinsky et al., 2021, Patterson et al., 2012). We focus on the Modern European Newts, which comprises; *Triturus*, *Lissotriton*, the NIO clade (*Neurergus*, *Ichthyosaura* & *Ommatotriton*) and *Calotriton*. Calculations are based on the total dataset and on chromosome-level datasets (Table S3).

We conduct the Dsuite analyses for the three different topologies discovered with the phylogenetic analyses earlier: the general species tree based on linkage groups 2-12, with *Lissotriton* being the closest relative to *Triturus*, followed by the NIO clade, then followed by *Calotriton* – named 'TOP1'), the alternate species tree as inferred with targets from chromosome 1 (with the NIO clade being closer related to *Triturus* than sister lineage *Lissotriton* – named 'TOP2'), and the deviant species tree as inferred with targets from only 1B (with both *Calotriton* and the NIO clade being more closely related to *Triturus* than sister lineage *Lissotriton* – named 'TOP3'). The topologies are provided to Dsuite in Newick format. We use *Euproctus* as the outgroup. The Dtrios function of Dsuite is used to calculate the D- and f4 admixture ratio statistics for all potential trio combinations. We apply Dsuite's Fbranch function on trios with significantly elevated D-values to calculate f-branch statistics, which are related to the f4-ratio. Finally, we construct heatmaps using Dsuite as well to visualize the results (Fig. 3, Table S3 and Figs. S36 – S86). We provide the scripts for this in our GitHub repository (https://github.com/Wielstra-Lab/Triturus_chr1_bloopergenes).

Determining the number of gene-flow events and their direction

The software Treemix allows us to infer maximum likelihood trees and details on admixture events from our target capture data (Pickrell and Pritchard, 2012), following the steps described in (Kalaentzis et al., 2024, https://github.com/Wielstra-Lab/banded_newts). As input we use the cleaned-up msVCF files that include the Modern European Newt samples with high quality SNPs. We perform more filtering to ensure files have no missing data using VCFtools v.0.1.16 (Danecek et al., 2011) and we remove any sites that are in linkage disequilibrium using PLINK v.1.9 (Chang et al., 2015). We then run the linkage disequilibrium-pruned data through Treemix, using *Calotriton* as an outgroup with 500 bootstraps, for one through five possible migration edges (m), and for an iteration of ten analyses. Then, we use the R package "OptM" (Fitak, 2021) to determine the optimal value of m using the Treemix output of all iterations (https://cran.r-project.org/package=OptM). Given that the variation in the standard deviations of the input data of our subsetted datasets was too low, OptM was not able to retrieve an optimal value for m for smaller subsets of the data (i.e., when including only markers that belong to a certain linkage group), so we decide to only include the overall target capture dataset. We visualize results using (adjusted versions of) the plotting_funcs.R script of Treemix (Fig. 4) and present the Evanno and Linear method outcomes by using the plot_optM function of OptM (Figs. S87 and S88). Again, we provide the scripts on GitHub (https://github.com/Wielstra-Lab/Triturus_chr1_bloopergenes).

Permit section

Triturus embryos are obtained by M.F. from his personal breeding colony. Housing and breeding salamanders as a private individual are not considered to require a license (BGBl. I S. 3125, 3126, 3750). Housing and breeding protocols comply to EU directive standards (EU directive annex III, section B, Table 9.1) and were reviewed by the Animal Welfare Body Leiden. As per EU legislation regarding the protection of animals used for scientific purposes (EU directive no. 2010/63/EU), sacrificing embryos that are not feeding independently does not qualify as an animal experiment. Sacrificing a *Triturus* individual for the Oxford Nanopore sequencing was approved by the Centrale Commissie Dierproeven (CCD; decision no. AVD1060020198065) after advice of the independent Dierexperimentencommissie Leiden (DEC Leiden).

Acknowledgements

Funding was provided by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (Grant Agreement No. 802759), and T.B. supported by DFG (INST 269/768-1). This work was performed using the compute resources from the Academic Leiden Interdisciplinary Cluster Environment (ALICE) provided by Leiden University. We are grateful for the invaluable feedback provided to us by Professor Roger K. Butlin from the University of Sheffield, and we thank Erik-Jan Bosch for creating the scientific illustrations of the salamanders under the open content License of Naturalis Biodiversity Center (© CC BY-NC-ND 4.0).

Author contributions

B.W., M.d.V., and J.F. conceived and designed the research. M.F. collected newt embryos. M.d.V. and J.F. performed the lab-work and pre-processing of the data. M.d.V. conducted the downstream analyses, with the assistance of O.P., C.v.d.P., T.B., and J.F.. M.d.V., B.W. and J.F. wrote the draft version of the paper, and all authors contributed to revising it.

Competing Interests

The authors declare no competing interests.

Additional information

We will provide the raw, Illumina sequencing reads used in this study publicly through BioProject (PRJNA1173497, PRJNA1171613 and PRJNA498336). Furthermore, the main scripts used for downstream analyses (Dsuite and Treemix) are provided through GitHub (https://github.com/Wielstra-Lab/Triturus_chr1_bloopergenes). All Supplementary Figures and Tables are accessible via Zenodo (https://zenodo.org/records/13991240).

References

ALONGE, M., LEBEIGLE, L., KIRSCHE, M., JENIKE, K., OU, S., AGANEZOV, S., WANG, X., LIPPMAN, Z. B., SCHATZ, M. C. & SOYK, S. 2022. Automated assembly scaffolding using RagTag elevates a new tomato system for high-throughput genome editing. *Genome Biology,* 23**,** 258.

ARNTZEN, J. W. 2024. Morphological and genetic diversification of pygmy and marbled newts, with the description of a new species from the wider Lisbon Peninsula (*Triturus*, Salamandridae). *Contributions to Zoology,* 93**,** 178-200.

BERDAN, E. L., BLANCKAERT, A., BUTLIN, R. K. & BANK, C. 2019. Muller's Ratchet and the Long-Term Fate of Chromosomal Inversions. *bioRxiv***,** 606012.

BERDAN, E. L., BLANCKAERT, A., BUTLIN, R. K., FLATT, T., SLOTTE, T. & WIELSTRA, B. 2022a. Mutation accumulation opposes polymorphism: supergenes and the curious case of balanced lethals. *Philosophical Transactions of the Royal Society B: Biological Sciences,* 377**,** 20210199.

BERDAN, E. L., BLANCKAERT, A., BUTLIN, R. K., FLATT, T., SLOTTE, T. & WIELSTRA, B. 2022b. Mutation accumulation opposes polymorphism: supergenes and the curious case of balanced lethals. *Philosophical Transactions of the Royal Society B,* 377**,** 20210199.

BLACK, D. & SHUKER, D. M. 2019. Supergenes. *Current Biology,* 29**,** R615-R617.

BROWN, T., ELEWA, A., IAROVENKO, S., SUBRAMANIAN, E., ARAUS, A. J., PETZOLD, A., SUZUKI, M., SUZUKI, K.-I. T., HAYASHI, T., TOYODA, A., OLIVEIRA, C., OSIPOVA, E., LEIGH, N. D., SIMON, A. & YUN, M. H. 2023. Sequencing and chromosome-scale assembly of the giant *Pleurodeles waltl* genome. *bioRxiv***,** 2022.10.19.512763.

CALLAN, H. G. & LLOYD, L. 1960. Lampbrush chromosomes of crested newts *Triturus cristatus* (Laurenti). *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences,* 243**,** 135-219.

CAMACHO, C., COULOURIS, G., AVAGYAN, V., MA, N., PAPADOPOULOS, J., BEALER, K. & MADDEN, T. L. 2009. BLAST+: architecture and applications. *BMC Bioinformatics,* 10**,** 421.

CHANG, C. C., CHOW, C. C., TELLIER, L. C., VATTIKUTI, S., PURCELL, S. M. & LEE, J. J. 2015. Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience,* 4**,** 7.

D'AMEN, M., VIGNOLI, L. & BOLOGNA, M. A. 2006. The normal development and the chromosome No. 1 syndrome in *Triturus carnifex carnifex* (Caudata, Salamandridae). *Italian Journal of Zoology,* 73**,** 325-333.

DANECEK, P., AUTON, A., ABECASIS, G., ALBERS, C. A., BANKS, E., DEPRISTO, M. A., HANDSAKER, R. E., LUNTER, G., MARTH, G. T., SHERRY, S. T., MCVEAN, G. & DURBIN, R. 2011. The variant call format and VCFtools. *Bioinformatics,* 27**,** 2156-2158.

DANECEK, P., BONFIELD, J. K., LIDDLE, J., MARSHALL, J., OHAN, V., POLLARD, M. O., WHITWHAM, A., KEANE, T., MCCARTHY, S. A., DAVIES, R. M. & LI, H. 2021. Twelve years of SAMtools and BCFtools. *Gigascience,* 10.

DAWSON, P. S. A. 1967. A Balanced Lethal System in the Flour Beetle *Tribolium castaneum*. *Heredity,* 22**,** 435-438.

DE VISSER, M. C., FRANCE, J., MCCARTNEY-MELSTAD, E., BUCCIARELLI, G. M., THEODOROPOULOS, A., SHAFFER, H. B. & WIELSTRA, B. in prep-a. NewtCap: an efficient target capture approach to boost genomic studies in Salamandridae (True Salamanders and Newts).

DE VISSER, M. C., VAN DER PLOEG, C., CVIJANOVIĆ, M., VUČIĆ, T., IVANOVIĆ, A., THEODOROPOULOS, A. & WIELSTRA, B. in prep-b. PAV-spotter: using signal cross-correlations to identify Presence/Absence Variation in target capture data

DOBZHANSKY, T. & PAVLOVSKY, O. 1955. An Extreme Case of Heterosis in a Central American Population of *Drosophila Tropicalis*. *Proceedings of the National Academy of Sciences,* 41**,** 289-295.

FAHRBACH, M. & GERLACH, U. 2018. The genus *Triturus*: History, Biology, Systematics, Captive Breeding.

FITAK, R. R. 2021. OptM: estimating the optimal number of migration edges on population trees using Treemix. *Biology Methods & Protocols,* 6**,** bpab017.

FRANCE, J., DE VISSER, M., PAULOUSKAYA, O., ARNTZEN, P., BABIK, W., CVIJANOVIĆ, M., IVANOVIĆ, A., SMITH, J. & VUČIĆ, T. in prep. Genomic evidence suggests the balanced lethal system in *Triturus* newts originated in an instantaneous speciation event.

GUAN, D., MCCARTHY, S. A., WOOD, J., HOWE, K., WANG, Y. & DURBIN, R. 2020. Identifying and removing haplotypic duplication in primary genome assemblies. *Bioinformatics,* 36**,** 2896-2898.

HILL, J., ENBODY, E. D., BI, H., LAMICHHANEY, S., LEI, W., CHEN, J., WEI, C., LIU, Y., SCHWOCHOW, D., YOUNIS, S., WIDEMO, F. & ANDERSSON, L. 2023. Low Mutation Load in a Supergene Underpinning Alternative Male Mating Strategies in Ruff (*Calidris pugnax*). *Molecular Biology and Evolution,* 40.

HILL, W. G. & ROBERTSON, A. 1966. The effect of linkage on limits to artificial selection. *Genetics Research,* 8**,** 269-294.

HORNER, H. A. & MACGREGOR, H. C. 1985. Normal development in newts *Triturus* and its arrest as a consequence of an unusual chromosomal situation. *Journal of Herpetology,* 19**,** 261-270.

JAMES, S. H. 1965. Complex hybridity in *Isotoma petraea*. I. The occurrence of interchange heterozygosity, autogamy and a balanced lethal system. *Heredity,* 20**,** 341-353.

JAY, P., WHIBLEY, A., FRÉZAL, L., RODRÍGUEZ DE CARA, M. Á., NOWELL, R. W., MALLET, J., DASMAHAPATRA, K. K. & JORON, M. 2018. Supergene evolution triggered by the introgression of a chromosomal inversion. *Current Biology,* 28**,** 1839-1845.

KALAENTZIS, K., KOSTER, S., ARNTZEN, J. W., BOGAERTS, S., FRANCE, J., FRANZEN, M., KAZILAS, C., LITVINCHUK, S. N., OLGUN, K. & DE VISSER, M. C. 2024. Phylogenomics Resolves the Ancient and Rapid Radiation of Banded Newts (Genus *Ommatotriton*). *[Available as SSRN Preprint]*.

KAY, T., HELLEU, Q. & KELLER, L. 2022. Iterative evolution of supergene-based social polymorphism in ants. *Philosophical Transactions of the Royal Society B,* 377**,** 20210196.

KAZILAS, C., DUFRESNES, C., FRANCE, J., KALAENTZIS, K., MARTINEZ-SOLANO, I., DE VISSER, M. C., ARNTZEN, J. W. & WIELSTRA, B. 2024. Spatial genetic structure in European marbled newts revealed with target enrichment by sequence capture. *Molecular Phylogenetics and Evolution,* 194**,** 108043.

KENT, T. V., UZUNOVIC, J. & WRIGHT, S. I. 2017. Coevolution between transposable elements and recombination. *Philosophical Transactions of the Royal Society B,* 372.

KIEREN, S., SPARREBOOM, M., HOCHKIRCH, A. & VEITH, M. 2018. A biogeographic and ecological perspective to the evolution of reproductive behaviour in the family Salamandridae. *Molecular Phylogenetics and Evolution,* 121**,** 98-109.

KUPPER, C., STOCKS, M., RISSE, J. E., DOS REMEDIOS, N., FARRELL, L. L., MCRAE, S. B., MORGAN, T. C., KARLIONOVA, N., PINCHUK, P., VERKUIL, Y. I., KITAYSKY, A. S., WINGFIELD, J. C., PIERSMA, T., ZENG, K., SLATE, J., BLAXTER, M., LANK, D. B. & BURKE, T. 2016. A supergene determines highly divergent male reproductive morphs in the ruff. *Nature Genetics,* 48**,** 79-83.

LETUNIC, I. & BORK, P. 2024. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic tree display and annotation tool. *Nucleic Acids Research*.

LI, H. 2011. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics,* 27**,** 2987-93.

LIN, Y. J. 1979. Chromosome distribution and catenation in *Rhoeo spathacea* var. *concolor*. *Chromosoma,* 71**,** 109-127.

LLAURENS, V., WHIBLEY, A. & JORON, M. 2017. Genetic architecture and balancing selection: the life and death of differentiated variants. *Molecular Ecology,* 26**,** 2430-2448.

MACGREGOR, H. C. & HORNER, H. 1980a. Heteromorphism for chromosome 1, a requirement for normal development in crested newts. *Chromosoma,* 76**,** 111-122.

MACGREGOR, H. C. & HORNER, H. 1980b. Heteromorphism for chromosome 1, a requirement for normal development in crested newts. *Chromosoma,* 76**,** 111-122.

MALINSKY, M., MATSCHINER, M. & SVARDAL, H. 2021. Dsuite - Fast D-statistics and related admixture evidence from VCF files. *Molecular Ecology Resources,* 21**,** 584-595.

MCCARTNEY-MELSTAD, E., MOUNT, G. G. & SHAFFER, H. B. 2016. Exon capture optimization in amphibians with large genomes. *Molecular ecology resources,* 16**,** 1084-1094.

MEILINK, W. R. M., DE VISSER, M., THEODOROPOULOS, A., FAHRBACH, M. & WIELSTRA, B. in prep. Determining zygosity with multiplex Kompetitive Allele-Specific PCR (mxKASP) genotyping.

MENDEL, G. 1865. Versuche uber pflanzen-hybriden. *Vortlegt in den Sitzungen.*

MIRARAB, S., RIVAS-GONZALEZ, I., FENG, S., STILLER, J., FANG, Q., MAI, U., HICKEY, G., CHEN, G., BRAJUKA, N., FEDRIGO, O., FORMENTI, G., WOLF, J. B. W., HOWE, K., ANTUNES, A., SCHIERUP, M. H., PATEN, B., JARVIS, E. D., ZHANG, G. & BRAUN, E. L. 2024. A region of suppressed recombination misleads neoavian phylogenomics. *Proceedings of the National Academy of Sciences,* 121**,** e2319506121.

ORTIZ, E. M. 2019. vcf2phylip v2. 0: convert a VCF matrix into several matrix formats for phylogenetic analysis.

PATTERSON, N., MOORJANI, P., LUO, Y., MALLICK, S., ROHLAND, N., ZHAN, Y., GENSCHORECK, T., WEBSTER, T. & REICH, D. 2012. Ancient admixture in human history. *Genetics,* 192**,** 1065-93.

PICKRELL, J. K. & PRITCHARD, J. K. 2012. Inference of population splits and mixtures from genome-wide allele frequency data. *PLoS Genetics,* 8**,** e1002967.

RANCILHAC, L., IRISARRI, I., ANGELINI, C., ARNTZEN, J. W., BABIK, W., BOSSUYT, F., KÜNZEL, S., LÜDDECKE, T., PASMANS, F., SANCHEZ, E., WEISROCK, D., VEITH, M., WIELSTRA, B., STEINFARTZ, S., HOFREITER, M., PHILIPPE, H. & VENCES, M. 2021a. Phylotranscriptomic evidence for pervasive ancient hybridization among Old World salamanders. *Molecular Phylogenetics and Evolution,* 155.

RANCILHAC, L., IRISARRI, I., ANGELINI, C., ARNTZEN, J. W., BABIK, W., BOSSUYT, F., KÜNZEL, S., LÜDDECKE, T., PASMANS, F., SANCHEZ, E., WEISROCK, D., VEITH, M., WIELSTRA, B., STEINFARTZ, S., HOFREITER, M., PHILIPPE, H. & VENCES, M. 2021b. Phylotranscriptomic evidence for pervasive ancient hybridization among Old World salamanders. *Molecular Phylogenetics and Evolution,* 155**,** 106967.

ROBINSON, J. T., THORVALDSDOTTIR, H., WINCKLER, W., GUTTMAN, M., LANDER, E. S., GETZ, G. & MESIROV, J. P. 2011. Integrative genomics viewer. *Nature Biotechnology,* 29**,** 24-6.

RUSCONI, M. 1821. *Amours des salamandres aquatiques: et developpement du tetard de ces salamandres depuis l'oeuf jusqu'a l'animal parfait,* Milano, Giusti.

SCHWANDER, T., LIBBRECHT, R. & KELLER, L. 2014. Supergenes and complex phenotypes. *Current Biology,* 24**,** R288-R294.

SESSIONS, S. K. 2008. Evolutionary cytogenetics in salamanders. *Chromosome Research,* 16**,** 183-201.

SESSIONS, S. K., MACGREGOR, H. C., SCHMID, M. & HAAF, T. 1988a. Cytology, embryology, and evolution of the developmental arrest syndrome in newts of the genus *Triturus* (Caudata: Salamandridae). *The Journal of Experimental Zoology,* 248**,** 321-334.

SESSIONS, S. K., MACGREGOR, H. C., SCHMID, M. & HAAF, T. 1988b. Cytology, embryology, and evolution of the developmental arrest syndrome in newts of the genus *Triturus* (Caudata: Salamandridae). *Journal of Experimental Zoology,* 248**,** 321-334.

SHAFIN, K., PESOUT, T., LORIG-ROACH, R., HAUKNESS, M., OLSEN, H. E., BOSWORTH, C., ARMSTRONG, J., TIGYI, K., MAURER, N., KOREN, S., SEDLAZECK, F. J., MARSCHALL, T., MAYES, S., COSTA, V., ZOOK, J. M., LIU, K. J., KILBURN, D., SORENSEN, M., MUNSON, K. M., VOLLGER, M. R., MONLONG, J., GARRISON, E., EICHLER, E. E., SALAMA, S., HAUSSLER, D., GREEN, R. E., AKESON, M., PHILLIPPY, A., MIGA, K. H., CARNEVALI, P., JAIN, M. & PATEN, B. 2020. Nanopore sequencing and the Shasta toolkit enable efficient de novo assembly of eleven human genomes. *Nature Biotechnology,* 38**,** 1044-1053.

SIMS, S. H., MACGREGOR, H. C., PELLATT, P. S. & HORNER, H. A. 1984. Chromosome 1 in crested and marbled newts (*Triturus*) - An extraordinary case of heteromorphism and independent chromosome evolution. *Chromosoma,* 89**,** 169-185.

SMITH, S. A. & O'MEARA, B. C. 2012. treePL: divergence time estimation using penalized likelihood for large phylogenies. *Bioinformatics,* 28**,** 2689-90.

STAMATAKIS, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics,* 30**,** 1312-3.

STANKOWSKI, S., ZAGRODZKA, Z. B., GARLOVSKY, M. D., PAL, A., SHIPILINA, D., CASTILLO, D. G., LIFCHITZ, H., LE MOAN, A., LEDER, E. & REEVE, J. 2024. The genetic basis of a recent transition to live-bearing in marine snails. *Science,* 383**,** 114-119.

STEINER, E. 1956. New Aspects of the Balanced Lethal Mechanism in *Oenothera. Genetics,* 41**,** 486-500.

STEINFARTZ, S., VICARIO, S., ARNTZEN, J. W. & CACCONE, A. 2007. A Bayesian Approach on Molecules and Behavior: Reconsidering Phylogenetic and Evolutionary Patterns of the Salamandridae with Emphasis on *Triturus* Newts. *Journal of experimental zoology. Part B, Molecular and developmental evolution,* 308B**,** 139-162.

STOLLE, E., PRACANA, R., LÓPEZ-OSORIO, F., PRIEBE, M. K., HERNÁNDEZ, G. L., CASTILLO-CARRILLO, C., ARIAS, M. C., PARIS, C. I., BOLLAZZI, M., PRIYAM, A. & WURM, Y. 2022. Recurring adaptive introgression of a supergene variant that determines social organization. *Nature Communications,* 13**,** 1180.

THIEN, L. B. 1969. Chromosome Translocations in *Gayophytum* (Onagraceae). *Evolution,* 23**,** 456-465.

THOMPSON, M. J. & JIGGINS, C. D. 2014. Supergenes and their role in evolution. *Heredity,* 113**,** 1-8.

VEITH, M., BOGAERTS, S., PASMANS, F. & KIEREN, S. 2018. The changing views on the evolutionary relationships of extant Salamandridae (Amphibia: Urodela). *PLoS ONE,* 13**,** 1-16.

VUCIC, T., DROBNJAKOVIC, M., AJDUKOVIC, M., BUGARCIC, M., WIELSTRA, B., IVANOVIC, A. & CVIJANOVIC, M. 2024. A staging table of Balkan crested newt embryonic development to serve as a baseline in evolutionary developmental studies. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*.

WALLACE, H. 1994. The balanced lethal system of crested newts. *Heredity,* 73**,** 41-46.

WANG, J., WURM, Y., NIPITWATTANAPHON, M., RIBA-GROGNUZ, O., HUANG, Y. C., SHOEMAKER, D. & KELLER, L. 2013. A Y-like social chromosome causes alternative colony organization in fire ants. *Nature,* 493**,** 664-8.

WICK, R. R., JUDD, L. M. & HOLT, K. E. 2019. Performance of neural network basecalling tools for Oxford Nanopore sequencing. *Genome Biology,* 20**,** 129.

WIELSTRA, B. 2020. Balanced lethal systems. *Current Biology,* 30**,** R742-R743.

WIELSTRA, B. & ARNTZEN, J. W. 2016. Description of a new species of crested newt, previously subsumed in *Triturus ivanbureschi* (Amphibia: Caudata: Salamandridae). *Zootaxa,* 4109**,** 73-80.

WIELSTRA, B., MCCARTNEY-MELSTAD, E., ARNTZEN, J. W., BUTLIN, R. K. & SHAFFER, H. B. 2019. Phylogenomics of the adaptive radiation of *Triturus* newts supports gradual ecological niche expansion towards an incrementally aquatic lifestyle. *Molecular Phylogenetics and Evolution,* 133**,** 120-127.

WIELSTRA, B., SALVI, D. & CANESTRELLI, D. 2021. Genetic divergence across glacial refugia despite interglacial gene flow in a crested newt. *Evolutionary Biology,* 48**,** 17-26.

WIELSTRA, B., SILLERO, N., VÖRÖS, J. & ARNTZEN, J. W. 2014. The distribution of the crested and marbled newt species (Amphibia: Salamandridae: *Triturus*) – an addition to the New Atlas of Amphibians and Reptiles of Europe. *Amphibia-Reptilia,* 35**,** 376-381.

ZHANG, D., RHEINDT, F. E., SHE, H., CHENG, Y., SONG, G., JIA, C., QU, Y., ALSTROM, P. & LEI, F. 2021. Most Genomic Loci Misrepresent the Phylogeny of an Avian Radiation Because of Ancient Gene Flow. *Systematic Biology,* 70**,** 961-975.

Figures

Fig 1. *Presence/absence variation matrix showing variation in 1A- and 1B-linked markers across the genus* Triturus. *For each* Triturus *species, three individuals with the 1A1A genotype, three with the 1A1B genotype, and three with the 1B1B genotype, were compared. Each column in the matrix represents a recovered 1A-linked (blue square) or 1B-linked marker (orange circle). A gap indicates no presence/absence was detected and a grey colour reflects uncertainty. The matrix is ordered based on the position of the markers in the* Pleurodeles *genome (scaffold 6; location in Mbp indicated at the bottom). From left to right there is a distinct 1B-block containing 40 markers, followed by a distinct 1A-block containing 30 markers, followed by a relatively small 'mixed' block with three 1A- and seven 1B-linked markers. At the far right, three markers are displayed that could not be mapped in* Pleurodeles *('Unclear'). A total of 35 1B-linked and 28 1A-linked markers are consistently found across all* Triturus *species. The* Triturus *phylogeny is based on 19,827 SNPs from 3,728 targets that are not linked to chromosome 1 (see also Fig. S35). [I] = Italy clade, [B] = Balkan clade.*

Fig 2. Topologies for the four clades that comprise the 'modern European newts' for each of the 12 linkage groups, including the recombining region of chromosome 1 (1*) and for chromosomes 1A and 1B, together comprising linkage group 1. Red numbers indicate three cases where the bootstrap support was below 80 (see Figs. S1-S18). The NIO clade is a consistently recovered monophyletic group that contains the three genera Neurergus, Ichthyosaura *and* Ommatotriton.

Fig 3. *F-branch (f_b) heatmaps for different subsets of markers/linkage group(s), showing signals of pervasive introgression in the past among 'modern European newts'. Colour shading reflects the intensity of excess allele-sharing (indicative of introgression) between tree branches on the y-and x-axes. Grey boxes reflect it is not possible to calculate f-branch statistics following the tree topology. Blue, dotted lines correspond to the internal tree branches (e.g. the line above* Lissotriton *and* Triturus *represents their common ancestor). (A) Heatmap based on markers positioned on* Triturus *linkage groups 2 through 12 (n=3,728 targets). (B, C, D, E) Summarized heatmaps for, respectively; 1A-linked markers only (n=28 targets), 1B-linked markers only (n=35 targets), markers positioned on the recombining part of chromosome 1 (n=329 targets), and all markers found on chromosome 1 (n=390 targets). A noticeable, strong signal for introgression between* Triturus *and the NIO clade is visible, except in the analysis including 1A-linked markers only. Abbreviations in figure B-E: T =* Triturus*, L =* Lissotriton*, N = the NIO clade (a consistently recovered monophyletic group that contains the three genera* Neurergus*,* Ichthyosaura *and* Ommatotriton*), and C =* Calotriton*. See also Table S3 and Figs. S36 – S86 for all other heatmaps produced.*

Fig 4. *TreeMix admixture graph for the 'modern European newts' with one migration edge. The arrow shows the inferred direction of the migration event, and the colour of the arrow indicates the migration weight. The axis at the bottom shows the genetic drift measured in the populations.*