Phosphorylation of HP1/Swi6 relieves competition with Suv39/Clr4 on nucleosomes and enables H3K9 trimethyl spreading.

Dana R Kennedy[1,2], Joel Lemiere[3], Catherine Tan[3,4], Eric Simental[1,2], Julian Braxton[5,6], Robert A Maxwell[7], Ahmed AA Amine[1], and Bassem Al-Sady[1*].

[1]Department of Microbiology and Immunology and GW Hooper Foundation, UCSF, [2]TETRAD graduate program, UCSF [3]Biomedical Sciences graduate program, UCSF, [4]Department of Cell and Tissue Biology, UCSF, [5]Chemistry and Chemical Biology graduate program, UCSF, [6]Institute for Neurodegenerative Diseases, UCSF, [7]The Vincent J. Coates Proteomics/Mass Spectrometry Core Laboratory, University of California, Berkeley, CA, USA.

*bassem.al-sady@ucsf.edu

ABSTRACT

Heterochromatin formation in Schizosaccharomyces pombe requires the spreading of histone 3 (H3) Lysine 9 (K9) methylation (me) from nucleation centers by the H3K9 methylase, Suv39/Clr4, and the reader protein, HP1/Swi6. To accomplish this, Suv39/Clr4 and HP1/Swi6 have to associate with nucleosomes both nonspecifically, binding DNA and octamer surfaces and specifically, via recognition of methylated H3K9 by their respective chromodomains. However, how both proteins avoid competition for the same nucleosomes in this process is unclear. Here, we show that phosphorylation tunes the nucleosome affinity of HP1/Swi6 such that it preferentially partitions onto Suv39/Clr4's trimethyl product rather than its unmethylated substrates. Preferential partitioning enables efficient conversion from di-to trimethylation on nucleosomes in vitro and H3K9me3 spreading in vivo. Together, our data suggests that phosphorylation of HP1/Swi6 creates a regime that relieves competition with the "read-write" mechanism of Suv39/Clr4 for productive heterochromatin spreading.

INTRODUCTION

Heterochromatin is a gene-repressive nuclear structure conserved across eukaryotic genomes[1]. Heterochromatin assembly requires seeding at nucleation sites and lateral spreading over varying distances to define a silenced domain[2]. In one highly conserved heterochromatic system, the spreading process requires at least two components: First, a "writer" enzyme, a suppressor of variegation 3-9 methyltransferase homolog (Suv39, Clr4 in *S. pombe*), which deposits Histone 3 lysine 9 methylation (H3K9me)[3]. Spreading by this H3K9 methylation "writer" depends on a positive feedback relationship in which the "writer" also contains a specialized histone-methyl binding chromodomain (CD) that recognizes its own product, H3K9me[4,5]. Second, spreading then further requires a "reader" protein[3,6,7], Heterochromatin Protein 1 (HP1, Swi6 in *S. pombe*), that also recognizes H3K9me2/3 via a CD[8].

How do HP1 proteins execute their essential function in heterochromatin spreading? One manner in which they do so is by directly recruiting the Suv39 methyltransferase to propagate H3K9 methylation[9–11]. Second, HP1 proteins oligomerize on H3K9me-marked chromatin, which has been invoked as a mechanism that supports spreading[12]. HP1 oligomerization also underlies its ability to undergo Liquid-Liquid Phase Separation (LLPS) *in vitro* on its own or with chromatin[13–15], and condensate formation *in vivo*[13,14,16,17]. This condensate formation may promote spreading by providing a specialized nuclear environment that concentrates HP1 and its effectors[18] and/or excludes antagonists of heterochromatin[13]. The silencing of heterochromatin by HP1 may be coupled to spreading by oligomerization, which likely promotes chromatin compaction and blocks RNA polymerase access[19,20]. Silencing may also require oligomerization-independent mechanisms like HP1's ability to bind RNA transcripts and recruit RNA turnover machinery[21,22].

However, these proposed mechanisms for HP1's role in spreading do not contend with a central problem, which is that HP1 and Suv39/Clr4 directly compete for the same substrate on multiple levels. This competition can be specific, as HP1 and Suv39/Clr4 have CDs that recognize the H3K9me2/3 chromatin mark[12,23]. It is also non-specific, as both HP1 and Suv39/Clr4 bind DNA and histone octamer surfaces of the nucleosome substrate[5,17,23–26]. How can HP1 promote H3K9 methylation spreading by Suv39/Clr4 but not get in its way? One explanation for managing the specific competition is an observed difference in methylation state preference. Clr4, for example, is more selective for the terminal trimethylated (H3K9me3) state than Swi6 or the other HP1 paralog in *S. pombe,* Chp2[23]. However, how the significant H3K9me3- independent nucleosome affinity of Clr4 and Swi6 is coordinated to avoid competition is not clear.

One possible way to regulate competition in spreading is through post-translational

modifications of HP1. For example, HP1a, HP1α, and Swi6 are phosphorylated by CKII protein

kinases[27–29]. Phosphorylation of HP1 across species has been shown to regulate multiple of its

biochemical activities including LLPS[13], specificity for H3K9me[13,30], and affinity for nucleic

acids[30]. In *S. pombe*, several Swi6 *in vivo* phosphorylation sites have been documented in the

N-terminal extension (NTE), the CD, and the hinge domain[27], which, when mutated, disrupt

transcriptional gene silencing[27]. While HP1 phosphorylation has been known to be important for

its function for 20 years[31], the mechanisms by which phosphorylation-induced biochemical

changes in HP1 direct its cellular activity and coordination with H3K9 "writers" remain unclear.

In this study, we focused on previously identified Swi6 phosphorylation target sites[27] and found

that two sites in particular, S18 and S24, are required for the spreading, but not nucleation, of

heterochromatin. Spreading defects in Swi6 S18/24A mutants arise inability to convert

H3K9me2 to H3K9me3 outside creation sites. We show biochemically that the primary role of

phosphorylation is to lower Swi6's overall chromatin affinity. This lowered affinity preferentially

partitions Swi6 onto H3K9me3 nucleosomes, rather than unmethylated nucleosomes, *in vitro*

and into heterochromatin foci, rather than the nucleoplasm, *in vivo.* It may appear counter-

intuitive that lowered affinity should have this effect. However, since phosphorylation also

increases Swi6's propensity to oligomerize, this ultimately reduces the Swi6 pool available to

bind unmethylated sites. We propose that phosphorylation of Swi6 frees up Clr4's substrates for

efficient trimethylation, and thus, spreading.

RESULTS

Serines 18 and 24 are necessary for heterochromatin spreading but not nucleation.

Previously, several phosphoserines in Swi6 have been shown to play a role in heterochromatin

gene silencing[27] (**Figure 1A**). To address whether the phosphorylation targets play a role in

nucleation and/or spreading of heterochromatin, we used our mating type locus (MAT)

heterochromatin spreading sensor (HSS[32,33]) (**Figure 1B**). The HSS allows us to separate

nucleation and spreading events at single-cell resolution via three separate transcriptional

reporters: "green" at nucleation sites, "orange" at spreading sites, and "red" in a euchromatic

site to control cell-to-cell noise[32,33]. Specifically, we used a MAT locus HSS with only the *cenH*

nucleator intact (MAT *ΔREIII* HSS[33]), which enables us to isolate spreading from one nucleator.

To query *swi6* serine-to-alanine (S-A) mutants in this background, we first replaced the *swi6*

open reading frame with the *ura4* gene (*swi6::ura4*). Using homologous recombination, we then

replaced the *ura4* cassette with either wild-type or S-A mutant *swi6* open reading frames

followed by a kanamycin resistance marker (**Figure 1B**). We based our S-A mutations on the

phosphoserines previously identified in *Shimada et al.,* which include S18, S24, S46, and S52 in

the NTE, S117 in the CD, and S192, S212, and S220 in the hinge (**Figure 1A**). Here, we

constructed the following S-A mutants: S18A and S24A (*swi6*[S18/24A]); S46A and S52A

(*swi6*[S46/52A]); S46A, S52A, S117A, S192A, S212A, and S220A, (*swi6*[S46/52/117-220A], "S18/S24

available"); and S18A, S24A, S117A, S192A, S212A, and S220A (*swi6*[S18/24/117-220A], "S46/S52

available"). These mutants are expressed at similar levels compared to wild-type as assessed

by western blot, using a polyclonal anti-Swi6 antibody (**Figure 1C,** further validated by

cytometry in **SFigure 4C**). Note that not all phospho-site mutants yield an observable band shift

by SDS-PAGE gel, even though Swi6 in these mutants is expected to retain phosphorylation at

other sites. This was previously observed[27] and is likely because the sequence context of a

phosphorylated residue determines whether or not it will result in a bandshift[34].

When analyzed by flow cytometry, *Δswi6* cells exhibit a silencing defect in which both the

nucleation (green ON) and spreading (orange ON) reporters are expressed (**Figure 1D**).

Conversely, wild-type *swi6* cells show robust silencing of both reporters as we reported prior[33]

(orange 11.1% ON, **Figure 1E**). Mutating only S46 and S52 to alanines (*swi6*[S46/52A]) largely

phenocopies wild-type *swi6* (orange 15% ON, **Figure 1F**). In contrast, mutation of serines at 18

and 24 (*swi6*[S18/24A]) resulted in the loss of spreading (orange 92.4% ON), while largely

maintaining proper nucleation (green off) (**Figure 1G, SFigure 1A-C**). Restoring S18 and S24,

while mutating the other 6 serines to alanines (*swi6*[S46/52/117-220A]) recovers much of the nucleation

and spreading observed in wild-type, though with a modest silencing loss at orange (orange

26.4% ON, **Figure 1H, SFigure 1D-F**). Thus, S18 and S24 play a dominant role in regulating

spreading, while other serines make a minor contribution. However, when only S46 and S52 are

available (*swi6*[S18/24/117-220A]), cells not only exhibit a loss of spreading (orange 94.3% ON) but

also a moderate loss of silencing at the nucleator (green shifted towards ON) (**Figure 1I**). This

loss of silencing approaches but is not as severe as the deletion of *ckb1*, the gene encoding a

crucial regulatory subunit of the CKII kinase. This indicates that while the complete loss of Swi6

phosphorylation disrupts spreading (orange 82-83% ON versus 2.3% in the wild-type control,

SFigure 1K, L), it also affects silencing at the nucleator. However, this defect is not nearly as

severe as in *Δswi6* (**Figure 1D**), highlighting the role of Swi6 phosphorylation primarily in

spreading. Overall, we interpret these results to indicate that NTE S18-52 phosphorylation

contributes to regulating spreading, with S18/24 as major and 46/52 as minor

contributors. Phosphorylation of serines in the Swi6 CD and hinge make a further minor

contribution to Swi6's overall silencing role, which is revealed only in the context of S18/24A.

Given the greater loss of silencing revealed by *Δckb1,* we speculate that there are additional

CKII target residues in Swi6, a notion confirmed by our *in vitro* Mass Spectrometry (see below),

and that their phosphorylation contributes to Swi6's silencing role at the nucleator.

Serines 18 and 24 are required for the spreading of H3K9me3 but not H3K9me2.

We next asked how phosphorylation of S18 and S24 contributes to the propagation of

heterochromatic histone marks. We used chromatin immunoprecipitation followed by

sequencing (ChIP-seq) to address how levels of the heterochromatic marks, Histone 3 lysine 9

di- and trimethylation (H3K9me2/me3), are affected in the context of wild-type *swi6*, *swi6*[S18/24A],

and *Δswi6* in the MAT *ΔREIII* HSS background containing the "green" and "orange" reporters

(**Figure 2A**). Consistent with prior work, we define H3K9me2 as the heterochromatin structural

mark[35] and H3K9me3 as the heterochromatin spreading and silencing mark[7,35,36]. We first

examined the MAT locus. Note that we cannot make definite statements about ChIP-seq signals

over the "green" and "orange" reporters themselves, as the reporter cassettes harbor

sequences that are duplicated 3-4 times in the genome[32], making ChIP-seq read assignment

ambiguous. Overall, H3K9me2 levels at the MAT locus dropped significantly in *Δswi6,*

consistent with prior work[7]; however, *swi6*[S18/24A] mutant maintained similar levels of H3K9me2 to

wild-type *swi6* (**Figure 2B,** top). Examining the distribution more closely, at the *cenH* nucleator,

only *Δswi6* showed a minor decline of H3K9me2 in some regions. To the left of *cenH,* H3K9me2

levels decreased in *Δswi6* but not in *swi6*[S18/24A] . To the right of *cenH,* H3K9me2 levels also

severely declined in *Δswi6,* while in *swi6*[S18/24A] they appear to drop moderately near *mat3M*, but

recovered to wild-type levels at IR-R. When examining H3K9me3, we observed a different

relationship: H3K9me3 patterns in *swi6*[S18/24A] much more closely mirrored *Δswi6*. Specifically, to

the left of *cenH,* H3K9me3 dropped to an intermediate level between wild-type and *Δswi6,* while

on the right of *cenH,* H3K9me3 levels closely matched *Δswi6 (***Figure 2B,** bottom). Importantly,

this behavior of H3K9me3 is consistent with our flow cytometry results (**Figure 1**), where

silencing is largely unaffected at "green" in *swi6*[S18/24A] , while "orange" was expressed.

We wanted to further examine if the observation of H3K9me3 loss in *swi6*[S18/24A] versus wild-type

swi6 held for other genome regions. When we analyzed the subtelomeric region (*tel IIR*) we

found that over the nucleation region *tlh2,* H3K9me2 levels are slightly elevated in *swi6*[S18/24A],

but then begin to drop ~6.4 kb to the left of *tlh2* (**Figure 2C,** top, red bar, and arrow)*.*

Interestingly, H3K9me3 levels drop closer to the *tlh2* nucleator than H3K9me2*;* the 95%

confidence interval of wild-type and *swi6*[S18/24A] separate at the left edge of *tlh2* (**Figure 2C,**

bottom). This observation at the *tlh2* nucleator suggests the conversion of H3K9me2 to

H3K9me3 is inhibited right as heterochromatin structures exit nucleation centers. We observed

the same trend at the left subtelomere of chromosome I (*tel IL*, **SFigure 2B**). At the

subtelomere, spreading distances outside nucleation sites are longer than at other loci, thus this

loss of H3K9me3 just outside *tlh2* has the opportunity to manifest as an H3K9me2 spreading

defect several kilobases downstream. This result is consistent with the requirement of

Suv39/Clr4 methyltransferases to bind H3K9me3 for H3K9 methylation spreading[5,23]. We note

that the left telomere of chromosome II contains no annotated nucleators in the published

sequence. Hence, we could not observe the same trend there (*tel IIL,* **SFigure 2C**).

A similar defect in H3K9me3 spreading also occurs at the pericentromere (*cenII*), specifically,

from the outer repeat (*otr*) into the inner repeat (*imr*) (**Figure 2D**, bottom versus top). However,

the distances are likely too short from nucleation centers in *otr* to observe a resulting loss of

H3K9me2 (**Figure 2D**). We note no distinguishable differences in H3K9me2 and H3K9me3 at

mei4, a well-studied heterochromatin island (**SFigure 1A**).

Together, our ChIP-seq data show that *swi6*[S18/24A] is deficient in the conversion of H3K9me2 to

me3 outside nucleation centers, which results in loss of silencing and ultimately, the loss of

H3K9me2 spreading, as evident for the subtelomere.

Swi6 phosphorylation increases oligomerization and decreases nucleosome affinity.

Next, we wanted to pinpoint the biochemical mechanisms that can account for the spreading

defects in *swi6*[S18/24A] (**Figure 1G, Figure 2**). HP1 oligomerization has been linked to spreading[12].

In turn, HP1's intranuclear dynamics have been linked to how it engages chromatin[37–40]. We

thus probed if and how phosphorylation may impact these two properties of Swi6.

We used Size Exclusion Chromatography followed by Multi-Angle Light Scattering (SEC-MALS)

to probe oligomerization, and fluorescence polarization to quantify H3K9me3 peptide and

nucleosome binding. To produce phosphorylated Swi6 (pSwi6), we co-expressed Swi6 with

Caesin Kinase II (CKII) in *E. coli* (**Figure 3A**). We used 2-dimensional Electron Transfer

Dissociation Mass Spectrometry (2D ETD-MS) to identify which residues in pSwi6 are

phosphorylated and used unphosphorylated Swi6 (unpSwi6) as a control (**Figure 3B**). We found

that only pSwi6, and not unpSwi6, has detectable phosphorylated peptides. The residues

phosphorylated in pSwi6 include several that were identified *in vivo* (**Figure 1A,** S18, S24, S46,

S52, S117, S212, S220 but not S192) and some additional sites not previously identified (S43,

S45, S165, S224, S227). This detection of additional CKII target sites is likely because of the

higher sensitivity achieved in our 2D-ETD-MS experiments from purified protein: 1. 2D-ETD-MS

better preserves phosphorylation sites compared to other methods and is highly sensitive. 2

Pure, *in vitro*-produced protein of high yield is likely to result in more detection events than *in*

vivo-derived protein.

SEC-MALS traces of uncrosslinked pSwi6 and unpSwi6 reveal both proteins are estimated to

be of similar dimer mass, 90.8 kDa and 100.4 kDa respectively (**SFigure 3A**). However, pSwi6

elutes before Swi6, a trend similar to phosphorylated HP1α[13]. There is also a small shoulder in

the pSwi6 trace, indicating a minor fraction of higher-order oligomers (**SFigure 3A**, grey arrow).

As previously published[12], Swi6 crosslinking leads to the appearance of higher molecular weight

species. We observed that crosslinked Swi6 and pSwi6 elute as apparent dimers (93.4 and 86.2

kDa, respectively) and tetramers (210.6 and 180.8 kDa, respectively) (**Figure 3D**). However,

only pSwi6 additionally forms octamers (365.2kDa) and possibly even larger oligomers, as

indicated by a broad shoulder (**Figure 3D**).

We next quantified the binding of pSwi6 to H3K9me0 and H3K9me3 peptides by fluorescence

polarization (**Figure 3D**). pSwi6 binds to H3K9me0 and H3K9me3 peptides with affinities (K_d) of

227.4µM and 2.45µM, respectively, revealing a ~93X specificity for H3K9me3 (**Figure 3F**).

While we could not determine the H3K9me0 peptide K_d for unpSwi6, the K_d for the H3K9me3

peptide was 8.17 µM (**Figure 3D**, **F**). Previously, the specificity for unpSwi6 was reported at

~130X[12], thus indicating little difference in H3K9me3 peptide specificity between the two

proteins. We note that consistent with previous reports on total cellular Swi6[35], recombinant

pSwi6 also shows a ~2.2X preference for H3K9me3 versus H3K9me2 peptides (**SFigure 3E**).

We next probed how phosphorylation affects nucleosome binding. We performed fluorescence

polarization with fluorescently labeled nucleosomes that are unmethylated (H3K9me0) or

trimethylated (H3K$_c$9me3)[41,12]. Phosphorylation had no impact on the specificity for the

H3K9me3 mark, consistent with the peptide observation (19.4X, vs. 19X for unpSwi6 or pSw6,

respectively, **Figure 3E**, **F**).

However, we observe a 12X difference in affinity to the nucleosome overall between pSwi6 and

Swi6 (**Figure 3F**). The H3K$_c$9me3 nucleosome affinity is 0.12 µM and 1.45 µM for unpSwi6 and

pSwi6, respectively, while the H3K9me0 affinity is 2.33 and 27.5 µM, respectively. We note the

affinity of pSwi6 to the H3K$_c$9me3 nucleosome is similar to its affinity to the H3K9me3 peptide,

binding only 1.7X tighter to the H3K$_c$9me3 nucleosome (1.45µM vs. 2.45µM). Instead, and

consistent with previous results, unpSwi6 binds 68X more tightly to the nucleosome than to the

tail (8.17 µM for the H3K9me3 tail versus 0.12µM for H3Kc9me3), which is thought to arise from

additional contacts beyond the H3 tail on the nucleosome.

Why would a 12X lower affinity towards the nucleosome substrate be advantageous for pSwi6's

function in spreading (**Figure 1**,**2**)? In the literature, the cellular abundance of Swi6 is measured

at 9000- 19,400 molecules per cell[42,43]. The estimated fission yeast nuclear volume of ~7mm^3

[44,45] then yields an approximate intranuclear Swi6 concentration of ~2.1 -4.6µM. Given our

measured nucleosome K$_d$s (**Figure 3F**), the intranuclear concentration of unpSwi6 would

theoretically be above its K$_d$ for both H3K9me0 and me3 nucleosomes. The concentration of

pSwi6 would exceed its K$_d$ for H3K$_c$9me3 but be significantly below (~10X) its K$_d$ for H3K9me0

nucleosomes. We cannot assume the same fraction of bound nucleosome from *in vitro*

measurements applies *in vivo*, because nucleosome concentrations in the cell (~10µM based

on accessible genome size and average nucleosome density[46,47]) greatly exceed what is used

in a binding isotherm. We can use a quadratic equation[48] (see methods) appropriate for these *in*

vivo regimes instead of a typical K$_d$ fit to estimate the fraction bound. As only 2% of the *S.*

pombe genome is heterochromatic, we approximate the total nucleosome concentration (10µM)

to reflect unmethylated nucleosomes. The small, methylated nucleosome pool will mostly be

bound by Swi6 irrespective of the phosphorylation state. However, we estimate that only 5% of

unmethylated nucleosomes would be bound by pSwi6, while this would be ~16% for unpSwi6.

At the high end of the Swi6 concentration estimate, this fraction bound would increase to 30% of

unmethylated nucleosomes. Further, we expect enhanced oligomerization of pSwi6 on

heterochromatin to reduce the free Swi6 pool (see discussion). Therefore, we predict that the

main function of phosphorylation is to limit the partitioning of Swi6 into the unmethylated pool,

confining it to heterochromatin.

One test of this prediction would be altered localization of wild-type and phosphorylation

defective Swi6 versions in the fission yeast nucleus. Across species, HP1 homologs have been

shown to localize into heterochromatic foci *in vivo* and form LLPS droplets *in vitro* [13,14,16,37].

Specifically, phosphorylation of the NTE in human HP1α is one driver of heterochromatin foci

formation[13,38]. We investigated whether the loss of phosphorylation sites that impair

heterochromatin spreading (**Figure 1**, **2**) impacted partitioning between heterochromatin foci

and regions outside these foci, likely representing H3K9 unmethylated nucleosomes. We C-

terminally tagged wild-type *swi6* and phospho-serine mutants at the native locus with super fold-

GFP (Swi6-GFP), as an N-terminal tag disrupt Swi6 dimerization and oligomerization[25]. We

crossed these strains into a background containing *sad1*:mKO2, a spindle pole body (SPB)

marker (**SFigure 4A**). We chose this background, as Sad1 denotes the position of

pericentromeric heterochromatin[49,50] and can help orient other heterochromatin sites relative to

it. We examined the following SF-GFP tagged mutant variants: *swi6*$^{S18/24A}$, *swi6*$^{S46/52A}$,

swi6$^{S46/52/117-220A}$ (S18/S24 available), *swi6*$^{S18/24/117-220A}$ (S46/S52 available) (**Figure 1G-I**) and

imaged these strains by confocal microscopy (**Figure 3G**, **SFigure 4B**). Largely, these

mutations do not impact either Swi6 accumulation (**SFigure 4C**), nuclear foci number (**SFigure

4D**), or position of the foci relative to the SPB[51] (**SFigure 4E**, **F**).

We next quantified the accumulation of Swi6-GFP in foci. Unlike foci number or spatial

arrangement, the average foci intensity for Swi6-GFP strains carrying the S18/24A mutations is

significantly decreased relative to wild-type Swi6-GFP (**Figure 3H**), while the nucleoplasmic

signal increases. Because total Swi6-GFP levels do not change in these mutants (**SFigure 4C**),

this result indicates that Swi6$^{S18/24A}$-GFP and Swi6$^{S18/24/117-220A}$-GFP molecules partition away

from heterochromatin foci. This finding is consistent with our prediction based on our *in vitro*

measurements and implies that Swi6 molecules that cannot normally be phosphorylated

partition onto unmethylated nucleosomes.

Swi6 phosphorylation facilitates the conversion of H3K9me2 to me3 by Clr4.

As unmethylated nucleosomes are Clr4's substrates, another prediction emerges. Since

unpSwi6 is more likely to bind unmethylated nucleosomes, Swi6 phosphorylation mutants may

interfere with Clr4 substrates, which could explain the defect in H3K9me2 to me3 conversion in

swi6$^{S18/24A}$ (**Figure 2**), the slowest transition catalyzed by Clr4[23]. For Swi6 phosphorylation to

prevent the conversion of H3K9me to me3, the Swi6 cellular pool would have to be mostly in the

phosphorylated state. To test this, we asked what fraction of Swi6 molecules in the cell are

phosphorylated at S18 and S24. We addressed this question by a quantitative western blot

approach, using two antibodies: a polyclonal Swi6 antibody[25] to detect all Swi6 molecules and a

phospho-serine antibody specific to phosphorylation at S18 and S24 (top blot vs. bottom blot,

respectively, **Figure 4A**). A standard curve of recombinant pSwi6 allowed us to quantify the total

pool of Swi6 molecules vs. those phosphorylated at S18 and S24. The *swi6*[S18/24A] mutant control

shows these phospho-serine antibodies are indeed specific (**Figure 4A**). We showed that the

majority of cellular Swi6 is phosphorylated at S18 and S24 (**Figure 4A** and **SFigure 5A**), 70%

and 100% across two biological replicate experiments.

We next tested if Swi6 phosphorylation directly impacted the ability of Clr4 to produce

H3K9me3. We incubated pSwi6 or unpSwi6 with Clr4 and monitored the conversion of the

H3K9me2 substrate to H3K9me3 under single turnover conditions[23] (**Figure 4B**). pSwi6 shows

an *in vitro* preference for H3K9me3 versus H3K9me2 peptides (**SFigure 3E**), suggesting that

phosphorylation may partition Swi6 towards H3K9me3 versus me0, but also, to some extent,

towards H3K9me3 versus me2.

We observed that the presence of unpSwi6 inhibits the conversion of H3K9me2 to H3K9me3 in

a concentration-dependent manner, but that this inhibition is significantly alleviated by pSwi6

(**Figure 4C** and **SFigure 5B,C**). Note we observe inhibition at the lowest concentration, 5μM,

which is near the estimated *in vivo* concentration of Swi6. When normalizing to H4 and fitting

H3K9me3 to k_{obs}, Clr4 methylation rates are significantly slowed in the presence of unpSwi6,

while pSwi6 reduces this inhibition (**Figure 4D**, **E**).

While these data could explain the H3K9me3-spreading defect, we observe for *swi6*[S18/24A]

(**Figure 2**), our *in vitro*-produced pSwi6 is phosphorylated at multiple residues. Given that S18

and S24 only represent around 1/6 of the detected phosphorylation sites (**Figure 3B**), we

cannot necessarily conclude whether the biochemical phenotypes we observe depend on S18

and S24 phosphorylation. To examine this, we expressed and purified a phospho-mutant

protein, pSwi6[S18/24A], in which S18 and S24 are mutated to alanines and co-expressed it with

CKII. pSwi6[S18/24A] is still phosphorylated to a similar degree as pSwi6, which is apparent by the

similar gel migration shift observed for both proteins (**SFigure 3B**,**C**). Upon phosphatase

treatment, pSwi6[S18/24A] and pSwi6 adopt the same migration pattern as unpSwi6 (**SFigure 3B**,

C). 2D ETD-MS analysis of pSwi6[S18/24A] additionally confirmed a similar phosphopeptide pattern

to pSwi6, though with small changes in phosphopeptide prevalence (**SFigure 3D**).

We examined nucleosome affinity of pSwi6[S18/24A] compared to pSwi6 via fluorescence

polarization and found that pSwi6[S18/24A] shows increased affinity towards both the H3K9me0 and

K_c9me3 nucleosomes, 4.5 and 2.6X, respectively (**Figure 4F**). This result is consistent with S18

and S24 phosphorylation sites acting to modulate Swi6's chromatin affinity. However, since the

change in affinity for pSwi6$^{S18/24A}$ is less than the 12X loss observed for unpSwi6 vs. pSwi6, this

implies that other phosphoserines also contribute to lowering nucleosome affinity.

Overall, this data suggests a model whereby Swi6 NTE phosphorylation, particularly at S18 and

S24, partitions Swi6 away from binding the unmethylated substrate of Clr4 *in vivo,* which is likely

enhanced by increased Swi6 oligomerization at heterochromatin sites. Together, both reduced

affinity and oligomerization mechanisms promote the H3K9me3 spreading reaction.

DISCUSSION

Previous work[27] identified key Swi6 phosphoserines that regulate transcriptional gene silencing.

In this work, we find that Swi6 phosphoserines 18 and 24 are required for heterochromatin

spreading, but not nucleation (**Figure 1**). Swi6 phosphorylation promotes oligomerization, and

tunes Swi6's overall chromatin affinity to a regime that allows Clr4 to access its substrate

(**Figure 3**), facilitating the conversion of dimethyl H3K9 to the repressive and spreading-

promoting trimethyl H3K9 state (**Figure 4**). This modulation of chromatin affinity *in vivo* restricts

Swi6 to heterochromatin foci (**Figure 3, Figure S4**), which suggests that phosphorylation of

HP1 molecules may be required for their concentration into the heterochromatic compartment.

Three central themes emerge from this work:

Swi6 phosphorylation decreases chromatin affinity, but not specificity.

Phosphorylation is known to regulate HP1's affinity with itself[13], DNA[30,38], and chromatin[30,38], but

in manners that are homolog-specific. For example, phosphorylation in the NTE of HP1α

induces LLPS, but not for HP1a in *Drosophila*, where phosphorylation instead regulates

chromatin binding[16,31,52]. Underlying this may be that CKII target sequences are not conserved

across HP1s, for example, HP1α is phosphorylated in a cluster of 4 serines at the NTE (S11-

14)[29], HP1a only at S15 in the NTE, and S202 C-terminal to the CSD[31], whereas we report here

Swi6 is phosphorylated by CKII in the NTE, CD, and hinge (**Figure 3B**).

Phosphorylation increases the affinity towards H3K9me3 and H3K9me0 peptides both for Swi6

(**Figure 3D**) and HP1α[38]. However, the impact on nucleosome specificity is different across

species. Our data here shows that phosphorylation of Swi6 does not affect its specificity for both

H3K9me0 and H3Kc9me3 nucleosomes (**Figure 3E**, **F**), but phosphorylation of HP1α and HP1a

was reported to increase its specificity for H3K9me3 nucleosomes[13,30]. Instead, Swi6

phosphorylation decreases overall nucleosome affinity for unmethylated and H3Kc9me3

nucleosomes to a similar degree, 11.8X and 12X respectively, in contrast to HP1α[30,53]. What

may explain these differences? Internal interactions between the NTE, CD, hinge, and CSD

work together to drive nucleosome binding[17,25]. We speculate that these domain interactions are

differentially impacted by 1. the unique phosphorylation patterns in different HP1 orthologs (see

above) and 2. divergence in Swi6 amino acid sequence and size of the NTE and hinge that

harbor most CKII target sites. Both these differences result in unique outcomes with respect to

nucleosome specificity and affinity in different HP1 orthologs. Cross-linking mass spectrometry

studies indicate that the NTE of HP1α[54], as well as Swi6[17], contact the C-terminus of

H2A.Z/H2A, respectively, H2B, and the core (HP1α) and tail (Swi6) of H3, among other

contacts. NTE phosphorylation may specifically decrease these contacts, leading to detachment

from the nucleosome core.

This overall decrease in affinity partitions pSwi6 in a different way than unpSwi6, restricting

access of pSwi6 to chromatin inside nuclear foci. This is supported by our imaging data (**Figure

3**, **SFigure 4**) but is also consistent with data from human HP1α[38] and *in vivo* diffusion

measurements in the *swi6 sm-1* mutant. This mutant likely disrupts NTE phosphorylation and

shows greater residence outside heterochromatin[39]. Further, it is likely that increased

oligomerization of pSwi6 additionally strengthens this partitioning onto heterochromatin (next

section). A separate consequence of this affinity decrease is the relief of competition with Clr4

for the nucleosome substrate (**Figure 4**, see third section below).

Swi6 phosphorylation increases oligomerization.

Swi6 has been shown to form dimers and higher-order oligomers. Swi6 oligomerization across

chromatin has been linked to heterochromatin spreading *in vivo*[12]. Here, we show that

phosphorylation increases the fraction of oligomeric states, revealing octamers and possibly

higher molecular weight species (**Figure 3**). Swi6 exists in a closed dimer that inhibits the

spreading competent state, or an open dimer that promotes oligomerization[25]. One way pSwi6

could form higher molecular weight oligomers is by phosphorylation shifting the equilibrium from

the closed dimer to the open dimer[25]. We speculate the following thermodynamic consequence

of phosphorylation on the nuclear Swi6 pool: Oligomerization will be driven at sites of high Swi6

accumulation, which is likely near its high-affinity H3K9me3 nucleosome target. If this is true,

oligomerization will reduce the pool of free Swi6 available to engage unmethylated nucleosomes

even further, and below the theoretical level we described above (~5%).

HP1 proteins, like Swi6, form foci *in vivo,* which are associated with condensate formation,

rooted in HP1 oligomerization[13,16,17]. The reduction of GFP-Swi6[S18/24A] in nuclear foci we

observe (**Figure 3**) may be due to defects in condensate formation, or simply that fewer Swi6

molecules are available to form heterochromatic condensates. As discussed above, we expect

defects in phosphorylation to steer Swi6 toward unmethylated chromatin sites. The reduction of

GFP signal in Swi6[S18/24A] mutant foci may thus be due to losing Swi6 molecules to the

nucleoplasmic space.

Phosphorylation of Swi6 enables H3K9 trimethylation by Clr4.

Achieving H3K9 trimethylation is essential for both gene silencing and heterochromatin

spreading by Suv39/Clr4 enzymes[5,23]. For heterochromatin spreading, this is due to the positive

feedback loop within Suv39/Clr4, which depends on binding trimethyl H3K9 tails via the

CD[23,35,55].

For Clr4, the conversion from H3K9me0 to me1 and H3K9me1 to me2 is 10X faster than the

conversion from H3K9me2 to me3[23]. This slow step requires significant residence time on the

nucleosome and is thus highly sensitive to factors promoting or antagonizing Clr4 substrate

access, as well as nucleosome density[56]. Clr4 and Swi6 both make extensive contacts with

nucleosomal DNA and the octamer core[17,24]. unpSwi6 and Clr4 affinity to H3K9me0

nucleosomes are very similar (1.8µM and 2.3µM for Clr4[23] and Swi6, respectively), but nuclear

Swi6 concentration (2-4 µM) is likely higher than the Clr4 concentration[57]. Thus, unpSwi6 would

compete and displace Clr4 from its substrate. However, pSwi6's affinity for the H3K9me0

nucleosome (28µM) is in a regime that is well above its predicted *in vivo* concentration. Any

residual competition between pSwi6 and Clr4 would be mitigated by this lower affinity and the

likely higher affinity of the Clr4 complex to its *in vivo* nucleosome substrate, driven by additional

chromatin modfications[58].

This lowered pSwi6 nucleosome affinity likely relieves the trimethylation inhibition we observe

for unpSwi6 (**Figure 4**). Therefore, we propose that a major outcome of Swi6 phosphorylation is

to clear nucleosome surfaces for Clr4 to access its substrate (**Figure 4G**). An alternative, and

non-exclusive, possibility is that the reduced affinity of pSwi6 to trimethyl nucleosomes may also

limit the ability of Clr4/Suv39 to spread across nucleosomes, which requires the engagement of

its CD[23]. While the K_c9me3 nucleosome K_d of pSwi6 is just below its predicted *in vivo*

concentration, the fraction bound at trimethylated nucleosomes *in vivo* would be expected to

somewhat lower than for unpSwi6. We note, that instead of starting with H3K9me0

nucleosomes, we examined the conversion of H3K9me2 to me3. pSwi6 may have increased

affinity to those H3K9me2 than H3K9me0 substrates[43]. However, it has been shown by us

(**SFigure 3E**) and others[35] that pSwi6 or Swi6 isolated from *S. pombe* cells, which is mostly

phosphorylated (**Figure 4A**), has a preference for H3K9me3 over H3K9me2[35]. This lower

H3K9me2 preference may still help pSwi6 distinguish between binding H3K9me2 versus me3

chromatin *in vivo*, and not just H3K9me0 versus H3K9me3.

Our *in vivo* data (**Figures 1**- **3**) reveals several serines in Swi6 contribute to spreading, but S18

and S24 have a dominant effect. The contribution of other serines is highlighted by 1. A change

in nucleosome affinity in pSwi6$^{S18/24A}$ that is 3-4X less than for unpSwi6 (**Figure 4F and Table**

1), and, 2. The additional phenotype of *swi6*$^{S18-220A}$ in gene silencing compared to *swi6*$^{S18/24A}$

(**Figure 1**, **SFigure 1**). Still, why might these two residues, when mutated, have a strong impact

on heterochromatin spreading? It is possible that phosphorylation of S18 and S24 plays a

disproportional role versus other residues in shifting the Swi6 from the closed to the open state.

Alternatively, it is possible that *in vivo*, phosphorylation at S18 and S24 are involved in the

recruitment of H3K9me3-promoting factors, including Clr3, and also other factors like

Abo1[36,59,60]. Prior work [27] has shown that Clr3 recruitment to heterochromatin is somewhat

compromised in *swi6*$^{S18-117A}$, while the recruitment of the anti-silencing protein Epe1 is

increased. While this loss of Clr3 and gain of Epe1 may be an indirect consequence of

compromised heterochromatin in *swi6*$^{S18-117A}$, it cannot be excluded that phosphorylation at S18

and S24 is necessary to help recruit Clr3 and/or exclude Epe1. This would provide another

mechanism for Swi6 to support trimethylation spreading by Clr4. Whether this is the case

requires further investigation.

Together, we believe that our work resolves a critical problem in heterochromatin biology, which

is how "writers" and "readers" promote heterochromatin spreading if they compete for the same

substrate surfaces. Phosphorylation of Swi6 tunes the partitioning of Swi6 between

unmethylated and methylated nucleosomes *in vivo,* such that Clr4 unmethylated substrates

remain largely unbound. Whether this phosphorylation is regulated temporally, at different

stages of heterochromatin formation, or spatially, at nucleation versus spreading sites, remains

to be investigated.

LIMITATIONS OF THIS STUDY

In this study, we examine how the phosphorylation of Swi6, especially at S18 and S24, impacts

its association with chromatin and interaction with Clr4/Suv39 and, ultimately, heterochromatin

spreading. We connect the loss of H3K9 trimethyl spreading and delocalization from

heterochromatin foci in *swi6*$^{S28/24A}$ mutants *in vivo* to an increased affinity for unmethylated

457 nucleosomes by unpSwi6. However, it is possible that phosphorylation at S18 and S24 is

458 important for the specific recruitment of Swi6 into heterochromatin foci by other factors. Our

459 study does not directly address this possibility. Our biochemical work does not provide direct

460 insight into why S18 and S24 are dominant *in vivo* compared to other phospho-serines. Further,

461 while we show that pSwi6 drives increased oligomerization, we have no direct evidence that

462 increased oligomerization via phosphorylation *in vivo* supports H3K9 trimethyl spreading. This

463 would require uncoupling phosphorylation from oligomerization, which we have not been able to

464 do so far. Our kinetic studies focus on the impact of Swi6 phosphorylation on the conversation

465 of H3K9me2 to me3. While Swi6 does prefer H3K9me3 over me2, we do not know if this

466 preference is sufficient *in vivo* to decrease the residence of pSwi6 on H3K9me2 enough not to

467 inhibit Clr4/Suv39 and support conversion to H3K9me3. Finally, while we show inhibition of this

468 conversion by unpSwi6, our study does not address whether unpSwi6 and Clr4/Suv39 occupy

469 the exact same surfaces on the nucleosome. Theoretically they could co-occupy the

470 nucleosome, and antagonism by unpSwi6 may utilize a mechanism other than

471 displacement/occlusion.

REFERENCES

1. Grewal, S. I. S. The molecular basis of heterochromatin assembly and epigenetic inheritance. *Mol. Cell* **83**, 1767–1785 (2023).

2. Hamali, B., Amine, A. A. A. & Al-Sady, B. Regulation of the heterochromatin spreading reaction by trans-acting factors. *Open Biol.* **13**, 230271 (2023).

3. Elgin, S. C. & Reuter, G. Position-effect variegation, heterochromatin formation, and gene silencing in Drosophila. *Cold Spring Harb Perspect Biol* **5**, a017780 (2013).

4. Zhang, K., Mosch, K., Fischle, W. & Grewal, S. I. S. Roles of the Clr4 methyltransferase complex in nucleation, spreading and maintenance of heterochromatin. *Nat Struct Mol Biol* **15**, 381–8 (2008).

5. Muller, M. M., Fierz, B., Bittova, L., Liszczak, G. & Muir, T. W. A two-state activation mechanism controls the histone methyltransferase Suv39h1. *Nat Chem Biol* **12**, 188–93 (2016).

6. Noma, K. *et al.* RITS acts in cis to promote RNA interference-mediated transcriptional and post-transcriptional silencing. *Nat Genet* **36**, 1174–80 (2004).

7. Hall, I. M. *et al.* Establishment and maintenance of a heterochromatin domain. *Science* **297**, 2232–2237 (2002).

8. Jacobs, S. A. & Khorasanizadeh, S. Structure of HP1 chromodomain bound to a lysine 9-methylated histone H3 tail. *Science* **295**, 2080–3 (2002).

9. Haldar, S., Saini, A., Nanda, J. S., Saini, S. & Singh, J. Role of Swi6/HP1 self-association-mediated recruitment of Clr4/Suv39 in establishment and maintenance of heterochromatin in fission yeast. *J Biol Chem* **286**, 9308–20 (2011).

10. Jenuwein, T. & Allis, C. D. Translating the histone code. *Science* **293**, 1074–80 (2001).

11.   Aagaard, L., Schmid, M., Warburton, P. & Jenuwein, T. Mitotic phosphorylation of

SUV39H1, a novel component of active centromeres, coincides with transient accumulation at

mammalian centromeres. *J. Cell Sci.* **113 (Pt 5)**, 817–829 (2000).

12.   Canzio, D. *et al.* Chromodomain-mediated oligomerization of HP1 suggests a

nucleosome-bridging mechanism for heterochromatin assembly. *Mol. Cell* **41**, 67–81 (2011).

13.   Larson, A. G. *et al.* Liquid droplet formation by HP1α suggests a role for phase

separation in heterochromatin. *Nature* **547**, 236–240 (2017).

14.   Sanulli, S. & J Narlikar, G. Liquid-like interactions in heterochromatin: Implications for

mechanism and regulation. *Curr. Opin. Cell Biol.* **64**, 90–96 (2020).

15.   Keenen, M. M. *et al.* HP1 proteins compact DNA into mechanically and positionally

stable phase separated domains. *eLife* **10**, (2021).

16.   Strom, A. R. *et al.* Phase separation drives heterochromatin domain formation. *Nature*

547, 241–245 (2017).

17.   Sanulli, S. *et al.* HP1 reshapes nucleosome core to promote phase separation of

heterochromatin. *Nature* **575**, 390–394 (2019).

18.   Holla, S. *et al.* Positioning Heterochromatin at the Nuclear Periphery Suppresses Histone

Turnover to Promote Epigenetic Inheritance. *Cell* **180**, 150-164.e15 (2020).

19.   Fischer, T. *et al.* Diverse roles of HP1 proteins in heterochromatin assembly and

functions in fission yeast. *Proc Natl Acad Sci USA* **106**, 8998–9003 (2009).

20.   Verschure, P. J. *et al.* In vivo HP1 targeting causes large-scale chromatin condensation

and enhanced histone lysine methylation. *Mol Cell Biol* **25**, 4552–64 (2005).

21.   Keller, C. *et al.* HP1(Swi6) Mediates the Recognition and Destruction of

Heterochromatic RNA Transcripts. *Mol. Cell* **47**, 215–27 (2012).

22. Motamedi, M. R. *et al.* HP1 proteins form distinct complexes and mediate

heterochromatic gene silencing by nonoverlapping mechanisms. *Mol. Cell* **32**, 778–90 (2008).

23. Al-Sady, B., Madhani, H. D. & Narlikar, G. J. Division of labor between the

chromodomains of HP1 and Suv39 methylase enables coordination of heterochromatin

spread. *Mol. Cell* **51**, 80–91 (2013).

24. Akoury, E. *et al.* Disordered region of H3K9 methyltransferase Clr4 binds the

nucleosome and contributes to its activity. *Nucleic Acids Res* **47**, 6726–6736 (2019).

25. Canzio, D. *et al.* A conformational switch in HP1 releases auto-inhibition to drive

heterochromatin assembly. *Nature* **496**, 377–81 (2013).

26. Shirai, A. *et al.* Impact of nucleic acid and methylated H3K9 binding activities of

Suv39h1 on its heterochromatin assembly. *Elife* **6**, (2017).

27. Shimada, A. *et al.* Phosphorylation of Swi6/HP1 regulates transcriptional gene silencing

at heterochromatin. *Genes Dev* **23**, 18–23 (2009).

28. Eissenberg, J. C., Ge, Y. W. & Hartnett, T. Increased phosphorylation of HP1, a

heterochromatin-associated protein of Drosophila, is correlated with heterochromatin

assembly. *J. Biol. Chem.* **269**, 21315–21321 (1994).

29. LeRoy, G. *et al.* Heterochromatin Protein 1 Is Extensively Decorated with Histone Code-

like Post-translational Modifications*. *Mol. Cell. Proteomics* **8**, 2432–2442 (2009).

30. Nishibuchi, G. *et al.* N-terminal phosphorylation of HP1alpha increases its nucleosome-

binding specificity. *Nucleic Acids Res* **42**, 12498–511 (2014).

31. Zhao, T., Heyduk, T. & Eissenberg, J. C. Phosphorylation site mutations in

heterochromatin protein 1 (HP1) reduce or eliminate silencing activity. *J. Biol. Chem.* **276**,

9512–9518 (2001).

32.   Greenstein, R. A. *et al.* Local chromatin context regulates the genetic requirements of the

heterochromatin spreading reaction. *PLoS Genet.* **18**, e1010201 (2022).

33.   Greenstein, R. A. *et al.* Noncoding RNA-nucleated heterochromatin spreading is

intrinsically labile and requires accessory elements for epigenetic stability. *eLife* **7**, e32948

(2018).

34.   O'Donoghue, L. & Smolenski, A. Analysis of protein phosphorylation using Phos-tag

gels. *J. Proteomics* **259**, 104558 (2022).

35.   Jih, G. *et al.* Unique roles for histone H3K9me states in RNAi and heritable silencing of

transcription. *Nature* (2017) doi:10.1038/nature23267.

36.   Yamada, T., Fischle, W., Sugiyama, T., Allis, C. D. & Grewal, S. I. S. The nucleation

and maintenance of heterochromatin by a histone deacetylase in fission yeast. *Mol. Cell* **20**,

173–85 (2005).

37.   Cheutin, T., Gorski, S. A., May, K. M., Singh, P. B. & Misteli, T. In Vivo Dynamics of

Swi6 in Yeast: Evidence for a Stochastic Model of Heterochromatin. *Mol Cell Biol* **24**, 3157–

3167 (2004).

38.   Hiragami-Hamada, K. *et al.* N-terminal phosphorylation of HP1{alpha} promotes its

chromatin binding. *Mol. Cell. Biol.* **31**, 1186–1200 (2011).

39.   Williams, J. F. *et al.* The condensation of HP1-α/Swi6 imparts nuclear stiffness.

2020.07.02.184127 Preprint at https://doi.org/10.1101/2020.07.02.184127 (2023).

40.   Biswas, S. *et al.* HP1 oligomerization compensates for low-affinity H3K9me recognition

and provides a tunable mechanism for heterochromatin-specific localization. *Sci. Adv.* **8**,

eabk0793 (2022).

41.   Simon, M. D. *et al.* The site-specific installation of methyl-lysine analogs into

recombinant histones. *Cell* **128**, 1003–12 (2007).

42.   Carpy, A. *et al.* Absolute proteome and phosphoproteome dynamics during the cell cycle

of Schizosaccharomyces pombe (Fission Yeast). *Mol. Cell. Proteomics MCP* **13**, 1925–1936

(2014).

43.   Sadaie, M. *et al.* Balance between Distinct HP1 Family Proteins Controls

Heterochromatin Assembly in Fission Yeast. *Mol Cell Biol* **28**, 6973–6988 (2008).

44.   Lemière, J., Real-Calderon, P., Holt, L. J., Fai, T. G. & Chang, F. Control of nuclear size

by osmotic forces in Schizosaccharomyces pombe. *eLife* **11**, e76075 (2022).

45.   Neumann, F. R. & Nurse, P. Nuclear size control in fission yeast. *J Cell Biol* **179**, 593–

600 (2007).

46.   Lantermann, A. B. *et al.* Schizosaccharomyces pombe genome-wide nucleosome

mapping reveals positioning mechanisms distinct from those of Saccharomyces cerevisiae.

Nat Struct Mol Biol **17**, 251-U15 (2010).

47.   Godde, J. S. & Widom, J. CHROMATIN STRUCTURE OF

SCHIZOSACCHAROMYCES-POMBE - A NUCLEOSOME REPEAT LENGTH THAT IS

SHORTER THAN THE CHROMATOSOMAL DNA LENGTH. *J Mol Biol* **226**, 1009–1025

(1992).

48.   Jarmoskaite, I., AlSadhan, I., Vaidyanathan, P. P. & Herschlag, D. How to measure and

evaluate binding affinities. *eLife* **9**, e57264 (2020).

49.   Hou, H. *et al.* Csi1 links centromeres to the nuclear envelope for centromere clustering. *J.*

Cell Biol. **199**, 735–744 (2012).

50.   Barrales, R. R., Forn, M., Georgescu, P. R., Sarkadi, Z. & Braun, S. Control of

heterochromatin localization and silencing by the nuclear membrane protein Lem2. *Genes*

Dev **30**, 133–48 (2016).

51.   Al-Sady, B., Greenstein, R. A., El-Samad, H. J., Braun, S. & Madhani, H. D. Sensitive

and Quantitative Three-Color Protein Imaging in Fission Yeast Using Spectrally Diverse,

Recoded Fluorescent Proteins with Experimentally-Characterized In Vivo Maturation

Kinetics. *PLoS One* **11**, e0159292 (2016).

52.   Zhao, T. & Eissenberg, J. C. Phosphorylation of heterochromatin protein 1 by casein

kinase II is required for efficient heterochromatin binding in Drosophila. *J. Biol. Chem.* **274**,

15095–15100 (1999).

53.   Nishibuchi, G. *et al.* Mitotic phosphorylation of HP1α regulates its cell cycle-dependent

chromatin binding. *J. Biochem. (Tokyo)* **165**, 433–446 (2019).

54.   Sokolova, V. *et al.* Structural mechanism of HP1α-dependent transcriptional repression

and chromatin compaction. *Struct. Lond. Engl. 1993* S0969-2126(24)00382–4 (2024)

doi:10.1016/j.str.2024.09.013.

55.   Murawska, M. *et al.* The histone chaperone FACT facilitates heterochromatin spreading

by regulating histone turnover and H3K9 methylation states. *Cell Rep.* **37**, 109944 (2021).

56.   Cutter DiPiazza, A. R. *et al.* Spreading and epigenetic inheritance of heterochromatin

require a critical density of histone H3 lysine 9 tri-methylation. *Proc. Natl. Acad. Sci. U. S. A.*

118, e2100699118 (2021).

57.   Iglesias, N. *et al.* Native Chromatin Proteomics Reveals a Role for Specific Nucleoporins

in Heterochromatin Organization and Maintenance. *Mol. Cell* **77**, 51-66.e8 (2020).

58.   Stirpe, A. *et al.* SUV39 SET domains mediate crosstalk of heterochromatic histone

marks. *eLife* **10**, e62682 (2021).

59.   Zofall, M., Sandhu, R., Holla, S., Wheeler, D. & Grewal, S. I. S. Histone deacetylation

primes self-propagation of heterochromatin domains to promote epigenetic inheritance. *Nat.*

Struct. Mol. Biol. **29**, 898–909 (2022).

60.   Dong, W. *et al.* Abo1 is required for the H3K9me2 to H3K9me3 transition in

heterochromatin. *Sci. Rep.* **10**, 6055 (2020).

61.   Torres-Garcia, S. *et al.* SpEDIT: A fast and efficient CRISPR/Cas9 method for fission

yeast. *Wellcome Open Res.* **5**, 274 (2020).

62.   Rougemaille, M., Shankar, S., Braun, S., Rowley, M. & Madhani, H. D. Ers1, a rapidly

diverging protein essential for RNA interference-dependent heterochromatic silencing in

Schizosaccharomyces pombe. *J. Biol. Chem.* **283**, 25770–25773 (2008).

63.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinforma. Oxf. Engl.* **30**, 2114–2120 (2014).

64.   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods*

9, 357–359 (2012).

65.   Li, H. *et al.* The sequence alignment/map format and SAMtools. *Bioinformatics* **25**,

2078–2079 (2009).

66.   Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A. & Manke, T. DeepTools: A flexible

platform for exploring deep-sequencing data. *Nucleic Acids Res* **42**, W187–W191 (2014).

67.   Lawrence, M., Gentleman, R. & Carey, V. rtracklayer: an R package for interfacing with

genome browsers. *Bioinforma. Oxf. Engl.* **25**, 1841–1842 (2009).

68.   Hahne, F. & Ivanek, R. Visualizing Genomic Data Using Gviz and Bioconductor.

Methods Mol. Biol. Clifton NJ **1418**, 335–351 (2016).

69.   GenomicRanges. *Bioconductor* http://bioconductor.org/packages/GenomicRanges/.

70.   Edelstein, A. D. *et al.* Advanced methods of microscope control using μManager

software. *J. Biol. Methods* **1**, e10 (2014).

71.   Edelstein, A., Amodaj, N., Hoover, K., Vale, R. & Stuurman, N. Computer control of

microscopes using μManager. *Curr. Protoc. Mol. Biol.* **Chapter 14**, Unit14.20 (2010).

72.   Ershov, D. *et al.* TrackMate 7: integrating state-of-the-art segmentation algorithms into

tracking pipelines. *Nat. Methods* **19**, 829–832 (2022).

73.   Tinevez, J.-Y. *et al.* TrackMate: An open and extensible platform for single-particle

tracking. *Methods San Diego Calif* **115**, 80–90 (2017).

74.   Liu, H., Lin, D. & Yates, J. R. Multidimensional separations for protein/peptide analysis

in the post-genomic era. *BioTechniques* **32**, 898, 900, 902 passim (2002).

75.   Washburn, M. P., Wolters, D. & Yates, J. R. Large-scale analysis of the yeast proteome

by multidimensional protein identification technology. *Nat. Biotechnol.* **19**, 242–247 (2001).

Acknowledgments

We would like to thank Geeta J. Narlikar for the discussion on theory and critical feedback on the manuscript. We thank Sigurd Braun, Daniele Canzio, and Lucy D. Brennan for valuable feedback and discussion. B.A-S. was supported by a National Institutes of Health grant R35GM141888 and a National Science Foundation grant 2113319. D.R.K. and C.T. were supported by a National Science Foundation Graduate Research Fellowships Grant No. 2034836. E.S. was supported by a grant from the Ford Foundation and National Institutes of Health supplement DP2GM123484-01S1. We acknowledge the UCSF PFCC (RRID:SCR_018206) for assistance in generating Flow Cytometry data. The research reported here was supported in part by the DRC Center Grant NIH P30 DK063720. We thank the Southworth lab for access and training to SEC-MALS equipment.

FIGURE LEGENDS

Figure 1: S18 and S24 in Swi6 are required for spreading, but not nucleation of

heterochromatin silencing. A. Overview of the Swi6 protein domain architecture and previously

identified (Shimada *et al.*) *in vivo* phosphorylation sites (green residue numbers). NTE: N-terminal

extension; CD: chromodomain (H3K9me binding); HINGE: unstructured hinge region; CSD:

chromo-shadow domain (dimerization and effector recruitment). **B.** Strategy for production of *swi6*

S-A mutants in the MAT *ΔREIII* HSS reporter background. **C.** Swi6 levels are not affected by S-A

mutations. Total extracts of *swi6* wild-type or indicated mutants were probed with an anti-Swi6

polyclonal antibody. *In vitro* purified Swi6 that was either phosphorylated (pSwi6) or not (unpSwi6)

is run as size controls. Note, not all mutant Swi6 proteins display a band shift even if they retain

phosphosites **D.-I.** 2-D Density hexbin plots examining silencing at nucleation "green" and

spreading "orange" reporter in *Δswi6,* wild-type, and indicated S-A mutants. The yellow box

indicates a "green" and "orange" regime consistent with silencing loss, and the magenta box

indicates a regime consistent with loss of spreading, but not nucleation. The dashed line indicates

the threshold for orange ON and the numbers the fraction of cells above the line.

Figure 2: Conversion from H3K9me2 to H3K9me3 is compromised outside nucleation

centers in S18 and S24 Swi6 mutants. A. Overview of the ChIP-seq experiments. **B-D.** ChIP-

seq signal visualization plots. The solid ChIP/input line for each genotype represents the mean of

three repeats, while the shading represents the 95% confidence interval. **B.** Plots of H3K9me2

(TOP) and H3K9me3 (BOTTOM) ChIP signal over input at the MAT *ΔREIII* HSS mating type locus

for wild-type (black), *swi6*[S18/24A] (blue), and *Δswi6* (gold). Signal over "green" and "orange"

reporters are greyed out, as reads from these reporters map to multiple locations within the

reference sequence, as all reporters contain control elements derived from the *ura4* and *ade6*

genes. **C.** H3K9me2 (TOP) and H3K9me3 (BOTTOM) plots as in A. for subtelomere IIR for wild-

type and *swi6*[S18/24A]. The red bar on the H3K9me2 plots indicates the distance from *tlh2* to where

H3K9me2 levels drop in *swi6*[S18/24A] relative to wild-type. Inset: a zoomed-in view proximal to *tlh2*

is shown for H3K9me2 and me3. The red arrows in the insets indicate the point of separation of

the 95% confidence intervals, which is significantly further telomere proximal for H3K9me3. **D.**

H3K9me2 (TOP) and H3K9me3 (BOTTOM) plots as in A. for centromere II for wild-type and

swi6[S18/24A]. Inset: the left side of the pericentromere.

Figure 3: Swi6 phosphorylation increases oligomerization and decreases nucleosome
binding, without affecting specificity. A. Production of phosphorylated Swi6 (pSwi6) in *E. coli.*
Casein Kinase II (CKII) is co-expressed with Swi6. After lysis and purification, the 6X His tag is
removed from the pSwi6 or unpSwi6 protein. **B.** Mass Spectrometry on pSwi6. Shown is a domain
diagram of Swi6. Phosphorylation sites identified in pSwi6 by 2D-ETD-MS are indicated and
grouped by detection prevalence in the sample. **C.** Size Exclusion Chromatography followed by
Multi-Angle Light Scattering (SEC-MALS) on EDC/NHS cross-linked unpSwi6 (black) and pSwi6
(green). Relative refractive index signals (solid lines, left y-axis) and derived molar masses (lines
over particular species, right y-axis) are shown as a function of the elution volume. [Swi6] was
100μM. **D.** Fluorescence polarization (FP) with fluorescein (star)- labeled H3 tail peptides (1-20)
and pSwi6 (green) or unpSwi6 (black) for H3K9me0 (open circles) and H3K9me3 (filled circles)
is shown. Error bars represent standard deviation. Binding was too low to be fit for unpSwi6 and
H3K9me0 peptides. **E.** FP with H3K9me0 (open circles) or H3K$_c$9me3 (MLA, filled circles)
mononucleosomes. Fluorescein (green star) is attached by a flexible linker at one end of the 147
bp DNA template. For D.&E., the average of three independent fluorescent polarization
experiments for each substrate is shown. Error bars represent standard deviation. **F.** Summary
table of affinities and specificities for D. and E. **G.** Representative maximum projection live
microscopy images of indicated Swi6-GFP / Sad1-mKO2 strains. **H.** Analysis of signal intensity in
Swi6-GFP foci in indicated strains. Wt Swi6, n=242; Swi6[S18/24A], n=251; Swi6[S18/24/117-220A], n=145;
Swi6[S46/52/117-220A], n=192. n, number of foci analyzed.

Figure 4: Swi6 phosphorylation mitigates inhibition of the Clr4-mediated conversion of
H3K9me2 to H3K9me3 A. Most Swi6 molecules in the cell are phosphorylated at S18 and S24.
Quantitative western blots against total Swi6 and phosphorylated Swi6 at S18/S24. A standard
curve of pSwi6 isolated as in Figure 3 is included in both blots. Total protein lysates from wild-
type *swi6* and *swi6*[S18/24A] strains were probed with a polyclonal anti-Swi6 antibody (α-Swi6) or an
antibody raised against a phosphorylated S18/S24 peptide (α-S18P-S24P). α-tubulin was used
as a loading control. One of two independent experiments is shown. L; ladder. **B.** Experimental
scheme to probe the impact of Swi6 on H3K9 trimethylation. **C.** Quantitative western blots on the
time-dependent formation of H3K9me3 from H3K9me2 mononucleosomes in the presence of
pSwi6 or unpSwi6. The same blots were probed with α-H3K9me3 and α-H4 antibodies as a
loading and normalization control. **D.** Single exponential fits of production of H3K9me3 tails over
time for indicated concentrations of unpSwi6 or pSwi6. **E.** plot of k_{obs} vs. [Swi6] (μM). **F.**

Fluorescence polarization with H3K9me0 (open circles) or H3K$_c$9me3 (MLA, filled circles)

mononucleosomes as in Figure 3E., with pSwi6 (green) or pSwi6$^{S18/24A}$ (magenta). Relative K$_d$

values in Table 1. Error bars represent standard deviation. **G.** Model of the impact of pSwi6 on

Clr4 activity. Top: pSwi6 does not engage with K3K9me0 nucleosomes, clearing the substrate for

Clr4, and has reduced interactions with the nucleosome core. Bottom: Swi6 binds H3K9me3 and

me0 nucleosomes, occluding Clr4 access.

Table 1. Relative affinities of pSwi6 and pSwi6$^{S18/24A}$ for H3K9me0 and H3K9me3 nucleosomes.

Table 2. Table of *S. pombe* strains used in this work.

Supporting Figure 1: Additional isolates demonstrating that S18 and S24 in Swi6 are

required for spreading, but not nucleation of heterochromatin silencing. 2-D Density hexbin

plots examining silencing at nucleation "green" and spreading "orange" reporter in the MAT *ΔREIII*

HSS for three additional isolates of **A.-C.** swi6$^{S18/24A}$ mutants, **C.-F.** swi6$^{S46/5/117-220A}$ ("S18/24

available"), and **G.-I.** swi6$^{S46/52A}$. **J-L.** As A.-I. but for the *Δckb1* mutant. An independent wild-type

isolate from the cross is shown alongside 2 *Δckb1* isolates.

Supporting Figure 2: H3K9me2 and H3K9me3 ChIP-seq plots in additional genomic loci in

wild-type or *swi6$^{S18/24A}$* A. H3K9me2 (TOP) and H3K9me3 (BOTTOM) plots as in Figure 2 at

mei4 for wild-type and *swi6$^{S18/24A}$*. **B.** H3K9me2 (TOP) and H3K9me3 (BOTTOM) plots as in

Figure 2 at *tel IL* for wild-type and *swi6$^{S18/24A}$*. **B.** H3K9me2 (TOP) and H3K9me3 (BOTTOM) plots

as in Figure 2 at *tel IIL* for wild-type and *swi6$^{S18/24A}$*.

Supporting Figure 3: Characterization of recombinant pSwi6 A. Size Exclusion

Chromatography followed by Multi-Angle Light Scattering (SEC-MALS) on uncrosslinked

unpSwi6 (black) and pSwi6 (green). Relative refractive index signals (solid lines, left y-axis) and

derived molar masses (lines over particular species, right y-axis) are shown as a function of the

elution volume. A migration shift is apparent in pSwi6, as well as a small shoulder of higher

molecular weight species (arrow). **B.** Calf Intestine Phosphatase (CIP) treatment of Swi6

examined in a 15% SDS-PAGE gel. unpSwi6, pSwi6, or pSwi6$^{S18/24A}$ were treated with (+) or

without (-) CIP or with heat-inactivated CIP (b). **C.** CIP treatment of Swi6 examined in a Phos-Tag

gel as in A. Blots of both gels were probed with an anti-Swi6 polyclonal antibody. **D.** Mass

Spectrometry on pSwi6$^{S18/24A}$. Shown is a domain diagram of Swi6$^{S18/24A}$. Phosphorylation sites

identified in pSwi6 $^{S18/24A}$ by 2D-ETD-MS are indicated and grouped by detection prevalence in
the sample.

Supporting Figure 4: Analysis of Swi6-GFP heterochromatin foci number and spatial
distribution. A. Strategy for production of GFP-tagged *swi6* S-A mutants in the *sad1:mKO2*
background. The wildtype *swi6* or S-A mutant gene from Figure 2A was cut with CRISPR/Cas9,
and the break was repaired with a cassette containing a super-folder GFP, *swi6* 3' sequence
homology, and a HygMX cassette. **B.** Representative maximum projection live microscopy
images of indicated Swi6$^{S46/52A}$-GFP /Sad1-mKO2 compared to the wild-type strain. **C.**
Quantification of Swi6-GFP signals by flow cytometry. The GFP signal of independent wild-type
or S-A mutant isolates compared to GFP- cells as measured by flow cytometry. **D.** Distribution of
nuclear foci in nuclei of indicated strains represented as relative frequency. Wt Swi6-GFP , n=85;
Swi6$^{S18/24A}$-GFP, n=94; Swi6$^{S18/24/117-220A}$-GFP, n=50; Swi6$^{S46/52/117-220A}$-GFP, n=82. **E.** distribution
of Swi6-GFP heterochromatin foci relative to Sad1-mKO2. overview: center-to-center distances
were measured in 3D from the peri-spindle pole body Sad1-mKO2 signal to all Swi6-GFP foci
identified in each nucleus. **F.** relative frequency histogram binning the distribution of Sad1-mKO2
to Swi6-GFP foci distances in indicated strains.

Supporting Figure 5: Additional replicates of Swi6 westerns from cell lysates and
nucleosome trimethylation. A. Independent repeat of α-Swi6 and α-S18P-S24P westerns as
in Figure 4A. **B.** A repeat of quantitative western blots querying time-dependent formation of
H3K9me3 from H3K9me2 mononucleosomes in the presence of pSwi6 or unpSwi6 (0 and 5µM
Swi6). **C.** A repeat of quantitative western blots querying time-dependent formation of H3K9me3
from H3K9me2 mononucleosomes in the presence of pSwi6 or unpSwi6 (15µM and 30µM Swi6).
B and C. Swi6 concentration time courses were collected at the same time; westerns were run on
separate days.

METHODS

Strain construction

To construct wild-type *swi6* and *swi6* phosphoserine mutants, the swi6 open reading frame

(ORF) was first deleted by integrating a *ura4* gene cassette in the MAT HSS background. A

plasmid, pRS316, was constructed containing 5' homology-*swi6* promoter-*swi6 (or swi6* S-A

mutant)- 3' UTR-*kan*-3' genome homology and linearized by PmeI double digest to replace the

ura4 cassette by genomic integration via homologous recombination. After transformation, cells

were plated on YES agar for 24 hours before replica plating on G418 selection plates. For the

Δ*ckb1* mutant, we crossed the deletion strain from our chromatin function library[32] to the MAT

ΔREIII HSS and selected Δ*ckb1* MAT ΔREIII HSS strains by random spore analysis on

HYG+G418 double selection. For Swi6-GFP fusions in the Sad1-mKO2 background, *swi6* wild-

type and *swi6* S-A mutant strains were first crossed with the *sad1::mkO2* strain to remove the

MAT HSS. Next, *swi6* and *swi6* S-A mutant ORFs were C-terminally fused to SF-GFP followed

by a hygromycin resistance marker by CRISPR/Cas9 editing as previously described[61].

Modifications were confirmed by gDNA extraction and PCR amplification of the 5' *swi6* to 3'

genome region downstream from the hygromycin marker. For all strain construction, isolates

were verified by genomic PCR.

Western blot

Proteins were separated on a 15% SDS-Page gel and transferred to a PVDF membrane

(Millipore) for 90 minutes at 100V and 4°C. Membranes were blocked overnight in 1:1 1X PBS:

Intercept PBS Blocking Buffer (LiCor). Next, membranes were incubated with either polyclonal

anti-Swi6 antibody[25] or anti-pSwi6 antibody (Rockland Immunochemicals, this study) diluted

1:1000 in 1:1 1X PBS, 0.2% Tween-20 (PBS-T): Intercept PBS Blocking Buffer overnight at 4°C

on a nutator. Anti-α-tubulin antibody was diluted 1:2000 and used as loading control.

Membranes were washed twice with PBS-T for 10 minutes followed by two washes for 5

minutes before incubation with secondary antibodies. Secondary fluorescent antibodies were

diluted either 1:10000 (anti-rabbit, 680 nm, Cell Signaling Technology 5366P, lot # 14) or 1:5000

(anti-mouse, 800 nm, Li-Cor, D10603-05) and were incubated with the membranes for 45

minutes at RT. Finally, membranes were washed 3 times with PBS-T for 10 minutes and once

with PBS for 10 minutes before imaging on a LiCor Odyssey CLx imager.

HSS Flow cytometry

Strains were struck out of a -80°C freezer onto YES plates. Recovered cells were grown in 200

µL of YES media in a 96-well plate overnight to saturation at 32°C. The next morning, cells were

diluted 1:25 in YES media into mid-log phase and analyzed by flow cytometry on an LSR

Fortessa X50 (BD Biosciences). Fluorescence compensation, data analysis, and plotting in R

were performed as described in Greenstein *et al.* 2022[33].

Chromatin immunoprecipitation followed by sequencing (ChIP -seq) sample collection and

library preparation

Cells were grown in YES media overnight to saturation (32°C, 225RPM shaking). The following

morning cells were diluted to OD 0.03, grown to OD 1, and 300×10^6 were fixed and frozen at -

80°C. Cells were processed for ChIP as described in Canzio *et al.* 2011 with the following

modifications: Three technical replicates were processed for ChIP-seq. After lysis, cells were

bead beat 10 rounds for 1 minute each round with 0.5 mm Zirconia/Silica beads (Cat No.

11079105z). Tubes were chilled on ice for 2 minutes between rounds. Lysates were then spun

down to isolate chromatin. The chromatin pellet was resuspended in 1.5 mL lysis buffer, moved

to a 15 mL Diagenode Bioruptor tube (Cat. No. C01020031) and sonicated with a Diagenode

Bioruptor Pico sonicator for a total of 35 cycles, 30 seconds on/ 30 seconds off, in the presence

of sonication beads (Diagenode, Cat. No. C03070001). Every 10 cycles tubes were vortexed.

Chromatin lysate was spun down for 30 minutes at 14000 RPM and 4°C. The lysate volume

was brought up to 900 µL. 45 µL was taken out to check shearing of the DNA. 40 µL was taken

out for input and kept at RT until the reverse crosslinking step. The remaining ~800 µL was

divided into 2 tubes to incubate with either 2 µL anti-H3K9me2 (Abcam 1120, Lot No. 1009758-

6) or 1 µg anti-H3K9me3 (Diagenode, Cat. No. C15500003 Lot No. 003) overnight on a tube

rotator at 4°C. The next morning, Protein A Dynabeads (Invitrogen, LOT 01102248) and M280

Streptavidin beads (Invitrogen, LOT 2692541) were washed twice with Lysis Buffer without

protease inhibitors. 20 µL Protein A Dynabeads beads were added to each anti-H3K9me2

sample, and 30 µL M-280 Streptavidin beads were added to each anti-H3K9me3 sample.

Beads were incubated with samples for 3 hours on a tube rotator at 4°C, and then washed with

700 µL cold buffers at RT on a tube rotator in the following order: 2X Lysis Buffer for 5 minutes,

2X High Salt Buffer for 5 minutes, 1X Wash Buffer for 5 minutes, and 1X TE (buffer recipes as

in[62]). Samples were incubated with 100 µL elution buffer (50 mM Tris pH 8.0, 10 mM EDTA,

1% SDS) for 20 minutes at 70°C in a ThermoMixer F1.5 (Eppendorf). Input samples were

brought up to 100 µL in TE with a final concentration of 1% SDS. Input and eluted samples were

then incubated overnight in a 65°C water bath with 2.5 µL 2.5 mg/mL Proteinase K (Sigma

Aldrich, Lot 58780500) for reverse crosslinking. Samples were purified with a PCR clean-up kit

(Machery-Nagel) and eluted in 100 µL 10 mM Tris pH 8.0. The quality and size of the DNA were

assessed by 4200 TapeStation instrument (Agilent). Next, libraries were prepared using Index

Primer Set 1 (NEBNext Multiplex Oligos for Illumina, E7335L, Lot 10172541), Ultra II FS DNA

Library Prep Kit for Illumina (E7805L, Lot 10202083). The manufacturer's protocol, "Protocol for

FS DNA Library Prep Kit (E7805, E6177) with Inputs ≤100 ng (NEB)", was used starting with

200 pg of DNA. PCR-enriched adaptor-ligated DNA was cleaned up using NEBNext sample

purification beads (E6178S, Lot 10185312, "1.5. Cleanup of PCR Reaction" in manufacturer's

protocol). Individual adaptor-ligated DNA sample concentrations were quantified using a Qubit 4

Fluorometer (Thermo Fisher), and the quality of the DNA was assessed by a 4200 TapeStation

instrument (Agilent). Libraries were pooled to equimolar quantities and sequenced using a

NextSeq 2000 P2 (400 million clusters) (Chan Zuckerberg Biohub San Francisco) (40bp read

length, paired-end).

ChIP-seq data analysis

Sequencing adaptors were trimmed from raw sequencing reads using Trimmomatic v0.39.

The S. pombe genome was downloaded from NCBI under Genome Assembly ASM294v2. The

MAT locus of chromosome II was edited to our custom HSS MAT locus, and the genome was

indexed using the bowtie2-build function of Bowtie2 v2.5.1[63]. Trimmed sequencing reads were

aligned to the genome using Bowtie2 v2.5.1 with flags [--local --very-sensitive-local --no-unal --

no-mixed --no-discordant --phred33 -I 10 -X 700][64]. Next, the resulting SAM files were converted

to BAM files using SAMtools v1.18[65] view function: -S -b ${base}.sam > ${base}.bam. The

resulting BAM files were further processed by removing low-quality alignments, PCR duplicates,

and multimappers, and retain properly aligned paired-end reads using SAMtools view with the

following flags: -bh -F 3844 -f 3 -q 10 -@ 4. The processed BAM files were then sorted and

indexed (SAMtools). Sorted, indexed BAM files were converted to bigWig coverage tracks using

deepTools v3.5.4[66]: bamCoverage: -b "$bam_file" -o "$filename_without_extension.bw" --

binSize 10 --normalizeUsing CPM --extendReads --exactScaling --samFlagInclude 64 --

effectiveGenomeSize 13000000. BigWig files normalized to input were generated using the

bigwigCompare tool (deepTools). Normalized bigWig files were loaded into R v4.3.0 using

rtracklayer v1.60.1[67] and processed for visualization as in Greenstein *et al.* 2022 with

modifications. The Gviz v1.44.2 (Bioconductor) DataTrack function was used to create a

visualization track of ChIP-seq signal in bigWig files for each genotype[68]. The Bioconductor

GenomicRanges package was used to create a GRanges object to store custom genomic

coordinates defined by a BEDfile[69]. Genomic annotations for signal tracks were created using

the AnnotationTrack (Gviz) function. The GenomeAxisTrack (Gviz) function generated a visual

reference (in megabases) to display the position of genomic annotations and signal tracks.

Finally, the plotTracks (Gviz) function was used to plot the DataTrack, AnnotationTrack, and

GenomeAxisTrack objects for visualization.

Swi6-GFP live cell imaging

Swi6-GFP/Sad1-mKO2 strains were struck out onto fresh YES 225 agar plates and incubated at

32°C for 3-5 days. Colonies were inoculated into liquid YES 225 medium (#2011, Sunrise

Science Production) and grown in an incubator shaker at 30°C, 250 rpm to an OD of 0.2 -0.6.

Cells were placed onto 2% agarose (#16500500, Invitrogen) pads in YES 225, covered with a

coverslip (#2850-22, thickness 1 ½, Corning), and sealed with VALAP for imaging. Cells were

imaged on a Ti-Eclipse inverted microscope (Nikon Instruments) with a spinning-disk confocal

system (Yokogawa CSU-10) and a Borealis illumination system that includes 488nm and 541nm

laser illumination and emission filters 525±25nm and 600±25nm respectively, 60X (NA: 1.4)

objectives, and an EM-CCD camera (Hamamatsu, C9100-13). These components were

controlled with μManager v. 1.41[70,71]. The temperature of the sample was maintained at 30°C by

a black panel cage incubation system (#748–3040, OkoLab). The middle plane of cells was first

imaged in brightfield and then two Z-stacks with a step size of 0.5μm were acquired in spinning-

disk confocal mode with laser illumination 488nm and 541nm (total of 9 imaging planes per

channel). The exposure, laser power, and EM gain for the Z-stacks were respectively 50ms /

1% / 800, and 200ms / 5% / 800. Between 9 and 12 fields of view were acquired per strain.

Image analysis

For each field of view, nuclei were manually cropped using Fiji. Cells containing multiple Sad1-

mK02 foci were discarded from this analysis. For each selected nucleus TrackMate was used to

determine the coordinates in space (X, Y, Z) of Sad1 and every Swi6 focus and their

fluorescence intensity[72,73]. Using a custom script on Jupiter Notebook in Python we then

automatically counted the number of Swi6 foci detected by TrackMate for each nucleus.

Additionally, we automated the calculation of the distance between Swi6 foci and the spindle

pole body by measuring the distance from each Swi6 focus to Sad1 within a given nucleus. For

Swi6 intensity measurements, a region of interest (ROI) outside of each nucleus was

automatically selected to measure the background intensity. This background intensity was then

used to correct Swi6 fluorescence signal by subtracting the average intensity of this ROI for a

given analyzed nucleus. Finally, we used a one-way ANOVA statistical test on Swi6 intensity

signal to determine differences between mutants.

Protein cloning and purification

Wildtype *swi6* open reading frame was cloned by ligation-independent cloning into vector 14B

(QB3 Berkeley Macrolab expression vectors). Vector 14B encodes an N-terminal 6xHis tag

followed by a TEV cleavage sequence. Wildtype Swi6 was expressed in BL21-gold (DE3)

competent cells. To produce Swi6$^{S18/24A}$, a gene block containing S18A/S24A Swi6 was cloned

into vector 14B using Gibson assembly. To isolate pSwi6 and pSwi6$^{S18/S24A}$, the respective

vectors were co-expressed with the catalytic subunits of Caesin Kinase II in pRSFDuet. All three

proteins were grown, harvested, and purified using a protocol adapted from [10] and modified as

follows: Cells were grown at 37°C until OD600 0.5-0.6 and induced with a final concentration of

0.4mM Isopropyl-β-D-thiogalactopyranoside. Induced cells were grown at 18°C overnight.

Harvested cells were resuspended in lysis buffer containing 1X PBS buffer pH 7.3, 300 mM

NaCl, 10% glycerol, 0.1% Igepal CA-630, 7.5 mM Imidazole, 1 mM Beta-Mercaptoethanol

(βME), with protease inhibitors. Resuspended cells were sonicated 2 seconds on /2 seconds off

at 40% output power for three 5-minute cycles. The lysate was centrifuged at 25,000xg for 25

minutes, and the supernatant was collected. Nickel NTA resin was equilibrated with lysis buffer.

The supernatant and resin were incubated for 1-2 hours and washed 3 times with 40 ml of lysis

buffer each time before the protein was eluted with 25 mM HEPES pH 7.5, 100 mM KCl, 10%

glycerol, 400 mM Imidazole, and 1 mM βME. The eluted protein was then dialyzed in TEV

cleavage buffer containing 25 mM HEPES pH 7.5, 100 mM KCl, and 1 mM βME and 6 mg TEV

protease. The following morning 3-6 mg of TEV protease was spiked in for about 1 hour to

ensure full cleavage. Nickel NTA resin was equilibrated with TEV cleavage buffer and the his-

tagged TEV was captured by the resin while Swi6 protein was isolated by gravity flow. Cleaved

protein was concentrated using a 10kDa MWCO concentrator and applied to a Superdex 200

Increase 10/300 GL size exclusion column equilibrated in storage buffer containing 25 mM

HEPES pH 7.5, 100 mM KCl, 10% glycerol, and 10 mM βME. Protein was concentrated, flash-

frozen in N_2 (liq), and stored at -80°C. Protein concentration was quantified against a BSA

standard curve on an SDS page gel and sypro red stain.

EDC/NHS crosslinking

unpSwi6 or pSwi6 was purified as described above. However, the storage buffer was 25 mM

HEPES pH 7.5 and 100 mM KCl for SEC-MALS. Protein, either 100 µM Swi6 or pSwi6, was

incubated with 2 mM EDC and 5 mM NHS in a total volume of 95 µL for 2 hours. The reaction

was quenched with a final concentration of 20 mM hydroxylamine.

Size-exclusion Chromatography coupled with Multi-Angle Light Scattering (SEC-MALS)

Crosslinked and uncrosslinked Swi6 and pSwi6 were filtered with 0.2 µm spin columns (Pall

Corporation, Ref. ODM02C34). For SEC, uncrosslinked and crosslinked proteins were injected

onto a KW-804 silica gel chromatography column (Shodex) in a volume of 50 µL at 100 µM. The

column was run using an ÄKTA pure FPLC (GE Healthcare Life Sciences) and equilibrated with

SEC-MALS storage buffer at a flow rate of 0.4 mL/min and temperature of 8°C. The SEC

column was connected in-line to a DAWN HELEOS II (Wyatt Technology) 18-angle light

scattering instrument and an Optilab T-rEX differential refractive index detector (Wyatt

Technology). Data was analyzed using ASTRA software (version 7.1.4.8, Wyatt Technology)

and graphed using GraphPad Prism software (version 9.5.1).

Fluorescence Polarization

Fluorescence Polarization binding measurements were conducted as described in Canzio *et al.*

2013 and modified as follows:

Peptide reaction buffer was 50 mM HEPES pH 7.5, 100 mM KCl, 10% glycerol, 0.01% NP-40,

and 2 mM βME. Fluoresceinated $H3_{1-20}$ K9me0 or $H3_{1-20}$ K9me3 peptide concentration was

fixed at 100 nM while Swi6, pSwi6, or $pSwi6^{S18/24A}$ protein concentration varied from 0-200 µM.

Mononuclesome reaction buffer was 20 mM HEPES pH 7.5, 80 mM KCl, 4 mM Tris, 0.2 mM

EDTA, 10% glycerol, 0.01% NP-40, 2 mM βME. H3K9me0 and $H3K_C9me3$ mononucleosomes

were reconstituted with fluorescein-labeled 601 DNA as described[12]. Nucleosome concentration

was fixed at 25 nM while Swi6, pSwi6, or $pSwi6^{S18/24A}$ protein concentrations varied from 0-200

µM. Both peptide and mononucleosome reaction volumes were 10 µL and measured in a

Corning 384 low-volume, flat bottom plates (product number 3820, LOT 23319016).

Fluorescence polarization was recorded using a Cytation 5 microplate reader (Biotek, λ_{ex}=

485/20nm, λ_{em}= 528/20nm) and Gen5 software (Biotek, version 3.09.07). Data was analyzed

and fit to a K_d equation using GraphPad Prism.

Single turnover kinetics

Clr4 protein was purified exactly as described[23]. H3K9me2 nucleosomes were purchased from Epicypher (#16-0324), and pSwi6 and unpSwi6 were purified as above. Single turnover reactions were carried out as follows: 5 μM Clr4 was preincubated 5 minutes with 1 mM final S-adenosyl-methionine (liquid SAM, 3 2mM, NEB #B9003S), and varying concentrations of pSwi6 or unP Swi6, at 25°C to reach equilibrium. 5μM Clr4 was chosen as the minimal Clr4 concentration to yield robust H3K9me3 signal under Single Turnover conditions. The reaction was started with the addition of H3K9me2 nucleosomes to 500nM final. Timepoints were stopped by boiling with SDS-Laemmli buffer. Samples were separated on 18% SDS-PAGE gel and probed for the presence of H3K9me3 (polyclonal, Active Motif #39161. lot 22355218-11) and H4 (Active Motif #39269 lot 31519002) as a loading control. Signals were quantified on a Li-Cor imager by using a dilution of H3K9me3 nucleosomes (Epicypher, #16-0315), establishing standard curves for H4 and H3K9me3. Rates were fit to a single exponential rise in GraphPad Prism software exactly as published[23].

Phosphatase treatments

1500 ng of Swi6, pSwi6, and pSwi6$^{S18/24A}$ were incubated for 20 minutes at 37°C with 50 U of Calf Intestinal Phosphatase (QuickCIP, NEB, M0525S) in 100 mM NaCl, 50 mM Tris-HCl, 10 mM MgCl$_2$, 1 mM dithiothreitol, pH 7.9. Reactions were stopped by boiling in SDS-Laemmli buffer. For reactions with inactivated CIP, 200U CIP was pre-incubated for 20 minutes at 80°C. 75 ng of Swi6, pSwi6, and pSwi6$^{S18/24A}$ that was either mock-treated, treated with active or inactivated CIP was separated on either a 15% SDS-PAGE gel or a SuperSep Phos-Tag gel (Fujifilm, 15.5%, 17 well, 100x100x6.6mm, Lot PAR5302). The Phos-Tag gel was washed with western transfer buffer with 10 mM EDTA to remove Zn^{2+} ions and then blotted and probed for Swi6 with Swi6 polyclonal antibody as above.

Mass Spectrometry

In-solution Trypsin/Lys C digested peptides were analyzed by online capillary nanoLC-MS/MS using several different methods. High resolution 1 dimensional LCMS was performed using a 25 cm reversed-phase column fabricated in-house (75 μm inner diameter, packed with ReproSil-Gold C18-1.9 μm resin (Dr. Maisch GmbH)) that was equipped with a laser-pulled nanoelectrospray emitter tip. Peptides were eluted at a flow rate of 300 nL/min using a linear gradient of 2–40% buffer B in 140 min (buffer A: 0.02% HFBA and 5% acetonitrile in water;

buffer B: 0.02% HFBA and 80% acetonitrile in water) in a Thermo Fisher Easy-nLC1200

nanoLC system. Peptides were ionized using a FLEX ion source (Thermo Fisher) using

electrospray ionization into a Fusion Lumos Tribrid Orbitrap Mass Spectrometer (Thermo Fisher

Scientific). Data was acquired in orbi-trap mode. Instrument method parameters were as

follows: MS1 resolution, 120,000 at 200 m/z; scan range, 350−1600 m/z. The top 20 most

abundant ions were subjected to higher-energy collisional dissociation (HCD) or electron

transfer dissociation (ETD) with a normalized collision energy of 35%, activation q 0.25, and

precursor isolation width 2 m/z. Dynamic exclusion was enabled with a repeat count of 1, a

repeat duration of 30 seconds, and an exclusion duration of 20 seconds.

Low-resolution, 1-dimensional LCMS was performed using a nano-LC column packed in a 100-

μm inner diameter glass capillary with an integrated pulled emitter tip. The column consisted of

10 cm of ReproSil-Gold C18-3 μm resin (Dr. Maisch GmbH)). The column was loaded and

conditioned using a pressure bomb. The column was then coupled to an electrospray ionization

source mounted on a Thermo-Fisher LTQ XL linear ion trap mass spectrometer. An Agilent

1200 HPLC equipped with a split line so as to deliver a flow rate of 1 ul/min was used for

chromatography. Peptides were eluted with a 90-minute gradient from 100% buffer A to 60%

buffer B. Buffer A was 5% acetonitrile/0.02% heptafluorobutyric acid (HBFA); buffer B was 80%

acetonitrile/0.02% HBFA. Collision-induced dissociation spectra were collected for each m/z.

Multidimensional protein identification technique (MudPIT) was performed as described[74,75].

Briefly, a 2D nano-LC column was packed in a 100-μm inner diameter glass capillary with an

integrated pulled emitter tip. The column consisted of 10 cm of ReproSil-Gold C18-3 μm resin

(Dr. Maisch GmbH)) and 4 cm strong cation exchange resin (Partisphere, Hi Chrom). The

column was loaded and conditioned using a pressure bomb. The column was then coupled to

an electrospray ionization source mounted on a Thermo-Fisher LTQ XL linear ion trap mass

spectrometer. An Agilent 1200 HPLC equipped with a split line so as to deliver a flow rate of 1

ul/min was used for chromatography. Peptides were eluted using a 4-step gradient with 4

buffers. Buffer (A) 5% acetonitrile, 0.02% heptafluorobutyric acid (HFBA), buffer (B) 80%

acetonitrile, 0.02% HFBA, buffer (C) 250mM NH4AcOH, 0.02% HFBA, (D) 500mM NH4AcOH,

0.02% HFBA. Step 1: 0-80% (B) in 70 min, step 2: 0-50% (C) in 5 min and 0- 45% (B) in 100

min, step 3: 0-100% (C) in 5 min and 0- 45% (B) in 100 min, step 4 0-100% (D) in 5 min and 0-

45% (B) in 160 min. Collision-induced dissociation (CID) spectra were collected for each m/z.

Data analysis: RAW files were analyzed using PEAKS (Bioinformatics Solution Inc) with the

following parameters: semi-specific cleavage specificity at the C-terminal site of R and K,

allowing for 5 missed cleavages, precursor mass tolerance of 15 ppm (3 Da for low-resolution

LCMS), and fragment ion mass tolerance of 0.5 Daltons. Methionine oxidation and

phosphorylation of serine, threonine, and tyrosine were set as variable modifications and

Cysteine carbamidomethylation was set as a fixed modification. Peptide hits were filtered using

a 1% false discovery rate (FDR). Phosphorylation occupancy ratio for amino acids was

determined by summing the count of unphosphorylated and phosphorylated amino acids

detected in the experiment. We only considered phospho-peptides detected more than once

and at least 2% minimal ion intensity.

1065

Estimate of in vivo nucleosome fractions bound

In vitro binding isotherms for nucleosomes (N) can be fit simply via $fraction\ Nbound =$

$\frac{[Swi6]}{[Swi6]+Kd}$. However, this assumes first that [N] << [Swi6] and << K_d, such that [Swi6] total $\approx$

[Swi6] free. In the nucleus, these assumptions do not hold. However, a quadratic equation[48] can

be used to estimate N bound, accounting for bound Swi6. In this case,

$frNbound = \frac{[N]total+[Swi6]total+Kd-\sqrt{([N]total+[Swi6]total+Kd)^2-4*[N]total*[Swi6]total}}{2*[N]total}$. To estimate the

fraction of unmethylated nucleosomes bound by pSwi6 or unpSwi6, we used K_ds from Figure

4F, total Swi6 concentrations of 2.1-4.6µM, and total nucleosome contraction estimate of

~10µM.

1075

1076

1077

1078   **Data availability statement:**

1079   ChIP-Seq data is deposited at NIH GEO Record GSE271394. The following secure token has

1080   been created to allow review, while it remains in private status: kvwbksygjhebfyj

Figure 1

SFigure 1

SFigure 2

SFigure 3

A

B

unpSwi6   pSwi6   pSwi6$^{S18/24A}$

– + b – + b – + b   CIP

p
unp

15% SDS PAGE

C

unpSwi6   pSwi6   pSwi6$^{S18/24A}$

– + b – + b – + b   CIP

Phos-tag gel

D

pSwi6$^{S18/24A}$

<30% prevalence
>30% prevalence

E

H3 tail binding
affinity, K_d (µM)

H3K9me2	H3K9me3
6.40 ± 0.41	2.94 ± 0.14

SFigure 4

A

Swi6 WT/*S-A*   KAN^R

WT/*S-A* SF-GFP   HYG^R

Swi6 WT/*S-A*   KAN^R

⊗

Sad1:mKO2

B

wt Swi6

S46/52A

C

GFP fluorescence

☐ GFP -
■ wt Swi6
■ S18/24A
■ S18/24/117/192-220A
■ S46/52A
■ S46/52/117/192-220A

D

Puncta count

***
****
ns
ns   ns   ****

■ wt Swi6
■ S18/24A
■ S18/24/117/192-220A
■ S46/52A
■ S46/52/117/192-220A

E

Sad1-mKO2

SPB

d

Swi6-GFP

d

nucleus

Z Y X

F

foci [%]

0.6

0.5

0.4

0.3

0.2

0.1

0.0

0≤d<1   1≤d<2   d≥2

distance to SPB [μm]

■ wt Swi6
■ S18/24A
■ S18/24/117/192-220A
■ S46/52A
■ S46/52/117/192-220A

SFigure 5

A

pSwi6 (ng)

437 219 109 55 27   L   WT   S18,24A

← Swi6

α−Swi6

α−S18P-S24P

α−tubulin

B

H3K9me3 oct.   0μM   unpSwi6

←H3K9me3
←H4

0   90   180   270   390

H3K9me3 oct.   5μM unpSwi6   5μM pSwi6

←H3K9me3
←H4

0   90   180   270   390   0   90   180   270   390

C

H3K9me3 oct.   15μM unpSwi6   15μM pSwi6

←H3K9me3
←H4

0   90   180   270   390   90   180   270   390

H3K9me3 oct.   30μM unpSwi6   30μM pSwi6

←H3K9me3
←H4

0   90   180   270   390   90   180   270   390

Supporting Table 1

	Mononucleosome binding	
	relative affinity	
	H3K$_c$9me0	H3K$_c$9me3
pSwi6	7.48 ± 1.07	**1 ± 0.1**
pSwi6$^{S18/24A}$	1.67 ± 0.15	0.38 ± 0.003
	specificity	
pSwi6	7.48	
pSwi6$^{S18/24A}$	4.39	

Supporting Table 2

Yeast strains used in this study

Identifier	Genotype	Figure; experiment	Source
PAS210	h+, *sad1*:mKO2:NATMX	Fig. 3G, H; SFig.4	Al-Sady et al 2016
PAS332	h90, cenH:: ade6p:SF-GFP (Kint2); mat3m(EcoRV):: ade6p:mKO2; ade6p:3xE2C:hygMX at Locus2; ΔREIII::REIII(Δs1, Δs2)	Fig. S1J	Greenstein et al 2018
PAS807	h90, cenH:: ade6p:SF-GFP (Kint2); mat3m(EcoRV):: ade6p:mKO2; ade6p:3xE2C:hygMX at Locus2; ΔREIII::REIII(Δs1, Δs2), *swi6::ura4*	Fig. 1C,D; Fig. 2B; SFig. A-C	This study
PAS814	h90, cenH:: ade6p:SF-GFP (Kint2); mat3m(EcoRV):: ade6p:mKO2; ade6p:3xE2C:hygMX at Locus2; ΔREIII::REIII(Δs1, Δs2), *swi6*:KANMX	Fig. 1C,E; Fig. 2B-D; SFig. A-C	This study
PAS851,858, 859, 860 (isolates)	h90, cenH:: ade6p:SF-GFP (Kint2); mat3m(EcoRV):: ade6p:mKO2; ade6p:3xE2C:hygMX at Locus2; ΔREIII::REIII(Δs1, Δs2), *swi6S18/24A*:KANMX	Fig. 1C,G; SFig.1A-C; Fig. 2B-D; SFig. A-C	This study
PAS852, 861,862, 863 isolates)	h90, cenH:: ade6p:SF-GFP (Kint2); mat3m(EcoRV):: ade6p:mKO2; ade6p:3xE2C:hygMX at Locus2; ΔREIII::REIII(Δs1, Δs2), *swi6S46/52,117-220A*:KANMX	Fig. 1C,H; SFig.1D-F	This study
PAS853, 864, 865, 866 (isolates)	h90, cenH:: ade6p:SF-GFP (Kint2); mat3m(EcoRV):: ade6p:mKO2; ade6p:3xE2C:hygMX at Locus2; ΔREIII::REIII(Δs1, Δs2), *swi6S46/52*:KANMX	Fig. 1C,F	This study
PAS854	h90, cenH:: ade6p:SF-GFP (Kint2); mat3m(EcoRV):: ade6p:mKO2; ade6p:3xE2C:hygMX at Locus2; ΔREIII::REIII(Δs1, Δs2), *swi6S18/24,117-220A*:KANMX	Fig. 1C,I	This study
PAS909	*sad1*:mKO2:NATMX; *swi6S46/52*:SF-GFP:HYGMX	SFig4B-F	This study
PAS910,911	*sad1*:mKO2:NATMX; *swi6S18/24,117-220A*:SF-GFP:HYGMX	Fig3G,H; SFig4C-F	This study
PAS913	*sad1*:mKO2:NATMX; *swi6*:SF-GFP:HYGMX	Fig3G,H; SFig4C-F	This study
PAS919	*sad1*:mKO2:NATMX; *swi6S18/24*:SF-GFP:HYGMX	Fig3G,H; SFig4C-F	This study
PAS922	*sad1*:mKO2:NATMX; *swi6S46/52,117-220A*:SF-GFP:HYGMX	Fig3G,H; SFig4C-F	This study
PAS1189	h90, cenH:: ade6p:SF-GFP (Kint2); mat3m(EcoRV):: ade6p:mKO2; ade6p:3xE2C:hygMX at Locus2; ΔREIII::REIII(Δs1, Δs2), *ckb1::*KANMX	Fig. S1K,L	This study