Extinction vortices are driven more by a shortage of beneficial mutations than by deleterious mutation accumulation

Walid Mawass[1,2], Joseph Matheson[1,3], Ulises Hernández[1], Jeremy Berg[2], Joanna Masel[1]

[1]Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, 85721, USA

[2]Department of Human Genetics, University of Chicago, Chicago, IL, 60637, USA

[3]Department of Ecology, Behavior, and Evolution, University of California San Diego, San Diego, CA, 92093, USA

Corresponding author: Joanna Masel

Abstract

Natural populations are increasingly at risk of extinction due to climate change and habitat loss or fragmentation. The long-term viability of small populations can be threatened by an "extinction vortex" — a positive feedback loop between declining fitness and declining population size. Two distinct mechanisms can drive an extinction vortex: i) ineffective selection in small populations allows deleterious mutations to fix, driving "mutational meltdown", and ii) fewer individuals produce fewer of the novel beneficial mutations essential for long-term adaptation, a mechanism we term "mutational drought". We measure the relative importance of each mechanism, on the basis of how sensitive beneficial vs. deleterious components of fitness flux are to changes in census population size near the critical population size at which fitness is stable. We derive analytical results given linkage equilibrium, complemented by simulations that capture the complex linkage disequilibria that emerge under high deleterious mutation rates. Even in the absence of environmental change, mutational drought can be nearly as important as mutational meltdown. Real populations must also adapt to a changing environment, making mutational drought more important. A partial exception is that mutational drought is somewhat less important when the beneficial mutation rate is high, although its contribution remains substantial. Linkage between deleterious and beneficial mutations drives the critical population size substantially higher and increase (albeit modestly) the importance of mutational drought.

Keywords: population persistence, extinction spiral, background selection, genetic hitchhiking, population genetics, effective population size, evolutionary rescue

Introduction

Climate change, habitat loss, and habitat fragmentation directly cause natural populations to shrink, or in some cases go extinct. To what degree the genetics of a population might contribute to its decline and eventual demise, and how, remains an open question (Keller and Waller 2002; Spielman et al. 2004; Frankham 2005; Fox and Reed 2011; Kardos et al. 2021; Kyriazis et al. 2021; Dussex et al. 2023). On short timescales, population size reduction results in the loss of genetic variation and the fixation of recessive alleles, both of which typically increase the chance of extinction. Here we consider longer-term population viability on the timescale of multiple allele fixation events. A positive feedback loop known as an "extinction vortex" can occur on these longer timescales, in which declining fitness makes populations smaller, causing their fitness to fall even faster (Giplin and Soulé 1986; Tanaka 2000; Fagan and Holmes 2006). We consider two distinct mechanisms by which an extinction vortex might operate. First, ineffective selection in small populations might allow a series of deleterious mutations to become fixed, in a process of "mutational meltdown". Second, fewer individuals entail fewer opportunities for the appearance of the novel mutations required for long-term adaptation, a process we call "mutational drought".

Mutational meltdown has received the most theoretical attention of the two (Lynch and Gabriel 1990; Gabriel et al. 1993; Lynch et al. 1995a; Lynch et al. 1995b; Poon and Otto 2000; Zeyl et al. 2001; Shaw and Baer 2011; Coron et al. 2013; Lansch-Justen et al. 2022; Olofsson et al. 2023). Mutational meltdown is a positive feedback loop between population size N and mean fitness $\overline{W}$ (Figure 1). As a population declines in size, deleterious mutations fix more easily due to random genetic drift (Figure 1, red arrow). This decreases mean fitness $\overline{W}$, causing the population size to decline further (Figure 1, upward arrow).

Figure 1. **An extinction vortex can be driven by the dynamics of either deleterious (mutational meltdown; red) or beneficial (mutational drought; blue) mutations**.

In the absence of beneficial mutations, even large populations melt down eventually. This is because, given a continuous distribution of deleterious selection coefficients, some are too small to be reliably purged by selection, even in large sexual populations; this has been called "Ohta's ratchet" (Ruan et al. 2020). Schultz and Lynch (1997) and Whitlock (2000) therefore modeled a low rate of beneficial mutation, sufficient in larger populations to offset deleterious fixation events. They calculated the fitness fluxes v_d and v_b from deleterious and beneficial fixation events, respectively. Each of the fluxes is given by $UN \int s p_{fix}(s) p(s) ds$, where $p_{fix}(s)$ is the probability of fixation of a new mutation with selection coefficient s. Unlike Lande (1998), who considered only reverse mutations, treatment of full distributions for both s_b and s_d enables compensatory asymmetry, i.e. many small effect deleterious fixations can be

offset by a smaller number of larger effect beneficial fixations (Whitlock 2000; Whitlock et al. 2003;

Matheson et al. 2023).

Schultz and Lynch (1997) emphasized the critical value of U_b/U_d (i.e. ratio of genome-wide

beneficial : deleterious mutation rates) that is needed to balance deleterious and beneficial fluxes, as a

function of population size N and degree of outcrossing. Here we follow Whitlock (2000), who studied

an essentially identical model, but inverted the relationship to instead emphasize a critical population

size N_{crit} as a function of U_b/U_d. Below N_{crit}, the population crosses the threshold into an extinction

vortex, i.e., a sustained and irreversible loss of fitness.

What was not noted in these previous treatments is that new beneficial mutations do not just offset

degradation, but they also create a second, distinct positive feedback loop (Figure 1, blue arrow).

Declining N reduces the number of beneficial mutations entering the population, determined by NU_b.

This slows the adaptive increase in $\bar{W}$, which feeds back into lower N (Figure 1, upward arrow). The

importance of this second feedback loop was previously briefly noted in the Discussion of Osmond and

Klausmeier (2017) as an incidental finding, within a study focused on the effect of fitness function shape

on extinction vortices in the context of a rapidly changing environment. We refer to this novel type of

extinction vortex as "mutational drought," and note that when U_d is high, it could be important even in

the absence of environmental change.

Here we systematically explore how much declining beneficial flux vs. increasing deleterious flux

contribute to falling net fitness flux, as habitat loss drives a population's size down toward N_{crit}.

Mathematically, we describe the sensitivities of both beneficial and deleterious fitness fluxes as their

derivatives with respect to population size, $\frac{dv}{dN}$, evaluated at N_{crit}. The "ben:del driver ratio"

$\left|\frac{dv_b}{dN}\middle/\frac{dv_d}{dN}\right|_{N_{crit}}$ quantifies the degree to which beneficial vs. deleterious mutations are more important in

precipitating an extinction vortex. A ratio < 1 indicates a deleterious driver of mutational meltdown, a

ratio > 1 indicates a beneficial driver of mutational drought, and a ratio equal to 1 represents equal

contributions.

Whitlock's (2000) simple model of fitness flux $v = UN \int sp_{fix}(s)p(s)ds$ assumes that each

mutation evolves independently. However, this assumption is unrealistic because of linkage disequilibria

(LD) between loci. Deleterious mutations cause background selection, and beneficial mutations cause

hitchhiking and clonal interference (Park and Krug 2007; Stephan 2010; Charlesworth and Jensen 2021).

These increase the fixation probability of deleterious mutations and decrease that of beneficial mutations.

Indeed, LD among deleterious mutations has been shown to accelerate mutational meltdown (Lynch et

al. 1995b).

Background selection among unlinked sites must be accounted for in the $U_d > 1$ regime in which it

is more powerful than background selection among linked sites (Matheson and Masel 2024). The

evidence for high $U_d > 1$ is best established for humans. Lesecque et al. (2012) obtained a lower bound

for human U_d by assuming that mutations are deleterious only in the 55% of the 6×10^9 diploid genome

that is not dominated by dead transposable elements shared with chimpanzees, and that this 55% of the

genome evolves 5.7% slower due to this constraint (Eőry et al. 2010). Combined with a conservative

point mutation rate of 1.16×10^{-8} (Wang and Obbard 2023), this yields an estimated rate of deleterious

mutations of $0.55 \times 6 \times 10^9 \times 0.057 \times 1.16 \times 10^{-8} = 2.2$ per replication. This estimate is conservative: some

mutations to transposable element regions are deleterious, 1.16×10^{-8} is the lower bound of a 95%

confidence interval (Wang and Obbard 2023), and non-point mutations and beneficial mutations are

neglected. Some therefore argue that the human deleterious mutation rate is as high as $U_d = 10$

(Kondrashov 2017). Humans have only a modestly higher point mutation rate than most other eukaryotes

(Wang and Obbard 2023), making it likely that $U_d > 1$ is common.

However, capturing the unlinked LD that is important in the $U_d > 1$ regime requires simulating the

entire genome rather than just a genomic region. This is computationally infeasible for most forward-

time simulators (e.g., SLiM, Haller and Messer 2019), because the number of segregating mutations to

be tracked scales with $U_d N$, which becomes too large. Here we conduct whole genome simulations using

a recently developed approach (Matheson et al. 2023) that assumes that recombination occurs only at

hotspots; we track only a summary of each "linkage block" (region between hotspots), avoiding the need

to track every segregating mutation.

Here we ask how important mutational drought is relative to mutational meltdown, as a function of

three main parameters: 1) U_d/U_b, 2) the presence or absence of environmental change, and 3) the

presence or absence of LD. We also ask how much LD affects N_{crit}, and how the reduction in both N_{crit}

and the usual N_e are influenced by background selection vs. hitchhiking vs. clonal interference when a

population is near the edge of long-term viability.

Methods

1. Without linkage disequilibrium (LD)

Fixation probability. Independent evolution of each site (i.e. no LD nor epistasis) enables an

analytical, one locus approach. Each copy of a mutation with fitness effect x affects fitness by a factor

of $(1 + x)$. To facilitate comparisons with simulation results, we capture co-dominance not via a

dominance coefficient $h = 0.5$, but by reducing fitness by $(1 + x)^2$ for homozygous mutants. The exact

probability of fixation $p_{fix}(s)$ of a new mutation under the Moran birth-death process is then

126
$$p_{fix}(s) = \frac{1 - e^{-2s}}{1 - e^{-4Ns}}$$

127   where the selection coefficient $s = \ln(1 + x)$ represents the difference in Malthusian fitness. We take

128   this from Equation 1 of McCandlish et al. (2015), adjusted for a diploid population of size N.

129   *Deleterious fitness flux*. Polymorphism data allows inference of the distribution of $2N_e x$, which

130   we divide by $2N_e$. We use the human distribution of fitness effects (DFE), because it has been well

studied, and is at least somewhat appropriate for large-bodied species commonly of conservation concern. We contrast two published DFEs: Boyko et al. (2008; α = 0.184, β = 8200) and Kim et al. (2017; α = 0.169, β = 1327.4). Both fit a gamma distribution $p(-2N_e x_d) = \frac{e^{-2N_e x_d/\beta} \, 2N_e x_d^{-1+\alpha} \beta^{-\alpha}}{\Gamma(\alpha)}$ where $\Gamma(\alpha)$ is the gamma function, α is the shape parameter, and β is the scale parameter. Selection coefficients are of larger magnitude in the Boyko et al. (2008) DFE than in the Kim et al. (2017) DFE, as can be seen by comparing the cumulative probability distributions (Figure S1).

The deleterious fitness flux, i.e. the rate population fitness changes due to deleterious fixations, is equal to the per-generation rate of new mutations $U_d N$, times their fixation probability $p_{fix}(s_d)$, times their homozygous impact on fitness $2x_d$ once fixed:

$$v_d = U_d N \int_{-\infty}^{0} p_{fix}\left(\ln(1 + \frac{2N_e x_d}{2N_e})\right) \frac{2N_e x_d}{N_e} \, p(2N_e x_d) \, d(2N_e x_d) \qquad (1)$$

We integrate with respect to $2N_e x_d$ rather than x_d for convenient usage of inferred DFEs. We rescale using the inferred N_e from either Boyko et al. (2008; N_e= 25636) or Kim et al. (2017; N_e= 11823), matching the DFE used.

We incorporated environmental change as a constant added to the deleterious flux, that varied across the range $[10^{-6} - 10^{-3.5}]$. Environmental change rates are expressed, for ease in interpretation, as the number of generations over which environmental change causes a 10% reduction in fitness; this is equal to $0.1/offset$. We note that these values are slow compared to the current rate of anthropogenic climate change, yet fast relative to historical mass extinction events on the geological time scale.

Beneficial fitness flux. We assume an exponential distribution with mean effect size $\bar{x}_b$:

$$p(x_b) = \frac{1}{\bar{x}_b} \, e^{-x_b/\bar{x}_b}$$

151   The beneficial fitness flux, i.e. the change in population fitness due to the fixation of beneficial mutations

152   is

$$v_b = U_b N \int\limits_{0}^{+\infty} p_{fix}(\ln(1 + x_b)) \, 2x_b \, p(x_b) \, dx_b \qquad (2)$$

Critical population size. We calculated the deleterious and beneficial fitness fluxes numerically

in Mathematica 12.3.0 (Supplementary Notebook), using Equations (1) and (2). We solve for N_{crit} as the

value of N for which $v_{net} = v_b + v_d = 0$. We then calculate the ben : del ratio as $\left|\frac{dv_b}{dN} \middle/ \frac{dv_d}{dN}\right|_{N_{crit}}$.

2. With LD

Simulation setup. Our forward time simulation method is described in detail elsewhere (Matheson et

al. 2023). Briefly, each genome consists of two haplotypes, divided into 23 chromosomes, further

segmented into 'linkage blocks'. The fitness impact of mutations within each linkage block is

summarized as $l_j = \prod_i (1 + x_i)$. The fitness of an individual combines the effects of all linkage blocks

as $w_i = \prod_{j=1}^{L}(l_{j,1}) \prod_{j=1}^{L}(l_{j,2})$ where $l_{j,1}$ and $l_{j,2}$ refer to the two copies of linkage block j in a diploid.

We employ a Moran model with a constant population size N, where individuals are chosen to reproduce

based on their fitness.

Recombination occurs only at hotspots between adjacent linkage blocks. We simulated two

recombination events per chromosome per meiosis (Pardo-Manuel de Villena and Sapienza 2001). We

used 100 linkage blocks per chromosome for computational efficiency, which previous work found to be

more than sufficient (Matheson et al. 2023).

We sampled the numbers of new deleterious and beneficial mutations from Poisson distributions

with means U_d and U_b, respectively. We sampled selection coefficients from the distributions described

in the previous section, rescaling in the case of x_d.

Simulations ran for $100N$ generations. Because we initialized simulations with a monomorphic

population, we excluded a burn-in phase when measuring fitness flux. Following Matheson et al. (2023),

we ended the burn-in phase 100 generations after a linear regression of the variance in fitness over the

last 200 generations produced a slope less than an arbitrary small value of $0.07/N$. Net fitness flux v_{net}

was estimated post burn-in as the slope $\hat{v}_{net}$ of log mean population fitness over time, without

decomposition into v_d and v_b.

Critical population size. N_{crit} is defined as the value of N for which $v_{net} = 0$, but $\hat{v}_{net}$ estimation

is too noisy to find N_{crit} using purely deterministic root-finding methods (specifically for small values

of N_{crit}). After using a standard root-finding approach to choose a set of points near N_{crit} at which to

evaluate $\hat{v}_{net}$, we therefore estimated N_{crit} as the root of a fitted linear regression, to cancel out the noise.

Specifically, we first bracketed N_{crit}, with N = 1500 and 6000 as initial guesses, progressing outward by

a factor of two as needed until we had $\hat{v}_{net}$ values both above and below zero. Then we iterated the

secant method until $\hat{v}_{net}$ changed by < 15% in the last iteration. Then we fitted a straight line to all

evaluated $\{N, \hat{v}_{net}\}$ data points for which N was within a factor of 3 from the final evaluated value. We

estimated N_{crit} as the zero for this fitted line.

Ben:del ratio. To calculate the ben:del driver ratio in our simulations, we needed to partition the

net fitness flux v_{net} into its deleterious and beneficial components, such that $v_{net} \sim v_b - v_d$. We

estimated v_d and v_b from the set of mutations whose time of final fixation lay between the end of the

burn-in phase and the end of the run. Tracking fixations does not exactly track mean population fitness

due to fluctuations caused by polymorphisms. For computational efficiency, the simulations described

so far summarize the fitness of many mutations per linkage block, making information about individual

fixed mutations inaccessible. We solve this using tree sequence recording (Kelleher et al. 2018), which

succinctly encodes the complete history of genetic lineages within a population. In simulations intended

to estimate the ben:del driver ratio, we used tree sequencing recording to track all non-neutral mutations,

including their time of fixation. We calculated $v_d = \frac{\sum_{i=1}^{n_d} 2 \ln (1+x_{di})}{G^*}$ and $v_b = \frac{\sum_{i=1}^{n_b} 2 \ln (1+x_{bi})}{G^*}$, where G^*

equals generations elapsed since the burn-in phase ended, and n_d and n_b represent the numbers of

deleterious and beneficial mutations that fixed during this time.

We numerically estimated the derivative of each fitness flux component, $\frac{dv_b}{dN}$ and $\frac{dv_d}{dN}$, as

200
$$\frac{dv}{dN} = \frac{v(N_{crit} + \varepsilon) - v(N_{crit} - \varepsilon)}{2\varepsilon},$$

where we chose $\varepsilon = 150$ individuals as a balance between reducing noise and avoiding curvature in v.

Results

Without LD, Whitlock's (2000) choice of $U_d/U_b = 1000$, combined with the deleterious DFE

that Kim et al. (2007) inferred for humans, yields $N_{crit} \simeq 1834$ (Figure 2A, solid blue and red lines

intersecting at dotted vertical line). Approximating the DFE as point masses at the beneficial and

deleterious means, Melissa and Desai's (2024) expression $N_{crit} = \frac{2}{(\bar{x}_b+|\bar{x}_d|)} \ln[(U_d|\bar{x}_d|)/(U_b\bar{x}_b)] =$

1821 is more accurate than Whitlock's (2000) $N_{crit} \approx \sqrt[3]{\frac{U_d}{64 \, \bar{x}_b{}^2 |\bar{x}_d| U_b}} = 1202$. Note that in both

expressions, N_{crit} depends on U_d and U_b only via their ratio. The Kim et al. (2007) DFE has $\bar{x}_d = -0.009$,

compared to Whitlock's (2000) choice of $\bar{x}_d = -0.02$ and -0.002; as an alternative, we used the DFE from

Boyko et al. (2008) DFE, whose larger $\bar{x}_d = -0.029$ yields $N_{crit} \simeq 1682$. The difference arises because

the Kim et al. (2007) DFE has more probability mass in the weakly selected regime where selection is

inefficient at purging deleterious alleles. A key reason for our high value of N_{crit} relative to Whitlock's

(2000) is that we are conservative with respect to the role of beneficial mutations, using lower $\bar{x}_b = 0.001$

than Whitlock's (2000) $\bar{x}_b = 0.02$ and 0.002.

Figure 2. Extinction vortices can be driven as much or more by mutational drought as by mutational meltdown. Vertical lines indicate N_{crit}, defined by the intersection of beneficial (blue) and deleterious and/or environmental (red) fitness fluxes in (A). Dotted vertical lines are for deleterious fitness flux alone, dashed vertical lines (and red dot-dashed line in (A)) also include the impact of environmental change in degrading fitness. Environmental degradation adds a constant amount to deleterious flux by in (A), and hence does not change the red line representing slope in (B). Continuous red and blue lines are analytical results given linkage equilibrium. $U_d = 2$, $U_d/U_b = 1000$, $\bar{x}_b = 0.001$, $\bar{x}_d = -0.009$ via the DFE from Kim et al. (2017), and environmental change degrades fitness by 3×10^{-5} per generation.

At N_{crit} in the absence of environmental change (dotted vertical line in Figure 2B), declining N

has 85% of the impact on beneficial flux as it has on the deleterious flux that is normally assumed the

sole cause of an extinction vortex according to mutational meltdown theory. When adaptation must

counter environmental change in addition to deleterious fitness flux, larger populations can enter the

extinction vortex, that is N_{crit} is higher (Fig. 2A illustrates the case of environmental change reducing

fitness by 10% over 30,000 generations, red dot-dashed line intersects with blue solid line to produce

dashed vertical line at $N_{crit} \simeq 2941$). In this scenario, the impact of declining N on beneficial flux is

larger than that on deleterious flux (blue line is above red along dashed vertical line in Figure 2B). In

other words, mutational drought is more important than mutational meltdown.

More broadly, the ben:del ratio is > 1 with $U_d/U_b = 1000$ when slow environmental change

causes fitness to decline by 10% over as many as ~20,000 generations (Figure 3A, right grey dot-dashed

vertical line). This falls only modestly to ~4000 generations when U_b is 10 times higher (Figure 3A, left

blue dotted vertical line). For $U_d = 2$, these slow rates of environmental decline correspond to only 0.19×

and 0.28× the rates of fitness decline due to deleterious fixations (Figure 3B, intersections with vertical

lines).

Figure 3. **Slow environmental degradation is sufficient to make mutational drought more important than mutational meltdown.** $\bar{x}_b$ = 0.001, U_d = 2, $\bar{x}_d$ = -0.009 via the DFE from Kim et al. (2017). The black star indicates the example of environmental change shown in Figure 2. Vertical lines represent where the ben:del ratio = 1, matching the color of the main curves. Diagonal black dashed line in (B) represents proportionate change in the x-axis and y-axis, corresponding to no increase in deleterious flux in the face of rapid adaptation. Red dashed lines in (C) indicate the asymptotes, i.e. no environmental change.

When beneficial mutations are scarce (higher U_d/U_b), the relative importance of mutational

drought is higher, with the ben:del ratio saturated by the $U_d/U_b \sim 1000$ value (Figure 4A) used in Fig. 2.

Lowering U_b has two opposing effects. First, it proportionately lowers the beneficial fitness flux v_b and

its derivative $\frac{dv_b}{dN}$ (i.e. it reduces the slope of the blue line in Figures 2A and consequently reduces the

height of the blue line in Figure 2B). Second, lower v_b moves the intersection N_{crit} to the right, i.e.

towards larger values of N, which lowers $\left.\frac{dv_d}{dN}\right|_{N_{crit}}$ without substantially changing $\left.\frac{dv_b}{dN}\right|_{N_{crit}}$ (Figure 2B,

red and blue lines, respectively). When beneficial mutations are scarce at high $U_d/U_b \geq 1000$, these two

effects are exactly balanced.

Using the higher mean deleterious effect size $\bar{x}_d$ of Boyko et al. (2008) only slightly decreases the

relative importance of mutational drought at higher $\bar{x}_d$, primarily when beneficial mutations are common

(Figure 4A, difference between lines is largest at low U_d/U_b). Larger beneficial effect sizes only slightly

increase the relative importance of mutational drought, again with a slightly larger effect when beneficial

mutations are more common (Figure 4B; see Figure S2 for change in N_{crit}).

Figure 4. **The mean sizes of (A) deleterious and (B) beneficial selection coefficients have negligible impact on the relative importance of mutational drought : mutational meltdown.** (A) Mutational drought becomes somewhat less important when beneficial mutations are abundant (low U_d/U_b). Analytical results are shown for the independent sites model. When not otherwise specified, $U_d = 2$, $\bar{x}_b = 0.001$. Results for different U_d are superimposable.

Including additional fitness degradation due to environmental change makes mutational drought more important. With environmental change, dependence on U_d and U_b does not simplify to dependence on their ratio, as in Figure 4A. We illustrate this in Figure 5 for one fast and one slow environmental change scenario, contextualized relative to mutational degradation rates in Figures 5B and 5E, and relative to N_{crit} in Figure 5C and 5F. Faster environmental degradation, fast enough to put the ben:del ratio above 1, makes mutational drought more rather than less important for lower U_d (Figure 5A). The ben:del ratio is only below 1 when relatively high U_b is combined with slow environmental change (Figure 5D), substantially slower than deleterious flux (Figure 5E).

Figure 5. (A) Faster environmental degradation makes mutational drought more rather than less important for lower U_d. (A) and (D) can be compared to the no-environmental change case in Figure 4A, where only the ratio U_d/U_b matters; reverse scaling of the U_b x-

axis, and different plausible x-axis ranges are displayed, to facilitate comparison. Dotted
vertical lines represent $U_d = 2$ and $U_b = 0.002$, matching the example in Figure 2. Dotted
horizontal lines in (C) and (F) indicate the N_{crit} with this parameter combination but no
environmental change, showing that our slow environmental change scenario (dashed lines,
fitness reduced 10^{-6} per generation) has negligible effect on N_{crit}. Faster environmental change
(solid lines) represents a fitness reduction of 3×10^{-5} per generation, matching the example in
Figure 2; the impact relative to deleterious flux is contextualized in (B) and (E), and relative
to N_{crit} in (C) and (F). Analytical results are shown for the independent sites model with $\bar{x}_b =$
0.001 and $\bar{x}_d = -0.009$ via the DFE from Kim et al. (2017). A higher mean size of deleterious
selection coefficients leads to ben:del values a little further from 1 (see Figure S3A and B).

The presence of LD given $U_d = 2$ also produces a (more modest 8%) increase in the ben:del

ratio, from 0.85 to 0.92, as beneficial flux becomes slightly more sensitive to N (Figure 6D, blue points

are above blue line). LD moves the $v_d = v_b$ intersection slightly higher in Figure 6C, corresponding to

higher mean fitness and larger fitness fluxes.

LD raises N_{crit} up to 4403 (Figure 6A, solid dots; Figure S4). The shifts in both beneficial and

deleterious flux curves are well captured by rescaling N to a form of N_e (constructed by enforcing equal

effective N_{crit} values), as seen by the close match between analytical lines and simulated points when

the x-axes of Figures 6A and 6B are rescaled to produce Figures 6C and 6D.

Figure 6. Linkage disequilibrium makes the role of mutational drought slightly more important compared to the linkage equilibrium case. Continuous red and blue lines are analytical results given linkage equilibrium, while dots are simulation results with LD. (A) and (B) are the same plots in Figure 2 with simulation results added in(A). In (C) and (D), the x-axis for simulation results is scaled such that the lines intersect at the same "effective" N_{crit}. $U_d = 2$, $U_d/U_b = 1000$, $\bar{x}_b = 0.001$, $\bar{x}_d = -0.009$ via the DFE from Kim et al. (2017). We note that the beneficial-fitness-flux-effective population size and the deleterious-fitness-flux-effective population size behave similarly to one another (C and D).

The reduction in N_e could be due to beneficial mutations interfering with each other (clonal

interference decreasing beneficial flux), and/or hitchhiking (beneficial mutations increasing deleterious

flux), and/or background selection (deleterious mutations both decreasing beneficial flux and increasing

deleterious flux). In Figure 6 without environmental change, background selection from deleterious

mutations on deleterious or neutral mutations (using eq. (4) in Matheson and Masel 2024) predicts a ratio

of $N_e/N = 0.82$, but we see a ratio of $N_{crit(analytical)}/N_{crit(simulation)} = 0.42$. A model of deleterious

mutations alone thus accounts for only $(1 - 0.82)/(1 - 0.42) = 30\%$ of the reduction in effective N_{crit}.

When we measure N_e the usual way, as the expected coalescent time, the factor reduction in N_e (Figure

7, blue triangles) matches the reduction in fitness fluxes and hence N_{crit} (Figure 7, black circles), but not

that predicted by deleterious mutations alone (Figure 7, red squares).

If low N_e were due to clonal interference, then the reduction in N_e should be stronger for high U_b;

but the (very weak) trend is in the opposite direction (Figure 7A). Since neither beneficial mutations

alone (clonal interference) nor deleterious mutations alone can explain our results, this points to an

interaction between deleterious and beneficial mutations. There are two kinds of interaction: beneficial

mutations increase the fixation probability of deleterious mutations (Charlesworth 1994; Peck 1994), and

deleterious mutations decrease the fixation probability of beneficial mutations (Good and Desai 2014;

Pénisson et al. 2017). From Figure 6, we deduce that both happen to the same degree. This interpretation

explains greater N_e reduction for large U_d (Figure 7B). It is also compatible with slightly less N_e

reduction for larger U_b (Figure 7A), where there is more competition between less-loaded vs. more-

loaded beneficial lineages.

Figure 7. **The degree to which effective population size is reduced below census size is more sensitive to the deleterious than to the beneficial mutation rate.** $N_{crit(analytical)}/N_{crit(simulation)}$ serves as an estimate of N_e/N. $\bar{x}_b = 0.001$, $U_d = 2$ (in A), $U_b = 0.02$ (in B), and $\bar{x}_d = -0.009$ via the DFE from Kim et al. (2017). Model of deleterious mutations alone is from Equation 4 in Matheson and Masel (2024).

Figure 8. **More recombination does not make N_{crit} converge to the expectation for independent loci (A), but this does not affect the ben:del ratio (C).** (B) More recombination makes the model of deleterious mutations alone a slightly worse fit. $\bar{x}_b = 0.001$, $U_b = 0.002$, $U_d = 2$, $\bar{x}_d = -0.009$ via the DFE from Kim et al. (2017), $L = 100$ per chromosome. Red dashed lines show analytical expectations in the absence of LD. Model of deleterious mutations alone in (B) is from Equation 4 in Matheson and Masel (2024).

We manipulate the amount of recombination in our simulations of LD by changing the number of

chromosomes, given two crossover events per chromosome plus independent segregation. More

recombination lowers N_{crit}, but realistic numbers of chromosomes are not sufficient to converge on the

value expected without LD (Figure 8A). However, 50 chromosomes, each with 100 linkage blocks, are

sufficient to obtain a ben:del ratio in line with independent-sites expectations, with more LD causing

only a modest increase in the relative importance of mutational drought (Figure 8C). Our model of

deleterious mutations alone, which poorly describes the effect of LD on N_{crit}, becomes somewhat worse

given more recombination (Figure 8B).

Discussion

Mutational meltdown describes an extinction vortex in which small populations cannot purge

slightly deleterious mutations, whose fixations handicap the population, making it still smaller. Here, we

find that a shortage of beneficial mutations appearing in smaller populations creates mutational drought,

a distinct kind of extinction vortex. Mutational drought is only slightly less significant than mutational

meltdown in the absence of environmental change, and becomes more important in the more realistic

scenario of a changing environment. Mutational drought is particularly significant if we assume that

beneficial mutations are rare, and to a less extent if there is LD. Beneficial mutations are thus either

common, or else their population-size-dependent scarcity is critical – in neither case can population

genetic models ignore beneficial mutations. Near the critical census population size, the reduction in

effective population size is best explained by interactions between beneficial and deleterious mutations.

The role of beneficial mutations in population persistence studied here plays out over much longer

timescales than processes of "evolutionary rescue" in previous population genetic (discrete) models,

although in line with the broader use of the term in the context of quantitative genetics models (see

Osmond and Klausmeier 2017). In the broad sense, evolutionary rescue refers to scenarios where genetic

adaptation struggles to keep up with ongoing environmental change, over an indefinitely long period of

time (Gomulkiewicz and Holt 1995; Gonzalez et al. 2013; Alexander et al. 2014; Carlson et al. 2014;

Bell 2017). When using the narrow sense of the term, population genetic models consider a population

whose birth rate is initially less than its death rate, e.g. following a single abrupt environmental change

that can be remedied by one or several beneficial mutation(s) (Orr and Unckless 2008; Martin et al. 2013;

Orr and Unckless 2014; Uecker et al. 2014; Uecker and Hermisson 2016; Wilson et al. 2017). Here we

use a population genetic model to consider evolutionary rescue in the broad sense more commonly treated

by quantitative genetic models, across an indefinitely long series of beneficial and deleterious fixations.

Given ongoing environmental change, we find that mutational drought, not mutational meltdown, is the

dominant process driving the extinction vortex. We furthermore find that mutational drought is nearly as

important, even in the absence of any environmental change.

Conservation management and species recovery programs assess the 'minimum viable

population' (MVP) and the factors influencing it (Shaffer 1981; Lacava and Hughes 1984; Reed et al.

2003; Traill et al. 2007). We follow Nabutanyi and Wittmann (2022) in defining N_{crit} as the lowest

population size that avoids an extinction vortex. This is a different kind of MVP to the usually considered

empirical predictor of probability of persistence after a specified period of time, e.g. 1000 years (Nunney

and Campbell 1993; Traill et al. 2007). Our N_{crit} concerns the ability to resupply the population with

new beneficial mutations to balance both the accumulation of deleterious mutations, and ongoing

environmental change, to achieve long-term viability. N_{crit} is the gateway to the extinction vortex below

which some combination of mutational meltdown and mutational drought contribute. Our N_{crit} is

relevant over the long timescales of multiple sweeps making our results applicable more clearly to species

than to populations, reliant on mutation rather than migration as source of new variants.

The reduction from N to N_e, at the realistically high $U_d = 2$ we consider, is primarily caused by

LD between beneficial and deleterious mutations, with a smaller contribution from LD among deleterious

mutations. A model of background selection among linked deleterious mutations alone, capturing only the smaller component (Matheson and Masel 2024) of the smaller effect, is sometimes considered as the best explanatory model of genetic diversity patterns (Johri et al. 2020; Murphy et al. 2022; Buffalo and Kern 2024), e.g., via one-locus models that use the B parameter to rescale N_e (B is the ratio of the effective population size under background selection to the neutral value; Charlesworth 2013). Our results here suggest that such a model will underestimate the degree to which LD reduces N_e and N_{crit} in the parameter value regime of interest to species of conservation concern.

The assessed extinction risk of several classes of vertebrate species in the Anthropocene (IUCN 2024) includes assessed long-term risks to environmental conditions (Pacifici et al. 2015). Current conservation efforts often focus on emergency revitalization programs with tools that might only achieve short-term persistence (e.g., Santidrián Tomillo et al. 2021), potentially creating trade-offs between long-term and short-term objectives (Rout et al. 2007). Short-term goals may be problematically prioritized under triage from restricted budgets (Bottrill et al. 2008; Vucetich et al. 2017) over achieving long-term sustainable populations (Willis and Birks 2006; Barnosky et al. 2017).

Some conservation geneticists advocate genetic rescue, whereby gene flow from a few introduced individuals helps small fragmented populations (Whiteley et al. 2015; Frankham et al. 2017; Bell et al. 2019). The genetic rescue paradigm aims to maximize genetic diversity to reduce the risk of extinction (Whiteley et al. 2015; Ralls et al. 2018; Ralls et al. 2020). However, gene flow from large, outbred populations may introduce too many recessive deleterious alleles to be easily purged, with deleterious effects that might quickly have devastating consequences within a small inbred population (Bijlsma et al. 2010; Kyriazis et al. 2021; Pérez-Pereira et al. 2022). Our results suggest that abandoning the genetic rescue paradigm, while potentially beneficial in the short term, might still be a mistake over longer timescales. Fortunately, according to simulations, the benefits of genetic rescue can be obtained while minimizing harm by steadily introducing migrants from other small inbred populations from which

410   recessive deleterious mutations are already purged (Kyriazis et al. 2021; Pérez-Pereira et al. 2022), as

411   has occurred naturally in Florida scrub-jays (Chen et al. 2016).

Data Availability

Analtyical results conducted in Wolfram Mathematica. Simulation code written in C, and graphs produced with R. Simulation code and scripts are available on GitHub at https://github.com/MaselLab/MutationLoad.

Conflict of interest

All authors declare no conflicts of interest.

References

Alexander HK, Martin G, Martin OY, Bonhoeffer S. 2014. Evolutionary rescue: linking theory for conservation and medicine. Evol Appl. 7(10):1161–1179. doi:10.1111/eva.12221.

Barnosky AD, Hadly EA, Gonzalez P, Head J, Polly PD, Lawing AM, Eronen JT, Ackerly DD, Alex K, Biber E, et al. 2017. Merging paleobiology with conservation biology to guide the future of terrestrial ecosystems. Science. 355(6325):eaah4787. doi:10.1126/science.aah4787.

Bell DA, Robinson ZL, Funk WC, Fitzpatrick SW, Allendorf FW, Tallmon DA, Whiteley AR. 2019. The Exciting Potential and Remaining Uncertainties of Genetic Rescue. Trends Ecol Evol. 34(12):1070–1079. doi:10.1016/j.tree.2019.06.006.

Bell G. 2017. Evolutionary Rescue. Annu Rev Ecol Evol Syst. 48(Volume 48, 2017):605–627. doi:10.1146/annurev-ecolsys-110316-023011.

Bijlsma R, Westerhof MDD, Roekx LP, Pen I. 2010. Dynamics of genetic rescue in inbred Drosophila melanogaster populations. Conserv Genet. 11(2):449–462. doi:10.1007/s10592-010-0058-z.

Bottrill MC, Joseph LN, Carwardine J, Bode M, Cook C, Game ET, Grantham H, Kark S, Linke S, McDonald-Madden E, et al. 2008. Is conservation triage just smart decision making? Trends Ecol Evol. 23(12):649–654. doi:10.1016/j.tree.2008.07.007.

Boyko AR, Williamson SH, Indap AR, Degenhardt JD, Hernandez RD, Lohmueller KE, Adams MD, Schmidt S, Sninsky JJ, Sunyaev SR, et al. 2008. Assessing the Evolutionary Impact of Amino Acid Mutations in the Human Genome. PLOS Genet. 4(5):e1000083. doi:10.1371/journal.pgen.1000083.

Buffalo V, Kern AD. 2024. A quantitative genetic model of background selection in humans. PLOS Genet. 20(3):e1011144. doi:10.1371/journal.pgen.1011144.

Carlson SM, Cunningham CJ, Westley PAH. 2014. Evolutionary rescue in a changing world. Trends Ecol Evol. 29(9):521–530. doi:10.1016/j.tree.2014.06.005.

Charlesworth B. 1994. The effect of background selection against deleterious mutations on weakly selected, linked variants. Genet Res. 63(3):213–227. doi:10.1017/S0016672300032365.

Charlesworth B. 2013. Background Selection 20 Years on: The Wilhelmine E. Key 2012 Invitational Lecture. J Hered. 104(2):161–171. doi:10.1093/jhered/ess136.

Charlesworth B, Jensen JD. 2021. Effects of Selection at Linked Sites on Patterns of Genetic Variability. Annu Rev Ecol Evol Syst. 52(Volume 52, 2021):177–197. doi:10.1146/annurev-ecolsys-010621-044528.

Chen N, Cosgrove EJ, Bowman R, Fitzpatrick JW, Clark AG. 2016. Genomic Consequences of
Population Decline in the Endangered Florida Scrub-Jay. Curr Biol. 26(21):2974–2979.
doi:10.1016/j.cub.2016.08.062.

Coron C, Méléard S, Porcher E, Robert A. 2013. Quantifying the Mutational Meltdown in Diploid
Populations. Am Nat. 181(5):623–636. doi:10.1086/670022.

Dussex N, Morales HE, Grossen C, Dalén L, Oosterhout C van. 2023. Purging and accumulation of
genetic load in conservation. Trends Ecol Evol. 38(10):961–969. doi:10.1016/j.tree.2023.05.008.

Eőry L, Halligan DL, Keightley PD. 2010. Distributions of Selectively Constrained Sites and
Deleterious Mutation Rates in the Hominid and Murid Genomes. Mol Biol Evol. 27(1):177–192.
doi:10.1093/molbev/msp219.

Fagan WF, Holmes EE. 2006. Quantifying the extinction vortex. Ecol Lett. 9(1):51–60.
doi:10.1111/j.1461-0248.2005.00845.x.

Fox CW, Reed DH. 2011. Inbreeding Depression Increases with Environmental Stress: An
Experimental Study and Meta-analysis. Evolution. 65(1):246–258. doi:10.1111/j.1558-
5646.2010.01108.x.

Frankham R. 2005. Genetics and extinction. Biol Conserv. 126(2):131–140.
doi:10.1016/j.biocon.2005.05.002.

Frankham R, Ballou JD, Ralls K, Eldridge M, Dudash MR, Fenster CB, Lacy RC, Sunnucks P. 2017.
Genetic Management of Fragmented Animal and Plant Populations. Oxford University Press.

Gabriel W, Lynch M, Bürger R. 1993. Muller's Ratchet and Mutational Meltdowns. Evolution.
47(6):1744–1757. doi:10.1111/J.1558-5646.1993.TB01266.X.

Giplin ME, Soulé ME. 1986. Minimum viable populations : Processes of species extinction. Conserv
Biol Sci Scarcity Divers.:19–34.

Gomulkiewicz R, Holt RD. 1995. When does Evolution by Natural Selection Prevent Extinction?
Evolution. 49(1):201–207. doi:10.2307/2410305.

Gonzalez A, Ronce O, Ferriere R, Hochberg ME. 2013. Evolutionary rescue: an emerging focus at the
intersection between ecology and evolution. Philos Trans R Soc B Biol Sci. 368(1610):20120404.
doi:10.1098/rstb.2012.0404.

Good BH, Desai MM. 2014. Deleterious Passengers in Adapting Populations. Genetics. 198(3):1183–
1208. doi:10.1534/genetics.114.170233.

Haller BC, Messer PW. 2019. SLiM 3: Forward Genetic Simulations Beyond the Wright–Fisher Model.
Mol Biol Evol. 36(3):632–637. doi:10.1093/molbev/msy228.

Johri P, Charlesworth B, Jensen JD. 2020. Toward an Evolutionarily Appropriate Null Model: Jointly
Inferring Demography and Purifying Selection. Genetics. 215(1):173–192.
doi:10.1534/genetics.119.303002.

Kardos M, Armstrong EE, Fitzpatrick SW, Hauser S, Hedrick PW, Miller JM, Tallmon DA, Funk WC. 2021. The crucial role of genome-wide genetic variation in conservation. Proc Natl Acad Sci. 118(48):e2104642118. doi:10.1073/pnas.2104642118.

Kelleher J, Thornton KR, Ashander J, Ralph PL. 2018. Efficient pedigree recording for fast population genetics simulation. PLOS Comput Biol. 14(11):e1006581. doi:10.1371/JOURNAL.PCBI.1006581.

Keller LF, Waller DM. 2002. Inbreeding effects in wild populations. Trends Ecol Evol. 17(5):230–241. doi:10.1016/S0169-5347(02)02489-8.

Kim BY, Huber CD, Lohmueller KE. 2017. Inference of the Distribution of Selection Coefficients for New Nonsynonymous Mutations Using Large Samples. Genetics. 206(1):345–361. doi:10.1534/genetics.116.197145.

Kondrashov AS. 2017. Crumbling Genome: The Impact of Deleterious Mutations on Humans. John Wiley & Sons.

Kyriazis CC, Wayne RK, Lohmueller KE. 2021. Strongly deleterious mutations are a primary determinant of extinction risk due to inbreeding depression. Evol Lett. 5(1):33–47. doi:10.1002/evl3.209.

Lacava J, Hughes J. 1984. Determining Minimum Viable Population Levels. Wildl Soc Bull 1973-2006. 12(4):370–376.

Lande R. 1998. Risk of population extinction from fixation of deleterious and reverse mutations. Genetica. 102(0):21–27. doi:10.1023/A:1017018405648.

Lansch-Justen L, Cusseddu D, Schmitz MA, Bank C. 2022. The extinction time under mutational meltdown driven by high mutation rates. Ecol Evol. 12(7):e9046. doi:10.1002/ece3.9046.

Lesecque Y, Keightley PD, Eyre-Walker A. 2012. A Resolution of the Mutation Load Paradox in Humans. Genetics. 191(4):1321–1330. doi:10.1534/genetics.112.140343.

Lynch M, Conery J, Bürger R. 1995a. Mutation Accumulation and the Extinction of Small Populations. Am Nat. 146(4):489–518.

Lynch M, Conery J, Bürger R. 1995b. Mutational Meltdowns in Sexual Populations. Evolution. 49(6):1067–1080. doi:10.1111/j.1558-5646.1995.tb04434.x.

Lynch M, Gabriel W. 1990. Mutation Load and the Survival of Small Populations. Evolution. 44(7):1725–1737. doi:10.1111/j.1558-5646.1990.tb05244.x.

Martin G, Aguilée R, Ramsayer J, Kaltz O, Ronce O. 2013. The probability of evolutionary rescue: towards a quantitative comparison between theory and evolution experiments. Philos Trans R Soc B Biol Sci. 368(1610):20120088. doi:10.1098/rstb.2012.0088.

Matheson J, Bertram J, Masel J. 2023. Human deleterious mutation rate implies high fitness variance, with declining mean fitness compensated by rarer beneficial mutations of larger effect. :2023.09.01.555871. doi:10.1101/2023.09.01.555871. [accessed 2024 Jan 31]. https://www.biorxiv.org/content/10.1101/2023.09.01.555871v1.

Matheson J, Masel J. 2024. Background Selection From Unlinked Sites Causes Nonindependent
Evolution of Deleterious Mutations. Genome Biol Evol. 16(3):evae050. doi:10.1093/gbe/evae050.

McCandlish DM, Epstein CL, Plotkin JB. 2015. Formal properties of the probability of fixation:
Identities, inequalities and approximations. Theor Popul Biol. 99:98–113.
doi:10.1016/j.tpb.2014.11.004.

Melissa MJ, Desai MM. 2024. A dynamical limit to evolutionary adaptation. Proc Natl Acad Sci.
121(4):e2312845121. doi:10.1073/pnas.2312845121.

Murphy DA, Elyashiv E, Amster G, Sella G. 2022. Broad-scale variation in human genetic diversity
levels is predicted by purifying selection on coding and non-coding elements. Nordborg M, Weigel D,
Nordborg M, editors. eLife. 12:e76065. doi:10.7554/eLife.76065.

Nabutanyi P, Wittmann MJ. 2022. Modeling minimum viable population size with multiple genetic
problems of small populations. Conserv Biol. 36(5):e13940. doi:10.1111/cobi.13940.

Nunney L, Campbell KA. 1993. Assessing minimum viable population size: Demography meets
population genetics. Trends Ecol Evol. 8(7):234–239. doi:10.1016/0169-5347(93)90197-W.

Olofsson P, Chipkin L, Daileda RC, Azevedo RBR. 2023. Mutational meltdown in asexual populations
doomed to extinction. J Math Biol. 87(6):88. doi:10.1007/s00285-023-02019-y.

Orr HA, Unckless RL. 2008. Population Extinction and the Genetics of Adaptation. Am Nat.
172(2):160–169. doi:10.1086/589460.

Orr HA, Unckless RL. 2014. The Population Genetics of Evolutionary Rescue. Pritchard JK, editor.
PLoS Genet. 10(8):e1004551. doi:10.1371/journal.pgen.1004551.

Osmond MM, Klausmeier CA. 2017. An evolutionary tipping point in a changing environment.
Evolution. 71(12):2930–2941. doi:10.1111/evo.13374.

Pacifici M, Foden WB, Visconti P, Watson JEM, Butchart SHM, Kovacs KM, Scheffers BR, Hole DG,
Martin TG, Akçakaya HR, et al. 2015. Assessing species vulnerability to climate change. Nat Clim
Change. 5(3):215–224. doi:10.1038/nclimate2448.

Pardo-Manuel de Villena F, Sapienza C. 2001. Nonrandom segregation during meiosis: the unfairness
of females. Mamm Genome. 12(5):331–339. doi:10.1007/s003350040003.

Park S-C, Krug J. 2007. Clonal interference in large populations. Proc Natl Acad Sci. 104(46):18135–
18140. doi:10.1073/pnas.0705778104.

Peck JR. 1994. A ruby in the rubbish: beneficial mutations, deleterious mutations and the evolution of
sex. Genetics. 137(2):597–606. doi:10.1093/genetics/137.2.597.

Pénisson S, Singh T, Sniegowski P, Gerrish P. 2017. Dynamics and Fate of Beneficial Mutations Under
Lineage Contamination by Linked Deleterious Mutations. Genetics. 205(3):1305–1318.
doi:10.1534/genetics.116.194597.

Pérez-Pereira N, Caballero A, García-Dorado A. 2022. Reviewing the consequences of genetic purging
on the success of rescue programs. Conserv Genet. 23(1):1–17. doi:10.1007/s10592-021-01405-7.

Poon A, Otto SP. 2000. Compensating for Our Load of Mutations: Freezing the Meltdown of Small Populations. Evolution. 54(5):1467–1479. doi:10.1111/j.0014-3820.2000.tb00693.x.

Ralls K, Ballou JD, Dudash MR, Eldridge MDB, Fenster CB, Lacy RC, Sunnucks P, Frankham R. 2018. Call for a Paradigm Shift in the Genetic Management of Fragmented Populations. Conserv Lett. 11(2):e12412. doi:10.1111/conl.12412.

Ralls K, Sunnucks P, Lacy RC, Frankham R. 2020. Genetic rescue: A critique of the evidence supports maximizing genetic diversity rather than minimizing the introduction of putatively harmful genetic variation. Biol Conserv. 251:108784. doi:10.1016/j.biocon.2020.108784.

Reed DH, O'Grady JJ, Brook BW, Ballou JD, Frankham R. 2003. Estimates of minimum viable population sizes for vertebrates and factors influencing those estimates. Biol Conserv. 113(1):23–34. doi:10.1016/S0006-3207(02)00346-4.

Rout TM, Hauser CE, Possingham HP. 2007. Minimise long-term loss or maximise short-term gain?: Optimal translocation strategies for threatened species. Ecol Model. 201(1):67–74. doi:10.1016/j.ecolmodel.2006.07.022.

Ruan Y, Wang H, Zhang L, Wen H, Wu C-I. 2020. Sex, fitness decline and recombination – Muller's ratchet vs. Ohta's ratchet. :2020.08.06.240713. doi:10.1101/2020.08.06.240713. [accessed 2024 Jan 31]. https://www.biorxiv.org/content/10.1101/2020.08.06.240713v1.

Santidrián Tomillo P, Wallace BP, Paladino FV, Spotila JR, Genovart M. 2021. Short-term gain, long-term loss: How a widely-used conservation tool could further threaten sea turtles. Biol Conserv. 261:109260. doi:10.1016/j.biocon.2021.109260.

Schultz ST, Lynch M. 1997. Mutation and Extinction: The Role of Variable Mutational Effects, Synergistic Epistasis, Beneficial Mutations, and Degree of Outcrossing. Evolution. 51(5):1363–1371. doi:10.1111/j.1558-5646.1997.tb01459.x.

Shaffer ML. 1981. Minimum Population Sizes for Species Conservation. BioScience. 31(2):131–134. doi:10.2307/1308256.

Shaw FH, Baer CF. 2011. Fitness-dependent mutation rates in finite populations. J Evol Biol. 24(8):1677–1684. doi:10.1111/j.1420-9101.2011.02320.x.

Spielman D, Brook BW, Frankham R. 2004. Most species are not driven to extinction before genetic factors impact them. Proc Natl Acad Sci. 101(42):15261–15264. doi:10.1073/pnas.0403809101.

Stephan W. 2010. Genetic hitchhiking versus background selection: the controversy and its implications. Philos Trans R Soc B Biol Sci. 365(1544):1245–1253. doi:10.1098/rstb.2009.0278.

Tanaka Y. 2000. Extinction of populations by inbreeding depression under stochastic environments. Popul Ecol. 42(1):55–62. doi:10.1007/s101440050009.

Traill LW, Bradshaw CJA, Brook BW. 2007. Minimum viable population size: A meta-analysis of 30 years of published estimates. Biol Conserv. 139(1):159–166. doi:10.1016/j.biocon.2007.06.011.

Uecker H, Hermisson J. 2016. The Role of Recombination in Evolutionary Rescue. Genetics. 202(2):721–732. doi:10.1534/genetics.115.180299.

Uecker H, Otto SP, Hermisson J. 2014. Evolutionary Rescue in Structured Populations. Am Nat.
183(1):E17–E35. doi:10.1086/673914.

Vucetich JA, Nelson MP, Bruskotter JT. 2017. Conservation Triage Falls Short Because Conservation
Is Not Like Emergency Medicine. Front Ecol Evol. 5. doi:10.3389/fevo.2017.00045. [accessed 2024
Aug 28]. https://www.frontiersin.org/journals/ecology-and-
evolution/articles/10.3389/fevo.2017.00045/full.

Wang Y, Obbard DJ. 2023. Experimental estimates of germline mutation rate in eukaryotes: a
phylogenetic meta-analysis. Evol Lett. 7(4):216–226. doi:10.1093/evlett/qrad027.

Whiteley AR, Fitzpatrick SW, Funk WC, Tallmon DA. 2015. Genetic rescue to the rescue. Trends Ecol
Evol. 30(1):42–49. doi:10.1016/j.tree.2014.10.009.

Whitlock MC. 2000. Fixation of New Alleles and the Extinction of Small Populations: Drift Load,
Beneficial Alleles, and Sexual Selection. Evolution. 54(6):1855–1861.

Whitlock MC, Griswold CK, Peters Andrew D, Whitlock M C, Griswold CK&, Peters A D. 2003.
Compensating for the meltdown: The critical effective size of a population with deleterious and
compensatory mutations.

Willis KJ, Birks HJB. 2006. What Is Natural? The Need for a Long-Term Perspective in Biodiversity
Conservation. Science. 314(5803):1261–1265. doi:10.1126/science.1122667.

Wilson BA, Pennings PS, Petrov DA. 2017. Soft Selective Sweeps in Evolutionary Rescue. Genetics.
205(4):1573–1586. doi:10.1534/genetics.116.191478.

Zeyl C, Mizesko M, De Visser JAGM. 2001. Mutational Meltdown in Laboratory Yeast Populations.
Evolution. 55(5):909–917. doi:10.1111/j.0014-3820.2001.tb00608.x.