Shared and unique lifetime stressor characteristics and brain networks predict
adolescent anxiety and depression

Yueyue Lydia Qu[1,2], Sidhant Chopra[3,4], Shijie Qu[1,2], Carrisa V. Cocuzza[5], Loïc Labache[5],
Clemens C.C. Bauer [6,7,8], Francesca Morfini[6,7], Susan Whitfield-Gabrieli[6,7,8], George M. Slavich[9],
Jutta Joormann[1,2], & Avram J. Holmes[5]

[1] Department of Psychology, Yale University, New Haven, CT, USA
[2] Wu Tsai Institute, Yale University, New Haven, CT, USA
[3] Orygen, Melbourne, VIC, Australia
[4] Centre for Youth Mental Health, The University of Melbourne, Melbourne, VIC, Australia
[5] Department of Psychiatry, Brain Health Institute, Rutgers University, Piscataway, NJ, USA
[6] Department of Psychology, Northeastern University, Boston, MA, USA
[7] Center for Cognitive & Brain Health, Northeastern University, Boston, MA, USA
[8] Department of Brain and Cognitive Sciences and McGovern Institute for Brain Research,
Massachusetts Institute of Technology, Cambridge, MA, USA
[9] Department of Psychiatry and Biobehavioral Sciences, University of California, Los Angeles,
CA, USA

Correspondence: Yueyue Lydia Qu, lydia.qu@yale.edu
Avram J. Holmes, avram.holmes@rutgers.edu

Key Words: Major life stressor characteristics, resting-state functional networks, longitudinal
prediction, anxiety, depression, adolescence

Abstract

Background

Exposure to major life stressors and aberrant brain functioning have been linked to anxiety and

depression, especially during periods of heighted functional brain plasticity, such as

adolescence. However, it remains unclear if specific characteristics of major life stressors and

functional network disruptions differentially predict anxiety and depression symptoms over time

and, if so, whether they act independently or jointly.

Methods

We collected baseline lifetime stressor exposure data and resting-state functional magnetic

resonance imaging data in a longitudinal sample of 107 adolescents enriched for anxiety and

depressive disorders. We examined five stressor characteristics: physical danger, interpersonal

loss, humiliation, entrapment, and role change/disruption. Anxiety and depression symptoms

were assessed at baseline, 6-month and 12-month follow-ups. Linear mixed effect models

tested if these stressor characteristics, functional connectivity within and between frontoparietal,

default, and ventral attention networks, and their interactions differentially predicted anxiety and

depression symptoms at 6-month and 12-month follow-ups.

Results

Greater lifetime severity of physical danger and humiliation prospectively predicted increased

anxiety symptoms at both follow-ups, whereas greater lifetime entrapment severity prospectively

predicted higher anxiety and depression symptoms. Only the effects of lifetime entrapment

severity were robust to including within- and between-network functional connectivity metrics

and other significantly predictive stressor characteristics. Lifetime entrapment severity more

strongly predicted anxiety symptoms in youth with higher default network connectivity. Greater

functional connectivity between frontoparietal and default networks prospectively predicted

increased depression symptoms.

Conclusions

Taken together, these results underscore the critical importance of using stressor characteristics

and functional connectivity jointly to study predictors for adolescent anxiety and depression.

Introduction

Exposure to major life stressors is a strong risk factor for the onset and subsequent
recurrence of affective disorders (1–5), especially in adolescence when there is increased brain
plasticity and heightened vulnerability to the emergence of psychopathology (6–10). However,
stressors come in many different forms and are diverse with respect to both their characteristics
and associations with clinical symptoms (11,12). Results from large-scale, prospective cohort
studies have suggested that certain characteristics of major life stressors may preferentially
increase risk for specific clinical outcomes. For instance, stressors that involve devaluation of
the self, such as interpersonal loss and humiliation, are theorized to preferentially heighten risk
for depression (13–17). In contrast, stressors marked by a threat to one's physical integrity,
such as danger, are theorized to be stronger predictors of anxiety (13,15,16,18). Stressors
characterized by feelings of failure without any means of escape, such as entrapment, predict
both anxiety and depression (13,19). These patterns may be explained by the cognitive content-
specificity hypothesis of anxiety and depression which posits that anxiety and depression can
be discriminated by distinct forms of negative beliefs (20) and which has been partially
supported (21,22). However, the potential neural mechanisms linking distinct stressor
characteristics with adolescent anxiety and depression remain unknown.

Major life stressors can induce long-lasting effects on neurobiological functioning (23–27),
particularly when they occur during childhood and adolescence (24,25,28–30). Prior neuroimaging
studies have associated changes in brain functioning during monetary reward and emotional face
tasks with exposure to life stressors (31–34). Altered brain functioning is also evident when
considering resting-state functional connectivity (RSFC) patterns within and between the
frontoparietal (35), default (36,37), and salience/ventral attention (38) networks among people
who were exposed to early-life stress and trauma (28,39–42). Because altered connectivity within
the frontoparietal, default and ventral attention networks are theorized to underlie dysfunctional
cognitive (43–45), self-referential (44–46) and salience processing (44,47) respectively in anxiety
and depression, it is possible that disruptions within and between these functional networks
moderate the process of stress exposure leading to psychopathology. Although prior research has
examined the extent to which intrinsic patterns of functional connectivity predict adolescent
anxiety and depression following general stress exposure (48,49), these studies have not
considered the interactions between neurobiological factors and life experience in differentiating
anxiety and depression. Longitudinal associations linking distinct dimensions of life stressor
exposure, network connectivity, and clinical symptoms thus remain to be established.

Present Study

      To address these gaps in knowledge, we first conducted hypothesis-driven analyses investigating if distinct characteristics of major life stressors occurring over the entire life course differentially predicted anxiety and depression symptoms at 6-month and 12-month follow-ups. We then explored if total lifetime severity of any stressor characteristics, RSFC metrics within and between the frontoparietal, default, and ventral attention networks, and their interactions predicted anxiety and depression symptoms at both follow-ups. These analyses were conducted in a longitudinal sample of adolescents recruited from school-based and hospital-based child treatment programs (50). Most participants had a current diagnosis of at least one anxiety or depressive disorder at the time of baseline assessment. Based on the prior research summarized above, we tested three hypotheses using linear mixed-effects (LME) models: (a) greater lifetime severity of physical danger at baseline would differentially predict higher anxiety symptoms at both follow-ups; (b) greater lifetime severity of interpersonal loss and humiliation at baseline would differentially predict higher depression symptoms at both follow-ups; (c) greater lifetime severity of entrapment at baseline would predict higher levels of both depression and anxiety symptoms at both follow-ups. In further exploratory analyses, we assessed whether any stressor characteristics and RSFC metrics within and between the three large-scale networks predict anxiety/depression symptoms at two 6-month follow-ups, and whether these RSFC metrics would moderate the strengths of prospective associations between stressor characteristics and anxiety/depression symptoms.

Methods and Materials

Participants

Data were collected from 215 adolescents (M_{age} = 15.44, range = 14-17 years old) enrolled in the Boston Adolescent Neuroimaging of Depression and Anxiety (BANDA) study (50) and assessed at 6-month intervals after the initial visit for up to 12 months. The present analyses were restricted to data from the baseline, 6-month follow-up and 12-month follow-up assessments. Resting-state functional magnetic resonance imaging (rsfMRI) data was available for 203 participants at baseline (50). Out of these 203 adolescents, 107 had available self-reported anxiety and depression symptoms at baseline, 6- and 12- month follow-up assessments and were included in the final analytical sample (**Table 1; Figure 1**). Of these 107 participants, 62% (n = 66) had a current diagnosis of at least one anxiety or depressive disorder. These diagnoses were given by a blinded, licensed clinical psychologist based on the Diagnostic and Statistical Manual of Mental Health Disorders, 5[th] Edition (DSM-5; 51) and reached moderate to substantial inter-rater agreement (50).

127 **Table 1.** Demographic characteristics for the final analytical sample ($n = 107$)

Baseline		
Demographics	*M/n*	*SD/%*
Age, mean ± SD	15.42	0.87
Sex, (% female)	66	61.68
Diagnostic group	*n*	*%*
Anxiety [a]	38	35.51
Control	41	38.32
Depression [b]	28	26.17
Race/ethnicity	*n*	*%*
African American	2	1.87
Asian	2	1.87
Hawaiian	1	0.93
White	102	95.33
Lifetime severity of stressor characteristics [c]	*M*	*SD*
Entrapment	11.21	10.30
Humiliation	7.60	7.18
Interpersonal loss	10.59	8.48
Physical danger	4.50	5.50
Role change/disruption	5.82	5.44
Clinical symptoms	*M*	*SD*
Anxiety symptoms [d]	24.36	17.86
Depression symptoms [e]	15.74	14.82
Six-month follow-up		
Clinical symptoms	*M*	*SD*
Anxiety symptoms [d]	23.81	17.71
Depression symptoms [e]	16.84	15.43
Twelve-month follow-up		
Clinical symptoms	*M*	*SD*
Anxiety symptoms [d]	21.35	15.61
Depression symptoms [e]	14.62	14.13

Note: [a] Anxiety = having a current diagnosis of at least one anxiety disorder and no depressive

disorder based on DSM-5. [b] Depression = having a current diagnosis of at least one depressive

disorder based on DSM-5. [c] The total lifetime severity for each social-psychological stressor

characteristic based on self-reported acute and chronic stressors from the Stress and Adversity

Inventory for Adolescents (STRAIN). Entrapment: 0-26. Humiliation: 0-35. Interpersonal loss: 0-

50. Physical danger: 0-65. Role change/disruption: 0-70 [d] Anxiety symptoms was computed by

summing the four anxiety subscales (Separation Anxiety Disorder, Social Phobia, Generalized

Anxiety and Panic Disorder) from the Revised Children's Anxiety and Depression Scale

(RCADS), ranging from 0 to 93. [e] Depression symptoms was computed by the total Mood and

Feelings Questionnaire (MFQ) score ranging from 0 to 66.

Figure 1. Schematic representation of the Boston Adolescent Neuroimaging of Depression and Anxiety (BANDA) study. Participants were recruited across three sites and underwent four study sessions, including in-person clinician evaluations, self-report measures, neuroimaging, and online and in-person follow-up assessments. The two clinical and imaging baseline visits occurred within two weeks from each other. Participants completed an online batter of self-report questionnaires 6 months after their second baseline visit. Finally, they went through on-site clinical evaluations and completed an additional batter of self-report questionnaires 12 months after their second baseline visit. Picture icons were downloaded from www.freepik.com.

Measures of social-psychological characteristics of life stressors

Exposure to acute and chronic stressors occurring over the entire life course was

assessed using the Stress and Adversity Inventory for Adolescents (STRAIN) (52). The STRAIN

measured the total lifetime severity for each social-psychological stressor characteristic,

including physical danger, interpersonal loss, humiliation, entrapment, and role

change/disruption, for each participant based on their self-reported acute and chronic stressors

(see https://www.strainsetup.com). Examples of stressors linked to each of these five

characteristics have been described elsewhere (52). Total lifetime severity scores for these five

characteristics were used to predict the levels of anxiety and depression symptoms at two six-

month follow-up assessments. Brief definitions for these stressor characteristics are as follows

(13,16,53,54):

Physical danger. The degree of potential future threat to one's physical safety that

might occur as a result of the stressor.

Interpersonal loss. Diminution of a sense of connectedness or well-being as a result of

a real or realistically imagined loss of a person by death or by separation.

Humiliation. The likelihood of a stressor rendering a person devalued in relation to

others or self, usually due to rejection or a sense of core failure.

Entrapment. Ongoing circumstances of marked difficulty of at least 6 months' duration

that the individual can reasonably expect to persist or get worse, with little or no possibility that a

resolution can be achieved as a result of anything that might reasonably be done.

Role change/disruption. Life transitions that involve addition, subtraction or change of

social roles.

The STRAIN has demonstrated excellent test-retest reliability, good concurrent and

discriminant validity, as well as predictive utility in relation to various clinical outcomes including

anxiety and depression (52,55).

Measures of anxiety and depression symptoms

Self-reported anxiety symptoms in this study were assessed by the Revised Children's

Anxiety and Depression Scale (RCADS) (56). The RCADS has exhibited excellent internal

consistency, test-retest reliability, convergent and discriminant validity (56). The RCADS have

six subscales: separation anxiety disorder, social phobia, generalized anxiety, panic disorder,

obsessive-compulsive disorder and low mood. Total anxiety symptoms for each participant were

computed by summing the four anxiety subscales (Separation Anxiety Disorder, Social Phobia,

Generalized Anxiety and Panic Disorder).

Self-reported depression symptoms were assessed by the Mood and Feelings
Questionnaire (MFQ) (57,58). Prior studies have found the MFQ to be a reliable and valid
measure of adolescent depression in both clinical and non-clinical samples across different
populations (59–62). We used the total MFQ score as the measure of depression symptoms for
each participant.
186
Neuroimaging
Data acquisition and processing
Functional and anatomical neuroimaging data were acquired at baseline assessment
using a 3-Tesla Siemens Prisma scanner with a 2D multi-band gradient-recalled echo-planar
imaging (EPI) sequence. Each participant underwent four 5.8-minute resting-state functional
MRI (rsfMRI) runs, consisting of two runs with opposite phase encoding directions (AP/PA).
Each rsfMRI scan was acquired using 2mm isotropic resolution and a TR of 800ms. Full details
of the acquisition protocol can be found elsewhere (63).
The acquired rsfMRI data then went through the previously established Human
Connectome Project (HCP) minimal preprocessing pipelines (64). Minimally preprocessed T1w
images (64) went through bias- and distortion- correction using the *PreFreeSurfer* pipeline and
registered to MNI space. Cortical surface reconstruction was conducted using FreeSurfer v5.2
using recon-all adapted for high-resolution images. The reconstructed surface meshes were then
registered to the Conte69 surface template (65). During preprocessing, the fMRI data were first
corrected for gradient-nonlinearity-induced distortions. The fMRI time series in each frame were
then realigned to the single-band reference image to correct for subject motion using rigid body
transformation (66,67) with FSL. The resulting single-band image underwent spline interpolation
to correct for distortions and was then registered to the T1w image (68). The registered fMRI
volumes then went through nonlinear registration to the Conte69 surface template (65) and
mapped to the standard CIFTI grayordinate coordinate space. Further details about the HCP
minimal preprocessing pipelines of structural and functional images can be found elsewhere (64).
The minimally preprocessed fMRI data for each run were then denoised using ICA+FIX (69,70)
pre-trained using HCP_hp2000.RData and aligned across participants using MSMAll multi-modal
surface registration (71,72).
211
Resting-state functional connectivity
We defined 400 cortical regions of interest (ROIs) using a previously validated atlas (73).
Resting-state functional connectivity (RSFC) was measured by Pearson's *r* correlations

between the mean time series of each pair of ROIs. The average FC matrix across all runs in

each participant was computed after applying Fisher Z-transformation and used for subsequent

analyses.

Specifically, this study focused on RSFC within and between the frontoparietal, default,

and ventral attention networks (**Figure 2**) according to the 17-network solution (74) as

predictors of anxiety and depression symptoms levels at two 6-month follow-ups. Within-

network connectivity was assessed by averaging the pairwise RSFC of all regions assigned to

that network, resulting in three within-network connectivity values per individual. "Between"

network connectivity was assessed by computing the pair-wise correlations of each ROI in one

network (e.g., frontoparietal) to each ROI in the other network (e.g., default) and averaging

across them, resulting in three between network connectivity values per individual.

Figure 2. The functional network organization of the human cerebral cortex revealed through intrinsic functional connectivity. Colors reflect regions estimated to be within the same network. Cortical regions-of-interest (ROIs) defined by Schaefer's parcellation (73) and assigned to frontoparietal (yellow), default (red), and salience/ventral attention (purple) networks (74).

Covariates

The following covariates were dummy coded, converted to factors and entered into each LME model: participant's race (White: Yes=1, No=0; African American: Yes=1, No=0; Asian: Yes=1, No=0; Hawaiian: Yes=1, No=0), ethnicity (Hispanic: Yes=1, No=0), sex (female=0, male=1), participant's current diagnostic group (Depression [having a current diagnosis of at least one depressive disorder]: Yes=1, No=0; Anxiety [having a current diagnosis of at least one anxiety disorder and no depressive disorder]: Yes=1, No=0; Control [having no current or lifetime diagnosis of any psychiatric disorder]: Yes=1, No=0). We included current diagnosis of depressive and anxiety disorders as covariates because there were significant differences in depression and anxiety symptoms severity as well as total lifetime severity of stressor characteristics across the three diagnostic groups (**Supplemental Result 1**). Participant's age at scan was entered as a continuous covariate in each LME model. We additionally included baseline depression symptoms in each LME model predicting anxiety symptoms and baseline anxiety symptoms in each LME model predicting depression symptoms as covariates to parse out unique predictors of each symptom (i.e., anxiety, depression) over time.

Statistical analyses

We constructed linear mixed-effect (LME) models using the *lme4* package in Rv4.2.0 (75) with restricted maximum likelihood estimation (REML) to test our hypotheses. Each hypothesis-driven LME model assessed if total lifetime severity for each stressor social-psychological characteristic (i.e., physical danger, interpersonal loss, humiliation, entrapment) at baseline predicted each symptom (i.e., anxiety, depression) at two 6-month follow-ups. We used the *cAIC4* package *v1.0* (76) to determine whether each of these models yielded lower conditional Akaike information criterion (cAIC) when fitted with a random intercept or with a random slope plus a random intercept. We also included covariates in each model to test if the fixed effects of these stressor characteristics were robust to the inclusion of potential confounders. Continuous predictors, covariates and outcome variables were standardized to make the beta estimates more interpretable and to avoid multicollinearity.

Finally, we included the total lifetime severity of all stressor characteristics, all RSFC metrics and their interactions in a single "unified LME model" predicting each symptom (i.e., anxiety, depression) at two 6-month follow-ups. In each unified LME model, we explored if total lifetime severity of any stressor characteristics, any RSFC metrics within and between the frontoparietal, default, and ventral attention networks, and their interactions emerged as significant predictors of each symptom. Each unified LME model was fitted either with a random

intercept or with a random intercept plus a random slope. Potential confounders were included

in each unified LME model and all continuous variables were standardized. As an example, the

formula for the unified LME model predicting prospective anxiety symptoms with a random slope

and intercept was:

$AnxietySx \sim (PhysicalDanger + InterpersonalLoss + Humiliation + Entrapment +$

$RoleReversal) * (RSFCwithinFPN + RSFCwithinDN + RSFCwithinVAN + RSFCbetwFPN\text{-}DN +$

$RSFCbetwFPN\text{-}VAN + RSFCbetwDN\text{-}VAN) + DepressionDx + AnxietyDx +$

$BaselineDepressionSx + Race + Ethnicity + BaselineAge + sex + time + (1 + time | Subject).$

The unified LME model predicting prospective depression symptoms with only a random

intercept was:

$DepressionSx \sim (PhysicalDanger + InterpersonalLoss + Humiliation + Entrapment +$

$RoleReversal) * (RSFCwithinFPN + RSFCwithinDN + RSFCwithinVAN + RSFCbetwFPN\text{-}DN +$

$RSFCbetwFPN\text{-}VAN + RSFCbetwDN\text{-}VAN) + DepressionDx + AnxietyDx +$

$BaselineAnxietySx + Race + Ethnicity + BaselineAge + sex + time + (1 | Subject).$

Since total lifetime severity scores for the five stressor characteristics were highly

intercorrelated (**Supplemental Table 1**), as expected, the resulting unified LME models may

exhibit high multicollinearity (VIF≥5) despite standardization of predictor and outcome variables.

Hence, for each unified LME model, we started with including all five stressor characteristics,

iteratively removing different subsets of one to four stressor characteristics, rerunning the model

and recomputing cAIC. We selected the optimal subset of stressor characteristics yielding the

lowest conditional Akaike information criterion (cAIC) for each unified LME model. The

conditional R^2 captured by each optimal unified LME model was determined using the

performance package (77) in Rv4.2.0.

 Results

 Total lifetime severity of stressor characteristics differentially predicts prospective

 anxiety and depression symptoms

Across all hypothesis-driven LME models predicting anxiety symptoms at two 6-month

follow-ups, the models fitted with random slopes plus random intercepts yielded lower cAICs

than the corresponding models fitted with only random intercepts (**Supplemental Table 3**). On

the other hand, all hypothesis-driven LME models predicting prospective depression symptoms

were singular when fitted with both random slopes and random intercepts (**Supplemental Table**

4). Therefore, we will focus on results from LME models fitted with random slopes and

intercepts when prospective anxiety symptoms is the predicted variable and focus on results

from LME models fitted with only random intercepts when prospective depression symptoms is

the predicted variable.

Consistent with our hypothesis that lifetime physical danger severity is a specific

predictor of anxiety, the LME analyses revealed that greater lifetime severity of physical danger

at baseline predicted higher anxiety ($\beta = 0.25,\ p = 3.63 \times 10^{-5}$; **Supplemental Table 4**) but not

depression symptoms ($\beta = -0.0079,\ p = 0.90$; **Supplemental Table 5**) at follow-up, after

accounting for potentially confounding covariates.

Failing to support our second hypothesis theorizing interpersonal loss as a specific

predictor of depression, the second set of hypothesis-driven LME models indicated that the

main effects of lifetime severity of interpersonal loss on both anxiety ($\beta = 0.039,\ p = 0.55$;

Supplemental Table 6) and depression symptoms were not robust to the inclusion of

potentially confounding covariates ($\beta = 0.057,\ p = 0.29$; **Supplemental Table 7**). These results

suggest that individual differences in the lifetime severity of interpersonal loss does not predict

prospective depression symptoms above and beyond anxiety symptoms and depression

diagnosis at baseline.

Contrary to our third hypothesis, the third set of hypothesis-driven LME models revealed

that greater lifetime severity of humiliation at baseline predicted higher anxiety ($\beta = 0.31,\ p =$

2.56×10^{-6}; **Supplemental Table 8**) but not depression symptoms ($\beta = 0.034,\ p = 0.62$;

Supplemental Table 9) at follow-up, after accounting for potentially confounding covariates.

These data suggest that lifetime severity of humiliation may be a specific risk factor for anxiety

rather than depression.

Finally, the fourth set of hypothesis-driven LME models revealed that greater total

lifetime severity of entrapment predicted both anxiety ($\beta = 0.39,\ p = 5.20 \times 10^{-8}$; **Supplemental**

Table 10) and depression symptoms at two 6-month follow-ups, after accounting for potentially

confounding covariates ($\beta = 0.15, \ p = 0.05$; **Supplemental Table 11**). These results suggest

that lifetime severity of entrapment may be a shared risk factor for both anxiety and depression.

Total lifetime severity of stressor characteristics, large-scale brain networks and

symptoms of anxiety and depression

Next, we sought to determine if prospective associations between stressor

characteristics and clinical symptoms are moderated by patterns of functional connectivity within

and between the large-scale brain networks theorized to underlie the expression of affective

illness. To this end, we used two unified LME models to explore if total lifetime severity of any

stressor characteristics, any RSFC metrics within and between the frontoparietal, default, and

ventral attention networks, and their interactions emerged as significant predictors of

prospective anxiety and depression symptoms. The unified LME model predicting prospective

anxiety symptoms yielded the lowest cAIC when fitted with both a random slope and intercept

and when total lifetime severity for physical danger and entrapment were entered as predictors

along with all RSFC metrics (**Supplemental Table 12**). The optimal unified model for anxiety

had a conditional R^2 of 0.873. The unified LME model predicting prospective depression

symptoms yielded the lowest cAIC when fitted with only a random intercept and when total

lifetime severity entrapment was the only stressor characteristic in the model (**Supplemental**

Table 13). The optimal unified model for depression had a conditional R^2 of 0.759. Below, we

focus on the results from the optimal unified LME models.

Results from the optimal unified model for anxiety revealed that total lifetime severity of

entrapment at baseline was positively associated with anxiety symptoms at two follow-up

assessments ($\beta = 0.36, \ p = 3.10 \times 10^{-4}$), and the association was stronger when RSFC within

the default network was higher (**Table 2; Figure 3A&C**). When considering depression, the

optimal unified model indicated that RSFC between the frontoparietal and default network

($\beta = 0.18, \ p = 8.20 \times 10^{-4}$) and total lifetime severity of entrapment stressor exposure ($\beta = 0.17,$

$p = 0.05$) at baseline were positively associated with depression symptoms levels at two follow-

up assessments (**Table 3; Figure 3B&D**). These results demonstrate that entrapment severity

remained to be a shared predictor for both anxiety and depression even after accounting for

main and interaction effects involving within- and between-network RSFC metrics, further

supporting its importance in predicting both anxiety and depression.

Table 2. Results of unified LME models testing if anxiety symptoms at two 6-month follow-ups can be predicted from total lifetime severity of physical danger, total lifetime severity of entrapment, and RSFC within and between functional networks at baseline

Predictor	Estimate	SE	95% CI		t	p
			LL	UL		
wFPN FC	-0.057	0.065	-0.17	0.053	-0.87	0.39
wVAN FC	0.076	0.063	-0.029	0.18	1.22	0.23
wDN FC	-0.024	0.060	-0.13	0.082	-0.40	0.69
bFPN-DN FC	-0.12	0.070	-0.24	-0.0014	-1.71	0.091
bFPN-VAN FC	0.089	0.059	-0.0096	0.19	1.52	0.13
bDN-VAN FC	-0.023	0.053	-0.11	0.066	-0.43	0.67
Physical danger	0.083	0.086	-0.062	0.23	0.97	0.33
Entrapment	**0.36**	**0.095**	**0.20**	**0.52**	**3.78**	**0.0031*****
Baseline depression symptoms	**0.41**	**0.093**	**0.25**	**0.57**	**4.42**	**3.14×10^{-5} ******
Diagnosis of Depression	-0.032	0.18	-0.34	0.28	-0.17	0.86
Diagnosis of Anxiety	**0.46**	**0.13**	**0.24**	**0.67**	**3.59**	**0.00057******
White	0.016	0.51	-0.84	0.87	0.032	0.97
African American	0.19	0.62	-0.86	1.24	0.30	0.76
Asian	-0.060	0.62	-1.10	0.98	-0.097	0.92
Ethnic group	-0.28	0.20	-0.62	0.065	-1.36	0.18
Baseline age	0.072	0.054	-0.021	0.17	1.34	0.18
Sex	**-0.30**	**0.11**	**-0.49**	**-0.11**	**-2.70**	**0.0085****
Time	**-0.088**	**0.037**	**-0.16**	**-0.015**	**-2.37**	**0.020***
wFPN FC:Physical danger	0.17	0.094	0.0076	0.33	1.77	0.080
wFPN FC:Entrapment	-0.15	0.086	-0.30	-0.010	-1.80	0.076
wVAN FC:Physical danger	-0.029	0.069	-0.15	0.088	-0.42	0.68
wVAN FC:Entrapment	0.012	0.076	-0.12	0.14	0.15	0.88
wDN FC:Physical danger	-0.10	0.064	-0.22	0.0074	-1.63	0.11
wDN FC:Entrapment	**0.27**	**0.077**	**0.14**	**0.40**	**3.52**	**0.00073******
bFPN-DN FC:Physical danger	-0.0039	0.12	-0.21	0.20	-0.033	0.97
bFPN-DN FC:Entrapment	-0.022	0.11	-0.21	0.16	-0.21	0.84
bFPN-VAN FC:Physical danger	-0.14	0.088	-0.29	0.0026	-1.65	0.10

bFPN-VAN FC:Entrapment	0.10	0.094	-0.058	0.26	1.07	0.29
bDN-VAN FC:Physical danger	0.045	0.086	-0.10	0.19	0.52	0.60
bDN-VAN FC:Entrapment	0.048	0.066	-0.063	0.16	0.73	0.47
Intercept	-0.024	0.51	-0.88	0.83	-0.047	0.96

364 Note. *$p < 0.05$; **$p < 0.01$; ***$p < 0.005$; ****$p < 0.001$. FC = functional connectivity. bFPN-DN =

365 between frontoparietal and default network, bDN-VAN = between default network and ventral

366 attention network. bFPN-VAN = between frontoparietal and ventral attention network.

367 wFPN = within frontoparietal network. wDN = within default network. wVAN = within ventral

368 attention network

Table 3. Results of the unified LME model testing if depression symptoms at two 6-month follow-ups can be predicted from total lifetime severity of entrapment and RSFC within and between functional networks at baseline

Predictor	Estimate	*SE*	95% CI		*t*	*p*
			LL	*UL*		
wFPN FC	0.089	0.065	-0.025	0.20	1.37	0.17
wVAN FC	-0.086	0.061	-0.19	0.021	-1.41	0.16
wDN FC	0.088	0.059	-0.015	0.19	1.50	0.14
bFPN-DN FC	**0.18**	**0.067**	**0.064**	**0.30**	**2.71**	**0.0082****
bFPN-VAN FC	-0.049	0.056	-0.15	0.050	-0.87	0.39
bDN-VAN FC	0.0066	0.053	-0.087	0.10	0.12	0.90
Entrapment	**0.17**	**0.086**	**0.023**	**0.32**	**2.02**	**0.046***
Baseline anxiety symptoms	**0.45**	**0.091**	**0.29**	**0.61**	**4.98**	**3.35×10^{-6}******
Diagnose of Depression	**0.58**	**0.16**	**0.31**	**0.86**	**3.71**	**0.000372******
Diagnosis of Anxiety	-0.028	0.14	-0.27	0.22	-0.20	0.84
White	-0.55	0.51	-1.45	0.34	-1.08	0.28
African American	0.50	0.64	-0.62	1.61	0.78	0.44
Asian	-0.60	0.63	-1.71	0.52	-0.94	0.35
Ethnic group	0.25	0.21	-0.11	0.62	1.22	0.22
Baseline age	**0.12**	**0.053**	**0.031**	**0.22**	**2.34**	**0.022***
Sex	-0.048	0.11	-0.25	0.15	-0.41	0.68
Time	-0.038	0.034	-0.11	0.029	-1.11	0.27
wFPN FC:Entrapment	-0.070	0.068	-0.19	0.048	-1.04	0.30
wVAN FC:Entrapment	-0.025	0.065	-0.14	0.089	-0.39	0.70
wDN FC:Entrapment	0.015	0.061	-0.092	0.12	0.25	0.80
bFPN-DN FC:Entrapment	0.056	0.086	-0.096	0.21	0.65	0.52
bFPN-VAN FC:Entrapment	-0.068	0.056	-0.17	0.031	-1.21	0.23
bDN-VAN FC:Entrapment	-0.0026	0.055	-0.098	0.093	-0.047	0.96
Intercept	0.39	0.51	-0.51	1.29	0.76	0.45

Note. *p < 0.05; **p < 0.01; ***p < 0.005; ****p < 0.001. FC = functional connectivity. bFPN-DN = between frontoparietal and default network, bDN-VAN = between default network and ventral attention network. bFPN-VAN = between frontoparietal and ventral attention network.

375   wFPN = within frontoparietal network. wDN = within default network. wVAN = within ventral

376   attention network

Figure 3. (A) The association between total lifetime severity of entrapment and the sums of
marginal fit and conditional residuals from the fitted LME models predicting anxiety symptoms
depression symptoms at two 6-month follow-ups; (B) The association between total lifetime
severity of entrapment and anxiety symptoms at different levels of functional connectivity within
the default network. The association between (C) total lifetime severity of entrapment, (D)
functional connectivity between the frontoparietal and default networks and the sums of
marginal fit and conditional residuals from the fitted LME models predicting depression
symptoms at two 6-month follow-ups. Different colors represent different participants.

Discussion

Despite a wealth of prior research examining associations between stress, neurobiology

and internalizing psychopathology, it remains unknown if the lifetime severity of distinct life

stressor characteristics differentially predicts future anxiety and depression in adolescence, and

if such prospective associations are moderated by large-scale network connectivity. To

investigate, we acquired functional neuroimaging at baseline and tracked a sample of

adolescents longitudinally for one year, more than half of whom were currently diagnosed with at

least one depressive or anxiety disorder at time of initial assessment. We first tested our

hypotheses associating lifetime severity of specific stressor characteristics differentially with

anxiety or depression symptoms at two 6-month follow-up. We then explored if distinct stressor

characteristics and RSFC within and between large-scale networks acted independently or jointly

to predict prospective anxiety and depression symptoms.

Results from the hypothesis-driven LME models revealed that (a) higher total lifetime

severity of physical danger and humiliation predicted higher levels of anxiety but not depression

symptoms; (b) higher total lifetime severity of entrapment predicted higher levels of both anxiety

and depression symptoms; and (c) total lifetime severity of interpersonal loss predicted neither

anxiety or depression symptoms. Results from the exploratory unified LME models additionally

included fixed and interaction effects of RSFC metrics within and between large-scale functional

networks and showed that (a) the main effects of higher total lifetime severity of entrapment still

predicted higher levels of both anxiety and depression symptoms at the two 6-month follow-ups;

(b) more positive RSFC between the frontoparietal and the default networks uniquely predicted

higher levels of depression symptoms at the two 6-month follow-ups; and (c) the association

between total lifetime severity of entrapment and anxiety symptoms at the two 6-month follow-

ups was more pronounced (i.e., more positive) among adolescents who had more positive

RSFC within the default network. These results thus support the importance of including major

life stressors, as well as both RSFC within and between large-scale functional networks, in

models aiming to predict changes in anxiety and depression in adolescence over time.

The results from the hypothesis-driven LME models are consistent with the prior studies

suggesting that exposure to major life stressors characterized by danger, which implies threat to

one's physical integrity, is a specific risk factor for anxiety (13,15,16,18). However, these data

did not reveal unique associations between loss, humiliation, and depression, as has been

previously theorized (13–17). Such a discrepancy may have arisen from differences in sample

characteristics. For example, although most prior studies have focused on adults recruited from

the community, we used an adolescent sample, around 60% of whom were already diagnosed

with anxiety and depressive disorders at the time of baseline assessment. Indeed, having formal

diagnoses of anxiety and depressive disorders are strong predictors of prospective anxiety and

depression symptoms (**Supplemental Tables 5-12**) and may have obscured the associations

between loss, humiliation and depression. Nevertheless, the fixed effects of total lifetime

entrapment severity were not obscured by including anxiety and depression diagnoses in the

LME models predicting prospective anxiety and depression symptoms, implying that entrapment

severity predicts both anxiety and depression regardless of diagnostic status.

These results are consistent with prior studies implying entrapment as a shared risk

factor for both anxiety and depression (13,19). Exposure to entrapment, which refers to chronic

difficulties that are unlikely to be resolved, has been associated with the onset of depressive (53)

or mixed major depression-generalized anxiety episode (13). Self-reported feelings of

entrapment have been associated with longitudinal changes in anxiety and depression

symptoms at 1-year follow-up in clinically depressed and healthy adolescents (19). Results from

the present study are thus consistent with these findings by showing that higher total lifetime

severity of entrapment exposure was associated with higher levels of anxiety and depression

symptoms at two 6-month follow-up assessments, highlighting entrapment as the most central

shared risk factor for anxiety and depression among all stressor characteristics we examined.

More broadly, they are consistent with a stressor characteristics perspective on stress, which

argues that the effects of stressors on health are not uniform across different of stressors (77).

In turn, results from the unified LME model predicting anxiety symptoms showed that

higher total lifetime severity of entrapment exposure at baseline prospectively predicted even

higher anxiety symptoms at the follow-ups for among participants who had more positive RSFC

within the default network. A meta-analysis found converging evidence that hyperconnectivity

within the default network was associated with anxiety symptoms across multiple anxiety

disorders, including social anxiety, generalized anxiety and panic disorders (78). Findings from

this meta-analysis are consistent with our results, where anxiety symptoms were computed by

summing across subscales related to separation anxiety, social phobia, generalized anxiety and

panic disorder symptoms. A few other studies have found that more positive intra-network

RSFC within the default network in anxiety disorder (79,80). Aspects of the default network,

including the posterior cingulate and the precuneus, are hypothesized to support internally-

related cognition (81–83) such as rumination. Therefore, although speculative, participants

displaying hyperconnectivity within the default network may exhibit higher tendency to engage in

anxiety-related internal rumination and experience higher levels of anxiety symptoms over time

after exposure to stressful life events (84,85).

Finally, results from the unified LME model predicting depression symptoms showed that
total lifetime entrapment severity did not interact with any RSFC metric, suggesting that
entrapment severity and RSFC within and between functional networks independently predicted
prospective depression symptoms. The model also revealed a positive fixed effect of RSFC
between the frontoparietal and the default networks in predicting prospective depression
symptoms. This finding is consistent with prior studies finding hyperconnectivity between the
default and the frontoparietal network in depressed adolescents and adults (45,86–88). Since
anti-correlation between the default and the frontoparietal networks is theorized to reflect
competition between internally-oriented and externally-oriented modes (89), the more positive
RSFC between the default and the frontoparietal networks may reflect difficulty disengaging
from internally-oriented thoughts to meet executive demands (90).
469
Strengths and Limitations
Several strengths and limitations of this work should be noted. In terms of strengths, we
leveraged a longitudinal sample of adolescents enriched for clinical diagnosis of depressive and
anxiety disorders with a narrow age range. This enabled us to examine associations between
exposure to major life stressors, neurobiology, and clinical symptoms during the developmental
stage characterized by heightened vulnerability to stress-related psychopathology (6–10). We
also assessed exposure to a variety of different types of theoretically relevant stressors across
the entire life course using distinct dimensions of social-psychological characteristics. Lastly, we
included all five social-psychological characteristics, all six RSFC metrics within and between the
three *a priori* functional networks, as well as the potential confounding variables into the same
LME model when predicting anxiety/depression symptoms at follow-up. This ensures that any
prospective association between a stressor characteristic or RSFC metric and clinical symptom
is robust to the inclusion of other inter-correlated stressor characteristics or RSFC metrics.
Several limitations should also be noted. First, the sample was relatively small, which
limits the statistical power to detect the complex associations between stressful life event
exposure, RSFC patterns, and clinical symptoms. Although this is a common issue with richly-
phenotyped, longitudinal datasets involving clinical samples, findings from the present study
should be validated by larger samples in future studies. Since the fMRI protocol of the current
dataset was harmonized with other HCP studies (63), other HCP datasets may provide valuable
resources for this purpose. Second, the vast majority (95.33%) of participants were White, and
additional research is needed to examine the generalizability of these findings to other
populations and demographic groups. Third, considering the limited sample size, our

hypothesis-driven analyses focused on RSFC within and between the default, frontoparietal and
ventral attention networks. RSFC involving other functional networks such as the dorsal
attention and the limbic networks as well as the subcortical structures such as the amygdala
have also been associated with anxiety and depression (45,91–94), and future high-powered
analyses would benefit from a whole-brain approach. Fourth, although we included a number of
covariates such as age, race and ethnicity, baseline symptoms, and diagnostic status in our
LME models to ensure that our findings could not be explained by these confounders, other
potential confounders such as medication use and family history of psychopathology were
unavailable in this dataset and may be relevant.
501
Conclusion
Through the use a longitudinal sample of adolescents, over half of whom met clinical
cut-offs for at least one depressive or anxiety disorder, the present analyses demonstrate that
greater total lifetime severity of entrapment exposure predicts higher levels of in anxiety and
depression symptoms at two subsequent 6-month follow-up time points. Lifetime entrapment
exposure prospectively predicted anxiety symptoms in participants with more positive default
network connectivity. Heightened RSFC between the frontoparietal and the default network
specifically predicted higher levels of depression symptoms at the two 6-month follow-ups.
These results imply that among all characteristics of major life stressors that we examined using
the STRAIN, entrapment may be the most important risk factor for both anxiety and depression.
Our results also suggest that more positive connectivity in the default network may be a specific
risk factor for anxiety when an individual is exposed to entrapment, and that more positive
connectivity between the frontoparietal and default network may be a specific risk factor for
depression.
516

Acknowledgments

G.M.S. was supported by grant OPR21101 from the California Governor's Office of Planning and Research/California Initiative to Advance Precision Medicine. S.C. was supported by the McKenzie Fellowship from the University of Melbourne. The study was made possible by National Institute of Mental Health grant R01 MH120080 to A.J.H. The findings and conclusions in this article are those of the authors and do not necessarily represent the views or opinions of these organizations, which had no role in designing or planning this study; in collecting, analyzing, or interpreting the data; in writing the article; or in deciding to submit this article for publication.

Disclosures

The authors declare no conflicts of interest with respect to this work.

References

1. Kessler RC (1997): The effects of stressful life events on depression. *Annu Rev Psychol* 48: 191–214.
2. Miloyan B, Joseph Bienvenu O, Brilot B, Eaton WW (2018): Adverse life events and the onset of anxiety disorders. *Psychiatry Res* 259: 488.
3. Kendler KS, Karkowski LM, Prescott CA (1999): Causal relationship between stressful life events and the onset of major depression. *American Journal of Psychiatry* 156: 837–841.
4. Francis JL, Moitra E, Dyck I, Keller MB (2012): The impact of stressful life events on relapse of generalized anxiety disorder. *Depress Anxiety* 29: 386–391.
5. Faravelli C, Pallanti S (1989): Recent life events and panic disorder. *Am J Psychiatry* 146: 622–626.
6. Paus T, Keshavan M, Giedd JN (2008): Why do many psychiatric disorders emerge during adolescence? *Nature Reviews Neuroscience 2008 9:12* 9: 947–957.
7. Merikangas KR, He JP, Burstein M, Swanson SA, Avenevoli S, Cui L, *et al.* (2010): Lifetime prevalence of mental disorders in U.S. adolescents: Results from the national comorbidity survey replication-adolescent supplement (NCS-A). *J Am Acad Child Adolesc Psychiatry* 49: 980–989.
8. Casey BJ, Getz S, Galvan A (2008): The adolescent brain. *Developmental Review* 28: 62–77.
9. March-Llanes J, Marqués-Feixa L, Mezquita L, Fañanás L, Moya-Higueras J (2017): Stressful life events during adolescence and risk for externalizing and internalizing psychopathology: a meta-analysis. *Eur Child Adolesc Psychiatry* 26: 1409–1422.
10. Larsen B, Luna B (2018): Adolescence as a neurobiological critical period for the development of higher-order cognition. *Neurosci Biobehav Rev* 94: 179–195.
11. Slavich GM (2019): Stressnology: The primitive (and problematic) study of life stress exposure and pressing need for better measurement. *Brain Behav Immun* 75: 3–5.
12. Cohen S, Murphy MLM, Prather AA (2019): Ten Surprising Facts About Stressful Life Events and Disease Risk. *Annu Rev Psychol* 70: 577.
13. Kendler KS, Hettema JM, Butera F, Gardner CO, Prescott CA (2003): Life Event Dimensions of Loss, Humiliation, Entrapment, and Danger in the Prediction of Onsets of Major Depression and Generalized Anxiety. *Arch Gen Psychiatry* 60: 789–796.
14. Farmer AE, McGuffin P (2003): Humiliation, loss and other types of life events and difficulties: a comparison of depressed subjects, healthy controls and their siblings. *Psychol Med* 33: 1169–1175.
15. Asselmann E, Wittchen HU, Lieb R, Höfler M, Beesdo-Baum K (2015): Danger and loss events and the incidence of anxiety and depressive disorders: a prospective-longitudinal community study of adolescents and young adults. *Psychol Med* 45: 153–163.
16. Finlay-Jones R, Brown GW (1981): types of stressful life event and the onset of anxiety and depressive disorders. *Psychol Med* 11: 803–815.
17. Keller MC, Neale MC, Kendler KS (2007): Association of different adverse life events with distinct patterns of depressive symptoms. *Am J Psychiatry* 164: 1521–1529.
18. Ayazi T, Lien L, Eide A, Swartz L, Hauff E (2014): Association between exposure to traumatic events and anxiety disorders in a post-conflict setting: a cross-sectional community study in South Sudan. *BMC Psychiatry* 14: 6.
19. Griffiths AW, Wood AM, Maltby J, Taylor PJ, Tai S (2014): The prospective role of defeat and entrapment in depression and anxiety: A 12-month longitudinal study. *Psychiatry Res* 216: 52–59.
20. Beck A (1976): *Cognitive Therapy and the Emotional Disorders*. New York, NY: International Universities Press.
21. Beck R, Perkins TS (2001): Cognitive content-specificity for anxiety and depression: A meta-analysis. *Cognit Ther Res* 25: 651–663.

22. Cho Y, Telch MJ (2005): Testing the cognitive content-specificity hypothesis of social anxiety and depression: An application of structural equation modeling. *Cognit Ther Res* 29: 399–416.
23. Bremner JD (2006): Traumatic stress: effects on the brain. *Dialogues Clin Neurosci* 8: 445.
24. Brieant AE, Sisk LM, Gee DG (2021): Associations among negative life events, changes in cortico-limbic connectivity, and psychopathology in the ABCD Study. *Dev Cogn Neurosci* 52: 101022.
25. Casement MD, Shaw DS, Sitnick SL, Musselman SC, Forbes EE (2015): Life stress in adolescence predicts early adult reward-related brain function and alcohol dependence. *Soc Cogn Affect Neurosci* 10: 416–423.
26. Hosseini-Kamkar N, Varvani Farahani M, Nikolic M, Stewart K, Goldsmith S, Soltaninejad M, *et al.* (2023): Adverse Life Experiences and Brain Function: A Meta-Analysis of Functional Magnetic Resonance Imaging Findings. *JAMA Netw Open* 6: e2340018–e2340018.
27. Stanton CH, Holmes AJ, Chang SWC, Joormann J (2019): From Stress to Anhedonia: Molecular Processes through Functional Circuits. *Trends Neurosci* 42: 23–42.
28. Fadel E, Boeker H, Gaertner M, Richter A, Kleim B, Seifritz E, *et al.* (2021): Differential alterations in resting state functional connectivity associated with depressive symptoms and early life adversity. *Brain Sci* 11: 591.
29. Grant MM, White D, Hadley J, Hutcheson N, Shelton R, Sreenivasan K, Deshpande G (2014): Early life trauma and directional brain connectivity within major depression. *Hum Brain Mapp* 35: 4815–4826.
30. Luo Q, Zou Y, Nie H, Wu H, Du Y, Chen J, *et al.* (2023): Effects of childhood neglect on regional brain activity and corresponding functional connectivity in major depressive disorder and healthy people: Risk factor or resilience? *J Affect Disord* 340: 792–801.
31. Gollier-Briant F, Marie-Laure Paillère-Martinot, Lemaitre H, Miranda R, Vulser H, Goodman R, *et al.* (2016): Neural correlates of three types of negative life events during angry face processing in adolescents. *Soc Cogn Affect Neurosci* 11: 1961–1969.
32. Sokołowski A, Folkierska-Żukowska M, Jednoróg K, Moodie CA, Dragan W (2020): The relationship between early and recent life stress and emotional expression processing: A functional connectivity study. *Cognitive, Affective, & Behavioral Neuroscience 2020 20:3* 20: 588–603.
33. Weinberg A, Kujawa A, Riesel A (2022): Understanding Trajectories to Anxiety and Depression: Neural Responses to Errors and Rewards as Indices of Susceptibility to Stressful Life Events. *https://doi.org/101177/09637214211049228* 31: 115–123.
34. Goldstein BL, Kessel EM, Kujawa A, Finsaas MC, Davila J, Hajcak G, Klein DN (2020): Stressful life events moderate the effect of neural reward responsiveness in childhood on depressive symptoms in adolescence. *Psychol Med* 50: 1548.
35. Vincent JL, Kahn I, Snyder AZ, Raichle ME, Buckner RL (2008): Evidence for a frontoparietal control system revealed by intrinsic functional connectivity. *J Neurophysiol* 100: 3328–3342.
36. Raichle ME, MacLeod AM, Snyder AZ, Powers WJ, Gusnard DA, Shulman GL (2001): A default mode of brain function. *Proc Natl Acad Sci U S A* 98: 676–682.
37. Buckner RL, Andrews-Hanna JR, Schacter DL (2008): The Brain's Default Network. *Ann N Y Acad Sci* 1124: 1–38.
38. Seeley WW, Menon V, Schatzberg AF, Keller J, Glover GH, Kenna H, *et al.* (2007): Dissociable Intrinsic Connectivity Networks for Salience Processing and Executive Control. *Journal of Neuroscience* 27: 2349–2356.
39. Sheynin J, Duval ER, Lokshina Y, Scott JC, Angstadt M, Kessler D, *et al.* (2020): Altered resting-state functional connectivity in adolescents is associated with PTSD symptoms and trauma exposure. *Neuroimage Clin* 26: 102215.

40. Long J, Huang X, Liao Y, Hu X, Hu J, Lui S, *et al.* (2014): Prediction of post-earthquake depressive and anxiety symptoms: a longitudinal resting-state fMRI study. *Scientific Reports 2014 4:1* 4: 1–10.
41. Philip NS, Sweet LH, Tyrka AR, Price LH, Bloom RF, Carpenter LL (2013): Decreased default network connectivity is associated with early life stress in medication-free healthy adults. *European Neuropsychopharmacology* 23: 24–32.
42. Harnett NG, van Rooij SJH, Ely TD, Lebois LAM, Murty VP, Jovanovic T, *et al.* (2021): Prognostic neuroimaging biomarkers of trauma-related psychopathology: resting-state fMRI shortly after trauma predicts future PTSD and depression symptoms in the AURORA study. *Neuropsychopharmacology 2021 46:7* 46: 1263–1271.
43. Cole MW, Repovš G, Anticevic A (2014, December 20): The frontoparietal control system: A central role in mental health. *Neuroscientist*, vol. 20. SAGE Publications Inc., pp 652–664.
44. Menon V (2011): Large-scale brain networks and psychopathology: a unifying triple network model. *Trends Cogn Sci* 15: 483–506.
45. Kaiser RH, Andrews-Hanna JR, Wager TD, Pizzagalli DA (2015): Large-Scale Network Dysfunction in Major Depressive Disorder: A Meta-analysis of Resting-State Functional Connectivity. *JAMA Psychiatry* 72: 603–611.
46. Berman MG, Peltier S, Nee DE, Kross E, Deldin PJ, Jonides J (2011): Depression, rumination and the default network. *Soc Cogn Affect Neurosci* 6: 548–555.
47. Manoliu A, Meng C, Brandl F, Doll A, Tahmasian M, Scherr M, *et al.* (2014): Insular dysfunction within the salience network is associated with severity of symptoms and aberrant inter-network connectivity in major depressive disorder. *Front Hum Neurosci* 7. https://doi.org/10.3389/fnhum.2013.00930
48. Gee DG, Casey BJ (2015): The impact of developmental timing for stress and recovery. *Neurobiol Stress* 1: 184–194.
49. Tottenham N, Galván A (2016): Stress and the adolescent brain: Amygdala-prefrontal cortex circuitry and ventral striatum as developmental targets. *Neurosci Biobehav Rev* 70: 217–227.
50. Hubbard NA, Bauer CCC, Siless V, Auerbach RP, Elam JS, Frosch IR, *et al.* (2024): The Human Connectome Project of adolescent anxiety and depression dataset. *Scientific Data 2024 11:1* 11: 1–15.
51. American Psychiatric Association (2013): *Diagnostic and Statistical Manual of Mental Disorders*, 5th ed. Retrieved October 12, 2024, from https://psychiatryonline.org/doi/book/10.1176/appi.books.9780890425596
52. Slavich GM, Stewart JG, Esposito EC, Shields GS, Auerbach RP (2019): The Stress and Adversity Inventory for Adolescents (Adolescent STRAIN): associations with mental and physical health, risky behaviors, and psychiatric diagnoses in youth seeking treatment. *J Child Psychol Psychiatry* 60: 998–1009.
53. Harris TO, Hepworth C, Brown GW (1995): Loss, humiliation and entrapment among women developing depression: a patient and non-patient comparison. *Psychol Med* 25: 7–21.
54. Slotter EB, Walsh CM (2017): All role transitions are not experienced equally: Associations among self-change, emotional reactions, and self-concept clarity. *Self and Identity* 16: 531–556.
55. Slavich GM, Shields GS (2018): Assessing Lifetime Stress Exposure Using the Stress and Adversity Inventory for Adults (Adult STRAIN): An Overview and Initial Validation. *Psychosom Med* 80: 17–27.
56. Chorpita BF, Yim L, Moffitt C, Umemoto LA, Francis SE (2000): Assessment of symptoms of DSM-IV anxiety and depression in children: a revised child anxiety and depression scale. *Behaviour Research and Therapy* 38: 835–855.
57. COSTELLO EJ, ANGOLD A (1988): Scales to Assess Child and Adolescent Depression: Checklists, Screens, and Nets. *J Am Acad Child Adolesc Psychiatry* 27: 726–737.

58. Angold A, Costello EJ, Messer SC, Pickles A (1995): Development of a short questionnaire for use in epidemiological studies of depression in children and adolescents. *Int J Methods Psychiatr Res* 5: 237–249.

59. Wood A, Kroll L, Moore A (1995): Properties of the mood and feelings questionnaire in adolescent psychiatric outpatients: a research note. *J Child Psychol Psychiatry* 36: 327–334.

60. Sund AM, Larsson B, Wichstrøm L (2001): Depressive symptoms among young Norwegian adolescents as measured by the Mood and Feelings Questionnaire (MFQ). *Eur Child Adolesc Psychiatry* 10: 222–229.

61. Burleson Daviss W, Birmaher B, Melhem NA, Axelson DA, Michaels SM, Brent DA (2006): Criterion validity of the Mood and Feelings Questionnaire for depressive episodes in clinic and non-clinic subjects. *J Child Psychol Psychiatry* 47: 927–934.

62. Thabrew H, Stasiak K, Bavin LM, Frampton C, Merry S (2018): Validation of the Mood and Feelings Questionnaire (MFQ) and Short Mood and Feelings Questionnaire (SMFQ) in New Zealand help‑seeking adolescents. *Int J Methods Psychiatr Res* 27. https://doi.org/10.1002/MPR.1610

63. Siless V, Hubbard NA, Jones R, Wang J, Lo N, Bauer CCC, *et al.* (2020): Image acquisition and quality assurance in the Boston Adolescent Neuroimaging of Depression and Anxiety study. *Neuroimage Clin* 26: 102242.

64. Glasser MF, Sotiropoulos SN, Wilson JA, Coalson TS, Fischl B, Andersson JL, *et al.* (2013): The minimal preprocessing pipelines for the Human Connectome Project. *Neuroimage* 80: 105–124.

65. Van Essen DC, Glasser MF, Dierker DL, Harwell J, Coalson T (2012): Parcellations and hemispheric asymmetries of human cerebral cortex analyzed on surface-based atlases. *Cereb Cortex* 22: 2241–2262.

66. Jenkinson M, Smith S (2001): A global optimisation method for robust affine registration of brain images. *Med Image Anal* 5: 143–156.

67. Jenkinson M, Bannister P, Brady M, Smith S (2002): Improved Optimization for the Robust and Accurate Linear Registration and Motion Correction of Brain Images. *Neuroimage* 17: 825–841.

68. Greve DN, Fischl B (2009): Accurate and Robust Brain Image Alignment using Boundary-based Registration. *Neuroimage* 48: 63.

69. Griffanti L, Salimi-Khorshidi G, Beckmann CF, Auerbach EJ, Douaud G, Sexton CE, *et al.* (2014): ICA-based artefact removal and accelerated fMRI acquisition for improved resting state network imaging. *Neuroimage* 95: 232–247.

70. Salimi-Khorshidi G, Douaud G, Beckmann CF, Glasser MF, Griffanti L, Smith SM (2014): Automatic denoising of functional MRI data: combining independent component analysis and hierarchical fusion of classifiers. *Neuroimage* 90: 449–468.

71. Robinson EC, Jbabdi S, Glasser MF, Andersson J, Burgess GC, Harms MP, *et al.* (2014): MSM: A new flexible framework for Multimodal Surface Matching. *Neuroimage* 100: 414–426.

72. Glasser MF, Smith SM, Marcus DS, Andersson JLR, Auerbach EJ, Behrens TEJ, *et al.* (2016): The Human Connectome Project's neuroimaging approach. *Nature Neuroscience 2016 19:9* 19: 1175–1187.

73. Schaefer A, Kong R, Gordon EM, Laumann TO, Zuo X-N, Holmes AJ, *et al.* (2018): Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cereb Cortex* 28: 3095–3114.

74. Yeo BTT, Krienen FM, Sepulcre J, Sabuncu MR, Lashkari D, Hollinshead M, *et al.* (2011): The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *J Neurophysiol* 106: 1125–1165.

75. Bates D, Mächler M, Bolker BM, Walker SC (2015): Fitting Linear Mixed-Effects Models Using lme4. *J Stat Softw* 67: 1–48.
76. Säfken B, Rügamer D, Kneib T, Greven S (2021): Conditional Model Selection in Mixed-Effects Models with cAIC4. *J Stat Softw* 99: 1–30.
77. Lüdecke D, Ben-Shachar MS, Patil I, Waggoner P, Makowski D (2021): performance: An R Package for Assessment, Comparison and Testing of Statistical Models. *J Open Source Softw* 6: 3139.
78. Pierce ZP, Black JM (2023): Default mode intranetwork resting-state functional connectivity is correlated with increased symptom severity in common anxiety disorders: A systematic review and meta-analysis. *J Affect Disord Rep* 14: 100674.
79. Rabany L, DIefenbach GJ, Bragdon LB, Pittman BP, Zertuche L, Tolin DF, *et al.* (2017): Resting-State Functional Connectivity in Generalized Anxiety Disorder and Social Anxiety Disorder: Evidence for a Dimensional Approach. *Brain Connect* 7: 289–298.
80. Li R, Shen F, Sun X, Zou T, Li L, Wang X, *et al.* (2023): Dissociable salience and default mode network modulation in generalized anxiety disorder: a connectome-wide association study. *Cerebral Cortex* 33: 6354–6365.
81. Leech R, Sharp DJ (2014): The role of the posterior cingulate cortex in cognition and disease. *Brain* 137: 12–32.
82. Nejad AB, Fossati P, Lemogne C (2013, October 10): Self-referential processing, rumination, and cortical midline structures in major depression. *Frontiers in Human Neuroscience*, vol. 7. Frontiers Media S. A. https://doi.org/10.3389/fnhum.2013.00666
83. Gusnard DA, Akbudak E, Shulman GL, Raichle ME (2001): Medial prefrontal cortex and self-referential mental activity: Relation to a default mode of brain function. *Proc Natl Acad Sci U S A* 98: 4259–4264.
84. Michl LC, McLaughlin KA, Shepherd K, Nolen-Hoeksema S (2013): Rumination as a mechanism linking stressful life events to symptoms of depression and anxiety: Longitudinal evidence in early adolescents and adults. *J Abnorm Psychol* 122: 339–352.
85. Ruscio AM, Gentes EL, Jones JD, Hallion LS, Coleman ES, Swendsen J (2015): Rumination predicts heightened responding to stressful life events in major depressive disorder and generalized anxiety disorder. *J Abnorm Psychol* 124: 17–26.
86. Ma C, Ding J, Li J, Guo W, Long Z, Liu F, *et al.* (2012): Resting-State Functional Connectivity Bias of Middle Temporal Gyrus and Caudate with Altered Gray Matter Volume in Major Depression. *PLoS One* 7: 45263.
87. Davey CG, Harrison BJ, Yücel M, Allen NB (2012): Regionally specific alterations in functional connectivity of the anterior cingulate cortex in major depressive disorder. *Psychol Med* 42: 2071–2081.
88. Connolly CG, Ho TC, Blom EH, LeWinn KZ, Sacchet MD, Tymofiyeva O, *et al.* (2017): Resting-state functional connectivity of the amygdala and longitudinal changes in depression severity in adolescent depression. *J Affect Disord* 207: 86–94.
89. Fransson P (2005): Spontaneous low-frequency BOLD signal fluctuations: An fMRI investigation of the resting-state default mode of brain function hypothesis. *Hum Brain Mapp* 26: 15.
90. Clare Kelly AM, Uddin LQ, Biswal BB, Castellanos FX, Milham MP (2008): Competition between functional brain networks mediates behavioral variability. *Neuroimage* 39: 527–537.
91. Cullen KR, Westlund MK, Klimes-Dougan B, Mueller BA, Houri A, Eberly LE, Lim KO (2014): Abnormal Amygdala Resting-State Functional Connectivity in Adolescent Depression. *JAMA Psychiatry* 71: 1138–1147.
92. Liu R, Wang Y, Chen X, Zhang Z, Xiao L, Zhou Y (2021): Anhedonia correlates with functional connectivity of the nucleus accumbens subregions in patients with major depressive disorder. *Neuroimage Clin* 30: 102599.

93. Sylvester CM, Corbetta M, Raichle ME, Rodebaugh TL, Schlaggar BL, Sheline YI, *et al.* (2012): Functional network dysfunction in anxiety and anxiety disorders. *Trends Neurosci* 35: 527–535.

94. Xu J, Van Dam NT, Feng C, Luo Y, Ai H, Gu R, Xu P (2019): Anxious brain networks: A coordinate-based activation likelihood estimation meta-analysis of resting-state functional connectivity studies in anxiety. *Neurosci Biobehav Rev* 96: 21–30.