FRUCTOSE ACTIVATES A STRESS RESPONSE SHARED BY METHYLGLYOXAL AND HYDROGEN PEROXIDE IN *STREPTOCOCCUS MUTANS*

Alejandro R. Walker, Danniel N. Pham, Payam Noeparvar, Alexandra M. Peterson, Marissa K. Lipp, José A. Lemos, and Lin Zeng[‡]

Department of Oral Biology, University of Florida College of Dentistry, Gainesville, Florida, USA.

Key words: Fructose, Methylglyoxal, Hydrogen Peroxide, F-1-P, Stress Response, Metal homeostasis, Gene Regulation.

[‡] Corresponding author:

Lin Zeng, PhD

1395 Center Drive

Gainesville, FL 32610

Email: lzeng@dental.ufl.edu

Phone: (352) 273-8868

Abstract

Fructose catabolism by *Streptococcus mutans* is initiated by three PTS transporters yielding fructose-1-phoshate (F-1-P) or fructose-6-phosphate. Deletion of one such F-1-P-generating PTS, *fruI*, was shown to reduce the cariogenicity of *S. mutans* in rats fed a high-sucrose diet. Moreover, a recent study linked fructose metabolism in *S. mutans* to a reactive electrophile species (RES) methylglyoxal. Here, we conducted a comparative transcriptomic analysis of *S. mutans* treated briefly with 50 mM fructose, 50 mM glucose, 5 mM methylglyoxal, or 0.5 mM hydrogen peroxide (H_2O_2). The results revealed a striking overlap between the fructose and methylglyoxal transcriptomes, totaling 176 genes, 61 of which were also shared with the H_2O_2 transcriptome. This core of 61 genes encompassed many of the same pathways affected by exposure to low pH or zinc intoxication. Consistent with these findings, fructose negatively impacted metal homeostasis of a mutant deficient in zinc expulsion and the growth of a mutant of the major oxidative stress regulator SpxA1. Importantly, fructose metabolism lowered culture pH at a faster pace, allowed better survival under acidic and nutrient-depleted conditions, and enhanced the competitiveness of *S. mutans* against *Streptococcus sanguinis*, although a moderated level of F-1-P might further boost some of these benefits. Conversely, several commensal streptococcal species displayed a greater sensitivity to fructose that may negatively affect their persistence and competitiveness in dental biofilm. In conclusion, fructose metabolism is integrated into the stress core of *S. mutans* and regulates critical functions required for survival and its ability to induce dysbiosis in the oral cavity.

Importance. Fructose is a common monosaccharide in the biosphere, yet its

overconsumption has been linked to various health problems in humans including

insulin resistance, obesity, diabetes, non-alcoholic liver diseases, and even cancer.

These effects are in large part attributable to the unique biochemical characteristics and

metabolic responses associated with the degradation of fructose. Yet, an understanding

of the effects of fructose on the physiology of bacteria and its implications to the human

microbiome is severely lacking. Here we performed a series of analyses on the gene

regulation of a dental pathogen *Streptococcus mutans* by exposing it to fructose and

other important stress agents. Further supported by growth, persistence, and

competition assays, our findings revealed the ability of fructose to activate a set of

stress-related functions that may prove critical to the ability of the bacterium to persist

and cause diseases both within and without of the oral cavity.

Introduction

Disruption in microbial homeostasis in human microbiomes, a condition termed

dysbiosis, has been associated with the development of many diseases that range from

GI tract disorders such as inflammatory bowel disease and pseudomembranous colitis

to oral infectious diseases such as dental caries, candidiasis, and periodontitis (1, 2).

Carbohydrate metabolism plays a vital role in oral health by contributing to the

biochemistry and ecology of oral microbiota. It is widely understood that

overconsumption of refined carbohydrates is a defining factor in the development of

dysbiosis through acid production and establishment of an acidogenic and aciduric

dental biofilm (3-6). Emerging evidence also supports the co-occurrence of dental caries

and periodontal diseases, pointing to potentially shared mechanisms in underlying

etiology (7-9). With the food industry shifting heavily toward the use of fructose and

fructose-releasing sugars in the past several decades and the concurrent pandemic of

metabolic syndrome, the impact of dietary fructose to oral and systemic health demands

urgent attention (10).

Fructose is a ubiquitous nutrient widely found in fruits, yet overconsumption of

fructose is often associated with adverse health outcomes in humans that include

insulin resistance, obesity, liver diseases, systemic inflammation, type-2 diabetes,

cogitative decline, and even cancer (11-15). According to the "fructose survival

hypothesis", fructose triggers a conserved cellular response in mammalian cells that

manifests as mitochondrial oxidative stress and reduced cellular ATP levels,

accompanied by behavioral changes that includes hunger, thirst, foraging, and weight

gain (16). While this response could have been limited and beneficial to animals and our

human ancestors, the negative effects of fructose have been greatly exacerbated by its

widespread use in Western diet.

Streptococci are the most abundant commensal species colonizing oral surfaces, and

most of them rely on carbohydrate fermentation, via the Embden–Meyerhof–Parnas

(EMP) pathway of glycolysis, to produce energy and precursors for various biogenesis

processes (3). Multiple fructose-transporting mechanisms exist in streptococci, each

harboring a fructose: phosphotransferase (PTS) permease that generates either

fructose-1-phosphate (F-1-P) or fructose-6-phosphate (F-6-P), which is further

phosphorylated into F-1,6-bP by their respective phosphofructokinases (17). Studies so

far have indicated that F-1-P metabolism contributes to *in vitro* fitness, biofilm

development, and virulence for pathogenic species such as *Streptococcus pyogenes,*

Streptococcus agalactiae, and *Streptococcus gordonii* (18-22). For the cariogenic

pathobiont *Streptococcus mutans*, the primary fructose pathway is encoded by the

fruRKI operon, encoding a negative regulator FruR, a PTS permease (FruI) that yields

F-1-P, and a F-1-P kinase (FruK) (17). Deletion of *fruI* in *S. mutans* UA159 resulted in

lower levels, though not complete lack of F-1-P when exposed to fructose (23) and

reduced the ability of the bacterium to colonize and induce dentin caries in rats fed a

high-sucrose diet (24). Based on genetic studies, expression of *fruRKI,* and its

homologue *fruRBA* in related streptococci, is inducible by F-1-P which acts allosterically

on FruR (17, 18, 25). Another fructose-PTS (EII^{Lev}), encoded by *levDEFG,* exists in

some of these oral streptococci, including *S. mutans, Streptococcus sanguinis* and *S.*

gordonii, which likely generates F-6-P and requires a four-component system LevQRST

for induction in response to extracellular fructose (26). *S. mutans* harbors yet another

fructose-PTS (*fruCD*) that is also predicted to generate F-1-P, although its contribution

to the overall fructose transport appears to be minor (17). Overall, the F-1-P pathway

encoded by *fruRBA/fruRKI* operon is conserved in most Gram-positive bacteria

encompassing *Bacillus, Enterococcus, Lactococcus, Lactobacillus, Clostridium, Listeria,*

and most *Streptococcus* species (27-29). Moreover, *S. mutans, S. gordonii,* and *S.*

sanguinis also harbor another fructose-related operon, *sppRA.* SppA has been

characterized as a hexose-phosphate phosphohydrolase that specializes in

dephosphorylating F-1-P, and F-6-P to a lesser extent; and SppR is a negative auto-

regulator whose activity is allosterically regulated by F-1-P (23).

A recent study involving fructose centered on the molecular mechanisms required for

oral streptococci to degrade a group of highly reactive electrophile species (RES),

including methylglyoxal and glyoxal (30), which are created by all life as byproducts of

glycolysis and serve as important immune effectors (31, 32). It has been reported that

S. mutans is significantly more resistant to methylglyoxal than several commensal

streptococci including *S. gordonii, S. sanguinis,* and members of the mitis group (30). A

zinc-containing protein GloA2 (SMU.1112c), predicted to be a paralogue to the main

methylglyoxal-degrading enzyme glyoxalase A (LguL or GloA) based on its sequence

and crystal structure, was identified in all oral streptococci examined so far (30). The

gloA2-null mutants of both *S. mutans* and *S. sanguinis* displayed increased chaining

and autolysis upon treatment by fructose which resembled the effect of exposure to

methylglyoxal, including its reversal by treatment with glutathione (GSH) which is known

to scavenge methylglyoxal before degradation by LguL (32). Interestingly, the

expression of *lguL* was inducible by growth on fructose, and the fructose-sensitivity of

gloA2 mutants was largely dependent on the integrity of the F-1-P-generating PTS, FruI

(30). Herein we further examined fructose-dependent gene regulation and physiology in

relation to streptococcal tolerance to important stressors including methylglyoxal and

hydrogen peroxide. Our findings revealed the integration of fructose-mediated response

with general stress response experienced during bacteria-host and inter-microbial

interactions.

Results

Comparative RNA-seq analyses identified a stress core shared by methylglyoxal,

fructose, and H_2O_2.

We previously performed an RNA-seq analysis on *S. mutans* cultured to the

exponential phase with 0.5% fructose (27.7 mM) in the medium (33). The strain used in

this study was a mutant deficient in several enzymes capable of degrading sucrose and

other carbohydrate polymers (34), and was later revealed to contain a spontaneous

mutation in a major peroxide-sensing regulator PerR (35). While this work provided us

important information regarding the state of the bacterium after growing on fructose,

these genetic deviations are of concern and the study could not capture the response

the cells present immediately after fructose intake, as metabolic regulation tends to

return to equilibrium after initial perturbance. To mimic the physiological impact to

bacteria when humans consume a large dose of fructose in food or drinks, we cultured

S. mutans strain UA159 (from ATCC) in a glucose-based medium to the exponential

phase and then treated the cells with 50 mM fructose for 30 min, which were

immediately processed for RNA-seq analysis in comparison to untreated controls.

Considering the similarity between glucose and fructose, the same UA159 cultures were

treated with 50 mM glucose in the same manner and analyzed as an additional control.

These concentrations of carbohydrates represent the physiological levels the microbiota

likely encounter periodically in oral cavity. The experiment was also repeated using the

same bacteria after treatment with 5 mM methylglyoxal for 30 min (30). In addition, we

obtained and reanalyzed the datasets of a similar RNA-seq study performed on UA159

cultures that were prepared in a glucose-based medium and treated with 0.5 mM H_2O_2

for 5 min (36) (see Fig. 1 for volcano plots, Fig. 2 for Venn diagrams, Fig. S1 for

principal component analyses, Fig. S2 for Gene Ontology term analyses, and

supplemental Tables S1 to S3 for other details).

Methylglyoxal. Treatment of UA159 with 5 mM of methylglyoxal differentially affected

the expression of a total of 474 genes when compared to the mock-treated control

group, with 271 being up-regulated and 203 being down-regulated (Fig. 2, Fig. S2).

Methylglyoxal is known to react with the reduced form of glutathione (GSH), which is

interconnected with other reductive potentials such as NADPH and ascorbate (37).

Treatment with methylglyoxal resulted primarily in several effects similar to oxidative

stress response (36): increased expression of genes involved in thiol homeostasis

including thioredoxins *trxAB*, GSH synthetase *gshAB* (38)*,* glutathione reductase *gor,*

and glutathione S-transferase *gst*; increased expression of the iron-sulfur (Fe-S) cluster

biosynthesis system *sufCDSUB* (39), the ascorbate metabolic cluster (SMU.271 to

SMU.275) (40), the reactive oxygen species (ROS) scavenging genes *sodA*, *ahpCF*,

and *tpx*, and genes whose products are involved in metal homeostasis: copper exporter

operon *copYAZ*, and iron-sequestering gene *dpr*. Also upregulated were several

transcription regulators known to be involved in stress management: *relS*, *perR*,

phoR/ycbL, *scnKR*, *rcrRPQ*, SMU.1097c, *vicKX*, *comR*, *comDE*, and *lexA*. Likely as the

need to restore the GSH/GSSG balance, treatment of methylglyoxal upregulated

several putative aldo/keto reductases: SMU.382c and SMU.383c, SMU.677 to

SMU.681, SMU.1040c, and SMU.1602 residing immediately upstream of *lguL* which

encodes the major methylglyoxal-degrading enzyme LguL/GloA (both up by 3~4-fold). It

is worth noting that most of these aldo/keto reductases, other than *lguL*, are absent in

related commensal streptococci. Also related to improved reducing capacity was

upregulation of several putative ribonucleotide reductases (41, 42): SMU.667 to

SMU.669c, SMU.991, and SMU.2070 to SMU.2074 (*nrdG*).

Interestingly, methylglyoxal treatment reduced the expression of iron/manganese

transporter *sloAB* (not *sloC*) and cognate *sloR* regulator. It also reduced the expression

of ferrous ion transporter genes *feoAB*. These changes were consistent with the

purported ability of GSH to buffer intracellular metal ions (43), as a release of these ions

would likely increase the expression of metal exporters (*copYA*) but reduce that of metal

importers (*sloABC* and *feoAB*). Another likely effect of methylglyoxal exposure is its

reaction with intracellular amino acids such as arginine and lysine. For this, we

observed increased expression of amino acid biosynthesis/transport operons, *argCJBD*

for arginine and *livFGMHK* for branched-chain amino acids. As a general response to

stressful stimuli, we observed elevated expression of genes required for DNA repair

(*exoA*, *end3*, *mutY*, and *uvrA*) and protein or peptide degradation (*clpC, clpE, pepO*,

and *pepP*). Conversely, there were also several stress management genes that showed

reduced expression: *mreCD*, *irvR*, *hrcA, grpE,* and *dnaK*. Most ribosomal proteins and

the translation initiation factor InfC also had lower mRNA levels, indicative of reduced

protein synthesis activity. There was likely a shift toward a more efficient, mixed-acid

fermentation as we saw greater expression by acetoin metabolic operon (*adhABCD-*

lplA), the related alpha-acetolactate metabolic genes *alsS* and *aldB*, NADH oxidase

(*nox*), and a pyruvate-formate lyase activating enzyme *pflA*. At the same time, the

genes encoding the two major enzymes of the PTS, *ptsH* and *ptsI*, along with the

transport and metabolic pathways for mannitol (EII^{Mtl}, SMU.1184c and SMU.1185),

cellobiose (EII^{Cel}, SMU.1596 to SMU.1600), sucrose (EII^{Scr}, SMU.1841 to SMU.1844),

and glucose (EII^{Man}, SMU.1877 to SMU.1879) all showed significantly reduced

expression.

Comparative analysis of methylglyoxal regulon with H_2O_2-mediated stress response.

During the analysis of the methylglyoxal-induced transcriptomic shift, it was immediately

apparent that many of these genes were also identified by a previous study on peroxide

stress response of *S. mutans* UA159 (36). A closer inspection of the peroxide stress

transcriptome identified a total of 115 genes (Supplemental Tables S1 and S4, Fig. 2A)

as being shared by methylglyoxal and H_2O_2, indicative of the common bacterial

mechanisms needed to deal with these two stressors. These aforementioned functions

included ROS scavenging, thiol and metal homeostasis, DNA repair and protein

processing, and energy metabolism. Out of these 115 genes, 100 showed upregulation

and 12 were downregulated; only 3 showed different polarity of change. Important

distinctions were also noted which encompassed 359 methylglyoxal-specific genes that

were missing in H_2O_2 regulon, and conversely 79 H_2O_2-responsive genes absent in the

methylglyoxal regulon (Supplemental Table S4). Briefly, absent in H_2O_2-induced

response included the reduced expression of the PTS genes, namely *ptsH, ptsI*, and

EII^{Mtl}, EII^{Cel}, EII^{Scr}, and EII^{Man} operons. Also absent were genes for ribosomal proteins,

lrgAB-lytST, *argCJBD* and agmatine deiminase operon (44)*,* several putative aldo/keto

reductase clusters, potassium transporters *trkAB*, the *nrd* operon SMU.2070 to

SMU.2074, *mreCD*, *hrcA-grpE-dnaK*, and transcription regulators including *mleR, relS*,

phoR/ycbL, *nagR*, *irvR*, *vicKX*, *comX,* and *ahrC.* Missing from the methylglyoxal regulon

were some of the H_2O_2-induced genes, including mutacin genes (*nlmABCD* and

nlmTE), *sloC*, *liaSR*, *levQRST*, *relA,* and the zinc import operon *adcBCR.* Interestingly,

while methylglyoxal induced expression of most of the competence regulators,

comCDE, comR, and *rcrR* (45)*,* and the late-competence operon *comYA-YI* (46), H_2O_2

reduced expression of two important *nlm* operons, one of which (*nlmABCD*) requires an

active *comDE* circuit for expression (47).

Significant overlap between fructose, methylglyoxal, and H_2O_2-mediated responses.

Treatment of UA159 with 50 mM fructose for 30 min differentially affected the

expression of a total of 449 genes, with 250 being up-regulated and 199 being down-

regulated (Fig. 2B, Fig. S2, and Table S3). Though not a critical point of this study,

these findings represent a striking difference in gene regulation compared to what was

previously observed in UA159 grown with 0.5% fructose (33). As a control, similar

treatment with 50 mM glucose for 30 min also upregulated 128 genes and down-

regulated 216 genes (Table S3). Significantly, there was a larger overlap between the

genes affected by fructose and methylglyoxal, totaling 176 genes (98 up and 62 down,

16 with opposite directions of change), compared to the 116-gene overlap between

glucose and methylglyoxal (38 up and 48 down, 30 with opposite directions of change).

Among the 176 genes shared by methylglyoxal and fructose, 61 overlapped the H_2O_2

regulon; among the 116 genes shared by methylglyoxal and glucose, only 24

overlapped with the H_2O_2 regulon (Table S5). Therefore, compared to glucose,

significantly more genes were shared by treatment of fructose with that of methylglyoxal

and H_2O_2. Comparative analysis identified a set of 44 genes that were similarly affected

by fructose, methylglyoxal, and H_2O_2, excluding 17 genes impacted by all four

treatments. Altogether, these 61 genes (Table S5 tab "Core genes") encoded for some

of the aforementioned functions such as thiol and metal homeostasis (*trxAB, gst, gor,*

sloA, dpr, and *sufCDSUB,*), enzymes for ROS scavenging (*sodA, ahpCF,* and *tpx*) and

restoration of redox balance (*nox*, ascorbate operon, *gapN*, and multiple *nrd* genes),

glucan-binding protein *gbpC*, proteins implicated in general stress management

including *clpE, clpL*, *uvrA, mutY,* SMU.503c (48), *exoA, lexA* (49), *pepP,* and *pepO* (50,

51), and transcription regulators such as *scnKR*, *perR,* and *msmR.* Again, nearly all 61

genes (except for 2) showed the same polarity of change under three treatments.

Notably, many genes shared by glucose and fructose, e.g., *sloA, sodA, gor, tpx, gst,*

trxAB, and *lexA*, showed significantly greater change due to the treatment by fructose

than by glucose of the same amount.

Also included in this group of genes was the CRISPR1-Cas system (SMU.1402c to

SMU.1406c) that was repressed by all four treatments, although the effect of H_2O_2 was

slightly below the 2-fold threshold (Table S1). Conversely affected was the 10-gene

CRISPR2-Cas cluster (SMU.1753c to SMU.1764c) that produced increased levels of

transcripts under all four conditions, despite that the induction of several genes by

methylglyoxal or glucose was below the threshold. An earlier study characterizing these

two systems in *S. mutans* suggested that CRISPR-Cas overall was required for

tolerance towards membrane, thermal, pH, and oxidative stressors (52). The differential

regulation of these two systems in response to carbohydrates, methylglyoxal, and H_2O_2

warrants further exploration. It is also worth noting that the relative impact of fructose to

both clusters was the highest among all four treatments.

Similarity between responses to fructose and glucose. Considering the effects of

carbohydrates in increasing osmolarity and acid production, it was not surprising to find

a substantial overlap between fructose- and glucose-responsive genes (total 165 genes,

80 genes when excluding methylglyoxal and H_2O_2; Supplemental Table S6), although

many (near 50%) encoded hypothetical proteins. Interestingly, the operon required for

malolactic fermentation (SMU.137-SMU.141) were most highly induced by either

treatment, with the mRNA levels increasing by 12~20-fold. Considering the inducibility

of this system by acidic pH (53, 54), a rapid drop in pH due to glycolytic activities likely

triggered this response. The agmatine deiminase (*aguA*) that has been shown to

contribute to acid tolerance through production of ammonia (44) was similarly

upregulated. Also induced by both carbohydrates were the majority of the genes/ORFs

included in the genomic island TnSmu1 (55), from SMU.196c to SMU.215c, showing an

increase in expression by 2~4-fold. Other notably upregulated genes included the

SMU.1602/*lguL* cluster, the zinc/cadmium-exporter *zccE* (56), the glucosyltransferase

gtfD, the fructosyltransferase *ftf*, a putative F-1-P kinase *pfk* (17), the lactose operon

(57), and the HPr kinase/phosphatase *hprK*. The fact that SMU.1602/*lguL* cluster was

induced by methylglyoxal, fructose, and glucose suggested that endogenous production

of methylglyoxal or related RES likely increased because of rapid influx of

carbohydrates. Like before, *zccE* expression was >2-fold higher when treated with

fructose compared to treatment by glucose. One cluster, SMU.1903c to SMU.1913c,

along with the orphan response regulator *gcrR* (58), showed reduced expression under

both treatments.

Genes affected only by fructose. A total of 168 genes were differentially affected by

fructose alone (Supplemental Table S6 tab "Fru only", Fig. S2A for enriched molecular

functions). In addition to what have been discussed thus far, treatment with 50 mM

fructose altered the expression of several pathways and additional genes required for

carbohydrate metabolism. Namely, fructose upregulated the operons of *sppRA*, *fruRKI*,

levDEFG, and the sorbitol PTS, however downregulated the nigerose PTS, the maltose

PTS (*malT* or *ptsG*), *pfl,* the *msm* cluster (59), *galK*, *gtfC,* the glycogen operon (*glg*)

(60), an α-glucan metabolic cluster (SMU.1564 to SMU.1571) (61), and importantly the

fructose-bisphosphate aldolase (*fbaA*, down 2-fold) which is responsible for cleaving F-

1,6-bP into dihydroxyacetone phosphate (DHAP) and glyceraldehyde 3-phosphate

(GADP). Expression of three additional clusters was notably affected by fructose alone.

First, SMU.1057 to SMU.1063, encoding SatCDE, Ffh, YlxM, and two ABC transporters,

were upregulated by 2- to 3-fold. This cluster has been implicated in protein

translocation and acid tolerance in *S. mutans* (62). Second, operon (SMU.1527 to

SMU.1534) which encodes for the F_1F_0-ATPase complex were downregulated by 2- to

3-fold. It was reported that the F_1F_0-ATPase, whose main function in *S. mutans* is

expelling excess proton to the environment, was upregulated in response to a shift from

steady state at pH 7 to pH 5 as part of the acid tolerance response (ATR) (63-65).

Despite the likelihood of an acidic stress under the treatment of both glucose and

fructose, there was no significant impact by glucose to this operon, and the treatment of

fructose resulted in a lower rather than higher expression of all 8 genes encoding the

F_1F_0-ATPase. Last, SMU.1734 to SMU.1745c, encoding for the pathway for fatty acid

biosynthesis and similarly involved in ATR, showed 2- to 3-fold reduction in expression

when treated with 50 mM fructose. While this effect was largely consistent with previous

findings made in *S. mutans* under a glucose shock at 200 mM (65), here again 50 mM

glucose failed to induce a similar response. Considering the importance of both

pathways in maintaining membrane integrity and overall fitness, these findings could

provide explanations to previously reported fructose-specific phenotypes such as

autolysis and increased biofilm development (23, 30).

RT-qPCR analysis. Detailed transcriptional analyses were carried out to validate and

further understand these findings, focusing on a list of genes relevant to bacterial fitness

and pathogenicity. Considering the dynamic nature of gene regulation, mRNA

abundance was compared in *S. mutans* UA159 treated with 50 mM fructose or 0.5 mM

H_2O_2, each for 5 min and 30 min. Also included were the original treatments (30 min) of

5 mM methylglyoxal and 50 mM glucose, as well as two control conditions of growth in

FMC-glucose and FMC-fructose. As shown in Table 1, mRNA levels of these 20 genes

in the 4 treatments used in RNA-seq study confirmed their respective impact, including

upregulation in functions such as radical scavenging, RES degradation, copper export,

thiol homeostasis, and stress-responsive regulators; and downregulation of *mreC*,

cellobiose PTS component *celB*, and mutacin gene *nlmA*. Thirty-min compared to 5-min

treatment with H_2O_2 generally gave more substantial activation of most genes tested.

Similarly, a time-dependent response to fructose was noted for genes *lguL*, *sloA*, *celB*,

lexA, *gshAB,* and *sppA*, although several did show signs of reversal in impact, namely

scnK, gbpC, nrdG, and SMU.667. To test the significance of F-1-P to gene regulation, a

fruI null mutant was treated with 50 mM fructose for 30 min. To our surprise, only 4 out

16 tested genes showed a reversal in impact due to loss of FruI, suggesting that FruI-

mediated effect plays only a partial role in the extensive transcriptomic reprogramming

triggered by the fructose treatment.

Fructose impacts metal homeostasis and Spx-dependent oxidative stress

response.

A likely effect of methylglyoxal on bacterial physiology was the reduction of

intracellular GSH/GSSG ratio and potential impact on metal homeostasis. Previously it

was demonstrated that a zinc/cadmium exporter ZccE was responsible for the high zinc

tolerance of *S. mutans*, and its expression required a cognate regulator ZccR (56).

Deletion of either *zccE* or *zccR* resulted in a peroxide sensitivity under high-zinc

conditions due to a disturbed zinc/manganese ratio which could be alleviated by

manganese supplementation. When we tested the growth phenotype of a *zccR* null

mutant in TV medium (66), which contains ~19 μM of zinc according to ICP-OES

analysis (56), the mutant nonetheless grew normally when supported by glucose as the

sole carbohydrate source (Fig. 3A). When fructose was substituted for glucose

however, $\Delta zccR$ showed a significant reduction compared to the WT UA159 in both

growth rate and the final optical density (Fig. 3B). Like reported before, addition of 25

μM MnSO$_4$ to the TV-fructose medium significantly alleviated the growth deficiency of

$\Delta zccR$. Importantly, after we introduced a *fruI* deletion into the *zccR* background, the

double mutant grew at levels comparable to the WT UA159 on either glucose or

fructose, without manganese supplementation (Fig. 3B). As *fruI* deletion was previously

shown to reduce intracellular F-1-P levels (23), these findings strongly supported the

theory that excess F-1-P levels negatively impacted metal homeostasis in *S. mutans*.

Since fructose-mediated transcriptome included a cohort of oxidative stress genes

primarily regulated by SpxA1 (67), we repeated this growth assay by including a

deletion mutant of *spxA1*. To our surprise, $\Delta spxA1$ grew at a faster rate in TV-fructose

than in TV-glucose (Fig. 4A). Considering the relatively high zinc content in TV medium,

we next switched to a chemically defined FMC medium that contained very little zinc but

sufficient manganese (around 110 μM). In FMC medium, $\Delta spxA1$ grew relatively well on

glucose but could not grow on fructose (Fig. 4B). When we introduced the same *fruI*

deletion into the $\Delta spxA1$ background, the growth deficiency on fructose was greatly

alleviated (Fig. 4C). At the same time, only modest benefit was observed on fructose

when a *levD* deletion was introduced into $\Delta spxA1$. Although the exact mechanisms

remain to be clarified, the medium-specific fructose sensitivity of $\Delta spxA1$ added further

support to the notion that F-1-P affects oxidative stress response and metal

homeostasis by interacting with Spx-mediated regulation.

To further test this hypothesis, a P*sodA::gfp* promoter-reporter fusion was established

in the WT UA159 background to assess the response of Spx-mediated regulation to the

presence of fructose; the superoxide dismutase (*sodA*) requires Spx proteins for optimal

expression (36). As shown in Fig. 5 and Fig. S3, the *sodA* promoter was activated by

the presence of fructose in a dose-dependent manner, from levels as low as 0.5 mM,

yet it only produced a modest and slightly delayed response to the presence of 0.5 mM

of H_2O_2. While 0.5 mM H_2O_2 significantly impeded the growth of *S. mutans* [Fig. S4 and

(68)], as much as 10 mM fructose showed little to now effect. To ascertain the nature of

fructose-derived signal in this effect, *fruI* and *levD* deletions were each introduced into

UA159 containing the P*sodA::gfp* fusion. Consistent with above observations, deletion

of *fruI* but not *levD* resulted in the loss of fructose-dependent activation of the *sodA*

promoter activity. As expected, deletion of *spxA1* also abolished the expression of *sodA*

promoter (36). These results not only confirmed the role of F-1-P in influencing Spx

regulation, but given the magnitude of impact, suggested that fructose may serve as a

primary signal for bacterial response to environmental changes.

F-1-P is the likely allosteric inducer for activating the *fruRKI* operon. To assess the

levels of fructose needed to induce the expression, we utilized a strain containing a

promoter-reporter fusion P*fruR::cat* (20). Bacteria were cultured in glucose-based FMC

medium and then exposed to different amounts of fructose for 30 min. CAT activities

from UA159/P*fruR::cat* suggested that at levels as low as 20 μM, fructose can

significantly induce the expression of the *fruRKI* operon (Table S7).

Fructose contributes to fitness during starvation, acidic pH, and competition with

peroxigenic streptococci.

Our previous studies have suggested that fructose, when used in excess under

laboratory conditions, may induce autolysis and cell death. Paradoxically, considering

the activation of much of the stress response core by fructose, we could anticipate

improved persistence under stressful conditions. To test this hypothesis, we subjected

S. mutans to a series of physiological tests by exposing it to fructose. First, strain

UA159 was cultivated in FMC with 20 mM glucose or fructose, and their CFU was

monitored while these cultures were continuously incubated without medium

refreshment for 11 days. While fructose produced lower CFU than glucose at day 1 of

the incubation as reported previously (23), it maintained the CFU of UA159 by at least 1

log higher than on glucose throughout the rest of the experiment (Fig. 6A). When 1 mM

fructose was used together with 19 mM glucose, the fructose-associated effects were

reduced and limited to the first two days of the incubation. We then repeated this

experiment using strain Δ*fruI* (Fig. 6B)*.* Echoing the findings in RT-qPCR analysis, Δ*fruI*

showed significantly higher persistence than the wild-type parent, especially when

fructose was used.

As a further test of acid tolerance under the fructose condition, we cultured UA159 for

20 h in a TV medium (FMC contains 10 mM phosphate buffer thus not ideal) supported

with different amounts of glucose or fructose and measured the resting pH of these

cultures. The results (Table 2) showed comparable growth yield but significantly lower

resting pH in TV-fructose cultures than in TV-glucose cultures. For comparison, we

conducted the same experiment on a health-associated commensal, *S. sanguinis* (SSA)

421 strain SK36. To our surprise, SK36 failed to produce a substantial growth on fructose,

422 especially for higher concentrations (100 or 200 mM). Similar findings were made when

423 this experiment was repeated using another *S. sanguinis* isolate BCC23 (69). Further

424 tests on a number of low-passage, clinical isolates of additional commensal streptococci

425 (70), encompassing the species of *S. gordonii, S. mitis, S. oralis, S. dentisani, S.*

426 *intermedius, S. sp.* A12*, S. parasanguinis,* and *S. cristatus*, showed similar sensitivity in

427 most but not all species (e.g., *S. intermedius*) tested (Table 2). On the contrary, tests on

428 6 additional WT *S. mutans* strains (Table S8) indicated that most of them were able to

429 grow comparably well on fructose, and lowered the environmental pH to similar levels to

430 that achieved on glucose. However, one isolate, OMZ175 of serotype *f* known for its

431 collagen-binding and invasive activities (71), did show a modest sensitivity to fructose

432 by having lower final OD_{600} and notably higher resting pH on fructose than glucose. We

433 then performed a pH drop assay to further test the notion that fructose could lower the

434 culture pH better than glucose. When a suspension of UA159 cells from an exponential

435 phase, fructose-grown culture was fed 50 mM of fructose without buffering, it lowered

436 the environmental pH at a notably faster rate, which ended at a significantly lower point,

437 than a glucose-grown culture of UA159 did in 50 mM glucose (Fig. 7A and Fig. S5). It

438 appeared that *S. mutans* UA159 catabolized fructose more rapidly than glucose and

439 was more acid-tolerant under fructose conditions. We also performed this assay using

440 BHI-grown bacteria for pH drops with glucose or fructose, which showed a similar, albeit

441 smaller difference between these two sugars. We conclude that while most *S. mutans*

442 strains tested by us appear well adapted to fermenting fructose, *S. sanguinis* and many

443 commensal streptococci may be less capable of tolerating high levels of fructose.

Next, *S. mutans* was subjected to a competition assay in a mixed-species planktonic

culture with the peroxigenic commensal *S. sanguinis*. Cultures from the exponential

phase were mixed at approximately 1:1 ratio between two species and diluted into the

FMC medium supplemented with 20 mM of glucose or fructose. After an overnight of

incubation, CFUs for each species were enumerated by plating on selective agar plates

(Fig. S6 includes CFU from mixed-species cultures and single-species controls). The

resultant competitive indices (Fig. 7B) indicated a greater competitiveness for *S. mutans*

UA159 in the presence of fructose than glucose, especially when used at higher

concentrations. It is perhaps worth noting that this competitiveness of UA159 favored by

fructose was at least partly attributable to the fructose sensitivity of its competitor, SK36,

a phenotype that could be species- and even strain-specific (Table 2). Interestingly,

deleting *spxB* in SK36 did not significantly change the competitive indices (Fig. S6E, 20

mM carbohydrates) whereas deleting *fruI* in UA159 further enhanced the

competitiveness of *S. mutans* against *S. sanguinis* SK36 when supported with fructose.

Since Δ*fruI* has a defect in fructose transport and grows more slowly than the WT on

fructose (17), these and earlier results (Fig. 6B) suggested that a certain optimal level of

F-1-P could benefit *S. mutans* when competing against commensal species capable of

producing inhibitory levels of H_2O_2.

463   **Discussion**

It is frequently observed in stress biology that different stressors result in overlapping

responses in bacteria (72). As an oral bacterium, *S. mutans* is frequently subjected to

environmental insults that include acidic pH, oxygen and H_2O_2, hyperosmolarity, thermal

fluctuation, metal ions, and nutrient starvation (73). In this study, we identified

methylglyoxal and fructose as two potential stress signals capable of inducing

streptococcal stress responses (Fig. 8). A stress core comprised of 61 genes were

shared by the regulons of fructose, methylglyoxal, and H_2O_2. This commonality in gene

regulation can be attributed to the overlapping effects exerted by these treatments. For

example, exposure to concentrated sugars likely leads to acidic stress, hyperosmolarity,

and increased RES production. Treatments by both RES and ROS are expected to

perturb intracellular redox pools that could in turn influence bacterial metal and thiol

homeostasis. Exposure to extreme pH, RES, and ROS can all lead to denaturation of

amino acids, proteins, lipids, and nucleic acids. Each of these outcomes likely activates

a central regulatory system to restore essential nutrients or repair damage. Not

surprisingly, 40 out of the 61 core stress genes (Table S5) have been previously

identified in *S. mutans* by an acid adaptation study in response to either a glucose

shock (200 mM) or a shift from neutral to acidic pH (65). Conversely, only 17 out of the

61 stress-related genes were affected by our treatment of 50 mM glucose. The

discrepancy between our study and earlier findings by others (65) is likely due to the

difference in sugar content and culture conditions, as 200 mM glucose and continuous

culturing were used before as opposed to 50 mM glucose and batch cultures herein.

Corroborating the transcriptomic overlap between fructose and acid tolerance,

phenotypic analyses suggested a superior ability for *S. mutans*, particularly certain

strains such as UA159, GS-5 and ST1 (Table S8), to lower environmental pH when

fructose was used as the supporting carbohydrate, especially at higher concentrations.

This ability to tolerate fructose appears to be limited or absent in many commensal

streptococcal species we have tested so far, with potential species and strain specificity,

which may have important implications to microbial ecology in oral biofilms.

Furthermore, several pathways triggered by hyperosmolarity in previously studies (74-

76), including genes *lguL, sodA, nox*, the *opcA* cluster (SMU.2116 to SMU.2119), the

opu/ffh/sat cluster, and potassium uptake mechanism *trkAB*, were also identified as part

of the fructose/methylglyoxal/H_2O_2 response core.

Theories established in mostly Gram-negative bacteria suggest that methylglyoxal

bypass, a function that converts DHAP into pyruvate via the activities of a methylglyoxal

synthase (*mgsA*), two glyoxalases (*gloAB*), and a D-lactate dehydrogenase (D-*ldh*),

serves an important function in modulating glycolysis under sugar excess and

phosphate limitation (32). Genes *mgsA* and D-*ldh* are absent in most lactic acid bacteria

(LAB) genomes (30). However, methylglyoxal and related RES can be produced from

DHAP spontaneously or from a variety of other mechanisms (77). The conservation of

glyoxalases in streptococci (78) and the transcriptomic overlap among fructose,

glucose, and methylglyoxal support the notion that methylglyoxal metabolism is an

integral part of bacterial physiology, and the sources of methylglyoxal could be both

endogenous and exogenous. This conclusion is consistent with an emerging theory in

bacterial pathophysiology termed the aldehyde hypothesis (79), where a group of

reactive metabolic intermediates (RES) from the host and pathogens may serve as

potential antimicrobial effectors. This theory has supports from studies on LAB

pathogens that engage host immunity, including *S. pyogenes* and *S. agalactiae*, which

required glyoxalases for fitness and virulence (31, 78). For oral streptococci, the effects

of fructose on methylglyoxal metabolism could involve several mechanisms. First,

multiple RES compounds can be generated during fructose metabolism, given the

presence of three phosphorylation products, F-1-P, F-6-P, and F-1,6-bP. Glucose

metabolism generates only F-6-P and F-1,6-bP. DHAP is a direct product of cleavage of

F-1,6-bP; cleavage of F-1-P by an aldolase produces glyceraldehyde (GA) and DHAP;

and similarly, cleavage of F-6-P yields G3P and dihydroxyacetone (DHA). These

metabolites are either themselves RES, e.g., GA, or in the case of DHAP and DHA, can

generate methylglyoxal due to a reaction that is spontaneous or enzymatic in nature

(MgsA). Second, fructose degradation may be regulated differently than glucose

degradation, which could affect the accumulation or degradation of metabolic

intermediates that are precursors of RES. Rapid phosphorylation of fructose by

fructokinase and the resultant drawdown of ATP is a major factor in fructose-dependent

metabolic impact in mammalian cells (16). Although in bacteria, carbohydrates are

phosphorylated by PTS at the expense of PEP instead of ATP, here we showed a

notably faster rate of glycolysis by *S. mutans* UA159 degrading fructose than glucose.

As a result, fructose influx may alter the metabolism of F-6-P and F-1,6-bP, impacting

RES production and gene regulation even in Δ*fruI* background. For example,

accumulation of F-6-P and its degradation into DHA (instead of phosphorylation into F-

1,6-bP) could presumably be amplified during high-fructose treatment, especially if F-

1,6-bP cleavage is impeded. In support of these reasonings, fructose transcriptome

reported downregulation of the F-1,6-bP aldolase *fbaA* (by 2-fold), and many common

genes shared by fructose and glucose treatments produced a greater amplitude of

change under the fructose condition. As part of the effort to understand the genetic

mechanisms of fructose-specific biology, we have identified a novel 5-gene cluster, well

conserved in *S. mutans* and most streptococci, which encodes a putative F-6-P

aldolase and a glycerol dehydrogenase (80, 81). Products of these genes could

contribute to metabolism of F-6-P and DHA, as suggested by a study in *Listeria innocua*

(82); and their overexpression was recently implicated in the hypervirulent phenotypes

of a dominant GAS variant in England (83). Moreover, most Gram-positive bacteria

maintain a novel LguL paralog, GloA2, whose deletion in *S. mutans* resulted in a

fructose-specific phenotype that resembled methylglyoxal exposure (30). Highly

pertinent to our research, an earlier animal study demonstrated that high fructose corn

syrup (HFCS, 45% glucose and 55% fructose), despite producing no glucans, can

induce dental caries in a rat model as effectively as sucrose (84). Likewise, *in vitro*

studies by others reported that HFCS induced greater demineralization of tooth enamel

compared to a similar treatment by sucrose (85).

In addition to influencing RES metabolism, F-1-P homeostasis appears to be an

essential physiological signal in most streptococci. The *fruRKI/fruRBA* operon required

for catabolizing fructose via the F-1-P pathway is highly conserved in most Gram-

positive bacteria. Several oral streptococci also harbor the *sppRA* operon for

moderating the F-1-P levels (23). We previously showed that a *fruK* deficient strain

accumulated significant levels of F-1-P even at the absence of fructose and was highly

defective in growth on fructose, suggesting an endogenous source of F-1-P (e.g.,

dephosphorylation of F-1,6,-bP) and the need to keep its level within an optimal range

(23). Notably, the *sppRA* promoter required higher levels of fructose (25 to 50 mM) for

induction (23), suggesting that *sppRA* are expressed only when F-1-P is above a certain

threshold to avoid futile cycling between F-1-P and fructose. Conversely, we have

determined herein that 0.5 mM fructose is capable of activating Spx-mediated regulation

and as little as 20 μM of fructose efficiently induced the expression of the *fruRKI* operon.

We also observed in *S. sanguinis* the induction of *fruRBA* expression by similar levels of

fructose or 25% human sera (Table S7). It is worth noting that for diabetic or

hyperglycemic individuals, blood fructose levels are often significantly elevated as a

function of the polyol pathway (16), capable of activating the *fruRKI/fruRBA* pathway

(86). These findings allow us to posit that fructose regulates streptococcal function at

both high (mM) and low (μM) levels, with the former contributing to fitness during the

feast-and-famine cycle of the oral cavity and the latter in human circulatory system.

While we only showed that fructose can benefit *S. mutans* at mM concentrations, we

have good reasons to speculate that a similar benefit applies under μM levels of

fructose. Several studies have suggested that *fruRKI/fruRBA* genes are required for

streptococcal virulence during extraoral infections (19, 22, 83), where only sub-mM

fructose is available, although the mechanisms remain uncharacterized. We further

posit that fructose may trigger a surveillance mechanism against host-derived ROS and

RES molecules, which primes bacterial stress response and additional fructose-specific

pathways (Fig 8). Aside from FruR, SppR, LevQRST, and LacR that were previously

shown to be responsive to fructose, herein we identified several transcription regulators

whose expression appeared uniquely responsive to fructose treatment, including *spxA2,*

glnR, hdrR, and a two-component system *spaRK.* Targeting these regulators and F-1-P-

specific metabolism, i.e., GloA2, could help us unravel the molecular mechanisms

required for cells to optimize stress management in response to fructose. With many

oral streptococci existing as dual-niche organisms and the widespread use of fructose in

food, this research has implications in both oral microbial homeostasis and systemic

health.

Lastly, it is worth reiterating that a fructose-PTS mutant (*fruI*) of *S. mutans* displayed

reduced colonization and cariogenicity in an earlier rat study under high-sucrose

condition (24). This study revealed an important connection between fructose and

sucrose in *S. mutans* physiology, as sucrose metabolism releases significant quantities

of fructose both intracellularly and extracellularly, the latter of which can be

reinternalized by the PTS (33, 34, 87). While it is tempting to speculate that fructose

released from sucrose might contribute to virulence of *S. mutans* and influence the

persistence of certain commensal streptococci, more in-depth investigation is needed to

fully understand the role of fructose in caries development and biofilm dysbiosis.

Concluding remarks. By establishing a stress regulon shared by fructose,

methylglyoxal, and H_2O_2, our study revealed a unique identity of fructose in

streptococcal physiology as a fermentable substrate, a stressor, and potentially an

important environmental signal. Gene regulations associated with an F-1-P homeostasis

contributed to bacterial fitness under stressful conditions, especially for the cariogenic

pathobiont *S. mutans*. The overlapping nature of stress response in general also

suggests that fructose metabolism may affect the overall resilience against other

stressors such as acidic pH, ROS, RES, toxic metals, starvation, and hyperosmolarity;

conditions to be expected in both oral cavity and the blood stream (Fig. 8). Considering

the widespread conservation of some of these genetic features, most of the fructose-

mediated effects discussed here are likely applicable to other lactic acid bacteria which

have high relevance in both oral and systemic health.

607   **Materials and Methods**

Bacterial strains and culture conditions. Brain heart infusion (BHI) medium (Difco

Laboratories, Detroit, MI) was used to maintain *S. mutans* and other streptococcal

strains (Table 3). Liquid BHI was used to prepare for starter cultures for most of the

experiments. Antibiotics were used, only when necessary, at the following

concentrations: kanamycin (Km) 1 mg/mL, erythromycin (Em) 10 µg/mL, and

spectinomycin (Sp) 1 mg/mL. For cultures requiring specific carbohydrates, a semi-

defined Tryptone-Vitamin (TV) medium (66) or a synthetic medium FMC (88) was used,

each supplemented with specified carbohydrates. Selection between TV and FMC was

made based on the requirements of each experiment. Specifically, FMC was selected

as the base medium when certain carbohydrates were added at very low levels, 1 mM

or lower, to avoid potential contaminations from ingredients of TV. Similarly, FMC was

used in promoter reporter study to reduce background fluorescence. TV and FMC also

contain significantly different buffering capacity and levels of nutritional and toxic metal

ions. Other than growth curves, all cultures were incubated without agitation at 37˚C in

an aerobic atmosphere supplemented with 5% of CO_2.

For growth curves, bacterial cultures were diluted 100-fold into a TV or FMC medium

containing specified carbohydrates, overlaid with 70 µL mineral oil, and maintained at

37˚C in a Bioscreen C lab system (Labsystems Oy, Helsinki, Finland) that recorded the

optical density of the cultures at 600 nm (OD_{600}) every hour. A brief lateral shaking was

included before each reading to disperse aggregates. To monitor the expression levels

of the *sodA* promoter over time using a P*sodA::gfp* reporter fusion, bacterial cultures

from the exponential phase were similarly diluted into an FMC medium containing

specified amounts of carbohydrates and H_2O_2 and overlaid with mineral oil. For 20 h,

the OD_{600} and the relative fluorescence units (RFU) of the cultures (excitation 485/20

nm, emission 528/20 nm) were recorded hourly using a Synergy 2 Multi-Mode reader

(BioTek, Winooski, VT). The RFU results were subtracted with the RFU readings from a

control strain, which was of the same genetic background but without the *gfp* promoter

fusion, before normalization against the corresponding OD_{600} of the culture (89).

Construction of genetic mutants. Genetic mutants of *S. mutans* were constructed

through allelic exchange strategy and natural transformation facilitated by the use of

competence-stimulating peptide (30). Recombinant DNA fragments used in this study

were generated by PCR amplification and ligated together with antibiotic cassettes via

Gibson Assembly (23). Primers used in DNA amplification and RT-qPCR are listed in

Table S9. All mutants used in this study have been validated by PCR followed by

Sanger sequencing. Promoter: reporter fusions were constructed using the plasmid

pMC340B by inserting a promoter of interest in front of a promoterless *gfp* (green

fluorescent protein) or *cat* (chloramphenicol acetyltransferase) gene, followed by

integration at a distal site (between *mtlA* and *phnA*) of the chromosome unrelated to

functions studied here (20).

RNA deep sequencing, transcriptomic analysis, and reverse transcriptase

quantitative PCR (RT-qPCR). To compare bacterial gene regulation in response to

fructose metabolism and exposure to methylglyoxal, *S. mutans* strain UA159 was

cultivated to early exponential phase (OD_{600} = 0.4) in a chemically defined medium FMC

(88) supplemented with 20 mM glucose. Bacteria were harvested by centrifugation

(3,800× *g*, room temperature, 10 min), resuspended in fresh FMC containing 50 mM

fructose, 50 mM glucose, or 5 mM methylglyoxal. An equal portion of the culture was

kept in the original FMC-glucose medium as control. All samples were returned to

incubation for 30 min before being harvested for RNA extraction.

RNA extraction and deep sequencing were carried out by following a previously

published protocol detailed elsewhere (90). Briefly, cells were disrupted using a

beadbeater in the presence of equal volume of acidic phenol:chloroform. After

centrifugation, the cell lysate was processed using the RNeasy minikit (Qiagen,

Germantown, MD) and an RNase-free DNase I solution (Qiagen) for extraction of total

RNA. The RNA samples were shipped on dry ice to SeqCenter (Pittsburgh, PA) for

quality check, cDNA preparation, and deep sequencing, yielding up to 12 million paired

end reads (2 × 51 bp) per sample. The analysis of the RNA-seq data was performed in

software R version 3.5.2 Eggshell Igloo as described previously (90). The compositional

matrix of the expression data was normalized with the voom (91) function from the R

package limma (92). The statistical analysis of the expression data (Table S10) was

conducted using the package edgeR (version 3.24.3). A false-discovery rate (FDR) of

0.05 and a fold-of-change of 2.0 were used as the cutoff values for identification of

genes with differential expression, consistent with the published work on H_2O_2

transcriptome in *S. mutans* (36). Findings from this analysis were validated by reverse

transcriptase quantitative PCR (RT-qPCR) (30) carried out on 20 selected genes,

following a previously published protocol detailed elsewhere (93, 94). The sequencing

data from this study have been deposited at NCBI Gene Expression Omnibus (GEO)

under the accession numbers GSE279080.

For comparative analysis under different treatments, we obtained the RNA-seq

datasets (GEO, GSE98526) of an analysis previously performed on strain UA159 that

was treated with 0.5 mM H_2O_2 for 5 min (36). Comparison between transcriptomes was

carried out in R statistical language using the package edgeR as done previously (95).

Long-term persistence assay. For comparison of the ability to persist under low-pH

and starvation conditions, the cultures of *S. mutans* were each diluted 1000-fold into an

FMC medium supplemented with 20 mM glucose, 20 mM fructose, or 19 mM glucose

and 1 mM fructose combined. The cultures were incubated at 37°C in an ambient

atmosphere supplemented with 5% CO_2 for a duration of 11 days. After one or two days

of incubation, a 30-second sonication at 100% power (FB120 water bath sonicator,

Fisher Scientific) was applied to all cultures to disperse bacterial chains and

aggregates, then a 100-μL aliquot of each culture was removed. The cultures were

returned to incubation without any other treatments, and the 100-μL aliquots were each

diluted decimally in sterile PBS and plated on BHI agar. After 2 days of incubation, the

CFU on each plate were enumerated. Plating was repeated at specified time points

thereafter till end of the experiment.

Mixed-species competition in planktonic cultures. For interspecies competition in

liquid cultures, differentially marked strains of *S. mutans* (UA159-Km and Δ*fruI::Km*) (96)

and *S. sanguinis* (MMZ1945, Em resistant) (97) were each cultured in BHI to the

exponential phase (OD_{600} = 0.5). An inoculum of approximately 10^6 CFU/ml of *S.*

mutans together with similar numbers of *S. sanguinis* were added to FMC

supplemented with 20, or 200 mM glucose or fructose, then cultured for 20 h in a 5%-

CO_2 environment at 37°C. At both the start and the end of the incubation, cultures were

sonicated for 15 sec, decimally diluted and plated on respective antibiotic plates to

enumerate CFU of both species. The competitive index was calculated by following this

formula: [SMU(t_{end})/SSA(t_{end})] / [SMU(t_{start})/SSA(t_{start})], with values >1 indicating SMU (*S.*

mutans) being more competitive than SSA (*S. sanguinis*), and vice versa.

pH drop and chloramphenicol acetyltransferase (CAT) assay. Glycolytic profiling via

pH drop (98) and CAT assay (99) were conducted according to previously published

protocols detailed elsewhere.

Statistics. Statistical analysis of data was carried out using the software of Prism

(GraphPad of Dotmatics, San Diego, CA)

Acknowledgements

This study was supported by a grant DE12236 from NIDCR and a startup fund from

University of Florida to LZ. JL was supported by grants DE032555 and DE019783 from

NIDCR. Parts of Fig. 8 were created using BioRender.com.

Fig. 1 Volcano plots illustrating transcriptomes of four treatments. *S. mutans*

UA159 was grown to the exponential phase in a synthetic FMC medium containing 20

mM glucose. The samples were treated for 30 min with 5 mM methylglyoxal, 50 mM

fructose, or 50 mM glucose before harvesting for mRNA sequencing. Differential

expression analyses were performed against untreated cells using edgeR, with cutoffs

for foldchange >2 and FDR <0.05. Upregulated genes are shown in red, and

downregulated ones in blue. Results of treatment by H_2O_2 (0.5 mM, 5 min) were from an

independent study (36).

Fig. 2 Transcriptomic overlaps among treatments of methylglyoxal, fructose,

glucose and H_2O_2. The Venn diagrams show the numbers of shared and unique genes

in the genome of UA159, separated into groups of upregulated and downregulated,

among treatments of (A) 5 mM methylglyoxal (MG) and 0.5 mM H_2O_2, (B) MG and 50

mM fructose (Fru), (C) MG and 50 mM glucose (Glc50), and (D) all four treatments.

Fig. 3 Fructose impacts metal homeostasis in Δ*zccR* deficient in zinc expulsion.

S. mutans strains UA159 and its mutant derivatives were diluted from exponential

phase cultures into TV supplemented with 20 mM glucose (A) or fructose (B). Mn^{2+} was

added at 25 μM ($MnSO_4$). Growth was monitored using a Synergy 2 plate reader

maintained at 37°C. Results are average of three biological replicates, each tested in

technical duplicates. Error bars denote standard deviations.

Fig. 4 Fructose impacts growth of *spxA1* in a medium-specific manner. *S. mutans*

strains UA159 and various mutant derivatives were each diluted from exponential phase

cultures into TV (A) or FMC (B, C) supplemented with 20 mM glucose (Glc) or fructose

(Fru). Growth was monitored using a Synergy 2 plate reader maintained at 37°C.

Results are average of three biological replicates, each tested in technical duplicates.

Error bars denote standard deviations.

Fig. 5 Induction of the *sodA* promoter by fructose. (A) Cultures of UA159 harboring

a P*sodA::gfp* fusion were diluted into FMC containing specified concentrations (in mM)

of glucose (G), fructose (F), or H_2O_2. The cultures were monitored in a Synergy 2 plate

reader for green fluorescence and optical densities (OD_{600}) for 20 hours. (B) The same

experiment was performed on various mutants derived from UA159/P*sodA::gfp,* using

FMC containing 20 mM glucose (G20) or fructose (F20). The relative fluorescence units

(RFU) of each culture were recorded as a measure of *sodA* promoter activity,

subtracted with RFUs of a control strain (the same genetic background but without the

gfp fusion) cultured under the same condition, and normalized against corresponding

OD_{600} values of the bacterial cultures. Results are the average from three biological

replicates, conducted in technical duplicates. Error bars denote standard deviations.

Fig. 6 Fructose promotes long-term survival of *S. mutans*. Cultures of *S. mutans*

UA159 (A, B) or Δ*fruI* (B) were diluted into FMC containing 20 mM of glucose (G20) or

fructose (F20), or a combination of 19 mM glucose and 1 mM fructose (G19/F1),

followed by incubation without medium refreshment for 11 days. At specified time

points, an aliquot of culture was removed for CFU enumeration. Each strain was

represented by 3 independent cultures, with results presenting average and standard

deviation. Statistical significance was assessed using (A) One-Way ANOVA followed

with Dunnett's comparisons against G20 samples, and (B) Two-Way ANOVA followed

by Tukey's multiple comparisons against G20 samples (*, $P<0.05$; **, $P<0.01$; ***,

$P<0.001$; ****, $P<0.0001$).

Fig. 7 Fructose enhances acidogenicity (A) and competitiveness (B) of *S. mutans*.

(A) pH drop assay. UA159 was cultured in TV with 20 mM glucose or fructose to

exponential phase, harvested and resuspended in a solution of 50 mM KCl and 1 mM

$MgCl_2$, with OD_{600} adjusted to 5.0. The pH was adjusted to 7.2 slowly with 0.1 N KOH to

consume intracellular carbohydrate storage. Immediately after addition of 50 mM

glucose or fructose, same as the sugar used in the TV medium, the pH of the

suspension was monitored continuously on a stirring plate for the next 60 min. Each

experiment was repeated three times (Fig. S5), with representative results shown here.

After conversion into proton concentrations, area under the curve (AUC, see inset) of

each sample was calculated and used to assess statistical significance using Student's

t-test (*, $P < 0.05$). (B) Cultures of *S. mutans* wild type UA159 or Δ*fruI* were mixed at 1:1

ratio with a differentially marked *S. sanguinis* strain, then diluted at 1:100 into FMC

containing glucose (G) or fructose (F), each used at 20 mM or 200 mM. After 24 h of

incubation, the cultures were sonicated and used for CFU enumeration. CFUs from both

the start and end of the incubation (Fig. S6) were used to calculate competitive indices.

Results were the average of three independent repeats, with error bars denoting

standard deviations. Statistical significance was assessed using Two-Way ANOVA

followed by Tukey's multiple comparisons (*, $P<0.05$; **, $P<0.01$).

Fig. 8 Fructose metabolism triggers stress response in streptococci. Fructose is

internalized by oral streptococci primarily through a F-1-P-generating PTS transporter,

FruI, but also through two additional PTS permeases, including EII^{Lev} (LevDEFG) that

generates F-6-P. Compared to the glucose pathway, fructose is likely catabolized more

rapidly via glycolysis, during which RES such as methylglyoxal is accumulated and

impacts cytoplasmic redox balance represented by the ratios of GSH/GSSG and

NADPH/NADP$^+$. Increased F-1-P kinase activities (denoted by a plus sign) and reduced

expression of F-1,6-bP aldolase (by a minus sign) may further contribute to RES

production through impeded F-6-P phosphorylation and increased cleavage of F-6-P

into DHA. Also affected are the pools of intracellular metal ions. Sensing these and

other not-yet-characterized physiological changes and specific metabolites, the

bacterium reprograms a suite of stress responsive genes. The outcomes of exposure to

fructose include not only elevated activities in processing RES, through the functions of

LguL/GloB and related protein GloA2, but importantly enhanced tolerance to multiple

environmental stressors including ROS, RES, acidic pH, toxic metals, and nutrient

depletion.

Reference

1. Baker JL, Mark Welch JL, Kauffman KM, McLean JS, He X. 2024. The oral microbiome: diversity, biogeography and human health. Nat Rev Microbiol 22:89-104.

2. Hsu CL, Schnabl B. 2023. The gut-liver axis and gut microbiota in health and liver disease. Nat Rev Microbiol 21:719-733.

3. Willenborg J, Goethe R. 2016. Metabolic traits of pathogenic streptococci. FEBS Lett 590:3905-3919.

4. Bowen WH, Burne RA, Wu H, Koo H. 2018. Oral Biofilms: pathogens, matrix, and polymicrobial interactions in microenvironments. Trends Microbiol 26:229-242.

5. Tanner ACR, Kressirer CA, Rothmiller S, Johansson I, Chalmers NI. 2018. The Caries Microbiome: Implications for Reversing Dysbiosis. Adv Dent Res 29:78-85.

6. Spatafora G, Li Y, He X, Cowan A, Tanner ACR. 2024. The Evolving Microbiome of Dental Caries. Microorganisms 12.

7. Stangvaltaite-Mouhat L, Skudutyte-Rysstad R, Ko H, Stankeviciene I, Aleksejuniene J, Puriene A. 2024. Co-occurrence of dental caries and periodontitis: multilevel modelling approach. BMC Oral Health 24:149.

8. Strauss F-J, Espinoza I, Stähli A, Baeza M, Cortés R, Morales A, Gamonal J. 2019. Dental caries is associated with severe periodontitis in Chilean adults: a cross-sectional study. BMC Oral Health 19:278.

9.　Romandini P, Marruganti C, Romandini WG, Sanz M, Grandini S, Romandini M.

2024. Are periodontitis and dental caries associated? A systematic review with

meta-analyses. J Clin Periodontol 51:145-157.

10.　Tappy L, Lê KA. 2010. Metabolic effects of fructose and the worldwide increase

in obesity. Physiol Rev 90:23-46.

11.　Azevedo-Martins AK, Santos MP, Abayomi J, Ferreira NJR, Evangelista FS.

2024. The Impact of Excessive Fructose Intake on Adipose Tissue and the

Development of Childhood Obesity. Nutrients 16.

12.　Gillespie KM, White MJ, Kemps E, Moore H, Dymond A, Bartlett SE. 2023. The

Impact of Free and Added Sugars on Cognitive Function: A Systematic Review

and Meta-Analysis. Nutrients 16.

13.　Siddiqui SH, Rossi NF. 2024. Acute Intake of Fructose Increases Arterial

Pressure in Humans: A Meta-Analysis and Systematic Review. Nutrients 16.

14.　Staltner R, Burger K, Baumann A, Bergheim I. 2023. Fructose: a modulator of

intestinal barrier function and hepatic health? Eur J Nutr 62:3113-3124.

15.　Ting KKY. 2024. Fructose-induced metabolic reprogramming of cancer cells.

Front Immunol 15:1375461.

16.　Johnson RJ, Lanaspa MA, Sanchez-Lozada LG, Tolan D, Nakagawa T, Ishimoto

T, Andres-Hernando A, Rodriguez-Iturbe B, Stenvinkel P. 2023. The fructose

survival hypothesis for obesity. Philos Trans R Soc Lond B Biol Sci

378:20220230.

17.　Wen ZT, Browngardt C, Burne RA. 2001. Characterization of two operons that

encode components of fructose-specific enzyme II of the

sugar:phosphotransferase system of *Streptococcus mutans*. FEMS Microbiol Lett

205:337-42.

18.   Loo CY, Mitrakul K, Voss IB, Hughes CV, Ganeshkumar N. 2003. Involvement of

an inducible fructose phosphotransferase operon in *Streptococcus gordonii*

biofilm formation. J Bacteriol 185:6241-54.

19.   Cao Q, Dong Y, Guo C, Ji S, Nie M, Liu G, Wan X, Lu C, Liu Y. 2023. *luxS*

contributes to intramacrophage survival of *Streptococcus agalactiae* by positively

affecting the expression of *fruRKI* operon. Vet Res 54:83.

20.   Zeng L, Chakraborty B, Farivar T, Burne RA. 2017. Coordinated regulation of the

EII^{Man} and *fruRKI* operons of *Streptococcus mutans* by global and fructose-

specific pathways. Appl Environ Microbiol 83:e01403-17.

21.   Jesionowski AM, Mansfield JM, Brittan JL, Jenkinson HF, Vickerman MM. 2016.

Transcriptome analysis of *Streptococcus gordonii* Challis DL1 indicates a role for

the biofilm-associated *fruRBA* operon in response to *Candida albicans*. Mol Oral

Microbiol 31:314-328.

22.   Valdes KM, Sundar GS, Vega LA, Belew AT, Islam E, Binet R, El-Sayed NM, Le

Breton Y, McIver KS. 2016. The *fruRBA* operon is necessary for group A

streptococcal growth in fructose and for resistance to neutrophil killing during

growth in whole human blood. Infect Immun 84:1016-31.

23.   Zeng L, Burne RA. 2019. Essential roles of the *sppRA* fructose-phosphate

phosphohydrolase operon in carbohydrate metabolism and virulence expression

by *Streptococcus mutans*. J Bacteriol 201:e00586-18.

24. Tanzer JM, Thompson A, Wen ZT, Burne RA. 2006. *Streptococcus mutans*:

fructose transport, xylitol resistance, and virulence. J Dent Res 85:369-73.

25. Gauthier L, Mayrand D, Vadeboncoeur C. 1984. Isolation of a novel protein

involved in the transport of fructose by an inducible phosphoenolpyruvate

fructose phosphotransferase system in *Streptococcus mutans*. J Bacteriol

160:755-63.

26. Zeng L, Wen ZT, Burne RA. 2006. A novel signal transduction system and

feedback loop regulate fructan hydrolase gene expression in *Streptococcus*

mutans. Mol Microbiol 62:187-200.

27. Ajdic D, McShan WM, McLaughlin RE, Savic G, Chang J, Carson MB, Primeaux

C, Tian R, Kenton S, Jia H, Lin S, Qian Y, Li S, Zhu H, Najar F, Lai H, White J,

Roe BA, Ferretti JJ. 2002. Genome sequence of *Streptococcus mutans* UA159, a

cariogenic dental pathogen. Proc Natl Acad Sci U S A 99:14434-9.

28. Xu P, Alves JM, Kitten T, Brown A, Chen Z, Ozaki LS, Manque P, Ge X, Serrano

MG, Puiu D, Hendricks S, Wang Y, Chaplin MD, Akan D, Paik S, Peterson DL,

Macrina FL, Buck GA. 2007. Genome of the opportunistic pathogen

Streptococcus sanguinis. J Bacteriol 189:3166-3175.

29. Makarewicz O, Lucas M, Brandt C, Herrmann L, Albersmeier A, Rückert C, Blom

J, Goesmann A, van der Linden M, Kalinowski J, Pletz MW. 2017. Whole

Genome Sequencing of 39 Invasive *Streptococcus pneumoniae* Sequence Type

199 Isolates Revealed Switches from Serotype 19A to 15B. PLoS ONE

12:e0169370.

30. Zeng L, Noeparvar P, Burne RA, Glezer BS. 2024. Genetic characterization of

glyoxalase pathway in oral streptococci and its contribution to interbacterial

competition. J Oral Microbiol 16:2322241.

31. Zhang MM, Ong CL, Walker MJ, McEwan AG. 2016. Defence against

methylglyoxal in Group A *Streptococcus*: a role for Glyoxylase I in bacterial

virulence and survival in neutrophils? Pathog Dis 74.

32. Cooper RA. 1984. Metabolism of methylglyoxal in microorganisms. Annu Rev

Microbiol 38:49-68.

33. Zeng L, Burne RA. 2016. Sucrose- and fructose-specific effects on the

transcriptome of *Streptococcus mutans*, as determined by RNA sequencing. Appl

Environ Microbiol 82:146-56.

34. Zeng L, Burne RA. 2013. Comprehensive mutational analysis of sucrose-

metabolizing pathways in *Streptococcus mutans* reveals novel roles for the

sucrose phosphotransferase system permease. J Bacteriol 195:833-43.

35. Kajfasz JK, Zuber P, Ganguly T, Abranches J, Lemos JA. 2021. Increased

oxidative stress tolerance of a spontaneously occurring *perR* gene mutation in

Streptococcus mutans UA159. J Bacteriol 203.

36. Kajfasz JK, Ganguly T, Hardin EL, Abranches J, Lemos JA. 2017. Transcriptome

responses of *Streptococcus mutans* to peroxide stress: identification of novel

antioxidant pathways regulated by Spx. Sci Rep 7:16018.

37. Noctor G, Foyer CH. 1998. ASCORBATE AND GLUTATHIONE: Keeping Active

Oxygen Under Control. Annu Rev Plant Physiol Plant Mol Biol 49:249-279.

38.   Zheng X, Zhang K, Zhou X, Liu C, Li M, Li Y, Wang R, Li Y, Li J, Shi W, Xu X.

919   2013. Involvement of *gshAB* in the interspecies competition within oral biofilm. J

920   Dent Res 92:819-24.

921   39.   Ellepola K, Huang X, Riley RP, Bitoun JP, Wen ZT. 2021. *Streptococcus mutans*

922   lacking *sufCDSUB* Is viable, but displays major defects in growth, stress

923   tolerance responses and biofilm formation. Front Microbiol 12:671533.

924   40.   Wu X, Hou J, Chen X, Chen X, Zhao W. 2016. Identification and functional

925   analysis of the L-ascorbate-specific enzyme II complex of the

926   phosphotransferase system in *Streptococcus mutans*. BMC Microbiol 16:51.

927   41.   Makhlynets O, Boal AK, Rhodes DV, Kitten T, Rosenzweig AC, Stubbe J. 2014.

928   *Streptococcus sanguinis* class Ib ribonucleotide reductase: high activity with both

929   iron and manganese cofactors and structural insights. J Biol Chem 289:6259-72.

930   42.   Rhodes DV, Crump KE, Makhlynets O, Snyder M, Ge X, Xu P, Stubbe J, Kitten T.

931   2014. Genetic characterization and role in virulence of the ribonucleotide

932   reductases of *Streptococcus sanguinis*. J Biol Chem 289:6273-87.

933   43.   Stewart LJ, Ong CY, Zhang MM, Brouwer S, McIntyre L, Davies MR, Walker MJ,

934   McEwan AG, Waldron KJ, Djoko KY. 2020. Role of Glutathione in Buffering

935   Excess Intracellular Copper in *Streptococcus pyogenes*. mBio 11.

936   44.   Griswold AR, Chen YY, Burne RA. 2004. Analysis of an agmatine deiminase

937   gene cluster in *Streptococcus mutans* UA159. J Bacteriol 186:1902-4.

938   45.   Seaton K, Ahn S-J, Burne RA. 2014. Regulation of competence and gene

939   expression in *Streptococcus mutans* by the RcrR transcriptional regulator. Mol

940   Oral Microbiol 30:147-59.

46.   Merritt J, Qi F, Shi W. 2005. A unique nine-gene *comY* operon in *Streptococcus*

mutans. Microbiology 151:157-66.

47.   Merritt J, Qi F. 2012. The mutacins of *Streptococcus mutans*: regulation and

ecology. Mol Oral Microbiol 27:57-69.

48.   Banu LD, Conrads G, Rehrauer H, Hussain H, Allan E, van der Ploeg JR. 2010.

The *Streptococcus mutans* serine/threonine kinase, PknB, regulates competence

development, bacteriocin production, and cell wall metabolism. Infect Immun

78:2209-20.

49.   Leung V, Ajdic D, Koyanagi S, Levesque CM. 2015. The formation of

Streptococcus mutans persisters induced by the quorum-sensing peptide

pheromone is affected by the LexA regulator. J Bacteriol 197:1083-94.

50.   Alves LA, Ganguly T, Harth-Chu EN, Kajfasz J, Lemos JA, Abranches J, Mattos-

Graner RO. 2020. PepO is a target of the two-component systems VicRK and

CovR required for systemic virulence of *Streptococcus mutans*. Virulence 11:521-

536.

51.   Underhill SAM, Shields RC, Burne RA, Hagen SJ. 2019. Carbohydrate and PepO

control bimodality in competence development by *Streptococcus mutans*. Mol

Microbiol 112:1388-1402.

52.   Serbanescu MA, Cordova M, Krastel K, Flick R, Beloglazova N, Latos A, Yakunin

AF, Senadheera DB, Cvitkovitch DG. 2015. Role of the *Streptococcus mutans*

CRISPR-cas systems in immunity and cell physiology. J Bacteriol 197:749-761.

53.   Lemme A, Sztajer H, Wagner-Döbler I. 2010. Characterization of *mleR*, a positive

963   regulator of malolactic fermentation and part of the acid tolerance response in

964   *Streptococcus mutans*. BMC Microbiol 10:58.

965   54.   Sheng J, Marquis RE. 2007. Malolactic fermentation by *Streptococcus mutans*.

966   FEMS Microbiol Lett 272:196-201.

967   55.   King S, Quick A, King K, Walker AR, Shields RC. 2022. Activation of TnSmu1, an

968   integrative and conjugative element, by an ImmR-like transcriptional regulator in

969   *Streptococcus mutans*. Microbiology (Reading) 168.

970   56.   Ganguly T, Peterson AM, Burkholder M, Kajfasz JK, Abranches J, Lemos JA.

971   2022. ZccE is a novel P-type ATPase that protects *Streptococcus mutans* against

972   zinc intoxication. PLoS Pathog 18:e1010477.

973   57.   Zeng L, Das S, Burne RA. 2010. Utilization of lactose and galactose by

974   *Streptococcus mutans*: transport, toxicity, and carbon catabolite repression. J

975   Bacteriol 192:2434-44.

976   58.   Sato Y, Yamamoto Y, Kizaki H. 2000. Construction of region-specific partial

977   duplication mutants (merodiploid mutants) to identify the regulatory gene for the

978   glucan-binding protein C gene *in vivo in Streptococcus mutans*. FEMS Microbiol

979   Lett 186:187-91.

980   59.   Tao L, Sutcliffe IC, Russell RR, Ferretti JJ. 1993. Transport of sugars, including

981   sucrose, by the *msm* transport system of *Streptococcus mutans*. J Dent Res

982   72:1386-90.

60.   Costa Oliveira BE, Ricomini Filho AP, Burne RA, Zeng L. 2021. The Route of

Sucrose Utilization by *Streptococcus mutans* Affects Intracellular Polysaccharide

Metabolism. Front Microbiol 12:636684.

61.   Sato Y, Okamoto-Shibayama K, Azuma T. 2013. The *malQ* gene is essential for

starch metabolism in *Streptococcus mutans*. J Oral Microbiol 5:21285.

62.   Kremer BH, van der Kraan M, Crowley PJ, Hamilton IR, Brady LJ, Bleiweis AS.

2001. Characterization of the *sat* operon in *Streptococcus mutans*: evidence for a

role of Ffh in acid tolerance. J Bacteriol 183:2543-52.

63.   Gong Y, Tian XL, Sutherland T, Sisson G, Mai J, Ling J, Li YH. 2009. Global

transcriptional analysis of acid-inducible genes in *Streptococcus mutans*: multiple

two-component systems involved in acid adaptation. Microbiology 155:3322-32.

64.   Baker JL, Faustoferri RC, Quivey RG, Jr. 2017. Acid-adaptive mechanisms of

Streptococcus mutans-the more we know, the more we don't. Mol Oral Microbiol

32:107-117.

65.   Baker JL, Abranches J, Faustoferri RC, Hubbard CJ, Lemos JA, Courtney MA,

Quivey R, Jr. 2015. Transcriptional profile of glucose-shocked and acid-adapted

strains of *Streptococcus mutans*. Mol Oral Microbiol doi:10.1111/omi.12110.

66.   Burne RA, Wen ZT, Chen YY, Penders JE. 1999. Regulation of expression of the

fructan hydrolase gene of *Streptococcus mutans* GS-5 by induction and carbon

catabolite repression. J Bacteriol 181:2863-71.

67.   Kajfasz JK, Rivera-Ramos I, Scott-Anne K, Gregoire S, Abranches J, Lemos JA.

2015. Transcription of Oxidative Stress Genes Is Directly Activated by SpxA1

and, to a Lesser Extent, by SpxA2 in *Streptococcus mutans*. J Bacteriol

197:2160-2170.

68. Kreth J, Merritt J, Shi W, Qi F. 2005. Competition and coexistence between

Streptococcus mutans and *Streptococcus sanguinis* in the dental biofilm. J

Bacteriol 187:7193-7203.

69. Culp DJ, Hull W, Bremgartner MJ, Atherly TA, Christian KN, Killeen M, Dupuis

MR, Schultz AC, Chakraborty B, Lee K, Wang DS, Afzal V, Chen T, Burne RA.

2020. *In vivo* colonization with candidate oral probiotics attenuates colonization

and virulence of *Streptococcus mutans*. Appl Environ Microbiol 87.

70. Velsko IM, Chakraborty B, Nascimento MM, Burne RA, Richards VP. 2018.

Species designations belie phenotypic and genotypic heterogeneity in oral

streptococci. mSystems 3:e00158-18.

71. Miller JH, Avilés-Reyes A, Scott-Anne K, Gregoire S, Watson GE, Sampson E,

Progulske-Fox A, Koo H, Bowen WH, Lemos JA, Abranches J. 2015. The

collagen binding protein Cnm contributes to oral colonization and cariogenicity of

Streptococcus mutans OMZ175. Infect Immun 83:2001-10.

72. Papadimitriou K, Alegria A, Bron PA, de Angelis M, Gobbetti M, Kleerebezem M,

Lemos JA, Linares DM, Ross P, Stanton C, Turroni F, van Sinderen D, Varmanen

P, Ventura M, Zuniga M, Tsakalidou E, Kok J. 2016. Stress physiology of lactic

acid bacteria. Microbiol Mol Biol Rev 80:837-90.

73. Abranches J, Zeng L, Kajfasz JK, Palmer SR, Chakraborty B, Wen ZT, Richards

VP, Brady LJ, Lemos JA. 2018. Biology of oral streptococci. *In* Fischetti VA,

Novick RP, Ferretti J, Portnoy DA, Braunstein M, Rood JI (ed), Gram-positive

pathogens doi:doi:10.1128/microbiolspec.GPP3-0042-2018. Microbiol Spectrum.

74.   Liu C, Niu Y, Zhou X, Zhang K, Cheng L, Li M, Li Y, Wang R, Yang Y, Xu X. 2013.

Hyperosmotic response of *Streptococcus mutans*: from microscopic physiology

to transcriptomic profile. BMC Microbiology 13:275.

75.   Abranches J, Lemos JA, Burne RA. 2006. Osmotic stress responses of

Streptococcus mutans UA159. FEMS Microbiol Lett 255:240-6.

76.   Kunii M, Arimoto T, Hasegawa T, Kuwata H, Igarashi T. 2014. Role of protease

maturation lipoprotein in osmoadaptation of *Streptococcus mutans*. FEMS

Microbiol Lett 356:45-52.

77.   Lee C, Park C. 2017. Bacterial responses to glyoxal and methylglyoxal: reactive

electrophilic species. Int J Mol Sci 18.

78.   Akbari MS, Joyce LR, Spencer BL, Brady A, McIver KS, Doran KS. 2025.

Identification of glyoxalase A in group B *Streptococcus* and its contribution to

methylglyoxal tolerance and virulence. Infect Immun doi:10.1128/iai.00540-

24:e0054024.

79.   Darwin KH, Stanley SA. 2022. The aldehyde hypothesis: metabolic intermediates

as antimicrobial effectors. Open Biology 12:220010.

80.   Doi Y. 2019. Glycerol metabolism and its regulation in lactic acid bacteria. Appl

Microbiol Biotechnol 103:5079-5093.

81.   Taylor ZA, Chen P, Noeparvar P, Pham DN, Walker AR, Kitten T, Zeng L. 2024.

Glycerol metabolism contributes to competition by oral streptococci through

production of hydrogen peroxide. J Bacteriol 206:e0022724.

82.   Monniot C, Zébré AC, Aké FMD, Deutscher J, Milohanic E. 2012. Novel Listerial

1051   Glycerol Dehydrogenase- and Phosphoenolpyruvate-Dependent

1052   Dihydroxyacetone Kinase System Connected to the Pentose Phosphate

1053   Pathway. J Bacteriol 194:4972-4982.

1054   83.   Li HK, Zhi X, Vieira A, Whitwell HJ, Schricker A, Jauneikaite E, Li H, Yosef A,

1055   Andrew I, Game L, Turner CE, Lamagni T, Coelho J, Sriskandan S. 2023.

1056   Characterization of emergent toxigenic M1(UK) *Streptococcus pyogenes* and

1057   associated sublineages. Microb Genom 9.

1058   84.   Xiao T, Xia W. 2013. Cariogenicity of high fructose corn syrup: an experimental

1059   study on rats. Chin J Cons Dent 23:446–449.

1060   85.   Sun M, Kang Q, Li T, Huang L, Jiang Y, Xia W-W. 2014. Effect of high-fructose

1061   corn syrup on *Streptococcus mutans* virulence gene expression and on tooth

1062   demineralization. Eur J Oral Sci 122:216-22.

1063   86.   Kawasaki T, Akanuma H, Yamanouchi T. 2002. Increased fructose concentrations

1064   in blood and urine in patients with diabetes. Diabetes Care 25:353-7.

1065   87.   Tanzer JM, Chassy BM, Krichevsky MI. 1971. Sucrose metabolism by

1066   *Streptococcus mutans*, SL-I. Biochim Biophys Acta 261:379-87.

1067   88.   Terleckyj B, Willett NP, Shockman GD. 1975. Growth of several cariogenic strains

1068   of oral streptococci in a chemically defined medium. Infect Immun 11:649-55.

1069   89.   Zeng L, Chen L, Burne RA. 2018. Preferred hexoses influence long-term memory

1070   in and induction of lactose catabolism by *Streptococcus mutans*. Appl Environ

1071   Microbiol 84:e00864-18.

90.   Chen L, Walker AR, Burne RA, Zeng L. 2020. Amino sugars reshape interactions

between *Streptococcus mutans* and *Streptococcus gordonii*. Appl Environ

Microbiol 87:e01459-20.

91.   Law CW, Chen Y, Shi W, Smyth GK. 2014. voom: Precision weights unlock linear

model analysis tools for RNA-seq read counts. Genome Biol 15:R29.

92.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015. limma

powers differential expression analyses for RNA-sequencing and microarray

studies. Nucleic Acids Res 43:e47.

93.   Zeng L, Walker AR, Lee K, Taylor ZA, Burne RA. 2021. Spontaneous mutants of

Streptococcus sanguinis with defects in the glucose-PTS show enhanced post-

exponential phase fitness. J Bacteriol 203:JB0037521.

94.   Schmittgen TD, Livak KJ. 2008. Analyzing real-time PCR data by the

comparative C_T method. Nat Protocols 3:1101-8.

95.   Ranjit DK, Moye ZD, Rocha FG, Ottenberg G, Nichols FC, Kim HM, Walker AR,

Gibson FC, 3rd, Davey ME. 2022. Characterization of a Bacterial Kinase That

Phosphorylates Dihydrosphingosine to Form dhS1P. Microbiol Spectr

10:e0000222.

96.   Chen L, Chakraborty B, Zou J, Burne RA, Zeng L. 2019. Amino sugars modify

antagonistic interactions between commensal oral streptococci and

Streptococcus mutans. Appl Environ Microbiol 85:e00370-19.

97.   Zeng L, Walker AR, Burne RA, Taylor ZA. 2022. Glucose phosphotransferase

system modulates pyruvate metabolism, bacterial fitness, and microbial ecology

in oral streptococci. J Bacteriol 205:e0035222.

98.   Moye ZD, Zeng L, Burne RA. 2014. Modification of gene expression and

virulence traits in *Streptococcus mutans* in response to carbohydrate availability.

Appl Environ Microbiol 80:972-985.

99.   Zeng L, Burne RA. 2008. Multiple sugar: phosphotransferase system permeases

participate in catabolite modification of gene expression in *Streptococcus*

mutans. Mol Microbiol 70:197-208.

100.   Huang X, Palmer SR, Ahn SJ, Richards VP, Williams ML, Nascimento MM, Burne

RA. 2016. A highly arginolytic *Streptococcus* species that potently antagonizes

Streptococcus mutans. Appl Environ Microbiol 82:2187-201.

1104

Table 1. Relative expression of selected genes in *S. mutans* UA159 WT and Δ*fruI*

(shaded) with specified treatments. After growing to mid-exponential phase (OD_{600} =

0.5) in FMC-Glc (20 mM glucose) at 37°C in 5% CO_2, bacterial cultures were each

treated with: Fru50, 50 mM fructose for 5 or 30 min; MG, 5 mM methylglyoxal for 30

min; H_2O_2, 0.5 mM hydrogen peroxide for 5 or 30 min; or Glc50, 50 mM glucose for 30

min. All values were relative to the untreated WT cultivated with FMC-Glc (second

column). Exponential phase cultures of UA159 in FMC-Fru (20 mM fructose) were

included as additional controls. Results are average from 3 biological replicates.

Asterisks denote statistical significance assessed by One-Way ANOVA ((*, P <0.05; **,

P <0.01; ***, P <0.001; ****, P <0.0001). NT, not tested.

Genes	FMC-Glc cultures	FMC-Fru cultures	FMC-Glc cultures treated with the following						
			Fru50 5 min	Fru50 30 min	Δ*fruI* Fru50 30 min	MG 30 min	H_2O_2 5 min	H_2O_2 30 min	Glc50 30 min
lguL	1.00	2.27**	2.32**	2.87***	6.66****	2.06*	0.84	5.68****	2.31**
phoR	1.01	1.80	NT	1.64	2.74**	2.68**	0.97	NT	1.47
nlmA	1.01	3.86***	0.41	0.52	0.62	0.24	0.14*	0.43	NT
sloA	1.02	0.29****	0.29****	0.10****	0.04****	0.35****	0.06****	0.20****	0.56***
sloC	1.01	1.66*	NT	NT	NT	0.79	0.45	NT	NT
celB	1.16	0.38*	0.10**	0.06**	4.36****	0.05**	0.22**	0.17**	0.44*
mreC	1.00	0.85	0.61****	0.47****	0.44****	0.07****	0.45****	0.48****	0.52****
lexA	1.02	3.25*	2.04	9.17****	9.03****	6.32****	2.77	11.70****	NT
scnK	1.00	4.78**	5.16***	3.26*	5.56***	6.06***	5.64***	8.41****	1.67
gbpC	1.00	7.62****	6.22****	3.00*	2.32	4.01***	2.12	11.36****	NT
tpx	1.00	3.28	14.64****	NT	NT	10.01****	6.26**	23.32****	NT
ahpF	1.01	5.54*	6.35**	5.62**	11.13****	11.54****	6.17**	11.32****	0.92
copY	1.00	1.06	0.84	1.72	2.27	10.78****	1.21	1.18	1.75
nrdG	1.00	1.76	1.67	0.59	0.41	9.43****	0.48	0.80	NT
gshAB	1.00	1.15	1.33**	2.62****	2.27****	1.24	0.95	1.66****	2.02****
sppA	1.00	NT	3.87***	36.92****	0.71	NT	NT	1.21	NT
SMU.127	1.00	5.37*	4.63*	NT	NT	4.87*	2.91	8.93****	NT
SMU.383c	1.00	0.98	1.96***	NT	NT	2.19****	0.34**	2.08***	NT
SMU.667	1.00	1.75	7.36****	2.97****	0.97	5.68****	2.54***	5.54****	1.76
SMU.679	1.06	2.01*	3.76****	2.64***	1.29	4.55****	1.19	3.32****	1.83

1115

1116 **Table 2**. pH and final OD_{600} of 20-h TV cultures.
1117

Strains	Sugars	Resting pH			Final OD_{600}		
		20 mM	100 mM	200 mM	20 mM	100 mM	200 mM
SMU_UA159	Glucose	4.92	4.93	4.93	0.83	0.72	0.57
	Fructose	4.76****	4.82**	4.81***	0.82	0.73	0.64****
SSA_SK36	Glucose	6.5	6.15	5.70	0.34	0.41	0.45
	Fructose	6.55	ND	ND	0.27***	0.03****	0.00****
SSA_BCC23	Glucose	6.09	6.01	5.93	0.43	0.37	0.31
	Fructose	5.76***	ND	ND	0.49*	0.20****	0.02****
SDE_BCC10	Glucose	5.48	5.08	5.49	0.63	0.69	0.43
	Fructose	ND	ND	ND	0.28****	0.14****	0.00****
SOR_BCC11	Glucose	5.72	5.65	5.72	0.56	0.52	0.4
	Fructose	5.09****	5.56	ND	0.66****	0.55	0.01****
SPA_BCC15	Glucose	6.53	6.45	6.38	0.12	0.11	0.09
	Fructose	ND	ND	ND	0.02***	0.00***	0.00**
SMI_BCC2	Glucose	4.75	4.73	4.77	0.74	0.65	0.56
	Fructose	4.87**	4.87***	4.87**	0.61****	0.54****	0.47****
SGO_BCC9	Glucose	4.95	4.90	4.91	0.81	0.66	0.54
	Fructose	4.87	5.03	5.97****	0.70****	0.68	0.46***
A12_BCC21	Glucose	5.62	5.37	5.48	0.65	0.56	0.46
	Fructose	5.16***	5.36	ND	0.64	0.54	0.28****
SCR_BCC41	Glucose	6.20	5.93	5.85	0.5	0.47	0.42
	Fructose	5.69****	5.95	ND	0.51	0.41	0.10****
SIN_BCA19	Glucose	5.28	5.12	5.18	0.66	0.58	0.48
	Fructose	5.11***	5.01*	5.14	0.61****	0.55**	0.48

Results show the averages of four independent cultures. Statistical significance was determined
using Two-Way ANOVA followed by Bonferroni's test for comparisons with glucose cultures of
the same concentration (*, $P <0.05$; **, $P <0.01$; ***, $P <0.001$; ****, $P <0.0001$). ND, not
determined due to lack of substantial growth (in red). SMU, *S. mutans*; SSA, *S. sanguinis*; SDE,
S. dentisani; SOR, *S. oralis*; SPA, *S. parasanguinis*; SMI, *S. mitis*; SGO, *S. gordonii;* A12*, S. sp.*
A12; SCR*, S. cristatus;* SIN*, S. intermedius* (70, 100).

Table 3. Bacterial strains used in this study (excluding clinical isolates used in Table 2).

1125

Strains	Relevant characteristics[a]	Source or reference
S. mutans UA159	*S. mutans* wild type, *perR*[+]	ATCC 700610
MMZ1654	UA159 *fruI::Km*	From UA159
MMZ2120	UA159 *fruI::Em*	From UA159
ΔzccR	UA159 *zccR::Sp*	(56)
MMZ2156	UA159 *zccR::Sp fruI::Km*	From ΔzccR
ΔspxA1	UA159 *spxA1::Sp*	(36)
MMZ2161	UA159 *spxA1::Sp fruI::Em*	From ΔspxA1
MMZ2162	UA159 *spxA1::Sp levD::Em*	From ΔspxA1
ΔspxA2	UA159 *spxA2::Em*	(36)
MMZ2158	UA159 *sodA::gfp* fusion (*Km*)	From UA159
MMZ2160	UA159 *spxA1::Sp sodA::gfp* fusion	From ΔspxA1
MMZ2159	UA159 *fruI::Em sodA::gfp* fusion	From MMZ2120
MMZ2173	UA159 *levD::Em sodA::gfp* fusion	From MMZ2158
MMZ799	UA159 *fruR::cat* fusion (*Km*)	(20)
UA159-Km	UA159 *gtfA::Km*	(96)
S. sanguinis SK36	*S. sanguinis* wild type	Kitten laboratory
MMZ1945	SK36 *gtfP::Em*	(97)

1126 [a] *Km, Em,* and *Sp*: resistance against kanamycin, erythromycin, and spectinomycin,
1127 respectively.

1128

1129

A

TV

B

FMC

C

FMC

A UA159/ P*sodA::gfp*

B P*sodA::gfp*: Δ*fruI*, Δ*levD*, Δ*spxA1*

A legend:
—□— G20　—✳— G19.5_F0.5　—□— G19_F1　—■— G10_F10　—△— G20_H₂O₂_0.5

B legend:
—□— UA159_G20　—△— *fruI_G20*　—○— *levD_G20*　—△— *spxA1_G20*
—■— UA159_F20　—▲— *fruI_F20*　—●— *levD_F20*　—▲— *spxA1_F20*