# Ice and air: Visualisation of freeze-thaw embolism and freezing spread in

# young *L. tulipifera* leaves

Kate M. Johnson[1], Muriel Scherer[2], Dominic Gerber[2], Robert W. Style[2], Eric R. Dufresne[2,3], Craig R.

Brodersen[1]

[1] Centre de Recerca Ecològica i Aplicacions Forestals (CREAF), Barcelona, Catalonia, Spain

[2] Eidgenössische Technische Hochschule (ETH), Department of Materials, Zürich, Switzerland

[3] Cornell University, Materials Science and Engineering, New York, USA

[4] Yale University School of the Environment, New Haven, Connecticut, USA

10
**Authors for correspondence**: Kate M. Johnson, kate.johnson@utas.edu.au , Craig R. Brodersen,

email: craig.brodersen@yale.edu

**Emails :** Kate M. Johnson: kate.johnson@utas.edu.au Muriel Scherer: muriel.scherer@mat.ethz.ch,

Dominic Gerber: domi.gerber@gmx.ch , Robert W. Style: robert.style@mat.ethz.ch Eric R Dufresne:

erd43@cornell.edu Craig R. Brodersen:  craig.brodersen@yale.edu

**Running title:** The spread of ice and air in young *L.tulipifera* leaves

**Wordcount:** 4309

**Figures:** 9

**Tables:** 0

**Supplementary data:** 7 Figures, 3 videos

**Highlight**

Both ice and air lead likely lead to death in young *L.tulipfera* leaves exposed to freezing, with

the spread of both governed by physical characteristics of these leaves.

**Abstract**

Spring freezing is an unforgiving stress for young leaves, often leading to death, with

consequences for tree productivity and survival. While both the plant water transport system

and living tissues are vulnerable to freezing, we do not know whether damage to one or both

of these systems causes death in leaves exposed to freezing. Whole saplings of

*Liriodendron tulipifera* were exposed to freezing and thawing trajectories designed to mimic

natural spring freezes. We monitored the formation of freeze-thaw xylem embolism and

damage to photosynthetic tissues to reveal a predictable progression of ice formation across

the leaf surface that is strongly influenced by leaf vein architecture, notably the presence or

absence of bundle sheath extensions. Our data also show that freeze-thaw embolism occurs

only in the largest vein orders where mean vessel diameter exceeds 30μm. With evidence of

both freeze-thaw embolism and damage to photosynthetic tissue, we conclude that this dual-

mode lethality may be common among other wide-vesseled angiosperm-leaves, potentially

playing a role in limiting geographic distributions, and show that bundle sheath extensions

may stall or even prevent freezing spread.

**Key words:** Embolism, Fluorescence, Freezing, Freeze-thaw embolism, Leaves, Optical

imaging, Spring-freezing.

**Introduction**

Freezing is one of the strongest determinants of plant distributions (Burke, Gusta et al. 1976,

Loehle 1998, Sakai and Larcher 2012, Muffler, Beierkuhnlein et al. 2016), owing to the

unforgiving and often lethal nature of freezing in plant tissues that are not specifically

equipped to avoid, resist or compartmentalise ice crystal formation (Burke, Gusta et al. 1976,

Stegner, Buchner et al. 2023). Some trees have adaptations to prevent or tolerate freezing

within their tissues year-round, while others, such as those in temperate regions, are

adapted to seasonal winter-freezing (Burke, Gusta et al. 1976, Wisniewski and Fuller 1999,

Sakai and Larcher 2012). Both the xylem and living leaf tissue are critical to consider when

assessing the effects of freezing, as damage to plant water transport or the sites of

photosynthesis jeopardise tree growth and survival (Brodribb, Brodersen et al. 2021).

Within branches and tree-trunks, xylem anatomy is known to influence the likelihood of

freezing damage. During freezing, gas bubbles become suspended in the xylem sap as the

crystallization process pushes dissolved gasses out of solution. Upon thawing, these

bubbles either dissolve back into solution or expand to fill xylem conduits and block water

flow depending on the pressure of the surrounding liquid (Hammel 1967, Sperry and Sullivan

1992, Hacke and Sauter 1996). Species with larger diameter xylem conduits have been

shown to be more vulnerable to 'freeze-thaw embolism' (Sperry and Sullivan 1992, Feild and

Brodribb 2001, Tyree and Zimmermann 2002, Cavender-Bares, Cortes et al. 2005, Sevanto,

Holbrook et al. 2012, Li, Luo et al. 2024), which may, in turn, at least partially explain

geographic range distributions based on the likelihood of freezing conditions (Sperry 1995).

There are a number of possible explanations for this. Larger vessels in stems have been

shown to embolise earlier than smaller vessels exposed to freezing (Davis, Sperry et al.

1999, Cavender-Bares, Cortes et al. 2005), and smaller conduits have been shown, in some

cases, to freeze at  higher velocity which lead to smaller air bubbles (Ewers 1985) which are

more easily dissolved upon thawing (Sevanto, Holbrook et al. 2012).

Winter deciduousness allows many angiosperm trees (and some gymnosperms) to avoid

leaf-level freezing damage, but does not protect spring growth, nor leaves that persist into

autumn. In particular, freezes after bud-burst in spring can have severe consequences for

deciduous trees (Burke, Gusta et al. 1976, Vitasse, Bottero et al. 2019), including strong

reductions in tree growth (Vitasse, Bottero et al. 2019).

Intracellular ice formation invariably leads to cell death by rupture of the cell membranes,

caused by internal nucleation of piercing by external ice crystals (Mazur 1969, Burke, Gusta

et al. 1976, Steponkus, Dowgert et al. 1983, Guy 1990). Extracellular ice formation can also

result in cell death by creating a water potential gradient which draws out cellular water to

feed the freezing-front (Guy 1990, Steponkus and Webb 1992), therefore imposing a

desiccation stress resembling that caused by drought (Ruelland, Vaultier et al. 2009, Vitra,

Lenz et al. 2017, Yang, Gerber et al. 2024).

Spring frost may be lethal in young leaves through at least two different modes that are not

mutually exclusive. First, ice formation in or around the living tissues poses a significant

threat because of the potential damage to cell walls and membranes leading to electrolyte

leakage, damage to organelles and the generation of reactive oxygen species (Wise 1995,

Lütz 2010). Second, embolism formation resulting from the freezing and thawing of the

xylem sap could lead to complete failure of the leaf vein network thereby cutting off the

supply of water to downstream tissues (Brodribb et al. 2021). While the effects of freeze-

thaw embolism have been well documented in stems (Sperry, Donnelly et al. 1988, Feild and

Brodribb 2001, Ashworth and Pearce 2002, Cobb, Choat et al. 2007, Mayr, Schmid et al.

2014, Robinson, Rennie et al. 2023), this has never been visualised in leaves.

The primary means of tracking the propagation of freezing damage through whole leaves

has been with high temporal resolution using multiple forms of thermal imaging (Larcher,

Meindl et al. 1991, Gusta, Wisniewski et al. 2004, Hacker and Neuner 2007, Kokin, Pennar

et al. 2018, Zhang, Han et al. 2023). Hacker and Neuner (2007) in particular provide key,

novel information about ice nucleation and freezing relative to leaf anatomy, showing ice

nucleation beginning in the veins and spreading through the mesophyll tracking leaf

venation, however this is one of few papers to consider the influence of leaf venation on ice-

spread. While thermal imaging with very high temporal resolution (Hacker and Neuner 2007;

25 images per second) and some optical imaging (Kokin, Pennar et al. 2018) have been

used to study leaf freezing; longer-term high resolution imaging of leaves during freezing and

thawing has not been conducted. Given that ice crystallization is often lethal in living tissues

and that leaf cells are also supplied by a vascular system that is vulnerable to freeze-thaw

embolism, one of the goals of the present study was to provide a detailed characterisation of

ice formation in leaf tissue by visualising freezing and thawing in the leaves of intact plants

using a new application of an existing method, as used in (Kane and McAdam 2024).

Here we visualise the effects of freezing and thawing on both the living cells and the xylem in

young leaves in *Liriodendron tulipifera*, a common deciduous tree in North America. To

detect possible embolism and track ice crystal formation in leaves we used a combination of

time-lapse imaging and chlorophyll fluorescence imaging to non-invasively study young

leaves of *L. tulipifera* saplings exposed to freezing conditions, with minimum air-

temperatures ranging from -1 to -5 °C. By capturing images of newly expanded leaves at

regular intervals throughout freezing and thawing we aimed to elucidate the pattern of

freezing spread in these leaves and capture any embolism that may occur using the Optical

Vulnerability Method of Brodribb et al. (2017). We then measured chlorophyll fluorescence of

leaf tissue before and after freezing-and-thawing as a proxy for tissue damage (Brodribb et

al. 2021). To compare the effect of freeze-thaw embolism vs. drought-induced embolism,

validating the capacity of the cavicams to detect freeze embolism, and isolating how freezing

may differentially impact leaf tissue, we tracked drought-induced embolism in leaves of the

same age, sourced form adult trees at various locations.

*L. tulipifera* has been shown to possess bundle sheath extensions (BSEs), structures that

connect the bundle sheath and the epidermis and influence a range of physiological process

in plants relating to both photosynthesis and hydraulics (Wylie 1943, Wylie 1952, Pray 1954,

Esau 1960 ; Fig S1, Zwieniecki, Brodribb et al. 2007).  Bundle sheath extensions

compartmentalise the leaf lamina restricting the movement of gases therefore allowing gas

exchange to occur independently across the leaf surface (Neger 1918), and have been

shown to have numerous physiological and biophysical functions (Karabourniotis, Bornman

et al. 2000, Zwieniecki, Brodribb et al. 2007, Buckley, Sack et al. 2011, Trifiló, Raimondo et

al. 2016, Kawai, Miyoshi et al. 2017). Notably, highly sclerified BSEs have been shown to

prevent the lateral prorogation of ice in the lamina in *Cinnamomum camphora*, yet this is the

only species in which this has been documented (Hacker and Neuner 2007, Barbosa,

Chitwood et al. 2019).  In *L. tulipifera* BSEs extend to both the upper and lower epidermises

in second order and third order veins (Pray 1954; Fig S1), while they extend only to the

abaxial epidermis in higher order veins (Pray 1954, Pieruschka, Schurr et al. 2006, Barbosa,

Chitwood et al. 2019). It is possible that even the non-sclerified bundle-sheath extensions in

*L. tulipifera* could act as a barrier to the ice crystallization front as it progresses across the

lamina, which led us to attempt to determine their influence in this species.

Based on the known physical processes that govern ice nucleation and growth in laminar

systems that superficially resemble leaves, we expected ice to first form in leaf regions with

low solute content, and therefore a higher freezing temperature, such as the midveins

containing numerous xylem conduits loaded with xylem sap. Ice would then propagate

outward across the lamina as temperature decreases below the freezing point of the

surrounding tissues with higher solute content. Given that the BSEs represent a physical

barrier that would slow the movement of water toward the ice nucleation front, we

hypothesised that BSEs would significantly influence the spatial pattern and progression of

ice across the lamina, which would be most noticeable in the higher order veins where BSEs

span both the upper and lower epidermis leading to patchy ice-spread. We developed two

possible working hypotheses for freeze-thaw embolism: First, freeze-thaw embolism would

occur upon thawing in a pattern resembling embolism caused by drought, where the lower

vein orders embolize first, followed by the higher order veins. Second, embolism would occur

only in veins with large vessel diameters (equal to or > 30 μm) based on research in stems

which shows a steep increase in freeze thaw embolism in conduits above this threshold

(Pittermann and Sperry 2006). We predicted that ice crystal formation would lead to

damaged cell walls and membranes and thus irreparable damage to the mesophyll, made

visible by significant decreases in chlorophyll fluorescence shortly after thawing. If freeze-

thaw embolism occurred, it would lead to cell death in any remaining un-frozen tissue but

over a longer time period.

## Materials and Methods

This study comprised three core experiments, whereby we first exposed saplings to freezing

and thawing cycles while monitoring leaf vein embolism and chlorophyll fluorescence. We

then compared those patterns of freeze-thaw embolism to those produced with drought-

induced. Finally, we examined the freezing and thawing process at much higher

magnification to determine the underlying physical processes.

*Experiment 1: Freezing whole saplings*

Saplings of *Liriodendron tulipifera* (~ 2 years old) were grown under glasshouse conditions

(day temperature of ~25-30 °C and night temperatures of ~ 15-20 °C) for 1.5 months, until

there were at least two fully expanded leaves per tree, before experimentation (conducted

between March and May 2022). A total of 13 well-watered trees were frozen to one of five

temperature minima (-1, -2, -3, -4 or -5 °C). We selected this range of temperatures to both

determine the temperature needed to fully freeze leaves, but also to test the effects of cold

temperatures that did not induce freezing. We used the Optical Vulnerability Method (OVT)

of Brodribb et al. (2017) to monitor both the formation of embolism but also the changes in

optical density of the mesophyll associated with freezing. We imaged a total of 24 leaves,

each tree was frozen as described below.

Saplings growing in pots were placed in an upright freezer (Model: FFFH20F2QWC,

Electrolux, Stockholm, Sweden) which was attached to a thermostat (Model: ITC-310T-B,

Inkbird Tech, Shenzhen, China) which was used to control the freezing trajectory. Two

adjacent fully expanded leaves were placed inside 'cavicam' imaging clamps

(docs.cavicams.com, Fig. 1). Briefly, a digital Raspberry Pi camera (Model: V2, 9132664;

Raspberry Pi Ltd., Cambridge, UK) was positioned to observe the adaxial leaf surface and

transmitted light is provided by a digitally controlled set of LED lights that only illuminate

during image capture (Brodribb, Bienaime et al. 2016). Cameras were set to capture images

every 5 minutes during freezing and thawing, a process that was programmed to occur over

20 hours (see details below). These cameras had slightly different magnifications, allowing

freezing to be captured at two different scales, but the results were still comparable with no

significant difference found in the embolism of percentage of freezing detected between the

cameras (ANOVA p> 0.05). A custom-built temperature logger 'envirologger' run by a data-

logger (Adafruit feather M0 Adalogger; Adafruit Industries LLC; New York, USA) was placed

in the freezer to track the internal conditions with a temperature/humidity sensor (Model:

AM2315 12C, Adafruit Industries LLC).

The freezing sequence and trajectory were based on winter/spring temperature data from a

weather station (Model: CR300; Campbell Scientific, Utah, USA) monitoring windspeed, air

temperature, relative humidity, precipitation, soil moisture and soil temperature, averaged

hourly over the measurement period) at nearby Yale Myers Forest (Union, CT, USA; 41° 57'

8.2944'' N, 72° 7' 26.688'' W), where *L. tulipifera* trees are known to grow, in order to mimic

the freezing conditions experienced by this species under natural conditions. A

representative night in 2019 where temperatures reached -5 ˚C was chosen as a reference.

The temperature data from Yale Myers Forest (recorded half-hourly) was averaged and

rounded to the nearest whole number to determine mean hourly temperatures. These

temperatures where then used to inform a 12-step programmed freezing sequence. This

resulted in a stepped freezing sequence that ran for approximately 20 hours (including an ~

6 hour period at a positive temperature at the end) mimicking an overnight freezing event

(Fig S2).

**Figure 1**: The set-up for monitoring freeze in *L. tulipifera* trees. Panel (A) shows the sapling inside the

freezer with cameras attached. The unit for measuring freezer air temperature is denoted by a yellow

box while the temperature probe attached to the device controlling the freeze trajectory is circled in

yellow.  In panel (B) the two cameras are distinguished by boxes, with the higher magnification

camera in blue (C) and lower magnification camera in red (D). Corresponding images taken by each

camera are shown below the respective cameras, in boxes of the same colours (blue; higher

magnification and red; lower magnification) with a 5mm scale bar in the bottom right-hand of each

image.

*Imaging: Chlorophyll Fluorescence*

To determine the physiological effects of freezing temperatures on leaves we used

chlorophyll fluorescence imaging, which measures the status of the photosynthetic

apparatus and its capacity for processing light. We performed dark-adapted Fv/Fm imaging

on leaves in the laboratory both before freezing and thawing to provide a baseline reference

measurement, and then again after freezing and thawing to determine how much the

photosystems were damaged (Mini Imaging-PAM M-series, Walz GmbH, Germany). Before

fluorescence measurements, a small piece of opaque tape was applied to a small section (~

0.5x 0.5 cm) of the leaf, then the leaves were wrapped in aluminium foil to exclude light. The

patch of opaque tape was used because initial testing showed that, as removal of aluminium

foil was required to align the leaf under the imaging PAM, an initial measure was needed to

maintain some leaf tissue in the dark-adapted state required for Fv/Fm measurements. The

tape was removed seconds before measurement to ensure that the leaf tissue beneath the

tape remained fully dark adapted until fluorescence was measured.

*Imaging: Freezing and thawing in the leaf mesophyll*

Image sequences were processed in FIJI software (Schindelin, Arganda-Carreras et al.

2012) to determine the spread and extent of ice crystallization during the experiments. The

tracing tool in FIJI was used to quantify the total leaf area frozen at each time step, which

was indicated by a change in the colour intensity of the pixels. Freezing was detected as

patches of brighter pixels in the leaf lamina, typically bounded by higher order veins.

To quantify the colour change in both mesophyll tissue that did and did not freeze, circular

Regions of Interest (ROIs) of 40x40 pixels were selected in regions of both mesophyll that

froze and that which did not in all leaves frozen to -4 ˚C. The mean grey scale value in these

ROIs was calculated at three time-points, at the start of image capture, at -4 ˚C (at freezing)

and immediately after thawing. The values at -4 ˚C and after thawing were expressed as a

percentage of the initial mean greyscale value (from the start of image capture) and the

average percentage of the initial value was calculated for both frozen and unfrozen

mesophyll. Mean Fv/Fm for unfrozen, frozen and thawed tissues was calculated from

measurements taken across all leaves that froze.

A similar method, utilising circular ROIs in FIJI, was used to track the change in mean
greyscale area in the midveins and mesophyll of all leaves which experienced freezing. In
each leaf, ROIs which fit within the diameter of the midvein were selected and added to the
ROI manager. This was repeated across the image stack or up to 250 images (where more
images were recorded), with the ROI shifted using 'edit>selection>specify' when the leaf
moved. The 'Interpolate ROI's' option through the ROI manager was then used to interpolate
between the selected ROIs for all images. The 'measure' function was then used to calculate
the mean greyscale area in the selected ROI for each image. This was then repeated for a
section of mesophyll adjacent to the midvein, using an ROI of the same area. The mean
greyscale values across the images were expressed as a percentage of the initial (unfrozen)
mean greyscale value in order to generate lines plots of changes in mean grey scale value
before, during and after freezing.

The freezing extent was calculated in the image where the maximum amount of tissue was
frozen, which corresponded with the minimum air temperature to which the leaves were
exposed. The frozen regions were selected using the tracing tool in FIJI (Schindelin,
Arganda-Carreras et al. 2012) and expressed as percentage of the total visible leaf area
within the field of view to calculate a percentage.

*Imaging: Embolism upon thawing*

We monitored the leaf vein network for the formation of embolism using the OVT (Brodribb,
Bienaime et al. 2016). Embolism was monitored by the cameras for 12-24 hours post
thawing to ensure that all embolism was captured. Freeze-thaw embolism was analysed
using the same methodology which is used to highlight drought-induced embolism with leaf
images analysed via an image subtraction, where each image is subtracted from the one
before to reveal the changes between the images. As there is a change in light transmission
as air rapidly replaces water in the xylem due to embolism, the image subtraction reveals
these changes which can then be visualised through time. A detailed description of the
methodology used to process images can be found here: https://www.opensourceov.org/.
While embolism was expected to occur upon thawing, cameras remained attached to plant
organs for ~6 hours to ensure that all embolism was captured.

*Experiment 2: Drought embolism in additional leaves*

Optical analysis of drought-induced embolism was undertaken in a total of six leaves from
branches of adult trees, one leaf was sourced from a tree located on the Yale University
campus, New Haven, Connecticut, USA (41°19'16.6"N 72°55'25.8"W), and the remaining
five leaves were sampled form across four trees located in North Hobart, Tasmania,
Australia (42°52'07.6"S 147°18'57.8"E). Young, fully expanded leaves (as a similar

developmental stage as those used in the freezing experiment) were chosen for imaging.

One leaf from each branch was placed within an imaging clamp and set to take images at 5-

minute intervals as the branches dried. Embolism was determined to have finished after 12

hours without embolism.  Images were analysed using an image subtraction, as described

above.

*Analysis: Embolism in Experiment 1 and Experiment 2*

The timing and extent of embolism were determined both in leaves exposed to freezing and

those exposed to drought. To visualise embolism through time, embolism was colour coded

according to time in both a leaf exposed to freeze and a leaf exposed to drought. This was

done using  FIJI (Schindelin, Arganda-Carreras et al. 2012) using the 'colour slices' function

in the 'OSOV toolbox', the toolbox and instructions are available here:

https://www.opensourceov.org/.

*Analysis: Embolism and vein order*

To determine how much the vein network embolised, the total length of the visible venation

network was calculated along with the total area of the embolised vein network using the

'segmented line' tool in image J and the embolised vein area was then expressed as a

percentage of the vein network. Total vein area was calculated from a raw image of each

leaf taken from the stack of images which captured during the drought or freezing treatment

while total embolised vein area was calculated form the 'Z- stack' showing all the embolism

which occurred during the treatment. Total embolised vein area was then expressed as a

percentage of total vein area visible with the optical cameras. As embolism was only

observed in the mid-rib and major veins in response to freeze, only these vein orders were

included in calculations of vein length and embolised vein length for frozen leaves. The raw

images of the vein network and Z-stacks of embolism from leaves exposed to drought were

segmented into four even quadrants using a macro in FIJI and the top right corner was

analysed for each leaf in order to provide a random sample of the vein network.

In *L.tulipifera* leaves, the midvein, second order and tertiary veins are deemed the 'major

veins' while the fourth, fifth and sixth and seventh order veins are considered the 'minor'

veins and make up the most of the total vein length (Pray 1954).

To quantify the contribution of the midvein and second order veins to the earliest embolism

is leaves exposed to drought, 25% (embolised area) of the total drought-induced embolism

was chosen as a reference value. This value was used to determine whether embolism

observed due to freezing and thawing matched early embolism during drought.  At this

percentage embolism, in all droughted leaves, we calculated both the area contribution of

these veins compared to higher vein orders, and the total length of midvein and second

order veins embolised as a percentage of the total embolised area of these veins at 100%

embolism. For area, the 'freehand sections' tool in FIJI was used to 'circle' all vein orders

expect for the midvein and second order veins and the area of these veins was calculated

using the 'measure' function and expressed as a percentage of the total cumulative

embolised area at 25% embolism. The resulting percentage of higher-order-veins was

subtracted from 100% to reveal the percentage contribution of the midvein and second order

veins to the embolised area at 25 % embolism. For vein length, the 'segmented line' tool was

used to calculate the length of the total embolised area of the midvein and second order

veins in a Z-stack of the cumulative embolism at 25% embolism. This was expressed as a

percentage total embolised midvein and second order vein length at 100% embolism.

## *Experiment 3: High resolution imaging of freezing*

Branches of adult *L. tulipifera* trees were excised from individuals on the EPFL campus,

Lausanne, Switzerland (46.5191° N, 6.5668° E) and transported to the ETH Honggerberg

Campus in Zurich. Small leaves were mounted between two glass microscope slides and

placed on a customised freezing stage beneath a compound microscope using confocal

microscopy (Yokogawa Spinning Disk and Nikon Eclipse, set up &software by 3i imaging).

The set-up used is described in (Gerber, Wilen et al. 2023) and the documentation is

available at: https://github.com/dogerber/temperature_gradient_microscopy_stage/tree/main.

A CF640 filter was used to visualise eaves, capturing the autofluorescence of chlorophyll

therefore highlighting the cells in the images.

## *Anatomical assessment: Xylem conduit diameter analysis*

Young *L. tulipifera* leaves were collected from trees and brought to the lab. Vein samples

from three leaves were used for the conduit diameter analysis. Midvein samples were

collected 1cm from the petiole lamina junction. Second and third order veins were collected

0.5 cm from the junction with the subtending lower order vein. Samples were mounted to a

freezing stage (BFS-3MP, Physitemp Instruments, USA) on a sliding microtome encased in

a dilute glucose solution. Thin sections were stained with a 0.5% toludine blue solution,

rinsed in DI water, and then mounted in water. Images were captured with a digital camera

mounted to a compound light microscope (BX43, Olympus Inc, Japan) using a 4x or 10x

objective lens. Images were then used to measure xylem vessel diameter in FIJI. Vessel

diameter was measured as the longest distance across the lumen.

*Data analysis*

Data were plotted and analysed using SIGMAPLOT v.12.5 (Systat Software Inc, San Jose,

California) and RStudio was used to perform Single factor ANOVAs to determine statistical

differences between groups in RStudio (2016). Xylem vessel diameter analysis was

performed in RStudio using the dplyr package (Yarberry and Yarberry 2021).

## Results

386 *Pattern of Freezing and thawing*

387 We observed mesophyll freezing in all leaves which were exposed to minimum air

388 temperatures of -4° C or -5° (total of 18 leaves). At -4° C greyscale pixel values increased in

389 the freezing regions that can be interpreted as a change from air to water in the intercellular

390 space as water is drawn out of the mesophyll cells to feed the freezing front (Fig. 2).

**Figure 2:** The average percentage of initial (pre-treatment) mean greyscale value (brightness) for all

18 leaves that froze, comparing tissue that did and did not freeze within the same leaves. These

values were calculated at -4 °C and immediately after thawing. Unfrozen tissue is shown in black,

while frozen tissue in shown in white. This was calculated for all 18 leaves that froze. Asterix's

indicate a statistically significant difference (ANOVA, P<0.05). As these values were calculated as a

percentage of the mean greyscale value at the first image in the image sequence, some values were

< 100% due to small changes in lighting during image capture.

While an increase in mean greyscale value, indicating freezing, was always detected in the

mesophyll, this was not consistently observed in the midvein (Fig 3). Freezing in the

mesophyll resulted in a 10-20% increase in mean greyscale brightness while freezing in the

veins incurred an increase in mean greyscale value of <5% (Fig 3). Only one of the eight

leaves frozen to -4°C showed a signal of freezing in the midvein, while six of the ten leaves

frozen to -5°C showed this signal of freezing.

**Figure 3:** Percentage of initial greyscale value over time for the midvein and mesophyll for 18 *L.*

*tulipifera* leaves frozen to -4°C or -5°C. The mean greyscale values for leaf midveins (blue) and

mesophyll adjacent to the midvein (orange) across 250 images (~20 hours) for all leaves frozen to -

4°C or -5°C (A). Increases in mean greyscale brightness in the mesophyll (orange) indicate mesophyll

freezing. Panel (B) shows the midvein and mesophyll mean greyscale traces for a representative

single leaf frozen to -4°C and panel (C) shows the same for a leaf frozen to -5°C with a black arrow

denoting the increase in mean greyscale brightness indicating freezing in the midvein. Panels (B) and

(C) serve to highlight two examples and do not describe the pattern observed in all leaves frozen to -

4°C and -5°C respectively. The signal and timing of mean greyscale increases varied across of leave

independent of temperature.

Freezing was first detected in mesophyll immediately adjacent to the midvein and 2° veins,

followed by ice propagation away from the vein into areoles bounded by higher order vines

(3-5° veins) (Fig. 4). Freezing began at ~ -4°C continuing until 0°C when thawing occurred

(Fig. 4). Freezing advanced across the leaf surface in a stepwise pattern between adjacent

areoles in discrete patches. Freezing was in some cases not perfectly spatially homogenous,

and in all but two leaves where 100% of the area frozen (16 out of 18 leaves) some areoles

bounded by third order veins remained unfrozen (Fig. 4, Vid. 1, Fig. S3).

**Figure 4:** Representative freezing and thawing sequence including images, mean greyscale value
and air temperature for a representative *L. tulipifera* leaf visualised using time-lapse imaging. Panel
(A) shows greyscale images of the leaf surface including the midvein and surrounding higher order
veins with the transect used to calculates mean greyscale values denotes by a yellow rectangle.
Mean greyscale values (B) across a 7mm mean greyscale profile between the midvein and a major
vein of *L. tulipifera* leaves measured at 5 stages of freezing in a leaf frozen to -5°C [pre-freeze (1):
black, freezing; blue (2) and yellow (3), peak freeze: light orange (4), thawed; dark orange (5)]. The
temperature at each stage is in the bottom right-hand corner of each image and panel (6) shows time
(minutes) vs temperature (°C). This figure shows the distinctive colour change which occurs with
freezing visually (A) and numerically (B), illustrating a situation where a patch in the centre of the
mean greyscale profile remains unfrozen (denoted by a black arrow in the 'Peak freeze' image in
panel (B).

Freezing at high magnification in young leaves from adult trees revealed a similar pattern,
where a freezing-front propagated through individual areoles, but then stalled upon reaching

a third-order vein boundary (Video 2). The freezing front can be observed progressing from

the bottom left corner towards the middle of the field of view where it can be seen moving

through cells before reaching a third-order vein boundary and then progressing to the bottom

right and moving towards the middle before reaching the vein boundary in the same manner

(Video 1).

Thawing, by contrast, did not appear to occur as a staged process, occurring over 5-10

minutes images when the temperature rose above 0 °C, Fig. 4). Thawing resulted in a

'darkening' of the previously lighter frozen patches under the optical cameras which used

transmitted light (Fig. 2, Fig. 4), while thawing corresponded with lightening in the high-

resolution video of thawing as this set-up utilised reflected light (Video 3). In the hours

following thawing, previously frozen tissue gained a red colouration (Fig. 5).

**Figure 5:** L. *tulipifera* leaves (n=8) at their maximum freezing extent (left hand column) and ~ 10
hours after thawing (right hand column), demonstrating the reddening of the tissue that occurred in
frozen tissue. Image pairs (A, B) and (C, D) show freezing and reddening I leaves frozen to an air
temperature of -4° C, while (E, F) and (G, H) show this in leaves frozen to -5 ° C (c). The 5mm scale
bar in (a) is true for all panels.

*Freezing extent*

Freezing began at air temperatures below -4 °C, with no visible evidence of freezing in
leaves exposed to air temperatures of -1°C, -2°C and -3°C. In leaves exposed to air
temperatures below -4°C (n = 8), an average of 58%± 5% of the leaf froze within the
camera's field of view. The mean percentage of the leaf turning red increased to 93% ±2% in
the ten leaves frozen to -5°C (; P< 0.000008, ANOVA; Fig. 6).

**Figure 6:** The percentage frozen leaf area in leaves frozen to air temperatures of -4°C and - 5°C.
panel (A) shows to total percentage frozen leaf tissue in all leaves frozen to -4°C (gold) and -5°C
(blue). Representative images of the freezing extent are also shown in both a leaf frozen to -4°C (B)
and a leaf frozen to -5°C (C). Arrows mark unfrozen tissue (black arrows) and frozen tissue (white
arrows) in both (B) and (C). The 5mm scale bar in (C) is true for (B) and (C) panels.

*Chlorophyll Fluorescence Imaging*

Chlorophyll fluorescence measured ~6 hours after freezing and thawing invariably
decreased in frozen tissue and largely remained high (close to 0.8 Fv/Fm) within leaves that
did not freeze and in unfrozen tissue within leaves that did freeze (Fig. 7). Mean Fv/Fm for

frozen tissue (0.349 ± 0.06) was significantly lower than that for unfrozen tissue (0.733 ±

0.21; P< 0.000003, ANOVA). In leaves exposed to -1°C -2°C or -3°C, where we observed no

freezing of the lamina, fluorescence remained > 0.7 Fv/Fm (Fig S4). Evidence of both Fv/Fm

depression and high Fv/Fm values in frozen and unfrozen tissue, respectively, were

measured in leaves exposed to temperatures of -4°C or -5°C. The 'patchy' nature of freezing

in the leaf mesophyll visible with the optical camera (Fig.6) was also visible through

chlorophyll fluorescence imaging (Figs. 7b, S5). A lack of fluorescence decline in leaves

where freezing was not observed, (exposed to -1°C, -2°C and -3°C air temperatures)

indicates that fluorescence values were not impacted by the time spent in the freezer or the

placement of cameras on leaves (Fig. S4).

**Figure 7**: Changes in chlorophyll fluorescence in response to negative temperatures in 24 leaves of
*L. tulipifera*. Panel (A) shows  Fv/Fm (photosynthetic capacity) in frozen (orange) and unfrozen (blue)
leave tissue against the minimum air temperature leaves were frozen to, along with a visual
representation of fluorescence in a leaf post-freeze (B).The dark blue rectangle in (B) shows the patch
of unfrozen tissue which was 'dark-adapted' before the fluorescence measurement, with a value of
0.715 Fv/Fm. Unfrozen tissues outside of this dark-adapted area are turquoise and light green,
indicating a higher fluorescence than the surrounding, darker unfrozen tissue. Point values of Fv/fm

are shown in eight circular areas of interest arranged diagonally across the image. The colour-scale

beneath panel (B) indicates fluorescence values.

*Tree Recovery*

While leaves exposed to air temperatures of -4°C or -5°C invariably died, all but one of the

13 trees exposed to these temperatures resprouted leaves (Fig. S7). Resprouting occurred

from dormant buds below the leaves frozen during the experiment (Fig. S7).

*Freeze-thaw vs. drought-induced embolism*

Freeze-thaw embolism was seen in six of the eight leaves frozen to -4°C, but not in any of

the leaves frozen to -5°C.  Embolism after freezing occurred in 1-3 events and were isolated

to the midvein and 2˚ veins (Fig. S6; Fig.8). We observed no freeze-thaw embolism in the

higher order veins, which was distinctly different than the pattern of embolism spread

resulting from drought (Fig. 8). Importantly, the clear visibility of embolism in higher order

veins in leaves exposed to drought means that the lack of freeze-thaw embolism in higher

order veins was not a failure of the optical cameras to detect these events.

Freeze-thaw embolism was confined to the midvein and second order veins, while drought

induced embolism was observed in all vein orders (Fig. 8). In leaves where freeze-thaw

embolism was observed, 77.6 ± 11.8 % of the midvein and second order veins embolised,

compared to 100% of these veins in drought-exposed leaves. In response to drought, the

percentage of embolism in tertiary veins and above was a similar to the percentage of

freeze-thaw embolised midveins and major veins 74.25 ± 7 %, including all vein orders in

drought-exposed leaves (adding major and second order veins) did not alter the percentage

of the vein network that was embolised substantially, with 78%± 1.4% of the total vein

network found to embolise in leaves exposed to drought. The midvein and second order

veins often embolised first in response to drought embolism (representing 83.56% ± 4% of

the total embolised area at $P_{25}$ and 76% ± 34% of the total embolised length of midveins and

second order veins at $P_{25}$ as a percentage of the total length at $P_{100}$). However, embolism in

these veins was accompanied, or shortly followed, by embolism in the minor veins as well

(Fig. 8) unlike in leaves exposed to freezing and thawing. The timescale across which

drought embolism and freeze thaw-embolised were also very different. While freeze-thaw

embolism events occurred within 5- 15 minutes, drought induced embolism occurred over a

range of 1- 5 days (Fig. 8).

**Figure 8:** Representative image sequences showing the difference between drought- and freeze-thaw-induced embolism in *L. tulipifera* leaves. Drought-induced embolism spread was detected in young leaves of adult trees and occurred over approximately 1 day, 25.42 hours (A, B, C). Freeze-thaw embolism detected in whole saplings occurred over 15 minutes (D,E,F) in *L. tulipifera* leaves. Panels a, b and c show the progression of drought-induced embolism in four steps of ~23.5 hours each, while the three images in panel (B) show the progression of freeze-thaw embolism in three steps of 5 minutes, where thawing began at 0 minutes. Embolisms are colour coded according to when they occurred, as indicated by the colour scale beneath the panels.

*Xylem vessel diameter*

We measured vessel lumen diameter distributions in the three lowest vein orders to determine whether this trait might influence our observations of embolism in this species given that previous work indicates that freeze-thaw embolism vulnerability increases when mean conduit diameter exceeds 30μm (Pittermann and Sperry 2003). We found a mean vessel diameter of 34.4± 6.96, 27.4± 6.95, and 7.45 ±1.85 for the midveins, second order veins, and third order veins, respectively (Fig. 9). We found that 72% and 33.3% of the vessels were greater than 30μm in the midveins and second order veins, respectively, but no vessels >30μm were found in high order veins.

**Figure 9:** Xylem vessel lumen diameter distributions for leaves of *L. tulipifera*. Distributions are shows for different vein orders (blue = midveins, green = second order veins, orange = high order veins). n = 3, mean diameter is statistically different for each group (ANOVA, p<0.05). Vertical dashed line shows the 30µm threshold, where conduits below this threshold are known to be less vulnerable to freeze-thaw embolism.

## Discussion

Freezing young *L. tulipifera* trees to air temperatures of -4 °C or -5 °C invariably led to visible freezing in the mesophyll, cell damage after thawing, and ultimately leaf death. Our visualisations of ice formation revealed that mesophyll freezing began adjacent to the midveins and second order veins before progressing in a step-wise pattern across areoles bounded by third order veins. These freezing patterns support our hypothesis that bundle-sheath extensions (BSEs) strongly influence how ice propagates and spreads through the lamina of *L.tulipifera* leaves. Freeze-thaw embolism occurred only in veins containing vessels with diameters 30 µm or higher, in contrast to drought-induced embolism which occurred across all vein orders. This finding supports our second hypothesis that freeze-thaw embolism is driven by conduit size, generally agreeing with research in stems (Pittermann and Sperry 2003), providing further support that conduit diameter is a major driver of freeze-thaw embolism across multiple plant organs.

*Mesophyll freezing and thawing*

Mesophyll freezing was associated with an increase in tissue brightness, which was also

recently shown by Kane and McAdam (2024). This increase in brightness can be explained

by the transition in optical density of the intercellular airspaces as the air is replaced with

liquid and then solid water during the freezing process. This is the same principle that

underpins the Optical Vulnerability Technique (OVT) for the observation of drought-induced

embolism (Brodribb, Bienaime et al. 2016, Brodribb, Carriqui et al. 2017). The OVT detects

embolism in leaves as a colour-change from light-coloured translucent xylem (water-filled) to

darker air-filled xylem, due to the change in optical density of the conduits (Brodribb, Carriqui

et al. 2017). Therefore, the transition from darker to lighter tissue at freezing temperatures

(Kane and McAdam 2024) can likely be explained by the movement of water/ ice into

previously air-filled spaces. As the refractive indexes of liquid water (1.33) and ice (1.31) are

very similar, the flooded intercellular space would appear brighter even after thawing, as we

saw here (Fig. 4). This colour change was reversed in the high-resolution freezing, whereby

freezing caused a darkening of tissue (Video 2). The occurrence of intercellular freezing is in

agreement with the idea of freezing as a dehydration stress in plants (Guy 1990, Steponkus

and Webb 1992, Ruelland, Vaultier et al. 2009, Vitra, Lenz et al. 2017, Yang, Gerber et al.

2024) whereby water is pulled out of living cells to feed the freezing front in the spaces

between cells. In the absence of intracellular freezing, this process is thought to cause tissue

death (Steponkus and Webb 1992).

We attribute the difference in the wave-like spread of freezing from around the midvein and

second order veins and the stepwise, 'patchy' freezing that occurred in mesophyll bound by

tertiary veins (Videos 1, 2) to the presence or absence of bundle-sheath extensions (BSEs).

previously published work shows that freezing initiated close to the midvein and second

order veins (Hacker and Neuner 2007). BSEs extend to both the upper and lower epidermis

in tertiary veins of *L. tulipifera* (Fig. S1), but are not present in the midvein or second order

veins. BSEs have been shown to be discontinuous or extend to only one leaf surface in

fourth order veins and higher in *L. tulipifera* (Pray 1954). The absence of BSEs in the two

lowest vein orders likely accounts for the apparently uninhibited spread of ice adjacent to the

midvein and second order veins while their presence may explain the independent and

patchy freezing or areoles bounded by tertiary veins containing BSEs (Fig S3). The

presence or absence of BSEs can therefore be used to explain the delay to freezing spread

at tertiary vein boundaries (Videos 1, 2), or complete blockage in others (Figs 4, 6, 7), where

BSEs are acting as physical barriers to the movement of the ice crystallisation.

*Freeze-thaw xylem embolism*

The most convincing driver of freeze-thaw embolism here may be xylem conduit diameter.
Freeze-thaw embolism in the lamina was confined to the two lowest vein orders (midvein
and second order veins), which also possess the largest vessels (Figs. 8, 9). We found
vessel lumen diameters exceeding 30μm in both vein orders showing embolism. In contrast,
we observed no embolism in veins with vessels <15μm in diameter veins. This observation
aligns with research in conifer stems which shows a sharp increase in vulnerability to freeze-
thaw embolism when conduit diameter exceeds 30μm (Pittermann and Sperry 2003) and is
consistent with the large body of research in woody plants which shows that larger diameter
conduits are at greater risk (Sperry and Sullivan 1992, Sperry, Nichols et al. 1994, Feild and
Brodribb 2001, Tyree and Zimmermann 2002, Pittermann and Sperry 2006, Choat, Medek et
al. 2011, Robinson, Rennie et al. 2023). This has been theorised (Sevanto, Holbrook et al.
2012) and been inferred to be the case in leaves (Ball, Canny et al. 2006) but has, until now,
not been visualised.

A reduction in xylem conduit diameter was one of the key adaptations that allowed flowering
trees to radiate into freezing environments (Zanne, Tank et al. 2014) and decreasing conduit
diameter with increasing probability of freezing temperatures has been observed at a global
scale (Sperry 1995, Sevanto, Holbrook et al. 2012). With a large population of vessels
<30μm, it is possible that if embolism of the lower vein orders was incomplete, the smaller
conduits which are theoretically less vulnerable to freeze-thaw embolism could maintain
water supply to these leaves. This may allow *L. tulipifera* leaves to resist freezing
temperatures provided that photosynthetic tissues are 'hardened' to freezing temperatures.
The extent of freezing damage in photosynthetic tissues may, however, outweigh any effects
of embolism, at least early in the growing season when leaves are not acclimated for cold
temperatures.

A mean greyscale signal of freezing in the midvein was observed in six out of the 10 leaves
frozen to -5°C but only one out of the eight frozen to -4°C. This is contrary to evidence from
thermal imaging which showed that freezing occurs first in the veins (Hacker and Neuner
2007). It is possible that vein freezing is stochastic, occurring base on spontaneous ice
nucleation which does not always occur. This may be explained by the nucleation by
substances other than water such as macromolecules or dust, termed 'heterogenous
nucleation'(Kanji, Ladino et al. 2017, Shardt, Isenrich et al. 2022). Another possibility is that
the inconsistency in the detection of freezing in the midvein may be related to the already
high transparency of the leaf venation in optical images making it difficult to detect the subtle
change in the reflective index when water freezes.

A notable factor that we did not test in this study is the speed of freezing and thawing. We

opted for a slow freeze-thaw trajectory which mimics field conditions but did not test different

speeds of freezing or thawing. Slower freezes are  thought to increase the likelihood of

freeze-thaw embolism (Sevanto, Holbrook et al. 2012) but slower thawing is thought the

decrease the loss of conductivity in the xylem (Langan, Ewers et al. 1997). Given the large

increase in percentage frozen area between -4°C and -5°C degrees it is possible that that

rapidity of freezing over this one degree temperature range precluded embolism formation

and that a slower freeze or thaw may have produced different results.

*Freeze-thaw vs. drought induced xylem embolism*

Differences in the mechanism behind embolism induced by drought vs. freeze-thaw

embolism in the leaf lamina likely explains the contrasting extent of embolism observed in *L.*

*tulipifera* leaves exposed to freezing and drought. The vulnerability of xylem conduits to air-

seeding under drought conditions has been linked to anatomical characteristics such as the

thickness of the pit membranes in the xylem wall (the site at which air enters, causing

embolism due to drought;Thonglim, Delzon et al. 2021, Thonglim, Bortolami et al. 2023) and

connectivity of the xylem network (Brodersen and Roddy 2016, Johnson, Brodersen et al.

2020, Mrad, Johnson et al. 2020). Combined, these anatomical features likely explain the

pattern of drought-induced embolism we observed in young *L. tulipifera* leaves whereby

embolism progressed through the major vein orders terminating in the higher order veins, a

pattern which is consistently observed across the leaves of woody and herbaceous

angiosperms (Johnson, Jordan et al. 2018, Tonet, Carins-Murphy et al. 2023). In contrast,

freeze-thaw embolism, likely caused by gas segregation and subsequent bubble expansion

(Sevanto, Holbrook et al. 2012), is thought to be strongly influenced by xylem conduit

diameter that determines the size of the resulting bubbles (Pittermann and Sperry 2006).

*What kills* L. *tulipifera leaves during a freeze?*

The freeze-induced embolism and damage to living cells detected here highlights that

freezing can damage both the water transport and photosynthetic systems in young *L.*

*tulipifera* leaves. It should also be noted however, that regardless of whether embolism

occurred, all leaves which showed evidence of mesophyll freezing subsequently died. Cell

damage and death was evident in all leaves exposed to temperatures below -4°C, clearly

evident in leaf redness immediately following thawing and then subsequent tissue browning

in the following days (Fig. S7). This lethal damage was also evident in the depression of

chlorophyll fluorescence (Fig. 7). Additionally, *L. tulipifera* leaves presented 'wilted and wet'

after thawing, a state described by Burke and Gusta et al (1976) in observations of young

spring leaves after frost events, attributed to the loss of semi permeability in the cell

membranes. The absence of embolism in the lamina of leaves frozen to -5°C indicates that
embolism in the lamina is not necessary to induce leaf-death.

## Conclusions

Freezing invariably killed *L. tulipifera* leaves. Death of living tissue was a constant,
sometimes accompanied by freeze-thaw embolism which occurred rapidly and was confined
to the veins with the largest conduits, in stark contrast to drought-induced embolism. This
supports research in stems showing that conduit diameter drives freeze-thaw embolism
vulnerability. The progression of freezing through the mesophyll in patches bounded by
tertiary veins implicates BSEs in driving lateral ice propagation. While freezing in the
mesophyll conclusively led to death in *L. tulipifera* leaves, a significant proportion of xylem
vessels were inherently vulnerable to freeze-thaw embolism, showing that spring frost
results in a dual mode of lethality in *L. tulipifera* leaves.

## Supplementary Data

**Figure S1:** Transverse light microscope section of the lamina of a *Liriodendron tulipifera* leaf
highlighting a bundle-sheath extension.

**Figure S2:** Comparison of natural and experimental freezing trajectories.

**Video 1:** The pattern of freezing and thawing detected in a leaf frozen to -5 ° C with time-lapse
imaging.

**Figure S3:** Examples of the 'patchy' nature of freezing observed in *Liriodendron tulipifera* leaves
frozen to air temperatures of both -4 ° C and -5 ° C.

**Figure S4:** Chlorophyll fluorescence (Fv/Fm) in *Liriodendron tulipifera* leaves which were placed in
the freezer but did not show visible signs of freezing.

**Figure S5:** Comparison of optically resolved freezing and cell damage shown through imaging
fluorescence in *Liriodendron tulipifera leaves*.

**Figure S6:** The total embolism overlayed onto initial images of each the six *Liriodendron tulipifera*
leaves in which embolism was observed.

**Video 2**: High resolution freezing in a *Liriodendron tulipifera* leaf.

**Video 3**: High resolution GIF of thawing of a *Liriodendron tulipifera* leaf.

**Figure S7:** *Liriodendron tulipifera* trees shortly after they were exposed to -4°C or -5°C freeze
treatments.

## Acknowledgements

We would like to thank Peter Johnson for his assistance in designing the 'Envirologger' monitoring systems.

## Author contributions

KMJ and CRB conceived and designed the study. KMJ and CRB conducted experiments and measurements using optical cameras and microscopy, while KMJ and MS designed and conducted the collection of high-resolution imagery with guidance from DG, RWS and ERD. KMJ and CRB wrote the manuscript with contributions from all authors.

## Conflict of Interest

None declared

## Funding Statement

KMJ was supported by a Fulbright Future Postdoctoral Fellowship Awarded by the Australian Fulbright Foundation and the Kinghorn Foundation. RWS, MS and DG were supported by Swiss National Science Foundation grant: 200021-212066.

## Data availability

The data will be made available by the corresponding authors upon reasonable request.

## References

(2016). R: A language and environment for statistical computing. R Foundation for Statistical Computing Vienna, Austria R Core Team.

Ashworth, E. N. and R. S. Pearce (2002). "Extracellular freezing in leaves of freezing-sensitive species." Planta **214**(5): 798-805.

Ball, M. C., M. J. Canny, C. X. Huang, J. J. G. Egerton and J. Wolfe (2006). "Freeze/thaw-induced embolism depends on nadir temperature: the heterogeneous hydration hypothesis." Plant, Cell & Environment **29**(5): 729-745.

Barbosa, M. A. M., D. H. Chitwood, A. A. Azevedo, W. L. Araújo, D. M. Ribeiro, L. E. P. Peres, S. C. V. Martins and A. Zsögön (2019). "Bundle sheath extensions affect leaf structural and physiological plasticity in response to irradiance." Plant, Cell & Environment **42**(5): 1575-1589.

Brodersen, C. R. and A. B. Roddy (2016). "New frontiers in the three-dimensional visualization of plant structure and function." Am J Bot **103**(2): 184-188.

Brodribb, T., C. R. Brodersen, M. Carriqui, V. Tonet, C. Rodriguez Dominguez and S. McAdam (2021). "Linking xylem network failure with leaf tissue death." New Phytologist **232**: 68-79.

Brodribb, T. J., D. Bienaime and P. Marmottant (2016). "Revealing catastrophic failure of leaf networks under stress." PNAS **113**(17): 4865-4869.

Brodribb, T. J., M. Carriqui, S. Delzon and C. Lucani (2017). "Optical Measurement of Stem Xylem Vulnerability." Plant Physiol **174**(4): 2054-2061.

Buckley, T. N., L. Sack and M. E. Gilbert (2011). "The Role of Bundle Sheath Extensions and Life Form
in Stomatal Responses to Leaf Water Status   " <u>Plant Physiology</u> **156**(2): 962-973.
Burke, M., L. Gusta, H. Quamme, C. Weiser and P. Li (1976). "Freezing and injury in plants." <u>Annual</u>
<u>Review of Plant Physiology</u> **27**(1): 507-528.
Cavender-Bares, J., P. Cortes, S. Rambal, R. Joffre, B. Miles and A. Rocheteau (2005). "Summer and
winter sensitivity of leaves and xylem to minimum freezing temperatures: a comparison of co-
occurring Mediterranean oaks that differ in leaf lifespan." <u>New Phytologist</u> **168**(3): 597-612.
Choat, B., D. E. Medek, S. A. Stuart, J. Pasquet-Kok, J. J. G. Egerton, H. Salari, L. Sack and M. C. Ball
(2011). "Xylem traits mediate a trade-off between resistance to freeze–thaw-induced embolism and
photosynthetic capacity in overwintering evergreens." <u>New Phytologist</u> **191**(4): 996-1005.
Cobb, A. R., B. Choat and N. M. Holbrook (2007). "Dynamics of freeze–thaw embolism in Smilax
rotundifolia (Smilacaceae)." <u>American Journal of Botany</u> **94**(4): 640-649.
Davis, S. D., J. S. Sperry and U. G. Hacke (1999). "The relationship between xylem conduit diameter
and cavitation caused by freezing." <u>American Journal of Botany</u> **86**(10): 1367-1372.
Esau, K. (1960). "Anatomy of Seed Plants." <u>Soil Science</u> **90**(2): 149.
Ewers, F. W. (1985). "Xylem' Structure and Water Conduction in Conifer Trees, Dicot Trees, and
Llanas." <u>IAWA Journal</u> **6**(4): 309-317.
Feild, T. S. and T. Brodribb (2001). "Stem water transport and freeze-thaw xylem embolism in
conifers and angiosperms in a Tasmanian treeline heath." <u>Oecologia</u> **127**(3): 314-320.
Gerber, D., L. A. Wilen, E. R. Dufresne and R. W. Style (2023). "Polycrystallinity Enhances Stress
Buildup around Ice." <u>Physical Review Letters</u> **131**(20): 208201.
Gusta, L. V., M. Wisniewski, N. T. Nesbitt and M. L. Gusta (2004). "The effect of water, sugars, and
proteins on the pattern of ice nucleation and propagation in acclimated and nonacclimated canola
leaves." <u>Plant Physiol</u> **135**(3): 1642-1653.
Guy, C. L. (1990). "Cold acclimation and freezing stress tolerance: role of protein metabolism."
<u>Annual review of plant biology</u> **41**(1): 187-223.
Hacke, U. and J. Sauter (1996). "Xylem dysfunction during winter and recovery of hydraulic
conductivity in diffuse-porous and ring-porous trees." <u>Oecologia</u> **105**: 435-439.
Hacker, J. and G. Neuner (2007). "Ice propagation in plants visualized at the tissue level by infrared
differential thermal analysis (IDTA)." <u>Tree Physiology</u> **27**(12): 1661-1670.
Hammel, H. T. (1967). "Freezing of Xylem Sap without Cavitation." <u>Plant Physiology</u> **42**(1): 55-66.
Johnson, K. M., C. R. Brodersen, M. R. Carins-Murphy, B. Choat and T. J. Brodribb (2020). "Xylem
embolism spreads by single-conduit events in three dry forest angiosperm stems." <u>Plant Physiology</u>
**184**: 212-222.
Johnson, K. M., G. J. Jordan and T. J. Brodribb (2018). "Wheat leaves embolized by water stress do
not recover function upon rewatering." <u>Plant Cell Environ</u> **41**(11): 2704-2714.
Kane, C. N. and S. A. M. McAdam (2024). "Spatial and Temporal Freezing Dynamics of Leaves
Revealed by Time-Lapse Imaging." <u>Plant, Cell & Environment</u> **n/a**(n/a).
Kanji, Z. A., L. A. Ladino, H. Wex, Y. Boose, M. Burkert-Kohn, D. J. Cziczo and M. Krämer (2017).
"Overview of Ice Nucleating Particles." <u>Meteorological Monographs</u> **58**: 1.1-1.33.
Karabourniotis, G., J. F. Bornman and D. Nikolopoulos (2000). "A possible optical role of the bundle
sheath extensions of the heterobaric leaves of Vitis vinifera and Quercus coccifera." <u>Plant, Cell &</u>
<u>Environment</u> **23**(4): 423-430.
Kawai, K., R. Miyoshi and N. Okada (2017). "Bundle sheath extensions are linked to water relations
but not to mechanical and structural properties of leaves." <u>Trees</u> **31**: 1227-1237.
Kokin, E., M. Pennar, V. Palge and K. Jürjenson (2018). "Strawberry leaf surface temperature
dynamics measured by thermal camera in night frost conditions."
Langan, S. J., F. W. Ewers and S. D. Davis (1997). "Xylem dysfunction caused by water stress and
freezing in two species of co-occurring chaparral shrubs." <u>Plant, Cell & Environment</u> **20**(4): 425-437.
Larcher, W., U. Meindl, E. Ralser and M. Ishikawa (1991). "Persistent Supercooling and Silica
Deposition in Cell Walls of Palm Leaves." <u>Journal of Plant Physiology</u> **139**(2): 146-154.

Li, Z., D. Luo, M. M. Ibrahim, E. Hou and C. Wang (2024). "Adaptive strategies to freeze-thaw cycles
in branch hydraulics of tree species coexisting in a temperate forest." Plant Physiology and
Biochemistry **206**: 108223.
Loehle, C. (1998). "Height growth rate tradeoffs determine northern and southern range limits for
trees." Journal of Biogeography **25**(4): 735-742.
Lütz, C. (2010). "Cell physiology of plants growing in cold environments." Protoplasma **244**(1): 53-73.
Mayr, S., P. Schmid, J. Laur, S. Rosner, K. Charra-Vaskou, B. Dämon and U. G. Hacke (2014). "Uptake
of Water via Branches Helps Timberline Conifers Refill Embolized Xylem in Late Winter    " Plant
Physiology **164**(4): 1731-1740.
Mazur, P. (1969). "Freezing Injury in Plants." Annual Review of Plant Biology **20**(Volume 20, 1969):
419-448.
Mrad, A., D. M. Johnson, D. M. Love and J.-C. Domec (2020). "The roles of conduit redundancy and
connectivity in xylem hydraulic functions." New Phytol.
Muffler, L., C. Beierkuhnlein, G. Aas, A. Jentsch, A. H. Schweiger, C. Zohner and J. Kreyling (2016).
"Distribution ranges and spring phenology explain late frost sensitivity in 170 woody plants from the
Northern Hemisphere." Global Ecology and Biogeography **25**(9): 1061-1071.
Neger, F. W. (1918). "Die Wegsamkeit der Laubblätter für Gase." Flora oder Allgemeine Botanische
Zeitung **111**: 152-161.
Pieruschka, R., U. Schurr, M. Jensen, W. F. Wolff and S. Jahnke (2006). "Lateral diffusion of CO2 from
shaded to illuminated leaf parts affects photosynthesis inside homobaric leaves." New Phytologist
**169**(4): 779-788.
Pittermann, J. and J. Sperry (2003). "Tracheid diameter is the key trait determining the extent of
freezing-induced embolism in conifers." Tree Physiology **23**(13): 907-914.
Pittermann, J. and J. S. Sperry (2006). "Analysis of Freeze-Thaw Embolism in Conifers. The Interaction
between Cavitation Pressure and Tracheid Size." Plant Physiology **140**(1): 374-382.
Pray, T. R. (1954). "Foliar Venation of Angiosperms. I. Mature Venation of Liriodendron." American
Journal of Botany **41**(8): 663-670.
Robinson, J. A., M. Rennie, M. Clearwater, D. J. Holland, A. K. van den Berg and M. Watson (2023).
"Examination of embolisms in maple and birch saplings utilising microCT." Micron **168**: 103438.
Ruelland, E., M.-N. Vaultier, A. Zachowski and V. Hurry (2009). Chapter 2 Cold Signalling and Cold
Acclimation in Plants. Advances in Botanical Research, Academic Press. **49:** 35-150.
Sakai, A. and W. Larcher (2012). Frost survival of plants: responses and adaptation to freezing stress,
Springer Science & Business Media.
Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden,
S. Saalfeld, B. Schmid, J.-Y. Tinevez, D. J. White, V. Hartenstein, K. Eliceiri, P. Tomancak and A.
Cardona (2012). "Fiji: an open-source platform for biological-image analysis." Nature Methods **9**(7):
676-682.
Sevanto, S., N. M. Holbrook and M. Ball (2012). "Freeze/Thaw-Induced Embolism: Probability of
Critical Bubble Formation Depends on Speed of Ice Formation." Frontiers in Plant Science **3**.
Shardt, N., F. N. Isenrich, B. Waser, C. Marcolli, Z. A. Kanji, A. J. deMello and U. Lohmann (2022).
"Homogeneous freezing of water droplets for different volumes and cooling rates." Physical
Chemistry Chemical Physics **24**(46): 28213-28221.
Sperry, J. S. (1995). 5 - Limitations on Stem Water Transport and Their Consequences. Plant Stems. B.
L. Gartner. San Diego, Academic Press: 105-124.
Sperry, J. S., J. R. Donnelly and M. T. Tyree (1988). "SEASONAL OCCURRENCE OF XYLEM EMBOLISM
IN SUGAR MAPLE (ACER SACCHARUM)." American Journal of Botany **75**(8): 1212-1218.
Sperry, J. S., K. L. Nichols, J. E. M. Sullivan and S. E. Eastlack (1994). "Xylem Embolism in Ring-Porous,
Diffuse-Porous, and Coniferous Trees of Northern Utah and Interior Alaska." Ecology **75**(6): 1736-
1752.

Sperry, J. S. and J. E. M. Sullivan (1992). "Xylem Embolism in Response to Freeze-Thaw Cycles and
Water Stress in Ring-Porous, Diffuse-Porous, and Conifer Species 1." Plant Physiology **100**(2): 605-
613.
Stegner, M., O. Buchner, M. Geßlbauer, J. Lindner, A. Flörl, N. Xiao, A. Holzinger, N. Gierlinger and G.
Neuner (2023). "Frozen mountain pine needles: The endodermis discriminates between the ice-
containing central tissue and the ice-free fully functional mesophyll." Physiologia Plantarum **175**(1):
e13865.
Steponkus, P. and M. Webb (1992). Freeze-induced dehydration and membrane destabilization in
plants. Water and Life: Comparative Analysis of Water Relationships at the Organismic, Cellular, and
Molecular Levels, Springer: 338-362.
Steponkus, P. L., M. F. Dowgert and W. J. Gordon-Kamm (1983). "Destabilization of the plasma
membrane of isolated plant protoplasts during a freeze-thaw cycle: The influence of cold
acclimation." Cryobiology **20**(4): 448-465.
Thonglim, A., G. Bortolami, S. Delzon, M. Larter, R. Offringa, J. J. B. Keurentjes, E. Smets, S. Balazadeh
and F. Lens (2023). "Drought response in Arabidopsis displays synergistic coordination between
stems and leaves." Journal of Experimental Botany **74**(3): 1004-1021.
Thonglim, A., S. Delzon, M. Larter, O. Karami, A. Rahimi, R. Offringa, J. J. B. Keurentjes, S. Balazadeh,
E. Smets and F. Lens (2021). "Intervessel pit membrane thickness best explains variation in embolism
resistance amongst stems of Arabidopsis thaliana accessions." Annals of Botany **128**(2): 171-182.
Tonet, V., M. Carins-Murphy, R. Deans and T. J. Brodribb (2023). "Deadly acceleration in dehydration
of Eucalyptus viminalis leaves coincides with high-order vein cavitation." Plant Physiology **191**(3):
1648-1661.
Trifiló, P., F. Raimondo, T. Savi, M. A. Lo Gullo and A. Nardini (2016). "The contribution of vascular
and extra-vascular water pathways to drought-induced decline of leaf hydraulic conductance."
Journal of Experimental Botany **67**(17): 5029-5039.
Tyree, M. T. and M. H. Zimmermann (2002). Xylem structure and the ascent of sap, Springer Science
& Business Media.
Vitasse, Y., A. Bottero, M. Cailleret, C. Bigler, P. Fonti, A. Gessler, M. Lévesque, B. Rohner, P. Weber,
A. Rigling and T. Wohlgemuth (2019). "Contrasting resistance and resilience to extreme drought and
late spring frost in five major European tree species." Global Change Biology **25**(11): 3781-3792.
Vitra, A., A. Lenz and Y. Vitasse (2017). "Frost hardening and dehardening potential in temperate
trees from winter to budburst." New Phytologist **216**(1): 113-123.
Wise, R. R. (1995). "Chilling-enhanced photooxidation: The production, action and study of reactive
oxygen species produced during chilling in the light." Photosynthesis Research **45**(2): 79-97.
Wisniewski, M. and M. Fuller (1999). Ice nucleation and deep supercooling in plants: new insights
using infrared thermography. Cold-Adapted Organisms: Ecology, Physiology, Enzymology and
Molecular Biology. R. Margesin and F. Schinner. Berlin, Heidelberg, Springer Berlin Heidelberg: 105-
118.
Wylie, R. B. (1943). "The Role of the Epidermis in Foliar Organization and its Relations to the Minor
Venation." American Journal of Botany **30**(4): 273-280.
Wylie, R. B. (1952). "The Bundle Sheath Extension in Leaves of Dicotyledons." American Journal of
Botany **39**(9): 645-651.
Yang, S., D. Gerber, Y. Feng, N. Bain, M. Kuster, L. de Lorenzis, Y. Xu, E. R. Dufresne and R. W. Style
(2024). "Dehydration drives damage in the freezing of brittle hydrogels." arXiv preprint
arXiv:2401.12871.
Yarberry, W. and W. Yarberry (2021). "Dplyr." CRAN recipes: DPLYR, stringr, lubridate, and regex in
R: 1-58.
Zanne, A. E., D. C. Tank, W. K. Cornwell, J. M. Eastman, S. A. Smith, R. G. FitzJohn, D. J. McGlinn, B. C.
O'Meara, A. T. Moles, P. B. Reich, D. L. Royer, D. E. Soltis, P. F. Stevens, M. Westoby, I. J. Wright, L.
Aarssen, R. I. Bertin, A. Calaminus, R. Govaerts, F. Hemmings, M. R. Leishman, J. Oleksyn, P. S. Soltis,

N. G. Swenson, L. Warman and J. M. Beaulieu (2014). "Three keys to the radiation of angiosperms
into freezing environments." <u>Nature</u> **506**(7486): 89-92.
Zhang, P., B. Han, P. Ding and J.-J. Zhu (2023). "An approach to quantify the free water content in
leaves of *Quercus acutissima* Carruth. with differential thermal analysis." <u>Plant Science Journal</u> **41**(5):
687-693.
Zwieniecki, M. A., T. J. Brodribb and N. M. Holbrook (2007). "Hydraulic design of leaves: insights from
rehydration kinetics." <u>Plant, Cell & Environment</u> **30**(8): 910-921.

1028

## Figure Legends

**Figure 1**: The set-up for monitoring freeze in *L. tulipifera* trees.

**Figure 2:** The average percentage of initial (pre-treatment) mean greyscale value (brightness) for all
18 leaves that froze, comparing tissue that did and did not freeze within the same leaves.

**Figure 3:** Percentage of initial greyscale value over time for the midvein and mesophyll for 18 *L.*
*tulipifera* leaves frozen to -4°C or -5°C.

**Figure 4:** Representative freezing and thawing sequence including images, mean greyscale value
and air temperature for a representative *L. tulipifera* leaf visualised using time-lapse imaging.

**Figure 5:** L. *tulipifera* leaves (n=8) at their maximum freezing extent (left hand column) and ~ 10
hours after thawing (right hand column), demonstrating the reddening of the tissue that occurred in
frozen tissue.

**Figure 6:** The percentage frozen leaf area in leaves frozen to air temperatures of -4°C and - 5°C.

**Figure 7**: Changes in chlorophyll fluorescence in response to negative temperatures in 24 leaves of
*L. tulipifera*.

**Figure 8:** Representative image sequences showing the difference between drought- and freeze-
thaw-induced embolism in *L. tulipifera* leaves.

**Figure 9:** Xylem vessel lumen diameter distributions for leaves of *L. tulipifera*.

1046