Genetic and Morphological Insights Reveal the Hidden and Colorful Diversity in *Oscarella* Sponges

E. Guiollot[1,2], D. Guillemain[3], E. Renard[1], C. Borchiellini[1] & Q. Schenkelaars[1*]

[1] Aix-Marseille Université, CNRS, IRD, Avignon Université, IMBE, UMR 7263, Station Marine d'Endoume,
Rue de la Batterie des Lions, 13007 Marseille, France
[2] Marine Biology Department, Centre Scientifique de Monaco, Monaco, 98000, Monaco
[3] Aix-Marseille Université, OSU Institut Pythéas, Campus de Luminy OCEANOMED, 163 avenue de
Luminy, 13009 Marseille, France
* Corresponding author : quentin.schenkelaars@imbe.fr

10
11
ABSTRACT

The identification and classification of new taxa are crucial for understanding biodiversity. However,
assigning samples to new taxa requires cautious and rigorous approaches. Historically, taxonomy has
heavily relied on morphological traits, which can be subjective and may not always reflect underlying
genetic divergence—particularly in organisms with few diagnostic morphological traits. A prime
example is the sponge genus *Oscarella* (Homoscleromorpha), where species delimitation is challenged
by the absence of spicules and limited cell diversity. Here, we address this gap by combining an
extensive genetic dataset (192 specimens, five markers) with systematic photographic documentation.
This approach enabled a robust assessment of *Oscarella* diversity in the Western Mediterranean,
resulting in the identification of four species new to science. Multigene phylogenetic analyses also
enabled us to propose an evolutionary scenario for color polymorphism. Moreover, our data
highlighted critical limitations in current methodologies for studying *Oscarella*, including the low
resolution of the widely used cytochrome c oxidase subunit I (cox1/COI) gene, the lack of genetic data
for many species, and insufficient information on their geographical distribution. These issues, which
mirror challenges across many taxa, highlight the urgent need for standardized genetic frameworks and
comprehensive datasets to improve taxonomic resolution for each taxon.

Key words: Porifera, Genetic markers, Species delimitation, Taxonomy, Biodiversity

INTRODUCTION

Marine biodiversity remains largely unknown or underestimated in many ecosystems, despite their undeniable ecological, economic, and cultural importance[1]. Monitoring and implementing appropriate local management of these ecosystems therefore remain major challenges. One reason is that marine habitats are less accessible than most continental ones, requiring oceanographic vessels or diving equipment. Furthermore, they are often subject to strict regulations. As a result, marine ecosystems lag behind in their taxonomic description. Another factor contributing to this gap is the recurrent clustering of cryptic, non-vertebrate species under the names of presumably cosmopolitan taxa[2–5]. This issue stems from the frequent lack of clear morphological traits in numerous invertebrate groups, and from the fact that, although the use of molecular data for species description has gained wider acceptance, it is still not universally endorsed. Among the marine invertebrates facing such challenges, sponges are a prime example[6,7].

Sponges (Porifera) are major components of many marine ecosystems. Together with other invertebrates such as corals and gorgonians (Cnidaria), they form structural frameworks that provide habitats for numerous organisms. Accordingly, many sponge species are considered ecosystem engineers, whose life cycles and persistence strongly influence the surrounding community. In the Mediterranean Sea, approximately 1,000 sponge species have been recorded, with a high level of endemism[8]. Among them, *Oscarella* species (Homoscleromorpha) play important roles, covering up to 37% of the substrate covered by sponges in some coralligenous ecosystems[9]. The first described *Oscarella* species was *O. lobularis* (Schmidt, 1862)[10]. Originally identified in the Mediterranean Sea, it has since been reported from various locations worldwide[11–30], and is therefore often regarded as cosmopolitan. To date, 25 *Oscarella* species are formally recognized worldwide[31] (**Table 1**), but their true diversity remains poorly understood, and their geographical distribution is still largely unresolved. In this context, the Bay of Marseille (Western Mediterranean Sea, France) can be considered the cradle of *Oscarella* research, as it is the area where species diversity has been most thoroughly documented, with seven species clearly identified (**Table 1**)[32,7,33].

Table 1: Chronological list of *Oscarella* accepted species according to Word Porifera Database. Species names underlined in blue indicate those found in the Bay of Marseille, Gulf of Lion, Western Mediterranean Sea. The table includes references for the original descriptions, along with the corresponding sampling locations for each species. Additionally, it provides information on whether informative images were included in the original descriptions and whether genetic data are available. The data presented in this study are highlighted in red. Note that, according to all phylogenetic studies, including this one, *Pseudocorticium jarrei* should be renamed *Oscarella jarrei*.

Year	Scientific Name	Reference	Original location	Picture*	Genetic data
1862	*Oscarella lobularis*	Schmidt, 1862	Mediterranean Sea, Adriatic Sea, Croatia	no	Gazave et al., 2010 (mitochondrion, *18S, 28S*) ; Ivanisevic et al., 2011 (*cox1*) ; Gazave et al., 2013 (*atp6, tatC*) ; Ereskovsky et al., 2017 (*rDNA*) ; Guiollot et al., (*nad1-atp8, epic12*)
1868	*Oscarella tuberculata*	Schmidt, 1868	Mediterranean Sea, Adriatic Sea, Croatia	?	Gazave et al., 2010 (mitochondrion) ; Ivanisevic et al., 2011 (*cox1*) ; Gazave et al., 2013 (*18S, 28S, atp6, tatC*) ; Ereskovsky et al., 2017 (*rDNA*) ; Guiollot et al., (*nad1-atp8, epic12*)
1876	*Oscarella cruenta*	Carter, 1876	North Atlantic Ocean, Cape St. Vincent, Portugal	no	no
1909	*Oscarella membranacea*	Hentschel, 1909	Indian Ocean, Shark Bay, Autralia	no	no
1909	*Oscarella tenuis*	Hentschel, 1909	Indian Ocean, Shark Bay, Autralia	no	no
1995	*Pseudocorticium jarrei*	Boury-Esnault et al., 1995	Mediterranean Sea, Gulf of Lion, France	yes	Gazave et al., 2010 (mitochondrion, *18S, 28S*) ; Ivanisevic et al., 2011 (*cox1*) ; Gazave et al., 2013 (*atp6, tatC*) ; Ereskovsky et al., 2017 (*rDNA*)
1996	*Oscarella viridis*	Muricy et al., 1996	Mediterranean Sea, Gulf of Lion, France	no	Gazave et al., 2010 (mitochondrion) ; Ivanisevic et al., 2011 (*cox1*) ; Gazave et al., 2013 (*18S, atp6, tatC*) ; Ereskovsky et al., 2017 (*rDNA*) ; Guiollot et al., (*nad1-atp8, epic12*)
1996	*Oscarella microlobata*	Muricy et al., 1996	Mediterranean Sea, Gulf of Lion, France	yes	Gazave et al., 2010 (mitochondrion, *18S, 28S*) ; Ivanisevic et al., 2011 (*cox1*) ; Gazave et al., 2013 (*atp6, tatC*) ; Ereskovsky et al., 2017 (*rDNA*) ; Guiollot et al., (*nad1-atp8, epic12*)
1996	*Oscarella imperialis*	Muricy et al., 1996	Mediterranean Sea, Gulf of Lion, France	no	no
2004	*Oscarella carmela*	Muricy and Pearse, 2004	North Pacific Ocean, Carmel Bay, California	yes	Kober et Nichols, 2006 (*28S*) ; Lavrov et al., 2008 (mitochondrion, *18S*) ; Gazave et al., 2013 (*atp6, tatC*) ; Ereskovsky et al., 2017 (*rDNA*)
2004	*Oscarella nigraviolacea*	Bergquist and Kelly 2004	Indian Ocean, Pemba Island, Tanzania	no	no
2004	*Oscarella ochreacea*	Muricy and Pearse, 2004	Indian Ocean, Madagascar	no	no
2004	*Oscarella stillans*	Bergquist and Kelly 2004	Sulu Sea, Honda Bay, Palawan	no	no
2006	*Oscarella malakhovi*	Ereskovsky, 2006	Japan Sea, Vostok Bay, Russia	yes	Gazave et al., 2010 (mitochondrion, *18S, 28S*) ; Gazave et al., 2013 (*atp6, tatC*) ; Ereskovsky et al., 2017 (*rDNA*)
2009	*Oscarella kamchatkensis*	Ereskovsky et al., 2009	North Pacific Ocean, Avacha Gulf, Russia	yes	Gazave et al., 2013 (*atp6, tatC, 18S, 28S*) ; Ereskovsky et al., 2017 (*rDNA*)
2011	*Oscarella balibaloi*	Pérez et al., 2011	Mediterranean Sea, Gulf of Lion, France	yes	Ivanisevic et al., 2011 (*cox1*) ; Gazave et al., 2013 (*18S, atp6, tatC*) ; Ereskovsky et al., 2017 (partial mitochondrion, *rDNA*) ; Guiollot et al., (*nad1-atp8, epic12*)
2013	*Oscarella bergenensis*	Gazave et al., 2013	North Sea, Bergen Fjords, Norway	yes	Gazave et al., 2013 (*18S, 28S, atp6, tatC*) ; Ereskovsky et al., 2017 (*rDNA*)
2013	*Oscarella nicolae*	Gazave et al., 2013	North Sea, Bergen Fjords, Norway	yes	Gazave et al., 2013 (*18S, 28S, atp6, tatC*) ; Ereskovsky et al., 2017 (*rDNA*)
2017	*Oscarella pearsei*	Ereskovsky et al., 2017	North Pacific Ocean, Monterey Bay, California	yes	Ereskovsky et al., 2017 (mitochondrion, rDNA)
2018	*Oscarella filipoi*	Pérez and Ruiz, 2018	Caribbean Sea, Martinique	yes	Pérez and Ruiz, 2018 (*cox1*)
2018	*Oscarella zoranja*	Pérez and Ruiz, 2018	Caribbean Sea, Martinique	yes	Pérez and Ruiz, 2018 (*cox1*)
2022	*Oscarella minka*	Ruiz et al., 2022	Caribbean Sea, Martinique	yes	Ruiz et al., 2022 (*cox1*)
2022	*Oscarella aurantia*	Stillitani et al, 2022	South Atlantic Ocean, Cabo Frio, Brazil	yes	Stillitani et al, 2022 (*cob, cox1*)
2022	*Oscarella carollineae*	Stillitani et al, 2022	South Atlantic Ocean, Cabo Frio, Brazil	yes	Stillitani et al, 2022 (*cob, cox1*)
2022	*Oscarella ruthae*	Stillitani et al, 2022	South Atlantic Ocean, Cabo Frio, Brazil	yes	Stillitani et al, 2022 (*cob, cox1*)
2024	*Oscarella veronicae sp. nov.*	present study	Mediterranean Sea, Gulf of Lion, France	yes	Guiollot et al., (partial mitochondrion, *atp6, cox1, tatC, nad1-atp8, epic12*)
2024	*Oscarella sabinae sp. nov.*	present study	Mediterranean Sea, Gulf of Lion, France	yes	Guiollot et al., (partial mitochondrion, *atp6, cox1, tatC, nad1-atp8, epic12*)
2024	*Oscarella lunettae sp. nov.*	present study	Mediterranean Sea, Gulf of Lion, France	yes	Guiollot et al., (partial mitochondrion, *atp6, cox1, tatC, nad1-atp8, epic12*)
2024	*Oscarella fragilis sp. nov.*	present study	Mediterranean Sea, Gulf of Lion, France	yes	Guiollot et al., (partial mitochondrion, *atp6, cox1, tatC, nad1-atp8, epic12*)

Species identification in sponges often relies on the description of their skeletons. The composition,
shape, and size of small mineral structures called spicules, as well as the density and reticulation of
organic fibers, vary greatly among species[34]. This approach, however, cannot be applied to *Oscarella*,
as these sponges lack a skeleton[8]. Consequently, species identification in this genus has traditionally
relied on morphology (*e.g.*, lobe shape, surface texture), cytological features (*e.g.*, cell diversity and
distribution), tissue consistency (fragile, soft, or cartilaginous), and color[7]. The limited number of traits,
their low variability, and their inherent subjectivity make reliable species delimitation challenging. Since
2013, molecular analyses have therefore been systematically incorporated into the description of new
species in this genus.
76
In the first genetic survey of *Oscarella*, two nuclear markers (*18S* and *28S ribosomal DNA*) and two
mitochondrial markers (*ATP synthase membrane subunit 6*, *atp6*, and *twin-arginine translocase*
component, *tatC*) were used to study a few samples collected worldwide[35]. While this approach was
sufficient to discriminate most species and describe two new taxa, it failed to efficiently distinguish *O.*
lobularis and *O. tuberculata*. In more recent studies, the *I3-M11* fragment of the widely used
mitochondrial marker *cytochrome c oxidase subunit 1* (*cox1/COI*) was applied[36–40]. Unfortunately, it also
failed to discriminate some closely related *Oscarella* species, such as *O. carollineae, O. filipoi* and *O.*
ruthae[36]. These results may reflect (i) that the mentioned taxa are in fact single species, (ii) that the
markers used are insufficiently informative to distinguish closely related species (*e.g.*, due to low
mutation rates or incomplete lineage sorting)[41], or (iii) that species identification based on morphology
and tissue consistency is not straightforward.
88
In this study, we collected 192 specimens of *Oscarella* spp. from eight locations across the Bay of
Marseille to conduct a comprehensive genetic survey of local diversity. Our barcoding approach
included (i) the three previously used mitochondrial protein-coding markers (*atp6*, *cox1*, and *tatC*), (ii)
a newly developed mitochondrial marker covering the end of the NADH dehydrogenase subunit 1 gene,
the beginning of the ATP synthase membrane subunit 8 gene, and the intergenic region (*nad1-atp8*),
and (iii) a newly developed nuclear marker, an Exon Primed Intron Crossing (EPIC) locus (*epic12*)[42,43].
We compared these five markers and identified the most suitable for implementing a barcoding
approach aimed at efficient species identification. In addition to species delimitation, partial
mitochondrial genomes were obtained to clarify phylogenetic relationships and trace the evolution of
color polymorphisms. Overall, this study represents the most comprehensive genetic investigation of
this genus to date, both in terms of specimen sampling and the number of genetic markers.
100
101
METHODS
Specimen collection
We collected 192 *Oscarella* spp. specimens representing different color morphs from multiple locations
across the Bay of Marseille (**Figure 1**). Each specimen was photographed (**Supplementary Figure 1**) and
then assigned a unique identifier incorporating the sampling location and sample number (*e.g.*,
3PP_01; **Supplementary Table 1**). Because our goal was to preserve the possibility of conducting
complementary analyses on the same individuals (*e.g.,* immunohistochemistry, *in situ* hybridization) in
addition to PCR amplification, we employed here a somewhat unconventional protocol previously
developed in *Hydra*[44]. Accordingly, specimens were fixed overnight at 4 °C in a solution containing 4%
paraformaldehyde (50% PBS-Tween 1‰, 25% seawater, 25% 16% paraformaldehyde aqueous
solution). The following day, samples were washed in PBS-Tween, gradually transferred to increasing
methanol concentrations, and finally stored in 100% methanol at −20 °C. When possible, one holotype
and one paratype have been deposited at the Muséum National d'Histoire Naturelle de Paris for each

new species. The corresponding accession numbers are provided in the Systematics section. All
additional samples are maintained at our home institution (IMBE). For many specimens, small pieces
of fresh tissue were also flash-frozen in liquid nitrogen and stored at −80 °C at the home institute
(IMBE).

Figure 1: Sampling locations across the Bay of Marseille and number of samples per location.

Marker amplification, sequencing, and haplotype networks

In a previous study, we demonstrated that a direct-PCR approach based on rapid mechanical grinding
with a pestle yields PCR products of quality comparable to standard extraction-based methods, while
requiring substantially less time and reducing costs[44]. For this reason, no DNA extraction was performed
in the present study. Instead, a small piece of fixed tissue (~3 mm²) was ground in 1 mL of water with
a pestle to serve as crude template. Mitochondrial (*atp6*, *cox1*, *nad1-atp8*, *tatC*) and nuclear (*epic12*)
markers were amplified using GoTaq® G2 Flexi DNA polymerase (Promega). PCR reactions were
performed in 20 µL volumes containing 1X buffer, 0.2 mM dNTPs, 1 µM primers, 1.5 mM MgCl₂, 0.5 U
GoTaq, and 3 µL of template. Amplifications were carried out for 30 cycles. Primer sequences, amplicon
sizes, and PCR annealing temperatures are listed in **Table 2**. PCR products were verified on 1% agarose
gels and submitted to Eurofins Genomics for Sanger sequencing. All sequences generated in this study
have been deposited in the NCBI database (**Supplementary Table 2**). Sequences of *nad1-atp8* and *tatC*
were aligned using MUSCLE[45], whereas other markers did not require any alignment because the
sequences are highly conserved, identical in length, and contain no indels. For these markers, we simply
trimmed the sequence ends to standardize their length. Haplotype networks were constructed using
the Minimum Spanning Network function in PopArt[46].

147 **Table 2: Information regarding PCR primers, Tm and amplicon size.**

Foward primer		Reverse primer		Tm	Amplicon size
Name	Sequence	Name	Sequence		
*Oscarella_atp6_Fw**	TGACAGCAGCATATTTTGATCAG	*Oscarella_atp6_Rv*	AACACATAGGCCTGTATTACAGC	45°C	700 bp
Oscarella_cox1_Fw	CCGCCATTGTGTGCATATCA	*Oscarella_cox1_Rv**	TCACCCTGCAAAACTATCAGC	45°C	1 440 bp
*Oscarella_i12_Fw**	CAATCAGACAAGAAAATGCTGAC	*Oscarella_i12_Rv*	TTGCTGTCNACNGGAATATCC	45°C	350 - 700 bp
Oscarella_mito_Fw	AGATTGCTGCGGGGATAAGA	*Oscarella_mito_Rv*	ATTCCGCGCTCATTTGGTTT	53°C	9 950 bp
Oscarella_nad1-atp8_Fw	CAGGATGGGCAAGTAACTCTAA	*Oscarella_nad1-atp8_Rv**	CTGATCAAAATATGCTGCTGTCA	45°C	1 100 bp
*Oscarella_tatC_Fw**	TGTTGGGTAGCTACGACGAA	*Oscarella_tatC_Rv*	GCCTACCATTAACCGAATCTCC	57°C	1 140 bp
*O.veronica_SNP_Fw**	GGTTTGCCCTTTCAGAAGCA	*O.veronica_SNP_Rv*	TAACGCCCCGATTAAGAGCA	45°C	388 bp

Mitochondrial genome amplification of new species, sequencing, and phylogenetic reconstruction

DNA was extracted from 25 mg of flash-frozen tissue using the DNeasy® Blood & Tissue Kit (Qiagen).
Long PCRs were performed with GoTaq® Long PCR Master Mix (Promega) containing 1X mix, 1 µM
primers, and 5 µg DNA. Primer sequences, amplicon sizes, and melting temperatures are provided in
Table 2. PCR amplicons were visualized on 0.5% agarose gels, purified using the Qiagen DNA extraction
kit, and sequenced via Oxford Nanopore Technologies services at Eurofins Genomics. Partial
mitochondrial genomes have been deposited in the NCBI database (**Supplementary Table 2**).
Amplicons encompassed eight mitochondrial protein-coding genes (*cox1*, *nad1*, *atp8*, *atp6*, *cox3*, *atp9*,
nad4, and *nad6*), which were concatenated for each species. Resulting sequences were aligned with
MUSCLE[45], and poorly aligned positions were removed using Gblocks with the following settings:
minimal conservation for a flank position 85%, maximal contiguous non-conserved positions 8 bp,
minimal block length 10 bp, and no gaps allowed in final blocks[47], retaining 98% of the original 6591
positions. The best-fit nucleotide substitution model was estimated using jModelTest[48]. Phylogenetic
reconstruction was performed in MEGA[49] using the Maximum Likelihood method with the General
Time Reversible model, a gamma-distributed substitution rate (four categories), and a proportion of
invariable sites (GTR + I + G). Branch support was assessed with 1000 bootstrap replicates.

165
166

RESULTS
Comprehensive genetic survey of *Oscarella* species in the Bay of Marseille

PCR amplification of the *atp6* marker using crude macerates as templates was successful for the
majority of specimens (95% amplification success). Consequently, *atp6* haplotypes were used as a
practical initial partitioning tool, at a stage where morphological identification was not yet possible.
Analysis of *atp6* sequences from 183 newly collected specimens identified 11 distinct *atp6* haplotypes,
labeled a to k (**Figure 2A**). Comparisons with NCBI reference mitochondrial genomes assigned *atp6*
haplotypes a and d to *O. tuberculata*, b to *O. lobularis*, g to *O. viridis*, and h to *O. microlobata*. In
contrast, none of the *atp6* sequences matched *O. balibaloi* or *P. jarrei*, so we incorporated available
mitochondrial genome sequences for these two species into the *atp6* network (*atp6* haplotypes l and
m, respectively). Species identification could not be determined for the remaining six *atp6* haplotypes.
Notably, *atp6* haplotypes e and j (corresponding to ten and one samples, respectively) branched
directly on the main *O. tuberculata* haplotype (*atp6* haplotype a) without additional connections
(**Figure 2A**).

To (i) validate the *atp6* data, (ii) assess whether the six unresolved *atp6* haplotypes represent new taxa,
and (iii) obtain genetic information for the nine specimens where *atp6* amplification failed, we
sequenced three additional mitochondrial markers in a subset of individuals: *nad1-atp8* (66% of
specimens), *tatC* (40%), and *cox1* (14%) (**Supplementary Figure 2**). Overall, the partitions obtained
from these markers—except for *cox1*—were either fully consistent or compatible with the *atp6* data
(**Figure 2B**, **Supplementary Table 1**, **Supplementary Figure 2**). For instance, *atp6* haplotype b and *tatC*

haplotype a were consistently associated, with all individuals carrying one also carrying the other (*i.e.*
consistent). Alternatively, only individuals with *atp6* haplotype c carried *nad1–atp8* haplotype e or m,
and all individuals with *nad1–atp8* haplotype e or m carried *atp6* haplotype c (*i.e.* compatible).
Specimens for which *atp6* amplification failed were identified as *O. balibaloi* using *nad1-atp8* and *tatC*,
while *atp6* haplotypes a, d, e, and j must be grouped together as *O. tuberculata* according to *nad1-*
atp8 and *tatC* partitioning. These clustering patterns were further corroborated by the newly
developed nuclear marker *epic12*, which yielded data consistent or compatible with the *atp6* haplotype
network (**Figure 2B**, **Supplementary Table 1**, **Supplementary Figure 2**). Amplification of *epic12* failed
for *O. microlobata*.
197
Combined analyses of *atp6*, *nad1-atp8, tatC* and *epic12* markers strongly suggest that individuals
showing different *atp6* haplotypes correspond to distinct lineages, likely representing separate species.
Indeed, none of these lineages share alleles for the mitochondrial markers (*atp6*, *nad1–atp8*, *tatC*) nor
for the nuclear marker (*epic12*), despite their sympatric distribution within the Bay of Marseille
(**Supplementary Table 1**). Importantly, the nuclear dataset includes a substantial sample size (*n* = 103
individuals), providing robust support for the absence of allele sharing. This pattern strongly suggests
that no ongoing gene flow occurs among these lineages, which is consistent with a classical biological
definition of species. Thus, at least ten *Oscarella* species inhabit the bay, including four new to science.
We propose the following names: *O. veronicae* sp. nov. (*atp6* haplotype c), *O. sabinae* sp. nov. (*atp6*
haplotype k), *O. fragilis* sp. nov. (*atp6* haplotype i), and *O. lunettae* sp. nov. (*atp6* haplotype f). Formal
descriptions with morphological and genetic diagnosis are provided in the *Systematics* section (**Figure**
3 and 4).

210
Figure 2: Species Delimitation Analysis of *Oscarella spp.* mapped onto a Maximum Likelihood (ML)
Phylogenetic Tree. (A) Haplotype network based on the *atp6* marker. Each haplotype (a, b, c, etc.) is
depicted as a circle, with the size of the circle proportional to the number of specimens represented.
Numbers between haplotypes indicate the number of mutations separating them. (B) Schematic
representation of species partitions derived from single-locus sequences, either mitochondrial (*atp6*,
nad1-atp8, etc.) or nuclear (*epic12*). The proportion of each color morphs within each species is also
provided (color). For instance, among the 104 *O. tuberculata* specimens collected, 39% were pink, 22%
yellow, 20% blue, 9% purple, 9% black, and 1% beige. Detailed specimen information is provided in
Supplementary Table 1 and haplotype networks are shown in **Supplementary Figure 2**. (C) Maximum
likelihood (ML) phylogenetic cladogram of *Oscarella* species based on eight mitochondrial protein-
coding genes (*cox1*, *nad1*, *atp8*, *atp6*, *cox3*, *atp9*, *nad4*, and *nad6*). Only nodes with bootstrap values
greater than 85% (1000 BS replicates) are shown. Dots on the tree indicate the inferred ancestral color
state of the common ancestor of *Oscarella* and the last common ancestor of *O. lobularis*, *O. sabinae*
sp. nov., *O. tuberculata*, and *O. veronicae* sp. nov. Note that *Pseudocorticium jarrei* is proposed to be
renamed *Oscarella jarrei,* in agreement with the present results and previous ones (**Table 1**).
226

Figure 3: Comparative external morphology of *Oscarella* species within the polychromatic complex.
(A to C') *Oscarella lobularis.* (D to F') *Oscarella tuberculata.* (G and G') *Oscarella sabinae* sp. nov. (H and H') *Oscarella veronicae* sp. nov. (A, B, C, D, E, F, G and H) Top-view photographs of the entire specimens. Scale: 1 cm. (A', B', C', D', E', F', G' and H') Close-up view of the lobes. Scale: 0.25 cm.

Phylogenetic relationships reveal a complex of four polychromatic species

Haplotype network analyses indicated that four *Oscarella* species are closely related. Specifically, *O. tuberculata* and *O. lobularis* consistently clustered with *O. sabinae* sp. nov. and *O. veronicae* sp. nov. (**Figure 2A, Supplementary Figure 2**). Regardless of the marker used (*atp6, nad1-atp8, tatC, cox1,* or *epic12*), genetic distances among haplotypes within this complex were very short compared to the distances between this complex and the nearest neighboring species (*e.g.*, 0.4% vs. 5.2% for *atp6*). Similarly, *O. viridis* and *O. lunettae* sp. nov. were closely related across all markers (*e.g.*, 1.6% for *atp6*).

To further resolve relationships among *Oscarella* species, we sequenced 9,874 bp of the mitochondrial genome for one specimen of each new species. The resulting phylogenetic tree strongly supported (BS = 100%) the close relationship of *O. tuberculata*, *O. lobularis*, *O. sabinae* sp. nov., and *O. veronicae*

245 sp. nov. (**Figure 2C, Supplementary Figure 3**). The analysis also confirmed that *O. lunettae* sp. nov. is
246 the sister species to *O. viridis* (BS = 100%), whereas *O. fragilis* sp. nov. formed a distinct lineage,
247 diverging before the *O. carmela* / *O. malakovi* clade, two North Pacific species (BS = 100%).

Figure 4: Comparative external morphology of yellow and orange *Oscarella* specimens. (A and A')
Oscarella fragilis sp. nov. (B to C') *Oscarella balibaloi*. (D to E') *Oscarella microlobata*. (F and F')
Oscarella viridis. (G and G') *Oscarella lunettae*. (H and H') *Oscarella veronicae* sp. nov. (A, B, C, D, E, F,
G and H) Top-view photographs of the entire specimens. Scale: 1 cm. (A', B', C', D', E', F', G' and H') Close-
up view of the lobes. Scale: 0.25 cm.

Given the importance of color for species identification in *Oscarella*, we examined color polymorphism
in each species (**Figure 2B, Supplementary Figure 1**). *O. tuberculata*, *O. lobularis*, and *O. veronicae* sp.
nov. exhibited a broad color range, including pink and blue morphotypes (**Figure 3**). Notably, the only
O. sabinae sp. nov. specimen was bright pink, whereas other *Oscarella* species displayed colors from
white to bright or rusty orange, including shades of light brown, beige, and yellow (**Figure 4**).
Polychromatic species with pink, violet/purple, or blue morphotypes formed a monophyletic group.

Consequently, color is a reliable criterion for distinguishing species within the complex from those
outside it (although yellow specimens also occur within the complex), but it is not sufficient on its own
for definitive species identification.
266
267
DISCUSSION
Towards a standardized approach for species identification in the *Oscarella*
By the late 1990s and early 2000s, PCR and sequencing were established as reliable tools for species
identification and discovery. In *Oscarella*, barcoding was first applied only in 2013, using mitochondrial
markers (*atp6* and *tatC*) and standard nuclear markers (18S, 28S)[35]. Despite some discrepancies, both
mitochondrial markers showed overall more consistent results with the *a priori* assignment of
individuals to putative species than the two nuclear markers. This may be due to ribosomal genes
existing in multiple tandem copies, and while they are expected to evolve in a concerted manner,
slightly divergent paralogues can persist within the same genome[50]. Amplification of these mixed
copies can thus complicate phylogenetic interpretation. Consequently, subsequent studies solely relied
on alternative mitochondrial markers such as *cox1 I3–M11* and *cob*[36–38,51] (**Table 1**). Nevertheless, *cox1*
failed to separate *O. lobularis* from *O. veronicae sp. nov.* in our study, as observed in other recent
studies for *O. carollineae*, *O. filipoi*, and *O. ruthae*[36]. This highlights the broader limitations of *cox1/COI*
for species delimitation in some taxa[52,53]. In the *Oscarella* context, *atp6*, *nad1-atp8*, and *tatC*
outperformed *cox1* in discriminating closely related species, even though we observed high *cox1*
amplification success rates. Furthermore, to bypass issues observed with standard nuclear markers
(28S and 18S), we developed the novel single-copy nuclear marker *epic12*[42,43], which can be reliably
amplified across relatively distant *Oscarella* species. Interestingly, this new nuclear marker provided
results strongly consistent with the mitochondrial markers. Accordingly, for reliable cross-study
comparisons and robust phylogenetic reconstruction, we recommend that future species descriptions
include partial mitochondrial genomes encompassing all relevant protein-coding markers, the nuclear
marker *epic12*, and photographic documentation. Achieving a uniform dataset will also require
resampling species for which genetic data and/or images are currently unavailable/incomplete, as
illustrated by *O. imperialis* in the Western Mediterranean Sea, and by many others worldwide (**Table
1**), which cannot currently be identified with confidence[54,55].
293
Expanding geographical knowledge beyond type localities
A major challenge in understanding *Oscarella* diversity is the poorly known distribution of each species.
For most of them, records are limited to their type localities. Exceptions such as *O. lobularis* and *O.
tuberculata*, first described in the Adriatic Sea (Croatia)[10,56], exhibit broader Mediterranean
distributions and have even been reported from the North Atlantic Ocean[11–30,57–61]. However, most of
these records lack supporting evidence (no genetic data and few photographs) and should therefore
be considered potential sightings rather than confirmed occurrences. Comprehensive genetic sampling
across the Mediterranean and beyond is essential to accurately map species distributions, inform
conservation efforts (endemic *vs*. cosmopolitan species), and reveal new taxa. Our study exemplifies
this approach, identifying four new species in the Bay of Marseille and increasing local *Oscarella* species
richness by 57%. This is particularly relevant for the Homoscleromorpha, which comprise only 136
accepted species globally (<1.4% of total sponge diversity)[11].
306
Evolutionary insights and biochemical mysteries of color polymorphism
Our results provide strong evidence that color polymorphism in *Oscarella* has a clear evolutionary
signal. *O. tuberculata* and *O. lobularis* are truly polychromatic species, with morphotypes ranging from
pink to blue and violet, whereas other formally described species examined here exhibit more subdued

hues such as cream, yellow, orange, or brown[7,32,33,35–38,62–65]. Phylogenetic analyses including the new
species suggest that remarkable color polymorphism arose within a single lineage uniting *O. tuberculata*, *O. lobularis*, *O. veronicae* sp. nov., and *O. sabinae* sp. nov., and thus represents a
tuberculata, *O. lobularis*, *O. veronicae* sp. nov., and *O. sabinae* sp. nov., and thus represents a
synapomorphy of this clade. By contrast, the orange *O. fragilis* sp. nov. and the yellow *O. lunettae* sp.
nov. fall outside this group, following the broader pattern. Exploring additional species such as the
Brazilian *O. ruthae* and *O. filipoi*, which also show signs of color polymorphism (although no blue
morphs have been observed)[36], could shed further light on whether this trait evolved once or multiple
times within the genus.
319
Despite the dazzling colors displayed by many sponges, the biochemical basis of their pigmentation
remains a largely uncharted territory. To date, most identified pigments are conventional specialized
metabolites—quinones, alkaloids, terpenes, or carotenoids—responsible for the more common yellow,
red, or black hues[66]. In contrast, true blue pigments are exceptionally rare across the tree of life, with
coloration in most animals arising instead from structural effects such as nanocrystals or microscopic
scales[67]. Only a handful of genuine blue pigments have been identified so far, including anthocyanins
in plants[68], GFP-like chromoproteins in cnidarians[69,70], and tunichrome peptides in tunicates[71]. The
presence of vivid blue morphotypes in *Oscarella* therefore raises a fascinating question. Deciphering
the molecular basis of this polymorphism could open new perspectives in both evolutionary biology
and natural product chemistry.
330
331
332   **SYSTEMATICS**
333
334   Phylum: Porifera Grant, 1836.
335   Class: Homoscleromorpha Bergquist, 1978.
336   Order: Homosclerophorida Dendy, 1905.
337   Family: Oscarellidae Lendenfeld, 1887.
338   Genus: *Oscarella* Vosmaer, 1884.
339
340   ***Oscarella fragilis* sp. nov. (Schenkelaars & Guiollot)**
341
342   **Holotype:** IMBE-A059-JAR01: Collected from France, Mediterranean Sea, Marseille, Jarre (43.19°N,
343   5.36°W), at a depth of 14-17 m by Dorian Guillemain on 2023-04-26 (JAR_01, **Figure 4A and A'**).
344   Specimen fixed in 4% paraformaldehyde and preserved in absolute methanol.
345
346   **Paratype**: IMBE-A144-TIB11: Collected from France, Mediterranean Sea, Marseille, Tiboulen du Frioul
347   (43.2799°N, 5.2860°W), at a depth of 30-35 m by Dorian Guillemain on 2023-06-28 (TIB_11,
348   **Supplementary Figure 1**). Specimen fixed in 4% paraformaldehyde and preserved in absolute
349   methanol.
350
351   **Molecular diagnosis:** Molecular analysis provides strong evidence for recognizing *O. fragilis* sp. nov. as
352   a distinct species, with its genetic sequences showing high divergence from all other known *Oscarella*
353   species (**Figure 2**). The most striking example of this is the *epic12* sequences, which do not align with
354   any haplotypes found in other species (*epic12* haplotype i and p, **Supplementary Figure 2**).
355
356   **Morphological diagnosis:** Although the entire surface of this species is lobed, the lobes form
357   transparent, very membranous lamellae, a feature that starkly contrasts with those of other species in
358   the genus (**Figure 3** and **4**). Furthermore, the tissue is more fragile than in *O. viridis* yet considered

delicate[7]. While *O. viridis* tends to crumble, *O. fragilis* sp. nov exhibits a soft, almost mucous-like consistency, making manipulation with forceps particularly challenging.

Morphological description: Like other *Oscarella* species, *O. fragilis* sp. nov. is an encrusting sponge characterized by its irregular growth forms. It covers areas of 15–20 cm² with a thickness of 15–25 mm. The specimens collected in this study exhibit distinct colorations. One is pale orange (JAR_01, **Figure 4A**), while the other is bright orange (TIB_11, **Supplementary Figure 1**). This color pattern is reminiscent of *O. balibaloi* (*e.g.* 3PP_03 and MEJ_11, **Figure 4B to C'**) and *O. microlobata* (*e.g.* JAR_14 and JAR_11, **Figure 4D to E'**).

Ecology and distribution: One specimen was found encrusting red algae, while the other was encrusted on another organism, possibly the bryozoan *Smittina cervicornis* (TIB_11, **Supplementary Figure 1**). The first specimen was collected at a depth of 14–17 m, while the second was found at 30–35 m. Given the limited data, it is currently premature to draw definitive conclusions about the ecology and distribution of this species.

Oscarella lunettae **sp. nov. (Schenkelaars & Guiollot)**

Holotype: *Muséum National d'Histoire Naturelle de Paris*, MNHN-IP-2019-3414, IMBE-A205-LEV04: Collected from France, Mediterranean Sea, Marseille, Levant (43.09288°N, 5.37365°W), at a depth of 40 m by Dorian Guillemain on 2024-03-07 (LEV_04, **Figure 4G and G'**). Specimen fixed in 4% paraformaldehyde and preserved in absolute methanol.

Paratype: *Muséum National d'Histoire Naturelle de Paris*, MNHN-IP-2019-3415, IMBE-A206-LEV05: Collected from France, Mediterranean Sea, Marseille, Levant (43.09288°N, 5.37365°W), at a depth of 40 m Dorian Guillemain on 2024-03-07 (LEV_05, **Supplementary Figure 1**). Specimen fixed in 4% paraformaldehyde and preserved in absolute methanol.

Other material examined: Station Marine d'Endoume, IMBE: three additional specimens collected from Levant (43.09288°N, 5.37365°W) and Tiboulen du Frioul (43.2799°N, 5.2860°W) (**Supplementary Table 1** and **Supplementary Figure 1**). Specimen fixed in 4% paraformaldehyde and preserved in absolute methanol.

Molecular diagnosis: Molecular data strongly support the recognition of *O. lunettae* sp. nov. as a distinct species, as its sequences show significant divergence from all currently available *Oscarella* data (**Figure 2**). Although the *epic12* sequence in *O. lunettae* has the same length as in most other species (51 bp), it does not align with them. In fact, *epic12* haplotype g (**Supplementary Figure 2**) does not share more than three consecutive residues with any same-length sequence. Instead, it more closely resembles *epic12* haplotype j (corresponding to *O. viridis*), which is longer (55 bp). Yet, the two sequences remain clearly distinct, differing by 13 diagnostic residues.

Morphological diagnosis: Although this species shares beige and orange tones with *O. balibaloi*, *O. microlobata*, and *O. fragilis* sp. nov. (**Figure 4**), *O. lunettae* sp. nov. exhibits a characteristic color gradient—from yellowish or cream at the base to more intensely colored tissue at the apex of the lobes—that mirrors the pattern typically observed in *O. lobularis* (**Figure 3 A to C'**). However, unlike *O. lobularis*, which is invariably blue, purple, or pink, *O. lunettae* sp. nov. consistently displays orange hues, making this gradient pattern and coloration combination diagnostic (**Figure 4G and G'**).

Description: Like other *Oscarella* species, *O. lunettae* sp. nov. is an encrusting sponge characterized by irregular growth forms. It covers areas up to 80 cm² with a thickness ranging from 10 to 20 mm. Numerous ostia, which are easily visible under low magnification, perforate the sponge surface (**Figure 4G and G'**), thus resembling *O. balibaloi* (**Figure 4B' and C'**), and *O viridis* (**Figure 4F'**). The consistency of *O. lunettae* is relatively soft but not fragile, broadly comparable to that of *O. lobularis*.

Ecology and distribution: Like other *Oscarella* species found in the Bay of Marseille, *O. lunettae* sp. nov. is a sciaphilous species, typically found in coralligenous environments. To date, it has been observed at only two locations: Levant and Tiboulen du Frioul. While *O. lobularis* and *O. tuberculata* are also present in these locations, *O. lunettae* has not been detected elsewhere, suggesting that it may occupy a slightly different ecological niche. All specimens were collected at depths between 30 and 40 m, suggesting that *O. lunettae* might thrive below the thermocline, where temperatures are more stable.

Oscarella sabinae **sp. nov. (Schenkelaars & Guiollot)**

Holotype: *Muséum National d'Histoire Naturelle de Paris*, MNHN-IP-2019-3416, IMBE-B007-MAI95: Collected from France, Mediterranean Sea, Marseille, Maire Island (43.2096°N, 5.3353°W), at a depth of 8-18 m by Dorian Guillemain on 2024-06-19 (MAI_95, **Figure 3G and G'**). Specimen fixed in 4% paraformaldehyde and preserved in absolute methanol. No other material has been examined.

Molecular diagnosis: Molecular analysis of *O. sabinae* sp. nov. revealed distinct haplotypes for all mitochondrial markers, differentiating it from other species within the complex (**Supplementary Figure 2**). Comparative analysis of partial mitochondrial genomes identified 19 diagnostic positions unique to this species, including two in *nad1–atp8* ($A_{193} \rightarrow G$ and $A_{251} \rightarrow G$). Likewise, two diagnostic substitutions were detected in the *epic12* haplotype o when compared to all other haplotypes from species within the complex ($G_{14} \rightarrow A$ and $T_{50} \rightarrow A$).

Morphological diagnosis: In the single specimen of *O. sabinae* sp. nov. examined to date, the lobes display irregular expansions that give the sponge a bright pink, cauliflower-like appearance (**Figure 3G and G'**). Although *O. veronicae* sp. nov. can also exhibit a cauliflower-like morphology (**Figure 3H and H'**), none of the specimens observed showed a similarly intense pink coloration. Indeed, the hue of *O. sabinae* sp. nov. is more reminiscent of the pink form of *O. lobularis* (**Figure 3A and A'**), yet the latter species never exhibits a cauliflower-like surface (n = 42 in this study, with many more examined in the laboratory).

Description: Like other *Oscarella* species, *O. sabinae* sp. nov. is an encrusting sponge characterized by irregular growth forms. The specimen collected measured 10-15 cm². While the specimen was extremely soft, it was not fragile.

Ecology and Distribution: Out of 192 samples collected, only one specimen was identified as *O. sabinae* sp. nov., indicating that this species may be rare or that the sampling area may not represent its primary habitat. Consequently, it is premature to detail the ecology and distribution of this species based on the available data.

***Oscarella veronicae* sp. nov. (Schenkelaars & Guiollot)**

Holotype: *Muséum National d'Histoire Naturelle de Paris*, MNHN-IP-2019-3417, IMBE-A088-MAI60: Collected from France, Mediterranean Sea, Marseille, Maire Island (43.2096°N, 5.3353°W), at a depth of 6-10 m by Dorian Guillemain on 2023-04-12 (MAI_35, **Figure 3H and H'**). Specimen fixed in 4% paraformaldehyde and preserved in absolute methanol.

Paratype: *Muséum National d'Histoire Naturelle de Paris*, MNHN-IP-2019-3418, IMBE-A081-MAI56: Collected from France, Mediterranean Sea, Marseille, Maire Island (43.2096°N, 5.3353°W), at a depth of 17-20 m by Dorian Guillemain on 2023-05-04 (MAI_56, **Supplementary Figure 1**). Specimen fixed in 4% paraformaldehyde and preserved in absolute methanol.

Other material examined: Station Marine d'Endoume, IMBE: 23 additional specimens collected from various locations across Marseille Bay, including 3PP (43.09°N, 05.36°W), Maïre Island (43.2096°N, 5.3353°W), Mugel (43.1001°N, 5.3618°W), Tiboulen du Frioul (43.2799°N, 5.2860°W), and Vesse (43.2026°N, 5.1539°W) (**Supplementary Table 1** and **Supplementary Figure 1**). Specimen fixed in 4% paraformaldehyde and preserved in absolute methanol.

Molecular diagnosis: Mitochondrial markers were found to differentiate *Oscarella veronicae* sp. nov. from related species, though they did not exhibit mutations exclusive to this species. However, comparison of partial mitochondrial genomes between *O. lobularis*, *O. tuberculata*, *O. veronicae* sp. nov., and *O. sabinae* sp. nov. identified a unique mutation specific to *O. veronicae* sp. nov. ($A_{7562} \rightarrow G$). This mutation was validated through PCR amplification across various specimens from each species (**Supplementary Figure 4**). In contrast, nuclear data provided a clearer distinction, with *epic12* haplotypes f and e both revealing four *O. veronicae* sp. nov. specific mutations despite the short length of the marker ($T_{29} \rightarrow C$, $G_{33} \rightarrow A$, $A_{38} \rightarrow G$, and $T48 \rightarrow C$) (**Supplementary Figure 2**).

Morphological diagnosis: Many specimens exhibited a cauliflower-like morphology closely resembling that of *O. sabinae* sp. nov. (**Figure 3G to H'**); however, unlike the latter, they never display a bright pink coloration. Instead, most specimens exhibit a blue hue that differs from other species within the complex, appearing relatively species-specific: shades tend toward cyan, whereas *O. lobularis* and *O. tuberculata* display blues that are more grayish or purplish (**Figure 3C and F**).

Description: Like other *Oscarella* species, *O. veronicae* sp. nov. is an encrusting sponge with irregular growth forms. It covers areas up to 20-30 cm² and reaches a thickness of 10-15 mm. Most specimens display a bright or light blue coloration (**Figure 3D**), although exceptions include yellow (2 out of 25 specimens) or dark pink like *O. tuberculata* (one specimen, TIB_09, **Supplementary Figure 1**). The sponge is neither soft nor cartilaginous in consistency.

Ecology and distribution: *O. veronicae* sp. nov. is a sciaphilous species consistently found in sympatry with *O. tuberculata* and *O. lobularis*. It inhabits the cavities of coralligenous walls or the entrances of submarine caves at depths ranging from 5 to 35 m. To date, no data are available to confirm the presence or absence of this species elsewhere in the Mediterranean Sea or along the French Atlantic coasts.

CONCLUSION

This study highlights the importance of integrating complementary genetic data and standardized methodologies in the identification and taxonomy of *Oscarella* species. By combining mitochondrial and nuclear markers with comprehensive photographic documentation, we refined species delimitation within this skeleton-less genus. Our results revealed four new species in the Bay of Marseille, underscoring both the hidden diversity of *Oscarella* and the evolutionary significance of color polymorphism. At the same time, we exposed major limitations in current practices, including the low resolution of the cytochrome c oxidase subunit I (cox1/COI) gene and the lack of genetic data for many accepted species. Addressing these gaps will require coordinated efforts to generate robust genetic frameworks, which are essential not only for reliable species identification but also for clarifying biogeographic patterns and evolutionary processes. Ultimately, advancing a standardized, data-rich taxonomy of *Oscarella* will strengthen our ability to assess, conserve, and understand marine biodiversity.

ADDITIONAL INFORMATION

Authors declare no competing financial or non-financial interests in relation to the work described.

FUNDING

This work was supported by the Mediterranean Institute of Biodiversity and Marine and Continental Ecology (IMBE) through the AOI C-POSCA and the French National Research Agency (ANR) through the JCJC program I_SEA_RAINBOWS (ANR-25-CE02-1622-01).

ACKNOWLEDGEMENTS

Data were generated through the Molecular and Cell Biology Facility of IMBE, Aix-Marseille Univ, Avignon Univ, CNRS, IRD, Marseille. We are grateful to the OSU Pytheas diving facility for their support in *Oscarella* sampling. We also warmly thank A. Chenuil-Maurel and N. Boury Esnault for insightful guidance on research design and discussions, and A. Baumel for helpful input on manuscript preparation.

DATA AVAILABILITY

The genetic datasets generated and analyzed during the current study are available in the NCBI repository, https://www.ncbi.nlm.nih.gov/. All accession numbers are provided in **Supplementary Table 1**. All other data generated during this study (pictures of animals) are included in this published article (and its Supplementary Information files).

REFERENCES

1. Luypaert, T., McCarthy, M. L., Hagan, J. & Poti, M. Status of Marine Biodiversity in the Anthropocene. in *YOUMARES 9 - The Oceans: Our Research, Our Future* (eds Jungblut, S., Liebich, V. & Bode-Dalby, M.) 57–82 (Springer Nature Switzerland AG, 2020).

548 2. Miglietta, M. P. & Pruski, S. Cryptic species in time and space: an assessment of cryptic diversity

549   within eight nominal species of Hydrozoa (Cnidaria). *Proc. Biol. Sci.* **290**, 20230851 (2023).

550 3. Stöhr, S., Weber, A. A.-T., Boissin, E. & Chenuil, A. Resolving the Ophioderma longicauda

551   (Echinodermata: Ophiuroidea) cryptic species complex: five sisters, three of them new –

552   Corrigendum. *Eur. J. Taxon.*

553   https://europeanjournaloftaxonomy.eu/index.php/ejt/article/view/863 (2020).

554 4. de Paula, T. S., Zilberberg, C., Hajdu, E. & Lôbo-Hajdu, G. Morphology and molecules on opposite

555   sides of the diversity gradient: Four cryptic species of the *Cliona celata* (Porifera, Demospongiae)

556   complex in South America revealed by mitochondrial and nuclear markers. *Mol. Phylogenet. Evol.*

557   **62**, 529–541 (2012).

558 5. Caputi, L. *et al.* Cryptic speciation in a model invertebrate chordate. *Proc. Natl. Acad. Sci.* **104**,

559   9364–9369 (2007).

560 6. Knapp, I. S., Forsman, Z. H., Williams, G. J., Toonen, R. J. & Bell, J. J. Cryptic species obscure

561   introduction pathway of the blue Caribbean sponge (Haliclona (Soestella) caerulea), (order:

562   Haplosclerida) to Palmyra Atoll, Central Pacific. *PeerJ* **3**, e1170 (2015).

563 7. Muricy, G., Boury-Esnault, N., Bezac, C. & Vacelet, J. Cytological evidence for cryptic speciation in

564   Mediterranean Oscarella species (Porifera, Homoscleromorpha). *Can J Zool* **74**, 881–896 (1996).

565 8. Soest, R. W. M. V. *et al.* Global Diversity of Sponges (Porifera). *PLoS ONE* **7**, e35105 (2012).

566 9. Çinar, M. E. *et al.* Coralligenous assemblages along their geographical distribution: Testing of

567   concepts and implications for management. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **30**, 1578–1594

568   (2020).

569 10. Schmidt, O. *Die Spongien Des Adriatischen Meeres*. (Wilhelm Engelmann, Leipzig, 1862).

570 11. Idan, T., Shefer, S., Feldstein, T. & Ilan, M. New discoveries in Eastern Mediterranean mesophotic

571   sponge grounds: updated checklist and description of three novel sponge species. *Mediterr. Mar.*

572   *Sci.* **22**, 270–284 (2021).

12. Marra, M. V. *et al.* Long-term turnover of the sponge fauna in Faro Lake (North-East Sicily,

Mediterranean Sea). *Ital. J. Zool.* **83**, 579–588 (2016).

13. Aloui-Bejaoui, N. & Afli, A. Functional diversity of the macro-invertebrate community in the port

area of Kerkennah Islands (Tunisia). *Mediterr. Mar. Sci.* **13**, 93–102 (2012).

14. Voultsiadou, E. Sponge diversity in the Aegean Sea: Check list and new information. *Ital. J. Zool.*

72, 53–64 (2005).

15. Kefalas, E., Castritsi-Catharios, J. & Miliou, H. The impacts of scallop dredging on sponge

assemblages in the Gulf of Kalloni (Aegean Sea, northeastern Mediterranean). *ICES J. Mar. Sci.* **2**,

402–410 (2003).

16. Mustapha, K. B., Zarrouk, S., Souissi, A. & Abed, A. E. Diversité des demosponges Tunisiennes.

INSTM Bull. Mar. Freshw. Sci. **30**, 55–78 (2003).

17. Cruz Simó, T. *Esponjas Marinas de Canarias*. (Consejería de Política Territorial y Medio Ambiente,

2002).

18. Boury-Esnault, N., Harmelin, J.-G., Ledoyer, M., Saldanha, L. & Zibrowius, H. Peuplement

benthique des grottes sous-marines de Sagres (Portugal, Atlantique Nord-Oriental). *Bol Mus Mun*

Funchal **6**, 15–38 (2001).

19. Corriero, G., Abbiati, M. & Santangelo, G. Sponges Inhabiting a Mediterranean Red Coral

Population. *Mar. Ecol.* **18**, 147–155 (1997).

20. Carballo, J. L. & Gracía-Gómez. Esponjas del Estrecho de Gibraltar y áreas próximas, con nuevas

aportaciones para la fauna iberica. *Cah Biol Mar* **35**, 193–211 (1994).

21. Soest, R. Affinities of the marine Demosponge fauna of the Cape Verde Islands and tropical West

Africa. *CFS Cour. Forschungsinstitut Senckenberg*

https://www.semanticscholar.org/paper/Affinities-of-the-marine-Demosponge-fauna-of-the-

Soest/94d53187448f2d2fe26a07b9ae1f41bc254cad37 (1993).

22. Boury-Esnault, N. & Lopes, M. T. Les démosponges littorales de l'archipel des Açores. *An. Inst.*

Océan. **61**, 149–225 (1985).

23. Pulitzer-Finali, G. A collection of Mediterranean Demospongiae (Porifera) with, in appendix, a list

of the Demospongiae hitherto recorded from the Mediterranean Sea. 'Annali del Museo civico di

storia naturale Giacomo Doria'. *Ann. Mus. Civ. Storia Nat. Giacomo Doria* **84**, 445–621 (1982).

24. Vacelet, J. *Inventaire de Spongiaires Du Parc National de Port-Cros (Var)*. 167–186 (1976).

25. Pouliquen, L. Les spongiaires des grottes sous-marines de la region de Marseille. Ecologie et

systematique. *Tethys* **3**, 717–758 (1972).

26. Boury-Esnault, N. Spongiaires de la zone rocheuse littorale de Banyuls-Sur-Mer - I. Ecologie et

répartition. *Vie Milieu* 159 (1971).

27. Descatoire, A. Peuplements sessiles de l'Archipel de Glénan. I. Inventaire spongiaires. *Vie Milieu*

177–209 (1969).

28. Borojevic, R., Cabioch, L. & Lévi, C. *Inventaire de La Faune Marine de Roscoff : Spongiaires*.

(Éditions de la station biologique de Roscoff, 1968).

29. Sara, M. Contributo alla conoscenza dei Poriferi del Mar Ligure. *Ann. Mus. Civ. Storia Nat. Genova*

70, 207–244 (1958).

30. Hernández, F. F. *Esponjas del litoral Asturias*. (Junta para Ampliación de Estudios e Investigaciones

Cientificas, 1918).

31. de Voogd, N. J. *et al.* World Porifera Database. https://doi.org/doi:10.14284/359.

32. Pérez, T. *et al.* Oscarella balibaloi, a new sponge species (Homoscleromorpha: Plakinidae) from

the Western Mediterranean Sea: cytological description, reproductive cycle and ecology. *Mar.*

Ecol. **32**, 174–187 (2011).

33. Boury-Esnault, N., Muricy, G., Gallissian, M.-F. & Vacelet, J. Sponges without skeleton: A new

Mediterranean genus of Homoscleromorpha (Porifera, Demospongiae). *Ophelia* **4**, 25–43 (1995).

34. Łukowiak, M. *et al.* The terminology of sponge spicules. *J Morphol* **283**, 1517–1545 (2022).

35. Gazave, E. *et al.* Systematics and molecular phylogeny of the family oscarellidae

(homoscleromorpha) with description of two new oscarella species. *PloS One* **8**, e63976 (2013).

36. Stillitani, D. *et al.* Solving a taxonomic puzzle: integrative taxonomy reveals new cryptic and

polymorphic species of Oscarella in south-eastern Brazil (Homoscleromorpha : Oscarellidae).

Invertebr. Syst. **36**, 714–750 (2022).

37. Ruiz, C., Ereskovsky, A. & Perez, T. New skeleton-less homoscleromorphs (Porifera,

Homoscleromorpha) from the Caribbean Sea: exceptions to rules are definitely common in

sponge taxonomy. *Zootaxa* **5200**, 128–148 (2022).

38. Perez, T. & Ruiz, C. Description of the first Caribbean Oscarellidae (Porifera: Homoscleromorpha).

Zootaxa **4369**, 501–514 (2018).

39. Erpenbeck, D., Hooper, J. N. A. & Wörheide, G. CO1 phylogenies in diploblasts and the 'Barcoding

of Life'— are we sequencing a suboptimal partition? *Mol. Ecol. Notes* **6**, 550–553 (2006).

40. Folmer, O., Black, M., Hoeh, W., Lutz, R. & Vrijenhoek, R. DNA primers for amplification of

mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mol. Mar.*

Biol. Biotechnol. **3**, 294–299 (1994).

41. Steenwyk, J. L., Li, Y., Zhou, X., Shen, X.-X. & Rokas, A. Incongruence in the phylogenomics era.

Nat. Rev. Genet. **24**, 834–850 (2023).

42. Gérard, K. *et al.* PCR survey of 50 introns in animals: cross-amplification of homologous EPIC loci

in eight non-bilaterian, protostome and deuterostome phyla. *Mar. Genomics* **12**, 1–8 (2013).

43. Chenuil, A. *et al.* An efficient method to find potentially universal population genetic markers,

applied to metazoans. *BMC Evol. Biol.* **10**, 276 (2010).

44. Schenkelaars, Q., Perez-Cortes, D., Perruchoud, C. & Galliot, B. The polymorphism of Hydra

microsatellite sequences provides strain-specific signatures. *PloS One* **15**, e0230547 (2020).

45. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput.

Nucleic Acids Res. **32**, 1792–1797 (2004).

46. Leigh, J. W. & Bryant, D. POPART: Full-feature software for haplotype network construction.

Methods Ecol. Evol. **6**, 1110–1116 (2015).

47. Talavera, G. & Castresana, J. Improvement of phylogenies after removing divergent and

ambiguously aligned blocks from protein sequence alignments. *Syst. Biol.* **56**, 564–577 (2007).

48. Posada, D. jModelTest: phylogenetic model averaging. *Mol. Biol. Evol.* **25**, 1253–1256 (2008).

49. Tamura, K., Stecher, G. & Kumar, S. MEGA11: Molecular Evolutionary Genetics Analysis Version

11. *Mol. Biol. Evol.* **38**, 3022–3027 (2021).

50. Krabberød, A. K., Stokke, E., Thoen, E., Skrede, I. & Kauserud, H. The Ribosomal Operon Database:

A Full-Length rDNA Operon Database Derived From Genome Assemblies. *Mol. Ecol. Resour.* **25**,

e14031 (2025).

51. Ivanišević, J., Thomas, O. P., Lejeusne, C., Chevaldonné, P. & Pérez, T. Metabolic fingerprinting as

an indicator of biodiversity: towards understanding inter-specific relationships among

Homoscleromorpha sponges. *Metabolomics* **7**, 289–304 (2011).

52. Serite, C. P. *et al.* Limitations of DNA barcoding in determining the origin of smuggled seahorses

and pipefishes. *Forensic Sci. Int. Anim. Environ.* **1**, 100006 (2021).

53. Chee, S. Y. Limitations of cytochrome oxidase I for the barcoding of Neritidae (Mollusca:

Gastropoda) as revealed by Bayesian analysis. *Genet. Mol. Res. GMR* **14**, 5677–5684 (2015).

54. Souza, M. R. de & Dawson, M. N. Redescription of Mastigias papua (Scyphozoa, Rhizostomeae)

with designation of a neotype and recognition of two additional species. *Zootaxa* **4457**, 520–536

(2018).

55. Freyhof, J., Özuluğ, M. & Saç, G. Neotype designation of Aphanius iconii, first reviser action to

stabilise the usage of A. fontinalis and A. meridionalis and comments on the family group names

of fishes placed in Cyprinodontidae (Teleostei: Cyprinodontiformes). *Zootaxa* **4294**, 573–585

(2017).

56. Schmidt, O. *Die Spongien Der Küste von Algier. Mit Nachträgen Zu Den Spongien Des Adriatischen*

Meeres (Drittes Supplement). (Wilhelm Engelmann, Leipzig, 1868).

57. Boury-Esnault, N., Sole-Cava, A. M. & Thorpe, J. P. Genetic and cytological divergence between

colour morphs of the Mediterranean sponge Oscarella lobularis Schmidt (Porifera,

Demospongiae, Oscarellidae). *J. Nat. Hist.* **26**, 271–284 (1992).

58. Harmelin, J.-G., Boury-Esnault, N. & Fichez, R. *Peuplement de La Grotte Sous-Marine de l'île de*

Bagaud (Parc National de Port-Cros, France, Méditerranée). 117–134 https://www.portcros-

parcnational.fr/fr/rapports-scientifiques/peuplement-de-la-grotte-sous-marine-de-lile-de-

bagaud-parc-national-de-port (2003).

59. Knittweis, L., Ereskovsky, A. V., Schembri, P. J., Borg, J. A. & Chevaldonne, P. A preliminary survey

of marine cave habitats in the Maltese Islands. https://doi.org/10.7423/XJENZA.2015.2.07 (2015)

doi:10.7423/XJENZA.2015.2.07.

60. Gerovasileiou, V. & Voultsiadou, E. Sponge diversity gradients in marine caves of the eastern

Mediterranean. *J. Mar. Biol. Assoc. U. K.* **96**, 407–416 (2016).

61. Idan, T. *et al.* Shedding light on an East-Mediterranean mesophotic sponge ground community

and the regional sponge fauna. *Mediterr. Mar. Sci.* **19**, 84–106 (2018).

62. Ereskovsky, A. V., Richter, D. J., Lavrov, D. V., Schippers, K. J. & Nichols, S. A. Transcriptome

sequencing and delimitation of sympatric Oscarella species (O. carmela and O. pearsei sp. nov)

from California, USA. *PloS One* **12**, e0183002 (2017).

63. Ereskovsky, A., Sanamyan, K. & Vishnyakov, A. A new species of the genus Oscarella (Porifera:

Homosclerophorida: Plakinidae) from the North-West Pacific. *Cah. Biol. Mar.* **50**, 369–381 (2009).

64. Ereskovsky, A. V. A new species of Oscarella (Demospongiae: Plakinidae) from the Western Sea of

Japan. *Zootaxa* **1376**, 37–51 (2006).

65. Muricy, G. & Pearse, J. A New Species of Oscarella (Demospongiae: Plakinidae) from California. in

(2004).

66. Bandaranayake, W. M. The nature and role of pigments of marine invertebrates. *Nat. Prod. Rep.*

23, 223–255 (2006).

67. Bagnara, J. T., Fernandez, P. J. & Fujii, R. On the blue coloration of vertebrates. *Pigment Cell Res.*

20, 14–26 (2007).

68. Alappat, B. & Alappat, J. Anthocyanin Pigments: Beyond Aesthetics. *Mol. Basel Switz.* **25**, 5500

(2020).

69. Tamayo-Nuñez, J. *et al.* aeBlue Chromoprotein Color is Temperature Dependent. *Protein Pept.*

Lett. **27**, 74–84 (2020).

70. Lukyanov, K. A. *et al.* Natural animal coloration can Be determined by a nonfluorescent green

fluorescent protein homolog. *J. Biol. Chem.* **275**, 25879–25882 (2000).

71. Marhamati, Z., Marhamatizadeh, M. H. & Mohebbi, G. Evaluation of the Physicochemical,

Antioxidant, and Antibacterial Properties of Tunichrome Released from Phallusia nigra Persian

Gulf Marine Tunicate. *J. Food Qual.* **2021**, 5513717 (2021).

709
710
711

712   **AUTHOR CONTRIBUTIONS**
713   DG conducted the majority of the sampling. QS captured the images. EG and QS prepared the samples
714   and performed all gene amplifications. QS carried out phylogenetic analysis and constructed haplotype
715   networks. QS, ER, CB, and EG contributed to writing the paper. QS supervised the entire project.
716