C9orf72 expansion creates the unstable folate-sensitive fragile site FRA9A

Mila Mirceta[1,2], Monika H.M. Schmidt[1,2], Natalie Shum[1,2], Tanya K. Prasolava[1], Bryanna Meikle[1,2], Stella Lanni[1], Mohiuddin Mohiuddin[1], Paul M. Mckeever[3], Ming Zhang[3-6], Minggao Liang[1,2], Ilse van der Werf[7], Stefaan Scheers[7], Patrick A. Dion[8,9], Peixiang Wang[1], Michael D. Wilson[1,2], Theresa Abell[10], Elliot A. Philips[10], Łukasz J. Sznajder[11,12], Maurice S. Swanson[11], Mustafa Mehkary[1,2], Mahreen Khan[1,2], Katsuyuki Yokoi[1], Christine Jung[13], Pieter J. de Jong[13], Catherine H. Freudenreich[10], Philip McGoldrick[3], Ryan K.C. Yuen[1,2], Agessandro Abrahão[14], Julia Keith[14], Lorne Zinman[14], Janice Robertson[3], Ekaterina Rogaeva[3], Guy A. Rouleau[8,9,15], R. Frank Kooy[7], Christopher E. Pearson[1,2]*

[1] Program of Genetics & Genome Biology, Peter Gilgan Centre for Research and Learning, The Hospital for Sick Children; Toronto, Canada.

[2] Department of Molecular Genetics, University of Toronto; Toronto, Canada.

[3] Tanz Centre for Research of Neurodegenerative Diseases, University of Toronto; Toronto, Canada.

[4] The First Rehabilitation Hospital of Shanghai, Department of Medical Genetics, School of Medicine, Tongji University; Shanghai, China.

[5] Institute for Advanced Study, Tongji University; Shanghai, China.

[6] Clinical Center for Brain and Spinal Cord Research, Tongji University; Shanghai, China.

[7] Department of Medical Genetics, University of Antwerp; Edegem, Belgium.

[8] Montreal Neurological Institute, McGill University; Montreal, Canada.

[9] Department of Neurology & Neurosurgery, McGill University; Montreal, Canada.

[10] Department of Biology, Tufts University; Medford, USA.

[11] Department of Molecular Genetics and Microbiology, Center for NeuroGenetics and the Genetics Institute, College of Medicine, University of Florida; Gainesville, USA.

[12] Currently at Department of Chemistry and Biochemistry, University of Nevada, Las Vegas, NV, USA.

[13] BACPAC Resource Center, Children's Hospital Oakland Research Institute; Oakland, USA.

[14] Sunnybrook Health Sciences Centre; Toronto, Canada.

[15] Department of Human Genetics, McGill University; Montreal, Canada.

* To whom correspondence should be addressed. Tel: (647) 388-5151; Email: cepearson.sickkids@gmail.com

Abstract

The hyper-unstable Chr9p21 locus, harbouring the interferon gene cluster, oncogenes and *C9orf72,* is linked to multiple diseases. *C9orf72* (GGGGCC)n expansions (*C9orf72*Exp) are associated with incompletely penetrant amyotrophic lateral sclerosis, frontotemporal dementia and autoimmune disorders. *C9orf72*Exp patients display hyperactive cGAS-STING-linked interferon immune and DNA damage responses, but the source of immuno-stimulatory or damaged DNA is unknown. Here, we show *C9orf72*Exp in pre-symptomatic and ALS-FTD patient cells and brains cause the folate-sensitive chromosomal fragile site, FRA9A. FRA9A centers on >33kb of *C9orf72* as highly-compacted chromatin embedded in an 8.2Mb fragility zone spanning 9p21, encompassing 46 genes, making FRA9A one of the largest fragile sites. *C9orf72*Exp cells show chromosomal instability, heightened global- and Chr9p-enriched sister-chromatid exchanges, truncated-Chr9s, acentric-Chr9s and Chr9-containing micronuclei, providing endogenous sources of damaged and immunostimulatory DNA. Cells from one *C9orf72*Exp patient contained highly-rearranged FRA9A-expressing Chr9 with Chr9-wide dysregulated gene expression. Somatic *C9orf72*Exp repeat instability and chromosomal fragility are sensitive to folate-deficiency. Age-dependent repeat instability, chromosomal fragility, and chromosomal instability can be transferred to CNS and peripheral tissues of transgenic *C9orf72*Exp mice, implicating *C9orf72*Exp as the source. Our results highlight unappreciated effects of *C9orf72* expansions that trigger vitamin-sensitive chromosome fragility, adding structural variations to the disease-enriched 9p21 locus, and likely elsewhere.

INTRODUCTION

Expansions of 30-7950 GGGGCC repeats in the *C9orf72* gene, which resides in 9p21 (*C9orf72*Exp), are the most common cause of ALS, FTD, and numerous seemingly unrelated diseases including inflammation dysregulation, autoimmune diseases, over-expression of type I *IFN* genes (1–6), and melanoma (7). Most *C9orf72*Exp carriers are asymptomatic at 40-years of age, some will develop amyotrophic lateral sclerosis (ALS), frontotemporal dementia (FTD), a co-occurring disease, or a mixture of these, while many can be disease-free into their 80s-90s (8,9). This incomplete penetrance suggests the involvement of genetic and environmental modifiers and/or cumulative events.

Many studies on *C9orf72*Exp disease have focused upon the expanded repeat RNA and repeat-associated non-AUG RAN-peptides, where many model systems are devoid of the endogenous chromosomal expanded repeat. Repeat expansions downregulate *C9orf72*Exp expression leading to reduced C9ORF72 protein levels (10, 11). *St*imulator of *In*terferon *G*enes (STING), which with cGAS activates the innate immune response following sensing of cytosolic double-stranded DNA that could arise from viruses, damaged mitochondria, or damaged chromosomes leaked from micronuclei. In *C9orf72*Exp individuals, decreased C9ORF72 leads to STING-hyperactivated *IFN* expression and inflammation in vulnerable neurons (12), but the source of the immunostimulatory DNA is unknown (4, 13, 14). Accumulating evidence supports a role of an activated DNA damage response (DDR) in *C9orf72*Exp disease (15–25), yet the source of the damaged DNA is unknown. Massive post-natal inter- and intra-tissue repeat length instability is evident in post-mortem tissues of *C9orf72*Exp individuals with the largest somatic expansions in the brain (26–30). Surprisingly, there have been no prior cytogenetic or chromosomal instability studies of *C9orf72*Exp. The recent discontinued clinical programs of antisense oligonucleotides targeting the sense-strand

C9orf72 mRNA, which successfully reduced the RAN-peptides, but resulted in no clinical

benefit or trended toward greater clinical decline for the cohort, highlight our poor

understanding of *C9orf72*Exp (31–36). A recent meeting focused on the post-trials path

forward for *C9orf72*Exp highlighted key areas needing attention: *"Critical information in our*

understanding of the biology around the C9orf72 expansion is still lacking, including what

molecular and cellular changes occur at presymptomatic stages…studies of genetics and

lifestyles might be useful" (31). Unknown biology of the *C9orf72*Exp has been highlighted as

a focus arena (31–36). Factors that may determine disease penetrance, clinical variation,

disease onset, progression and severity, are indispensable for family life decisions as well as

clinical trial design (6, 9, 31, 37–41).

Genomic context is important, as *C9orf72*Exp occurs on a risk haplotype, which could

influence repeat length instability, leading to reduced transcription and its retention of the

intron containing the expanded GGGGCC repeat (42–45) (Fig.1A). Many of the co-occurring

*C9orf72*Exp symptoms can arise independent of a *C9orf72*Exp and are associated with other

9p21 genes. For example, the type I interferon (*IFN*) cluster of 17 genes, including *IFNk*

adjacent to *C9orf72*, are linked to autoimmune diseases (e.g., dysregulated inflammation,

lupus, rheumatoid arthritis, diabetes, multiple sclerosis) (1, 3–5, 7). The enhancer-rich 9p21

gene desert is associated with *IFN* signaling (46, 47). *CDKN2A/B*, *EMICERI*, and *IFN*s are

linked to melanoma (48), a cancer recently associated with *C9orf72*Exp (7). *LINGO2*,

TOPORS, *APTX*, and *DNAJA1* have been associated with neurodevelopmental,

neurodegenerative, and motoneuron disease (49).

The 9p21 locus is hyper-prone to various recurrent structural rearrangements (50–54) (Fig.

1A). 9p deletion/duplication syndrome includes 9p21.2 as a breakpoint hotspot (55, 56).

Instability in cancers is suspected to involve chromosomal fragile sites although the hyper-

unstable 9p21 has no mapped fragile site (57). "Rare" fragile sites can be present in as few as

a single individual, as initially reported for FRA9A(58), to ≤5% of individuals and are linked

to partially penetrant neurological disorders, as reviewed (59). Most rare fragile sites are folate-

sensitive fragile sites (FSFS), induced in the absence of folate or by the antimetabolite 5-

fluorodeoxyuridine (FUdR). Ten of the ~30 FSFS have been mapped to gene-specific expanded

(CGG)n repeats, including FRAXA, FRAXE, FRAXF, FRA2A, FRA7A, FRA10A, FRA11A,

FRA11B, FRA12A, and FRA16A. FSFS are prone to DNA breaks making them mutation

hotspots — mutations that can alter clinical presentation, which is well-characterized for the

CGG-expanded FRAXA/*FMR1*, reviewed in (59). Through >75 years of study, many isolated

case reports, diverse mutation forms (varying CGG expansion sizes, contiguous gene deletions,

gene duplications, intra- and inter-chromosomal recombination, point mutations), and *FMR1*

mosaic epimutations have been identified as the cause of numerous seemingly unrelated

diseases, many incompletely penetrant (59). Individual case studies with deep clinical,

neuropathological and molecular phenotyping can be informative of disease etiology. Thus,

instability of FSFS can impact disease manifestation and clinical presentation. Our objective

was to better understand the biology of the endogenous expanded *C9orf72* repeat in the context

of 9p21 in presymptomatic and disease states. We assessed whether the expanded *C9orf72*

repeat, GGGGC*CGG*GGC*CGG*GGC*CGG*GGCC…, which contains the CGG-motif common

to all mapped FSFS, is a FSFS.

 Figure 1

 Figure 1. The 9p21 locus, *C9orf72*Exp cells, repeat lengths, CpG-methylation, and disease state.

(A) 9p21 locus is enriched with disease associations and is a mutation hotspot, see text. (B) ALS/FTD

families with Southern blot vertically aligned for each individual. Females are represented by circles,

males by squares, disease state is as per the insert legend. Colors of text and boxes around symbols

indicate the expansion, methylation, and affected status for each sample. Percentage of cells showing

FRA9A expression are indicated for each individual in the pedigree. "P#" refers to patient number, "E",

"M", and "A" refer to *E*xpanded, *M*ethylated, and *A*ffected, is summarized at right and this coding

system is used throughout the text. Southern blot of *C9orf72* (GGGGCC)n repeat expansion (Methods)

used *Eco*RI/*Bam*HI to release the repeat-containing fragment. (C) Schematic of Southern blot probe

location, CpG-methylation status at the GGGGCC repeat and CpG-islands (filled boxes denote

methylation). For raw CpG methylation data see Figure S2.

RESULTS

9p21 shares sequence and epigenetic features with (CGG)n FSFS

The *C9orf72* locus and each of the CGG-expanded FSFS share many sequence and epigenetic features (Fig. ED1A-D). Nine of the 10 CGG-FSFS plus *C9orf72* colocalize to boundaries between topologically associating domains and were enriched with CTCF sites and CpG-islands (Fig. ED1B-D). Thus, *C9orf72* and CGG-FSFS share with 22 other disease-associated repeat expansion loci these epigenetic features (60).

Variations in *C9orf72* repeat expansion size, CpG-methylation, and disease state permit assessment of contribution to fragility

To study the connection between *C9orf72*Exp and fragility, we characterized lymphoblastoid cells from *C9orf72*Exp families, where some carriers were presymptomatic and some presented with ALS or ALS/FTD (Fig. 1B). Sizing *C9orf72* expansions by Southern blots shows repeat length heterogeneity within and between individuals, with expansions ranging from ~300 to ~4400 repeats (Fig. 1B-C, Table ED1). All individuals carried (GGGGCC)2 on the non-expanded allele.

Most of our cell lines were CpG methylated in the expanded repeat, with varying levels in flanking CpG-islands, and no methylation in the non-expanded allele (Fig. 1C; Fig. ED2), consistent with previous findings of aberrant methylation and levels of methylation mosaicism in *C9orf72*Exp carriers (61, 62). We found an absence of methylation, likely arising from methylation mosaicism, in cells of a presymptomatic *C9orf72*Exp (P1) and ALS-affected individuals (P7 & P8). No obvious differences were observed between cells from presymptomatic and affected *C9orf72*Exp individuals. Methylation status was stable, consistent with longitudinally collected samples from *C9orf72*Exp carriers (63). Control cells from individuals without *C9orf72*Exp had no methylation at *C9orf72*. Thus, our *C9orf72*Exp

cell line collection permitted assessing fragility in the presence/absence of methylation and

with/without disease of donors (Fig. 1B, see right).

C9orf72 expansion is the molecular cause of a FSFS at 9p21 (FRA9A)

Fragility was assessed cytogenetically (Fig. 2A-D). FUdR-induced fragile sites were observed

and localized to 9p21 by fluorescent _in situ_ hybridization (FISH), in all eight _C9orf72_Exp cell

lines. Fragility was only observed in _C9orf72_Exp cells, with FUdR-treatment and only on one

Chr9, consistent with FRA9A (58). Fragility presented in various forms including gaps

(isochromatid and chromatid), constrictions, and despiralized regions (Fig. 2A). Full

metaphase spreads and zoomed-in versions, with greater resolution are available upon request.

With the exception of the despiralized form, these are similar to forms of FRAXA (59). The

despiralized Chr9 was similar to the despiralized heterochromatin in immunodeficiency,

centromeric instability, and facial anomalies (ICF) syndrome (64). Lengthening of the

despiralized FRA9A is evident when compared to the non-fragile Chr9 from the same

metaphase spread (Fig. 2A, control Chr9 in lower left is from the same metaphase as the

leftmost stretched FRA9A). These cytogenetic observations reveal a fragile site, FRA9A, at

9p21.

We next quantified fragility. Established FXS/FRAXA diagnostic guidelines (65, 66) dictate

that fragile site frequencies of 3% or 5% within 100 metaphases yields confidence levels of

95% or 99%. Typically, FXS individuals show fragility at 3-15%, but never >50% of CGG

expanded ChrXs (66, 67). We observed 9p21 fragility at 7-14% across _C9orf72_Exp cell lines

(Fig. 2E). Fragility was evident in _C9orf72_Exp cells of presymptomatic and ALS/FTD-affected

individuals. Chromosomal assignment and band location of fragility was confirmed by

Giemsa-banding, trypsinization, and karyotyping (Fig. ED3). Fragility at 9p21 was observed

on only one of the Chr9 homologs in a given metaphase of any one _C9orf72_Exp cell line,

consistent with each being heterozygously expanded. Fragility at 9p21 was only detected with

FUdR treatment. Similar levels of fragility arose using folate-free media (M199) as for FUdR,

revealing induction by different methods of folate stress (Fig. 2C, Fig. 2E). Fragility could not

be induced without *C9orf72* expansions in two control cell lines (100 metaphases each). Thus,

fragility depends on folate perturbation and the presence of the expanded repeat. As few as 300

repeats (P3, 9%) could express FRA9A at similar levels as 4400 repeats (P4, 8%). The localized

fragility, limited to one Chr9, in *C9orf72*Exp cells is consistent with previous FS mapping to

autosomes that concluded fragility arises on the expanded allele (68, 69). We validated that the

C9orf72 repeat expansion is the molecular cause of FRA9A fragility through gene-transfer

experiments (below).

We further mapped fragility using seven FISH probes spanning the outer edges of 9p21, in

eight *C9orf72*Exp cell lines, >100 metaphases each, with probe signals being telomeric,

spanning, or centromeric to the break (Fig. 2B, 2D & 2F). Breaks were detected across 9p21.1-

9p21.3 chromosomal bands, ~8.2Mb, with ≥83% of breaks at a ~1Mb stretch of 9p21.2

encompassing *C9orf72* (Fig. 2D). Splitting of the FISH signal to both sides of the break was

evident for the GGGGCC-containing probe 4, supporting that the expanded repeat causes

fragility (Fig. 2C). Probes telomeric to *C9orf72* preferentially fluoresced telomeric of the break,

with the same trend for centromeric probes (Fig. 2F). The telomeric boundary could be

considered at probe 2, which straddles 9p21.2-9p21.3, with fluorescence both telomeric and

spanning fragility. A centromeric boundary could not be defined, as probe 7 was detected

telomeric, spanning, and centromeric to breaks, suggesting fragility extended centromeric of

probe 7. Additional centromeric probes, being pericentromeric, hampered break detection.

Thus, within the limits of FISH resolution, FRA9A maps most intensely to *C9orf72*, but has a

wide zone of breakage (summarized in Fig. 2D). FRA9A's fragility core of enriched breakage

includes *C9orf72* covering ~500kb (plus the repeat), spanning the *C9orf72*Exp risk haplotype,

where breaks can extend ~2Mb and ~6.2Mb telomerically and centromerically (Table ED2).

This makes FRA9A one of the largest fragile sites, spanning ~8.2Mb over 9p21.1-21.3,

encompassing 46 genes (Fig. 1A), similar in size to FRA1H, FRA3B, FRA4F, FRA7B,

FRA9E, FRA11G, and FRAXF, spanning up to 12.2Mb over multiple chromosome bands,

encompassing 10-110 genes (Table ED2) (70–73). Moreover, CpG-methylation of

*C9orf72*Exp was not required, as fragility was observed in cells +/- methylation (Fig. 2E).

Thus, neither symptomatic state of blood donors nor CpG-methylation were required for

FRA9A expression. That we detected FRA9A fragility at 9p21 in *C9orf72*Exp, lymphoblastoid

cells, confirms previous observations that FRAXA expression was similarly detected

lymphoblastoid cells as in freshly-collected peripheral blood lymphocyte cultures (74). Thus,

FRA9A and FRAXA are distinct from the fragility at 11q23, which depends upon the EBV

encoded protein EBNA1 (75–78).

 Figure 2

Figure 2. *C9orf72*Exp causes FRA9A folate-sensitive fragile site that maps across 9p21.

(A) Fragile site forms (indicated), with examples of non-fragile chromosomes from the same

metaphase spreads. Showing DAPI-stained and paired FISH-probed chromosomes from FUdR-treated

cells. 9q-arm FISH probe (9q32~34.11; RP11-696J10) in green. FISH probe #2 at *C9orf72* (panel D-F)

in red. Magnification is the same for all chromosome images within each panel, facilitating comparison

within the panel. Control Chr9 in upper left of panel A is from the same metaphase as the leftmost

isochromatid Chr9. Full metaphase spreads can be provided upon request. Notably, the control "No

fragile site" Chr9 in lower left of panel A is from the same metaphase as the leftmost

"Despiralized/stretched" Chr9, revealing the lengthening of the stretched Chr9. (B) Representative

example of the same chromosome (same magnification) with or without indicated FISH probes. (C)

Two representative examples (same magnification) of isochromatids with FISH signal of *C9orf72*-

containing probe 4 split over the fragile site. (D) 9p21 location of FISH probes with regional summary

of fragile site location. (E) Quantification of fragility for ≥100 metaphases under two folate-stressed

conditions. (D) Cells were described in Figure 1. (F) Fragile site breakage regions calculated from

number of breaks relative to FISH signal found centromeric, telomeric, or spanning across probes used.

Percentage of telomerically located FISH signal relative to the break is on the bottom row. Frequencies

of probes found spanning or on each side the break were used to calculate the percent breakage within

each region in panel D.

206 ***C9orf72*Exp allele assumes a compact chromatin conformation, which is enhanced by**

207 **expansion size and CpG-methylation**

208 Since constricted fragile sites on metaphase chromosomes may be the result of localized

209 unusual chromatin condensation, we next determined whether the mutant *C9orf72*Exp locus

210 assumes an unusual chromatinized state relative to the non-mutant allele using micrococcal

211 nuclease (MNase) accessibility, which is widely used as an indication of regional chromatin

212 compaction. We were able to electrophoretically resolve MNase accessibility on the mutant

213 from the non-mutant alleles by Southern blotting. MNase can directly reveal both inaccessible

214 and accessible regions for each allele, while sequencing-based methods of chromatin

215 accessibility, like ATAC-seq, must impute inaccessibility and cannot resolve alleles.

216 Permeabilized cells were exposed to MNase, which preferentially cuts DNA between

217 nucleosomes; DNA was de-proteinized, isolated, restriction digested releasing the repeat-

218 containing fragment, electrophoretically resolved, and detected by Southern blotting (Fig. 3A-

219 B, blue probe). Over-digestion by MNase to mono-, di-, and tri-nucleosome sized DNA

220 fragments was evident by ethidium staining (Fig. 3C, left panel). A striking resistance to

221 MNase accessibility is seen exclusively for the *C9orf72*Exp allele, contrasting the non-

222 expanded allele, which serves as an internal equimolar control, completely degraded at low

223 MNase concentrations (Fig. 3C, compare lanes 4 & 5). The expanded allele resisted digestion

224 at much higher MNase concentrations (Fig. 3, compare lanes 5-10). MNase resistance extended

225 beyond the expansion into the flanking regions of the mutant allele. Increasing concentrations

226 of MNase (>50 Units) led to partial and progressive degradation of the expanded allele to a

227 distinct shorter size, losing ~1 kb of flanks from the full-length *Eco*RI-*Bam*HI restriction

228 fragment, which subsequently digested the probed region (Fig. 3C, schematic). Conversely, on

229 the non-mutant allele, the same flanking regions and non-expanded repeat were completely

digested at low MNase concentrations, along with the rest of the genome (<10 Units, Fig. 3C,

lanes 5-10). The poor digestion of the *C9orf72*Exp mutant allele relative to the internal

sequence control of the equimolar non-mutant allele, strongly suggests the formation of an

unusual chromatinized state of the mutant *C9orf72* allele. Control experiments confirmed that

the inaccessibility of MNase to the *C9orf72*Exp allele was due to its chromatinized state, and

not due to an inherent inability of MNase to cleave the expanded non-chromatinized

(GGGGCC)n sequence (Fig. ED4A & ED4B, Methods). Six cell lines showed MNase

resistance of the *C9orf72*Exp allele, regardless of presymptomatic or symptomatic state of the

blood donors (Fig. 3C-E).

Mapping the boundaries of MNase-resistance using multi-probe and multi-restriction

digests revealed the extent and distribution of MNase inaccessibility over the *C9orf72*Exp

allele. Maximal MNase resistance spanned a total of >33kb for an allele with 4400 repeats (P4),

with MNase inaccessibility extending ~3300bp upstream and ~3700bp downstream of the

repeat (summarized in Fig. 3A to scale, Fig. ED4C, see Methods). Mapping in six *C9orf72*Exp

cell lines showed little variation of the absolute boundaries of MNase inaccessibility with

expansion size or presence of disease. Together, these data confirm that the *C9orf72*Exp allele

assumes a large region of compact chromatin. Maximal MNase inaccessibility covers the

upstream CpG-island, the *C9orf72* promoter, transcription start sites (sense and anti-sense), the

expanded repeat, cryptic splice sites C1-4, the downstream CpG-island, beyond exon 2, and

falls within FRA9A's fragility core.

Repeat length and CpG-methylation enhanced MNase-resistance. Analysis of a cell line that

harbored a mosaic heterogenous mixture of expansion sizes ranging between ~300–4400 repeat

units demonstrated that repeat expansion size directly correlates with MNase resistance (Fig.

3D, P4). Longer expansions were more resistant to MNase than shorter expansions, which were

more resistant than the non-expanded alleles (Fig. 3D, lanes 3 & 4, quantified in Fig. 3D). CpG-

methylation exacerbated inaccessibility. Two cell lines with similar expansion sizes (~770

repeats) with or without aberrant methylation at the mutant *C9orf72* (P6 [isolated single cell

clone] and P7, respectively) showed that methylation further enhanced MNase resistance of the

expanded allele (Fig. 3E, ED4A-iii). Larger expansions and methylation enhance chromatin

compaction of the *C9orf72* locus.

We assessed chromatin compaction of post-mortem brains of two *C9orf72*Exp individuals,

previously deep-phenotyped for lifestyle choices, clinical, neuropathological, and molecular

biomarkers (29, 79). One was asymptomatic for ALS or FTD, with a distinct ~70 repeats in

blood, no signs of neurodegeneration, overexpression of *C9orf72*/C9ORF72, and devoid of

TDP-43 neuropathology through to the age of 90-years. The other was his daughter with 300-

3000 repeats in blood who had ALS at age 57 and died at 59 and autopsy showing

neurodegeneration and TDP-43 inclusions (29) (summarized in Table ED3). The asymptomatic

father had a strikingly heterogenous range of 13-3500 and 13-1500 repeats in the frontal cortex

and cerebellum, respectively, with detection of CpG-methylation likely triggered by large (>70

repeats) expansions (Fig. 3F, lanes 0). In contrast, the affected daughter showed very large

distinct expansions of mostly ~4400 and ~1600 CpG-methylated repeats in the frontal cortex

and cerebellum, respectively, where the intensity of the expansions matched that of the non-

expanded allele (Fig. 3F, lanes 0). Chromatin was MNase inaccessible on only the expanded

C9orf72 alleles in both brain regions of both individuals, while the non-expanded (GGGGCC)2

allele was rapidly digested by MNase (Fig. 3F). MNase inaccessibility for both individuals

appeared more severe in the cerebellum relative to the frontal cortex, which contrasts the

expectation that the longer expansions in the frontal cortex would make it more MNase

resistant, but may reflect the longer expansions and/or the increased levels of deacetylated and

trimethylated histone modifications in the frontal cortex over the cerebellum, relative to the

acetylated histones on the non-expanded allele of *C9orf72*Exp carriers (80). The shorter

expansions in the father's brain were more MNase sensitive than the larger expansions.

Quantifying MNase inaccessibility between asymptomatic and symptomatic brains is

hampered by the repeat length variations they presented, which we have shown to affect

chromatin accessibility (Fig. 3D). Expression of the *C9orf72* transcript in the asymptomatic

*C9orf72*Exp father was dramatically increased 2.5-7-fold in the frontal cortex, cerebellum,

spinal cord, blood, and four other tissues compared to the affected daughter's tissues, whose

C9orf72 levels were twice less than an individual devoid of expansions (29, 79). We suggest

that the increased transcription in the father's brain arose from the shorter range of expansion

sizes (13-1200 repeats) which were less MNase resistant than the longer expansions, which is

consistent with both the over expression in the father's blood, which contained predominantly

70 repeats, and with the low transcription in the daughter's brain that contained only very large

expansions. This interpretation is consistent with the 2.5-5-fold increased expression of

intermediate *C9orf72* repeats in human brains, blood, and knockin iPSC-derived neural

progenitor cells (63, 81). This parallels the 5- to 10-fold increased expression from the

premutation CGG expansions at the *FMR1* repeat in hemizygous males, which also show

compact chromatin (82, 83). We conclude that *C9orf72* expansions and flanks assume a highly

compact chromatin in cells and brain regions of *C9orf72*Exp carriers. MNase resistance is

evident on *all C9orf72*Exp alleles in a given sample, which differs from the 7-14% of

*C9orf72*Exp cells cytogenetically showing FRA9A-Chr9s, like FXS, only a portion of the

CGG-expanded ChrXs show FRAXA (66, 67). In this manner, MNase resistance might be

considered a proxy for fragility. Towards assessing whether MNase resistance was linked to

the unusual chromatin compaction of cytogenetic fragility or merely linked to expanded

repeats, we assessed the MNase accessibility of the *HTT* locus in Huntington's disease patient

cells with (CAG)21/180, which is known to not be a cytogenetically detectable fragile site (84).

MNase treatment and Southern blotting resolved the mutant and non-mutant *HTT* alleles and

both showed the same MNase accessibility when most of the genome was sensitive (Fig. ED5,

lanes 3-5), suggesting that MNase inaccessibility does not extend to this non-fragile mutant

locus. As part of another study, we extended the MNase inaccessibility to the CGG-expanded

FMR1 gene the molecular cause for the fragile site FRAXA. We find MNase resistance of only

the CGG expanded allele (in cells from male and female carriers of the CGG expansion) (not

shown). That the CGG expanded *FMR1*, which forms FRAXA, but not the non-fragile CAG

expanded *HTT*, forms an MNase resitant region supports our suggestion that MNase

inaccessibility of the *C9orf72*Exp can be considered a proxy for chromosomal fragility. We

suggest that the MNase resistance of the *C9orf72*Exp locus in human brains is a reflection of

chromosomal fragility (Fig. 3F).

Figure 3

Figure 3. *C9orf72*Exp locus is MNase-inaccessible.

(A) Schematic of the region of unusual chromatin organized as determined by MNase accessibility analysis in *C9orf72*-ALS patient cells and brain regions. *C9orf72* (GGGGCC)n-N repeat, CpG-islands (hatched boxes), long-2kb and intermediate 435bp promoters (green and red arrows, respectively), exons 1a (yellow dot) and 1b (orange dot), sense and antisense transcription start sites, restriction sites with distances from the beginning and end of the (GGGGCC)N tract, and probes for upstream (green) and downstream (blue) mapping of MNase accessibility. Cryptic splice site transcripts C1-C4 derived from the *C9orf72*Exp allele are indicated with their distance from the end of the repeat tract (42, 45). (B) Workflow for MNase assay. (C) MNase treatment of chromatinized DNA from a cell line (P6) with a large methylated *C9orf72*Exp that had undergone single-cell cloning (scc) to isolate a single expanded allele, ensuring a single expansion size and facilitating clarity. Ethidium bromide (EtBr) staining of the MNase-treated DNA reveals over-digestion of the DNA to mono-, di-, tri-nucleosomal, sized fragments. The same DNA was digested with *Eco*RI/*Bam*HI, to release the repeat fragment having 697 bp upstream and 1714 bp downstream of the repeat, then assessed by Southern blotting using the blue probe. Resistance of only the expanded allele occurs, even at high concentration of MNase (200U). Red "R" indicates the region with reduced MNase accessibility that is about to lose the probed region indicated in the schematic at right. (D) Sample P3 with a mixture of heterogeneous repeat allele lengths shows that with increasing repeat size, MNase accessibility is decreased, as quantified by densitometric analysis for each allele size (colours coordinate between graph and Southern blot). In panels D-F the repeat was released with *Afl*II/*Pci*I, leaving 404 bp upstream and 689 bp downstream allowing increased resolution. NB, van Blitterswijk uses *Xba*I/*Xba*I to release the repeat, leaving 1751 bp upstream and 608 bp downstream of the repeat, the greater amount of flanking sequence reduces electrophoretic resolution. (E) Methylation is associated with increased MNase inaccessibility as assessed in cell lines (P6 (scc) and P7) containing similar sized repeats but differing methylation status. The non-expanded allele is digested at the same rate in both lines. (F) MNase treatment of post-mortem (orbito)frontal cortex and cerebellum from an asymptomatic 90-year-old father and his ALS-FTD-affected daughter, previously deep-phenotyped for lifestyle choices, clinical, neuropathological, and molecular

343   biomarkers (29, 79) (see Table ED3). MNase concentrations are doubled between each lane (2.5U in

344   lane 2 up to 40U). Expanded allele in both tissues is MNase resistant relative to the equimolar non-

345   expanded allele, which serves as an internal control. See Figure ED4 for MNase controls and mapping

346   experiments.

Increased Chr9-containing micronuclei in *C9orf72*Exp cells

Micronuclei (MN) are indicators of chromosomal double-strand breaks (DSBs) including fragile sites (85–88), where MN often harbor broken chromosomes to be eliminated or reintegrated. Nuclear buds (NBuds) are chromatin-containing protrusions linked to the nucleus by a nucleoplasmic stalk (86, 88). Both MN and NBuds can arise from chromosome breaks. NBuds can become MN (86). These subnuclear structures can be enriched with tandem gene amplifications and satellite repeats (64, 89, 90). Cytoplasmic release of MN DNA triggers an immune response via DNA sensing by cGAS-STING that activates *IFN* expression (85), a pathway hyperactivated in vulnerable neurons of *C9orf72*Exp carriers (4, 12–14). Given these connections, we assessed MN and NBuds and whether they contained Chr9 in *C9orf72*Exp cells. *C9orf72*Exp cells showed elevated MN/NBud levels compared to control cells, which were further increased with folate stress (Fig. 4A). More NBuds over MNs contained Chr9 in *C9orf72*Exp cells compared to control cells. The modest increase in MN with Chr9 could be due to loss of fragile chromosome-containing MN during folate-perturbation, as suggested for FRAXA cells (Fig. 4B) (91, 92). Depending upon whether the breakpoint is centromeric or telomeric of *C9orf72*, the presence of the *C9orf72*-containing FISH signal alone or the 9q-arm FISH signal alone, supports subnuclear inclusion of an acentric or truncated Chr9, respectively, while the presence of both *C9orf72*- and q-arm signals supports the inclusion of truncated Chr9 or the full-Chr9 (Fig. 4B). An absence of either FISH signal could reflect an absence of any Chr9, or an acentric Chr9 broken telomeric of *C9orf72*. When Chr9 signals were present in MN/NBuds, the proportion of the *C9orf72* signal alone, 9q-arm alone, or the two together occurred at 1:1:5, respectively. Increased Chr9-containing MN/NBuds were evident in *C9orf72*Exp cells of presymptomatic and symptomatic individuals. Cumulatively, these data support increased chromosomal damage in *C9orf72*Exp cells.

We also observed increased nuclear blebs (NBlebs) in *C9orf72*Exp compared to control

cells (Fig. 4A). NBlebs are chromatin-containing nuclear herniations (86, 88, 93, 94), without

an obvious constriction between the nucleus and the protruding nuclear material. NBlebs may

in some instances lead to NBuds, which may lead to MN (86). NBlebs are hallmarks of the

DNA damage accumulating progeroid syndromes and can be enriched with genomic regions

with perturbed chromatin and DNA damage (γH2AX) (93–97). In *C9orf72*Exp cells, NBlebs

contained Chr9 at higher levels than in cells without expansions (Fig. ED6A), which is

consistent with reports of NBlebs being enriched with regions having altered chromatin

compaction and damaged DNA (93–95).

 Figure 4

Figure 4. *C9orf72*Exp cells display chromosomal instability.

(A) Quantification of micronuclei (MN) and nuclear buds (NBud) in binucleated cells without FISH,

left, and with *C9orf72* and 9q-FISH probes, middle, in WT and *C9orf72*Exp cells (P6 (ssc) and P7)

containing similar sized repeats (~770) but differing methylation status. Data are means of two

independent experiments. Dots show data of each experiment. Statistical analyses using Fisher's exact

test (*P<0.05, **P<0.01, ***P<0.001). Representative images of MN and NBuds in cytokinesis-

blocked cells treated with FUdR for 24 hours, that contain no FISH signal (hollow arrows), a *C9orf72*-

signal (red arrows), or 9q-signal (green arrows). (B) Depending upon site of breakage, *C9orf72*-FISH

probes can have varying interpretations, left. Example of *bona fide* chromosome breaks in *C9orf72*Exp

line P4 with acentric terminal *C9orf72*-containing fragment in same metaphase spread as a truncated-

Chr9 that is free of *C9orf72*-signal, middle. Example of truncated-Chr9 with *C9orf72*-signal, and

control Chr9 from same spread, right. (C) Levels of global sister chromatid exchanges (SCEs) in WT

and *C9orf72*Exp cells +/-FUdR. Methylation/clinical status is as per Figure 1B. Each datapoint

represents a single metaphase. At least 100 mitotic cells were analyzed per condition in each

experiment. Statistical analyses were performed by student's t-test (P<0.01). NS: not significant. See

also Figure ED5B-E. (D) *C9orf7*Exp line P8 with complex highly rearranged derivative-Chr9,

schematic, see text and Figure ED7A. (E) Example of derivative-Chr9 and control Chr9 in same spread

with *C9orf72*- (red) and 9q-FISH probes (green) (100%=100/100 metaphases). (F) Only the derivative-

Chr9 expresses FRA9A coincident with the *C9orf72*-FISH signal, revealing derivative-Chr9 to be the

*C9orf72*Exp-Chr9, (+FUdR, 7%=7/100, 99% confidence). (G) Gene expression (binned/gene) along

Chr9 in *C9orf72* Exp cells with trisomic (red dots) and monosomic (blue dots) in duplicated and deleted

regions of the derivative-Chr9 (P8). Individual interactive html files with exact expression levels for

each gene are available (see data availability statement). See Figure ED7B-D for statistical significance

of differential gene expression.

*C9orf72*Exp and double-strand breakage

Direct support that fragile sites are prone to DNA breakage is the cytogenetic manifestation of

a truncated chromosome or acentric chromosome fragment broken at the fragile site (98, 99).

We observed truncated Chr9s broken at 9p21, where the truncated p21→pter acentric arm,

containing the *C9orf72*-FISH signal, remained within the *same* metaphase spread (Fig. 4B).

We observed numerous instances of truncated Chr9s with the *C9orf72* signal at the broken end,

in the absence of their acentric p21→pter acentric arms, which had likely been lost. 9p21-

truncated Chr9s without *C9orf72*-signal were also observed, consistent with FRA9A's broad

~8.2Mb breakage zone (Fig. 4B, see truncated Chr9 broken centromeric and telomeric of the

C9orf72-FISH signal). Our cytogenetic findings, coupled with the increased Chr9-containing

MN/NBuds, is convincing evidence that double-strand DNA breaks occur at FRA9A,

consistent with the γ-H2AX foci in cells, spinal cord, and brains of *C9orf72*Exp carriers (23,

24).

We further explored the susceptibility of GGGGCC repeats to double-strand DNA breaks

using a yeast system that quantifies rates of chromosome end-loss (100). A (GGGGCC)60

inserted into a yeast chromosome was extremely unstable, yielding mixed populations with 15-

60 repeat units (Fig. ED7A-D). Rates of breakage showed a significant length-dependent

increase: Tracts of 35-60 repeats broke 6-16-fold compared to (GGGGCC)15, which incurred

breaks at 10-fold higher rates than a (CAG)85 tract analyzed in the same assay (Fig. ED7B and

Fig. ED7C). Thus, even short GGGGCC repeats can be highly prone to DNA breakage.

Instability of a (GGGGCC)70 tract in humans was reported (29).

Sister-chromatid exchange and recombinant events in *C9orf72* expansion cells

An increased incidence of sister-chromatid exchanges (SCEs) occurs at rare and common

fragile sites (101). We analyzed SCE formation in *C9orf72*Exp. Unexpectedly, *C9orf72*Exp

cells exhibit an increase, up to 2-fold, in spontaneous genome-wide SCEs/cell (~7 up to 10-

15/metaphase), compared to control cells without expansions (Fig. 4C). Increased SCEs were

evident in *C9orf72*Exp cells from presymptomatic and symptomatic individuals (Fig. 4C,

compare P1, P5 with P6, all greater than WT). The increased SCEs did not require FUdR. This

increase is significant, yet modest relative to 10-fold increased SCE/metaphase in individuals

with the chromosome instability Bloom syndrome. Bloom individuals present high SCEs (102–

105), increased MN (103, 106), a self-DNA cGAS-STING-mediated *IFN* over-expression

(106), and often succumb to cancer before age 30 due to a coincident 10-fold increased

mutation rate (105, 107, 108). Chemical and genetic factors can drive SCEs (105, 109, 110).

In all instances, SCEs predominantly occur at common (111–113) and rare fragile sites (101,

114–117), regardless of whether the fragile site is being induced or not (113, 118). Consistent

with this, the SCEs levels in *C9orf72*Exp cells were mildly but not significantly increased by

FUdR (Fig. 4C). Through FISH mapping SCEs were significantly enriched on Chr9p, where

C9orf72 resides (Fig. ED6B-D). Thus, the GGGGCC-expanded FRA9A, like the CGG-

expanded FRAXA, the AT-repeat-expanded FRA16B, and the EBNA1-motif repeat 11q23

fragile site, incurs increased SCEs (76, 78, 101, 117, 119). These findings extend the forms of

genetic instability and the sources of DNA damage in *C9orf72*Exp cells to include global and

focal SCEs.

Unexpectedly, one of the *C9orf72*Exp cell lines (P8) showed complex rearrangements

restricted to one Chr9, clearly evident in every metaphase (100/100 observed) (Fig. 4D-F).

Specifically, one Chr9 showed a duplication of nearly the entire 9q-arm (9qter-q21.11 fused to

9p23, Fig. 4C), a terminal deletion of 9pter-p23, and duplications of 9p23, 9p12, 9p11.2, 9q13,

9q21.11, and deletions in 9q13, making a derivative-Chr9 (see CNV and whole-genome

sequencing analysis, Fig. 4D-E, Fig. ED8A, Tables ED4 and ED5). Breakpoint pattern was

reminiscent to 9p deletion/duplication syndrome (55, 120). Importantly, only the derivative-

Chr9 expressed FRA9A with FUdR in 7% (7/100) metaphases (Fig. 4F), indicating it arose

from the *C9orf72*Exp Chr9. Strikingly every metaphase showed the same rearrangements,

which were evident prior to FUdR treatment. In our previous examinations of thousands of

metaphases from control cells, we have not observed such homogeneity of rearrangements,

making it highly unlikely that this derivative-Chr9 arose by cell line establishment or culturing.

It is also highly unlikely that this individual, who was neurotypical until presenting with bulbar

onset ALS at age 56, had inherited this complex set of rearrangements constitutionally in all

tissues, as congenitally such rearrangements would be incompatible with neurotypical life (55,

120). Rather than inherited, it is likely the rearrangements arose somatically in the blood of the

*C9orf72*Exp individual, which is supported by our observation of similar rearrangements of

the *C9orf72* transgene-containing Chr6 in mouse tissues (below). Whole genome sequencing

revealed the derivative-*C9orf72*Exp-Chr9 presented a unique state with ≥ 13 breakpoints

clustered on 9p, with direct and inverted junctions, confined to a single chromosome, as such,

fulfilling the criteria of chromothripsis (121–123). It is possible the FRA9A-expressing

*C9orf72*Exp-Chr9 isolated in a micronucleus, was shattered and reassembled as the derivative-

Chr9 (124). A single experimentally targeted double-strand break can induce chromothripsis

to the targeted chromosome (125, 126). The derivative-Chr9 may have arisen by stepwise

alterations. The origin of the derivative-*C9orf72*Exp-Chr9 could not be discerned as the patient

was deceased. The complex chromosomal rearrangements of the mutant *C9orf72*Exp FRA9A-

expressing Chr9 could have downstream impact.

While we cannot exclude the possibility that the highly rearranged derrivarive-Chr9 in the

*C9orf72*Exp cell line (P8) arose during cell line establishment or passage, it is difficult to argue

away the coincidences that the rearrangements we observe were limited to only one Chr9 – the

Chr9 with the *C9orf72*Exp expressing FRA9A (Fig. 4D-F), and that the breakpoints were non-

random but aligned with the breakpoints of 9p-duplication/deletion syndrome (Fig. 4G) (55).

We also observed numerous forms of chromosome rearrangements of murine chromosomes

harboring the human *C9orf72* expansion, including one reminiscent of the der-Chr9 we

observed in the human cell (see murine cytogenetics described below). Moreover, the

cytogenetic observation of thousands of human EBV-transformed cells have not presented such

chromosomal instability (59), asides from the EBV-induced fragility at 11q23, which depends

upon the EBV protein EBNA1 (75–78).

Gene expression across most of the derivative-Chr9 is dysregulated. Expression analysis by

RNA-seq comparing P8 to other *C9orf72*Exp cells, revealed trisomic and monosomic

expression levels of many genes corresponding to the large duplicated and deleted regions (Fig.

4G, see red and blue dots, for significance see Fig. ED8B-D, an interactive html file detailing

mis-expressed   genes   can   be   accessed   here: https://data.cyverse.org/dav-

anon/iplant/home/ljsznajder/FRA9A/Chromosome9.html), also presented in Table ED4.

Amongst the dysregulated genes along the der-Chr9 in P8, many are also dysregulated in

vulnerable neurons of *C9orf72*Exp ALS patients, including ALS- and inflammation-linked

genes   (https://www.biorxiv.org/content/10.1101/2023.12.22.573083v1,   Table   ED4).

Breakpoint and copy number changes analyzed in P8 are presented in Table ED5.

Cell culture in FUdR increases repeat instability

We tested whether repeat length instability could be generated in *C9orf72*Exp patient cells

under fragile site inducing conditions. We observed that expansion tracts shifted towards

shorter repeat lengths over culture time in FUdR-treated cells (Fig. ED9). In contrast, under

uninduced conditions, the expanded repeat showed a general shift towards *longer* repeat sizes

or the more predominant repeat length (Fig. ED9). FUdR affected the length of only the mutant

*C9orf72*Exp allele. FUdR-induced changes were consistent in three different *C9orf72*Exp cell

lines and did not appear to depend upon methylation or clinical status (Fig. ED9). These results

support repeat size variations being generated under folate-perturbing fragile site inducing

conditions.

Tissue-specific somatic repeat expansions, chromosomal fragility, and chromosomal

instability in *C9orf72*Exp mice

Tissue-specific somatic repeat instability, FRA9A chromosome fragility, and chromosome

instability, was recapitulated in *C9orf72*Exp-Transgenic(Tg) mice (*C9orf72*Exp-Tg) harboring

a single copy of the human *C9orf72* with a GGGGCC expansion integrated into murine Chr6

(127). Using this mouse as a tool for instability, we isolated cells from various organs of

*C9orf72*Exp-Tg and control non-Tg mice, aged 5-months, when cell proliferation in the CNS

and peripheral organs had ceased at 2-months of age (128–130) (see Methods). To retain as

close a representation of cell types, harvested tissues were cultured for short periods sufficient

for chromosome preparation (1-4 weeks, with no storage). Tissues from three mice inheriting

~620-690 repeats (tail at two-weeks, aged 5-months) showed a heterogenous smear of 586-

1488 repeat sizes, with some of the greatest expansions in the CNS (Fig. 5A). This supports

massive post-natal inter- and intra-tissue repeat expansions in non-proliferating murine tissues,

as observed in *C9orf72*Exp humans (26–30). Expansions were evident as early as 2-weeks

(Shum *et al.*, in preparation).

Chromosomal fragility of the *C9orf72*Exp transgene occurred in all tissues at frequencies

of 5% to >35% (99% confidence, Fig. 5Bi-iii). Fragility presented as despiralized regions,

constrictions, chromatid, and isochromatid, breaks, with split transgene-specific *C9orf72* FISH

signals, as we observed in human *C9orf72*Exp cells (Fig. 5Bi). Only chromosomes from the

transgenic mice showed FUdR-induced fragile sites, where fragility localized to the expanded

repeat of the integrated transgene, but never to the non-transgenic Chr6 (Fig. ED10). Thus,

FRA9A fragility can be transferred to this Tg-mouse thereby validating the *C9orf72* repeat

expansion as the molecular cause of FRA9A.

Chromosomal instability was evident on the *C9or72*Exp-Tg-containing Chr6, including

acentrics, truncations, duplications, translocations, deletions, and multi-branched

chromosomes (Fig. 5C). In one instance the *C9orf72*Exp-Tg translocated itself from murine

Chr6 to another chromosome (Fig. 5C), and in another the *C9orf72*Exp-Tg was deleted (Fig.

5C). We observed a rearranged chromosome with a large duplication (Fig. 5C), reminiscent of

the derivative-*C9orf72*Exp-Chr9 in ALS patient, P8 (Fig. 4D-G). Chromosomal fragility and

instability were observed in all tissues of all three *C9or72*Exp-Tg mice.

While rearranged chromosomes were rare relative to fragility, neither were ever observed

in the non-transgenic murine Chr6 at 6qE3, the site of transgene integration, or in any control

non-transgenic tissues (three replicate mice). The absence of fragility at 6qE3 in control non-

transgenic tissues is consistent with 6qE3 not being an endogenous fragile site in mice (131).

Together our data support the ability of the *C9orf72*Exp-Transgene to incur post-natal somatic

tissue-specific repeat expansions, display chromosomal fragility and chromosomal instability,

as in humans. Our findings provide insight into fragile site expression and rearrangement in

the CNS and periphery.

Figure 5

Figure 5. *C9orf72* **repeat instability, chromosomal fragility/instability in** *C9orf72***Exp-Tg-**

mice

Southern blots for *C9orf72* transgene in C9-500 mouse tissues. C9-500 mouse tissues were

sized for repeat size via Southern blot in three *C9orf72*Exp Tg-mice. The tail was the most

stable tissue. (A) Somatic repeat instability by Southern blots in tissues of three 5-month-old

*C9orf72*Exp-Transgenic mice harboring a single copy of the human *C9orf72* gene with a

GGGGCC expansion integrated into murine Chr6 (127). (B) Cells isolated from tissues of the

mice shown in panel A were grown for 1-4 weeks sufficient for metaphase spreads for fragility

analysis with 1μM FUdR with FISH analysis. A murine-specific green probe identified mouse

Chr6, and human-specific red probe identified the *C9orf72*Exp transgene. Pictured are FSFS

in the CNS (i), the periphery (II), and quantified in panel iii. (C) Chromosomal instability at

the *C9orf72*Exp transgene was assessed in the same tissues used in panel C, using the same

FISH probes.

DISCUSSION

Here we demonstrate that *C9orf72*Exp is the molecular cause of the rare folate-sensitive FRA9A, cytogenetically observed more than 4 decades ago (58, 132). This reveals previously unconsidered avenues to study genotype-phenotype relationships for co-occurring disease states associated with *C9orf72*Exp and attributes linked to the 9p21 region. First, folate-sensitivity suggests a possible gene-environment interaction. Second, FRA9A may serve as a DNA source to activate DDR and *IFN* pathways. Third, FRA9A chromosomal instability and global SCEs may increase somatic mutations at 9p21 and beyond.

Disease penetrance & gene-environment

Gene-environment interactions have long been suspected for ALS, FTD, and the incompletely penetrant co-occurring phenotypes of *C9orf72*Exp (133). Folate-sensitive induction of *C9orf72*Exp repeat instability, brain region-specific repeat expansions, FRA9A fragility, chromosomal instability, and methylation-sensitive chromatin compaction, which we have shown can arise in cells and tissues of both presymptomatic and symptomatic individuals, may have clinical implications. Lifestyle, including nutritional choices, of *C9orf72*Exp carriers can affect disease onset and progression (40). Proper levels of the linked vitamins B9 (folate) and B12 are protectors from ALS (134). Individuals with folate deficiencies, nutritionally-, pharmaceutically-, or genetically-induced, are associated with chromosomal fragility, localized chromosomal rearrangements (135–144), megaloblastic/pernicious anemia, ALS- and FTD-like symptoms (145, 146), reviewed (59). In such cases these presentations were reversed upon B12 supplementation – indicating that chromosomal instability was induced by folate-stress. Somatic GGGGCC repeat instability, FRA9A fragility, and global increase of SCEs depend upon *C9orf72* expansion and are folate-sensitive and can occur in cells from presymptomatic *C9orf72*Exp individuals. *C9orf72*Exp carriers, in folate-deficient states, may be predisposed to

repeat instability, FRA9A fragility, and chromosomal instability, which if exacerbated over

time may impact gene dysregulation, altering the susceptibility of phenotypic variations

amongst *C9orf72*Exp carriers. Fragile sites and SCEs can be induced in the blood and myeloid

cells by lifestyle choices, including cigarette smoking, alcohol consumption, and diet (59, 147–

151). Tobacco and alcohol consumption by presymptomatic *C9orf72*Exp carriers affected

disease age-of-onset (40). Various clinically used agents (152), including some considered for

use in ALS/FTD, *C9orf72*Exp (153), and fragile X syndrome (154), like phenytoin or

decitabine can induce fragile sites (59, 155), induce sister-chromatid exchanges (156, 157), and

modify chromosome condensation (158, 159). Age-dependent ongoing repeat expansions in

the brain have been suggested as a contributor to disease (26, 160, 161), could be folate

sensitive. Additionally, CpG-methylation of *C9orf72*Exp (160, 162) and acceleration of

genome-specific DNA methylation age (epigenetic clock) are modifiers of disease onset and

progression in *C9orf72*Exp carriers (163, 164). Chromatin compaction of *C9orf72*Exp is

sensitive to the degree of expansion and CpG-methylation (Fig. 3). Efforts to prophylactically

slow the epigenetic clock are a central focus (165–167). The *C9orf72* expansion was recently

reported to be bound by the chromatin remodeler DAXX (168). Coincidentally, DAXX, is

known to associate with satellite and telomere repeats and its downregulation led to a global

increase in MNase sensitivity (169–171). Environmental contributions to *C9orf72*Exp diseases

must be considered.

FRA9A as a DNA source to activate the DDR and *IFN* pathways

Both a spontaneous activated DNA damage response (DDR) (15–25) and a hyperactivated *IFN*

cGAS-STING-associated immune response occur in *C9orf72*Exp patient cells and brains (4,

12–14). The source of the damaged DNA or DNA-immunostimulant to activate either DDR or

IFN response is unknown. DDR in cells and brains of *C9orf72*Exp, evident as increased γ-

H2AX, phosphorylated-ATM, cleaved PARP-1, and 53BP1, is coincident with DSBs (22–24).

We reveal that DSBs arise at *C9orf72*Exp via heightened FRA9A fragility, broken/rearranged

Chr9s, increased genome-wide SCEs, increased MN, and Chr9-containing MN. The increased

C9ORF72 protein levels in multiple tissues of the asymptomatic father shown to have massive

somatic expansions (Fig. 3F and (29, 79)) may suppress an autoimmune response by increasing

STING levels. Models of the *C9orf72*Exp-linked DDR activation involve RAN-peptide

induced p53-target gene activation, inhibition, or mislocalization of the DSB DNA repair

proteins ATM, 53BP1, TDP-43, FUS, Ku80, and PARP-1 (15–25). Other non-RAN-peptide

DDR paths have received little attention.

The unusual >33kb of compact chromatin at FRA9A's core fragility zone (Fig. 3), could be

susceptible to DSBs, which would contribute to DDR activation. Supporting this possibility is

that experimentally-induced locus-specific compacted chromatin of an artificial ~10kb stretch

of ~250 tandem arrays of LacO, which becomes a fragile site, has been demonstrated to activate

a DDR in the absence of damage (172–175). A contribution of FRA9A fragility to DDR can

act in parallel to RAN-peptide induced DDR (23, 24). Disruption of higher-order chromosomal

organization may also contribute to DDR and *IFN* responses. Long-range chromosomal

interactions involving 9p21 are activated in the inflammatory response. All *IFN* genes at 9p21

are transcriptionally inactive until immunostimulated by viral infection or cytosolic self-DNAs,

whereupon dramatic chromatin (176) and multiple intra-9p21, inter-arm (9p21-9q34.2), and

inter-chromosomal (9p21-4p14 and 9p21-18q21.1) interactions arise to activate *IFN* cluster

expression and inflammation (47, 177). Reduced levels of the C9ORF72 protein in *C9orf72*Exp

patients, being unable to supress STING, leads to a sustained hyperactivation of *IFN* in brains

(178). Upon FRA9A breakage the *C9orf72*Exp-Chr9 activates DDR and is then sequestered to

MN whose contents can be released to the cytosol to activate *IFN* through cGAS-STING, and

this sequestration may disrupt other inter- and intra-Chr9 interactions (179). That disruption of

intra-/inter-chromosomal associations can drive DDR and *IFN* cascades is supported by the

experimental disruption of such interactions leading to chromosome sequestration to MN (180,

181). CGG expansions at *FMR1*, the cause of FRAXA, disrupt intra- and inter-chromosome

interactions (182). The 2-fold increase in global SCEs in *C9orf72*Exp cells we observe may

contribute to DDR and MN. MN enriched with the DSB repair protein, TDP-43, have been

detected in brains of ALS patients (183). Our suggestion that DNA damage induced by FRA9A

fragility can serve as a DNA-immunostimulant in the CNS is consistent with reports of *in vivo*

DNA damage by ATM-deficiency or brain trauma that activates an *IFN* response in motor

neurons (184–186). Our working model that the *C9orf72*Exp acts as a chromatin compacted

FSFS and is an inducer of DNA damage at many locations including on Chr9, is suited to non-

cycling neurons.

We suggest that the cytoplasmic release of DNA from FRA9A-enhanced MN, may further

tip the balance by exacerbating the stimulation of the aberrantly hyperactive cGAS-STING

initiated *IFN* expression in *C9orf72*Exp carriers (4, 13, 14) – which may explain the co-morbid

partially-penetrant autoimmunity (1–6). FRA9A broken-Chr9s and increased global SCEs

might also further exacerbate the DDR cascade, as DNA-immunostimulation enhances sense

and antisense transcription from *C9orf72* (42) whose RAN-peptide products would feedback

upon the DDR by suppressing DSB DNA repair proteins ATM, p53, TDP-43, FUS, Ku80, and

PARP-1 (15–25). The neuroinflammatory and autoimmune phenotypes of *C9orf72*Exp carriers

is complex (14), and the effects of either FRA9A altered chromatin, or MNs containing broken

Chr9s upon the *IFN* response is unknown.

FRA9A chromosomal instability, mosaic somatic mutations, and 9p21-associated diseases

A reasonable prediction from our findings is that the *C9orf72*/FRA9A fragile site may, like

other fragile sites, predispose to mosaic deletions, duplications, or rearrangements resulting in

altered expression of genes on 9p21 or throughout Chr9. Such mosaic chromosomal variations

may vary between tissues, accumulate with age, and lead to phenotypic variability. In the case

of *FMR1*/FRAXA, phenotypic variability can arise through mosaic genetic or chromosomal

instability (deletions, duplications, and rearrangements) extending beyond the *FMR1* CGG-

repeat, as revealed through decades of isolated case reports (59). Similar phenomena may arise

at the mutant *C9orf72*, as suggested by our observation of truncated-Chr9s broken at *C9orf72*,

acentric-Chr9s, chromosomal instability of the *C9orf72*Exp allele, and heightened levels of

genome-wide SCEs. Detection of somatic mosaic levels of chromosomal instability will likely

require more focused studies. For example, the dramatic Chr9 rearrangements, evident by FISH

in every metaphase of the ALS/*C9orf72*Exp line P8, escaped detection by repeat-primed PCR

and by Southern blot assessment of the repeat size (compare Fig. 1B with Fig. 4F). Indeed,

mosaic mutations at various loci (reviewed in (187–189)) including *FMR1* (190–195) required

focused attention. Mosaic mutations may be confined to specific cell types, as recently revealed

for Purkinje cells with large somatic CAG-expansions, amongst stable tracts in the cerebellum

of Huntington's disease patients (196, 197). One might expect an increase in somatic mutation

rates throughout the genome in *C9orf72*Exp patients, as the increased 2-fold global SCEs we

observed may have a coincident increase in mutation rate, similar to the 10-fold increased

mutation rates in BLM individuals who show a 10-fold increase of SCEs (102, 103, 107, 108).

*C9orf72*Exp cells from presymptomatic individuals show folate-sensitive repeat instability,

FRA9A fragility, chromosomal instability, MNase-resistance, increased levels of

micronuclei/NBuds, and global sister chromatid exchanges. Long-term display of these

features may exacerbate disease-onset and affect penetrance. Our findings support that the

*C9orf72*Exp patients may incur mosaic mutations at under-appreciated levels, and recent

findings support increased somatic mutations in brains of *C9orf72*Exp individuals

(https://www.biorxiv.org/content/10.1101/2023.11.30.569436v1).

Knowledge gaps remain for FRA9A/*C9orf72*. In particular, the co-occurrence of many

diverse phenotypes in *C9orf72* expansion carriers is unclear. That FRA9A encompasses the

IFN cluster, *CDKN2A*, *IFNk*, *LINGO2*, *TOPORS*, *APTX*, *DNAJA1*, and *SMU1* genes, each

independently associated with symptoms that can co-occur in *C9orf72*Exp carriers, provide

new insights into the possible involvement of chromosomal instability (Fig. 1A). An increase

in ALS occurring in cancer survivors, particularly melanoma, has been repeatedly reported

(7, 198–202). While a correlation of *C9orf72*Exp in cancer predisposition has not been

studied in large populations, *C9orf72*Exp carriers have a significantly higher risk of

melanoma (7). *C9orf72*Exp carriers may succumb to cancer prior to onset or awareness of

ALS. That both *CDKN2A* and the *IFN* gene cluster, frequently deleted in melanoma, are

encompassed in FRA9A's fragility zone introduces the possible involvement of *C9orf72*Exp.

Mosaic somatic mutations that extend beyond the *C9orf72* repeat, may affect expression of

genes encompassed by FRA9A or beyond. Thus, revisiting the *C9orf72*/9p21 locus with the

awareness that it is, and can share the attributes of, a chromosomal fragile site should provide

key insights into the genotype-phenotype relationships for the many and varied presentations

associated with the 9p21 region.

 MATERIAL AND METHODS

In silico **datasets and analyses**

Sequence feature and epigenetic datasets used in Figure 1 and Figure ED2 are described as

follows. Repeat annotations and %GC tracks were obtained from the UCSC genome browser

(hg19). Wavelet-summarized repliseq data (ENCODE) was obtained in bigWig format from

GEO:GSE34399. Erythroblast SNS-seq data and replication origins were obtained in bedGraph

and bed format, respectively, from GEO:GSE6197. Imputed %DNA methylation for GM12878

was obtained from the roadmap epigenomics data portal:

(https://egg2.wustl.edu/roadmap/data/byFileType/signal/consolidatedImputed/DNAMethylS

BS/E116-DNAMethylSBS.imputed.FractionalMethylation.signal.bigwig). Pre-calculated

2nd-order DNA shape features were obtained from the Rohs lab GBShape database.

Cell culture and fragile site induction

All cytogenetic studies were performed using Epstein-Barr virus transformed lymphoblastoid

cell lines from *C9orf72*-ALS patients, expansion carriers, and control individuals provided by

Dr. Guy Rouleau (McGill University). One of the patients, P4, displayed signs of FTD. All

patients provided written informed consent. Cells were cultured in RPMI 1640 media

supplemented with 15% FBS (ThermoFisher Scientific), 2 mM L-glutamine, Pen/Strep (100

IU/mL, 100 µg/mL), and 1 mM sodium pyruvate. Two modes of FSFS induction were used to

demonstrate fragility at 9p21.2: i) 24 hours with 0.1 µM 5-fluorodeoxyuridine (FUdR) or ii)

14 days with Media 199 (Gibco, Cat #: 11150-059) supplemented with 2% FCS, 2 mM L-

glutamine, and Pen/Strep (100 IU/mL, 100 µg/mL). Cells in 199 media were treated with 100

nM methotrexate for 16 hours, washed and then treated with 0.01 µM bromodeoxyuridine

(BrdU) for a total of 3 hours prior to harvest. All cells were treated with 6 µg/mL ethidium

bromide (EtBr) and 0.05 µg/mL KaryoMAX (Gibco; Cat #: 15212012) prior to harvest. Cells

were then centrifuged and resuspended in 0.075 M KCl solution for 30 minutes at 37ºC

followed by several washes with fixative solution (3:1 methanol:acetic acid). Long-term FUdR

culture experiments were performed in either RPMI 1640 media (as described above) or RPMI

1640 media + 0.1 µM FUdR for a total of 8 weeks. Media and FUdR was replaced every 2-3

days. Cells were harvested at 1, 2, 4, 6, 8-week time points for DNA purification and Southern

blot was run as described below.

725   Metaphase chromosome and FISH analysis

Slide preparation for both cytogenetic and FISH analysis of metaphase spreads were made from

fixed cell suspensions using a Model CDS 5 Cytogenetic Drying chamber according to standard

methods. For 199 media conditions, breaks and gaps were quantified on total Giemsa stained

metaphases and subsequently G-banded to karyotype each metaphase and confirm fragile site

on 9p-arm. SpectrumGreen & SpectrumOrange-conjugated dUTP probes (RP11-15P13, RP11-

274I21, RP11-1149M23, RP11-491J7, RP11-945G22, RP11-1079P2, RP11-360B21, RP11-

696J10) from RPCI-11 human BAC library were generated by The Centre for Applied

Genomics (TCAG, PGCRL, Toronto, Canada) using the Vysis nick translation kit (Cat #07J00-

001; Abbott Molecular, Illinois, USA) in accordance with manufacturer's instructions.

Chromosomes were counterstained with DAPI. One hundred metaphases per sample were

scored for each condition using either brightfield microscopy, Olympus IX81 quorum spinning

disk confocal microscope, or Zeiss Axioplan 2 fluorescence microscope. Images were collected

using a CCD camera (Hamamatsu C9100-13 EM-CCD, Hamamatsu, Japan) (Princeton

Instruments Pentamax, Roper Scientific, New Jersey, USA) and analyzed using Volocity

(Perkin Elmer, Massachusetts, USA) or CytoVision (Leica Biosystems, Germany) software.

741   Southern blot analysis of *C9orf72* repeat expansion size

Southern blot analysis was performed to size *C9orf72* repeat expansions as this method is the

gold standard for sizing large (GGGGCC)n repeat expansions (203). Southern blotting

provides an estimate of non-bias expansion sizes (204). We used Southern blotting method

(203, 205) with 241 bp probe, which anneals 153 bp upstream of the (GGGGCC)n repeat tract,

and used selected restriction-endonuclease digestions of genomic DNA (as noted in the figures)

– which permit sensitive detection of repeat size and length heterogeneity at molar levels. The

241 bp probe was produced by PCR using primers (Fwd: 5'–

AGAACAGGACAAGTTGCCCC–3' and Rev: 5'–AACACACACCTCCTAAACCC–3') as

published (203, 205). For mapping boundaries of MNase accessibility an additional probe was

used (see below). Molecular weight markers were used to estimate the size of repeats.

Distances between molecular weight markers were measured using ImageLab software

(BioRad) and plotted against the known number of base pairs for each marker. A best-fit linear

regression was calculated in Microsoft Excel and the equation of this line was used to calculate

the size of each band on the blot. Different restriction enzyme combinations were used to cut

the genomic DNA, as noted in each figure, hence resulting in different flank repeat sizes.

Methylation analysis of the *C9orf72* repeat

Methylation analyses were done in blinded experiments using the same DNA prep as for

Southern blot. Genomic DNA was bisulfite converted using the EZ DNA Methylation-

Lightning™ Kit (Zymo). We estimated the number of methylated CpG-sites at the CpG-island

5′ of the (GGGGCC)n repeat using bisulfite sequencing as reported previously (61). To

estimate the methylation of the (GGGGCC)n repeat itself, we used a qualitative GGGGCC-

methylation assay, sensitive enough to detect repeat methylation in a mixture containing ~5%

high-methylated DNA standard (60). We always observed methylation mosaicism for

expanded repeat samples, revealed by presence of the amplification product of the expansion

in both methylated (blue) and unmethylated (green) channels (206).

Micronucleus and nuclear bud detection

Micronuclei and nuclear bud analyses were done as previously described with some

modifications (86, 88, 207). Briefly, cells were treated with 0.5 µM FUdR for a total of 24

hours, with cytochalasin B (4 µg/mL) being added 3 hours after addition of FUdR. Cells were

briefly incubated in 0.075 M KCl, fixed with Carnoy's fixative and slides with sparsely spread

cells were prepared using a Model CDS 5 Cytogenetic Drying chamber (Thermotron,

Michigan, USA). SpectrumGreen (RP11-696J10) & SpectrumOrange (RP11-672M4)-

conjugated dUTP probes from RPCI-11 human BAC library were generated by The Centre for

Applied Genomics (TCAG, PGCRL, Toronto, Canada) using the Vysis nick translation kit (Cat

#07J00-001; Abbott Molecular, Illinois, USA) in accordance with manufacturer's instructions.

Nuclei were counterstained with DAPI. At least 950 binucleated cells were scored per

condition for the presence of micronuclei and nuclear buds and Chr9 signals, as previously

described (88). Two blinded and independent experiments were carried out for each data asset.

Statistical analyses were done using Fisher's exact test for the combined values of micronuclei

and nuclear bud events as they are believed to arise from the same phenomenon (88, 208).

Images were captured using an Olympus BX61 fluorescence microscope. Images were

collected using a CCD camera (Hamamatsu C9100-13 EM-CCD, Hamamatsu, Japan)

(Princeton Instruments Pentamax, Roper Scientific, New Jersey, USA) and analyzed using ASI

SpotScan software (Applied Spectral Imaging, California, USA).

Sister-chromatid exchange (SCE) assay

Cells were incubated in the presence of 10 µM BrdU for two cell cycle periods and pulsed with

0.1 µg/mL of KaryoMAX (Gibco; Cat #: 15212012) for the last 2 hours before being harvested.

FUdR (0.1 µM) was added for the last cell cycle. Harvested cells were treated with 0.075 M

KCl for 20 min and subsequently fixed with methanol:acetic acid (3:1) for at least 30 minutes.

Cells were fixed onto glass slides. Dried slides were incubated with 10 µg of Hoechst 33258

per ml in phosphate buffer (pH 6.8) for 20 minutes, followed by rinsing with MacIlvaine

solution (164 mM Na_2HPO_4, 16 mM citric acid [pH 7.0]). Slides were irradiated with UV light

for 2 hours and incubated in 2X SSC solution at 62°C for 1 h before staining with 5% Giemsa

solution and subsequent microscopy. At least 100 metaphase cells were evaluated for SCE

events genome-wide per genotype. For Chr9-specific SCEs, slides were destained with xylene

solution followed by methanol:acetic acid (3:1) and then standard FISH procedures, as

described above were followed using SpectrumOrange-dUTP conjugated RP11-696J10 probe.

At least 30 of the same Giemsa-stained metaphases were relocalized using fluorescence

microscopy and imaged to allow for quantification of SCEs specifically on the 9p- and 9q-

arms.

Construction of plasmids and yeast strains

An *Xba*I-*Hin*dIII fragment containing a region of the human *C9orf72* gene containing an

expanded (GGGGCC)n repeat and 547 bp of flanking sequence (from pcDNA3.1(+)c9-200+

obtained from R.H. Brown; CF stock #435) was cloned into the MCS of plasmid pEP1 to create

plasmids pEP1-(GGGGCC)n. pEP1 was made by cloning the *ADE2* gene from pRS402 into

the *Bsa*AI and *Nsi*I sites of pHZ1; pHZ1 was made by cloning the G4T4 sequence into the *Sma*I

site of pYIP5. The CAG repeat was cloned into pEP1 *Xho*I and *Xba*I sites. Repeat lengths in

plasmid clones were confirmed by sequencing. Repeat lengths of the (GGGGCC)n plasmids

were approximately 16, 29, and 36 for plasmids 504, 484, and 485, respectively; exact number

of repeats varied with different sequencing runs. Plasmids were linearized with *Xcm*I and

recombined into the *URA3* marked yeast artificial chromosome (YAC) CF1 in strain VPS105

(CFY #289) to create the new GGGGCC-*ADE2-URA3* and CAG-*ADE2-URA3* YACs (see Fig.

ED5A & ED5D), which were confirmed for YAC structure by PCR. CAG tract lengths were

confirmed by sequencing of PCR amplicons but (GGGGCC)n tract lengths could only be

determined approximately by Southern blot since amplification from genomic DNA was

unsuccessful, and are therefore reported to the nearest 5 bp.

YAC fragility assay

End loss of the repeat-*ADE2-URA3* YACs was determined by an established protocol (100)

except that 1 ml of an overnight culture inoculated with a specific colony was plated on media

containing FOA and a low amount of adenine but lacking leucine (FOA-Leu low Ade) to

determine the number of mutants, and a dilution was plated on yeast complete media lacking

leucine (YC-Leu) for a total cell count. Rates were determined by the method of the median.

Another 1 ml of the same culture was used to isolate genomic DNA for Southern blot to assess

repeat length. For reasons not completely understood, but that could include reduction of

transcription through the repeat tract, the insertion of the *ADE2* marker resulted in a lower rate

of FOA^R compared to previously published values for the CAG-85 YAC without the *ADE2*

gene. End loss was confirmed in a subset of colonies (usually one from each FOA-Leu plate)

by PCR (retention of the sequence to the left of the G_4T_4 telomere seed and loss of sequence to

the right of the repeat tract; Fig. ED5F).

Southern blot - yeast

Yeast genomic DNA was digested with *Nhe*I and *Xba*I restriction enzymes, separated by

horizontal gel electrophoresis, and blotted to a nylon membrane using a standard Southern

protocol. Either the Roche Molecular Weight Marker VI or VII was run as a size marker. For

the preparation of the probe, a 301 bp fragment adjacent to the (GGGGCC)n repeat was PCR

amplified from DNA of plasmid #485 as template, using Primers #1402 (ALSsouthernFOR)

and #1403 (ALSsouthernREV). Probes were labeled using the DIG High-Prime DNA Labeling

and Detection Starter Kit I (Cat #: 11745832910; Roche). Repeat tract estimation was

performed as previously described (Southern blot analysis of repeat expansion size).

Chromatin accessibility *in cellulo* with MNase

Human lymphoblast cell lines from *C9orf72*-expansion carriers were assessed for MNase

accessibility, as outlined in Figure 3. Briefly, cell pellets were washed in ice-cold 1X PBS, re-

pelleted, and permeabilized by ice-cold NP-40 lysis buffer (10 mM Tris pH 7.4, 10 mM NaCl,

3 mM MgCl$_2$, 0.5% NP-40, 0.15 mM spermine, 0.5 mM spermidine) for 5 min on ice. Nuclei

were collected at 200 Xg (10 minutes, 4ºC), washed in 1X MNase digestion buffer (10X stock:

100 mM Tris pH 7.4, 150 mM NaCl, 600 mM KCl, 5 mM spermidine, 1.5 mM spermine), re-

pelleted, and subsequently resuspended in 1X MNase digestion buffer supplemented with 1mM

CaCl$_2$. Treatment followed with MNase for 5 min. at 22ºC, at differing concentrations (refer to

Fig. 3, ED7 – all serial dilutions). All digestion reactions were stopped with an equal volume

of *Stop* solution (0.625 mg/mL proteinase K, 20 mM EDTA, 20 mM EGTA, 1% SDS diluted

in 1X MNase buffer) and incubated at 37°C for a minimum of 1 hour. Reaction conditions were

optimized to ensure over-digestion by MNase, which diminishes MNase cleavage preference.

DNA was purified by phenol/chloroform extraction, RNase A treatment, and ethanol

precipitation. A small amount of purified DNA was assessed for completeness of MNase

digestion, by its conversion to mono-, di-, tri-nucleosome sized fragments as resolved by

electrophoresis stained with ethidium bromide and visualized by UV imaging (Fig. 3B, left

panel; ED7A). DNA samples were analyzed by Southern blot as described.

Chromatin accessibility in post-mortem tissues with MNase

We conducted MNase accessibility analysis using post-mortem brain tissues (orbito-frontal

cortex and cerebellum) of an asymptomatic 90-year-old male and his 59-year-old ALS-FTD-

affected daughter, previously deep-phenotyped (summarized in Table ED3) (29, 79). The

asymptomatic male carried 70-repeat *C9orf72* expansion in the blood and was deceased at 90-

years of age. Tissue was Dounce homogenized using 1X MNase digestion buffer, 10% NP-40

was added to final concentration 0.5%, incubated 5 minutes on ice, filtered through a 40 μm

strainer, and centrifuged. Nuclei pellets were washed in MNase digestion buffer and incubated

with different concentrations of MNase. Remaining steps were performed as described above.

Digestibility of *C9orf72* BAC DNA *in vitro* with MNase

A bacterial artificial chromosome (BAC) containing ~160 (GGGGCC)•(GGCCCC) repeats,

prepared in the lab of Piet de Jong, was restricted by *Nhe*I and *Xho*I giving rise to a shorter

fragment of approximately 1750 base pairs, which was gel-purified. Half (by volume) of the

purified product underwent CpG-methylation by *M.Sss*I methyltransferase supplemented with

640 µM S-adenosylmethionine (New England Biolabs) and purified using the QIAEX II kit

(Qiagen). Completeness of methylation was checked by the inability to be digested by the

methyl-sensitive *Hpa*II but successful digestion by the methyl-insensitive *Msp*I. Subsequently,

in parallel, equal concentrations (~100 ng/reaction) of the non-methylated and methylated

fragments were treated with increasing levels of MNase (0U – 0.1U – 1U – 5U – 25 U) for 5

minutes at 22ºC in 1X MNase digestion buffer. The MNase reaction was halted by addition of

6X purple loading dye (New England Biolabs) and electrophoresed immediately on a 1%

agarose gel at 50 V for 20 minutes, followed by 100 V for 100 minutes. Gene Ruler Mix

(Thermo Scientific) was loaded in the first lane. Electrophoresed products were stained with

ethidium bromide, visualized by UV imaging (Fig. ED8B, left hand side), and DNA fragments

were transferred to a positively-charged nylon membrane, and probed with a 177 bp probe,

annealing just upstream of the GGGGCC repeat tract. To maximize our ability to assess that

loss of signal corresponded to MNase digestion of the GGGGCC repeat fragment, we

developed the 177 bp probe which anneals immediately adjacent to the repeat. (The 177 probe

anneals 7 bp upstream of the GGGGCC repeat tract, as opposed to the 241 bp probe, with

anneals 153 bp upstream). The 177 bp probe was produced by PCR using primers (Fwd: 5'–

GAGCAGGTGTGGGTTTAGGA–3' and Rev: 5'–CGACTCCTGAGTTCCAGAGC–3'). A

phosphor image of the Southern blot was obtained using the Typhoon system (Fig. ED8B, right

hand side), and the rate of digestion of probed sequence was calculated using densitometric

analysis in ImageJ. The percentage of DNA resistant to MNase (therefore, the fraction of

undigested 'starting fragment' BAC DNA) was plotted as a function of MNase units of activity

(concentration) in GraphPad Prism 8 (Fig. ED8B).

MNase controls

The MNase resistance we observe is not a reflection of the sequence preference of MNase or

the sequence composition of the *C9orf72* region. The MNase resistance spanning the mutant

C9orf72 locus supports altered chromatin packaging of the non-repetitive flanks, as the 241 bp

probe anneals 153 bp upstream of the (GGGCC)n repeat, and resistance is still evident (Fig.

7). The relative resistance of the expanded allele compared to the internal control of the non-

mutant *C9orf72* allele, provides strong support for altered chromatin accessibility. MNase is

known to display a cleavage sequence preference, preferring 5′-NpT and 5′-NpA sites(209,

210), yet is capable of digesting G-C only sequences and can digest most calf thymus DNA

with over-digestion, which diminishes MNase cleavage preference(209). All of our nuclear

digests were completed to over-digestion. The *C9orf72* sequences flanking the repeat, that are

probed herein, contain an extensive number of the preferred 5′-NpT and 5′-NpA sites (Fig.

ED2B, green font). That protein-free genomic DNA from *C9orf72* expansion carriers is

digested at an equal rate relative to the internal equimolar control of the non-expanded *C9orf72*

allele, supports that the flanks can be digested (Fig. ED7A). The MNase digest of the purified

(GGGGCC)160 repeat DNA further confirms that the pure GC-rich repeat can be digested by

MNase as evidenced by the ethidium bromide-stained gel (Fig. ED7B). As such, the reduced

digestion of the expanded allele in its native chromatinized state represents a chromatin-

mediated inaccessibility.

To rule-out possible artefacts such as differences in the sensitivity between ethidium

bromide staining and Southern probe hybridization, or loss of DNA during Southern transfer,

or an uneven transfer or membrane retention of DNA, an aliquot of the same samples were run

on a high-resolution gel, ethidium-stained, hybridized with the *C9orf72*-specific probe,

imaged, and then the blot was stripped and re-probed using radiolabeled total genomic DNA

as the probe (Fig. ED7A). The hybridization pattern from total genomic DNA probe is very

similar to the ethidium bromide-stained gel: most of the genome has been digested to

completion to mono-, di-, tri-, and tetra-nucleosome sized fragments (Fig. ED7A). In contrast,

hybridization with the *C9orf72* probe shows dramatic MNase resistance of the expanded allele,

while the non-expanded allele appears to be digested at the same rate as the rest of the genome.

Control MNase digestions of protein-free genomic DNA of unmethylated and methylated

C9orf72-ALS samples, compared to the chromatinized samples of the same cell lines showed

that MNase digestion occurs similarly for both alleles at extremely low MNase concentrations

(0.01 Units) in protein-free conditions, yet the expanded allele is strongly resistant in

chromatinized samples, supporting that this resistance is due to an unusual chromatin state (Fig.

ED7B-C).

There is a striking difference in MNase digestion between the ethidium-stained bulk

genomic DNA, and the Southern blot detected *C9orf72* expanded allele, which is internally

controlled for by its non-expanded homolog (Fig. 3). See above notes and Figure ED7 for

controls. The *C9orf72* region showed reduced presence in the mono-, di-, and tri-nucleosomal

digested regions, relative to the rest of the genome (compare Southern blot with ethidium stain,

Fig. 3B).

Control MNase digests using cloned *C9orf72* expanded (GGGGCC)160 repeat and flanks

(+/-CpG-methylation) were performed to assess the impact of MNase sequence selectivity and

the influence of CpG-methylation on digestibility. As shown in Figure ED7B, the expanded

repeat itself can be completely digested by MNase, and under the conditions used, the effect of

sequence and CpG-methylation had limited effects on MNase digestibility, as previously

reported (209, 210). This is consistent with the extensive number of the MNase-preferred 5′-

NpT and 5′-NpA sites (209, 210) in the *C9orf72* sequences flanking the repeat, that are probed

in the Southern blot (Fig. ED2B, green font). These controls, coupled with the equal rate of

MNase digestion of the protein-free *C9orf72* expanded allele in genomic DNA relative to the

internal control of the non-expanded *C9orf72* allele (Fig. ED7A-ii), demonstrates that the

expanded repeat and flanks can be digested. Thus, the inaccessibility of MNase to the

chromatinized *C9orf72* expanded allele and flanks (where the Southern probe binds), is due to

its unusual chromatin compaction, and is not due to an inherent inability of MNase to cleave

the expanded (GGGGCC)n DNA sequence.

Further mapping of the MNase accessibility in the chromatinized *C9orf72* expanded cells,

with and without aberrant methylation, was performed. Essentially, for large expansions the

mutant *C9orf72* allele is less accessible to MNase ~3,300 bp upstream and ~2,500 bp

downstream of the expanded repeat. Decreased accessibility is further exacerbated by aberrant

methylation (detailed in Fig. ED7A-iii). The absolute boundaries of decreased MNase

accessibility varied with repeat expansions.

Mapping MNase accessibility boundaries

We mapped the boundaries of MNase resistance in patient cell lines with no expansions (WT),

expansions (P1, P7), and expansions with methylation (P3, P6), using a series of post-MNase

restriction digests upstream and downstream of the repeat; fragments were detected by

Southern blotting using either the 241 bp or 177 bp probe, as indicated. Upstream mapping was

carried out with a post-MNase genomic digestion using *Ban*I, which cleaves just downstream

of the repeat, in combination with either one of a series of restriction digests progressing along

the upstream flank, as far as 3335bp upstream (*Afl*II, *Eco*RI, *Pvu*II, *Sph*I) (Fig. ED7C). These

samples were assessed by Southern blot, detected with the 177 bp probe (Fig. ED7C upper

panel, schematic). For each restriction fragment, the non-expanded allele signal was

completely lost while the expanded allele was either resistant or partially digested to a shorter,

distinct, nearly repeat-only containing fragment. This might suggest that the repeat tract itself

is the most resistant to MNase accessibility. The amount of MNase-resistant starting material

decreases progressively with increasing distance upstream from the repeat (Fig. ED7C). Thus,

the MNase resistance of the mutant allele extends upstream of the repeat at least ~1252 bp to

the *Pvu*II site, where some of the full-length fragment is still inaccessible to MNase, but

diminishes further upstream ~3335 bp to the *Sph*I site (Fig. ED7C). Downstream mapping was

conducted using a post-MNase genomic digestion with *Afl*II, which cleaves upstream of the

repeat and beside the region detected by the 241 bp probe. *Afl*II digestion was paired with one

of a series of restriction digests progressing along the downstream flank, as far as 3750 bp

downstream (*Ban*I, *Bst*YI, *Bam*HI, *Pvu*II, or *Psi*I) (Fig. ED7C lower panel, schematic). For

each restriction fragment, the expanded allele was resistant and progressively digested to a

shorter, distinct nearly repeat-only containing fragment, again suggesting that the repeat tract

is the most MNase resistant region. Resistance in the downstream region was more evident for

the methylated allele, and for larger expansions. For the larger methylated sample, P3 (~3000

repeats), the MNase resistance of the mutant allele extends downstream of the repeat at least

~2440bp to the *Pvu*II site, where some of the full-length fragment is still inaccessible to MNase,

but diminishes further downstream ~3750 bp to the *Psi*I site (Fig. ED7C). For the shorter

methylated expansion P6 (~770 repeats), the MNase resistance of the mutant allele extends

downstream of the repeat at least ~1714bp to the *Bam*HI site, where some of the full-length

fragment is still inaccessible to MNase, but diminishes further downstream ~3750bp to the *Psi*I

site (Fig. ED7C). Thus, the MNase resistance of the mutant allele extends at least ~2440 bp

downstream from the repeat to the *Pvu*II site, where some of the full-length starting fragment

is still resistant downstream ~3,750 bp to the distal *Psi*I site (Fig. ED7C). The methylated and

expanded allele was more resistant than the non-methylated expanded allele. In summary, for

large expansions some of the mutant *C9orf72* alleles can be resistant to MNase ~1250 bp,

diminishing through to ~3300 bp upstream and downstream resistance is ~2440 bp,

diminishing through to ~3700 bp away from the expanded repeat, and resistance is further

exacerbated by aberrant methylation.

Southern blot - *HTT*

Southern blotting for the *HTT* allele was followed as described (211), modifying the probe to

be 226 bp, which anneals 143 bp upstream of the (CAG)n repeat tract, and used selected

restriction-endonuclease digestions of genomic DNA – permitting sensitive detection of repeat

size and length heterogeneity at molar levels. The 226 bp probe was produced by PCR using

primers (Fwd: 5'-CTTGCTGTGTGAGGCAGAAC-3' and Rev: 5'-

CGCAGGTAAAAGCAGAACCT-3'). Molecular weight markers were run alongside the DNA

(Gene Ruler Mix, Thermo Scientific, and Roche Molecular Weight Marker II).

Chromatin accessibility *in cellulo* with MNase

An HD patient fibroblast cell line, with an *HTT* expansion of (CAG)21/180 (GM09197) was

assessed for MNase accessibility at *HTT*. Cell pellets were washed in ice-cold 1X PBS, re-

pelleted, and permeabilized by ice-cold NP-40 lysis buffer (10 mM Tris pH 7.4, 10 mM NaCl,

3 mM MgCl$_2$, 0.5% NP-40, 0.15 mM spermine, 0.5 mM spermidine) for 5 min. on ice. Nuclei

were collected at 200 *xg* (10 minutes, 4ºC), washed in 1X MNase digestion buffer (10 X stock:

100 mM Tris pH 7.4, 150 mM NaCl, 600 mM KCl, 5 mM spermidine, 1.5 mM spermine), re-

pelleted, and subsequently resuspended in 1X MNase digestion buffer supplemented with 1mM

CaCl$_2$. Treatment followed with MNase for 5 min. at 22ºC, at differing concentrations (0.5-

20U). All digestion reactions were stopped with an equal volume of *Stop* solution (0.625

mg/mL proteinase K, 20 mM EDTA, 20 mM EGTA, 1% SDS diluted in 1X MNase buffer)

and incubated at 37°C for a minimum of 1 hour. Reaction conditions were optimized to ensure

over-digestion by MNase, which diminishes MNase cleavage preference. DNA was purified

by phenol/chloroform extraction, RNase A treatment, and ethanol precipitation. A small

amount of purified DNA was assessed for completeness of MNase digestion, by its conversion

to mono-, di-, tri-nucleosome sized fragments as resolved by electrophoresis stained with

ethidium bromide and visualized by UV imaging. 12 ug of each DNA sample was then

restriction digested to completion to release the repeat-containing fragment and flanking

regions – this increases resolution by electrophoresis. Following neutral transfer, the *HTT*

containing regions were detected by Southern blot as described above.

RNA-seq library preparation, sequencing, and analysis

Total RNA from the lymphoblastoid cell lines was isolated using the hot TRIzol extraction

protocol (212). Briefly, samples dissolved in TRIzol were incubated in a thermomixer set to

55°C and 1,000 rpm for 10 min. After cooling to ambient temperature, RNA was isolated by

using TRIzol Reagent and the Direct-zol RNA MiniPrep Kit with DNase treatment according

to manufacturer's provided protocol (Zymo Research). RNA was quantitated using Qubit RNA

BR Assay Kit (Thermo Fisher Scientific). Strand-specific, rRNA-depleted RNA-seq libraries

were prepared using the KAPA Stranded RNA-seq Kit with RiboErase HMR (Kapa Biosystems) per manufacturer's instructions, except for the use of custom Illumina-compatible index primers to allow multiplexing. Library size distribution was assessed using the High Sensitivity NGS Fragment Analysis Kit (DNF-747) on a Fragment Analyzer (Agilent). 2×150 paired-end sequencing was performed using an Illumina NextSeq 500. Reads were aligned to the genome assembly GRCh38 (hg38). Salmon was used for transcript expression quantification and differential gene expression analysis was performed using DESeq2(213, 214). A heat map was generated in R using the ggplot2 package. Graph was generated and statistical analysis was performed by using GraphPad Prism 8.

Developing murine Tissue-Derived Cell Lines

Mouse tissues for cytogenetic analysis of chromosomes followed published protocols (128–131, 215–220). Mouse care, sacrifice, and tissue harvesting was done on the same day, including three wild type and three transgenic mice, all 5 months of age. Mice aged 5-months, when cell proliferation in the CNS and peripheral organs had ceased at 2-months of age (128–130). Ear, lung, tail, cerebrum, cerebellum, brain stem, kidney, liver, and spleen were harvested. A section of each tissue was manually macerated then treated with DMEM + Trypsin-EDTA for 1 hour in a 37°C incubator. Cells were collected, spun down, then seeded in a T25 flask with DMEM media supplemented with 20% FBS and 2 mM L-Glutamate. Two flasks were prepared: One with 1% Pen/Strep added, and one without. Once confluent, cells were seeded in a new T25 in DMEM (supplemented with 15% Fetal Bovine Serum, Penicillin/Streptomycin (100 IU/mL, 100 µg/mL), and 2 mM L-Glutamate to begin fragile site induction. To avoid cell type selection and cell type loss due to an inability to adapt to cell culture, harvested tissues were purposely cultured for very short terms following tissue harvesting. When possible, cells were grown to 80% confluency. Avoiding high cell densities, or long periods at confluency, has been shown to diminish FRAXA expression (217). Tissues

were harvested and immediately made into cultures. These cultures were grown and induced

to express the fragile sites and cytogenetically assessed. There was neither transit delays nor

storage of cell lines until fragile sites were induced in metaphase. All procedures, including

colony housing and tissue harvesting, were done in-house. Precautions were taken to avoid

delays following tissue harvesting, tissue freezing, extended culturing and storage of cells, as

these have been noted to diminish fragile site expression (217–220).

REFERENCES

1. Atanasio,A., Decman,V., White,D., Ramos,M., Ikiz,B., Lee,H.-C., Siao,C.-J., Brydges,S., LaRosa,E., Bai,Y., *et al.* (2016) C9orf72 ablation causes immune dysregulation characterized by leukocyte expansion, autoantibody production, and glomerulonephropathy in mice. *Sci Rep*, **6**, 23204.

2. Breevoort,S., Gibson,S., Figueroa,K., Bromberg,M. and Pulst,S. (2022) Expanding Clinical Spectrum of C9ORF72-Related Disorders and Promising Therapeutic Strategies: A Review. *Neurol Genet*, **8**, e670.

3. Fredi,M., Cavazzana,I., Biasiotto,G., Filosto,M., Padovani,A., Monti,E., Tincani,A., Franceschini,F. and Zanella,I. (2019) C9orf72 Intermediate Alleles in Patients with Amyotrophic Lateral Sclerosis, Systemic Lupus Erythematosus, and Rheumatoid Arthritis. *Neuromolecular Med*, **21**, 150–159.

4. McCauley,M.E., O'Rourke,J.G., Yáñez,A., Markman,J.L., Ho,R., Wang,X., Chen,S., Lall,D., Jin,M., Muhammad,A.K.M.G., *et al.* (2020) C9orf72 in myeloid cells suppresses STING-induced inflammation. *Nature*, **585**, 96–101.

5. Miller,Z.A., Sturm,V.E., Camsari,G.B., Karydas,A., Yokoyama,J.S., Grinberg,L.T., Boxer,A.L., Rosen,H.J., Rankin,K.P., Gorno-Tempini,M.L., *et al.* (2016) Increased prevalence of autoimmune disease within C9 and FTD/MND cohorts: Completing the picture. *Neurol Neuroimmunol Neuroinflamm*, **3**, e301.

6. van der Ende,E.L., Jackson,J.L., White,A., Seelaar,H., van Blitterswijk,M. and Van Swieten,J.C. (2021) Unravelling the clinical spectrum and the role of repeat length in C9ORF72 repeat expansions. *J Neurol Neurosurg Psychiatry*, **92**, 502–509.

7. Tábuas-Pereira,M., Almendra,L., Almeida,M.R., Durães,J., Pinho,A., Matos,A., Negrão,L., Geraldo,A. and Santana,I. (2019) Increased risk of melanoma in C9ORF72 repeat expansion carriers: A case-control study. *Muscle Nerve*, **59**, 362–365.

8. Chiò,A., Mazzini,L., D'Alfonso,S., Corrado,L., Canosa,A., Moglia,C., Manera,U., Bersano,E., Brunetti,M., Barberis,M., *et al.* (2018) The multistep hypothesis of ALS revisited: The role of genetic mutations. *Neurology*, **91**, e635–e642.

9. Van Wijk,I.F., Van Eijk,R.P.A., Van Boxmeer,L., Westeneng,H.-J., Van Es,M.A., Van Rheenen,W., Van Den Berg,L.H., Eijkemans,M.J.C. and Veldink,J.H. (2024) Assessment of risk of ALS conferred by the GGGGCC hexanucleotide repeat expansion in C9orf72 among first-degree relatives of patients with ALS carrying the repeat expansion. *Amyotroph Lateral Scler Frontotemporal Degener*, **25**, 188–196.

10. Waite,A.J., Bäumer,D., East,S., Neal,J., Morris,H.R., Ansorge,O. and Blake,D.J. (2014) Reduced C9orf72 protein levels in frontal cortex of amyotrophic lateral sclerosis and frontotemporal degeneration brain with the C9ORF72 hexanucleotide repeat expansion. *Neurobiol Aging*, **35**, 1779.e5-1779.e13.

11. Xiao,S., MacNair,L., McLean,J., McGoldrick,P., McKeever,P., Soleimani,S., Keith,J., Zinman,L., Rogaeva,E. and Robertson,J. (2016) C9orf72 isoforms in Amyotrophic Lateral Sclerosis and Frontotemporal Lobar Degeneration. *Brain Res.*, **1647**, 43–49.

12. Marques,C., Held,A., Dorfman,K., Sung,J., Song,C., Kavuturu,A.S., Aguilar,C., Russo,T.,
Oakley,D.H., Albers,M.W., *et al.* (2024) Neuronal STING activation in amyotrophic lateral
sclerosis and frontotemporal dementia. *Acta Neuropathol*, **147**, 56.

13. Pang,W. and Hu,F. (2023) C9ORF72 suppresses JAK-STAT mediated inflammation. *iScience*, **26**,
106579.

14. Masrori,P., Beckers,J., Gossye,H. and Van Damme,P. (2022) The role of inflammation in
neurodegeneration: novel insights into the role of the immune system in C9orf72 HRE-
mediated ALS/FTD. *Mol Neurodegener*, **17**, 22.

15. Wang,Y., Liu,L., Chen,H., Yang,Y., Mu,C., Ren,H., Liu,Y., Yu,L., Fang,Q., Wang,G., *et al.* (2023)
Disrupted phase behavior of FUS underlies poly-PR-induced DNA damage in amyotrophic
lateral sclerosis. *Hum Mol Genet*, **33**, 64–77.

16. Braems,E., Bercier,V., Van Schoor,E., Heeren,K., Beckers,J., Fumagalli,L., Dedeene,L., Moisse,M.,
Geudens,I., Hersmus,N., *et al.* (2022) HNRNPK alleviates RNA toxicity by counteracting DNA
damage in C9orf72 ALS. *Acta Neuropathol*, **144**, 465–488.

17. Konopka,A. and Atkin,J.D. (2022) DNA Damage, Defective DNA Repair, and Neurodegeneration in
Amyotrophic Lateral Sclerosis. *Front Aging Neurosci*, **14**, 786420.

18. Maor-Nof,M., Shipony,Z., Lopez-Gonzalez,R., Nakayama,L., Zhang,Y.-J., Couthouis,J., Blum,J.A.,
Castruita,P.A., Linares,G.R., Ruan,K., *et al.* (2021) p53 is a central regulator driving
neurodegeneration caused by C9orf72 poly(PR). *Cell*, **184**, 689-708.e20.

19. Andrade,N.S., Ramic,M., Esanov,R., Liu,W., Rybin,M.J., Gaidosh,G., Abdallah,A., Del'Olio,S.,
Huff,T.C., Chee,N.T., *et al.* (2020) Dipeptide repeat proteins inhibit homology-directed DNA
double strand break repair in C9ORF72 ALS/FTD. *Mol Neurodegener*, **15**, 13.

20. Nihei,Y., Mori,K., Werner,G., Arzberger,T., Zhou,Q., Khosravi,B., Japtok,J., Hermann,A.,
Sommacal,A., Weber,M., *et al.* (2020) Poly-glycine-alanine exacerbates C9orf72 repeat
expansion-mediated DNA damage via sequestration of phosphorylated ATM and loss of
nuclear hnRNPA3. *Acta Neuropathol*, **139**, 99–118.

21. Lopez-Gonzalez,R., Yang,D., Pribadi,M., Kim,T.S., Krishnan,G., Choi,S.Y., Lee,S., Coppola,G. and
Gao,F.-B. (2019) Partial inhibition of the overactivated Ku80-dependent DNA repair pathway
rescues neurodegeneration in C9ORF72-ALS/FTD. *Proc Natl Acad Sci U S A*, **116**, 9628–9633.

22. Konopka,A. and Atkin,J.D. (2018) The Emerging Role of DNA Damage in the Pathogenesis of the
C9orf72 Repeat Expansion in Amyotrophic Lateral Sclerosis. *Int J Mol Sci*, **19**, E3137.

23. Walker,C., Herranz-Martin,S., Karyka,E., Liao,C., Lewis,K., Elsayed,W., Lukashchuk,V., Chiang,S.-C.,
Ray,S., Mulcahy,P.J., *et al.* (2017) C9orf72 expansion disrupts ATM-mediated chromosomal
break repair. *Nat Neurosci*, **20**, 1225–1235.

24. Farg,M.A., Konopka,A., Soo,K.Y., Ito,D. and Atkin,J.D. (2017) The DNA damage response (DDR) is
induced by the C9orf72 repeat expansion in amyotrophic lateral sclerosis. *Hum Mol Genet*,
26, 2882–2896.

25. Lopez-Gonzalez,R., Lu,Y., Gendron,T.F., Karydas,A., Tran,H., Yang,D., Petrucelli,L., Miller,B.L.,
Almeida,S. and Gao,F.-B. (2016) Poly(GR) in C9ORF72-Related ALS/FTD Compromises

Mitochondrial Function and Increases Oxidative Stress and DNA Damage in iPSC-Derived
Motor Neurons. *Neuron*, **92**, 383–391.

26. van Blitterswijk,M., DeJesus-Hernandez,M., Niemantsverdriet,E., Murray,M.E., Heckman,M.G.,
Diehl,N.N., Brown,P.H., Baker,M.C., Finch,N.A., Bauer,P.O., *et al.* (2013) Association between
repeat sizes and clinical and pathological characteristics in carriers of C9ORF72 repeat
expansions (Xpansize-72): a cross-sectional cohort study. *The Lancet. Neurology*, **12**, 978–
988.

27. Nordin,A., Akimoto,C., Wuolikainen,A., Alstermark,H., Jonsson,P., Birve,A., Marklund,S.L.,
Graffmo,K.S., Forsberg,K., Brännström,T., *et al.* (2015) Extensive size variability of the
GGGGCC expansion in C9orf72 in both neuronal and non-neuronal tissues in 18 patients with
ALS or FTD. *Human Molecular Genetics*, **24**, 3133–3142.

28. Akimoto,C., Volk,A.E., van Blitterswijk,M., Van den Broeck,M., Leblond,C.S., Lumbroso,S.,
Camu,W., Neitzel,B., Onodera,O., van Rheenen,W., *et al.* (2014) A blinded international
study on the reliability of genetic testing for GGGGCC-repeat expansions in C9orf72 reveals
marked differences in results among 14 laboratories. *J Med Genet*, **51**, 419–424.

29. Xi,Z., van Blitterswijk,M., Zhang,M., McGoldrick,P., McLean,J.R., Yunusova,Y., Knock,E.,
Moreno,D., Sato,C., McKeever,P.M., *et al.* (2015) Jump from pre-mutation to pathologic
expansion in C9orf72. *Am J Hum Genet*, **96**, 962–970.

30. Xi,Z., Yunusova,Y., van Blitterswijk,M., Dib,S., Ghani,M., Moreno,D., Sato,C., Liang,Y.,
Singleton,A., Robertson,J., *et al.* (2014) Identical twins with the C9orf72 repeat expansion
are discordant for ALS. *Neurology*, **83**, 1476–1478.

31. Sattler,R., Traynor,B.J., Robertson,J., Van Den Bosch,L., Barmada,S.J., Svendsen,C.N.,
Disney,M.D., Gendron,T.F., Wong,P.C., Turner,M.R., *et al.* (2023) Roadmap for C9ORF72 in
Frontotemporal Dementia and Amyotrophic Lateral Sclerosis: Report on the C9ORF72
FTD/ALS Summit. *Neurol Ther*, **12**, 1821–1843.

32. Cammack,A.J., Balendra,R. and Isaacs,A.M. (2024) Failure of C9orf72 sense repeat-targeting
antisense oligonucleotides: lessons learned and the path forward. *Brain*, **147**, 2607–2609.

33. Petri,S. (2024) Targeting C9orf72 in people with ALS. *Lancet Neurol*, **23**, 850–852.

34. van den Berg,L.H., Rothstein,J.D., Shaw,P.J., Babu,S., Benatar,M., Bucelli,R.C., Genge,A.,
Glass,J.D., Hardiman,O., Libri,V., *et al.* (2024) Safety, tolerability, and pharmacokinetics of
antisense oligonucleotide BIIB078 in adults with C9orf72-associated amyotrophic lateral
sclerosis: a phase 1, randomised, double blinded, placebo-controlled, multiple ascending
dose study. *Lancet Neurol*, **23**, 901–912.

35. Wave Life Sciences Announces Topline Results from Phase 1b/2a FOCUS-C9 Study of WVE-004
for C9orf72-associated Amyotrophic Lateral Sclerosis and Frontotemporal Dementia - Wave
Life Sciences.

36. Tran,H., Moazami,M.P., Yang,H., McKenna-Yasek,D., Douthwright,C.L., Pinto,C., Metterville,J.,
Shin,M., Sanil,N., Dooley,C., *et al.* (2022) Suppression of mutant C9orf72 expression by a
potent mixed backbone antisense oligonucleotide. *Nat Med*, **28**, 117–124.

37. Saracino,D. and Le Ber,I. (2022) How can we define the presymptomatic C9orf72 disease in
1179   2022? An overview on the current definitions of preclinical and prodromal phases. *Rev*
1180   *Neurol (Paris)*, **178**, 426–436.

1181   38. Saracino,D. and Le Ber,I. (2021) Clinical Update on C9orf72: Frontotemporal Dementia,
1182   Amyotrophic Lateral Sclerosis, and Beyond. *Adv Exp Med Biol*, **1281**, 67–76.

1183   39. Mandrioli,J., Zucchi,E., Martinelli,I., Van der Most,L., Gianferrari,G., Moglia,C., Manera,U.,
1184   Solero,L., Vasta,R., Canosa,A., *et al.* (2023) Factors predicting disease progression in
1185   C9ORF72 ALS patients. *J Neurol*, **270**, 877–890.

1186   40. Westeneng,H.-J., van Veenhuijzen,K., van der Spek,R.A., Peters,S., Visser,A.E., van Rheenen,W.,
1187   Veldink,J.H. and van den Berg,L.H. (2021) Associations between lifestyle and amyotrophic
1188   lateral sclerosis stratified by C9orf72 genotype: a longitudinal, population-based, case-
1189   control study. *Lancet Neurol*, **20**, 373–384.

1190   41. Querin,G., Biferi,M.G. and Pradat,P.-F. (2022) Biomarkers for C9orf7-ALS in Symptomatic and
1191   Pre-symptomatic Patients: State-of-the-art in the New Era of Clinical Trials. *J Neuromuscul*
1192   *Dis*, **9**, 25–37.

1193   42. Rizzu,P., Blauwendraat,C., Heetveld,S., Lynes,E.M., Castillo-Lizardo,M., Dhingra,A., Pyz,E.,
1194   Hobert,M., Synofzik,M., Simón-Sánchez,J., *et al.* (2016) C9orf72 is differentially expressed in
1195   the central nervous system and myeloid cells and consistently reduced in C9orf72, MAPT
1196   and GRN mutation carriers. *Acta Neuropathol Commun*, **4**, 37.

1197   43. Niblock,M., Smith,B.N., Lee,Y.-B., Sardone,V., Topp,S., Troakes,C., Al-Sarraj,S., Leblond,C.S.,
1198   Dion,P.A., Rouleau,G.A., *et al.* (2016) Retention of hexanucleotide repeat-containing intron
1199   in C9orf72 mRNA: implications for the pathogenesis of ALS/FTD. *Acta Neuropathol Commun*,
1200   **4**, 18.

1201   44. Sznajder,Ł.J., Thomas,J.D., Carrell,E.M., Reid,T., McFarland,K.N., Cleary,J.D., Oliveira,R.,
1202   Nutter,C.A., Bhatt,K., Sobczak,K., *et al.* (2018) Intron retention induced by microsatellite
1203   expansions as a disease biomarker. *Proc Natl Acad Sci U S A*, **115**, 4234–4239.

1204   45. Ben-Dor,I., Pacut,C., Nevo,Y., Feldman,E.L. and Reubinoff,B.E. (2021) Characterization of C9orf72
1205   haplotypes to evaluate the effects of normal and pathological variations on its expression
1206   and splicing. *PLoS Genet*, **17**, e1009445.

1207   46. Sarkar,M.K., Hile,G.A., Tsoi,L.C., Xing,X., Liu,J., Liang,Y., Berthier,C.C., Swindell,W.R., Patrick,M.T.,
1208   Shao,S., *et al.* (2018) Photosensitivity and type I IFN responses in cutaneous lupus are driven
1209   by epidermal-derived interferon kappa. *Ann Rheum Dis*, **77**, 1653–1664.

1210   47. Harismendy,O., Notani,D., Song,X., Rahim,N.G., Tanasa,B., Heintzman,N., Ren,B., Fu,X.-D.,
1211   Topol,E.J., Rosenfeld,M.G., *et al.* (2011) 9p21 DNA variants associated with coronary artery
1212   disease impair interferon-γ signalling response. *Nature*, **470**, 264–268.

1213   48. Joung,J., Engreitz,J.M., Konermann,S., Abudayyeh,O.O., Verdine,V.K., Aguet,F., Gootenberg,J.S.,
1214   Sanjana,N.E., Wright,J.B., Fulco,C.P., *et al.* (2017) Genome-scale activation screen identifies a
1215   lncRNA locus regulating a gene neighbourhood. *Nature*, **548**, 343–346.

49. Ahel,I., Rass,U., El-Khamisy,S.F., Katyal,S., Clements,P.M., McKinnon,P.J., Caldecott,K.W. and
West,S.C. (2006) The neurodegenerative disease protein aprataxin resolves abortive DNA
ligation intermediates. *Nature*, **443**, 713–716.

50. Lim,G., Karaskova,J., Beheshti,B., Vukovic,B., Bayani,J., Selvarajah,S., Watson,S.K., Lam,W.L.,
Zielenska,M. and Squire,J.A. (2005) An integrated mBAND and submegabase resolution tiling
set (SMRT) CGH array analysis of focal amplification, microdeletions, and ladder structures
consistent with breakage-fusion-bridge cycle events in osteosarcoma. *Genes Chromosomes*
Cancer, **42**, 392–403.

51. Fletcher,J.A., Gebhardt,M.C. and Kozakewich,H.P. (1994) Cytogenetic aberrations in
osteosarcomas. Nonrandom deletions, rings, and double-minute chromosomes. *Cancer*
Genet Cytogenet, **77**, 81–88.

52. Giollant,M., Bertrand,S., Verrelle,P., Tchirkov,A., du Manoir,S., Ried,T., Mornex,F., Doré,J.F.,
Cremer,T. and Malet,P. (1996) Characterization of double minute chromosomes' DNA
content in a human high grade astrocytoma cell line by using comparative genomic
hybridization and fluorescence in situ hybridization. *Hum Genet*, **98**, 265–270.

53. Satinover,D.L., Vance,G.H., Van Dyke,D.L. and Schwartz,S. (2001) Cytogenetic analysis and
construction of a BAC contig across a common neocentromeric region from 9p.
Chromosoma, **110**, 275–283.

54. Jee,K.J., Kim,Y.T., Kim,K.R., Aalto,Y. and Knuutila,S. (2001) Amplification at 9p in cervical
carcinoma by comparative genomic hybridization. *Anal Cell Pathol*, **22**, 159–163.

55. Sams,E.I., Ng,J.K., Tate,V., Claire Hou,Y.-C., Cao,Y., Antonacci-Fulton,L., Belhassan,K., Neidich,J.,
Mitra,R.D., Cole,F.S., *et al.* (2022) From karyotypes to precision genomics in 9p deletion and
duplication syndromes. *HGG Adv*, **3**, 100081.

56. Shapiro,J.A. (2021) How Chaotic Is Genome Chaos? *Cancers (Basel)*, **13**, 1358.

57. Bignell,G.R., Greenman,C.D., Davies,H., Butler,A.P., Edkins,S., Andrews,J.M., Buck,G., Chen,L.,
Beare,D., Latimer,C., *et al.* (2010) Signatures of mutation and selection in the cancer
genome. *Nature*, **463**, 893–898.

58. Sutherland,G.R., Jacky,P.B., Baker,E. and Manuel,A. (1983) Heritable fragile sites on human
chromosomes. X. New folate-sensitive fragile sites: 6p23, 9p21, 9q32, and 11q23. *Am J Hum*
Genet, **35**, 432–437.

59. Mirceta,M., Shum,N., Schmidt,M.H.M. and Pearson,C.E. (2022) Fragile sites, chromosomal
lesions, tandem repeats, and disease. *Front Genet*, **13**, 985975.

60. Sun,J.H., Zhou,L., Emerson,D.J., Phyo,S.A., Titus,K.R., Gong,W., Gilgenast,T.G., Beagan,J.A.,
Davidson,B.L., Tassone,F., *et al.* (2018) Disease-Associated Short Tandem Repeats Co-localize
with Chromatin Domain Boundaries. *Cell*, **175**, 224-238.e15.

61. Xi,Z., Zhang,M., Bruni,A.C., Maletta,R.G., Colao,R., Fratta,P., Polke,J.M., Sweeney,M.G.,
Mudanohwo,E., Nacmias,B., *et al.* (2015) The C9orf72 repeat expansion itself is methylated
in ALS and FTLD patients. *Acta Neuropathol*, **129**, 715–727.

62. Xi,Z., Zinman,L., Moreno,D., Schymick,J., Liang,Y., Sato,C., Zheng,Y., Ghani,M., Dib,S., Keith,J., *et*
al. (2013) Hypermethylation of the CpG island near the G4C2 repeat in ALS with a C9orf72
expansion. *Am J Hum Genet*, **92**, 981–989.

63. Jackson,J.L., Finch,N.A., Baker,M.C., Kachergus,J.M., DeJesus-Hernandez,M., Pereira,K.,
Christopher,E., Prudencio,M., Heckman,M.G., Thompson,E.A., *et al.* (2020) Elevated
methylation levels, reduced expression levels, and frequent contractions in a clinical cohort
of C9orf72 expansion carriers. *Mol Neurodegeneration*, **15**, 7.

64. Xu,G.L., Bestor,T.H., Bourc'his,D., Hsieh,C.L., Tommerup,N., Bugge,M., Hulten,M., Qu,X.,
Russo,J.J. and Viegas-Péquignot,E. (1999) Chromosome instability and immunodeficiency
syndrome caused by mutations in a DNA methyltransferase gene. *Nature*, **402**, 187–191.

65. Hecht,F. and Sutherland,G.R. (1984) Detection of the fragile X chromosome and other fragile
sites. *Clin Genet*, **26**, 301–303.

66. Veenema,H., Beverstock,G.C., de Koning,T., Pearson,P.L. and van de Kamp,J.J. (1988) The fragile
X-chromosome: an evaluation of the results in a routine cytogenetic laboratory in the period
1981-1986. *Clin Genet*, **33**, 410–417.

67. Cantú,E.S. and Jacobs,P.A. (1984) Fragile (X) expression: relationship to the cell cycle. *Hum Genet*,
67, 99–102.

68. Yu,S., Mangelsdorf,M., Hewett,D., Hobson,L., Baker,E., Eyre,H.J., Lapsys,N., Le Paslier,D.,
Doggett,N.A., Sutherland,G.R., *et al.* (1997) Human chromosomal fragile site FRA16B is an
amplified AT-rich minisatellite repeat. *Cell*, **88**, 367–374.

69. Nancarrow,J.K., Kremer,E., Holman,K., Eyre,H., Doggett,N.A., Le Paslier,D., Callen,D.F.,
Sutherland,G.R. and Richards,R.I. (1994) Implications of FRA16A structure for the mechanism
of chromosomal fragile site genesis. *Science*, **264**, 1938–1941.

70. Bosco,N., Pelliccia,F. and Rocchi,A. (2010) Characterization of FRA7B, a human common fragile
site mapped at the 7p chromosome terminal region. *Cancer Genet Cytogenet*, **202**, 47–52.

71. Callahan,G., Denison,S.R., Phillips,L.A., Shridhar,V. and Smith,D.I. (2003) Characterization of the
common fragile site FRA9E and its potential role in ovarian cancer. *Oncogene*, **22**, 590–601.

72. Curatolo,A., Limongi,Z.M., Pelliccia,F. and Rocchi,A. (2007) Molecular characterization of the
human common fragile site FRA1H. *Genes Chromosomes Cancer*, **46**, 487–493.

73. Rozier,L., El-Achkar,E., Apiou,F. and Debatisse,M. (2004) Characterization of a conserved
aphidicolin-sensitive common fragile site at human 4q22 and mouse 6C1: possible
association with an inherited disease and cancer. *Oncogene*, **23**, 6872–6880.

74. Abruzzo,M.A., Hunt,P.A., Mayer,M., Jacobs,P.A., Wang,J.C. and Erbe,R.W. (1986) A comparison of
fragile X expression in lymphocyte and lymphoblastoid cultures. *Am J Hum Genet*, **38**, 533–
539.

75. Sutherland,G.R., Baker,E. and Callen,D.F. (1988) A BrdU-enhanceable fragile site or viral
modification site at 11q23.1 in lymphoblastoid cultures. *Cytogenet Cell Genet*, **47**, 201–203.

76. Seki,N., Tsuji,H., Takahashi,E., Yamauchi,M., Saito,T., Hashimoto,T., Yamamoto,K. and Hori,T.
(1992) Induction of a BrdU-enhanceable fragile site-like lesion and sister chromatid
exchanges at 11q23.1 in EBV-transformed lymphoblastoid cell lines. *Cytogenet Cell Genet*,
61, 95–98.

77. Sutherland,G.R. and Baker,E. (2003) Forgotten fragile sites and related phenomena. *Cytogenet*
Genome Res, **100**, 89–91.

78. Li,J.S.Z., Abbasi,A., Kim,D.H., Lippman,S.M., Alexandrov,L.B. and Cleveland,D.W. (2023)
Chromosomal fragile site breakage by EBV-encoded EBNA1 at clustered repeats. *Nature*,
616, 504–509.

79. McGoldrick,P., Zhang,M., van Blitterswijk,M., Sato,C., Moreno,D., Xiao,S., Zhang,A.B.,
McKeever,P.M., Weichert,A., Schneider,R., *et al.* (2018) Unaffected mosaic C9orf72 case:
RNA foci, dipeptide proteins, but upregulated C9orf72 expression. *Neurology*, **90**, e323–
e331.

80. Belzil,V.V., Bauer,P.O., Prudencio,M., Gendron,T.F., Stetler,C.T., Yan,I.K., Pregent,L., Daughrity,L.,
Baker,M.C., Rademakers,R., *et al.* (2013) Reduced C9orf72 gene expression in c9FTD/ALS is
caused by histone trimethylation, an epigenetic event detectable in blood. *Acta*
Neuropathol, **126**, 895–905.

81. Cali,C.P., Patino,M., Tai,Y.K., Ho,W.Y., McLean,C.A., Morris,C.M., Seeley,W.W., Miller,B.L.,
Gaig,C., Vonsattel,J.P.G., *et al.* (2019) C9orf72 intermediate repeats are associated with
corticobasal degeneration, increased C9orf72 expression and disruption of autophagy. *Acta*
Neuropathol, **138**, 795–811.

82. Tassone,F., Hagerman,R.J., Taylor,A.K., Gane,L.W., Godfrey,T.E. and Hagerman,P.J. (2000)
Elevated Levels of FMR1 mRNA in Carrier Males: A New Mechanism of Involvement in the
Fragile-X Syndrome. *The American Journal of Human Genetics*, **66**, 6–15.

83. Eberhart,D.E. and Warren,S.T. (1996) Nuclease sensitivity of permeabilized cells confirms altered
chromatin formation at the fragile X locus. *Somat. Cell Mol. Genet.*, **22**, 435–441.

84. Beverstock,G.C., Mol,A. and Wienhofer,E. (1985) Absence of significant autosomal lesions in
Huntington's disease. *Ann Hum Genet*, **49**, 283–290.

85. Krupina,K., Goginashvili,A. and Cleveland,D.W. (2021) Causes and consequences of micronuclei.
Curr Opin Cell Biol, **70**, 91–99.

86. Utani,K., Okamoto,A. and Shimizu,N. (2011) Generation of micronuclei during interphase by
coupling between cytoplasmic membrane blebbing and nuclear budding. *PLoS One*, **6**,
e27233.

87. Fenech,M., Kirsch-Volders,M., Natarajan,A.T., Surralles,J., Crott,J.W., Parry,J., Norppa,H.,
Eastmond,D.A., Tucker,J.D. and Thomas,P. (2011) Molecular mechanisms of micronucleus,
nucleoplasmic bridge and nuclear bud formation in mammalian and human cells.
Mutagenesis, **26**, 125–132.

88. Fenech,M. (2007) Cytokinesis-block micronucleus cytome assay. *Nat Protoc*, **2**, 1084–1104.

89. Shimizu,N., Itoh,N., Utiyama,H. and Wahl,G.M. (1998) Selective entrapment of
extrachromosomally amplified DNA by nuclear budding and micronucleation during S phase.
J Cell Biol, **140**, 1307–1320.

90. Guttenbach,M. and Schmid,M. (1994) Exclusion of specific human chromosomes into micronuclei
by 5-azacytidine treatment of lymphocyte cultures. *Exp Cell Res*, **211**, 127–132.

91. Bjerregaard,V.A., Garribba,L., McMurray,C.T., Hickson,I.D. and Liu,Y. (2018) Folate deficiency
drives mitotic missegregation of the human FRAXA locus. *Proc Natl Acad Sci U S A*, **115**,
13003–13008.

92. Duncan,A.M. (1986) Enhanced sensitivity of lymphoblastoid cells from individuals carrying the
mutation for the fragile X syndrome to the clastogenic effects of FUdR. *Mutat Res*, **173**, 201–
205.

93. Shah,P., Wolf,K. and Lammerding,J. (2017) Bursting the Bubble - Nuclear Envelope Rupture as a
Path to Genomic Instability? *Trends Cell Biol*, **27**, 546–555.

94. Srivastava,N., Nader,G.P. de F., Williart,A., Rollin,R., Cuvelier,D., Lomakin,A. and Piel,M. (2021)
Nuclear fragility, blaming the blebs. *Curr Opin Cell Biol*, **70**, 100–108.

95. Stephens,A.D., Liu,P.Z., Banigan,E.J., Almassalha,L.M., Backman,V., Adam,S.A., Goldman,R.D. and
Marko,J.F. (2018) Chromatin histone modifications and rigidity affect nuclear morphology
independent of lamins. *Mol Biol Cell*, **29**, 220–233.

96. Bercht Pfleghaar,K., Taimen,P., Butin-Israeli,V., Shimi,T., Langer-Freitag,S., Markaki,Y.,
Goldman,A.E., Wehnert,M. and Goldman,R.D. (2015) Gene-rich chromosomal regions are
preferentially localized in the lamin B deficient nuclear blebs of atypical progeria cells.
Nucleus, **6**, 66–76.

97. Niedernhofer,L.J., Gurkar,A.U., Wang,Y., Vijg,J., Hoeijmakers,J.H.J. and Robbins,P.D. (2018)
Nuclear Genomic Instability and Aging. *Annu Rev Biochem*, **87**, 295–322.

98. Fitchett,M. and Seabright,M. (1984) Deleted X chromosomes in patients with the fragile X
syndrome. *J Med Genet*, **21**, 373.

99. Verdyck,P., Berckmoes,V., De Vos,A., Verpoest,W., Liebaers,I., Bonduelle,M. and De Rycke,M.
(2015) Chromosome fragility at FRAXA in human cleavage stage embryos at risk for fragile X
syndrome. *Am J Med Genet A*, **167A**, 2306–2313.

100. Polleys,E.J. and Freudenreich,C.H. (2018) Methods to Study Repeat Fragility and Instability in
Saccharomyces cerevisiae. *Methods Mol Biol*, **1672**, 403–419.

101. Wenger,S.L., Hennessey,J.C. and Steele,M.W. (1987) Increased sister chromatid exchange
frequency at Xq27 site in affected fragile X males. *Am J Med Genet*, **26**, 909–914.

102. van Wietmarschen,N., Merzouk,S., Halsema,N., Spierings,D.C.J., Guryev,V. and Lansdorp,P.M.
(2018) BLM helicase suppresses recombination at G-quadruplex motifs in transcribed genes.
Nat Commun, **9**, 271.

103. Chan,K.L., Palmai-Pallag,T., Ying,S. and Hickson,I.D. (2009) Replication stress induces sister-
chromatid bridging at fragile site loci in mitosis. *Nat Cell Biol*, **11**, 753–760.

104. Chaganti,R.S., Schonberg,S. and German,J. (1974) A manyfold increase in sister chromatid
exchanges in Bloom's syndrome lymphocytes. *Proc Natl Acad Sci U S A*, **71**, 4508–4512.

105. Cunniff,C., Bassetti,J.A. and Ellis,N.A. (2017) Bloom's Syndrome: Clinical Spectrum, Molecular
Pathogenesis, and Cancer Predisposition. *Mol Syndromol*, **8**, 4–23.

106. Gratia,M., Rodero,M.P., Conrad,C., Bou Samra,E., Maurin,M., Rice,G.I., Duffy,D., Revy,P.,
Petit,F., Dale,R.C., *et al.* (2019) Bloom syndrome protein restrains innate immune sensing of
micronuclei by cGAS. *J Exp Med*, **216**, 1199–1213.

107. Suzuki,T., Yasui,M. and Honma,M. (2016) Mutator Phenotype and DNA Double-Strand Break
Repair in BLM Helicase-Deficient Human Cells. *Mol Cell Biol*, **36**, 2877–2889.

108. Warren,S.T., Schultz,R.A., Chang,C.C., Wade,M.H. and Trosko,J.E. (1981) Elevated spontaneous
mutation rate in Bloom syndrome fibroblasts. *Proc Natl Acad Sci U S A*, **78**, 3133–3137.

109. Wenger,S.L. (1995) Chemical induction of sister chromatid exchange at fragile sites. *Cancer
Genet Cytogenet*, **85**, 72–74.

110. Kumar,J.V., Saraswathi,T., Ranganathan,K., Umadevi,K., Joshua,E. and Rooban,T. (2012) Sister
chromatid exchanges in smokers and smokers with alcohol habit. *J Oral Maxillofac Pathol*,
16, 338–342.

111. Glover,T.W. and Stein,C.K. (1987) Induction of sister chromatid exchanges at common fragile
sites. *Am J Hum Genet*, **41**, 882–890.

112. Fuster,C., Mirò,R., Templado,C., Barrios,L. and Egozcue,J. (1988) Can sister chromatid
intercrossings be considered as prelesions? *Hum Genet*, **79**, 179–180.

113. Gaddini,L., Pelliccia,F., Limongi,M.Z. and Rocchi,A. (1995) Study of the relationships between
common fragile sites, chromosome breakages and sister chromatid exchanges. *Mutagenesis*,
10, 257–260.

114. Gregory,P., Wang,N. and Howard-Peebles,P.N. (1986) Analysis of sister chromatid exchanges in
fra (X) individuals. *Am J Med Genet*, **23**, 563–566.

115. Branda,R.F., Arthur,D.C., Woods,W.G., Danzl,T.J. and King,R.A. (1984) Folate metabolism and
chromosomal stability in the fragile X syndrome. *Am J Med*, **77**, 602–611.

116. Suleĭmanova,D.G. and Kuleshov,N.P. (1987) [Spontaneous and induced chromosome instability
in patients with fragile X syndrome]. *Genetika*, **23**, 504–509.

117. Schmid,M., Feichtinger,W. and Haaf,T. (1987) The fragile site (16)(q22). II. Sister chromatid
exchanges. *Hum Genet*, **76**, 365–368.

118. Hirsch,B. (1991) Sister chromatid exchanges are preferentially induced at expressed and
nonexpressed common fragile sites. *Hum Genet*, **87**, 302–306.

119. Lukusa,T., Meulepas,E., Fryns,J.P., Van den Berghe,H. and Cassiman,J.J. (1991) 'Spontaneous'
FRA16B is a hot spot for sister chromatid exchanges. *Hum Genet*, **87**, 583–586.

120. Recalcati,M.P., Bellini,M., Norsa,L., Ballarati,L., Caselli,R., Russo,S., Larizza,L. and Giardino,D. (2012) Complex rearrangement involving 9p deletion and duplication in a syndromic patient: genotype/phenotype correlation and review of the literature. *Gene*, **502**, 40–45.

121. Korbel,J.O. and Campbell,P.J. (2013) Criteria for inference of chromothripsis in cancer genomes. *Cell*, **152**, 1226–1236.

122. Kinsella,M., Patel,A. and Bafna,V. (2014) The elusive evidence for chromothripsis. *Nucleic Acids Res*, **42**, 8231–8242.

123. Krupina,K., Goginashvili,A. and Cleveland,D.W. (2024) Scrambling the genome in cancer: causes and consequences of complex chromosome rearrangements. *Nat Rev Genet*, **25**, 196–210.

124. Zhang,C.-Z., Spektor,A., Cornils,H., Francis,J.M., Jackson,E.K., Liu,S., Meyerson,M. and Pellman,D. (2015) Chromothripsis from DNA damage in micronuclei. *Nature*, **522**, 179–184.

125. Leibowitz,M.L., Papathanasiou,S., Doerfler,P.A., Blaine,L.J., Sun,L., Yao,Y., Zhang,C.-Z., Weiss,M.J. and Pellman,D. (2021) Chromothripsis as an on-target consequence of CRISPR-Cas9 genome editing. *Nat Genet*, **53**, 895–905.

126. Urnov,F.D. (2021) CRISPR-Cas9 can cause chromothripsis. *Nat Genet*, **53**, 768–769.

127. Pattamatta,A., Nguyen,L., Olafson,H.R., Scotti,M.M., Laboissonniere,L.A., Richardson,J., Berglund,J.A., Zu,T., Wang,E.T. and Ranum,L.P.W. (2021) Repeat length increases disease penetrance and severity in C9orf72 ALS/FTD BAC transgenic mice. *Hum Mol Genet*, **29**, 3900–3918.

128. Semenov,M. (2021) Proliferative Capacity of Adult Mouse Brain. *Int J Mol Sci*, **22**, 3449.

129. Bordiuk,O.L., Smith,K., Morin,P.J. and Semënov,M.V. (2014) Cell proliferation and neurogenesis in adult mouse brain. *PLoS One*, **9**, e111453.

130. Lui,J.C. and Baron,J. (2011) Mechanisms limiting body growth in mammals. *Endocr Rev*, **32**, 422–440.

131. Sanz,M.M., Jenkins,E.C., Brown,W.T., Davisson,M.T., Kevin,M.J., Roderick,T.H., Silverman,W.P. and Wisniewski,H.M. (1986) Mouse chromosome fragility. *Am J Med Genet*, **23**, 491–509.

132. Kähkönen,M., Tengström,C., Alitalo,T., Matilainen,R., Kaski,M. and Airaksinen,E. (1989) Population cytogenetics of folate-sensitive fragile sites. II. Autosomal rare fragile sites. *Hum Genet*, **82**, 3–8.

133. Vasta,R., Chia,R., Traynor,B.J. and Chiò,A. (2022) Unraveling the complex interplay between genes, environment, and climate in ALS. *EBioMedicine*, **75**, 103795.

134. Goncharova,P.S., Davydova,T.K., Popova,T.E., Novitsky,M.A., Petrova,M.M., Gavrilyuk,O.A., Al-Zamil,M., Zhukova,N.G., Nasyrova,R.F. and Shnayder,N.A. (2021) Nutrient Effects on Motor Neurons and the Risk of Amyotrophic Lateral Sclerosis. *Nutrients*, **13**, 3804.

135. Cingam,S.R., Koshy,N., Veillon,D. and Peddi,P. (2017) Reversal of isolated 20q deletion with vitamin B12 replacement in a patient with pernicious anaemia. *BMJ Case Rep*, **2017**, bcr2016218689.

136. Wollman,M.R., Penchansky,L. and Shekhter-Levin,S. (1996) Transient 7q- in association with
megaloblastic anemia due to dietary folate and vitamin B12 deficiency. *J Pediatr Hematol*
Oncol, **18**, 162–165.

137. Goh,K. (1981) Vitamin B12 deficiency in an 18p--patient. *Arch Pathol Lab Med*, **105**, 164.

138. Parmentier,S., Meinel,J., Oelschlaegel,U., Mohr,B., Ehninger,G., Schaich,M. and Platzbecker,U.
(2012) Severe pernicious anemia with distinct cytogenetic and flow cytometric aberrations
mimicking myelodysplastic syndrome. *Ann Hematol*, **91**, 1979–1981.

139. Chintagumpala,M.M., Dreyer,Z.A., Steuber,C.P. and Cooley,L.D. (1996) Pancytopenia with
chromosomal fragility: vitamin B12 deficiency. *J Pediatr Hematol Oncol*, **18**, 166–170.

140. Maltby,E.L. and Higgins,S. (1987) Folate sensitive site at 10q23 and its expression as a deletion.
J Med Genet, **24**, 299.

141. Morel,C.F., Duncan,A.M.V. and Désilets,V. (2005) A fragile site at 10q23 (FRA10A) in a
phenytoin-exposed fetus: a case report and review of the literature. *Prenat Diagn*, **25**, 318–
321.

142. De Leon-Luis,J., Santolaya-Forgas,J., May,G., Tonk,V., Shelton,D. and Galan,I. (2005) Prenatal
diagnosis of FRA10A: a case report and literature review. *Am J Med Genet A*, **136**, 63–65.

143. Ozisik,Y.Y., Meloni,A.M., Stone,J.F., Sandberg,A.A. and Surti,U. (1994) Spontaneous expression
of the chromosome fragile site at 10q23 in leiomyoma. *Cancer Genet Cytogenet*, **74**, 73–75.

144. Sarafidou,T., Kahl,C., Martinez-Garay,I., Mangelsdorf,M., Gesk,S., Baker,E., Kokkinaki,M.,
Talley,P., Maltby,E.L., French,L., *et al.* (2004) Folate-sensitive fragile site FRA10A is due to an
expansion of a CGG repeat in a novel gene, FRA10AC1, encoding a nuclear protein.
Genomics, **84**, 69–81.

145. Sashindran,V.K., Aggarwal,V. and Khera,A. (2022) Prevalence of Vitamin B12 deficiency in
elderly population (>60 years) presenting with dementia to outpatient department. *Med J*
Armed Forces India, **78**, 94–98.

146. Pinto,W.B.V. de R., Souza,P.V.S. de, Rogério,R.M., Pedroso,J.L. and Barsottini,O.G.P. (2014)
Vitamin B12 deficiency mimicking neuroimaging features of motor neuron disease. *Arq*
Neuropsiquiatr, **72**, 85.

147. Chen,A.T., Reidy,J.A., Annest,J.L., Welty,T.K. and Zhou,H.G. (1989) Increased chromosome
fragility as a consequence of blood folate levels, smoking status, and coffee consumption.
Environ Mol Mutagen, **13**, 319–324.

148. Barale,R., Chelotti,L., Davini,T., Del Ry,S., Andreassi,M.G., Ballardin,M., Bulleri,M., He,J.,
Baldacci,S., Di Pede,F., *et al.* (1998) Sister chromatid exchange and micronucleus frequency
in human lymphocytes of 1,650 subjects in an Italian population: II. Contribution of sex, age,
and lifestyle. *Environ Mol Mutagen*, **31**, 228–242.

149. Maffei,F., Fimognari,C., Castelli,E., Stefanini,G.F., Forti,G.C. and Hrelia,P. (2000) Increased
cytogenetic damage detected by FISH analysis on micronuclei in peripheral lymphocytes
from alcoholics. *Mutagenesis*, **15**, 517–523.

150. Demirhan,O. and Tastemir,D. (2008) Cytogenetic effects of ethanol on chronic alcohol users.
Alcohol Alcohol, **43**, 127–136.

151. Ueda,K., Sakai,C., Ishida,T., Morita,K., Kobayashi,Y., Horikoshi,Y., Baba,A., Okazaki,Y.,
Yoshizumi,M., Tashiro,S., *et al.* (2023) Cigarette smoke induces mitochondrial DNA damage
and activates cGAS-STING pathway: application to a biomarker for atherosclerosis. *Clin Sci*
(Lond), **137**, 163–180.

152. Hesdorffer,C.S. and Longo,D.L. (2015) Drug-Induced Megaloblastic Anemia. *N Engl J Med*, **373**,
1649–1658.

153. Czuppa,M., Dhingra,A., Zhou,Q., Schludi,C., König,L., Scharf,E., Farny,D., Dalmia,A., Täger,J.,
Castillo-Lizardo,M., *et al.* (2022) Drug screen in iPSC-Neurons identifies nucleoside analogs as
inhibitors of (G4C2)n expression in C9orf72 ALS/FTD. *Cell Rep*, **39**, 110913.

154. Chiurazzi,P., Pomponi,M.G., Pietrobono,R., Bakker,C.E., Neri,G. and Oostra,B.A. (1999)
Synergistic effect of histone hyperacetylation and DNA demethylation in the reactivation of
the FMR1 gene. *Hum Mol Genet*, **8**, 2317–2323.

155. Lamparska,K., Clark,J., Babilonia,G., Bedell,V., Yip,W. and Smith,S.S. (2012) 2'-
Deoxyriboguanylurea, the primary breakdown product of 5-aza-2'-deoxyribocytidine, is a
mutagen, an epimutagen, an inhibitor of DNA methyltransferases and an inducer of 5-
azacytidine-type fragile sites. *Nucleic Acids Res*, **40**, 9788–9801.

156. Feichtinger,W. and Schmid,M. (1989) Increased frequencies of sister chromatid exchanges at
common fragile sites (1)(q42) and (19)(q13). *Hum Genet*, **83**, 145–147.

157. Rodríguez-Reyes,R. and Morales-Ramírez,P. (2011) The in vivo induction of sister chromatid
exchange by the demethylating agent 5-aza-2'-deoxycytidine. *Mutagenesis*, **26**, 551–554.

158. Haaf,T., Ott,G. and Schmid,M. (1986) Differential inhibition of sister chromatid condensation
induced by 5-azadeoxycytidine in human chromosomes. *Chromosoma*, **94**, 389–394.

159. Davidson,S., Crowther,P., Radley,J. and Woodcock,D. (1992) Cytotoxicity of 5-aza-2'-
deoxycytidine in a mammalian cell system. *Eur J Cancer*, **28**, 362–368.

160. Gijselinck,I., Van Mossevelde,S., van der Zee,J., Sieben,A., Engelborghs,S., De Bleecker,J.,
Ivanoiu,A., Deryck,O., Edbauer,D., Zhang,M., *et al.* (2016) The C9orf72 repeat size correlates
with onset age of disease, DNA methylation and transcriptional downregulation of the
promoter. *Mol. Psychiatry*, **21**, 1112–1124.

161. Kojak,N., Kuno,J., Fittipaldi,K.E., Khan,A., Wenger,D., Glasser,M., Donnianni,R.A., Tang,Y.,
Zhang,J., Huling,K., *et al.* (2024) Somatic and intergenerational G4C2 hexanucleotide repeat
instability in a human C9orf72 knock-in mouse model. *Nucleic Acids Res*, **52**, 5732–5755.

162. Russ,J., Liu,E.Y., Wu,K., Neal,D., Suh,E., Irwin,D.J., McMillan,C.T., Harms,M.B., Cairns,N.J.,
Wood,E.M., *et al.* (2015) Hypermethylation of repeat expanded C9orf72 is a clinical and
molecular disease modifier. *Acta neuropathologica*, **129**, 39–52.

163. Zhang,M., Tartaglia,M.C., Moreno,D., Sato,C., McKeever,P., Weichert,A., Keith,J., Robertson,J.,
Zinman,L. and Rogaeva,E. (2017) DNA methylation age-acceleration is associated with
disease duration and age at onset in C9orf72 patients. *Acta Neuropathol.*, **134**, 271–279.

164. Zhang,M., McKeever,P.M., Xi,Z., Moreno,D., Sato,C., Bergsma,T., McGoldrick,P., Keith,J.,
Robertson,J., Zinman,L., *et al.* (2020) DNA methylation age acceleration is associated with
ALS age of onset and survival. *Acta Neuropathol*, **139**, 943–946.

165. Galkin,F., Kovalchuk,O., Koldasbayeva,D., Zhavoronkov,A. and Bischof,E. (2023) Stress, diet,
exercise: Common environmental factors and their impact on epigenetic age. *Ageing Res*
Rev, **88**, 101956.

166. Fitzgerald,K.N., Hodges,R., Hanes,D., Stack,E., Cheishvili,D., Szyf,M., Henkel,J., Twedt,M.W.,
Giannopoulou,D., Herdell,J., *et al.* (2021) Potential reversal of epigenetic age using a diet and
lifestyle intervention: a pilot randomized clinical trial. *Aging (Albany NY)*, **13**, 9419–9432.

167. Sae-Lee,C., Corsi,S., Barrow,T.M., Kuhnle,G.G.C., Bollati,V., Mathers,J.C. and Byun,H.-M. (2018)
Dietary Intervention Modifies DNA Methylation Age Assessed by the Epigenetic Clock. *Mol*
Nutr Food Res, **62**, e1800092.

168. Liu,Y., Huang,Z., Liu,H., Ji,Z., Arora,A., Cai,D., Wang,H., Liu,M., Simko,E.A.J., Zhang,Y., *et al.*
(2023) DNA-initiated epigenetic cascades driven by C9orf72 hexanucleotide repeat. *Neuron*,
111, 1205-1221.e9.

169. Rapkin,L.M., Ahmed,K., Dulev,S., Li,R., Kimura,H., Ishov,A.M. and Bazett-Jones,D.P. (2015) The
histone chaperone DAXX maintains the structural organization of heterochromatin domains.
Epigenetics Chromatin, **8**, 44.

170. He,Q., Kim,H., Huang,R., Lu,W., Tang,M., Shi,F., Yang,D., Zhang,X., Huang,J., Liu,D., *et al.* (2015)
The Daxx/Atrx Complex Protects Tandem Repetitive Elements during DNA Hypomethylation
by Promoting H3K9 Trimethylation. *Cell Stem Cell*, **17**, 273–286.

171. Scott,W.A., Dhanji,E.Z., Dyakov,B.J.A., Dreseris,E.S., Asa,J.S., Grange,L.J., Mirceta,M.,
Pearson,C.E., Stewart,G.S., Gingras,A.-C., *et al.* (2021) ATRX proximal protein associations
boast roles beyond histone deposition. *PLoS Genet*, **17**, e1009909.

172. Burgess,R.C., Burman,B., Kruhlak,M.J. and Misteli,T. (2014) Activation of DNA damage response
signaling by condensed chromatin. *Cell Rep*, **9**, 1703–1717.

173. Jacome,A. and Fernandez-Capetillo,O. (2011) Lac operator repeats generate a traceable fragile
site in mammalian cells. *EMBO Rep*, **12**, 1032–1038.

174. Soutoglou,E., Dorn,J.F., Sengupta,K., Jasin,M., Nussenzweig,A., Ried,T., Danuser,G. and
Misteli,T. (2007) Positional stability of single double-strand breaks in mammalian cells. *Nat*
Cell Biol, **9**, 675–682.

175. Soutoglou,E. and Misteli,T. (2008) Activation of the cellular DNA damage response in the
absence of DNA lesions. *Science*, **320**, 1507–1510.

176. Freaney,J.E., Zhang,Q., Yigit,E., Kim,R., Widom,J., Wang,J.-P. and Horvath,C.M. (2014) High-
density nucleosome occupancy map of human chromosome 9p21-22 reveals chromatin
organization of the type I interferon gene cluster. *J Interferon Cytokine Res*, **34**, 676–685.

177. Apostolou,E. and Thanos,D. (2008) Virus Infection Induces NF-kappaB-dependent
interchromosomal associations mediating monoallelic IFN-beta gene expression. *Cell*, **134**,
85–96.

178. Pang,W. and Hu,F. (2021) Cellular and physiological functions of C9ORF72 and implications for ALS/FTD. *J Neurochem*, **157**, 334–350.

179. Quinodoz,S.A., Ollikainen,N., Tabak,B., Palla,A., Schmidt,J.M., Detmar,E., Lai,M.M., Shishkin,A.A., Bhat,P., Takei,Y., *et al.* (2018) Higher-Order Inter-chromosomal Hubs Shape 3D Genome Organization in the Nucleus. *Cell*, **174**, 744-757.e24.

180. Jagannathan,M. and Yamashita,Y.M. (2017) Function of Junk: Pericentromeric Satellite DNA in Chromosome Maintenance. *Cold Spring Harb Symp Quant Biol*, **82**, 319–327.

181. Jagannathan,M., Cummings,R. and Yamashita,Y.M. (2018) A conserved function for pericentromeric satellite DNA. *Elife*, **7**, e34122.

182. Malachowski,T., Chandradoss,K.R., Boya,R., Zhou,L., Cook,A.L., Su,C., Pham,K., Haws,S.A., Kim,J.H., Ryu,H.-S., *et al.* (2023) Spatially coordinated heterochromatinization of long synaptic genes in fragile X syndrome. *Cell*, **186**, 5840-5858.e36.

183. Droppelmann,C.A., Campos-Melo,D., Moszczynski,A.J., Amzil,H. and Strong,M.J. (2019) TDP-43 aggregation inside micronuclei reveals a potential mechanism for protein inclusion formation in ALS. *Sci Rep*, **9**, 19928.

184. Fritsch,L.E., Ju,J., Gudenschwager Basso,E.K., Soliman,E., Paul,S., Chen,J., Kaloss,A.M., Kowalski,E.A., Tuhy,T.C., Somaiya,R.D., *et al.* (2022) Type I Interferon Response Is Mediated by NLRX1-cGAS-STING Signaling in Brain Injury. *Front Mol Neurosci*, **15**, 852243.

185. Song,X., Ma,F. and Herrup,K. (2019) Accumulation of Cytoplasmic DNA Due to ATM Deficiency Activates the Microglial Viral Response System with Neurotoxic Consequences. *J Neurosci*, **39**, 6378–6394.

186. Quek,H., Luff,J., Cheung,K., Kozlov,S., Gatei,M., Lee,C.S., Bellingham,M.C., Noakes,P.G., Lim,Y.C., Barnett,N.L., *et al.* (2017) Rats with a missense mutation in Atm display neuroinflammation and neurodegeneration subsequent to accumulation of cytosolic DNA following unrepaired DNA damage. *J Leukoc Biol*, **101**, 927–947.

187. Lupski,J.R. (2023) Somatic cell structural variant mutagenesis and neurologic disease. *Cell Genom*, **3**, 100376.

188. Schuy,J., Grochowski,C.M., Carvalho,C.M.B. and Lindstrand,A. (2022) Complex genomic rearrangements: an underestimated cause of rare diseases. *Trends Genet*, **38**, 1134–1146.

189. Campbell,I.M., Shaw,C.A., Stankiewicz,P. and Lupski,J.R. (2015) Somatic mosaicism: implications for disease and transmission genetics. *Trends Genet*, **31**, 382–392.

190. Coffee,B., Ikeda,M., Budimirovic,D.B., Hjelm,L.N., Kaufmann,W.E. and Warren,S.T. (2008) Mosaic FMR1 deletion causes fragile X syndrome and can lead to molecular misdiagnosis: a case report and review of the literature. *Am J Med Genet A*, **146A**, 1358–1367.

191. Gómez-Rodríguez,M.J., Morales-Conejo,M., Arteche-López,A., Sánchez-Calvín,M.T., Quesada-Espinosa,J.F., Gómez-Manjón,I., Palma-Milla,C., Lezana-Rosales,J.M., Pérez de la Fuente,R., Martin-Ramos,M.-L., *et al.* (2022) Fragile X Syndrome Caused by Maternal Somatic Mosaicism of FMR1 Gene: Case Report and Literature Review. *Genes (Basel)*, **13**, 1609.

192. Gonçalves,T.F., dos Santos,J.M., Gonçalves,A.P., Tassone,F., Mendoza-Morales,G., Ribeiro,M.G.,
Kahn,E., Boy,R., Pimentel,M.M.G. and Santos-Rebouças,C.B. (2016) Finding FMR1 mosaicism
in Fragile X syndrome. *Expert Rev Mol Diagn*, **16**, 501–507.

193. Hensel,C.H., Vanzo,R.J., Martin,M.M., Ling,L., Aliaga,S.M., Bui,M., Francis,D.I., Twede,H.,
Field,M.H., Morison,J.W., *et al.* (2019) Abnormally Methylated FMR1 in Absence of a
Detectable Full Mutation in a U.S.A Patient Cohort Referred for Fragile X Testing. *Sci Rep*, **9**,
15315.

194. Prawer,Y., Hunter,M., Cronin,S., Ling,L., Aliaga Vera,S., Fahey,M., Gelfand,N., Oertel,R.,
Bartlett,E., Francis,D., *et al.* (2018) Prenatal Diagnosis of Fragile X Syndrome in a Twin
Pregnancy Complicated by a Complete Retraction. *Genes (Basel)*, **9**, 287.

195. Pandelache,A., Francis,D., Oertel,R., Dickson,R., Sachdev,R., Ling,L., Gamage,D. and Godler,D.E.
(2021) Detection of Cryptic Fragile X Full Mutation Alleles by Southern Blot in a Female and
Her Foetal DNA via Chorionic Villus Sampling, Complicated by Mosaicism for
45,X0/46,XX/47,XXX. *Genes (Basel)*, **12**, 798.

196. Mätlik,K., Baffuto,M., Kus,L., Deshmukh,A.L., Davis,D.A., Paul,M.R., Carroll,T.S., Caron,M.-C.,
Masson,J.-Y., Pearson,C.E., *et al.* (2023) Cell Type Specific CAG Repeat Expansions and
Toxicity of Mutant Huntingtin in Human Striatum and Cerebellum. *bioRxiv*,
10.1101/2023.04.24.538082.

197. Pressl,C., Mätlik,K., Kus,L., Darnell,P., Luo,J.-D., Paul,M.R., Weiss,A.R., Liguore,W., Carroll,T.S.,
Davis,D.A., *et al.* (2023) Selective Vulnerability of Layer 5a Corticostriatal Neurons in
Huntington's Disease. *bioRxiv*, 10.1101/2023.04.24.538096.

198. Albert,S.M. (2010) Neurodegenerative disease and cancer: a critical role for melanoma?
Neuroepidemiology, **35**, 305–306.

199. Baade,P.D., Herrero Hernández,E., Freedman,D.M., Smithers,B.M. and Fritschi,L. (2010) No role
for melanoma treatment in the association between melanoma and amyotrophic lateral
sclerosis or Parkinson's disease. *Neuroepidemiology*, **35**, 303–304.

200. Baade,P.D., Fritschi,L. and Freedman,D.M. (2007) Mortality due to amyotrophic lateral sclerosis
and Parkinson's disease among melanoma patients. *Neuroepidemiology*, **28**, 16–20.

201. Freedman,D.M., Curtis,R.E., Daugherty,S.E., Goedert,J.J., Kuncl,R.W. and Tucker,M.A. (2013) The
association between cancer and amyotrophic lateral sclerosis. *Cancer Causes Control*, **24**,
55–60.

202. Freedman,D.M., Travis,L.B., Gridley,G. and Kuncl,R.W. (2005) Amyotrophic lateral sclerosis
mortality in 1.9 million US cancer survivors. *Neuroepidemiology*, **25**, 176–180.

203. Akimoto,C., Volk,A.E., van Blitterswijk,M., Van den Broeck,M., Leblond,C.S., Lumbroso,S.,
Camu,W., Neitzel,B., Onodera,O., van Rheenen,W., *et al.* (2014) A blinded international
study on the reliability of genetic testing for GGGGCC-repeat expansions in C9orf72 reveals
marked differences in results among 14 laboratories. *J Med Genet*, **51**, 419–424.

204. Van Mossevelde,S., van der Zee,J., Cruts,M. and Van Broeckhoven,C. (2017) Relationship
between C9orf72 repeat size and clinical phenotype. *Curr Opin Genet Dev*, **44**, 117–124.

205. DeJesus-Hernandez,M., Mackenzie,I.R., Boeve,B.F., Boxer,A.L., Baker,M., Rutherford,N.J.,
Nicholson,A.M., Finch,N.A., Flynn,H., Adamson,J., *et al.* (2011) Expanded GGGGCC
hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD
and ALS. *Neuron*, **72**, 245–256.

206. Takahashi,E., Hori,T. and Murata,M. (1988) Population cytogenetics of rare fragile sites in
Japan. *Hum Genet*, **78**, 121–126.

207. Migliore,L., Coppedè,F., Fenech,M. and Thomas,P. (2011) Association of micronucleus
frequency with neurodegenerative diseases. *Mutagenesis*, **26**, 85–92.

208. Fenech,M., Kirsch-Volders,M., Natarajan,A.T., Surralles,J., Crott,J.W., Parry,J., Norppa,H.,
Eastmond,D.A., Tucker,J.D. and Thomas,P. (2011) Molecular mechanisms of micronucleus,
nucleoplasmic bridge and nuclear bud formation in mammalian and human cells.
Mutagenesis, **26**, 125–132.

209. Dingwall,C., Lomonossoff,G.P. and Laskey,R.A. (1981) High sequence specificity of micrococcal
nuclease. *Nucleic Acids Res*, **9**, 2659–2673.

210. Hörz,W. and Altenburger,W. (1981) Sequence specific cleavage of DNA by micrococcal nuclease.
Nucleic Acids Res, **9**, 2643–2658.

211. Bastepe,M. and Xin,W. (2015) Huntington Disease: Molecular Diagnostics Approach. *Curr Protoc
Hum Genet*, **87**, 9.26.1-9.26.23.

212. Nutter,C.A., Bubenik,J.L., Oliveira,R., Ivankovic,F., Sznajder,Ł.J., Kidd,B.M., Pinto,B.S., Otero,B.A.,
Carter,H.A., Vitriol,E.A., *et al.* (2019) Cell-type-specific dysregulation of RNA alternative
splicing in short tandem repeat mouse knockin models of myotonic dystrophy. *Genes Dev*,
33, 1635–1640.

213. Love,M.I., Huber,W. and Anders,S. (2014) Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol*, **15**, 550.

214. Patro,R., Duggal,G., Love,M.I., Irizarry,R.A. and Kingsford,C. (2017) Salmon provides fast and
bias-aware quantification of transcript expression. *Nat Methods*, **14**, 417–419.

215. Angelosanto,F.A. (1995) Tissues other than bone marrow that can be used for cytogenetic
analyses. *Environ Mol Mutagen*, **25**, 338–343.

216. Bayani,J. and Squire,J.A. (2004) Preparation of cytogenetic specimens from tissue samples. *Curr
Protoc Cell Biol*, **Chapter 22**, Unit 22.2.

217. Krawczun,M.S., Lele,K.P., Jenkins,E.C. and Brown,W.T. (1986) Fragile X expression increased by
low cell-culture density. *Am J Med Genet*, **23**, 467–473.

218. Fonatsch,C. and Schwinger,E. (1983) Frequency of fragile X chromosomes, fra(X), in
lymphocytes in relation to blood storage time and culture techniques. *Hum Genet*, **64**, 39–
41.

219. Brookwell,R., Daniel,A., Turner,G. and Fishburn,J. (1982) The fragile X(q27) form of X-linked
mental retardation: FUdR as an inducing agent for fra(X)(q27) expression in lymphocytes,
fibroblasts, and amniocytes. *Am J Med Genet*, **13**, 139–148.

220. Daniel,A., Ekblom,L. and Phillips,S. (1984) Constitutive fragile sites 1p31, 3p14, 6q26, and 16q23
and their use as controls for false-negative results with the fragile(X). *Am J Med Genet*, **18**,
483–491.

1673

ACKNOWLEDGEMENTS

This study is dedicated to the memory of Dr. Stephen T. Warren (1953-2021), a pioneer and advocate of Fragile X syndrome, a mentor, and friend. We thank Robert H. Brown for providing a YAC-*C9orf72* clone, John Volpe, Shamima Keka Islam for counting micronuclei, Lazar Joksimovic, and Raymond Wong for his expertise in the preparation of metaphase spreads. We thank Olivia Beck and Lieve Wiericx for sharing their expertise in cytogenetic fragile site recognition and Edwin Reyniers and Stephen W. Scherer, Jeff MacDonald, Mehdi Zarrei, Mehdi Layeghifard, and Bhooma Thiruvahindrapuram for help with the der-Chr9 data, and Michael Fenech for support regarding micronuclei, NBuds, and NBlebs.

DATA AVAILABILITY

Datasets generated during the current study supporting its findings have been deposited or made available as indicated:

- RNASeq gene expression data - available on Gene Expression Ombinus (GEO), accession # [to be inserted]
- Whole genome sequencing data in the form of CRAM files – deposited to NCBI's Seqeunce Read Archive (SRA), accession # [to be inserted]

INCLUSION AND ETHICS STATEMENT

All collaborators of this study who have fulfilled the criteria for authorship required by Nature journals are included as authors. Each author's contribution is detailed in the Author Contributions list. Expectations for collaborators were agreed upon ahead of the study's initiation, and authorship placement was agreed upon by each collaborator. This study was conducted in collaboration with researchers and physicians from distinct cultural, ethnic, and racial backgrounds. The findings of this study apply to members of the population globally, as well as locally in the regions where patient recruitment occurred. The research does not result

1698 in stigmatization, incrimination, discrimination, or personal risk to participants who provided

1699 samples. Local and regional research publications relevant to our study was considered and

1700 included in our bibliography.

1701 Hospital for Sick Children Research Ethics Board oversight and approval for the herein study

1702 was obtained. REB for human tissue, cell and animal studies: #1000056560.

AUTHOR CONTRIBUTIONS

Conceptualization: CEP and RFK

Fig. 1: MM, MHMS, CEP, PAD, GAR, EK, MZ

Fig. 2: MM, IvdW, SS, RFK, CEP

Fig. 3: MHMS, TKP, CEP (sample collection by LZ, JK, AA, PMcG, PMcK, JR)

Fig. 4: MM*, MHMS, NS, SL, KY, MM†, MK, RKCY, PMcK, JR, LS, MSS, CEP

Fig. 5: NS, BM (training by MM*, MHMS, TKP)

Fig. ED1: ML, MDW, CEP

Fig. ED2: MM*, MZ, ER, MHMS

Fig. ED3: MM*, IvdW, SS, RFK

Fig. ED4: MHMS

Fig. ED5: MM*

Fig. ED6: MM*, MM†, SL, NS, KY, MHMS, CEP

Fig. ED7: TA, EAP, CHF

Fig. ED8: RKCY, RFK, MM*, LS

Fig. ED9: NS

Fig. ED10: BM (training by MM*)

Table ED1: MHMS

Table ED2: MM* and CEP

Table ED3: LZ, JK, AA, PMcG, PMcK, JR, CEP

Table ED4: LS, MK, PMcK, RKCY, JR, and CEP

Table ED5: MK, RKCY

BAC DNA for provided by: CJ, PJdeJ

ALS patient samples provided by: PAD, GAR

Autopsy tissues prepared by: LZ, PMK, JR

Intellectual input and experimental design: CEP, RFK, EC, RW, MDW, MSS, ER

Supervision: CEP and RFK

Writing – original draft: MM*, CEP

Writing – review & editing: CEP, MHMS, NS, MSS, MM†

All authors provided feedback on the manuscript.

MM* = Mila Mirceta

MM† = Mohiuddin Mohiuddin

FUNDING

This work was supported by the ALS Canada-Brain Canada 2020 Discovery Grant

[HSC0005599 to C.E.P., E.R., G.A.R.]; ALS Canada Doctoral Research Award 2016 [to

M.H.M.S.]; the Weston Family Foundation [TR202268 to C.E.P.]; the Canadian Institutes of

Health Research [NDD-144223 to C.E.P.]; the W. Garfield Weston Foundation [CNV0001757

to C.E.P.]; the National Institutes of Health [GM122880 to C.H.F.]; and the National Natural

Science Foundation of China [82071430 to M.Z., E.R.]; James Hunter and Family ALS

Initiative [J.R. and E.R.]; Canadian Consortium on Neurodegeneration in Aging [to G.A.R. and

E.R.].

Funding for open access charge: The Canadian Institutes of Health Research

COMPETING INTERESTS

The authors declare no competing interests.

SUPPLEMENTARY INFORMATION

Supplementary information is available for this paper by request.

MATERIALS AVAILABILITY

Correspondence and requests for materials should be addressed to Dr. Christopher E. Pearson

(mail to: cepearson.sickkids@gmail.com). Mouse models presented in this study are

commercially available from Jackson Laboratories (FVB/NJ-Tg(C9orf72)500Lpwr/J).