Mechanisms of *P. aeruginosa* resistance to Type VI Secretion System attacks

Alejandro Tejada-Arranz[1], Annika Plack[1], Minia Antelo-Varela[1,2], Andreas Kaczmarczyk[1], Alexander

Klotz[1,3], Urs Jenal[1] and Marek Basler[1*]

[1]Biozentrum, University of Basel, Switzerland.

[2]Current affiliation: Proteomics Unit, Gulbenkian Institute for Molecular Medicine, Lisbon, Portugal.

[3]Current affiliation: Department of Biosystems Science and Engineering, ETH Zurich, Basel,

Switzerland.

*Correspondence should be addressed to: marek.basler@unibas.ch

Abstract

The Type VI Secretion Systems (T6SSs) is a molecular nanomachine that injects toxic effector proteins

into the environment or neighbouring cells, and thus plays an important role in interbacterial competition

and host antagonism during infection. *Pseudomonas aeruginosa* is an opportunistic bacterial pathogen

that encodes three different T6SS. The H1-T6SS delivers toxins into aggressive bacteria in response to

attacks mediated by the their own T6SS. This suggests that *P. aeruginosa* has the ability to survive T6SS

assaults. However, the resistance mechanisms are poorly characterized. In this work, we performed a

CRISPRi screen to identify pathways involved in resistance to T6SS effectors of *Acinetobacter baylyi*

and *Vibrio cholerae*. We show that members of the GacA/GacS TCS regulon, such as the *mag* operon,

and GacA-independent factors, such as the outer membrane protein OprF, confer resistance to T6SS

toxins. We show that outer membrane anchoring to the peptidoglycan is crucial for resistance against

T6SS attacks, as well as for resistance to different antibiotics, suggesting a link between general T6SS

resistance and antibiotic resistance.

Introduction

P. aeruginosa is a Gram-negative, ubiquitous, opportunistic pathogen that can colonize open wounds and the respiratory airways, and is particularly problematic for patients suffering from cystic fibrosis[1]. *P. aeruginosa* possesses three Type VI Secretion Systems (T6SS), molecular nanoweapons that deliver toxic effectors to neighbouring cells[2]. The first one, called H1-T6SS, plays a predominantly antibacterial role, while the H2- and H3-T6SSs are important for internalization and colonization of a eukaryotic host, although they also deliver trans-kingdom effectors that target both eukaryotic and bacterial cells[2,3].

Generally, T6SSs are composed of a membrane complex and a baseplate that trigger the assembly of an Hcp tube that is loaded with a tip complex, composed of a VgrG trimer and PAAR proteins, as well as a set of effectors that can be fused to Hcp, VgrG or PAAR or can be loaded separately[4]. The Hcp tube is surrounded by a sheath that, upon contraction, propels the Hcp tube as well as the tip complex and all loaded effectors to the extracellular space or into a neighbouring cell. Bacteria are typically protected from their own effectors by a set of dedicated immunity proteins that are encoded adjacently to the effectors[5]. These immunity proteins confer highly specific immunity, with few examples of cross-protection to related effectors with similar mechanisms of action[6].

Interestingly, the H1-T6SS of *P. aeruginosa* was reported to respond to incoming attacks from neighbouring bacteria, regardless of whether they are sister cells or members of a different species[7,8]. Specifically, the H1-T6SS comprises a sensor module, TagQRST, that senses membrane damage and triggers the assembly of a retaliatory H1-T6SS through post-translational modifications[7]. Furthermore, it was proposed that this retaliation strategy is a poor evolutionary strategy, as it prevents *P. aeruginosa* from attacking first and forces it to resist incoming T6SS attacks[9]. Here, we hypothesize that *P. aeruginosa* must be equipped to survive incoming T6SS attacks, at least until it is able to strike back, suggesting the existence of specific mechanisms that protect against attacks with foreign T6SS effectors even in the absence of an active H1-T6SS.

Acinetobacter baylyi ADP1 and *Vibrio cholerae* 2740-80 are two laboratory models that are frequently used to study T6SS dynamics. They are both Gram-negative environmental bacteria that each encode one antibacterial T6SS cluster with multiple effectors that is highly active *in vitro*. *A. baylyi* ADP1 WT

contains four described effectors[10]: Tpe1 (a predicted metallopeptidase without identified antimicrobial

activity), Tae1 (a PG-targeting effector), Tse2 (of unknown mechanism of action) and Tle1 (a lipase).

V. cholerae 2740-80 WT also has four effectors[11]: TseH and VgrG3 (two PG-targeting effectors), TseL

(a lipase) and VasX (a pore-forming toxin). Thus, they have different types of effectors, and the

resistance mechanisms of *P. aeruginosa* against these two organisms might differ.

In this work, we competed H1-T6SS-deficient *P. aeruginosa* PAO1 carrying a CRISPRi plasmid library

targeting all *P. aeruginosa* genes with T6SS-positive or –negative *Acinetobacter baylyi* ADP1 or *Vibrio*

cholerae 2740-80. Deep sequencing of the plasmid library revealed genes required to resist incoming

T6SS attacks. Interestingly, most of these genes belong to the regulon of the GacS/GacA two component

system, including the previously identified *arc1* and *arc3* clusters and *pa3267*, *aas*[12], as well as the *mag*

operon. Furthermore, we found that the PG-anchoring function of the outer membrane protein OprF also

contributes to T6SS resistance. Our data suggest that these pathways are general protection mechanisms

against multiple types of effectors and thus differ from specific immunity proteins.

Results

P. aeruginosa resists T6SS attacks in the absence of its H1-T6SS

The H1-T6SS of *P. aeruginosa* can promptly and efficiently retaliate to incoming T6SS attacks, eliminating the competing bacterium before it can efficiently kill *P. aeruginosa* [7,9,13]. Therefore, we constructed a mutant lacking *tssM1*, a structural membrane component of the H1-T6SS, that is unable to assemble its H1-T6SS and retaliate to incoming attacks. We observed that the H1-T6SS is indeed important to efficiently resist attacks from *A. baylyi* ADP1 (Figure 1A and Supplementary Figure 1A), but not from *V. cholerae* 2740-80 (Figure 1B and Supplementary Figure 1B). Furthermore, although this strain is more susceptible than its parental to attacks from the T6SS of *A. baylyi* ADP1, it is still far more resistant than *E. coli* (Figure 1A and 1C and Supplementary Figures 1A and 1C), as evidenced by the almost complete elimination of *E. coli* when competed in a 10:1 attacker:prey ratio and the strong killing observed in a 1:1 ratio. This suggests that *P. aeruginosa* possesses protective mechanisms against T6SS attacks.

Identification of T6SS-protective pathways

In order to identify new T6SS-protective pathways, we introduced the dCas9 protein from *Streptococcus pasteurianus* into *P. aeruginosa* Δ*tssM1* strain and performed a CRISPRi screen for genes important for T6SS protection using a previously described sgRNA library published elsewhere (ref). Specifically, we competed this T6SS-deficient *P. aeruginosa* strain carrying the sgRNA plasmid library with T6SS-active or inactive mutants (lacking *tssM*) of *A. baylyi* ADP1 or *V. cholerae* 2740-80. We performed two rounds of 3h long incubation of predator and prey at 1:1 and 10:1 ratios, respectively (Figure 2A). The sgRNA loci of the recovered *P. aeruginosa* cells were PCR-amplified (before competition and after the first and second rounds of competition) and the amplicons were sequenced by Illumina sequencing. Samples from competition with T6SS-active or inactive strains were compared in order to identify the pathways that, when knocked down, lead to higher sensitivity to T6SS attacks.

After one and two rounds of competition, plasmids encoding sgRNAs targeting a number of *P. aeruginosa* genes were depleted (Figure 2BC and Supplementary Figure 2A and 2B). The same hits were identified in both cases, although fold-changes were stronger after two rounds of competition. This

included known T6SS resistance-related genes such as *gacA*, *gacS*, *aas*, *arc1A* or *arc3A*[12] as well as

novel genes such as the *mag* operon or *oprF*. Interestingly, most of these genes are likely under the

control of the GacA/GacS two component system (TCS), as the encoded proteins, together with

components of the H1- and H2-T6SS, were less abundant in the proteome of a Δ*gacA* strain (Figure 2D,

Supplementary Figure 2C, and Supplementary Table 1). Overall, this indicates that GacA/GacS

regulates pathways involved in resistance to T6SS attacks.

GacA is a key regulator involved in T6SS resistance

As many of the hits of our screening are regulated by GacA, we next assessed whether a Δ*gacA* strain

was able to resist T6SS attacks from *A. baylyi* and *V. cholerae*. Indeed, the Δ*gacA* strain was 100-fold

more susceptible to T6SS attacks from *A. baylyi* and *V. cholerae* than a strain lacking the H1-T6SS

(Figure 2E and Supplementary Figure 2D), indicating that GacA-regulated genes other than the H1-

T6SS are required for T6SS resistance.

To get more insights in the mechanisms of resistance, we competed *P. aeruginosa* Δ*gacA* strain with *A.*

baylyi strains lacking individual effectors (Figure 2F and Supplementary Figure 2E). We previously

showed that these *A. baylyi* strains are able to assemble its T6SS as efficiently as the parental strain[10].

Interestingly, the Δ*gacA* strain is killed similarly when single effectors are absent, although Tse2 might

play the most predominant role, as an *A. baylyi* strain lacking *tse2* kills *P. aeruginosa* less efficiently

(Figure 2F).

Furthermore, we also competed *P. aeruginosa* Δ*gacA* with *V. cholerae* strains encoding individually

inactivated effectors through point mutations that were previously reported[8,14], and we found that the

inactivation of each effector leads to similar levels of killing of the Δ*gacA* strain (Figure 2G and

Supplementary Figure 2F). Taken together, our results suggest that the GacA regulon contains genes

that contribute to the resistance to different effector classes, in addition to lipase effectors as previously

reported[12].

The *mag* operon is important for resistance against PG-targeting effectors

Next, we assessed how individual genes from the GacA regulon contribute to T6SS resistance, and in particular the *mag* operon that was a clear hit in our CRISPRi screen (Figure 2B). In *P. aeruginosa*, the *mag* operon is composed of 6 genes, *magABCDEF*. MagD displays weak homology with type II α2-macroglobulin[15], homologs of eukaryotic α-macroglobulins, whereas the rest of the genes of the operon are important for the folding and stability of MagD[16]. Eukaryotic α2-macroglobulins are innate immunity proteins that protect against proteases by a trapping mechanism[17–19]. However, bacterial type II α2-macroglobulins lack key residues that conform the active site of eukaryotic α2-macroglobulins[16], and their function is thus unclear.

We constructed a strain lacking both *tssM1* and *magD*, and competed it against different mutants of *V. cholerae*. We performed quantitative killing assays and found that MagD is indeed important for resistance to T6SS attacks, and that it protects mainly against the PG-targeting effector VgrG3 (Figure 3A and Supplementary Figure 3). To confirm this result, we also performed a CPRG-based lysis assay[10], a colorimetric assay where lysis is measured based on the conversion rate of the membrane impermeable CPRG dye by LacZ enzyme that is released upon lysis of *P. aeruginosa*. This assay also indicated a VgrG3-dependent lysis of a Δ*magD* mutant (Figure 3B). Additionally, although only *magE* was identified in the *A. baylyi* competition experiment, *magD* also confers a mild protection against the PG-targeting effector Tae1 (Figure 3C). Thus, MagD encoded by the *mag* operon specifically protects against PG-targeting effectors from both *V. cholerae* and *A. baylyi*.

Arc1A, Arc3A and Aas contribute to resistance against lipase effectors

We also assessed the role of *arc1A*, *arc3A* and *pa3267* (*aas*), in T6SS resistance. Arc1A and 3A were previously shown to be important for resistance against the T6SS of *B. thailandensis*. Specifically, Arc3A protects against lipase effectors, and the effector class that Arc1A protects against was not clearly elucidated[12]. PA3267 is homologous to the lysophospholipid transporter LplT and the Aas acyltransferase from *E. coli*, where it was shown to be important for defense against phospholipase attacks[20]. We found that a *P. aeruginosa* mutant lacking either *arc1A*, *arc3A* or *aas* is indeed more sensitive to T6SS attacks from an *A. baylyi* carrying only the lipase Tle1 effector (Figure 4) in a CPRG

145 lysis assay. This indicates that *aas* together with the previously identified *arc3A* and *arc1A* play a role

146 in resistance to lipases.

147 Anchoring of the outer membrane to the PG is required for T6SS resistance and is genetically

148 linked to GacA/S

OprF is the main outer membrane porin of *P. aeruginosa*, and it has been reported to perform many

different roles including channelling small solutes, maintenance of outer membrane integrity, binding

and adhesion, biofilm formation, or outer membrane vesicle biogenesis[21]. OprF is folded in two

conformers, a minor population (around 5%) that is folded into a one-domain structure with a channel

that acts as porin; and a major population (around 95%) that is folded into two domains including a PG-

anchoring domain. It was suggested that this conformation is responsible for anchoring of the OM to the

PG[22,23], and that a Δ*oprF* strain displays defects in the OM[24,25]. To understand how sgRNA mediated

inhibition of the *oprF* expression resulted in lower resistance to T6SS attacks, we constructed a *P.*

aeruginosa Δ*oprF* strain and competed it against *V. cholerae* and *A. baylyi* attackers. We show that

when the attacker strains carry full effector sets, or when single effectors are inactivated, the Δ*oprF*

strain is more sensitive than the parental strain (Fig. 5A, B and Supplementary Figure 4A, B).

Importantly, Δ*oprF* strain displays a growth defect in LB, which is exacerbated in LB without salt

(Supplementary Fig. 4C), suggesting that the strain is sensitive to osmotic shock as previously shown[24].

Furthermore, complementation *in trans* with a wt copy of *oprF* in a pPSV35 plasmid fully restores both

growth and resistance to T6SS attacks (Fig. 5C and Supplementary Fig. 4D and E). The R296E *oprF*

variant, that was previously shown to be unable to bind PG in *E. coli*[26,27], fails to complement either of

these phenotypes (Fig. 5C and Supplementary Fig. 4D and E). Specifically, the OprF-R296E variant

fails to complement the survival defect when faced with *A. baylyi*, and only partially resists attacks from

V. cholerae (Fig. 5C and Supplementary Fig. 4D and E).

Interestingly, we observed different colony sizes when the *P. aeruginosa* Δ*tssM1* Δ*oprF* strain was

grown in LB without salt, suggesting the appearance of revertants (Figure 6A). We isolated four of these

clones and analysed their growth, cell morphology and ability to compete with *V. cholerae*. These

revertants displayed wild type-like growth on LB without salt (Figure 6B), as well as wild-type cellular

morphology (Figure 6C), as shown by the absence of cell ghosts that are frequent in the Δ*tssM1* Δ*oprF*

strain. However, the revertants were still susceptible to T6SS attacks from *V. cholerae* at a 1:1 ratio

(Figure 6D and Supplementary Fig. 4F).

We performed whole genome sequencing of these revertants and found that all of them carry the same

inactivating 431bp deletion within the *gacS* gene, suggesting that the GacA/S is inactive in these strains.

We validated this by deletion of *gacA* on a Δ*oprF* strain, which restores the growth defect of this strain

(Figure 6E). Thus, GacA/S is involved in causing the growth and morphology defect of the Δ*oprF* strain,

which explains why these revertants are still sensitive to T6SS attacks, as they lack a functional GacA/S

that is important for resistance to such attacks (Figure 2). Our results also show that there is a genetic

link between *oprF* and the GacA/S.

Immunity strategies are species-specific

While the previous study from Ting *et al.*[12] identified three operons, *arc1*, *arc2* and *arc3*, involved in

resistance to T6SS from *Burkholderia thailandensis*, only two of them, *arc1* and *arc3* appeared in our

CRISPRi screening. Likewise, the two novel resistance factors that we identified, the *mag* operon and

oprF, were not identified in the previous transposon screen. In order to verify whether these mechanisms

are important to resistance to other pathogens, we also tested the resistance of our strains to *B.*

thailandensis attacks (Fig. 7 and Supplementary Figure 5). Although *magD* is dispensable for resistance

to the T6SS of *B. thailandensis*, both *gacA* and *oprF* play a role, albeit to a lesser extent than in the case

of *A. baylyi* or *V. cholerae*. Thus, some T6SS-protective mechanisms, such as those in the GacA regulon,

are specific to certain species or certain effector types, others, such as the OprF-mediated mechanisms,

appear to be generic resistance mechanisms.

T6SS immunity strategies can confer resistance to antibiotics

In order to determine whether T6SS resistance mechanisms could also be also employed as antibiotic

resistance mechanisms, we explored the susceptibility of the T6SS-sensitive strains to 30 antibiotics of

different classes. We found that the Δ*oprF* strain is more sensitive to antibiotics such as tetracycline

(Figure 8A) and piperacillin (Figure 8B), while it is more resistant to others like colistin (Figure 8C),

among others (Supplementary Figure 6). Furthermore, the Δ*magD* strain is more resistant to the cell

membrane-targeting antibiotic chlorhexidine (Figure 8D and Supplementary Figure 7). This suggests

that general T6SS immunity mechanisms such as the ones mediated by OprF or MagD also contribute

to the antibiotic susceptibility profile of *P. aeruginosa*.

Discussion

P. aeruginosa is able to promptly sense and retaliate to T6SS attacks from other species[7]. However, the mechanisms that allow *P. aeruginosa* to withstand the initial attack are not well understood. Only one recent report[12] determined three operon involved in resistance to T6SS attacks from *B. thailandensis*, called *arc1*, *arc2* and *arc3*, and identified that *arc3* is particularly important to resist the delivery of lipase effectors, likely through the detoxification of lysophospholipids that are generated as a result of the activity of these lipases. However, the effector types that *arc1* and *arc2* confer resistance against, and whether there are other mechanisms involved in the resistance to other effector types, is not known. Here, we performed a CRISPRi screen to identify novel factors involved in T6SS attack resistance in *P. aeruginosa*, using the laboratory models *A. baylyi* and *V. cholerae* as aggressor species. These two species contain different effector sets[10,11]: both of them have at least one lipase (Tle1 in *A. baylyi* and TseL in *V. cholerae*) and at least one PG-targeting effector (Tae1 in *A. baylyi* and VgrG3 and TseH in *V. cholerae*); *A. baylyi* encodes two other effectors of unknown activity (Tpe1 and Tse2); and *V. cholerae* encodes one membrane-targeting pore-forming effector VasX.

In our screen, we identified known factors such as the *arc1* and *arc3* clusters, as well as *gacS* and *gacA* of the GacA/S TCS that regulates the expression of said clusters[12] in addition to the expression of the H1- and H2-T6SS[30,31] (Figure 2B-C). We also identified novel factors that contribute to resistance, the *mag* operon, that is also regulated by GacA, underscoring the importance of this transcriptional regulator for the survival of *P. aeruginosa* in polymicrobial communities. Furthermore, we found other factors that are now known to be regulated by GacA (Figure 2E), such as the outer membrane protein OprF.

Through quantitative killing assays and kinetic CPRG-based lysis assays, we could validate that these factors are indeed involved in resistance to T6SS attacks (Figure 2-5). In agreement with Ting *et al.*[12], we also observed that *arc3* is important specifically to resist to lipases (Figure 4). Here, we also observe that *arc1* as well as *aas* are required for full resistance against the lipase effector of *A. baylyi*. In their study, Ting *et al.* did not determine the effector type that *arc1* is active against and, although they hypothesized that *aas* is active against lipases[12], they did not find a clear phenotype against *B.*

thailandensis. This suggests that the specific catalytic mechanism of the lipases in question might be

relevant for the mechanism of action of the corresponding resistance mechanisms.

Our results also show that the GacA regulon is important for the resistance against effector classes other

than lipases. We found that the *mag* operon, that is regulated by GacA, is particularly important to resist

against certain PG-targeting effectors such as VgrG3 from *V. cholerae* and Tae1 from *A. baylyi.*

Interestingly, the central gene of this operon, *magD*, displays homology with human α2-

macroglobulins[15], innate immunity factors that defend against proteases through a trapping

mechanism[17–19]. However, the catalytic residues of this protein family are not conserved in bacterial type

II α2-macroglobulins that *magD* belongs to[16], raising questions about the true function of this protein

family in bacteria. Here, we found that MagD contributes to resistance to T6SS PG-targeting effectors,

although the exact molecular mechanism remains to be elucidated. Interestingly, MagD was previously

found to co-immunoprecipitate with TssM1[15], a structural transmembrane component of the H1-T6SS

of *P. aeruginosa*, suggesting that these two components might co-localize where a foreign T6SS attack

was detected and contribute to the defense against the injected effectors (in the case of MagD) and the

assembly of a counterattack H1-T6SS machine (in the case of TssM1).

P. aeruginosa also possesses other resistance mechanisms that are not known to be regulated by GacA

(Figure 2D), such as the outer membrane protein (OMP) OprF. OprF is the most abundant OMP in *P.*

aeruginosa, and carries out multiple functions including an outer membrane porin as well as anchoring

the outer membrane to the PG[21]. Here, we found that the anchoring of the OM to the PG contributes to

resistance to multiple T6SS effector classes (Figure 5A and B and Supplementary Figures 4A and B).

We further confirmed this phenotype by complementation with a wild-type or mutant variant of OprF

that no longer binds PG (Figure 5C), and found that this variant is unable to promote survival against *A.*

baylyi and only partially improves survival against T6SS-active *V. cholerae*, underscoring the

importance of OM integrity for survival to T6SS attacks.

Our study also provides the first evidence for the genetic relationship between *oprF* and the *gacA/S*

TCS. A strain lacking *oprF* displays a growth defect in LB that is more pronounced in LB without salt

(Supplementary Figure 4C). In LB without salt, this strain spontaneously reverts to wild-type-like

growth at high frequency (Figure 6A and B) and, although these colonies also display WT-like

morphology (Figure 6C), they are still unable to survive T6SS attacks from *V. cholerae* (Figure 6D).

Our whole genome sequencing analysis showed that these revertants carry a loss-of-function deletion

of *gacS*, which explains why these strains are susceptible to T6SS attacks. Indeed, the GacA/S TCS is

responsible for the growth defect of the Δ*oprF* strain, since deletion of *gacA* restores this defect (Figure

6E). Further research is needed in order to determine why the expression of the GacA regulon leads to

a growth defect in the Δ*oprF* genetic background.

Next, we also assessed whether our newly identified resistance mechanisms also play a role against *B.*

thailandensis E264. As expected from a transposon screen performed by Ting et al.[12], *magD* does not

play a role in defending from attacks from *B. thailandensis*. However, both GacA and OprF are

important for this (Figure 7 and Supplementary Figure 5). Interestingly, the *arc2* operon identified by

Ting et al.[12] did not appear in our screen. Taken together, our results show that *P. aeruginosa* possesses

an array of T6SS resistance mechanisms, frequently under the control of the GacS/A TCS, that help it

survive attacks from a variety of aggressor species that it can encounter in the environment. However,

these resistance mechanisms might only protect against certain species or against effectors with specific

modes of action. We hypothesize that *P. aeruginosa* encodes further T6SS resistance mechanisms, for

instance against the pore-forming effector VasX from *V. cholerae*, since a Δ*gacA* strain is more

susceptible to attacks with VasX than a WT strain (Figure 2G), indicating that the GacA regulon contains

other resistance mechanisms that are yet to be identified.

Interestingly, other general stress response factors such as those previously found to protect *E. coli* and

P. aeruginosa[32,33] were not highlighted in our screen. In the case of *P. aeruginosa*, those factors were

identified upon overexpression of TseL in the host periplasm, a condition that is likely more deleterious

for *P. aeruginosa* than a T6SS attack where only a few molecules of TseL may be injected at the same

time[32]. This would suggest that members of the GacA regulon mount a first layer of defence and that,

when they are overwhelmed, general stress response mechanisms of the cell come into play. Performing

a similar screen in a Δ*gacA* genetic background where dedicated protection mechanisms are not

expressed might highlight such general stress response mechanisms.

We also wondered whether general T6SS resistance mechanisms could be regarded as general

mechanisms that could also be involved in resistance to other antimicrobials such as antibiotics. We

screened a panel of 30 antibiotics against the WT, Δ*oprF* and Δ*magD* strains and found that the Δ*oprF*

is indeed more sensitive to antibiotics of different classes such as the beta-lactam piperacillin and the

ribosome-targeting tetracycline, among others (Figure 8A-C and Supplementary Figure 6). These

phenotypes cannot be explained by the porin function of *oprF*[21], since the strain should be more resistant

to these antibiotics if *oprF* was required for drug uptake. Furthermore, the Δ*oprF* is more resistant to

colistin. It is known that colistin kills Gram-negative bacteria by targeting LPS and disrupting the

OM[34,35]. Most mechanisms of resistance to colistin have been linked to LPS modification[34]. More

research is needed to clarify why an *oprF* mutant, that displays defects in OM stability, would be more

resistant to colistin. Interestingly, the Δ*magD* strain was more resistant to the cell membrane-targeting

antibiotic chlorhexidine[36] (Figure 8D and Supplementary Figure 7). More research is needed to

understand why this strain can deal better with the stress imposed by this antiseptic molecule. However,

our findings serve as proof-of-concept that the study of the mechanisms of resistance to the T6SS can

lead to valuable insights about antimicrobial resistance in bacteria.

In conclusion, in this work we identified multiple novel resistance mechanisms in *P. aeruginosa* against

T6SS attacks from *A. baylyi* and *V. cholerae*, which are also involved in resistance to classical

antimicrobials. In the future, further research will be needed to determine whether these resistance

mechanisms play a role in the survival of *P. aeruginosa* strains in naturally-occurring polymicrobial

communities, and particularly in the context of clinical infections.

Materials and methods

Bacterial strains and growth conditions

Bacterial strains used in this study can be found in Supplementary Table S2. Bacteria were grown in LB at 37°C unless otherwise stated. *E. coli* DH5α λpir was used as cloning strain and *E. coli* SM10 λpir was used for conjugation of *P. aeruginosa* and *V. cholerae*. Antibiotics used were streptomycin (100 μg/mL for *V. cholerae* and *A. baylyi* on solid media; and 50 μg/mL for *A. baylyi* in liquid media), gentamicin (20 μg/mL for *P. aeruginosa* and *E. coli*), kanamycin (50 μg/mL for *A. baylyi*), carbenicillin (150 μg/mL for *E. coli* and *V. cholerae* and 100 μg/mL for *P. aeruginosa*), irgasan (20 μg/mL for *P. aeruginosa*) and trimethoprim (200 μg/mL for *B. thailandensis* and 50 μg/mL for *E. coli*).

Mutant construction

All plasmids used in this study are listed in Supplementary Table S3, and primers in Supplementary Table S4. *P. aeruginosa* mutants were constructed as previously described using the pEXG2 suicide plasmid[37]. Briefly, 800bp homology arms were amplified by PCR using primers (a list of all primers can be found in Supplementary Table S3) with 20bp stretches that were homologous to the *Hind*III and *Xba*I restriction sites of the pEXG2 plasmid. PCR products were inserted into linearized pEGX2 plasmid by Gibson assembly[38] (NEB, E2611L). The resulting vectors were transformed into chemically-competent *E. coli* DH5α λpir, selected on LB plates with 20 μg/mL gentamicin and verified by colony PCR and Sanger sequencing. The purified plasmids were then transformed into chemically-competent *E. coli* SM10 λpir and the resulting strains were used for conjugation of *P. aeruginosa* PAO1. Selection of *P. aeruginosa* transconjugants was performed on LB with 20 μg/mL gentamicin and 20 μg/mL irgasan. The pEXG2 backbone was then curated by restreaking on LB plates without salt supplemented with 20 μg/mL irgasan and 10% sucrose, which were incubated for 36h at room temperature. Colonies were screened by colony PCR with flanking primers.

pPSV35-derived plasmids were introduced into *P. aeruginosa* by electroporation of 20-200ng of plasmid into 1 mL of overnight stationary phase cultures washed twice with sterile water using a Bio-Rad GenePulser with the following settings: 25 μF, 400 Ω, 1.8 kV. All steps were performed at room

temperature. 1 mL of LB medium was added immediately after the pulse and cells were incubated 1.5-

2h at 37°C with shaking before plating on LB plates containing 20 μg/mL gentamicin.

The chromosomal insertion of *dCas9* in the attnT7 site was performed as previously described (ref).

Briefly, an 18-22 h old culture of *P. aeruginosa ΔtssM1* in LB without salt was washed twice with sterile

water and resuspended in 100 μL sterile water. The bacterial suspension was electroporated with two

plasmids at the same time, pUC18T:TNT7:dCas9Spas and pTNS2 that contains the Tn7 integrase and

is unable to replicate in *P. aeruginosa*. The electroporation was performed as described above. The cells

were recovered for 1h in LB at 37°C and 100 μL were plated on LB agar plates with 20 μg/mL

gentamicin to select for transconjugants that integrated *dCas9* into the chromosome. The next day,

colonies were grown in LB no salt and, as above, electroporated with plasmid pFLP2 containing the

flippase that removes the backbone of the pUC18T plasmid. After recovery, 100 μL were plated on LB

agar plates with 100 μg/mL carbenicillin and incubated overnight at 37°C. The resulting colonies were

restreaked on LB no salt + 10% sucrose to counterselect pFLP2. The successful insertion of *dCas9* was

verified by PCR and the resulting strain was phenotypically validated by expression of an *ftsZ*-targeting

sgRNA that led to cell filamentation as described (ref).

V. cholerae mutants were constructed in a similar manner, using the pWM91 suicide plasmid and the

XbaI and HindIII restriction sites. *V. cholerae* transconjugants were selected on LB plates with 150

μg/mL carbenicillin and 100 μg/mL streptomycin. The pWM91 backbone was curated by restreaking

on LB plates without salt supplemented with 100 μg/mL streptomycin and 10% sucrose.

A. baylyi mutants were constructed by natural transformation as previously described[10], using a

kanamycin resistance, streptomycin counterselectable, cassette and 800bp homology arms.

B. thailandensis were conjugated with pMLBAD plasmid as previously described[39].

Quantitative killing assays

Overnight cultures of *P. aeruginosa*, *V. cholerae*, *E. coli* and *B. thailandensis* were grown in LB at 37°C

and 200 rpm without antibiotics unless *P. aeruginosa* or *B. thailandensis* contained plasmids. *A. baylyi*

overnight cultures were grown at 30°C in similar conditions without antibiotics. Day cultures were

launched in LB at the following dilutions: 1:100 for *P. aeruginosa*, 1:50 for *A. baylyi* and *B.*

thalandensis, 1:200 for *E. coli* and 1:400 for *V: cholerae*. After 2.5-3h, when all cultures reached an

OD_{600nm} of 0.7-1.4, cultures were spun down at 12,000g for 2 min at room temperature. The cultures

were concentrated to OD 10 with LB and 1:10 prey:attacker mixes were prepared. Five µL of the

resulting mixes were spotted on LB agar plates, in technical triplicates, and allowed to dry. The plates

were incubated at 37°C for 3-4h and then the spots were excised and resuspended in 500 µL liquid LB

by vortexing. Ten-fold serial dilutions were performed and 5 µL of the resulting suspensions were

spotted on selective plates. As selective agents, we used irgasan 20 µg/mL for *P. aeruginosa*,

streptomycin 100 µg/mL for *V. cholerae* and *A. baylyi*, gentamicin 20 µg/mL for *E. coli* and 200 µg/mL

trimethoprim for *B. thalandensis*. Plates were incubated overnight at 37°C and the surviving CFUs were

counted the next day and technical triplicates were averaged. The experiments were performed with at

least three biological replicates and plotted with Prism.

CRISPRi library electroporation

A *P. aeruginosa* Δ*tssM1* strain containing *dCas9* at the attnT7 site was grown overnight in LB without

salt and 4 mL were electroporated with 2 µL of the CRISPRi library (ref) as described above. After

electroporation, the bacterial suspension was resuspended in 1 mL LB and transferred to a 1 L

Erlenmeyer flask containing 100 mL LB and allowed to recover for 1-1.5h at 37°C while shaking. Next,

20 µg/mL gentimicin were added and the culture was incubated overnight at 37°C while shaking. This

protocol led to an electroporation efficiency of $3.8 \cdot 10^6$ cells/mL, which allows to achieve a good

coverage of the ca.83,000 sgRNA library.

CRISPRi screening competition

Three 100 mL day cultures in LB+20 µg/mL gentamicin+1mM IPTG were launched by inoculation of

1 mL at OD_{600nm} 1 from the library electroporation culture. In parallel, 20 mL day cultures were launched

from overnight cultures of *V. cholerae* and *A. baylyi* strains. The cultures were incubated for around 3h

until an OD_{600nm} of 0.7-1.4 was reached. Five mL of each *P. aeruginosa* culture were taken, pelleted and

frozen at -80°C until further processing (time 0 before the competition started). For each replicate, the

cultures were concentrated to OD 10 and mixed in a 10:1 *V. cholerae*:*P. aeruginosa* ratio or in a 1:1

A.baylyi:*P. aeruginosa* ratio and 90 μL of the resulting mixes were spotted on LB+1mM IPTG plates.

P. aeruginosa alone was also spotted (no co-incubation control). The spots were allowed to dry and

were then incubated at 37^{o}C for 3h. Then, the spots were excised and resuspended in 7 mL LB by

vortexing and pipetting. The resuspended bacteria was spun down for 10 min at 3,900g and resuspended

in 7 mL LB. Five mL were pelleted and frozen at -80^{o}C until further processing (samples after the first

round of competition). The remaining 2 mL were used to measure the OD$_{600nm}$ and inoculate new day

cultures in 100 mL LB + 20 μg/mL gentamicin + 1 mM IPTG with 1 mL OD 1. New day cultures of *V.*

cholerae and *A. baylyi* were launched similarly as earlier. The same process was repeated for a second

round of competition and the resulting samples were again pelleted and frozen at -80^{o}C until further

processing.

CRISPRi sequencing library preparation

Pellets were thawed and resuspended in 500 μL 50 mM NaOH. The samples were incubated for 10 min

at 50^{o}C and 1 μL was used as template for a PCR reaction using Phusion DNA Polymerase (M0530S,

NEB) as described (ref) using primer pairs oATA402+403, oATA404+405, oATA406+407 and

oATA408+409 for a fourth of the samples each. The PCR cycles were performed as follows: 98^{o}C for

3min, 35 cycles of 98^{o}C denaturation for 10 s, 63^{o}C annealing for 10 s and 72^{o}C elongation for 10 s,

followed by a final extension at 72^{o}C for 1 min and hold at 10^{o}C until further processing. The PCR

products were run on a 2% agarose gel and purified from gel using a NucleoSpin Gel and PCR Clean-

up kit (Macherey-Nagel). The DNA concentration of the samples was measured and samples were

diluted to 1 ng/μL in 10 μL, of which 5 μL were used as template for the indexing PCR.

Indexing was performed with the kit IDT for Illumina Nextera DNA UD Index Set C (ref 20026934)

with a PCR mix as follows: 5 μL DNA template, 5 μL primer mix, 5 μL Phusion GC Buffer, 0.5 μL

10mM dNTPs, 0.25 μL Phusion DNA polymerase (M0530S, NEB) and 9.25 μL H$_2$O. The following

program was used: 98^{o}C for 3 min, followed by 8 cycles of 98^{o}C denaturation for 10 s, 55^{o}C annealing

for 10 s and 72^{o}C elongation for 10 s, and then a final extension at 72^{o}C for 1 min and hold at 10^{o}C until

further processing. The PCR products were purified using 1.8 volumes of NEBNext Sample Purification

Beads (E7767S, NEB). Samples were resuspended in 20 μL H$_2$O and a quality check was run on a

TapeStation 4150 device (Agilent) using DNA ScreenTape chips (Agilent) as recommended by the

manufacturer. Sequencing was performed as previously described (ref) by Illumina sequencing at the

Genomics Facility Basel (ETH Zürich, University of Basel) using a NextSeq SR 81/10 kit.

Data analysis was performed using the European Galaxy server[40] using the previously described

MAGeCK pipeline[41] using median normalization, FDR-adjusted threshold of 0.25, FDR as p-value

adjustment method and 10 as remove zero threshold. The median method as used for the gene log-fold

change. Data was plotted using Prism.

Sample preparation for proteomics

Overnight cultures of *P. aeruginosa* were grown at 37^{o}C and 200 rpm. Day cultures in LB were launched

(1:100 dilution of the overnight) and incubated for 2.5-3h at 37^{o}C and 200 rpm, until they reached an

OD_{600nm} of 0.7-1.4. They were concentrated to an OD of 10 and 20 μL were spotted on LB plates. The

spots were allowed to dry and were then incubated at 37^{o}C for 3 h. The spots were then resuspended in

500 μL phosphate-buffered saline (PBS), pelleted by centrifugation at 12,000g for 2 min and stored at -

80^{o}C until further processing. Cell pellets were resuspended in lysis buffer (5% Sodium dodecyl sulfate

(SDS), 10mM tris(2-carboxyethyl) phosphine (TCEP), 100 mM Triethyloammonium bicarbonate

(TEAB)) followed by incubation at 95°C for 10 min. Cells were disrupted by ultrasonication using the

PIXUL system (Active Motif) for 20 min with default settings (pulse 50 cycles, PRF 1 kHz, burst rate

20 Hz)) and the protein content was determined by tryptophan-based fluorescence assay (Infinite M

Plex, Tecan). Sample alkylation was performed by addition of 20 mM iodoacetamide (IAA) and

incubation at 25°C for 30 min with gentle shaking. 10 μg of protein lysate was digested (trypsin 1/100,

w/w; Promega) and purified using the SP3 protocol[42]. Peptide concentration was determined using an

UV-based assay (Infinite M Nano, Tecan). Samples were then resuspended at a final concentration of

250 ng/μL.

SWATH-MS data acquisition

Samples were separated on a Dionex UltiMate 3000 system (ThermoFisher Scientific) coupled online

to an Orbitrap Exploris 480 mass spectrometer (ThermoFisher Scientific). In-house packed 20 cm, 75

μm ID capillary column with 1.9 μm Reprosil-Pur C18 beads (Dr. Maisch, Ammerbuch, Germany) was

used. The column temperature was maintained at 60 °C using an integrated column oven interfaced

online with the mass spectrometer. Formic acid (FA) 0.1% was used to buffer the pH in the two running

buffers used. The total gradient time was 60 min and went from 2% to 12% acetonitrile (ACN) in 5 min,

followed by 45 min to 35%, and 10 min at 50%. This was followed by a washout by 95% ACN, which

was kept for 20 min, followed by re-equilibration of 0.1% FA buffer. Flow rate was kept at 300 nL/min.

Spray voltage was set to 2500 V, funnel RF level at 40, and heated capillary at 275 °C. For DIA

experiments full MS resolutions were set to 120,000 and full MS normalized AGC target was 300%

with an IT of 45 ms. Mass range was set to 350–1400. Normalized AGC target value for fragment

spectra was set at 1000%. In all, 63 windows of 9 Da were used with an overlap of 1 Da. Resolution

was set to 15,000 and IT to 22 ms. Normalized CE was set at 28%. All data were acquired in centroid

mode using positive polarity and advanced peak determination was set to on.

SWATH-MS data analysis

For data processing and protein identification, raw data were imported into SpectroNaut

(16.1.220730.53000, Biognosys) and analyzed with directDIA. Searches were carried out against a fasta

file including the proteomes of *P. aeruginosa* (UP000002438). Cysteine carbamidomethylation was set

as fixed modification. Methionine oxidation, methionine excision at the N terminus were selected as

variable modifications. Results were filtered for a 1% false discovery rate (FDR) on spectrum, peptide,

and protein levels. Relative protein abundances and pairwise comparisons were calculated using the

MSstats package[43] with default settings selected.

CPRG lysis assay

Overnight cultures of *P. aeruginosa* and *V. cholerae*, were grown in LB at 37°C and 200 rpm without

antibiotics. *A. baylyi* overnight cultures were grown at 30°C in similar conditions. Day cultures were

launched in LB at the following dilutions: 1:100 for *P. aeruginosa*, 1:50 for *A. baylyi* and 1:400 for *V:*

cholerae. After 2.5-3h, when all cultures reached an OD_{600nm} of 0.7-1.4, cultures were spun down at

12,000g for 2 min at room temperature. The cultures were concentrated to OD 1 and mixed in a 1:2

prey:attacker ratio. Three μL of the resulting suspensions were spotted in wells of a flat-bottom 96-well

plate that had been extemporaneously prepared with 150 μL LB with 1% agar containing 0.1 mM IPTG

(isopropyl-β-D-thiogalactoside) and 20 µg/mL CPRG (red-β-D-galactopyranoside) and pre-dried for 20

min at room temperature next to the Bunsen burner. The plates were incubated in an Epoch2 plate reader

(BioTek) at 30^{o}C, and the OD_{575nm} was measured every 10 min. The curves were plotted with Prism.

Flow cytometry

466 Bacteria with different fluorescently labelled T6SS sheaths (TssB1-mNG or TssB2-mCh2) were grown

467 to mid-exponential phase at 37^{o}C from an overnight culture in LB. Prior to measurement, the samples

468 were diluted 100x in PBS and were analysed in a BD Fortessa flow cytometer, with 950V green and

469 950V red laser in low flow rate mode. Data were analysed using FlowJo v10, and the median

470 fluorescence intensity (MFI) was plotted using Prism.

Growth curves

472 Overnight cultures of *P. aeruginosa* were diluted to OD_{600nm} 1 in LB, and used to inoculate wells, in

473 triplicate, of a flat-bottom 96-well plate containing LB or LB without NaCl, with or without antibiotics,

474 at a starting OD_{600nm} of 0.05. They were incubated at 37^{o}C in an Epoch2 plate reader (BioTek) with

475 orbital shaking at 240cpm and the OD_{600nm} was measured every 10 min. If antibiotic treatment was

476 performed, antibiotics were added after 1.5h of incubation. The resulting curves were plotted in Prism.

Fluorescence microscopy

Overnight cultures of *P. aeruginosa* were diluted 100-fold in LB and incubated for around 3h at 37^{o}C

and 200 rpm. They were then pelleted on a tabletop centrifuge at 12.000g for 2 min at room temperature

and resuspended to OD 10 with LB containing 4 µg/mL FM4-64 (T13320, ThermoFisher Scientific)

and incubated for 10 min at room temperature in the dark. One µL of the cell suspensions were then

placed on agar pads (1/3 LB, 2/3 PBS with 1% agarose) and imaged on a Nikon Ti-E inverted motorized

microscope with Perfect Focus System and Plan Apo 200x Oil Ph3 DM (NA 1.4) objective lens. Spectra

X light engine (Lumencor) and ET-mCherry (Chroma 49008) filter sets were used to excite and filter

fluorescence and a sCMOS camera pco.edge 4.2 (PCO, Germany, pixel size 65nm) and VisiView

software (Visitron Systems, Germany) were used. Imaging was carried out at 30 ^{o}C by an Okolab T-unit

(Okolab).

Whole genome sequencing

Genomic DNA was extracted from overnight bacterial cultures in 1.5mL LB following the instructions of the GenElute Bacterial Genomic DNA kit (NA2110, Sigma). DNA concentration was measured using the Qubit dsDNA Quantitation Broad Range Assay kit (Q32850, ThermoFisher Scientific) and barcoding was performed according to the manufacturer's instructions using the Rapid Barcoding Kit 24 V14 (SQK-RBK114.24, Oxford Nanopore Technologies). Samples were sequenced on a MinION flow cell (R10.4.1, FLO-MIN114, Oxford Nanopore Technologies). Basecalling was performed using Dorado software from Nanopore and the genomes were compared using the Snakemake evo-genome-analysis pipeline that is publicly available at https://github.com/mmolari/evo-genome-analysis/. This pipeline uses the raw reads from each sample and maps it to the reference using Minimap2[44], extracting genomic changes including single nucleotide polymorphisms, gaps, insertions, clips and rearrangements.

Statistics

When only two means were compared, we performed unpaired t-tests. When multiple means were compared, we used rdinary one-way analysis of variance (ANOVA) with Dunnett's multiple comparison test, using GraphPad Prism version 9.3.1. Unless otherwise stated, data are represented as mean ± standard deviation (SD). *p-value<0.05, **p-value<0.005, ***p-value<0.0005, ****p-value<0.00005, ns – p-value>0.05.

Acknowledgements

This work was supported by the National Center of Competence in Research AntiResist funded by the Swiss National Science Foundation (51NF40_180541). We thank Mattia Zampieri for providing the antibiotic collection used in this study.

References

1. Malhotra, S., Hayes, D. & Wozniak, D. J. Cystic fibrosis and pseudomonas aeruginosa: The host-microbe interface. *Clin. Microbiol. Rev.* **32**, (2019).

2. Sana, T., Berni, N. & Bleves, S. The T6SSs of Pseudomonas aeruginosa Strain PAO1 and Their Effectors: Beyond Bacterial-Cell Targeting. *Front. Cell. Infect. Microbiol.* **6**, 61 (2016).

3. Swart, A. L. *et al.* Pseudomonas aeruginosa breaches respiratory epithelia through goblet cell invasion in a microtissue model. *Nat. Microbiol.* **9**, 1725–1737 (2024).

4. Wang, J., Brodmann, M. & Basler, M. Assembly and subcellular localization of bacterial type VI secretion systems. *Annu. Rev. Microbiol.* **73**, 621–638 (2019).

5. Yang, X., Long, M. & Shen, X. Effector–Immunity Pairs Provide the T6SS Nanomachine its Offensive and Defensive Capabilities. *Mol. A J. Synth. Chem. Nat. Prod. Chem.* **23**, (2018).

6. Zhang, H., Gao, Z. Q., Wei, Y., Xu, J. H. & Dong, Y. H. Insights into the cross-immunity mechanism within effector families of bacteria type VI secretion system from the structure of StTae4-EcTai4 complex. *PLoS One* **8**, (2013).

7. Basler, M., Ho, B. T. & Mekalanos, J. J. Tit-for-tat: Type VI secretion system counterattack during bacterial cell-cell interactions. *Cell* **152**, 884–894 (2013).

8. George, M., Narayanan, S., Tejada-Arranz, A., Plack, A. & Basler, M. Initiation of H1-T6SS dueling between Pseudomonas aeruginosa. *MBio* **15**, (2024).

9. Smith, W. P. J. *et al.* The evolution of tit-for-tat in bacteria via the type VI secretion system. *Nat. Commun.* **11**, (2020).

10. Ringel, P. D., Hu, D. & Basler, M. The Role of Type VI Secretion System Effectors in Target Cell Lysis and Subsequent Horizontal Gene Transfer. *Cell Rep.* **21**, 3927–3940 (2017).

11. Crisan, C. V. & Hammer, B. K. The Vibrio cholerae type VI secretion system: toxins, regulators and consequences. *Environmental Microbiology* vol. 22 4112–4122 (2020).

535   12.   Ting, S. Y. *et al.* Discovery of coordinately regulated pathways that provide innate protection
536   against interbacterial antagonism. *Elife* **11**, (2022).

537   13.   Hersch, S. J., Manera, K. & Dong, T. G. Defending against the Type Six Secretion System:
538   beyond Immunity Genes. *Cell Rep.* **33**, 108259 (2020).

539   14.   Liang, X. *et al.* An onboard checking mechanism ensures effector delivery of the type VI
540   secretion system in Vibrio cholerae. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 23292–23298 (2019).

541   15.   Robert-Genthon, M. *et al.* Unique features of a Pseudomonas aeruginosa α2-macroglobulin
542   homolog. *MBio* **4**, (2013).

543   16.   Zouhir, S. *et al.* Assembly of an atypical α-macroglobulin complex from Pseudomonas
544   aeruginosa. *Sci. Rep.* **8**, (2018).

545   17.   Rehman, A. A., Ahsan, H. & Khan, F. H. α-2-Macroglobulin: a physiological guardian. *J. Cell.*
546   *Physiol.* **228**, 1665–1675 (2013).

547   18.   Marrero, A. *et al.* The crystal structure of human α2-macroglobulin reveals a unique molecular
548   cage. *Angew. Chem. Int. Ed. Engl.* **51**, 3340–3344 (2012).

549   19.   Meyer, C., Hinrichs, W. & Hahn, U. Human α2-macroglobulin--another variation on the venus
550   flytrap. *Angew. Chem. Int. Ed. Engl.* **51**, 5045–5047 (2012).

551   20.   Lin, Y. *et al.* The phospholipid-repair system LplT/Aas in Gram-negative bacteria protects the
552   bacterial membrane envelope from host phospholipase A2 attack. *J. Biol. Chem.* **293**, 3386–
553   3398 (2018).

554   21.   Chevalier, S. *et al.* Structure, function and regulation of Pseudomonas aeruginosa porins.
555   *FEMS Microbiol. Rev.* **41**, 698–722 (2017).

556   22.   De Mot, R. & Vanderleyden, J. The C-terminal sequence conservation between OmpA-related
557   outer membrane proteins and MotB suggests a common function in both gram-positive and
558   gram-negative bacteria, possibly in the interaction of these domains with peptidoglycan. *Mol.*
559   *Microbiol.* **12**, 333–334 (1994).

560   23.   Koebnik, R. Proposal for a peptidoglycan-associating alpha-helical motif in the C-terminal

561   regions of some bacterial cell-surface proteins. *Mol. Microbiol.* **16**, 1269–1270 (1995).

562   24.   Rawling, E. G., Brinkman, F. S. L. & Hancock, R. E. W. Roles of the carboxy-terminal half of

563   Pseudomonas aeruginosa major outer membrane protein OprF in cell shape, growth in low-

564   osmolarity medium, and peptidoglycan association. *J. Bacteriol.* **180**, 3556–3562 (1998).

565   25.   Gotoh, N., Wakebe, H., Yoshihara, E., Nakae, T. & Nishino, T. Role of protein F in

566   maintaining structural integrity of the Pseudomonas aeruginosa outer membrane. *J. Bacteriol.*

567   **171**, 983 (1989).

568   26.   Ishida, H., Garcia-Herrero, A. & Vogel, H. J. The periplasmic domain of Escherichia coli outer

569   membrane protein A can undergo a localized temperature dependent structural transition.

570   *Biochim. Biophys. Acta - Biomembr.* **1838**, 3014–3024 (2014).

571   27.   Deghelt, M. *et al.* The outer membrane and peptidoglycan layer form a single mechanical

572   device balancing turgor. *bioRxiv* 2023.04.29.538579 (2023) doi:10.1101/2023.04.29.538579.

573   28.   Duchene, M., Barron, C., Schweizer, A., Von Specht, B. U. & Domdey, H. Pseudomonas

574   aeruginosa outer membrane lipoprotein I gene: molecular cloning, sequence, and expression in

575   Escherichia coli. *J. Bacteriol.* **171**, 4130–4137 (1989).

576   29.   Mathelié-Guinlet, M., Asmar, A. T., Collet, J. F. & Dufrêne, Y. F. Lipoprotein Lpp regulates

577   the mechanical properties of the E. coli cell envelope. *Nat. Commun. 2020 111* **11**, 1–11

578   (2020).

579   30.   Brencic, A. & Lory, S. Determination of the regulon and identification of novel mRNA targets

580   of *Pseudomonas aeruginosa* RsmA. *Mol. Microbiol.* **72**, 612–632 (2009).

581   31.   Allsopp, L. P. *et al.* RsmA and AmrZ orchestrate the assembly of all three type VI secretion

582   systems in Pseudomonas aeruginosa. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 7707–7712 (2017).

583   32.   Kamal, F. *et al.* Differential Cellular Response to Translocated Toxic Effectors and Physical

584   Penetration by the Type VI Secretion System. *Cell Rep.* **31**, 107766 (2020).

33.   Hersch, S. J. *et al.* Envelope stress responses defend against type six secretion system attacks

independently of immunity proteins. *Nat. Microbiol.* **5**, 706–714 (2020).

34.   Yin, C., Alam, M. Z., Fallon, J. T. & Huang, W. Advances in Development of Novel

Therapeutic Strategies against Multi-Drug Resistant Pseudomonas aeruginosa. *Antibiot. 2024,*

Vol. 13, Page 119 **13**, 119 (2024).

35.   Sabnis, A. *et al.* Colistin kills bacteria by targeting lipopolysaccharide in the cytoplasmic

membrane. *Elife* **10**, (2021).

36.   Mcdonnell, G. & Russell, A. D. Antiseptics and disinfectants: activity, action, and resistance.

Clin. Microbiol. Rev. **12**, 147–179 (1999).

37.   Rietsch, A., Vallet-Gely, I., Dove, S. L. & Mekalanos, J. J. ExsE, a secreted regulator of type

III secretion genes in Pseudomonas aeruginosa. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 8006–8011

(2005).

38.   Gibson, D. G. *et al.* Enzymatic assembly of DNA molecules up to several hundred kilobases.

Nat. Methods **6**, 343–345 (2009).

39.   Plum, M. T. W. *et al.* Burkholderia thailandensis uses a type VI secretion system to lyse

protrusions without triggering host cell responses. *Cell Host Microbe* **32**, 676-692.e5 (2024).

40.   Abueg, L. A. L. *et al.* The Galaxy platform for accessible, reproducible, and collaborative data

analyses: 2024 update. *Nucleic Acids Res.* **52**, W83–W94 (2024).

41.   Li, W. *et al.* Quality control, modeling, and visualization of CRISPR screens with MAGeCK-

VISPR. *Genome Biol.* **16**, 281 (2015).

42.   Hughes, C. S. *et al.* Single-pot, solid-phase-enhanced sample preparation for proteomics

experiments. *Nat. Protoc.* **14**, 68–85 (2019).

43.   Choi, M. *et al.* MSstats: an R package for statistical analysis of quantitative mass spectrometry-

based proteomic experiments. *Bioinformatics* **30**, 2524–2526 (2014).

44.   Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–3100

(2018).

Figure legends.

Figure 1. *P. aeruginosa* is more resistant to T6SS from *A. baylyi* and *V. cholerae* than *E. coli*. CFU

counts showing the survival of *P. aeruginosa* when competed against *A. baylyi* (*Ab*) (A) and *V. cholerae*

(*Vc*) (B) strains that have or not an active T6SS, in a 10:1 attacker:prey ratio where *P. aeruginosa* is the

prey. (C) CFU counts showing the survival of *E. coli* (*Ec*) when competed against *A. baylyi* and *V.*

cholerae in 10:1 and 1:1 attacker:prey ratios where *E. coli* is the prey.

Figure 2. CRISPRi screening for *P. aeruginosa* knockdowns that are sensitive to T6SS attacks. (A)
Schematic of the experimental design. *P. aeruginosa* lacking *tssM1* and carrying the sgRNA library was
competed against T6SS-positive or –negative *A. baylyi* and *V. cholerae* for 1 or 2 rounds of competition
lasting 3h each. Bacteria were grown to mid-exponential phase before each competition round. (B)
Negative-selection Volcano plot showing genes important for resistance to T6SS from *V. cholerae*. In
red are previously identified genes and in yellow genes newly identified in this study. (C) Negative-

selection Volcano plot showing genes important for resistance to T6SS from *A. baylyi*. (D) Volcano plot

showing proteins that are differentially regulated in a Δ*gacA* strain compared to wild-type *P. aeruginosa*.

In red are components of the H1-T6SS, in pink components of the H2-T6SS, in blue previously

identified members of the GacA regulon that are linked to resistance to T6SS attacks of *B. thailandensis*

and in yellow the newly identified hits from the *mag* operon. *oprF* is also highlighted as a non-

differentially regulated gene. (E) CFU counts showing the survival of different *P. aeruginosa* strains

when competed against T6SS-positive or negative *A. baylyi* and *V. cholerae*. (F) CFU counts showing

the survival of *P. aeruginosa* Δ*gacA* when competed against *A. baylyi* lacking different T6SS effectors.

(G) CFU counts showing the survival of *P. aeruginosa* Δ*gacA* when competed against *V. cholerae*

carrying inactivated versions of different T6SS effectors.

Figure 3. MagD is important for resistance against certain PG-targeting effectors. (A) CFU counts showing the survival of different *P. aeruginosa* strains when competed against *V. cholerae* carrying different inactivated effectors. (B) CPRG lysis assay showing the lysis levels over time of different *P. aeruginosa* strains when competed against *V. cholerae*. (C) CPRG lysis assay showing the lysis levels over time of different *P. aeruginosa* strains when competed against *A. baylyi* carrying only its PG-targeting effector, Tae1.

Figure 4. Arc1A, Arc3A and Aas are important for resistance to lipase effectors. CPRG lysis assay

showing the lysis levels over time of different *P. aeruginosa* strains when competed against *A. baylyi*

carrying only its lipase effector, Tle1.

Figure 5. Anchoring of the PG to the outer membrane is important for T6SS resistance. (A) CFU

counts showing the survival of different *P. aeruginosa* strains with or without *oprF* when competed

against *V. cholerae* strains carrying inactivated T6SS effectors. (B) CFU counts showing the survival of

different *P. aeruginosa* strains when competed against *A. baylyi* strains lacking single T6SS effectors.

(C) CFU counts showing the survival of different *P. aeruginosa* strains complemented with wild-type

or mutant variants of *oprF* when competed against *V. cholerae* and *A. baylyi*.

Figure 6. Δ*oprF* revertants grow normally and recover wild-type morphology but fail to resist T6SS attacks. (A) Image of a *P. aeruginosa* strain lacking *tssM1* and *oprF* growing on an LB no salt plate after overnight incubation at 37°C. (B) Growth curves of parental and revertant strains of *P. aeruginosa* Δ*tssM1* Δ*oprF*. (C) Fluorescence microscopy images of parental and revertant strains of *P. aeruginosa* lacking *tssM1* and *oprF*. (D) CFU counts showing the survival of the different *P. aeruginosa* revertants when competed against T6SS-active or –inactive *V. cholerae*. (E) Growth curve of wild-type and mutant *P. aeruginosa* strains lacking *oprF* and *gacA* or not.

Figure 7. T6SS resistance mechanisms are species- or effector-specific. CFU counts showing the survival of *P. aeruginosa* strains lacking different T6SS resistance mechanisms when competed with *B. thailandensis* (*Bt*) in a 10:1 attacker:prey ratio where *P. aeruginosa* is the prey.

Figure 8. General T6SS resistance mechanisms also influence antibiotic resistance. Growth curves of *P. aeruginosa* wild-type or Δ*oprF* with or without 20 μM tetracycline hydrochloride (A), piperacillin sodium salt (B) or colistin sulfate (C). (D) Growth curves of *P. aeruginosa* wild-type or Δ*magD* with or without 20 μM chlorhexidine.

    Supplementary Figure legends.

    Supplementary Figure 1. CFU counts showing the survival of the attacker when competing against *A.*

    baylyi (*Ab*) (A) and *V. cholerae* (*Vc*) (B) strains that have or not an active T6SS, in a 10:1 attacker:prey

    ratio where *P. aeruginosa* is the prey. (C) CFU counts showing the survival of *A. baylyi* and *V. cholerae*

    when competed against *E. coli* (*Ec*) in 10:1 and 1:1 attacker:prey ratios where *E. coli* is the prey.

Supplementary Figure 2. Log fold change of each sgRNA targeting the statistically significant hits of the CRISPRi screening where *P. aeruginosa* was competed against *V. cholerae* (A) or *A. baylyi* (B). (C) Mean Fluorescence Intensity (MFI) of Green Fluorescent Protein (GFP) of wild type, ΔretS and ΔgacA strains where the sheath protein TssB1 of the H1-T6SS is labelled with mNeongreen where indicated. (D) CFU counts showing the survival of *A. baylyi* and *V. cholerae* when competed against *P. aeruginosa* strains in a 10:1 attacker:prey ratio where *P. aeruginosa* is the prey. (E) CFU counts of *A. baylyi* strains lacking different T6SS when competed against *P. aeruginosa* ΔgacA in a 10:1 ratio. (F) CFU counts of *V. cholerae* strains carrying different T6SS inactivated effectors when competed against *P. aeruginosa* ΔgacA in a 10:1 ratio.

689 **Supplementary Figure 3.** CFU counts of *V. cholerae* strains carrying different T6SS inactivated

690 effectors when competed against *P. aeruginosa* with or without *magD* in a 10:1 ratio.

Supplementary Figure 4. (A) CFU counts of *V. cholerae* strains carrying different T6SS inactivated

effectors when competed against *P. aeruginosa* with or without *oprF* in a 10:1 ratio. (B) CFU counts of

A. baylyi strains lacking different T6SS when competed against *P. aeruginosa* with or without *oprF* in

a 10:1 ratio. (C) Growth curve of *P. aeruginosa* Δ*tssM1* and Δ*tssM1* Δ*oprF* strains in LB and LB without

salt. (D) Growth curve of *P. aeruginosa* Δ*tssM1* Δ*oprF* strains complemented *in trans* with wild-type

or an R296E variant of *oprF* in LB without salt. (E) CFU counts of *V. cholerae* and *A. baylyi* strains

with or without an active T6SS when competed against *P. aeruginosa* with different *oprF* variants in a

10:1 ratio. (F) CFU counts of *V. cholerae* when competed against *P. aeruginosa* Δ*tssM1* Δ*oprF*

revertants in a 10:1 ratio.

702 **Supplementary Figure 5.** CFU counts of *B. thailandensis* with active or inactive T6SS-1 when

703 competed against *P. aeruginosa* strains lacking different T6SS resistance mechanisms in a 10:1 ratio.

C

Supplementary Figure 6. Growth curves of *P. aeruginosa* strains treated with a DMSO control (black curves) or different antibiotics (red curves) at a concentration of 20 µM. The grey arrow indicates the point at which antibiotics were added. (A) Curves for the wild-type parental strain. (B) Curves for the Δ*oprF* strain. (C) Curves for the Δ*magD* strain. The treatments that show different behaviours compared to the parental strain are highlighted with a yellow background.