Foxi1 regulates multiple steps of mucociliary development and ionocyte specification through transcriptional and epigenetic mechanisms

Sarah Bowden[1,2,3,&], Magdalena Maria Brislinger-Engelhardt[1,2,4,&], Mona Hansen[1,4,&], Africa Temporal-Plo[1,2,3], Damian Weber[1,2], Sandra Hägele[1,2], Fabian Lorenz[2,5], Tim Litwin[2,5], Clemens Kreutz[2,5], Peter Walentek[1,2,3,4,#]

[1] Internal Medicine IV, Medical Center - University of Freiburg, Hugstetter Strasse 55, 79106 Freiburg, Germany.
[2] CIBSS Centre for Integrative Biological Signalling Studies, University of Freiburg, Schänzlestrasse 18, 79104 Freiburg, Germany.
[3] IMPRS-IEM International Max Planck Research School of Immunobiology, Epigenetics and Metabolism, Max Planck Institute of Immunobiology and Epigenetics, Stübeweg 51, 79108 Freiburg, Germany.
[4] SGBM Spemann Graduate School for Biology and Medicine, University of Freiburg, Albertstrasse 19A, 79104 Freiburg, Germany.
[5] IMBI Institute of Medical Biometry and Statistics, Medical Center - University of Freiburg, Stefan-Meier-Strasse 26, 70104, Freiburg, Germany.
[&] These authors contributed equally and are listed in alphabetical order.
[#] Corresponding author: peter.walentek@medizin.uni-freiburg.de

Key words:

Xenopus epidermis, development, mucociliary, ionocytes, multipotent progenitors

Abstract

Foxi1 is a master regulator of ionocytes (ISCs / INCs) across species and organs. Two subtypes of ISCs exist, and both α- and β-ISCs regulate pH- and ion-homeostasis in epithelia. Gain and loss of FOXI1 function are associated with human diseases, including Pendred syndrome, male infertility, renal acidosis and cancers. Foxi1 functions were predominantly studied in the context of ISC specification, however, reports indicate additional functions in early and ectodermal development. Here, we re-investigated the functions of Foxi1 in *Xenopus laevis* embryonic mucociliary epidermis development and found a novel function for Foxi1 in the generation of Notch-ligand expressing mucociliary multipotent progenitors (MPPs). We demonstrate that Foxi1 has multiple concentration-dependent functions: At low levels, Foxi1 confers ectodermal competence through transcriptional and epigenetic mechanisms, while at high levels, Foxi1 induces a multi-step process of ISC specification and differentiation. We further describe how *foxi1* expression is affected through auto- and Notch-regulation, how Ubp1 and Dmrt2 regulate ISC subtype differentiation, and how this developmental program affects Notch signaling as well as mucociliary patterning. Together, we reveal novel functions for Foxi1 in *Xenopus* mucociliary epidermis formation, relevant to our understanding of vertebrate development and human disease.

Introduction

The Forkhead-box transcription factor Foxi1 is a master regulator of ionocytes (ISCs) in

the lung, kidney, inner ear and epididymis across vertebrates (Hulander et al., 2003;

Pou Casellas et al., 2023). ISCs regulate ion homeostasis through the expression of an

array of transmembrane solute carriers (e.g. Pendrin encoded by *slc26a4* and Anion

exchanger 1 encoded by *slc4a1*) and pH-regulators (e.g. vacuolar (v)H$^+$ATPase

encoded by *atp6* subunit genes and Carbonic anhydrase encoded by *ca* genes). α-ISCs

and β-ISCs are subtypes with apical vs. basolateral vH$^+$ATPase localization and

differential transporter expression of *slc26a4* and *slc4a1* (Quigley et al., 2011). In ISCs

of the mammalian airway mucociliary epithelium, Foxi1 also regulates the expression of

cystic fibrosis transmembrane conductance regulator (*CFTR*), thereby controlling

chloride secretion and mucus properties (Montoro et al., 2018; Plasschaert et al., 2018;

Scudieri et al., 2020). Mutations in *FOXI1* and its transcriptional target solute carriers

cause Pendred syndrome and hearing loss, male infertility, and distal renal tubular

acidosis (Blomqvist et al., 2004; Blomqvist et al., 2006; Hulander et al., 2003; Yang et

al., 2007). In contrast, Foxi1 overexpression is found in cancer subtypes, e.g. in

chromophore renal cell carcinoma (chRCC) and in pulmonary large cell carcinoma

(LCC) (Lindgren et al., 2017; Simbolo et al., 2024; Skala et al., 2020; Yamada et al.,

2022).

The *Xenopus* embryonic epidermis is a popular model for vertebrate mucociliary

epithelia development and disease studies (Walentek and Quigley, 2017). It forms

Foxi1-dependent α-ISCs and β-ISCs as well as multiciliated cells (MCCs), secretory

cells (small secretory cells (SSCs) and goblet cells) and basal stem cells, similar to the

mammalian airway epithelium (Brooks and Wallingford, 2014; Haas et al., 2019; Hayes

et al., 2007; Quigley et al., 2011; Walentek, 2022; Walentek et al., 2014). Knockdown of

foxi1 in *Xenopus* caused a loss of epidermal ISCs as expected, but also induced

defective ciliation in MCCs suggesting potential Foxi1 functions in other cell types

(Dubaissi and Papalopulu, 2011; Mir et al., 2007). Foxi1 plays additional early roles in

Xenopus ectoderm development: In the blastula and gastrula, *foxi1* is initially activated

throughout the entire ectoderm where it is required as a determinant counteracting

vegetal mesendoderm-inducing factors (Mir et al., 2007; Suri et al., 2005). In neurula

stages, Foxi1 is required for placode formation in the neural plate border and for the

specification of ISCs in the epidermis (Dubaissi and Papalopulu, 2011; Maharana and

Schlosser, 2018; Quigley et al., 2011).

How Foxi1 regulates different aspects of *Xenopus* mucociliary epidermis development

from ectoderm to ISC-subtype specification, and how it could affect development of

additional mucociliary epidermis cell types remains unresolved. Understanding the

range of Foxi1 functions in this important model for mucociliary biology could provide

additional information regarding unappreciated Foxi1 functions in development and

human disease.

Here, we re-examined the roles of Foxi1 in *Xenopus* mucociliary epidermis

development. We found that Foxi1 acts in a concentration-dependent manner: At low

levels, Foxi1 marks multipotent mucociliary progenitors (MPPs) and establishes

ectodermal identity, at least in part through regulation of chromatin accessibility. At high

levels, Foxi1 induces ISC specification and subtype differentiation through Ubp1 and

Dmrt2. High Foxi1 levels are achieved by positive auto-regulation and inhibited by

Notch signaling. In this work, we elucidate novel concentration-dependent Foxi1

transcriptional and epigenetic functions in mucociliary development and ISC

specification.

Results

Specification and differentiation of ISCs is a multi-step process

Previous work has defined a core ISC gene set in the *X. laevis* mucociliary epidermis

(Quigley and Kintner, 2017). We generated epidermal mucociliary organoids from

animal cap explants and conducted bulk RNA-sequencing (RNA-seq) analysis of core

ISCs gene expression during epidermis development (Brislinger-Engelhardt et al., 2023;

Haas et al., 2019; Walentek, 2018). Z-scores of normalized counts (TPM) of ISC

transcripts were clustered to reveal dynamic co-expression (**Fig. 1A**). Five clusters

clearly separated along developmental time, with cluster I being the only set of genes

displaying strong expression during very early and late developmental stages, but not

during cell fate specification stages (st. 10-16). Cluster II contained *foxi1*, the Notch

ligand *dll1* and the cell cycle regulator *gadd45g*. Cluster III contained *ca12* and the

transcription factor *ubp1*, which was shown to induce ISC formation upon

overexpression in the epidermis (Quigley et al., 2011). Cluster IV contained multiple

transcription factors, including *tfcp2l1*, required for ISC formation in mouse kidney

(Werth et al., 2017). Cluster V was dominated by *slc26a4* and *atp6*-subunit expression

during later differentiation of ISCs (st. 20–32). These data indicated that ISCs develop

using a multi-step process.

To validate the core ISC gene set, we used manipulations to deplete or enrich ISCs in

mucociliary organoids and RNA-seq at the beginning of cell fate specification (st. 10), at

the end of cell fate specification (st. 16), during maturation and intercalation of ISCs (st.

25), and in the mature epithelium (st. 32) (Walentek and Quigley, 2017). As previously

described (Quigley et al., 2011), increased Notch signaling (Notch intracellular domain

(*nicd*) mRNA injections) inhibited core ISC gene expression, while inhibition of Notch

signaling (injection of dominant-negative suppressor of hairless/RBPJ (*suh-dbm*)

mRNA) promoted core ISC gene expression (**Fig. S1A,B**). Inhibition of Notch signaling

in combination with MCC inhibition (by co-injection of *dominant-negative mcidas* (*dn-*

mcidas) mRNA (Stubbs et al., 2012)) further increased core ISC gene expression,

reflecting stronger overproduction of ISCs. However, expression of *tfcp2l1*, *atp6v0d1.L*

and *csta.L* were reduced in these conditions suggesting that they might be expressed

more in MCCs than in ISCs (**Fig. S1C**). Notch repression of *foxi1* by *nicd* was

substantial but not statistically significant (**Fig. S1A**), potentially reflecting a Notch-

independent function of Foxi1.

We used single-cell RNA-seq (scRNA-seq) data from at late *Xenopus laevis* epidermis

stages containing mature ISCs (Aztekin et al., 2019). These data confirmed high levels

of *foxi1*, *ubp1* and *atp6*-subunit expression in both ISC subtypes, while *slc4a1* was

specifically expressed in α-ISCs and *slc26a4* in β-ISCs, as previously described (**Fig.**

S1D). Interestingly, the transcription factor *dmrt2* (cluster IV) was exclusively expressed

by α-ISCs (**Fig. S1D**), and loss of kidney α-ISCs was recently reported in *Dmrt2*

knockout mice (Wu et al., 2024).

To test if *foxi1*, *ubp1* and *dmrt2* contribute differentially to ISC-subtype formation, we

knocked down each factor using morpholino oligonucleotides (MOs) and analyzed ISC

marker expression in the mature epidermis at st. 30-32 by whole-mount *in situ*

hybridization (WMISH). Knockdown of *foxi1* led to a loss of pan-ISC marker *atp6v1e1*,

which could be rescued and expanded by co- and over-expression of *foxi1* mRNA,

respectively (**Fig. 1B,C**). *foxi1* knockdown also strongly reduced expression of *ubp1*,

dmrt2, *slc24a6* and *slc4a1* (**Fig. 1C,D**), indicating a loss of both ISC subtypes in the

mucociliary epidermis. In contrast, knockdown of *ubp1* specifically reduced expression

of the β-ISC marker *slc26a4*, while *dmrt2* loss lead to inhibition of α-ISC-specific *slc4a1*

expression (**Fig. 1E-H**). Importantly, *ubp1* and *dmrt2* MOs did not abolish pan-ISC

expressed *foxi1* and *atp6v1e1* (**Fig. 1E-H**).

Taken together, these data suggest that Foxi1 determines ISC fate commitment, while

Ubp1 and Dmrt2 regulate ISC-subtype differentiation in a multi-step process.

Foxi1 has multiple concentration-dependent functions

In addition to ISC-specification, an earlier role for Foxi1 as ectodermal determinant

during blastula stages in *Xenopus* was described (Suri et al., 2005). However, our

experiments investigating ISC formation after *foxi1* MO did not indicate a loss of

epidermal identity.

To confirm this, we injected *foxi1* MO together with membrane GFP to identify targeted

cells, and analyzed cell type composition and morphology in the mature mucociliary

epidermis at st. 32 by immunofluorescence (IF) confocal microscopy (Walentek, 2018).

As previously described, we also observed reduced ciliation in MCCs after *foxi1*

knockdown (Dubaissi and Papalopulu, 2011) as well as strongly reduced ISC and

increased MCC numbers, with little effect on secretory cell types (SSCs and goblet

cells) or epidermal identity (**Fig. 2A**). This indicated that *foxi1* knockdown reduced Foxi1

levels enough to inhibit ISC specification, but not strong enough to prevent ectoderm

specification.

Next, we caused stronger depletion of Foxi1 using higher doses of *foxi1* MO. This

treatment induced delamination of cells in gastrula stage embryos (**Fig. S2A**), as

previously described (Mir et al., 2007), and frequent formation of skin lesions at st. 32

that could be rescued by co-injection of *foxi1* mRNA (**Fig. 2B**).

This suggests that low levels of Foxi1 are sufficient for ectoderm identity, while high

Foxi1 levels are required for ISC specification.

Foxi1 induces multipotent mucociliary progenitors

Next, we analyzed *foxi1* expression by WMISH. In early blastula/gastrula (st. 9/10)

stages *foxi1* is expressed at low levels in patches throughout the prospective ectoderm,

which start to resolve at st. 12 with individual cells strongly increasing *foxi1* expression

by st. 16, resulting in a salt-and-pepper pattern of individual cells by st. 32, representing

individual ISCs (**Fig. S2B**). Hence, *foxi1* seems to be initially expressed in more

epidermal cells than just in developing ISCs. To test if *foxi1* is expressed broadly in

epidermal multipotent mucociliary progenitors (MPPs), we generated a fluorescent

reporter (*foxi1::gfp-utrophin*) (**Fig. S2C,D**) using a previously characterized *foxi1*

promoter fragment (Cha et al., 2012).

First, we injected embryos with *foxi1::gfp-utrophin* and analyzed reporter activity at st.

32 by IF. Indeed, GFP signals were detected in ISCs, MCCs and SSCs, and even some

goblet cells expressed GFP at low levels (**Fig. 2C**). In contrast, a well-characterized

Mcidas/Foxj1-regulated promoter construct (Tasca et al., 2021) driving mScarletI

fluorescence (*α-tub::mscarletI*) was expressed predominantly in MCCs (**Fig. S3A,B**).

Next, we confirmed that reporter expression dynamics resemble endogenous *foxi1*

expression patterns during epidermis development using WMISH (**Fig. 2D, S3C**) and

GFP expression by IF (**Fig. 2E, S4A**). Reporter-driven *gfp* transcripts were detected at

st. 9 - 32, starting with non-epithelial low-level expression at st. 9/10, which increased

by st. 12/16 in deep and superficial layer cells, and at st. 32 expression was found

predominantly in epithelial layer cells (**Fig. 2D, S3C**). GFP-fluorescent cells were

detected from st. 10 onwards, predominantly in deep layer cells, but also in some cells

of the outer epithelial layer (**Fig. 2E, S4A**). During st. 12-16, an increasing number of

cells became GFP(+), including intercalating differentiating cells (**Fig. 2E, S4A**). During

st. 20-32, the number of GFP(+) cells decreased and fluorescent cells were

progressively confined to the epithelial outer cell layer - however, basal positioned

GFP(+) cells were detected even at st. 32 (**Fig. 2E, S4A**).

Together these data support the conclusion that *foxi1* is initially expressed in MPPs

during mucociliary development.

Foxi1(+) multipotent progenitors express Notch ligands during cell fate

specification

Dll1 expression was assigned to ISCs by Quigley and colleagues, similar to Foxi(+) cells

in the zebrafish skin and mammalian kidney (Janicke et al., 2007; Mukherjee et al.,

2020; Quigley and Kintner, 2017). This is supported by scRNA-seq data from *X.*

tropicalis development (Briggs et al., 2018) showing that *dll1* is transiently enriched in

differentiating ISCs (**Fig. S4B**). However, another study observed *dll1* expression

overlapping with different cell markers during patterning stages in the *Xenopus*

epidermis (Cibois et al., 2015). Our temporal expression analysis indicated that Cluster

II contained very early ISC genes, including *foxi1* and *dll1*, likely representing the MPPs

and early ISC differentiation stages (**Fig. 1A**), in line with both published observations.

To address if *dll1* (and *dlc*; Brislinger-Engelhardt et al., 2023) expression is part of the

differentiation program across mucociliary cell types or specific to MPPs and early ISCs,

we tested whether master transcription factors inducing cell types of the mucociliary

epidermis were able to induce Notch ligands prematurely. We overexpressed *foxi1* for

MPPs/ISCs, *mcidas* and *foxj1* for MCCs, *foxa1* for SSCs or *ΔN-tp63* for basal cells.

Only *foxi1* robustly induced *dll1* and *dlc* (**Fig. 3A, S4C**), and conversely, depletion of

Foxi1 prevented *dll1* expression during cell fate specification stages (**Fig. 3B**). These

results suggested that *dll1* is expressed in *foxi1*(+) MPPs and terminated by cell fate

induction of MCCs, SSCs and basal cells, but not in ISCs, which maintain *foxi1*

expression.

This raised the question how *dll1* expression is terminated during ISC differentiation.

The *X. tropicalis* scRNA-seq data revealed that *dll1* enrichment to ISCs is lost once

ubp1 is expressed (Briggs et al., 2018) (**Fig. S4B**). To test if Ubp1 terminates *dll1*

expression, we knocked down *ubp1* and analyzed embryos at the end of cell fate

specification (st. 16) by WMISH. *foxi1* was maintained and *dll1* expression was

prolonged after loss of Ubp1 (**Fig. 3C**). Analysis of cell type composition at st. 32 by IF

in *ubp1* morphants further revealed reduced MCC and SSC numbers as well as

appearance of intercalating cells with ambiguous morphology, likely representing

incompletely differentiated ISCs (**Fig. 3D**). These results demonstrated that

manipulating *foxi1* and ISC differentiation leads to dysregulated Notch dynamics during

mucociliary development, which can affect MCCs and other epidermal cell types.

Together, these data demonstrate that Notch ligands are expressed by *foxi1*(+) MPPs

during mucociliary patterning, and that Ubp1-dependent differentiation of ISCs or

induction of MCC, SSC and basal cell fates terminate Notch ligand expression.

Feedback- and auto-regulation during ISC specification

Feedback regulation of *dll1* by Notch signaling was suggested in *Xenopus* epidermis

development (Deblandre et al., 1999). RNA-seq analysis of ligand and receptor

expression after Notch manipulations confirmed that gain of Notch signaling suppresses

dll1, while blocking Notch increases and prolongs *dll1* expression (**Fig. S1A,B**).

Knockdown of *dll1* or *notch1* and subsequent analysis of cell type composition by IF at

st. 32 revealed a strong increase in ISCs, which was reversed by co-injection of *nicd*

(**Fig. S5A**), confirming that ISC specification is negatively regulated by Notch signaling

through Dll1 and Notch1. Furthermore, unilateral knockdown of *dll1* by different (low to

high) MO concentrations, and analysis of cell type markers between the targeted and

untargeted control side at st. 16/17 by WMISH confirmed that modifying Dll1 levels

sufficed to affect all cell types (**Fig. S5B**). Together, these data indicate that MPPs,

ligand production and ISC specification are negatively regulated by Notch feedback.

This raised the question how MPPs can achieve high *foxi1* expression levels required

for specification of ISC fate and Ubp1-/Dmrt2-dependent differentiation of mature ISC

subtypes. One potential mechanism for conferring robust cell fate decisions is positive

auto-regulation, and Foxi1 could activate its own expression using core Foxi motifs

previously identified in the *foxi1* promoter (Cha et al., 2012). To test this, we deleted a

Foxi2 binding region (**Fig. S2C,D**) and analyzed reporter activity at st. 32, i.e. long after

foxi2 expression is terminated. This strongly decreased reporter activity (**Fig. 4A**),

suggesting that core Foxi motifs are also used by Foxi1 to maintain its expression

through auto-regulation.

To verify that Foxi1 can also activate its own promoter without contributions from Foxi2,

we injected *foxi1::gfp-utrophin* vegetally to target the prospective mesendoderm, which

lacks maternally deposited *foxi2* (Cha et al., 2012). Analysis of reporter-only injected

cells (marked by membrane RFP) in hemisected embryos at st. 11 showed no reporter

activity in endodermal cells, while co-injection of *foxi1* mRNA led to ectopic activation of

the reporter (**Fig 4B**).

In summary, Foxi1 induces first Dll1-expressing MPPs, which increase Notch levels

during cell fate specification stages and terminates ISC production. Progenitors

experiencing low Notch levels activate high *foxi1* expression through auto-regulation to

induce ISC fate, and *dll1* expression is terminated in differentiating ISCs by Ubp1.

Foxi1 regulates genome accessibility in the epidermis

We wondered how early Foxi1 expression could have such a profound impact on ectodermal development and formation of mucociliary MPPs given that it is expressed at low levels. Besides its effects countering mesendoderm induction through transcriptional activation of ectodermal genes in early *Xenopus* embryos (Suri et al., 2005) Foxi1 has been shown to remain bound to condensed chromatin during mitosis, to remodel nucleosome structure and to alter the transcriptional ground state of cells in zebrafish embryos (Yan et al., 2006). This suggested a pioneer-like function for Foxi1.

To test if Foxi1 affects chromatin state and genomic accessibility in *Xenopus* epidermal development, we performed assays for transposase-accessible chromatin with sequencing (ATAC-seq) on st. 10 control epidermal organoids and organoids after knockdown of *foxi1*. This revealed a dramatic reduction in accessible chromatin regions after loss of Foxi1 (control: 311,328, *foxi1*MO: 146,640) (**Fig.5A**,**B**). In Foxi1-depleted organoids, 53.7% of accessible regions (169,077 peaks) were lost, 45.2% were maintained (142,251 peaks), and 1.1% were gained (4,389 peaks) (**Fig. 5B**). Next, we investigated which transcription factor binding motifs were enriched in regions lost, maintained or gained after *foxi1* MO. We found that motifs for factors with known functions in *Xenopus* ectodermal development were enriched in regions that lost accessibility after *foxi1* knockdown: e.g. Tfap2a and Tfap2c, Hic1, Rbfox2, Zac1 that regulate neural and neural crest formation as well as Tp63, which regulates epidermal basal stem cells, and Pitx1, required for cement gland formation (**Fig. 5C**) (Giudetti et al., 2014; Luo et al., 2005; Ma et al., 2007; Ray and Chang, 2020; Schweickert et al.,

2001; Zhang et al., 2006). In contrast, regions that remained open were enriched for

mesendodermal transcription factors (e.g. Gata6, Tbxt, MyoD), and regions that gained

in accessibility were enriched in pluripotency factors (e.g. Brn1, Oct4) (**Fig. 5C**) (Afouda

et al., 2005; Hopwood et al., 1989; Jerabek et al., 2014; Mistri et al., 2015; Smith et al.,

1991). Together, these data support a function for Foxi1 in regulating accessible

chromatin state during ectodermal development in *Xenopus*.

Finally, we wondered how loss of Foxi1 affects chromatin accessibility in regions

harboring important genes for mucociliary epidermis development. First, we inspected a

region around the *krt12.4* (epidermal keratin) gene on chromosome 9/10.S, which

revealed strongly reduced accessibility and indicated a loss of epidermal competence

(**Fig. 5D**) (Wills et al., 2010). Next, we inspected genomic loci containing genes

associated with MPPs (*dll1.L*), ISCs (*ubp1.L* and *dmrt2.S*), and mucociliary

development (*foxj1.L* and *tp63.L*) (Deblandre et al., 1999; Haas et al., 2019; Quigley

and Kintner, 2017; Quigley et al., 2011). In all cases, we found reduced accessibility

(**Fig. 5D, Fig. S6A-D**). Together, these data support a broad function for Foxi1 in

regulating ectodermal and mucociliary developmental potential.

In conclusion, Foxi1 regulates chromatin accessibility required for ectoderm

development and mucociliary epidermis patterning. This provides an additional

explanation as to how Foxi1 acts as an ectodermal determinant in early *Xenopus*

development.

Discussion

This work revealed that Foxi1 regulates multiple crucial steps during *Xenopus*

mucociliary epidermis development through transcriptional and epigenetic mechanisms.

Initially, low-level *foxi1* expression is activated in the prospective ectoderm by

maternally deposited *foxi2* (Cha et al., 2012). This low-level expression is required to

regulate chromatin accessibility for pro-ectodermal transcription factors (e.g. Tfap2a/c),

mucociliary regulators (e.g. Tp63) as well as mediators of thyroid hormone, retinoic acid

and TGFβ signaling (e.g. Thrb, Rar-a, Smad4) that were described to regulate

ectodermal development (Cibois et al., 2015; Edri et al., 2023; Haas et al., 2019;

Hoffman et al., 2007; Tasca et al., 2021). Since baseline *foxi1* expression is Foxi2- but

not Notch-dependent, Notch over-activation does not alter ectodermal identity in line

with previous reports (Cha et al., 2012; Deblandre et al., 1999).

Furthermore, our data support previous findings that loss of Foxi1 leads to acquisition of

mesendodermal fates, as loci enriched for pro-mesendodermal transcription factors

(e.g. Gata6, Tbxt, MyoD) remain accessible in the absence of Foxi1 (Afouda et al.,

2005; Hopwood et al., 1989; Smith et al., 1991). It is attractive to speculate that this is

possible, because multiple transcription factors enriched in the maintained fraction of

peaks (e.g. Gata and Sox family members) are known factors with pioneer activity (Hou

et al., 2017; Tremblay et al., 2018), which could compensate for the loss of Foxi1.

After inducing ectodermal identity at low levels, Foxi1 then further increases its own

expression through auto-regulation in mucociliary multipotent progenitors (MPPs),

inducing Notch ligand expression (*dll1* and *dlc*) required for Notch-mediated mucociliary

cell fate decisions into ISCs, MCCs, secretory cells and basal stem cells (Brislinger-

Engelhardt et al., 2023). MPPs not exposed to elevated Notch levels further increase

foxi1 expression through auto-regulation, and high levels of Foxi1 induce ISC fate, in

line with the known role of Foxi1 as master transcription factor for ionocytes across

vertebrate tissues (Pou Casellas et al., 2023).

ISC subtype selection and differentiation is a multi-step process. In the *Xenopus*

epidermis, Ubp1 is expressed first in both ISC subtypes, which terminates Notch ligand

expression in differentiating ISCs. Ubp1 drives differentiation of β-ISCs, while Dmrt2

drives α-ISC differentiation. Interestingly, Dmrt2 has recently been shown to be required

for α-ISCs in the mouse kidney, while not Ubp1, but the related grainyhead-like

transcription factor Tfcp2l1 is employed in mammalian kidney β-ISCs (Quigley et al.,

2011; Werth et al., 2017; Wu et al., 2024). This could explain why Dll1 expression is

terminated in *Xenopus* epidermal ISCs, but remains active in mammalian kidney ISCs

(also called INCs) (Mukherjee et al., 2020; Werth et al., 2017).

Similarly to Ubp1, our results suggest that master transcription factors for other

mucociliary epidermal cell types also terminate Dll1/Dlc ligand expression in MPPs.

Together, this system provides a robust Notch feedback-regulated developmental

program for mucociliary epidermis development, with Foxi1 as a central player that acts

through transcriptional and epigenetic mechanisms.

Finally, our finding that Foxi1 drives an MPP state during mucociliary epidermis

development could serve as a starting point to better understand the role of Foxi1 in

certain cancers, e.g. chromophore renal cell carcinoma (chRCC) and in pulmonary large

cell carcinoma (LCC), both highly associated with Foxi1 activity (Lindgren et al., 2017;

Simbolo et al., 2024; Skala et al., 2020; Yamada et al., 2022).

References

Afouda, B.A., A. Ciau-Uitz, and R. Patient. 2005. GATA4, 5 and 6 mediate TGFbeta maintenance of
endodermal gene expression in Xenopus embryos. *Development*. 132:763-774.

Aztekin, C., T.W. Hiscock, J.C. Marioni, J.B. Gurdon, B.D. Simons, and J. Jullien. 2019. Identification of a
regeneration-organizing cell in the Xenopus tail. *Science*. 364:653-658.

Blomqvist, S.R., H. Vidarsson, S. Fitzgerald, B.R. Johansson, A. Ollerstam, R. Brown, A.E. Persson, G.G.
Bergstrom, and S. Enerback. 2004. Distal renal tubular acidosis in mice that lack the forkhead
transcription factor Foxi1. *J Clin Invest*. 113:1560-1570.

Blomqvist, S.R., H. Vidarsson, O. Soder, and S. Enerback. 2006. Epididymal expression of the forkhead
transcription factor Foxi1 is required for male fertility. *EMBO J*. 25:4131-4141.

Briggs, J.A., C. Weinreb, D.E. Wagner, S. Megason, L. Peshkin, M.W. Kirschner, and A.M. Klein. 2018. The
dynamics of gene expression in vertebrate embryogenesis at single-cell resolution. *Science*. 360.

Brislinger-Engelhardt, M.M., F. Lorenz, M. Haas, S. Bowden, A. Tasca, C. Kreutz, and P. Walentek. 2023.
Temporal Notch signaling regulates mucociliary cell fates through Hes-mediated competitive de-
repression. *bioRxiv*.

Brooks, E.R., and J.B. Wallingford. 2014. Multiciliated cells. *Curr Biol*. 24:R973-982.

Buenrostro, J.D., P.G. Giresi, L.C. Zaba, H.Y. Chang, and W.J. Greenleaf. 2013. Transposition of native
chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding proteins
and nucleosome position. *Nat Methods*. 10:1213-1218.

Cha, S.W., M. McAdams, J. Kormish, C. Wylie, and M. Kofron. 2012. Foxi2 is an animally localized
maternal mRNA in Xenopus, and an activator of the zygotic ectoderm activator Foxi1e. *PLoS*
One. 7:e41782.

Cibois, M., G. Luxardi, B. Chevalier, V. Thome, O. Mercey, L.E. Zaragosi, P. Barbry, A. Pasini, B. Marcet,
and L. Kodjabachian. 2015. BMP signalling controls the construction of vertebrate mucociliary
epithelia. *Development*. 142:2352-2363.

Community, T.G. 2024. The Galaxy platform for accessible, reproducible, and collaborative data
analyses: 2024 update. *Nucleic Acids Research*. 52:W83-W94.

Danecek, P., J.K. Bonfield, J. Liddle, J. Marshall, V. Ohan, M.O. Pollard, A. Whitwham, T. Keane, S.A.
McCarthy, R.M. Davies, and H. Li. 2021. Twelve years of SAMtools and BCFtools. *GigaScience*. 10.

Deblandre, G.A., D.A. Wettstein, N. Koyano-Nakagawa, and C. Kintner. 1999. A two-step mechanism
generates the spacing pattern of the ciliated cells in the skin of Xenopus embryos. *Development*.
126:4715-4728.
Dubaissi, E., and N. Papalopulu. 2011. Embryonic frog epidermis: a model for the study of cell-cell
interactions in the development of mucociliary disease. *Dis Model Mech*. 4:179-192.
Dubaissi, E., K. Rousseau, R. Lea, X. Soto, S. Nardeosingh, A. Schweickert, E. Amaya, D.J. Thornton, and N.
Papalopulu. 2014. A secretory cell type develops alongside multiciliated cells, ionocytes and
goblet cells, and provides a protective, anti-infective function in the frog embryonic mucociliary
epidermis. *Development*. 141:1514-1525.
Edri, T., D. Cohen, Y. Shabtai, and A. Fainsod. 2023. Alcohol induces neural tube defects by reducing
retinoic acid signaling and promoting neural plate expansion. *Front Cell Dev Biol*. 11:1282273.
Esmaeili, M., S.A. Blythe, J.W. Tobias, K. Zhang, J. Yang, and P.S. Klein. 2020. Chromatin accessibility and
histone acetylation in the regulation of competence in early development. *Dev Biol*. 462:20-35.
Feng, J., T. Liu, B. Qin, Y. Zhang, and X.S. Liu. 2012. Identifying ChIP-seq enrichment using MACS. *Nat*
Protoc. 7:1728-1740.
Fisher, M., C. James-Zorn, V. Ponferrada, A.J. Bell, N. Sundararaj, E. Segerdell, P. Chaturvedi, N. Bayyari,
S. Chu, T. Pells, V. Lotay, S. Agalakov, D.Z. Wang, B.I. Arshinoff, S. Foley, K. Karimi, P.D. Vize, and
A.M. Zorn. 2023. Xenbase: key features and resources of the Xenopus model organism
knowledgebase. *Genetics*. 224.
Giudetti, G., M. Giannaccini, D. Biasci, S. Mariotti, A. Degl'innocenti, M. Perrotta, G. Barsacchi, and M.
Andreazzoli. 2014. Characterization of the Rx1-dependent transcriptome during early retinal
development. *Dev Dyn*. 243:1352-1361.
Haas, M., J.L. Gomez Vazquez, D.I. Sun, H.T. Tran, M. Brislinger, A. Tasca, O. Shomroni, K. Vleminckx, and
P. Walentek. 2019. DeltaN-Tp63 Mediates Wnt/beta-Catenin-Induced Inhibition of
Differentiation in Basal Stem Cells of Mucociliary Epithelia. *Cell Rep*. 28:3338-3352 e3336.
Harland, R.M. 1991. In situ hybridization: an improved whole-mount method for Xenopus embryos.
Methods Cell Biol. 36:685-695.
Hayes, J.M., S.K. Kim, P.B. Abitua, T.J. Park, E.R. Herrington, A. Kitayama, M.W. Grow, N. Ueno, and J.B.
Wallingford. 2007. Identification of novel ciliogenesis factors using a new in vivo model for
mucociliary epithelial development. *Dev Biol*. 312:115-130.
Heinz, S., C. Benner, N. Spann, E. Bertolino, Y.C. Lin, P. Laslo, J.X. Cheng, C. Murre, H. Singh, and C.K.
Glass. 2010. Simple combinations of lineage-determining transcription factors prime cis-
regulatory elements required for macrophage and B cell identities. *Mol Cell*. 38:576-589.
Hoffman, T.L., A.L. Javier, S.A. Campeau, R.D. Knight, and T.F. Schilling. 2007. Tfap2 transcription factors
in zebrafish neural crest development and ectodermal evolution. *J Exp Zool B Mol Dev Evol*.
308:679-691.
Hopwood, N.D., A. Pluck, and J.B. Gurdon. 1989. MyoD expression in the forming somites is an early
response to mesoderm induction in Xenopus embryos. *Embo j*. 8:3409-3417.
Hou, L., Y. Srivastava, and R. Jauch. 2017. Molecular basis for the genome engagement by Sox proteins.
Semin Cell Dev Biol. 63:2-12.
Hulander, M., A.E. Kiernan, S.R. Blomqvist, P. Carlsson, E.J. Samuelsson, B.R. Johansson, K.P. Steel, and S.
Enerback. 2003. Lack of pendrin expression leads to deafness and expansion of the
endolymphatic compartment in inner ears of Foxi1 null mutant mice. *Development*. 130:2013-
2025.
Janicke, M., T.J. Carney, and M. Hammerschmidt. 2007. Foxi3 transcription factors and Notch signaling
control the formation of skin ionocytes from epidermal precursors of the zebrafish embryo. *Dev*
Biol. 307:258-271.

Jerabek, S., F. Merino, H.R. Schöler, and V. Cojocaru. 2014. OCT4: dynamic DNA binding pioneers stem
cell pluripotency. *Biochim Biophys Acta*. 1839:138-154.
Lindgren, D., P. Eriksson, K. Krawczyk, H. Nilsson, J. Hansson, S. Veerla, J. Sjolund, M. Hoglund, M.E.
Johansson, and H. Axelson. 2017. Cell-Type-Specific Gene Programs of the Normal Human
Nephron Define Kidney Cancer Subtypes. *Cell Rep*. 20:1476-1489.
Luo, T., Y. Zhang, D. Khadka, J. Rangarajan, K.W. Cho, and T.D. Sargent. 2005. Regulatory targets for
transcription factor AP2 in Xenopus embryos. *Dev Growth Differ*. 47:403-413.
Ma, L., J.C. Hocking, C.L. Hehr, C. Schuurmans, and S. McFarlane. 2007. Zac1 promotes a Müller glial cell
fate and interferes with retinal ganglion cell differentiation in Xenopus retina. *Dev Dyn*. 236:192-
202.
Maharana, S.K., and G. Schlosser. 2018. A gene regulatory network underlying the formation of pre-
placodal ectoderm in Xenopus laevis. *BMC Biol*. 16:79.
Mir, A., M. Kofron, A.M. Zorn, M. Bajzer, M. Haque, J. Heasman, and C.C. Wylie. 2007. Foxl1e activates
ectoderm formation and controls cell position in the Xenopus blastula. *Development*. 134:779-
788.
Mistri, T.K., A.G. Devasia, L.T. Chu, W.P. Ng, F. Halbritter, D. Colby, B. Martynoga, S.R. Tomlinson, I.
Chambers, P. Robson, and T. Wohland. 2015. Selective influence of Sox2 on POU transcription
factor binding in embryonic and neural stem cells. *EMBO Rep*. 16:1177-1191.
Montoro, D.T., A.L. Haber, M. Biton, V. Vinarsky, B. Lin, S.E. Birket, F. Yuan, S. Chen, H.M. Leung, J.
Villoria, N. Rogel, G. Burgin, A.M. Tsankov, A. Waghray, M. Slyper, J. Waldman, L. Nguyen, D.
Dionne, O. Rozenblatt-Rosen, P.R. Tata, H. Mou, M. Shivaraju, H. Bihler, M. Mense, G.J. Tearney,
S.M. Rowe, J.F. Engelhardt, A. Regev, and J. Rajagopal. 2018. A revised airway epithelial
hierarchy includes CFTR-expressing ionocytes. *Nature*. 560:319-324.
Mukherjee, M., J. DeRiso, M. Janga, E. Fogarty, and K. Surendran. 2020. Foxi1 inactivation rescues loss of
principal cell fate selection in Hes1-deficient kidneys but does not ensure maintenance of
principal cell gene expression. *Dev Biol*. 466:1-11.
Plasschaert, L.W., R. Zilionis, R. Choo-Wing, V. Savova, J. Knehr, G. Roma, A.M. Klein, and A.B. Jaffe.
2018. A single-cell atlas of the airway epithelium reveals the CFTR-rich pulmonary ionocyte.
Nature. 560:377-381.
Pou Casellas, C., C. Pleguezuelos-Manzano, M.B. Rookmaaker, M.C. Verhaar, and H. Clevers. 2023.
Transcriptomic profile comparison reveals conservation of ionocytes across multiple organs. *Sci
Rep*. 13:3516.
Quigley, I.K., and C. Kintner. 2017. Rfx2 Stabilizes Foxj1 Binding at Chromatin Loops to Enable
Multiciliated Cell Gene Expression. *PLoS Genet*. 13:e1006538.
Quigley, I.K., J.L. Stubbs, and C. Kintner. 2011. Specification of ion transport cells in the Xenopus larval
skin. *Development*. 138:705-714.
Ramírez, F., D.P. Ryan, B. Grüning, V. Bhardwaj, F. Kilpert, A.S. Richter, S. Heyne, F. Dündar, and T.
Manke. 2016. deepTools2: a next generation web server for deep-sequencing data analysis.
Nucleic Acids Research. 44:W160-W165.
Ray, H., and C. Chang. 2020. The transcription factor Hypermethylated in Cancer 1 (Hic1) regulates
neural crest migration via interaction with Wnt signaling. *Dev Biol*. 463:169-181.
Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S.
Saalfeld, B. Schmid, J.Y. Tinevez, D.J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, and A.
Cardona. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods*. 9:676-
682.
Schweickert, A., H. Steinbeisser, and M. Blum. 2001. Differential gene expression of Xenopus Pitx1,
Pitx2b and Pitx2c during cement gland, stomodeum and pituitary development. *Mech Dev*.
107:191-194.

Scudieri, P., I. Musante, A. Venturini, D. Guidone, M. Genovese, F. Cresta, E. Caci, A. Palleschi, M. Poeta,
F. Santamaria, F. Ciciriello, V. Lucidi, and L.J.V. Galietta. 2020. Ionocytes and CFTR Chloride
Channel Expression in Normal and Cystic Fibrosis Nasal and Bronchial Epithelial Cells. *Cells*. 9.
Simbolo, M., G. Centonze, A. Gkountakos, V. Monti, P. Maisonneuve, S. Golovco, G. Sabella, A. Del
Gobbo, S. Gobbo, S. Ferrero, A. Fabbri, C. Pardo, G. Garzone, N. Prinzi, S. Pusceddu, A. Testi, L.
Rolli, A. Mangogna, L. Bercich, M.R. Benvenuti, E. Bria, S. Pilotto, A. Berruti, U. Pastorino, C.
Capella, M. Infante, M. Milella, A. Scarpa, and M. Milione. 2024. Characterization of two
transcriptomic subtypes of marker-null large cell carcinoma of the lung suggests different origin
and potential new therapeutic perspectives. *Virchows Arch*. 484:777-788.
Sive, H.L., R.M. Grainger, and R.M. Harland. 2007a. Animal Cap Isolation from Xenopus laevis. *CSH
Protoc*. 2007:pdb.prot4744.
Sive, H.L., R.M. Grainger, and R.M. Harland. 2007b. Xenopus laevis In Vitro Fertilization and Natural
Mating Methods. *CSH Protoc*. 2007:pdb.prot4737.
Sive, H.L., R.M. Grainger, and R.M. Harland. 2010. Microinjection of Xenopus embryos. *Cold Spring Harb
Protoc*. 2010:pdb.ip81.
Skala, S.L., X. Wang, Y. Zhang, R. Mannan, L. Wang, S.P. Narayanan, P. Vats, F. Su, J. Chen, X. Cao, J.
Siddiqui, P. Argani, M.P. Cieslik, T.J. Giordano, A.M. Chinnaiyan, S.M. Dhanasekaran, and R.
Mehra. 2020. Next-generation RNA Sequencing-based Biomarker Characterization of
Chromophobe Renal Cell Carcinoma and Related Oncocytic Neoplasms. *Eur Urol*. 78:63-74.
Smith, J.C., B.M. Price, J.B. Green, D. Weigel, and B.G. Herrmann. 1991. Expression of a Xenopus
homolog of Brachyury (T) is an immediate-early response to mesoderm induction. *Cell*. 67:79-
87.
Stubbs, J.L., I. Oishi, J.C. Izpisúa Belmonte, and C. Kintner. 2008. The forkhead protein Foxj1 specifies
node-like cilia in Xenopus and zebrafish embryos. *Nat Genet*. 40:1454-1460.
Stubbs, J.L., E.K. Vladar, J.D. Axelrod, and C. Kintner. 2012. Multicilin promotes centriole assembly and
ciliogenesis during multiciliate cell differentiation. *Nat Cell Biol*. 14:140-147.
Suri, C., T. Haremaki, and D.C. Weinstein. 2005. Xema, a foxi-class gene expressed in the gastrula stage
Xenopus ectoderm, is required for the suppression of mesendoderm. *Development*. 132:2733-
2742.
Tasca, A., M. Helmstadter, M.M. Brislinger, M. Haas, B. Mitchell, and P. Walentek. 2021. Notch signaling
induces either apoptosis or cell fate change in multiciliated cells during mucociliary tissue
remodeling. *Dev Cell*. 56:525-539 e526.
Tremblay, M., O. Sanchez-Ferras, and M. Bouchard. 2018. GATA transcription factors in development
and disease. *Development*. 145.
Walentek, P. 2018. Manipulating and Analyzing Cell Type Composition of the Xenopus Mucociliary
Epidermis. *Methods Mol Biol*. 1865:251-263.
Walentek, P. 2022. Signaling Control of Mucociliary Epithelia: Stem Cells, Cell Fates, and the Plasticity of
Cell Identity in Development and Disease. *Cells Tissues Organs*. 211:736-753.
Walentek, P., T. Beyer, C. Hagenlocher, C. Muller, K. Feistel, A. Schweickert, R.M. Harland, and M. Blum.
2015. ATP4a is required for development and function of the Xenopus mucociliary epidermis - a
potential model to study proton pump inhibitor-associated pneumonia. *Dev Biol*. 408:292-304.
Walentek, P., T. Beyer, T. Thumberger, A. Schweickert, and M. Blum. 2012. ATP4a is required for Wnt-
dependent Foxj1 expression and leftward flow in Xenopus left-right development. *Cell Rep*.
1:516-527.
Walentek, P., S. Bogusch, T. Thumberger, P. Vick, E. Dubaissi, T. Beyer, M. Blum, and A. Schweickert.
2014. A novel serotonin-secreting cell type regulates ciliary motility in the mucociliary epidermis
of Xenopus tadpoles. *Development*. 141:1526-1533.

Walentek, P., and I.K. Quigley. 2017. What we can learn from a tadpole about ciliopathies and airway
diseases: Using systems biology in Xenopus to study cilia and mucociliary epithelia. *Genesis*. 55.
Werth, M., K.M. Schmidt-Ott, T. Leete, A. Qiu, C. Hinze, M. Viltard, N. Paragas, C.J. Shawber, W. Yu, P.
Lee, X. Chen, A. Sarkar, W. Mu, A. Rittenberg, C.S. Lin, J. Kitajewski, Q. Al-Awqati, and J. Barasch.
2017. Transcription factor TFCP2L1 patterns cells in the mouse kidney collecting ducts. *Elife*. 6.
Wills, A.E., V.M. Choi, M.J. Bennett, M.K. Khokha, and R.M. Harland. 2010. BMP antagonists and FGF
signaling contribute to different domains of the neural plate in Xenopus. *Dev Biol*. 337:335-350.
Wu, S.T., Y. Feng, R. Song, Y. Qi, L. Li, D. Lu, Y. Wang, W. Wu, A. Morgan, X. Wang, Y. Xia, R. Liu, S.I.
Alexander, J. Wong, Y. Zhang, and X. Zheng. 2024. Foxp1 Is Required for Renal Intercalated Cell
Differentiation and Acid-Base Regulation. *J Am Soc Nephrol*. 35:533-548.
Yamada, Y., D. Belharazem-Vitacolonnna, H. Bohnenberger, C. Weiss, N. Matsui, M. Kriegsmann, K.
Kriegsmann, P. Sinn, K. Simon-Keller, G. Hamilton, T. Graeter, G. Preissler, G. Ott, S. Scholch, N.
Nakajima, A. Yoshizawa, H. Haga, H. Date, R.K. Thomas, I. Petrini, G. Giaccone, P. Strobel, and A.
Marx. 2022. Pulmonary cancers across different histotypes share hybrid tuft cell/ionocyte-like
molecular features and potentially druggable vulnerabilities. *Cell Death Dis*. 13:979.
Yan, J., L. Xu, G. Crawford, Z. Wang, and S.M. Burgess. 2006. The forkhead transcription factor Foxl1
remains bound to condensed mitotic chromosomes and stably remodels chromatin structure.
Mol Cell Biol. 26:155-168.
Yang, T., H. Vidarsson, S. Rodrigo-Blomqvist, S.S. Rosengren, S. Enerback, and R.J. Smith. 2007.
Transcriptional control of SLC26A4 is involved in Pendred syndrome and nonsyndromic
enlargement of vestibular aqueduct (DFNB4). *Am J Hum Genet*. 80:1055-1063.
Zhang, Y., T. Luo, and T.D. Sargent. 2006. Expression of TFAP2beta and TFAP2gamma genes in Xenopus
laevis. *Gene Expr Patterns*. 6:589-595.

Acknowledgments:

We thank: S. Schefold and A. Andricek for expert technical help; L. Kodjabachian and
team, T. Manke and W. Deboutte, T. Kwon for support and discussions; Xenbase,
EXRC for Xenopus resources; Light Imaging Center Freiburg, BiMiC and Aqua Core for
microscope/animal resources; B. Grüning and the Freiburg Galaxy Team for
bioinformatics platform and support. This study was supported by the Deutsche
Forschungsgemeinschaft (DFG) under the Emmy Noether and Heisenberg
Programmes (grant WA3365/2-1 and WA3365/5-1), by DFG SFB1453 NephGen
(Project ID 431984000), by DFG/ANR grant WA3365/4-1, and by the NHLBI through a
Pathway to Independence Award (K99HL127275) to PW; and under Germany's
Excellence Strategy (CIBSS – EXC-2189 – Project ID 390939984) to PW and CK.

Author contribution:

SB: epigenetics; MMBE: cell fates and Notch; MOH: reporter studies; ATP, DW, SH,
PW: experimental support; SB, MMBE, MOH, PW: experimental design, planning,
analysis and interpretation of data; FL, TL, CK: crucial discussion and mathematical
modeling; PW, SB: bioinformatics. PW: study design and supervision, coordinating

collaborative work, manuscript preparation with input from all authors. SB, MMBE, MOH
contributed equally and can list themselves as first co-first authors.

Material and Mathods:

<u>Animal experiments:</u>
Wild-type *Xenopus laevis* were obtained from the European Xenopus Resource Centre
(EXRC) at University of Portsmouth, School of Biological Sciences, UK, or Xenopus 1,
USA. Frog maintenance and care was conducted according to standard procedures in
the AquaCore facility, University Freiburg, Medical Center (RI_00544) and based on
recommendations provided by the international Xenopus community resource centers
NXR (RRID:SCR_013731) and EXRC as well as by Xenbase (http://www.xenbase.org/,
RRID:SCR_003280)(Fisher et al., 2023). This work was done in compliance with
German animal protection laws and was approved under Registrier-Nr. G-18/76 and G-
22/43 by the state of Baden-Württemberg.
609
<u>Data availability:</u>
NGS datasets are available via NCBI GEO, ATAC-seq datasets (# pending), mRNA-seq
datasets were generated in previous studies (GSE130448, GSE215373, GSE215419,
GSE262944)(Brislinger-Engelhardt et al., 2023; Haas et al., 2019). Imaging and
quantification data are available to the scientific community upon request to
peter.walentek@medizin.uni-freiburg.de.

<u>Manipulation of *Xenopus* Embryos:</u>
X. laevis eggs were collected and in vitro-fertilized, then cultured and microinjected by
standard procedures (Sive et al., 2007b; Sive et al., 2010). Embryos were injected with
Morpholino oligonucleotides (MOs, Gene Tools), mRNAs or plasmid DNA at two-cell to
eight-cell stage using a PicoSpritzer setup in 1/3x Modified Frog Ringer's solution (MR)

with 2.5% Ficoll PM 400 (GE Healthcare, #17-0300-50), and were transferred after
injection into 1/3x MR containing Gentamycin. Drop size was calibrated to about 7–8nL
per injection.
625
626   Embryos injected with hormone-inducible constructs of (GFP-ΔN-tp63-GR and MCI-GR)
627   (Haas et al., 2019; Stubbs et al., 2012) were treated with 10μM Dexamethasone
628   (Sigma-Aldrich/Merck #D4902) in ethanol from eight-cell stage until fixation. Ultrapure
629   Ethanol (NeoFroxx #LC-8657.3) was used as vehicle control.
630
631   Morpholino oligonucleotides (MOs) were obtained from Gene Tools targeting *dll1*,
632   *dmrt2*, *foxi1*, *notch1* and *ubp1* and were used at doses as indicated in the list below.
633
mRNAs encoding *nicd (*100 ng/μl*)* (Deblandre et al., 1999), *foxi1 (25-100*ng/μl*)* (this
study using primers listed below into pCS107), *mcidas* (100ng/μl) (Stubbs et al., 2012),
foxj1 (100ng/μl) (Stubbs et al., 2008), *foxa1* (100ng/μl) (Dubaissi et al., 2014; Walentek
et al., 2014), *ΔN-tp63* (100ng/μl) (Haas et al., 2019) were injected together with
membrane-gfp or *membrane-rfp* (at 50ng/μL) or *h2b-rfp* (at 30ng/μL) as lineage tracers.
All mRNAs were prepared using the mMessage Machine kit using Sp6 (Invitrogen
#AM1340) supplemented with RNAse Inhibitor (Promega #N251B).
641
642   The *foxi1::gfp-utrophin*, *foxi1Δ1::gfp-utrophin* and *a-tub::mscarletI* plasmids were
643   purified using the Pure Yield midiprep kit (Promega #A2492) and injected at 5 ng/μl.
644
645   Morpholino Sequences and doses:

Name	Sequence	Concentration Range
dll1.L MO	5'-CCCATGTTGTCTGATATGCGATTG-3'	0.5-2 pmol
dll1.S MO	5'-AGGCACTGCTGTCCCATGTTG-3'	0.5-2 pmol
dmrt2 MO	5'-GTGCCTTCATCTCGTACATCTCCAG-3'	1.5 pmol
foxi1MO	5'-GTGCTTGTGGATCAAATGCACTCAT-3'	1.5-3 pmol
notch1 MO	5'-GCACAGCCAGCCCTATCCGATCCAT-3'	4 pmol
ubp1 MO	5'-TTGGGTCGGACAGGTACAATAATCC-3'	2-3 pmol

646
647   Full length *foxi1* cloning (3'- 5'):

Name	Sequence
Cla1-XLfoxi1-F	AAAAAAATCGATATGAGTGCATTTGATCCACAAGC
XLfoxi1-Sal1-R	AAAGTCGACTTATACTTCTGTACCTTCTCTG

648
649
foxi1.S reporter construct cloning and experiments:
To generate the *foxi1.S::gfp-utrophin* reporter construct, genomic DNA was prepared
from *X. laevis* using the phenol/chloroform DNA purification (ThermoFisher #15593031
and associated protocol). A 2.7 kb fragment (Fig.S2D) of the *foxi1.S* promoter was
cloned using Easy-A Hi-Fi Cloning Enzyme (Agilent #600404) and primers listed in the
table below. The PCR fragment was ligated using the pGEM-T Easy Vector System

(Promega #A1360). The *foxi1.S* promoter sequence was subcloned into *a-tub::gfp-*
utrophin (used in (Tasca et al., 2021)) after removal of the *a-tub* promoter sequence
using HiFi DNA Assembly (NEB #E2621S) and Q5 High-Fidelity DNA Polymerase (NEB
#M0491S) kits. *foxi1Δ1::gfp-utrophin* reporter version (Fig. S2C,D) was generated
using Q5 High-Fidelity DNA Polymerase and primers listed in the table below. The *a-*
tub::mscarletI reporter was generated by replacing the *gfp-utrophin* sequence in *a-*
tub::gfp-utrophin by the *mscarletI* sequence using HiFi DNA Assembly and Q5 High-
Fidelity DNA Polymerase and primers listed below. Final construct sequences were
analyzed by whole-plasmid nanopore sequencing.
665
666 Cloning primers *foxi1.S* reporter (3'- 5'):

Name	Sequence
gDNA-Prom_foxi1_F	GCATAATGAATCCCAAGTGTACTG
gDNA-Prom_foxi1_R	GAAGCAATCGTTTAGAGACAGG
Foxi1prom_F	CGCTATTACGCCAGTCGACCGCATAATGAATCCCAAGTG
Foxi1prom_R	CAATTCGAATCGATGGGATCAGTTAAAGCTAGCAGGTC
MS2_rmATUB_F	GATCCCATCGATTCGAATTG
MS2_rmATUB_R	GGTCGACTGGCGTAATAG
Q5foxi1Del1_F	TCTGTAGCTGATGTCTATAATC
Q5foxi1Del1_R	TACCACTGTGTGACTCAG
MS2_rmGFPUtro_F	GTACAAGTAACCTCTAGAACTATAGTGAGTC
MS2_rmGFPUtro_R	TGCTCACCATGGTTTGGATCAATTCGAATC
mScarletI_F	GATCCAAACCATGGTGAGCAAGGGCGAG
mScarletI_R	GTTCTAGAGGTTACTTGTACAGCTCGTCCATG

667
Whole mount *in situ* hybridization and sections:
For antisense *in situ* hybridization probes*, slc26a4*, *slc4a1*, *ubp1* and *dmrt2* fragments
were cloned from whole-embryo cDNAs derived from stages between 3 and 30 using
primers listed below (ISH-primers). All sequences were verified by Sanger sequencing.
In addition, the following, previously published probes were used: *foxi1* (Quigley et al.,
2011), *foxj1* and *mcidas* (Stubbs et al., 2008; Stubbs et al., 2012), *foxa1* (Walentek et
al., 2014), *tp63* (Haas et al., 2019), *atp6v1e1* (Walentek et al., 2015) and *dll1* (Tasca et
al., 2021).
676
Embryos were fixed in MEMFA (100mM MOPS pH7.4, 2mM EGTA, 1mM MgSO4, 3.7%
(v/v) Formaldehyde) overnight at 4°C and stored in 100% Ethanol at -20°C until used.
DNAs were purified using the PureYield Midiprep kit (Promega #A2492) and were
linearized before in vitro synthesis of anti-sense RNA probes using T7 or Sp6
polymerase (Promega, #P2077 and #P108G), RNAse inhibitor and dig-labeled rNTPs
(Roche, #3359247910 and 11277057001). Embryos were in situ hybridized according to
(Harland, 1991), bleached after staining with BM Purple (Roche #11442074001) and
imaged. Sections were made after embedding in gelatin-albumin with Glutaraldehyde at
50-70μm as described in (Walentek et al., 2012).
686
687 Probe cloning primers (5'-3'):

Name	Sequence

ISH-dmrt2-F	CAAACCAGTGTATCAGAGAC
ISH-dmrt2-R	ACTCCTTTCCTAAGAAGCAG
ISH-ubp1-F	ATTCCTGAAGCAAGAAGACC
ISH-ubp1-R	GAGAATGTGAATCCCATGAG
ISH-slc26a4-F	GATTCATACCACCTATGACAC
ISH-slc26a4-R	TTCCAATAGTTCCCTAGATTCC
ISH-slc4a1-F	ATCTCCTATCTCACCTTTCAC
ISH-slc4a1-R	ATCCATCTGTCTGTCTTCTC
ISH-anti-GFP-F	ATGGTGAGCAAGGGCGAGG
ISH-anti-GFP-R	CTTGTACAGCTCGTCCATGCCATGCCGAGAGTG

688
689
Evaluation of WMISH staining and morphological evaluations:
Embryos were staged according to Nieuwkoop and Faber (1994) Normal Table of
Xenopus laevis (Daudin). Garland Publishing Inc, New York ISBN 0-8153-1896-0. For
the *foxi1* expression stage series wt embryos from multiple batches were mixed and at
least 5 embryos per stage were assessed.
Images of embryos after *in situ* hybridization and corresponding sections were imaged
using a Zeiss AxioZoom setup, Zeiss AxioImager.Z1 or Zeiss Stemi508 with
Axiocam208-color, and images were adjusted for color balance, brightness and contrast
using Adobe Photoshop.
699
In Fig.1B,C expression strength were categorized in normal, reduced, strongly reduced
or increased. In Fig. 1D-G, 3C, S5B, expression level differences observed between the
uninjected control sides and manipulated sides of embryos were scored in whole mount
embryos, while depicted sections are shown for clarity. In Fig. 3A and S4C induction of
expression was scored. In Fig. 3B *dll1* expression in the ventral epidermis was analyzed
as normal or less (number of dots and expression intensity).
706
707 For analyses in Fig.2A and S2A, embryos injected with high dose of *foxi1* MO, cell
708 morphology and cell size were evaluated for Fig. S2A (and delamination was confirmed
709 in hemisected embryos) and skin lesions were evaluated for Fig. 2A.
710
Immunofluorescence staining and sample preparation:
Whole *Xenopus* embryos, were fixed at indicated stages in 4% paraformaldehyde at
4°C overnight or 2h at room temperature, then washed 3x 15min with PBS, 2x 30min in
PBST (0.1% Triton X-100 in PBS), and were blocked in PBST-CAS (90% PBS
containing 0.1% Triton X-100, 10% CAS Blocking; ThermoFischer #00-8120) for 30min-
1h at RT. A detailed protocol was described in (Walentek, 2018).
717
Mouse anti-Acetylated-α-tubulin (Sigma/Merck #T6793) primary antibody (1:1000) was
used to mark cilia / MCCs, Rabbit Anti-serotonin (Sigma/Merck #S5545) primary
antibody (1:500) was used to mark SSCs applied at 4°C overnight. Secondary
antibodies AlexaFlour-405-labeled goat anti-mouse (Invitrogen # A30104) and
AlexaFlour 405-labeled goat anti-rabbit antibody (Invitrogen #A31556) were used for 2 h
at RT (1:250). Antibodies were applied in 100% CAS Blocking (ThermoFischer #00-

8120). Actin was stained by incubation (30-120 min at room temperature) with
AlexaFluor 405-labeled Phalloidin (1:800 in PBSt; Invitrogen #A30104), mucus-like
compounds were stained by incubation (overnight at 4°C) with AlexaFluor-594- or -647-
labeled or PNA (1:500-1000 in PBSt; Molecular Probes #L32459 and #L32460).
728
729
<u>Fluorescence imaging, image processing and analysis:</u>
Confocal imaging was conducted using either a Zeiss LSM880 or a Zeiss LSM980
microscope and Zeiss Zen software in the LIC and BiMiC imaging facilities. Confocal
images were adjusted for channel brightness/contrast, Z-stack projections were
generated and cell types were quantified based on their morphology using ImageJ
(Schindelin et al., 2012). For analyses in Fig. 3D and S5A a detailed protocol for
quantification of *Xenopus* epidermal cell type composition was published (Walentek,
2018).
738
For analysis and comparison of fluorescent reporter construct activity on confocal
micrographs (Fig. 4A) in ImageJ, z-projections were performed using the "sum-slices"
function. Induction of reporter expression in the endoderm (Fig. 4B) embryos were
imaged using a Zeiss AxioImager.Z1 with Axiocam208-color camera. Induction was
scored as positive when GFP fluorescence was detected in the vegetal half of the
gastrula embryo. In some controls, activity was observed in involuting or animally
positioned mesoderm, where maternal *foxi2* deposition occurs. Fluorescent intensities
were color-coded using the function "lookup tables -> fire" (8 bit) in ImageJ.
747
748   <u>RNA- and ATAC-sequencing on *Xenopus* mucociliary organoids and bioinformatics</u>
749   <u>analysis:</u>
750   Manipulations and bulk mRNA-seq used in this paper were generated and published
751   here: (Brislinger-Engelhardt et al., 2023; Haas et al., 2019). scRNA-seq datasets were
752   published here: (Aztekin et al., 2019; Briggs et al., 2018).
753
For Fig. 1A, data from (Brislinger-Engelhardt et al., 2023) were used, TPM values from
.L and .S allo-allels were added, and the resulting matrix was clustered using Z-values
per line and galaxy.eu (ggplot2_heatmap2/3.1.3.1+galaxy). For Fig. S1A-C, log2-fold
changes were calculated using galaxy.eu (DeSeq2/2.1.3+galaxy) and visualized using
(ggplot2_heatmap2/3.1.3.1+galaxy). For Fig. S1 D, the online tool associated with
(Aztekin et al., 2019) was used (marionilab.cruk.cam.ac.uk/XenopusRegeneration). For
Fig. S4B, the online tool associated with (Briggs et al., 2018) was used
(kleintools.hms.harvard.edu/tools/currentDatasetsList_xenopus_v2.html)   to   extract
lineage-enriched transcripts and the heatmap was generated using galaxy.eu
(ggplot2_heatmap2/3.1.3.1+galaxy).
764
For ATAC-seq sample generation, injected and control embryos were cultured until st.
8. Animal caps were dissected in 1x Modified Barth's solution (MBS) and transferred to
0.5x MBS + Gentamycin (Sive et al., 2007a). 2 organoids per condition and replicate
were collected in PBS and ATAC-seq was performed as described in (Buenrostro et al.,
2013; Esmaeili et al., 2020). In short: Embryos were injected bilaterally in the animal

hemisphere at the two-cell stage with 3pmol *foxi1* MO or remained uninjected, animal
caps were prepared at st. 8, and organoids were collected upon the appearance of the
dorsal lip in control embryos cultured in parallel to the organoids (st. 10). Organoids
were transferred from MBS plates into a 1.5 mL low-bind microcentrifuge tube
(Eppendorf #0030108051) containing 1 mL of ice-cold 1x PBS. Samples were spun at
500 g at 4 ºC in the centrifuge for five minutes before removing the PBS and repeating
the wash step with fresh ice-cold 1x PBS. 50 µL of ice-cold lysis buffer (10 mM Tris pH
7.4, 10 mM NaCl, 3 mM MgCl2, 0.1% (w/v) Igepal CA-630) and pipetted to break up
samples. Samples were centrifuged at 500 g for 10 min at 4 ºC and pellets were
resuspended in 25 µL TD Buffer, 2.5 µL TDE1 Enzyme and 22.5 µL Nuclease-Free
water (Illumina #20034198). Samples were pelleted to mix and incubated on a
ThermoMixer at 37 ºC, 700 rpm for 30 min. Following incubation, the samples were
cleaned with MinElute Reaction Cleanup Kit (Qiagen, #28204), following manufacturer
instructions and eluted into 11 µL Buffer EB.
784
Libraries were prepared in collaboration with the NIG, University Medical Center
Göttingen. Quality was assessed with the Agilent Fragment Analyzer and prepared with
the ATAC-seq Kit (Active Motif, #53150). Samples were sequenced in triplicate on an
Illumina NovaSeq6000 with 150 nucleotide paired-end reads, totaling 50 million reads
per sample.
Raw sequencing files were assessed for quality using FastQC (v0.11.9, Andrews, S.
FastQC A Quality Control tool for High Throughput Sequence Data.
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/), and adapter sequences
were removed with TrimGalore (v0.6.7, https://doi.org/10.5281/zenodo.7598955). Data
were aligned to the *X. laevis* genome assembly v9.2 using BWA-MEM (v0.7.17,
https://arxiv.org/abs/1303.3997). Mitochondrial reads were removed using Samtools
(v1.21) (Danecek et al., 2021), and peak calling was performed with the callpeak
function of MACS2 (v2.2.7.1) (Feng et al., 2012). Differential analysis was performed
with the bdgdiff function of MACS2 and Venn diagrams were generated with
VennDiagram v1.7.3 in R v4.4.1. Heatmaps showing the average ATAC-seq signal
were generated using deepTools (v3.5.4) (Ramírez et al., 2016). Peaks were annotated
for the nearest *X. laevis* gene and transcription factor binding motifs with Homer (v4.11)
(Heinz et al., 2010), Plant-specific transcription factors were manually excluded from the
lists of transcription factors.. Bioinformatic analyses were performed on the Galaxy /
Europe platform (usegalaxy.eu) (Community, 2024). ATAC-seq data generated for this
study was deposited at NCBI GEO under (#pending).

806
<u>Quantification and statistical evaluation:</u>
Stacked bar graphs were generated in Microsoft Excel. Heatmaps and Venn diagrams
were generated using the Galaxy Europe platform (usegalaxy.eu) and R.
Sample sizes for all experiments were chosen based on previous experience and used
embryos derived from at least two different females. No randomization or blinding was
applied.
813
814 <u>Use of shared controls:</u>

815 For some of the *in situ* and IF experiments shared controls were used in multiple
816 graphs; all experiments are listed in the supplemental experiment log.
836 **Figure legends:**
837

Figure 1: Foxi1, Ubp1 and Dmrt2 differentially regulate ionocyte development
(A) Temporal expression analysis of core ISC genes. Heatmap of line-normalized z-
scores of TPMs (transcripts per million reads) derived from mRNA-seq of *Xenopus*
mucociliary organoids over the course of development (st. 9 - 32). **(B-G)** Knockdown of
ISC transcription factors (*foxi1* MO, *ubp1* MO, *dmrt2* MO) and analysis of effects by
WMISH at st. 29 - 32 against *atpv1e1* and *foxi1* (pan-ISC markers), *ubp1* and
slc25a4/pendrin (β-ISC markets), and *dmrt2* and *slc4a1/ae1* (α-ISC markers).
Representative images **(B,D,E,G)** and quantification **(C, F, H)** of results. n = number of
embryos analyzed per condition. Rescues (co-injection of *foxi1* mRNA) depicted in
(B,C) were scored as normal (norm.), reduced (red.), strongly reduced (s.red.) and
increased (incr.). Color code is shown in **(B)**. In conditions depicted in **(D, E, G)**,
expression levels were scored as less, more or equal expression as compared to the
uninjected control side. Color code is shown in **(F)**.
851
852

Figure 2: Foxi1 acts in a concentration-dependent manner to specify multipotent
progenitors
(A) Immunofluorescence confocal micrographs (IF) from control (ctrl.) and *foxi1*
morphants (*foxi1* MO; low concentration) at st. 32 stained for Acetylated-α-tubulin (Ac.-
α-tub., cilia, grey), F-actin (Actin, cell borders and morphology, grey), and mucus (PNA,
magenta). Targeted cells were identified by membrane GFP expression (memGFP,
green). Location of insets is indicated by dashed yellow box in upper panels.

Quantification of cell type composition is depicted as pie-charts, goblet cells (blue), ISCs
(yellow), MCCs (green) and SSCs (red). n embryos (above chart) and n quantified cells
(in/left of chart). **(B)** Brightfield images and quantification of st. 30-32 embryos.
Uninjected controls (ctrl.), *foxi1* morphants (*foxi1* MO; high concentration) and
morphants co-injected with *foxi1* mRNA (rescue) are depicted. Skin lesions (dashed
yellow outline) were quantified. **(C,D,E)** Analysis of *foxi1::gfp-utrophin* reporter (green)
injected embryos by IF **(C,E)** and WMISH against *gfp* **(D)**. **(C)** IF for Acetylated-α-tubulin
(Ac.-α-tub., cilia, grey), F-actin (Actin, cell borders and morphology, grey), and serotonin
(SSCs, grey) at st. 32. Targeted cells were identified by nuclear RFP expression (H2B-
RFP, blue). Magnifications of intercalating GFP(+) cell types are shown in insets.
Location of insets is indicated by dashed yellow boxes in left panels. **(D)** Sections of
epidermal locations from embryos depicted in Fig. S3C show *gfp* expressing cells in the
epidermis at key stages of mucociliary development (st. 10, 16, 25, 32). **(E)** IF for
foxi1::gfp-utrophin reporter (green) and F-actin (Actin, cell borders and morphology,
magenta) at st. 10.5 - 32 on hemistected embryos. Targeted cells were identified by
membrane RFP expression (mRFP, grey). Additional stages and full images shown in
Fig. S4A.
877
878
Figure 3: Foxi1 induces and Ubp1 terminates Notch ligand expression
(A,B,C) Manipulation of mucociliary cell fate transcription factors *foxi1* and *ubp1*
(ISCs/MPPs), *mcidas* and *foxj1* (MCCs), *foxa1* (SSCs) and *ΔN-tp63* (basal cells) and
analysis of effects by WMISH at st. 9 **(A)**, st. 11 **(B)** and st. 16 **(C)** against *dll1* (Notch
ligand) and *foxi1* (MPP/ISC marker). **(A)** Representative images of control (ctrl.) and
manipulated embryos (animal views) after mRNA overexpression of transcription factors
to test premature induction of *dll1*. Quantification of results and effects on *dlc* are shown
in Fig. S4C. Embryos were scored as induced or non-induced expression. **(B)**
Representative images of control (ctrl.) and *foxi1* morphants (*foxi1* MO) (ventral views)
to test effects on *dll1* expression. Quantification of results is shown in lower panel.
Locations of insets are indicated by dashed yellow box in upper panels. Embryos were
scored as normal or reduced (less) expression of *dll1*. **(C)** Representative images of
section embryos after unilateral knockdown of *ubp1* (*ubp1* MO). Expression of markers
was scored as more, less or equal to uninjected control (ctrl.) side. Locations of insets
are indicated by dashed yellow box in left panels. **(D)** IF of control (ctrl.) and *ubp1*
morphants (*ubp1* MO) at st. 32 stained for Acetylated-α-tubulin (Ac.-α-tub., cilia, grey),
F-actin (Actin, cell borders and morphology, grey), and mucus (PNA, magenta).
Targeted cells were identified by membrane GFP expression (memGFP, green).
Location of insets is indicated by dashed yellow box in upper panels. Quantification of
cell type composition is depicted as pie-charts, goblet cells (blue), ISCs (yellow), MCCs
(green) and SSCs (red). n embryos (above chart) and n quantified cells (in/left of chart).
900
901
Figure 4: Foxi1 regulates its own expression
(A) IF of embryos injected with *foxi1::gfp-utrophin* or *foxi1Δ1::gfp-utrophin* reporters
(green) at st. 32 stained for Acetylated-α-tubulin (Ac.-α-tub., cilia, grey), F-actin (Actin,
cell borders and morphology, grey), and serotonin (SSCs, grey) at st. 32. Targeted cells

were identified by nuclear RFP expression (H2B-RFP, blue). Right panels show false-
color of GFP fluorescence intensity. **(B)** Brightfield and epifluorescence images of
hemisected st. 11 gastrula embryos injected vegetally with *foxi1::gfp-utrophin* (green),
membrane RFP (memRFP; magenta) as control (*memRFP*) or with additional co-
injection of *foxi1* mRNA (*foxi1 + memRFP*). Right panels show false-color of GFP
fluorescence intensity. Induction was scored as positive when GFP was detected in
areas below the equator (mesendoderm). Ctrl. n = 7 induced, 26 non-induced; *foxi1*
mRNA = 26 induced, 11 non-induced.
914
915
Figure 5: Foxi1 regulates mucociliary epidermal competence through epigenetic
means
(A) Profiles of ATAC-Seq normalized accessibility around peak center ±1 kb in controls
(ctrl.) and foxi1 morphant (*foxi1*MO) organoids. (**B**) Venn diagram of peaks present in
uninjected organoids (grey) and *foxi1* MO-injected organoids (purple). (**C**) Top 15
transcription factor binding motifs predicted in sets of peaks with lost, maintained or
gained accessibility after *foxi1* MO. **(D)** Distribution of accessible regions around
epidermal genes *krt12.4* and *dll1.* Lost, maintained and gained tracks as generated by
MACS2 bdgdiff analysis and visualized in IGV. Turquoise track = control (ctrl.) and
purple track = morphant (*foxi1* MO). n = 2 organoids per condition and replicate. 3
replicates.
927
928
Figure S1: Notch regulation of ISC genes and ISC-subtype markers
(A,B,C) Effects of Notch gain (*nicd*; **A**), Notch loss (*suh-dbm*; **B**) and Notch and MCC
loss (*suh-dbm + dn-mcidas*; **C**) on core ISC gene expression in key developmental
stages (st. 10, 16, 25, 32). Heatmaps depict log2-fold change values derived from
DEseq2. Asterisks indicate statistical significant (adj-p value < 0.05) changes. **(D)**
Boxplots of ISC gene expression from scRNA-seq data published in Aztekin et al.,
2019. Visualization was generated using the published online tool:
marionilab.cruk.cam.ac.uk/XenopusRegeneration.
937
938
Figure S2: Early Foxi1 functions, expression and reporter construct
(A) Representative brightfield images of controls (ctrl.) and embryos (animal views) after
foxi1 MO (*foxi1*MO; high concentration) injection at st. 8. Morphants showed enlarged
cells and delamination of animal cells into the blastocoel. Quantification of results
shown in the right graph. Delamination events were scored based on morphological
analysis. n = number of embryos. **(B)** WMISH expression analysis of *foxi1* across
mucociliary epidermis development stages (st. 9 - 32). St. 9, 10 = animal views; st. 12,
16 = ventral views; st. 25, 32 = lateral views. Bottom row panels = magnified views of
epidermal areas. **(C,D)** Generation and promoter sequences of *foxi1::gfp-utrophin* or
foxi1Δ1::gfp-utrophin reporters. **(C)** Schematic representation of cloned genomic *foxi1.S*
promoter locus (grey box) and position of Foxi2 binding region determined in Cha et al.,
2012 (black outlined box). **(D)** Promoter sequence with indicated predicted core Foxi
binding-motifs (yellow) and Foxi2 binding region (bold, underscored).

952
953
Figure S3: Characterization of the foxi1 reporter
(A,B) IF of embryos injected with *foxi1::gfp-utrophin* (green) and α-tub.::mscarletI
(magenta) reporters at st. 32 stained for Acetylated-α-tubulin (Ac.-α-tub., cilia, grey), F-
actin (Actin, cell borders and morphology, grey), and serotonin (SSCs, grey) in **(A)**; or
for Acetylated-α-tubulin (Ac.-α-tub., cilia, grey) and F-actin (Actin, cell borders and
morphology, grey), in **(B)**. In **(B)** targeted cells were identified by nuclear RFP
expression (H2B-RFP, magenta). **(C)** WMISH expression analysis of *foxi1::gfp-utrophin*
(stained for *gfp* transcripts) across mucociliary epidermis development stages (st. 9 -
32). St. 9, 10 = animal views; st. 12, 16 = ventral views; st. 25, 32 = lateral views.
Bottom row panels = magnified views of epidermal areas. Related to sections shown in
Fig. 2D.
965
966
Figure S4: Foxi1 expression in mucociliary development and Notch ligand
induction
(A) IF for *foxi1::gfp-utrophin* reporter (green) and F-actin (Actin, cell borders and
morphology, magenta) at st. 10.5 - 32 on hemistected embryos. Targeted cells were
identified by membrane RFP expression (mRFP, grey). Related to sections shown in
Fig. 2E. **(B)** Heatmap of mucociliary marker gene enrichment differentiation inlneages
from scRNA-seq data published in Briggs et al., 2018. Values were derived using the
published   online   tool:
kleintools.hms.harvard.edu/tools/currentDatasetsList_xenopus_v2.html. NNE = non-
neural ectodermal precursors; BC – basal cells; ISC = ionocytes; MCC = multiciliated
cells; SSC = small secretory cells; GB = outer-layer goblet cells. **(C)** Representative
images of st. 9 control (ctrl.) and manipulated embryos (animal views) after mRNA
overexpression of transcription factors to test premature induction of *dlc*. Quantification
of results and effects on *dll1* (yellow) and *dlc* (blue) graphs. Embryos were scored as
induced or non-induced expression. Related to Fig. 3A.
982
983
Figure S5: Effects of Notch manipulation of mucociliary development
(A) IF of control (ctrl.), *dll1* (*dll1* MO) and *notch1* (*notch1* MO) morphants and
morphants after Notch gain of function (co-injected with *nicd*) at st. 32 stained for
Acetylated-α-tubulin (Ac.-α-tub., cilia, grey), F-actin (Actin, cell borders and morphology,
grey), and mucus (PNA, magenta). Targeted cells were identified by membrane GFP
expression (memGFP, green). Quantification of cell type composition is depicted as pie-
charts, goblet cells (blue), ISCs (yellow), MCCs (green) and SSCs (red). n embryos
(above chart) and n quantified cells (in/left of chart). **(B)** WMISH for cell type markers
(*foxi1* = ISC; *foxj1* = MCCs; *foxa1* = SSCs; basal cells = *ΔN-tp63*) at st. 16/17 in controls
and after knockdown of *dll1* (*dll1* MO). + = low dose, ++ = medium dose, +++ = high
dose. Sections of embryos allow detailed comparison between injected (*dll1* MO) and
uninjected control (ctrl.) sides of embryos. Graphs: Quantification of results. Expression
levels were scored more, equal, less expression on the injected vs. uninjected sides. n
= number of embryos.

998
999
Figure S6: Foxi1 is required for genomic accessibility of mucociliary genes
(A-D) Distribution of accessible regions around genes required for development and cell
fates specification in the embryonic mucociliary epidermis of *Xenopus*. Lost, maintained
and gained tracks as generated by MACS2 bdgdiff analysis and visualized in IGV. **(A)**
ubp1.L; **(B)** *dmrt2.S;* **(C)** *foxj1.L;* and **(D)** *tp63.L.* Turquoise track = control (ctrl.) and
purple track = morphant (*foxi1* MO). n = 2 organoids per condition and replicate. 3
replicates.