Regulation of sensorimotor serial learning in speech production by motor compensation rather than sensory error

Yuhan Lu[1,2,†], Xiaowei Tang[3,†], Zhenyan Xiao[2,4], Anqi Xu[5], Junxi Chen[3**], Xing Tian[1,2,4*]

[1]Shanghai Key Laboratory of Brain Functional Genomics (Ministry of Education), School of Psychology and Cognitive Science, East China Normal University, Shanghai 200062, China

[2]NYU-ECNU Institute of Brain and Cognitive Science, New York University Shanghai, Shanghai 200062, China

[3]Department of Neurosurgery, Guangzhou Sanjiu Brain Hospital, Guanzhou 510510, China

[4]Shanghai Frontiers Science Center of Artificial Intelligence and Deep Learning; Division of Arts and Sciences, New York University Shanghai, Shanghai 200122, China

[5]School of Humanities and Social Sciences, Harbin Institute of Technology, Shenzhen 518055, China

[†]These authors have contributed equally to this work

Corresponding:

*Xing Tian, E-mail: xing.tian@nyu.edu
Division of Arts and Sciences,
New York University Shanghai, Shanghai 200122, China.

**Junxi Chen, E-mail: chenjunxi2006@163.com
Department of Neurosurgery,
Guangzhou Sanjiu Brain Hospital, Guanzhou 510510, China.

Abstract

Motor control is essential for organisms to efficiently interact with the environment by maintaining accurate action and adjusting to future changes. Speech production, one of the most complex motor behaviors, relies on a feedback control process to detect sensory errors and trigger updates in a feedforward control process that implements compensations. However, the specific contributions of these critical processes in sensorimotor learning during continuous vocal production remain debated. Here, we used two experimental designs in five experiments to dissociate these mechanisms. First, we employed a serial-dependence design with randomized pitch perturbations, dissociating the influences of sensory errors and motor compensation on subsequent vocalizations on a trial-by-trial basis. We found that motor compensation, rather than sensory errors, predicted the compensatory responses in the subsequent trials, suggesting instantaneous serial learning mediated by updates in the feedforward process. This compensation-driven serial learning was generalized across productions of different vowel categories. Second, we further implemented a serial-dependence adaptation design in a sentential context, where auditory perturbation occurred only on a preceding syllable. Any learning effects in its subsequent syllable without pitch perturbation would reflect changes in the speech motor representation. Our results consistently revealed that compensation in the preceding syllable predicted pitch changes in the subsequent syllable, but only when the two adjacent syllables were embedded within a word boundary. Collectively, the study provides ecological-valid evidence supporting that error-based motor compensation, incorporating cognitive and linguistic constraints, directly regulates the speech motor representation and mediates the instantaneous serial learning in successive actions.

Keywords: speech motor control, feedforward, feedback, internal forward model, inverse model, sensory error, motor compensation, sensorimotor learning, serial dependence

Introduction

The accuracy and precision of motor output, as a result of efficient interplay among motor, sensory and cognitive systems, are crucial for human survival and social interactions.[1-3] In the domain of speaking, for example, the control of articulators relies on real-time integration among the language system, speech motor system and sensory feedback from the external environment.[4-7] Specifically, the brain anticipates the sensory consequences of articulatory movements via internal forward models,[8-13] continuously compares the predictions to sensory feedback in the feedback process (**Fig. 1A**, left panel).[14-16] Experimentally introduced auditory perturbations (e.g., pitch shifts) create mismatches between expected and heard feedback, effectively serving as sensory errors that trigger the inverse model and update speech motor representation in the feedforward process to correct articulation in real-time (**Fig. 1A**, left panel).[17,18] Notably, such online compensatory adjustments can propagate to influence subsequent productions[19-21], reflecting a form of rapid, trial-by-trial sensorimotor learning.

It remains heavily debated, however, how these serial adjustments contribute to sensorimotor learning via feedback and feedforward control mechanisms.[5,22-25] One possibility is that serial adjustments reflect changes in the feedback control process, whereby sensory errors encountered in a previous utterance recalibrate the sensory system's sensitivity to subsequent errors.[26,27] Supporting this hypothesis, recent findings show that exposure to unreliable auditory feedback down-weights auditory error signals and leads to reduced compensatory response in subsequent production.[19,28] Alternatively, another possibility posits that serial learning reflects updates to the feedforward process, and motor compensation acts not only as an immediate corrective response, but also leaves short-term traces in the speech sound map or articulator map (as proposed in the DIVA model).[22,25,29,30] These traces can persist beyond the removal of perturbation, leading to relatively long-lasting adaptive changes that shape subsequent productions even in the absence of feedback perturbation and sensory errors.[20]

Yet, disentangling the contribution of feedback and feedforward processes to sensorimotor learning remains a major challenge, as sensory errors and motor compensations are tightly coupled. Standard auditory feedback perturbation and adaptation paradigms struggle to differentiate the two because of their consistent co-occurrence across trials.[31] To overcome these obstacles, we employed a serial-dependence paradigm that leveraged trial-by-trial variability to dissociate the effects of past sensory error and motor compensation on future speech behavior, allowing us to test feedback and feedforward hypotheses (**Fig. 1A**, right panel).[32,33] In Experiments 1 and 2 with a serial-dependency perturbation design, if previous sensory errors did not influence compensation in the current trial, this would suggest that the 'one-shot' serial learning occurs either in the subsequent inverse model or within the speech motor representation. In Experiment 3 with a serial-dependency adaptation design, perturbation was introduced in the previous trial but not in the current trial; the inverse model would not operate due to the absence of perturbation. If behavioral changes were still observed despite the absence of perturbation, this would indicate that serial learning effects are due to updates in the speech motor representation within the feedforward process. Our results in five experiments consistently show that prior motor compensation, rather than sensory error, predicts subsequent vocal production. The serial learning

98 effects from past perturbed production adapt to the current production without
99 perturbation. These results provide evidence indicating that motor compensation is not
100 only a reflexive response to sensory perturbation, but also drives updates in speech
101 motor representation, forming an instantaneous plasticity under the linguistic constraints
102 during continuous speaking.

Results

Experiment 1: Trial-to-trial learning in repeated production of identical vowel

In the first experiment, we used the simplest setting where participants repeatedly
produced a single vowel continuously for 2.5 sec ($N = 14$). When their auditory feedback
(**Fig. 1B**, blue lines) was unexpectedly downward or upward perturbed, subjects could
compensate for the shift in speech output (orange lines). Through a long sequence of
trials in which six levels of perturbation amount were randomly applied to each trial (**Fig.**
1C), we separately measured how the serial changes of the current trial depended on
the sensory error and motor compensation of the immediately preceding trial.
Throughout the study, we defined each within-trial perturbation and compensation as C_t
and P_t, and its preceding-1-trial perturbation and compensation as C_{t-1} and P_{t-1}.

Figure 1. Study design. A, Schematics of a processing model of speech production
control. Feedback and feedforward processes interactively mediate the control of speech
production. When auditory feedback is perturbed and is inconsistent with the predicted
consequence from an internal forward model, sensory error is generated and triggers the
inverse model in the feedback control. The output of the feedback control further
updates the programming of speech-motor representation to compensate for the
auditory perturbation. The auditory perturbation and resultant motor compensation could
leave traces in either feedback and/or feedforward control processes that mediate
sensorimotor learning, even on a trial-by-trial basis. We separately use two paradigms,
firstly to test whether sensorimotor learning arises from the changes in the feedback or
feedforward processes, and secondly to test whether the learning can be attributed to
the updates in the inverse model or in the speech motor representation. **B**, Real-time
pitch perturbation during vowel production. The pitch of auditory feedback is artificially
shifted (blue) from the utterance recorded in the microphone (orange). The shifted
auditory feedback is sent to earphones, which causes compensation in the opposite
direction of the pitch shift (the upward change in the orange line of the utterance after
the downward pitch shift in the blue line). **C**, The trial sequence of the experiment.
Participants undergo 320 trials each day for three consecutive days. The perturbation
amounts and directions are randomly applied to each trial.

We first measured motor responses to perturbation in the current trial (i.e., the
relationship between C_t and P_t). We averaged across trials with the same amount of

perturbation. It demonstrated that motor compensations were opposed to the direction of
pitch perturbation, started around 150 ms after the onset of the perturbation in the
conditions of -300 cents, -180 cents, -60 cents, and 60 cents pitch shift (**Fig. 2A**, from
left to right, respectively, $ts(13)$ = 36.13, 56.36, 109.85, and -92.86, ps = 0.043, 0.021,
0.008, and 0.024, cluster-based permutation test), yet this effect was not significant in
the condition of 180-cent and 300-cent perturbations ($ps > 0.203$). To compare
conditions, we selected a 150~250 ms window that was commonly used to quantify the
amount of motor compensation.[21,34] Compensation amounts did not significantly differ
across upward and downward perturbations ($p > 0.067$, Friedman's chi-square test, FDR
corrected; **Fig. 2B**).

Next, we tested the feedforward and feedback hypotheses by separately measuring the
dependency of preceding-1-trial sensory error or motor compensation on its subsequent
trial. Specifically, we characterized whether C_t (in each trial) would follow or oppose C_{t-1}
or P_{t-1} (in its preceding trial). Therefore, we plotted C_t as a function of C_{t-1} and P_{t-1},
respectively (**Fig. 2CD**). The averaged data across trials (grey dots) and participants
(black lines) demonstrated that functions of C_t to C_{t-1} yielded a derivative-of-Gaussian-
shaped (DoG) curve (ps = 0.0001 for both, two-sided bootstrap, FDR corrected) (**Fig.
2C**). That is, C_t in the current trial followed C_{t-1} in the preceding trial, especially around 0;
if C_{t-1} were approaching the extreme condition (far from 0), the following effect became
weak.

We further combined trials with the same P_t to isolate the dependency of P_{t-1} (in the
preceding trial) on C_t (in the current trial). The main effect of P_{t-1} was not significant ($\chi^2(5)$
< 6.65, $p > 0.247$; Friedman's chi-square test, FDR corrected). We further built a linear
mixed-effect model to validate the contribution of P_t, P_{t-1}, and C_{t-1} to the C_t. Consistently,
the results revealed a significantly negative coefficient of P_t (same as **Fig. 2A**) and a
significantly positive coefficient of C_{t-1} (same as **Fig. 2C**) ($t(11,680)$ = -21.8 and 3.03, p =
0.0001 and 0.004, respectively, one-sample t-test, FDR corrected; **Fig. 2E**). No effect
was found for P_{t-1} (same as **Fig. 2D**) ($t(11,680)$ = 1.55, p = 0.160; **Fig. 2E**). These results
suggested that the preceding sensory error only contributed to the instant computation
for compensation, but not directly to the future production. It also suggested that updates
in the feedforward control process, not the feedback process, regulated subsequent
production.

Figure 2. Serial changes of the immediately preceding trial on the compensatory
responses of the current trial. **A**, Pitch traces of motor compensation in response to
upward or downward perturbations. The orange lines denote the averaged pitch trace
across participants. The red horizontal bars denote the times of the significant effects of
compensation ($p < 0.05$). **B**, Temporal averaged compensation in the periods of interest
(150-250 ms after the perturbation onset). In **C** and **D**, we plot compensation of the
current trial, C_t, as a function of compensation in the immediately preceding trial, C_{t-1},
and as a function of the perturbation in the immediately preceding trial, P_{t-1}, respectively.
C, In the left plots, each dot represents one trial and each black curve represents one
participant. The right plots show the relationship between preceding and current trials
after averaging across trials and participants. Color dots projecting to the same sign on
the x and y axes indicate that C_t is in the same direction as C_{t-1}. **D**, C_t as a function of P_{t-1}
does not show significant effects, suggesting the current compensation is not influenced
by the perturbation amount of the immediately preceding trial. **E**, The effects of current
perturbation (P_t), preceding perturbation (P_{t-1}) and preceding compensation (C_{t-1}) on
current compensation (C_t) revealed by a generalized linear mixed-effect model.
Regression coefficients for each predictor are extracted from the model. Current
perturbation and preceding compensation exert opposite and attractive influences,
respectively, on the current compensation. Error bars denote SEM across trials. In **A-D**,
shaded areas and error bars denote the standard error of mean (SEM) across
participants. *$p < 0.05$, **$p < 0.005$.

Experiment 2: Trial-to-trial learning across the production of different vowels

Speech production involved variable vowel categories and thus in Experiment 2, we
explored whether the serial effect of C_{t-1} remained across different vowels. In Experiment
2a, participants ($N = 17$) were asked to produce four different vowels that were randomly
presented in each trial, i.e., 啊 (/a1/), 咿 (/i1/), 呃 (/e1/), and 哦 (/o1/). We observed a
similar serial effect of C_{t-1} on C_t ($p = 0.0001$, two-sided bootstrap, FDR corrected; **Fig.
3A**), but not for the effect of P_{t-1} (**Fig. S1**). When comparing experiments 2a and 1, the

amplitude of the DoG curve was significantly lower when participants produced different
vowels than when they produced the same vowel (p = 0.036, two-sided bootstrap, FDR
corrected; **Fig. 3C**). The reduction of serial learning can be caused by different vowels or
by the uncertainty of next vowel. To exclude the latter possibility, we conducted
Experiment 2b where the four vowels were presented in a fixed order; that is,
participants would anticipate the identity of the next vowel. We found a consistent serial
effect caused by C_{t-1} (N = 12; p = 0.0001, two-sided bootstrap, FDR corrected; **Fig. 3B**).
Importantly, the amplitude of the DoG curve during fixed-order vowel production was
similar to that during random-order (p = 0.307, two-sided bootstrap, FDR corrected; **Fig.
3C**), suggesting that serial learning of motor compensation was reduced due to different
vowels, instead of uncertainty of next production.

Figure 3. Serial effects across different vowel categories. A and **B**, C_t as a function
of C_{t-1} when participants produce randomly-presented vowels (**A**) and when participants
produce expectedly-presented vowels (**B**). In both experiments, significant serial
changes to the current motor compensation are observed. **C**, The amount of
compensation, characterized by the amplitude of the DoG curve, shows differences
between experiments 1, 2a, and 2b, where participants produce the same vowel,
different vowels with uncertainty, and different vowels with certainty, respectively. The
learning effects are weaker when producing different vowels, compared with those of the
same vowels. Error bars denote one SD of the bootstrap distribution. *p < 0.05, **p <
0.005. See also Figure S1.

Experiment 3: inter-syllabic learning in sentence production within and across
word boundary

Experiments 1 and 2 demonstrated that previous motor compensations drove trial-to-trial
updates in current speech production, indicating that experience influenced future
productions via the feedforward, rather than feedback, control process. However, it
remains unclear whether this compensation-driven learning reflects updates in the
inverse model or direct modifications to the speech motor representation itself (**Fig. 1A**).
To resolve this ambiguity, we employed an adaptation paradigm in which auditory
perturbation was selectively applied to an initial syllable, while leaving the subsequent
syllable unperturbed. If trial-to-trial learning relies on updating the inverse model to
correct sensory errors, then no adaptive changes should emerge during the unperturbed
syllable since the inverse model would not be activated without sensory error.
Alternatively, if previous compensations directly reshape the speech motor

representation, adaptive changes should persist in subsequent unperturbed syllables
regardless of sensory error.

We manipulated the auditory feedback during the continuous production of sentences at
different positions of syntactic structures in two separate experiments (N = 15 and 13 in
Experiment 3a and 3b, respectively). During sentence production, auditory pitch
perturbation was randomly applied to either the second, third, or fourth syllable in each
trial (**Fig. 4AC**). We quantified compensation in the perturbed syllable and its serial
changes in the next syllable, which could be within or across the word boundaries
(shown as grey rectangles in insets; **Fig. 4BD**). We showed that upward pitch
perturbation drove a significant compensatory response around 200 ms after
perturbation onset, compared with the pitch trace without perturbation (t(14) < -40.30, p
< 0.009, cluster-based permutation test; **Fig. S2**). We extracted 200-300 ms mean pitch
as compensation amount in the perturbed syllable and as serial-learning amount in its
next syllable. Only when the two syllables formed a word, the compensation in the
preceding syllable influenced the production of the current syllable (right plot in **Fig. 4B:**
r = 0.09, p = 0.003; left plot in **Fig. 4D**: r = 0.22, p = 1 × 10^{-11}; Spearman correlation).
The correlation coefficients were larger in the within-word-boundary condition than in the
across-word-boundary condition (**Fig. 4B**: p = 0.180 and 0.026 for position 2 vs 4 and for
position 3 vs 4, respectively; **Fig. 4D**: p = 0.010 and 0.0005 for position 2 vs 3 and for
position 2 vs 4, respectively; one-sided bootstrap, FDR corrected). In contrast, auditory
perturbation in the preceding syllable did not contribute to inter-syllabic learning (**Fig.
S3**). These results strongly suggested that compensation-driven learning persisted in the
current vocal production even without any perturbation and compensation, and the serial
learning was constrained by word boundary. It also suggested that past compensations
directly modified speech motor representations for the instantaneous, trial-to-trial
learning.

Figure 4. Inter-syllabic learning persists only within the word boundary. A and **C**, Experiment 3a and 3b. Participants are asked to produce three 5-syllable sentences that have the same syntactic structure in each experiment. The position of the two-syllable noun is either at the end (Exp. 3a) or in the middle of the sentence (Exp. 3b). Each character is presented on the screen for 500 ms, followed by a blank for 500 ms. **B** and **D**, Relations between the compensation amount of the preceding syllable (x-axis) and serial-learning amount of the current syllable (y-axis) in Exp. 3a and 3b. The mean pitch averaged across 200-300 ms is separately extracted from the perturbed syllable (black dots in the insert plots) and the subsequent syllable (the dot pointed by a black arrow). The grey-shaped rectangle denotes the two-syllabic word. Only when the two syllables are within a word boundary, significant inter-syllabic learning is observed. **p < 0.005. See also Figures S2 and S3.

Discussion

We investigated how previous sensory error and motor compensation regulated trial-to-trial learning. We showed that motor compensation in the preceding trial influenced speech production in the next trial, whereas sensory errors did not have any effects. Crucially, this compensation-driven serial learning was modulated by linguistic contexts -- persisting across different vowels in a string of production and across vowels embedded within the word boundary. These results suggested that instantaneous sensorimotor learning was driven by motor compensation and was constrained by the linguistic system in continuous speech.

The study introduces a serial-dependence paradigm in the domain of speech production that dissociates the influence of previous sensory error and motor compensation on the current vocal production. We observed that when the compensation in trial t-1 was

around zero -- despite the presence of sensory error -- the compensation in trial t also
returned to baseline. Whereas, when the compensation in trial t-1 was positive or
negative, compensation in trial t followed the same direction. These observations are
consistent in all three experiments (**Figs. 2C, 3AB**), and can even be found in individual
subjects. Moreover, the compensation-driven serial learning was also observed in the
production of syllables without feedback perturbation (**Fig. 4**). This evidence suggests
that instantaneous plasticity in speech production emerges primarily through immediate
updates in the feedforward control process, initiated by preceding compensatory
responses. Using a novel serial dependency paradigm, this study extended
sensorimotor learning in a relatively longer time scale[32,35,36] to trial-by-trial instantaneous
plasticity.

The distinction between sensory error and motor correction on driving adaptation has
been discussed in the motor control literature. For example, studies on reaching have
shown that behavioral adaptation occurs even when online motor corrections are
unavailable, suggesting that sensory errors suffice to update subsequent movements.[37]
Speaking, however, differs from limb movements in many aspects, including the
automatic and involuntary nature of sensorimotor mappings,[38,39] as well as its mandatory
interplay with higher-level linguistic processing. It is possible that speech corrective
signals are generated in response to sensory errors in the brain and drive adaptive
changes in subsequent productions, regardless of overt compensatory behavior.
Moreover, the storage-based hierarchical yet fixed mappings between the linguistic and
motor systems may reinforce the feedforward process during control and learning in
speaking, as demonstrated in the results of Exp. 3. Rather than removing compensatory
behavior entirely, the current study directly tests whether the amount of compensation in
one trial influences the next, providing new insight into the trial-by-trial dynamics of
sensorimotor learning in speech.

The relationship between within-trial compensation and next-trial learning is not linear; it
is more likely a DoG-shaped curve. Specifically, if the previous compensation is large,
the serial changes in the current trial become smaller, suggesting a dynamic adjustment
process during speech planning.[5,6,40] Furthermore, this process is also constrained by
the similarity of production targets between trials. If the target in the next trial changes to
a different vowel, the trial-to-trial learning becomes weaker (**Fig. 3C**). Rochet-Capellan
et al. [41] tested whether sensorimotor learning during training transfers to new utterances
with different sounds and demonstrated that it is highly dependent on the specific
utterances. That is, the experience in the motor domain becomes less useful when the
next utterance is different.

Lastly, the extension of compensation-driven serial learning into sentence production
reveals that serial learning only exists within word boundary (**Fig. 4**). In Experiment 3,
participants may simply read out aloud each visually presented word without syntactic
processing. If this were true, inter-syllabic learning, as observed in Experiments 1 and 2,
would occur between every two syllables. However, the observed significant effects of
syntactic structures are inconsistent with the alternative 'reading-only' hypothesis.
Moreover, the extensive training on the three 5-syllabic sentences ensured that
participants were extremely familiar with the materials and retrieved the simple sentential
structure at the visual presentation of the first word. All the experimental settings and

positive results support the hypothesis regarding the role of syntactic constraints in
compensation-driven serial learning.

The results of within-word-boundary inter-syllabic learning suggest that words serve as a
central unit in speech planning. Motor compensation in the preceding syllable can be
updated to the next within-word syllable. However, any compensation-based experience
is not available for the syllables across words. Psycholinguistic studies, such as those by
Levelt [42], propose that during speech production, words are retrieved from the mental
lexicon as a whole unit, followed by phonological encoding into syllabic and phonetic
representations. The entire course of speech production at the linguistic level would
interact with sensorimotor processes via internal forward models and regulate
feedforward processes at both featural and temporal domains,[11,13,43-47] to achieve
efficient online control. The observed serial effects within word structure suggest that the
programming and regulation of speech production are presumably in the size of the
encoding unit within the word structure, which is consistent with the hypothesis of
interaction between linguistic and sensorimotor systems in speech production control.[4]

The study showed that previous motor compensation can modulate the subsequent
vocal pitch production; whether this compensation-driven serial learning can be
generalized to other speech domains, such as formants[19] and intensity[11] that are
controlled by distinct vocal apparatus and may be mediated by potentially different
mechanisms, requires further investigation. Considering a broader linguistic context in
speech motor control is challenging and relevant studies are rare.[48,49] In this study, we
used pitch as a representative dimension and carefully selected different combinations
of consonants and vowels in continuous speech to pioneer the investigation. Future
studies should be carried out with caution and careful controls. For example, pitch
contours in continuous speech can be potentially interfered with by consonantal
perturbations[50] or sentence intonation[51]. During the production of non-sonorant
consonants, the vibration of vocal folds will be affected, leading to a break and a carry-
over rising or falling in the pitch contours.[52]

In summary, by implementing a novel serial-dependence paradigm in the domain of
speaking, we found distinct functions of sensory error and motor compensation in
sensorimotor learning in continuous speech. Motor compensation drives updates in the
feedforward control process and regulates trial-to-trial learning that was constrained by
linguistic contexts, while sensory error only contributed to the instant computation of
compensation. These results suggest that the history of motor processes significantly
contributes to sensorimotor learning and control. Systematic interplay among cognitive,
motor, and sensory systems regulates behaviors.

Materials and methods

Participant

In five experiments, i.e., Experiments 1, 2a, 2b, 3a, and 3b, we recruited 14 participants
(9 females, 18-27 years old, mean 21 years old), 17 participants (11 females, 19-32
years old, mean 23 years old), 12 participants (8 females, 20-27 years old, mean 22
years old), 15 participants (9 females, 19-25 years old, mean 22 years old), and 13
participants (7 females, 18-26 years old, mean 21 years old), respectively. They were
college students with self-reported normal hearing and speech. All participants were

right-handed and were native Chinese speakers. Each volunteer only participated in one
experiment. Written informed consent was obtained from each participant before the
experiment. The experimental procedures were approved by the Research Ethics
Committee at the East China Normal University and New York University Shanghai.

Procedure in Experiment 1

Participants were asked to produce the vowel /a/ in a total of 960 trials over three days.
Each trial began with a white fixation cross, displayed at the center of the screen for a
randomized duration of 0.8–1.2 s (uniform distribution), signaling participants to be
prepared for vocalization. When the cross turned green, participants were instructed to
vocalize no more than 500 ms and sustain the /a/ sound with a steady and natural tone
for 2.5 s until the cross disappeared. A pitch perturbation for a duration of 500 ms was
randomly introduced 0.8–1.2 s after vocal onset. The direction (i.e., upward or downward)
and magnitude (i.e., 60, 180, or 300 cents) of the pitch shift were randomly assigned
across the 960 trials for each participant. At the end of each trial, participants received
visual feedback: "Good," "Too slow," or "Too short." Participants completed 95.86% ±
0.02% (s.d.) of trials that met criteria. Each trial was followed by a silent interval with a
random duration of 1–2 s (uniform distribution) before the next trial began.

Participants completed 320 trials in 10 blocks each day, with self-paced rest periods of
at least 1 minute between blocks. At the beginning of each session, participants
underwent a training block of 10 trials. The mean pitch from these trials was used to
define the pitch range in the Audapter toolbox (see *Auditory feedback perturbation*
below).

Procedure in Experiment 2a

The procedure for Experiment 2a was identical to Experiment 1, except participants
were required to produce four different vowels without semantic meaning—啊 (/a1/), 咿
(/i1/), 呃 (/e1/), and 哦(/o1/)—randomly assigned in each trial. A white letter
corresponding to the syllable was presented at the center of the screen, and upon
turning green, participants were instructed to vocalize a vowel. Participants completed
94.82% ± 0.11% of trials that met the criteria.

Procedure in Experiment 2b

402 The procedure for Experiment 2b was identical to Experiment 2a, except the order of
403 four vowels were presented in a fixed order, i.e., /i1/-/a1/-/e1/-/o1/ in every four trials so
404 that participants would not anticipate the next vowel production. Participants completed
405 97.55% ± 0.03 of trials that met the criteria.

Procedure in Experiment 3a

The procedure for Experiment 3a followed that of Experiment 1, with two differences.
First, participants completed 300 trials in a single day. Second, they were required to
produce sentences with the same syntactic structure rather than isolated vowels. In each
trial, one of three five-syllable sentences (i.e., ˙ ˙ ˙ 猫咪 [ma1 ma1 lu1 mao1 mi1], 丫丫
邀巫医 [ya1 ya1 yao1 wu1 yi1], 依依摸 ˙ ˙ [yi1 yi1 mo1 wu1 ya1]) was randomly
selected (uniform distribution), with each syllable appearing sequentially at the center of

the screen for 500 ms, followed by a 500 ms blank screen. The whole duration for a
sentence production was 5 s. During vocalization, a whole-utterance pitch perturbation
(upward or downward perturbation at 60, 180, 300 cents, randomly assigned) was
applied to the second, third, or fourth word during vocalization (uniformed distribution). In
20% of the sentences, no perturbation was applied. The speech materials and task were
carefully chosen to avoid the effect of consonantal obstruction[53] and the effect of
articulators' dynamics on pitch contour, respectively. Participants completed 94.44% ±
0.07% of trials that met the criteria.

Before the main experiment, participants completed a training session in which they
produced each of the three sentences four times. They were explicitly instructed on the
content and syntactic structure of the sentences. Additionally, they were instructed to
vocalize each syllable within 500 ms of its appearance and to avoid coarticulation
between words. Participants could repeat the training session upon request.

Procedure in Experiment 3b

The procedure for Experiment 3b was the same as Experiment 3a, except that the three
sentences were replaced with another three with a different syntactic structure, i.e., 邀依
琳摸ˇ [yao1 yi1 yi1 mo1 ying1], 拉ˇˇ腌ˇ [la1 ma1 ma1 yan1 ya1], and ˇ丫丫溜猫
[yue1 ya1 ya1 liu1 mao1], where the two-syllable noun was in the position of second and
third character in the sentences. Participants completed 97.36% ± 0.03% of trials that
met the criteria.

Auditory feedback perturbation

Participants' speech was recorded by a Shure beta 58A microphone that was placed 5
centimeters away from their mouths. The acoustic signals from the microphone were
pre-amplified and then provided to an audio interface (MOTU Microbook IIc), collected at
a 48k Hz sampling rate. The recorded speech was real-time manipulated (i.e.,
perturbation) in the audio interface, and the processed signal was split and sent both to
the computer and to the headphones. The speech processing in the audio interface was
performed using the Audapter toolbox.[54] The pitch of speech in each trial was shifted
using the time-domain method. A 10-ms linear ramp was added at boundaries to
attenuate possible abrupt jumps in the speech. A priori pitch range was set as +/-50 Hz
relative to the mean pitch of a vocal /a/ for each participant. The rest of the parameters
were set as default including algorithm, frame length, and cepstral liftering.

Acoustic analysis

In all experiments, we analyzed signalIn, the speech output from participants. In
experiments 1 and 2, we first removed silence from the recordings of produced vowels in
each trial using the *vad* function. Visual inspection was performed to make sure noise
did not affect the voice activity detection. Pitch tracks (at a 100-Hz sampling rate) were
measured using the Praat-based script ProsodyPro.[55] Default parameters were used.
Perturbation onset in the pitch trace was obtained by a customized automatic algorithm
combined with manual inspection. To convert hertz to cent scale, we defined baseline
pitch by taking the mean of pitch trace in the -400 to -200 ms before perturbation onset.
The pitch trace was transformed according to the formula:

$cents = 1200 * \log_2(\frac{pitch}{pitch_{baseline}})$. A trial rejection was performed if the pitch trace was

varied beyond its perturbation amount minus 100 cents (for downward-perturbation trials)

or plus 100 cents (for upward-perturbation trials). To extract perturbation and

compensation amounts for DoG-curve fitting and linear mixed effect model, the pitch

trace in the 50-80 ms and the 150-250 ms windows were averaged, respectively, for

each trial.[21] Compensation amounts for all trials and all participants were corrected by

subtracting the grand average. This correction ensures that the DoG-curve fitting starts

from zero, allowing its amplitude to be accurately measured and compared across

experiments.

In Experiment 3, we applied the same pipeline except that only the perturbed syllable

and its subsequent syllable were extracted to measure their pitch trace. Based on the

visual inspection of pitch contours (**Fig. S2**), Mandarin tone 1 produced in isolation was

characterized by a high-level tone and showed a pitch contour that typically involved a

slight rising intonation at the onset, lasting around 0-150 ms, before stabilizing into a

steady high pitch.[56] Therefore, we calculated the compensation amount in the perturbed

syllable and serial-learning amount in the following syllable using mean pitch in the 200-

250 ms time window, with baseline pitch in the 150-200 ms time window.

Statistical analysis of compensatory responses

For participant-level analysis, trials with the same perturbation condition were averaged,

regardless of vowel categories in Experiments 2 and 3. To assess whether there was a

compensatory response to perturbation, a two-sided cluster-based permutation test[57]

was used to test averaged pitch traces against perturbation amount or between

perturbed and normal conditions (**Fig. 1A** and **S2**). A one-sided one-sample t-test was

used to test motor compensation extracted from time of interests against 0 (**Fig. 2B**).

Statistical analysis of trial-to-trial learning effect

To explore the serial effect of preceding-1-trial motor compensation amount on the

current motor response, C_t was expressed as a function of C_{t-1}. For positive values of

this C_t or C_{t-1}, participants compensated upward in response to perturbation in the

current or preceding-1-trial. Consequently, data points in the scatter plot (**Fig. 2C** and

3AB) that had x- and y- values in the same sign indicated that C_t was in the direction of

C_{t-1}. We assess the trial-to-trial learning effect by polling the compensation amount of all

trials and fitting the DoG-shape curve, a well-found psychophysical function in previous

serial-dependence studies.[32,36] The DoG was given as $y = x * a * w * c * e^{-(wx)^2}$, where

a was the curve amplitude, *w* was the curve width, and *c* was the constant $\sqrt{2}/e^{-0.5}$. To

assess the serial effect of preceding motor compensation, *x* was C_{t-1}, and width

parameter *w* of the curve was constrained to a range of plausible values (*w* = 0.019-

0.071, corresponding to the curve peak between 10 and 36 perturbation pitch difference).

The model fitting used the function *fmincon* with default parameters. We employed a

bias-corrected and accelerated bootstrap to evaluate the significance of the DoG models

and to compare DoG models between conditions. We applied a fitted DoG model to the

pooled group data and generated a permutation distribution for the DoG-curve amplitude

parameter.[58,59] This permutation process was repeated 2,000 times. For a one-sided

comparison of curve amplitude, if the data population in one condition was greater than

A% of the data population in the other condition, the significance level was (100A +
1)/2,001. For a two-sided comparison, the significance level was (200A + 1)/2,001. A
false discovery rate (FDR) correction was applied.

To explore the serial effect of sensory error on C_t, C_t was expressed as a function of P_{t-1}.
We use a Friedman's chi-square test for multiple condition comparison, i.e., controlling
the current perturbation as the same to explore the effect of preceding perturbation on
the current compensation (**Fig. 2D**). Spearman correlation was used to find the relation
between C_{t-1}/P_{t-1} and inter-syllabic learning within and across syntactic boundaries (**Fig.**
4BD and **S3**). To compare the correlation coefficient in within- and across-boundary
conditions, a bias-corrected and accelerated bootstrap was used.[60]

Generalized linear mixed-effect model

To quantitatively test the relations of C_t to P_t, P_{t-1} and C_{t-1}, we built a generalized linear
mixed-effect models (**Fig. 2E** and **S1**). The model assumed that P_t, P_{t-1} and C_{t-1} together
influenced C_t. The function 'fitglme' was performed and accounted for group-level fixed
effects and random effects because of vocal pitch variation across participants. The
model was specified as $C_{t\ ij} = f(\beta_{0j} + \beta_{1j}P_{t\ ij} + \beta_{2j}P_{t-1\ ij} + \beta_{3j}C_{t-1\ ij} + 1|subjects)$,
where i indicated participant index and j indicated j^{th} trial. The regression coefficient of
each predictor, i.e., β, was accessed by the t-statistics against the null hypothesis test
that the coefficient was equal to 0. FDR correction was applied.

Data and code availability: All data and code that support the findings of this study is
available in Mendeley Data: 10.17632/pybdtwxvzz.2.

Acknowledgments: We thank Yuan Xie and Bao Hong for their thoughtful comments
on previous versions of the manuscript, Tingting Wang and Huixuan Liu for assisting in
collecting data. This study is supported by the National Natural Science Foundation of
China 32271101, Program of AI-Driven Initiative to Promote Research Paradigm Reform
and Empower Disciplinary Advancement by Shanghai Municipal Education Commission
(SMEC), Program of Introducing Talents of Discipline to Universities, Base B16018, and
NYU Shanghai Boost Fund to X.T.

Author contributions: Y.L., X.T., and J.C. designed the experiment. Y.L., W.T., and
Z.X. collected the data, Y.L., Z.X., A.X., and W.T., performed the analyses, Y.L., and X.T.
drafted, reviewed, and corrected the manuscript. X.T. and J.C. supervised the project.

Declaration of interests: The authors declare no competing financial interests.

Declaration of generative AI and AI-assisted technologies in the writing process:
During the preparation of this work, the authors used ChatGPT to proofread texts. After
using this tool or service, the authors reviewed and edited the content as needed and
took full responsibility for the content of the publication.

References

1. Rosenbaum, D.A. (2009). Human motor control (Academic press).
2. Ivanenko, Y.P., Poppele, R.E., and Lacquaniti, F. (2006). Motor control programs and walking.
Neuroscientist 12, 339-348. 10.1177/1073858406287987.
3. Olivier, E., Davare, M., Andres, M., and Fadiga, L. (2007). Precision grasping in humans: from
motor control to cognition. Curr Opin Neurobiol 17, 644-648. 10.1016/j.conb.2008.01.008.

4.   Hickok, G. (2012). Computational neuroanatomy of speech production. Nat Rev Neurosci *13*, 135-
145. 10.1038/nrn3158.
5.   Houde, J.F., and Nagarajan, S.S. (2011). Speech production as state feedback control. Front Hum
Neurosci *5*, 82. 10.3389/fnhum.2011.00082.
6.   Guenther, F.H. (2016). Neural control of speech (Mit Press).
7.   Tian, X., and Poeppel, D. (2010). Mental imagery of speech and movement implicates the
dynamics of internal forward models. Front Psychol *1*, 166. 10.3389/fpsyg.2010.00166.
8.   Kawato, M. (1999). Internal models for motor control and trajectory planning. Curr Opin
Neurobiol *9*, 718-727. 10.1016/s0959-4388(99)00028-8.
9.   Wolpert, D.M., and Ghahramani, Z. (2000). Computational principles of movement neuroscience.
Nat Neurosci *3 Suppl*, 1212-1217. 10.1038/81497.
10.   Schubotz, R.I. (2007). Prediction of external events with our motor system: towards a new
framework. Trends in Cognitive Sciences *11*, 211-218. 10.1016/j.tics.2007.02.006.
11.   Tian, X., Ding, N., Teng, X.B., Bai, F., and Poeppel, D. (2018). Imagined speech influences perceived
loudness of sound. Nature Human Behaviour *2*, 225-234. 10.1038/s41562-018-0305-8.
12.   Li, S., Zhu, H., and Tian, X. (2020). Corollary Discharge Versus Efference Copy: Distinct Neural
Signals in Speech Preparation Differentially Modulate Auditory Responses. Cereb Cortex *30*, 5806-
5820. 10.1093/cercor/bhaa154.
13.   Li, Y., Luo, H., and Tian, X. (2020). Mental operations in rhythm: Motor-to-sensory transformation
mediates imagined singing. PLoS Biol *18*, e3000504. 10.1371/journal.pbio.3000504.
14.   Eliades, S.J., and Tsunada, J. (2018). Auditory cortical activity drives feedback-dependent vocal
control in marmosets. Nat Commun *9*, 2540. 10.1038/s41467-018-04961-8.
15.   Niziolek, C.A., Nagarajan, S.S., and Houde, J.F. (2013). What does motor efference copy represent?
Evidence from speech production. J Neurosci *33*, 16110-16116. 10.1523/JNEUROSCI.2137-
13.2013.
16.   Chang, E.F., Niziolek, C.A., Knight, R.T., Nagarajan, S.S., and Houde, J.F. (2013). Human cortical
sensorimotor network underlying feedback control of vocal pitch. Proc Natl Acad Sci U S A *110*,
2653-2658. 10.1073/pnas.1216827110.
17.   Lametti, D.R., Nasir, S.M., and Ostry, D.J. (2012). Sensory preference in speech production
revealed by simultaneous alteration of auditory and somatosensory feedback. J Neurosci *32*,
9351-9358. 10.1523/JNEUROSCI.0404-12.2012.
18.   Larson, C.R., Altman, K.W., Liu, H., and Hain, T.C. (2008). Interactions between auditory and
somatosensory feedback for voice F0 control. Exp Brain Res *187*, 613-621. 10.1007/s00221-008-
1330-z.
19.   Tang, D.L., Parrell, B., and Niziolek, C.A. (2022). Movement variability can be modulated in speech
production. J Neurophysiol *128*, 1469-1482. 10.1152/jn.00095.2022.
20.   Houde, J.F., and Jordan, M.I. (1998). Sensorimotor adaptation in speech production. Science *279*,
1213-1216. 10.1126/science.279.5354.1213.
21.   Hantzsch, L., Parrell, B., and Niziolek, C.A. (2022). A single exposure to altered auditory feedback
causes observable sensorimotor adaptation in speech. Elife *11*. 10.7554/eLife.73694.
22.   Tourville, J.A., and Guenther, F.H. (2011). The DIVA model: A neural theory of speech acquisition
and production. Lang Cogn Process *26*, 952-981. 10.1080/01690960903498424.
23.   Parrell, B., and Houde, J. (2019). Modeling the Role of Sensory Feedback in Speech Motor Control
and Learning. J Speech Lang Hear Res *62*, 2963-2985. 10.1044/2019_JSLHR-S-CSMC7-18-0127.
24.   Kim, K.S., Gaines, J.L., Parrell, B., Ramanarayanan, V., Nagarajan, S.S., and Houde, J.F. (2023).
Mechanisms of sensorimotor adaptation in a hierarchical state feedback control model of speech.
PLoS Comput Biol *19*, e1011244. 10.1371/journal.pcbi.1011244.
25.   Manes, J.L., Bullock, L., Meier, A.M., Turner, R.S., Richardson, R.M., and Guenther, F.H. (2024). A
neurocomputational view of the effects of Parkinson's disease on speech production. Frontiers in
Human Neuroscience *18*. 10.3389/fnhum.2024.1383714.
26.   Tang, D.-l., Parrell, B., Beach, S.D., and Niziolek, C.A. (2025). The Brain's Sensitivity to Sensory
Error Can Be Modulated by Altering Perceived Variability. The Journal of Neuroscience *45*,
e0024242024. 10.1523/jneurosci.0024-24.2024.

27.   Sitek, K.R., Mathalon, D.H., Roach, B.J., Houde, J.F., Niziolek, C.A., and Ford, J.M. (2013). Auditory
594   cortex processes variation in our own speech. PLoS One 8, e82925.
595   10.1371/journal.pone.0082925.
596   28.   Niziolek, C.A., and Parrell, B. (2021). Responses to Auditory Feedback Manipulations in Speech
597   May Be Affected by Previous Exposure to Auditory Errors. J Speech Lang Hear Res 64, 2169-2181.
598   10.1044/2020_JSLHR-20-00263.
599   29.   Kawato, M., Furukawa, K., and Suzuki, R. (1987). A hierarchical neural-network model for control
600   and learning of voluntary movement. Biol Cybern 57, 169-185. 10.1007/BF00364149.
601   30.   Guenther, F.H. (2006). Cortical interactions underlying the production of speech sounds. J
602   Commun Disord 39, 350-365. 10.1016/j.jcomdis.2006.06.013.
603   31.   Raharjo, I., Kothare, H., Nagarajan, S.S., and Houde, J.F. (2021). Speech compensation responses
604   and sensorimotor adaptation to formant feedback perturbations. J Acoust Soc Am 149, 1147.
605   10.1121/10.0003440.
606   32.   Fischer, J., and Whitney, D. (2014). Serial dependence in visual perception. Nat Neurosci 17, 738-
607   743. 10.1038/nn.3689.
608   33.   Cicchini, G.M., Mikellidou, K., and Burr, D.C. (2024). Serial Dependence in Perception. Annu Rev
609   Psychol 75, 129-154. 10.1146/annurev-psych-021523-104939.
610   34.   Burnett, T.A., Freedland, M.B., Larson, C.R., and Hain, T.C. (1998). Voice F0 responses to
611   manipulations in pitch feedback. J Acoust Soc Am 103, 3153-3161. 10.1121/1.423073.
612   35.   Fritsche, M., Spaak, E., and de Lange, F.P. (2020). A Bayesian and efficient observer model
613   explains concurrent attractive and repulsive history biases in visual perception. Elife 9.
614   10.7554/eLife.55389.
615   36.   Liberman, A., Fischer, J., and Whitney, D. (2014). Serial dependence in the perception of faces.
616   Curr Biol 24, 2569-2574. 10.1016/j.cub.2014.09.025.
617   37.   Tseng, Y.W., Diedrichsen, J., Krakauer, J.W., Shadmehr, R., and Bastian, A.J. (2007). Sensory
618   prediction errors drive cerebellum-dependent adaptation of reaching. J Neurophysiol 98, 54-62.
619   10.1152/jn.00266.2007.
620   38.   Assaneo, M.F., Ripolles, P., Orpella, J., Lin, W.M., de Diego-Balaguer, R., and Poeppel, D. (2019).
621   Spontaneous synchronization to speech reveals neural mechanisms facilitating language learning.
622   Nat Neurosci 22, 627-632. 10.1038/s41593-019-0353-z.
623   39.   Krakauer, J., Naber, C., Niziolek, C.A., and Parrell, B. (2024). Divided Attention Has Limited Effects
624   on Speech Sensorimotor Control. Journal of Speech, Language, and Hearing Research 67, 4358-
625   4368. doi:10.1044/2024_JSLHR-24-00098.
626   40.   Houde, J.F., and Chang, E.F. (2015). The cortical computations underlying feedback control in
627   vocal production. Curr Opin Neurobiol 33, 174-181. 10.1016/j.conb.2015.04.006.
628   41.   Rochet-Capellan, A., Richer, L., and Ostry, D.J. (2012). Nonhomogeneous transfer reveals
629   specificity in speech motor learning. J Neurophysiol 107, 1711-1717. 10.1152/jn.00773.2011.
630   42.   Levelt, W.J.M. (1989). Speaking (MIT press). 10.7551/mitpress/6393.001.0001.
631   43.   Tian, X., and Poeppel, D. (2015). Dynamics of self-monitoring and error detection in speech
632   production: evidence from mental imagery and MEG. J Cogn Neurosci 27, 352-364.
633   10.1162/jocn_a_00692.
634   44.   Tian, X., and Poeppel, D. (2013). The effect of imagination on stimulation: the functional
635   specificity of efference copies in speech processing. J Cogn Neurosci 25, 1020-1036.
636   10.1162/jocn_a_00381.
637   45.   Indefrey, P., and Levelt, W.J. (2004). The spatial and temporal signatures of word production
638   components. Cognition 92, 101-144. 10.1016/j.cognition.2002.06.001.
639   46.   Hartsuiker, R.J., and Kolk, H.H. (2001). Error monitoring in speech production: a computational
640   test of the perceptual loop theory. Cogn Psychol 42, 113-157. 10.1006/cogp.2000.0744.
641   47.   Nozari, N., and Novick, J. (2017). Monitoring and Control in Language Production. Current
642   Directions in Psychological Science 26, 403-410. 10.1177/0963721417702419.
643   48.   Patel, R., Niziolek, C., Reilly, K., and Guenther, F.H. (2011). Prosodic adaptations to pitch
644   perturbation in running speech. J Speech Lang Hear Res 54, 1051-1059. 10.1044/1092-
645   4388(2010/10-0162).

49. Patel, R., Reilly, K.J., Archibald, E., Cai, S., and Guenther, F.H. (2015). Responses to Intensity-
Shifted Auditory Feedback During Running Speech. J Speech Lang Hear Res *58*, 1687-1694.
10.1044/2015_JSLHR-S-15-0164.
50. Ohala, J.J. (1974). A mathematical model of speech aerodynamics. Annual Report of the Institute
of Phonetics University of Copenhagen *8*, 11-22.
51. Daneš, F. (1960). Sentence intonation from a functional point of view. Word *16*, 34-54.
52. Xu, Y. (1997). Contextual tonal variations in Mandarin. Journal of Phonetics *25*, 61-83. DOI
10.1006/jpho.1996.0034.
53. Hanson, H.M. (2009). Effects of obstruent consonants on fundamental frequency at vowel onset
in English. J Acoust Soc Am *125*, 425-441. 10.1121/1.3021306.
54. Cai, S., Boucek, M., Ghosh, S.S., Guenther, F.H., and Perkell, J.S. (2008). A system for online
dynamic perturbation of formant trajectories and results from perturbations of the Mandarin
triphthong/iau. Proceedings of the 8th ISSP, 65-68.
55. Xu, Y. (2013). ProsodyPro—A tool for large-scale systematic prosody analysis. (Laboratoire Parole
et Langage, France).
56. Tseng, C.-y. (1981). An acoustic phonetic study on tones in Mandarin Chinese (Brown University).
57. Maris, E., and Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. J
Neurosci Methods *164*, 177-190. 10.1016/j.jneumeth.2007.03.024.
58. Lu, Y.H., Jin, P.Q., Ding, N., and Tian, X. (2023). Delta-band neural tracking primarily reflects rule-
based chunking instead of semantic relatedness between words. Cerebral Cortex *33*, 4448-4458.
10.1093/cercor/bhac354.
59. Lu, Y., Jin, P., Pan, X., and Ding, N. (2022). Delta-band neural activity primarily tracks sentences
instead of semantic properties of words. Neuroimage *251*, 118979.
10.1016/j.neuroimage.2022.118979.
60. Efron, B., and Tibshirani, R.J. (1994). An Introduction to the Bootstrap (CRC press).
10.1201/9780429246593.
672