Functional analysis of the *Nematostella* Wnt/β-catenin destruction complex provides insight into the evolution of a critical regulatory module in a major metazoan signal transduction pathway

Authors: Hongyan Sun[1,†], Brian M. Walters[1], Radim Zidek[2], Mark Q. Martindale[2] and Athula H. Wikramanayake[1*]

Affiliations:

[1]Department of Biology, University of Miami, Coral Gables, FL, 33146

[2]Whitney Laboratory for Marine Bioscience, 9505 Ocean Shore Blvd. St. Augustine, FL 32080-8610

[†]Current address: Pennington Biomedical Research Center, Baton Rouge, LA, 70808

[*]Corresponding author (athula@miami.edu)

Key Words: Wnt/β-catenin signaling, Axin, Adenomatous Polyposis Coli, evolution of signal transduction pathways, Wnt/β-catenin destruction complex, *Nematostella*, Cnidaria, Mnemiopsis, Ctenophora

Abstract

The genesis of signaling pathways likely drove metazoan evolution, but the origins of

these pathways and how they acquired signaling activity is poorly understood. Here, we

studied functional evolution of the Wnt/β-catenin (cWnt) pathway destruction-complex

(DC) that regulates β-catenin signaling. Bilaterian DC function requires β-catenin

binding to Axin and APC proteins, and Axin-APC heterodimerization. However,

bioinformatic analyses predicted that Axin and APC-like homologs in early-branching

non-bilaterians lack important previously defined bilaterian β-catenin binding domains,

questioning if they have functional cWnt-DCs. We demonstrate that both Axin and APC

proteins in the cnidarian *Nematostella vectensis* (a representative of the sister taxon to

bilaterians) can regulate cWnt signaling indicating an active cWnt-DC. Using *in vitro*

analyses, we show that NvAxin binds Nvβ-catenin weakly despite lacking the conserved

bilaterian Axin β-catenin-binding motif (βcatBM). Using AlphaFold3, we identified two

predicted βcatBM-like sequences in NvAxin, one within the Axin-RGS domain and

another towards the C-terminus. Similar analysis of placozoan, poriferan, and

ctenophore Axin identified single βcatBM-like sequences located within Axin-RGS. We

show that ctenophore Axin and β-catenin do not interact and changing a conserved

leucine on NvAxin-βcatBM-like motifs to resemble the ctenophore sequence abolished

NvAxin-Nvβ-catenin interactions. We propose that an ancestral Axin-RGS sequence

acquired low-affinity β-catenin binding early in metazoan evolution, followed by motif

duplication in the cnidarian-bilaterian ancestor. In bilaterians, the duplicated βcatBM

evolved higher-affinity for β-catenin, while the ancestral sequence was lost. Our results

demonstrate how phylogenetic insights, AI tools and functional assays can be used to

reconstruct the evolution of complex signaling pathways.

Introduction

The early development and diversification of multicellular animals (metazoans) is

regulated by a small set of conserved signal transduction pathways (Gerhart 1999; Pires-

daSilva and Sommer 2003; Richards and Degnan 2009; Babonis and Martindale 2017).

It has been hypothesized that the evolution of these signaling pathways was critical for

the evolution and radiation of the metazoan last common ancestor, or "urmetazoan",

over 600 million years ago (Pires-daSilva and Sommer 2003; Rokas 2008; Richards and

Degnan 2009; Babonis & Martindale, 2017; Richter and King 2013; Ros-Rocher et al.

2021). Despite the critical roles of these pathways in metazoan development and

evolution, relatively little is known about their origins or their functional activity in early

emerging taxa (Pires-daSilva and Sommer 2003; Babonis & Martindale, 2017; Richards

and Degnan 2009). One major constraint to advancing this research has been the

difficulty in the care and handling of embryos of early emerging (non-bilaterian) model

systems to experimentally manipulate signal transduction pathways. However, the

advancement of methodology to molecularly manipulate development in cnidarians

(e.g., sea anemones, corals, "jelly fish"), the closest outgroup to the Bilateria (Fig. 1A),

and foundational knowledge of Wnt/β-catenin (cWnt) pathway function in these

animals and has established cnidarians as one of the best non-bilaterian experimental

systems to functionally understand the direction of evolutionary change in the cWnt

signaling pathway (Wikramanayake et al. 2003; Lee et al. 2007; Layden et al. 2016;

Lebedeva et al. 2021; Houliston et al. 2022; Holzem et al. 2024).

The cWnt signaling pathway (Fig. 1B) is a broadly deployed signaling pathway

that is restricted to the Metazoa, and cWnt signaling plays diverse roles during

evolution, development and homeostasis (Petersen and Reddien 2009; Richards and

Degnan 2009; Loh et al. 2016; Babonis and Martindale 2017; Richter et al. 2018;

Holzem et al. 2024). Extensive studies of cWnt pathway function in bilaterian models

have led to a relatively advanced level of understanding of the critical regulatory nodes

in this pathway, hence making it a particularly useful model to investigate how

metazoan signaling pathways evolved and acquired signaling competence. For example,

like many of the metazoan signal transduction pathways, the cWnt pathway is modular

in its organization, with distinct protein complexes regulating transduction of signals

from the cell surface to the nucleus (MacDonald et al. 2009; Stamos and Weis 2013;

Gammons and Bienz 2018; Liu et al. 2022). One critical cytoplasmic regulatory node in

the cWnt pathway is the β-catenin destruction complex (DC) which is essential for

regulating cWnt signaling through the control of cytoplasmic levels of β-catenin, the

primary transducer of cWnt signaling in the nucleus (Fig. 1B) (Roberts et al. 2011;

Stamos and Weis 2013; Gammons and Bienz 2018). Evolution of the DC was critical for

the evolution of cWnt signaling because it likely provided a scaffold for exaptation of β-

catenin for a signaling function. Moreover, the DC also may have provided a central

regulatory node for context-dependent spatial and temporal control of this pathway

during its wide deployment in bilaterians during development and homeostasis

(Petersen and Reddien 2009; Valenta et al. 2012; Loh et al. 2016; Liu et al. 2022;

Holzem et al. 2024).

Axin is a key scaffolding protein in the bilaterian DC which contains multiple

binding sites that mediate its direct association with proteins in the DC including β-

catenin, the two serine/threonine protein kinases Casein Kinase 1α (CK-1α) and

Glycogen Synthase Kinase 3β (GSK-3β), the phosphoprotein Disheveled (Dvl), and APC,

the product of the Adenomatous Polyposis Coli gene (Fig. 1C) (Ikeda et al. 1998; Kishida

et al. 1998; Fagotto et al. 1999; Valenta et al. 2012; Stamos and Weis 2013; Pronobis et

al. 2015; van Kappel and Maurice 2017; Qiu et al. 2024). A detailed model for negative

regulation of β-catenin in the cWnt pathway by the DC has emerged from extensive

functional studies done in vertebrate and invertebrate model systems (Roberts et al.

2011; Stamos and Weis 2013; van Kappel and Maurice 2017; Liu et al. 2022; Kremer et

al. 2010; Peterson-Nedry et al. 2008). This model proposes that β-catenin is recruited

to Axin through a conserved binding motif (βcatBM) and this binding mediates the

sequential phosphorylation of β-catenin by CK-1α and GSK-3β leading to its

ubiquitylation and proteasome-mediated degradation (Xing et al. 2003; Kimelman and

Xu 2006; Roberts et al. 2011; Stamos and Weis 2013; van Kappel and Maurice 2017;

Gammons and Bienz 2018; Ranes et al. 2021). The transfer of phosphorylated β-catenin

to the proteasome requires APC, the second scaffolding protein in the DC. APC in

bilaterians is a relatively large protein with multiple domains that have been shown to

mediate its interactions with Axin, β-catenin, and over 100 other proteins (Fig. 1D)

(Roberts et al. 2011; Nelson and Näthke 2013; van Kappel and Maurice 2017; Abbott

and Näthke 2023). Strikingly, APC has multiple distinct binding sites for β-catenin that

have different affinities for the protein (Ha et al. 2004; Kimelman and Xu 2006;

Roberts et al. 2011; Pronobis et al. 2017; Ranes et al. 2021). Current models postulate

that following β-catenin phosphorylation on Axin, one of the 20 amino acid repeats on

APC is phosphorylated and subsequently phosphorylated β-catenin is transferred to this

domain on APC prior to being targeted to the proteasome for degradation (Kimelman &

Xu, 2006; Roberts et al. 2011; Ranes et al. 2021). Regulation of the DC has been subject

to extensive experimental interrogation because of its critical role in regulating cWnt

signaling, and the discovery that over 90% of human colorectal cancers have mutations

in either APC or other components of the DC that can lead to misregulation of cWnt

signaling (Polakis 1995; Kinzler and Vogelstein 1996; Fodde et al. 2001; Schneikert and

Behrens 2007; Kohler et al. 2008). Oncogenic human APC mutations are frequently

seen in the "mutational cluster region" (MCR) that delete many of the well-characterized

β-catenin binding sites and the three SAMP domains on the protein that mediate Axin-

APC heterodimerization and are required for DC function (Fig. 1D) (Kinzler and

Vogelstein 1996; Fodde et al. 2001; Schneikert and Behrens 2007; Kohler et al. 2008;

Roberts et al. 2011; Ranes et al. 2021).

The cWnt pathway is deeply conserved with homologs of components of the basic

pathway found in all metazoan phyla (Richards and Degnan 2009; Babonis and

Martindale 2017; Holzem et al. 2024), but ongoing genomic sequencing studies in early

branching metazoan clades have raised interesting fundamental questions about the

origin of signaling competence in the cWnt pathway. One of the first observations that

raised the possibility that the non-bilaterian cWnt pathway may not function like the

bilaterian cWnt pathway came from a phylogenetic analysis of Wnt pathway

components in two early emerging taxa, the poriferan (sponge) *Amphimedon*

queenslandica and the cnidarian (sea anemone) *Nematostella vectensis* (Richards and

Degnan, 2009; Adamska et al. 2010). This work showed that protein domains known to

be important in DC proteins in bilaterians were absent in these two species. For

example, the Axin proteins from *A. queenslandica* and *N. vectensis* appeared to lack the

highly conserved βcatBM seen in bilaterian Axin that mediates its interaction with β-

catenin (Richards and Degnan 2009; Adamska et al. 2010) (Fig. 1C). We used the

webtool SMART (Simple Modular Architecture Research Tool: http://smart.embl-

heidelberg.de) to confirm that Axin from a ctenophore and a placozoan also lack the

βcatBM suggesting its bilaterian origin (Fig. 1C). Additionally, APC proteins in these

taxa were considerably shorter than the bilaterian homologs, lacking many of the critical

protein-protein interaction domains seen in bilaterians (Richards and Degnan 2009;

Adamska et al. 2010) (Fig. 1D), and the corresponding genes have recently been

classified as "*APC-like*" (Adamska et al. 2010; Abbott and Näthke 2023) suggested that

the Axin and APC-like proteins in these early diverging taxa may not be part of the DC,

or that these proteins may be interacting in different ways to regulate DC function.

These observations provided a fundamental opportunity to better understand the

evolution of the regulation of a major metazoan signaling pathway and to reconstruct

the ancestral state of a critical regulatory node in the cnidarian-bilaterian last common

ancestor (LCA).

In this study we carried out a detailed functional analysis of the cWnt DC in the

cnidarian *Nematostella* to determine how it functions in the absence of critical domains

in Axin and APC required for bilaterian DC function. We show that *Nematostella* Axin

and APC-like proteins use novel protein-protein interactions to form a cWnt DC

suggesting that these proteins may have a level of promiscuity not reported in the

bilaterian cWnt DC. Additionally, using *in vitro* and *in vivo* protein-protein interaction

assays along with AI-enabled prediction we trace the evolutionary history of the critical

βcatBM on the bilaterian Axin protein from pre-bilaterian antecedents. We functionally

validate the regulation of a non-bilaterian DC function showing that cWnt signaling can

achieve deeply conserved developmental outcomes despite significant differences in the

regulation of the DC compared to bilaterians. Our results indicate that the key

scaffolding proteins in the cWnt DC in the cnidarian-bilaterian last common ancestor

were "promiscuous" which provides insight into the mechanisms leading to the

evolvability of the signaling pathways in bilaterians promoting the genesis of novel body

plans.

Results

The conserved β-catenin binding motif in bilaterian Axin proteins is not

detectable in non-bilaterian Axins

Structure-function studies in bilaterians have defined conserved regions on the

Axin protein that allow it to bind to APC, GSK-3β, and β-catenin in the DC thereby

promoting β-catenin phosphorylation and targeting it for degradation, maintaining the

pathway in an inactive state (Ikeda et al. 1998; Kishida et al. 1998; Fagotto et al. 1999;

Valenta et al. 2012; Stamos and Weis 2013; Pronobis et al. 2015; van Kappel and

Maurice 2017; Qiu et al. 2024) (Fig. 1C; Supplementary Table 1). In addition, Axin

proteins have a defined domain that binds to "activated" Disheveled when the cWnt

pathway is stimulated by upstream signals and a more poorly defined region that

interacts with CK-1α (Liu et al. 2002; MacDonald et al. 2009; Gammons and Bienz

2018; Qiu et al. 2024) (Fig. 1C; Supplementary Table 1). These Axin domains play a

critical role in recruiting components of the DC to regulate β-catenin stability (Xing et

al. 2003; Mendoza-Topaz et al. 2011; Stamos and Weis 2013; Qiu et al. 2024). A

previous bioinformatic analysis showed that the βcatBM is absent in *Amphimedon*

(Porifera) and *Nematostella* (Cnidaria) Axin proteins (Adamska et al. 2010). To

determine if this motif was also absent in the other two non-bilaterian taxa (Placozoa

and Ctenophora), the functional domains of Axin proteins in the four non-bilaterian

taxa and some selected bilaterian taxa were annotated using the webtool SMART

(Simple Modular Architecture Research Tool: http://smart.embl-heidelberg.de) and the

Axin proteins were aligned using Clustal Omega. This analysis showed that while the

βcatBM was identified in all bilaterian Axins, we could not identify this motif in Axin

proteins from the four non-bilaterian taxa (Supplementary Table 1). These observations

indicated that the βcatBM may have appeared in Axin after the divergence of the last

common ancestor of Bilateria and Cnidaria, or that the domain had diverged during

bilaterian evolution to render it undetectable in non-bilaterian Axin using the

bioinformatic tools we used. These observations also raised the possibility that in the

absence of a βcatBM non-bilaterian Axin would not be able to directly regulate β-catenin

stability in the cWnt DC.

NvAxin and NvAPC-like regulate cWnt signaling in Nematostella embryos

We have previously shown that a bilaterian Axin protein can disrupt early

development in *Nematostella* (Kumburegama et al. 2011). But to the best of our

knowledge, a functional role for a non-bilaterian Axin in regulating cWnt signaling has

not been examined. To study Axin function in *Nematostella*, we first examined the

expression pattern of *NvAxin* in eggs and early embryos using whole mount in situ

hybridization (WMISH) (Supplementary Fig. 1). WMISH revealed that *NvAxin* was

expressed ubiquitously at the egg, cleavage and blastula stages (Supplementary Fig. 1A-

C). However, starting at the early gastrula stage, *NvAxin* displayed an asymmetrical

expression pattern with the mRNA mostly restricted to the future oral pole and this

expression persisted through the planula stage (Supplementary Fig. 1D-F). This

expression pattern suggested a function for the NvAxin protein in regulating early

pattern formation in *Nematostella*.

We next asked if disrupting *NvAxin* function affected cWnt signaling during early

Nematostella development. It is now well established that cWnt signaling regulates cell

fates along the animal-vegetal axis in *Nematostella* (Wikramanayake et al. 2003; Lee et

al. 2007; Röttinger et al. 2012; Lebedeva et al. 2021; Wijesena et al. 2022). Blocking β-

catenin signaling using a dominant-negative form of NvTcf that prevents β-catenin from

interacting with cis-regulatory elements of cWnt target genes leads to the disruption of

normal gastrulation and downregulation of genes in cell types derived from animal half

blastomeres—the endomesoderm and pharynx (Röttinger et al. 2012) (Supplementary

Fig. 2). To determine if NvAxin regulates cWnt signaling and the nuclearization of β-

catenin in embryos, we co-expressed NvAxin-GFP with Nvβ-catenin-mCherry by mRNA

injection into zygotes and examined developing embryos for Nvβ-catenin-mCherry

expression at the blastula stage. Our results showed that while control embryos co-

injected with *GFP* and *Nvβ-catenin-mCherry* mRNA exhibited nuclearization of Nvβ-

catenin-mCherry protein on one side of the embryo, embryos co-expressing either

NvAxin-GFP or Axin-GFP from the sea urchin *Strongylocentrotus purpuratus* (SpAxin-

GFP) and Nvβ-catenin-mCherry did not display any nuclear localization of Nvβ-catenin-

mCherry (Fig. 2A-C). This result indicated that NvAxin overexpression blocks

nuclearization of Nvβ-catenin-mCherry, presumably by mediating phosphorylation and

destabilization of β-catenin within the DC.

Earlier studies had demonstrated that overexpression of sea urchin Axin, NvDvl-

DIX (a dominant-negative form of Dvl for cWnt signaling), a repressive form of β-

catenin and dominant-negative NvTcf produced a similar phenotype in *Nematostella*

embryos where the overexpression of the proteins blocked cWnt signaling, but primary

archenteron invagination still occurred (Lee et al. 2007; Kumburegama et al. 2011;

Röttinger et al. 2012). These studies noted that while the embryos in which cWnt was

downregulated initially formed an endodermal epithelium, this epithelium appeared to

disintegrate by the late gastrula stage (Kumburegama et al. 2011; Röttinger et al. 2012).

Consistent with these previous observations, overexpression of NvAxin in embryos by

mRNA injection into zygotes produced embryos that underwent primary archenteron

invagination, and these embryos failed to maintain a normal gut epithelium compared

to embryos expressing GFP (Fig. 2D-F).

We next investigated if manipulating Axin levels in *Nematostella* embryos had an

effect on endogenous gene expression consistent with a function for the protein in the

cWnt DC. We reasoned that overexpressing NvAxin would lead to a downregulation of

cWnt-dependent gene expression while knockdown or knockout of NvAxin would lead

to an increased expression of these genes. Consistent with these predictions, assaying

NvAxin- or SpAxin-overexpressing *Nematostella* embryos for gene expression in syn-

expression domains (Röttinger et al. 2012) (see Fig. 2G) using qPCR showed a

downregulation of genes expressed in the central domain and central ring (see Fig. 2G)

and an upregulation of genes expressed in the apical domain compared to control

embryos (Fig 2H,I). Conversely, knockdown of NvAxin using antisense morpholinos led

to an upregulation of genes expressed in the central domain, central ring, and external

ring while genes expressed in the apical domain were downregulated (Supplementary

Fig. 3A). To compare the changes in gene expression following knockdown of NvAxin

with the changes in gene expression caused by knockdown of NvAPC-like, we injected

antisense morpholinos against NvAPC-like into zygotes and used qPCR to analyze gene

expression in these embryos. Consistent with the effect we noted in NvAxin knockdown

embryos, we saw an upregulation of genes expressed in the central domain and the

central ring in NvAPC-like MO injected embryos (Supplementary Fig. 3B).

To use an additional strategy to assess the effect of Axin downregulation on cWnt

signaling in *Nematostella* embryos we carried out a CRISPR/Cas9-mediated NvAxin

knockout experiment. To compare the molecular phenotypes between *NvAPC-like* and

NvAxin knockout embryos we also designed an *NvAPC-like* single guide RNA (gRNA) to

introduce a mutation into *NvAPC-like* in *Nematostella* embryos. The *NvAxin* and

NvAPC gRNAs and Cas9 enzyme were injected into zygotes and animals were observed

when control Cas9-only injected animals were at the four-tentacle polyp stage. We

noted that *Nematostella* polyps developing from zygotes injected with *Axin* gRNA/Cas9

and *APC-like* gRNA/Cas9 had supernumerary tentacles (Kraus et al. 2016) (Fig. 3Aa,

Ab, Ac). Animals injected with *Axin* plus *APC-like* gRNA with Cas9 typically showed a

more severe phenotype with additional ectopic tentacles present throughout the animal

(Fig. 3Ad). The NvAxin gRNA-mediated genome editing was validated by sequencing

around the NvAxin gRNA excise sites in four single NvAxin gRNA-injected embryos and

one Cas9-injected embryo (Supplementary Fig. 4) and by using WMISH to detect

NvAxin expression in CRISPR/Cas9-mediated knockout embryos compared with Cas9-

injected control embryos at 24 hpf (Fig. 3B). WMISH showed that 81.9% of the control

embryos showed strong *NvAxin* expression at the animal pole; 13.7% showed detectable

but weaker asymmetrical *NvAxin* expression, and 4.4% showed no detectable signal

(n=204) (Fig. 3Ba). By contrast, 72.5% of NvAxin gRNA/Cas9-injected embryos showed

no detectable *Axin* expression, 24.4% showed weak *Axin* expression, and only 3.1%

showed a signal like that seen in the control (n=131) (Fig. 3Bb). The embryos with weak

NvAxin signals may be due to incomplete excision of NvAxin or the generation of

mosaic embryos where the gene was only excised in some nuclei. The percentage of

successful excisions using CRISPR/Cas9 noted in this study are consistent with other

studies that have used CRISPR/Cas9 to carry out gene editing in *Nematostella* embryos

(Kraus et al. 2016; Servetnick et al. 2017).

To evaluate the effect of CRISPR/Cas9-mediated *Axin* and *APC-like* knockout on

cWnt-dependent gene expression during early embryonic development, we used qPCR

analysis and the results revealed that the genes expressed in the central domain and

central ring were all upregulated indicating that knockout of *NvAxin* and *NvAPC-like*

led to increased cWnt signaling consistent with a function for NvAxin and NvAPC-like in

the cWnt DC (Fig. 3C,D). The qPCR analysis also revealed that the apical domain genes

Fz5 and *Six3* were downregulated, consistent with previous studies showing that global

upregulation of cWnt signaling leads to the downregulation of genes expressed in the

apical domain (Röttinger et al. 2012) (Fig. 3C,D). Notably, *NvWnt8* and *Nvlhx6* (a gene

expressed in the external ring) exhibited a stronger upregulation in both *NvAxin* gRNA

and *NvAPC-like* gRNA-injected embryos than other genes expressed at the animal pole,

suggesting that these genes are particularly sensitive to levels of cWnt signaling. Hence,

disruption of either one of the two core scaffold proteins in the *Nematostella* DC likely

resulted in the strong activation of *Wnt8* and *lhx6* expression. Alternatively, *NvWnt8*

and *Nvlhx6* might be repressed by cWnt outside their normal domain of expression and

are ectopically expressed when cWnt is downregulated. In sum, these results provided

strong evidence that Axin plays a crucial role in negatively regulating cWnt signaling in

Nematostella embryos and indicated that NvAxin has a role in the cWnt DC even

though it lacks a distinct β-catenin binding motif. Additionally, these experiments

showed that NvAPC-like regulates cWnt signaling even though it lacks most of the

domains shown be required for APC function in the DC in bilaterians. Our work also

corroborated the work of Kraus et al. (2016) who previously showed that NvAPC-like

regulates cWnt signaling in *Nematostella*.

Nematostella Axin overexpression disrupts cWnt signaling in sea urchin

embryos

We next sought to determine if *Nematostella* Axin could regulate cWnt signaling

in a bilaterian. Extensive work done in sea urchin embryos has shown that the cWnt

pathway plays a critical role in specifying and patterning the anterior-posterior (AP)

axis, and blocking β-catenin nuclearization in early embryos produces a unique

anteriorized phenotype hence making these embryos useful for assaying cWnt signaling

(Wikramanayake et al. 1998; Logan et al. 1999; Sun et al. 2021). When we expressed sea

urchin β-catenin-mCherry (Spβ-catenin-mCherry) in sea urchin embryos by mRNA

injection into zygotes, live imaging showed typical nuclearization of Spβ-catenin-

mCherry at the vegetal pole (Wikramanayake et al. 1998; Logan et al. 1999; Sun et al.

2021) (Fig. 4Aa). However, when Spβ-catenin-mCherry was co-expressed with NvAxin

or SpAxin in sea urchin embryos there was complete elimination of Spβ-catenin-

mCherry nuclearization (Fig. 4Ab, Ac). Moreover, consistent with previous observations

in sea urchin embryos where blocking Spβ-catenin nuclearization by SpAxin

overexpression led to severe anteriorization of sea urchin embryos (Sun et al. 2021) (Fig.

4Ba,Bb), overexpression of NvAxin in sea urchin embryos also led to a phenotype

indistinguishable from the phenotype generated from SpAxin overexpression (Fig. 4Bc).

Importantly, the anteriorized embryos were not able to recover even when control

animals were at the pluteus larval stage (Fig. 4Ba'-c') (Wikramanayake et al. 1998; Sun

et al. 2021). Consistent with the anteriorized phenotype, analysis of gene expression in

sea urchin embryos overexpressing NvAxin and SpAxin showed that all seven

endomesoderm gene markers tested were significantly downregulated, while two

neuroectoderm gene markers were upregulated (Fig. 4C, D). These experiments

demonstrated that NvAxin is a strong regulator of cWnt signaling in a bilaterian, even

though it lacks the conserved βcatBM.

The NvAxin GSK-3β binding domain is required for its function in the

cWnt pathway

In bilaterians, the βcatBM on Axin is critical for the protein to interact with and

target β-catenin for degradation during activation of cWnt signaling (Ikeda et al. 1998;

Xing et al. 2003; Stamos and Weis 2013). Bioinformatic analysis using SMART

indicated that NvAxin lacks the βcatBM but it has the APC- (RGS), GSK- (GID), and

Dvl- (DAX) binding domains, which are also found in the Axin proteins of other non-

bilaterians (Supplementary Table 1). Since our studies showed that NvAxin could

regulate β-catenin nuclearization in the sea urchin embryo, we used a rescue assay that

we had previously developed in sea urchins (Sun et al. 2021) to identify the domains on

NvAxin required for β-catenin regulation in the cWnt pathway. This rescue assay takes

advantage of the unique "posteriorized" phenotype produced in sea urchin embryos

when levels of Axin in the embryo are reduced using an anti-sea urchin Axin

morpholino, which can be rescued by overexpressing the wildtype Axin protein (Sun et

al. 2021) (Fig. 5A-C). We showed that while injection of a control MO has no effect on

embryo development, injections of an anti-sea urchin (*Lytechinus variegatus*, Lv) Axin

MO produces a posteriorized phenotype, and additionally, while *GFP* RNA control

injections have no effect on development, *NvAxin* RNA injection anteriorizes sea urchin

embryos (Fig. 5Ba-d'). To determine if NvAxin could rescue sea urchin embryos

posteriorized by injection of the LvAxin MO, we co-injected the LvAxin MO with full-

length *NvAxin* mRNA. Examination of these embryos showed that while the embryos

co-injected with the LvAxin MO and *GFP* mRNA were posteriorized as expected (Fig.

5Ca,a'), the embryos co-injected with the LvAxin MO and *NvAxin* mRNA were

completely rescued (Fig. 5Cb,b'). Similarly, co-injection of the LvAxin morpholino with

NvAxinΔRGS mRNA or *NvAxinΔDAX* mRNA efficiently rescued the posteriorized

phenotype induced by the LvAxin morpholino in sea urchins (Fig. 5Cc-d'). However, co-

injection of the LvAxin MO with *NvAxinΔGID* mRNA failed to rescue the phenotype

(Fig. 5Ce,e') indicating the requirement of the GSK-3β binding domain (GID) for

NvAxin to function in the cWnt DC. This result is consistent with our observations in

sea urchin embryos, where the SpAxinΔGID construct was also unable to rescue Axin

knockdown embryos (Sun et al. 2021). We conclude that the GID domain is likely the

most crucial domain of NvAxin for regulating β-catenin stability in the cWnt DC.

Previous studies in bilaterians have shown that overexpression of the Axin GID

peptide alone is sufficient to activate cWnt signaling (Hedgepeth et al. 1999; Sun et al.

2021). The underlying mechanism for this is still unclear but it has been suggested that

the GID domain can bind to endogenous GSK-3β and prevent it from interacting with

Axin (Hedgepeth et al. 1999; Chou et al. 2006; Sun et al. 2021). Our NvAxin rescue

experiments in sea urchins using Axin domain-deletion constructs clearly showed that

the GID domain is required for NvAxin's activity in the cWnt destruction complex (Fig.

5Ce,e'). GSK-3β binding to Axin is a critical step in assembling the DC in bilaterians. To

begin to assess if the NvAxin-GID (NvGID) domain was involved in regulating GSK-3β

in *Nematostella* we first overexpressed an NvGID-GFP construct in embryos by

microinjection of mRNA into zygotes. When control animals were at the polyp stage,

the NvGID-GFP overexpressing embryos had a morphology resembling *Nematostella*

embryos overexpressing an activated form of β-catenin (Röttinger et al. 2012) (Fig. 6A).

Analysis of germ layer-specific markers at the blastula stage using qPCR showed that

endomesoderm gene markers and pharyngeal markers were highly upregulated while

the ectoderm gene markers were downregulated (Fig. 6B). These results are consistent

with the notion that as seen in bilaterians, Axin-GID overexpression can activate cWnt

signaling and provides additional evidence that NvAxin scaffolds a critical DC

component similar to bilaterian Axin.

The residues within the GID domain mediating interaction between Axin and

GSK-3β are highly conserved in bilaterians and all four non-bilaterian taxa (Kremer et

al. 2010) (Supplementary Fig. 5). The conservation of the NvGID suggested that the

phenotype elicited by overexpression of the peptide in *Nematostella* was due to it

directly binding to NvGSK-3β and preventing NvGSK-3β from binding to NvAxin. To

validate this predicted interaction, we mixed NvGID-GFP and NvGSK-3β-mCherry

proteins made using *in vitro* transcription and translation (TnT) and

immunoprecipitated (IP) NvGID-GFP using anti-GFP antibodies. Western blot analysis

of the IP samples using anti-mCherry antibodies showed a clear interaction between

NvGID-GFP and NvGSK-3β-mCherry *in vitro* (Fig. 6C). These results demonstrated

that NvGID directly interacts with NvGSK-3β and further supports the notion that

NvAxin has an activity consistent with it being part of a DC in *Nematostella*.

It is important to note that the morphological and molecular phenotypes

generated from upregulating cWnt signaling by overexpression of NvGID, and NvAxin

and NvAPC-like knock down/knock out using morpholinos and CRISPR/Cas-9 show

some significant differences (compare Fig. 3 with Fig. 6). When NvAxin-GID was

overexpressed in *Nematostella* embryos, the embryos did not form a polyp stage like

controls, but rather had what appeared to be a hollow and expanded swollen appearance

(Fig. 6A). Gene expression analysis showed that these embryos had elevated expression

of genes expressed in the central domain, central ring and external ring and reduction in

expression of genes expressed in the apical domain. These embryos resemble the

phenotypes generated when ectopically activating cWnt signaling by overexpressing a

constitutively activate form of β-catenin (Röttinger et al. 2012). Both these methods are

likely to activate early cWnt signaling thus impacting early cell fate specification in

Nematostella embryos. It is likely that there is Axin and APC-like mRNA and protein in

the egg and early embryo and hence, when using morpholinos and CRISPR/Cas-9 to

knock down or knock out Axin and APC-like, these methods do not impact the

maternally loaded molecules. We suggest that the Axin and APC-like knock-

down/knock-out phenotypes are due to elevated cWnt signaling affecting oral-aboral

axis patterning, a relatively late event, and not early cell fate specification.

In vitro and in vivo protein-protein interaction assays indicate that the

Nematostella Destruction Complex proteins have binding activities

previously undefined in bilaterians

Previous studies in bilaterian taxa such as vertebrates and dipterans have shown

interactions between Axin-APC, Axin-β-catenin, APC-β-catenin, APC-GSK-3β and Axin-

GSK-3β *in vitro* and *in vivo* and in general, the protein binding interfaces for these

interactions have been well defined (Ikeda et al. 1998; Kishida et al. 1998; Dajani et al.

2003; Ha et al. 2004; Peterson-Nedry et al. 2008; Kremer et al. 2010; Roberts et al.

2011; Pronobis et al. 2015; Ranes et al. 2021). These studies have also demonstrated

that while there is a single defined β-catenin binding domain on bilaterian Axin, there

are multiple β-catenin binding domains on APC that have distinct sequence motifs

(Roberts et al. 2011; Pronobis et al. 2015; van Kappel and Maurice 2017; Pronobis et al.

2017). The experimental analyses that we have done in this study showed that both

NvAxin and NvAPC-like regulate cWnt in *Nematostella*. Hence, the lack of critical

domains required for cWnt DC activity in NvAxin and NvAPC-like prompted us to

investigate if these proteins could still form a functional DC perhaps through protein

binding domains previously uncharacterized in bilaterians. To explore this possibility,

we synthesized GFP-tagged NvAxin and SpAxin, and mCherry-tagged Nvβ-catenin, Spβ-

catenin, NvGSK-3β and SpGSK-3β using *in vitro* TnT and then mixed the various

protein pairs and used an anti-GFP antibody to IP the GFP tagged Axin proteins. We

then probed Western blots of the immunoprecipitated proteins with anti-mCherry

antibodies to evaluate any interaction between the protein pairs. As expected, this assay

revealed a robust interaction between sea urchin Axin-GFP and sea urchin β-catenin-

mCherry (Fig. 7A, Lane 5). We also noted a weaker but persistent interaction between

NvAxin and Nvβ-catenin and NvAxin and Spβ-catenin proteins in vitro (Fig. 7A, Lane

1,2). This was unexpected since NvAxin does not have the well-defined βcatBM seen in

bilaterian Axin (Fig. 1C).

We next asked if sea urchin Axin (which has the conserved βcatBM) could

interact with Nvβ-catenin. To our surprise, despite sea urchin Axin containing a

conserved βcatBM and a demonstrated ability to strongly interact with sea urchin β-

catenin *in vitro,* it only displayed a weaker interaction with Nvβ-catenin in this assay

(Compare Fig. 7A, Lane 4 and Lane 5). These results suggested that the Axin interacting

domains on Nvβ-catenin (the armadillo repeats) had diverged from the corresponding

domains on bilaterian β-catenin, and hence it was unable to interact strongly with a

bilaterian Axin. *Nematostella* β-catenin has all 12 ARM domains seen in bilaterians and

each of its ARM repeats has a high level of homology to corresponding ARM domains in

bilaterian β-catenin indicating a strong conservation over 700 million years of evolution

(Schneider et al. 2003). Hence, any changes in Nvβ-catenin protein that affects how it

interacts with a bilaterian Axin βcatBM would likely be more subtle amino acid changes

that affects binding affinity between the proteins. In sum, these experiments showed

that NvAxin can directly interact with *Nematostella* and sea urchin β-catenin, but based

on the *in vitro* assay, these interactions appeared considerably weaker than the

interaction between sea urchin Axin and sea urchin β-catenin.

We then investigated if NvAxin could directly interact with NvGSK-3β using the

in vitro pull-down assay. As predicted by the presence of a conserved GID on NvAxin

and our previous observation that the NvGID could interact with NvGSK-3β, Western

blotting of the IP products showed a clear interaction between NvAxin-GFP and NvGSK-

3β-mCherry (Fig. 7A, Lane 3). Likewise, our results showed that sea urchin Axin-GFP

interacted strongly with sea urchin GSK-3β-mCherry (Fig. 7A, Lane 6). It is notable that

the Axin GID domain is highly conserved, including in ctenophores (Supplementary Fig.

5), indicating that the Axin-GSK-3β interaction evolved over 700 mya.

In bilaterians, Axin and APC proteins heterodimerize to form the scaffold for the

β-catenin DC. Bilaterian APC/APC-like proteins have a complex domain structure that

allows them to interact with several proteins including Axin and β-catenin. NvAPC-like

is smaller in size than the vertebrate or dipteran proteins, and it lacks many of the

domains shown to bind to β-catenin and Axin in bilaterians (Fig. 1D; Roberts et al. 2011;

Pronobis et al. 2015; Pronobis et al. 2017; Ranes et al. 2021). When we analyzed

NvAPC-like domain structure using the webtool SMART (Simple Modular Architecture

Research Tool), we noted that it had seven conserved armadillo repeats and two poorly

characterized repeat regions at the N-terminus. NvAPC-like also lacked the SAMP

domains that are critical for bilaterian Axin-APC interaction (Pronobis et al. 2015;

Pronobis et al. 2017) (Fig. 1D). To determine if NvAPC-like (tagged with mCherry)

could directly interact with NvAxin, Nvβ-catenin and NvGSK (tagged with GFP), we

carried out the *in vitro* protein-protein interaction assay previously described. We

noted that NvAPC-like-mCherry robustly interacted with Nvβ-catenin-GFP in this assay

(Fig. 7B). We were also able to detect interactions between NvAPC-like-mCherry and

NvGSK-3β-GFP and NvAPC-like-mCherry and NvAxin-GFP *in vitro* (Fig. 7B).

The *in vitro* assay that we used in the experiments described above provides

insight into potential direct interactions between components of the cWnt DC in

Nematostella embryos. However, in those cases where we saw weak interactions *in*

vitro, it is possible that in the *in vivo* environment, post-translational modifications or

bridging by other proteins in the embryo lead to higher affinity interactions (Ha et al.

2004; Kimelman and Xu 2006; Pronobis et al. 2015; Ranes et al. 2021). To validate the

results that we saw in the *in vitro* experiments with an *in vivo* and *ex vivo* approach, we

adopted the split Click Beetle Green (CBG) luciferase system (Villalobos et al. 2010)

(Supplementary Fig. 6). Axin and β-catenin proteins from *Nematostella* and the sea

urchin *S. purpuratus* were tagged with split CBG luciferase (Villalobos et al. 2010). All

possible pairs of split-luciferase-tagged Axin and β-catenin proteins from the same or

different species, were expressed in *Nematostella* embryos by mRNA injection into

zygotes. The luminescence of split luciferase was measured in live embryos or in

embryo lysates. Normalized kinetic curves from embryonic lysates and live embryos are

shown in Supplemental Fig. 6. Consistent with the *in vitro* studies, SpAxin + Spβ-

catenin had the highest signal in both embryonic lysates and live embryos, while NvAxin

+ Spβ-catenin had clear but relatively lower signal (Supplemental Fig. 6A, B). However,

SpAxin did not appear to interact with Nvβ-catenin in either embryonic lysates or live

embryos (11 % and 10 % of SpAxin + Nvβ-catenin signal, respectively). This result is

inconsistent with the *in vitro* pull-down experiments where we noted a weak interaction

between SpAxin and Nvβ-catenin (Fig. 7, Lane 4). The reason for this discrepancy is not

known at this time. Interestingly, NvAxin + Nvβ-catenin yielded a relatively low signal

(27% of NvAxin + Nvβ-catenin intensity) in live embryos but virtually none in lysates

(13 %). These observations suggest that this interaction requires the presence of intact

cellular and molecular complexes that are lost in lysates. However, overall, the

interactions between NvAxin + Nvβ-catenin, NvAxin + Spβ-catenin, and SpAxin + Spβ-

catenin were consistent between the *in vitro* and *in vivo* approaches and gives credence

to the notion that the interaction between NvAxin and Nvβ-catenin is weaker than the

interaction between SpAxin and Spβ-catenin. Moreover, the *in vivo* and *ex vivo*

experiments confirmed that NvAxin and Nvβ-catenin, and SpAxin and Nvβ-catenin are

only capable of relatively weak interactions most likely discounting a major *in vivo* role

for bridging proteins or post-translational modifications to enhance these interactions.

AlphaFold software predicts novel protein-binding interfaces between

cWnt Destruction Complex components in Nematostella

The unexpected direct protein-protein interactions we noted in the *in vitro* assays

prompted us to use AlphaFold3 to determine if the AI-based program would be able to

predict the binding interfaces between the respective protein pairs. As expected,

AlphaFold analysis predicted the interactions between SpAxin-Spβ-catenin and

SpAxin-SpGSK-3β based on complementary domains seen in orthologs from other

bilaterian proteins (Supplementary Fig. 7A, B). Moreover, as expected from the

conserved domains, AlphaFold also predicted an interaction between NvAxin and

NvGSK-3β (Supplementary Fig. 7C). The AlphaFold analysis also indicated possible

interactions between NvAxin-Nvβ-catenin and NvAPC-like-Nvβ-catenin based on the

iPTM and PTM values (Supplementary Fig. 7D, E). This was unexpected because

NvAxin and NvAPC-like lack the β-catenin-interaction domains extensively

characterized in bilaterians (Roberts et al. 2011; Pronobis et al. 2015; Pronobis et al.

2017; Ranes et al. 2021). In contrast, even though the *in vitro* assays showed

interactions between NvAxin-NvAPC-like and NvAPC-like-NvGSK-3β, these

interactions were not supported by AlphaFold analysis. (Supplementary Fig. 7F, G).

This may be due to the NvAPC-like being a relatively large protein and in some such

cases AlphaFold is not as efficient in predicting protein-protein interactions (Bryant et

al. 2022; Shor and Schneidman-Duhovny 2024).

When we used AlphaFold3 and ChimeraX (Meng et al. 2023) to analyze

predicted and visualized interactions sites between the NvAPC-like and Nvβ-catenin,

we identified an α-helix interaction site within the ARM 708-753aa region (first ARM

domain) of NvAPC-like. This region was predicted to interact with Nvβ-catenin with

high iPTM and PTM values (Supplementary Fig. 7H). Notably, this ARM sequence is

conserved across *H. sapiens*, *D. melanogaster*, *S. purpuratus* and *N. vectensis* (amino

acid identity > 41%). We did not analyze the potential interactions of the other six

NvAPC-like ARM domains with Nvβ-catenin. Finally, we analyzed NvAPC-like protein

structure for a potential interaction with the NvAxin-RGS domain and AlphaFold

strongly predicted an interaction between the NvAxin-RGS and Repeat 1 of NvAPC-like

(Supplementary Fig. 7I) but not Repeat 2 (not shown).

The persistent, albeit weak, *in vitro* and *in vivo* interactions we noted between

NvAxin and Nvβ-catenin (Fig. 7, Supplementary Fig. 6) and the prediction of an

interaction between these proteins by AlphaFold3 analysis (Supplementary Fig. 7D)

were unexpected. AlphaFold 3 and ChimeraX software predicted two possible regions of

interaction between NvAxin and Nvβ-catenin, one located within the NvAxin-RGS

domain, and the other closer to the C-terminus of NvAxin (Figure 7C; Supplementary

Fig. 8A,B). Alignment of the predicted NvAxin-Nvβ-catenin interaction site on NvAxin

with the highly conserved βcatBM in bilaterian Axins using Clustal Omega and Jalview

showed some highly conserved amino acids between these sequences (Supplementary

Fig. 8F). The amino acid identity between the sequence in NvAxin-RGS and the second

sequence closer to the C-terminus, and the bilaterian βcatBM ranged from 20-26% and

11-28% respectively and we designated these sequences as "βcatBM-like". Significantly,

the leucines at positions 119 and 434 in the βcatBM-like sequences corresponds to a

leucine residue known to be essential for the interaction between bilaterian Axin and β-

catenin (Kremer et al. 2010) (Supplementary Fig. 8F). Additionally, Kremer et al.

(2010) identified a histidine residue at position 504 in *Drosophila* Axin as being critical

for β-catenin interaction with Axin. In *Nematostella* Axin, this corresponding residue is

missing, possibly accounting for the weak interaction between NvAxin and Nvβ-catenin.

We then examined Axin sequences from several other cnidarians to ask if similar

duplicated βcatBM-like sequences were present in the Axin proteins. Interestingly, the

predicted Axin-β-catenin binding sites are all located within the RGS domain

(Supplementary Fig. 9A) and C-terminal (Supplementary Fig. 9B) of Axin and are highly

conserved in each of these taxa. We also carried out AlphaFold and ChimeraX analysis

of predicted interactions between Axin and β-catenin in *Tricoplax adhaerens* (a

placozoan), *A. queenslandica* (a poriferan) and *Mnemiopsis leidyi* (a ctenophore). This

analysis predicted a single putative interaction site between Axin and β-catenin in these

species (Supplementary Fig. 8C,D,E,F). Alignment between the identified βcatBM-like

sequence on the ctenophore MlAxin-RGS and the bilaterian βcatBM did not reveal

conservation of the leucine residue; instead, an aspartic acid was identified

(Supplementary Fig. 8F). However, the identified βcatBM-like sequences on AqAxin-

RGS and TaAxin-RGS retained the conserved leucine associated with functional binding

(Kremer et al. 2010) (Supplementary Fig. 8F). The amino acid identities between the

βcatBM-like sequences in the RGS domain of TaAxin, AqAxin and MlAxin, and the

bilaterian βcatBM ranged from 16-33%, 12-25% and 21-28% respectively. We used

AlphaFold software to determine if the RGS domains of bilaterian Axins, including all

the bilaterian species listed in Supplementary Table 1, contained any putative β-catenin

interaction motifs, but we were not able to detect any such sequences (not shown).

Ctenophores are now widely believed to be the earliest-emerging metazoan group

(Dunn et al. 2008; Schultz et al. 2023). Consistent with this notion, the βcatBM-like

sequence on MlAxin identified using AlphaFold3 and ChimeraX lacked the conserved

leucine residue that is present in Axins from the other three non-bilaterian taxa. This

leucine along with a conserved histidine have been implicated as the two critical amino

acids involved in Axin-β-catenin interaction in bilaterians. The absence of both the

leucine and the histidine from the βcatBM-like sequence in ctenophore Axin indicated

that it would not be able to directly interact with ctenophore β-catenin. Consistent with

this idea MlAxin and Mlβ-catenin did not show any direct interactions when tested

using the *in vitro* TnT assay (Supplementary Fig. 8G). We next addressed if the

conserved leucine residues seen in the two βcatBM-like sequences in NvAxin were

responsible for the "weak" binding of NvAxin and Nvβ-catenin. We used site-directed

mutagenesis to change the conserved leucine in each of the βcatBM-like sequences in

NvAxin to aspartic acid to resemble the ctenophore βcatBM-like sequence and tested the

ability of these mutant NvAxin proteins to interact with Nvβ-catenin. Our analysis

showed that mutating the conserved leucine on each of the two βcatBM-like sequences

on NvAxin to aspartic acid reduced interaction with Nvβ-catenin compared to the wild-

type NvAxin βcatBM-like sequence, and mutating both leucines to aspartic acid

completely abolished interactions between NvAxin and Nvβ-catenin (Fig. 8C). In sum,

these results suggest that the bilaterian βcatBM evolved as a sequence on the Axin-RGS

with poor ability to interact with β-catenin and during pre-bilaterian evolution this

sequence acquired a critical leucine sequence that permitted weak interactions with β-

catenin.

Discussion

Studies on the evolution of metazoan signaling transduction pathways have been

impeded by the limited understanding of their roles in early emerging taxa and the

challenges to doing functional studies in these animals. We took advantage of the ability

to experimentally manipulate cWnt signaling in *Nematostella* embryos and the *in vitro*

assays we developed to identify direct protein interactions, coupled with the use of AI

tools to study the evolution of the cWnt destruction complex. We show that

Nematostella Axin and APC—which lack many of the domains required for bilaterian

DC function—can effectively regulate the stability of cytoplasmic β-catenin using

protein-protein interaction interfaces that to the best of our knowledge were previously

undefined in bilaterians. We also traced the evolution of the critical bilaterian Axin β-

catBM and provide evidence that it first appeared as a short β-catBM-like sequence

embedded in the Axin-RGS domain in the earliest branching metazoans. Our analysis

further indicates that during pre-bilaterian evolution the Axin β-catBM gained a leucine

residue that permitted weak direct interactions between Axin and β-catenin. The β-

catBM evolved in bilaterians through duplication of the β-catBM-like sequence in the

cnidarian-bilaterian last common ancestor (LCA) with the duplicated sequence inserted

towards the C-terminus of Axin. Our observations also suggest that in the Bilaterian

lineage the duplicated β-catBM-like sequence gained a higher affinity for β-catenin

through acquisition of additional amino acids, including a highly conserved histidine,

and the original β-catBM-like sequence was lost, most likely through drift. Overall, this

work has provided novel insight into the functional evolution of a major metazoan signal

transduction pathway.

NvAxin and NvAPC-like can assemble a functional destruction complex using novel protein binding interfaces

Axin and APC-like in cnidarians and other non-bilaterians lack the domains defined in bilaterians for assembling a functional cWnt DC. Hence, to investigate if the *Nematostella* DC could regulate cWnt signaling we carried out a series of experiments using readouts for cWnt signaling that we have previously established (Wikramanayake et al. 2003; Lee et al. 2007; Kumburegama et al. 2011; Röttinger et al. 2012; Sun et al. 2021). Our results showed that both NvAxin and NvAPC play a role in cWnt signaling, and moreover, we showed that NvAxin can also regulate cWnt signaling in sea urchin embryos (Fig. 4). We also used an *in vitro* protein-protein interaction assay to determine if NvAxin and NvAPC are able to directly interact with each other and with NvGSK-3β and Nvβ-catenin. This assay permitted us to identify previously undefined interactions between DC components. For example, our *in vitro* experiments showed that as predicted from NvAxin structure, this protein binds robustly to NvGSK-3β. This assay also allowed us to detect weak interactions between NvAxin and Nvβ-catenin that were unexpected because we were not able to identify a primary sequence that resembled the bilaterian β-catBM on NvAxin using SMART software. But using AI tools we were able to identify two β-catBM-like sequences on NvAxin. The *in vitro* assay also allowed us to show that NvAPC-like binds robustly to Nvβ-catenin and this interaction was also unexpected since NvAPC-like lacks the multiple domains that have been shown to bind β-catenin in bilaterians (Adamska et al. 2010; Roberts et al. 2011; Nelson and Näthke 2013; Pronobis et al. 2017; Abbott and Näthke 2023) (Fig. 1). Moreover, our *in vitro* experiments showed that NvAxin and NvAPC-like can interact directly even in the absence of the SAMP domains in NvAPC-like. Previous studies in *Drosophila* have

shown that DmAPC2 can interact with Axin in the absence of the SAMP domains

suggesting that some bilaterian APC proteins may also contain protein interaction

interfaces that have not been precisely defined previously (Roberts et al. 2011). We

suggest that in *Nematostella*, the interaction between NvAxin and NvAPC-like *in vivo*

would likely bring β-catenin and GSK-3β together for phosphorylation of the β-catenin

degron targeting the protein for degradation.

Other structural features of NvAPC-like indicated additional differences between

the *Nematostella* and bilaterian cWnt DCs in the molecular mechanisms targeting β-

catenin to the proteasome. Previous work has shown a clear role for NvAPC-like in

regulating cWnt signaling in *Nematostella* (Kraus et al. 2016) and our results

corroborated that work. However, the structure of NvAPC-like predicts that this protein

would be unable to target β-catenin to the proteasome. For example, APC proteins from

Drosophila and vertebrates have several 15 and 20 amino acid repeats that can bind to

β-catenin and in addition, these proteins have 2-3 SAMP domains that are involved in

dimerizing with Axin (Polakis 1995; Yang et al. 2006; Roberts et al. 2011; Nelson and

Näthke 2013; Pronobis et al. 2015; Abbott and Näthke 2023) (Fig. 1D). The second 20aa

repeat and an adjacent sequence are particularly significant because current models

propose that these domains are critical for transferring phosphorylated β-catenin from

Axin to APC prior to being moved to the proteasome for degradation (Kimelman and Xu

2006; Roberts et al. 2011; Pronobis et al. 2015). Strikingly, the NvAPC-like protein lacks

any detectable 15aa and 20aa repeats or the SAMP domains but has seven predicted

ARM repeats and two poorly characterized repeat regions (Fig. 1D). Hence, it appears

that NvAPC-like resembles a severely truncated bilaterian APC protein that would likely

be oncogenic in humans, and one that would not be predicted to regulate β-catenin

degradation (Polakis 1995; Kinzler and Vogelstein 1996; Fodde et al. 2001; Schneikert

and Behrens 2007). Despite these apparent structural deficits in cnidarian DC

scaffolding proteins experimental data in a number of cnidarian species has shown that

they can tightly regulate β-catenin protein stability in embryos. For example, when a β-

catenin-GFP fusion protein is expressed by mRNA injection in *Nematostella* and other

cnidarian embryos the protein is initially broadly expressed, but it is then cleared from

the vegetal pole blastomeres of the embryos and very tightly restricted to nuclei of the

animal pole blastomeres (Wikramanayake et al. 2003; Lee et al. 2007; Momose and

Houliston 2007; Momose et al. 2008; Kumburegama et al. 2011; Röttinger et al. 2012).

Hence, our observations raise important questions as to whether NvAPC-like has a role

in shuttling phosphorylated β-catenin to the proteasome, or if cnidarians have a

different mechanism to carry out this critical function. These are important questions

that need to addressed in future studies.

Another striking observation from our phylogenetic analysis of the Axin and

APC-like/APC proteins is the differential acquisition of domains between these proteins

during evolution. While we have not studied Axin domain evolution beyond the four

main domains (RGS, GID, βcatBM and DAX), Axin does not seem to have gained

additional domains other than the βcatBM with a high-affinity for β-catenin. In

contrast, the APC-like/APC proteins seem to have acquired many additional domains in

the bilaterian lineage likely through gene fusion (see Fig. 1) (Adamska et al. 2010;

Nelson and Näthke 2013; Abbott and Näthke 2023). This is particularly striking since

Axin and APC-like/APC most likely co-evolved in the cWnt DC and it would be a

reasonable prediction that changes in one protein would likely lead to changes in its

partner protein. Some of the additional domains acquired by bilaterian APC are likely

domains that are required for other functions of APC through its interactions with over

100 other previously identified proteins (Nelson and Näthke 2013). It is possible that

modeling the co-evolution of Axin and APC-like/APC may lead to some insights into the

evolutionary forces constraining the size of Axin while allowing the dramatic increase in

domain complexity and size in bilaterian APC-like/APC.

Reconstructing the evolution of the bilaterian Axin β-catenin binding motif

Our experimental work combined with the use of AI-enabled AlphaFold software

allowed us to propose a plausible mechanism for the evolution of the bilaterian Axin

βcatBM from pre-bilaterian antecedents. Insights into how the Axin βcatBM evolved

came from AlphaFold's identification of the NvAxin sequences underlying the weak but

consistent interactions we noticed between NvAxin and Nvβ-catenin *in vitro* and *in*

vivo. AlphaFold software predicted that two βcatBM-like sequences on NvAxin with

similarity to the conserved bilaterian βcatBM could potentially interact weakly with

Nvβ-catenin. But surprisingly one sequence was embedded within the NvAxin-RGS

domain and the other was closer to the C-terminus of Axin where the βcatBM is found in

bilaterian Axin (Fig. 8; Supplementary Fig. 8). Using AlphaFold we found a βcatBM-

like sequence embedded in the Axin-RGS domains in a placozoan, a sponge and three

ctenophores but we could not identify a second similar sequence closer to the C-

terminus in these taxa as we did in Axin from *Nematostella* and several other cnidarians

(Fig. 8; Supplementary Fig. 8,9). In addition, we could not identify a βcatBM-like

sequence in any bilaterian Axin-RGS. Within the two βcatBM-like sequences in NvAxin

and the single βcatBM-like sequence in TaAxin-RGS and AqAxin-RGS, there is a

conserved leucine residue, also seen in the bilaterian Axin βcatBM, that has been shown

to be critical for interaction between Axin and β-catenin in *Drosophila* (Kremer et al.

2010) (Fig. 8; Supplementary Fig. 8C,D,F). But the βcatBM-like sequences of NvAxin,

TaAxin and AqAxin lack a highly conserved histidine that has been shown to be critical

for Axin-β-catenin interactions in *Drosophila* (Kremer et al. 2010) (Supplementary Fig.

8C,D,F). Our analysis showed that the βcatBM-like sequence in the *Mnemiopsis* Axin-

RGS lacked both the conserved leucine and histidine residues required for interaction of

the *Drosophila* βcatBM with β-catenin and we showed that MlAxin and Mlβ-catenin

could not directly interact *in vitro* (Supplementary Fig. 8E,F).

It is now established that ctenophores are likely the earliest emerging metazoan

taxon (Dunn et al. 2008; Schultz et al. 2023). Hence, based on metazoan phylogeny we

can infer that the putative ancestral βcatBM-like sequence that was present in the Axin-

RGS domain in the "urmetazoan" acquired a leucine residue in the βcatBM-like

sequence prior to the split of the poriferans from a last common ancestor (LCA) to

placozoans, cnidarians and bilaterians. An alternative scenario is that the leucine was

present in the Axin βcatBM-like urmetazoan and was lost in ctenophore Axin.

Subsequently, after splitting from an LCA with cnidarians, bilaterian Axin acquired the

critical histidine residue, and perhaps others, in the βcatBM-like sequence which

provided an increased affinity between Axin and β-catenin. These data help reveal the

molecular changes that occurred prior to the stabilization of Wnt/β-catenin regulation

that appears in the majority of bilaterian clades.

Evolution of the bilaterian cWnt destruction complex

The cWnt pathway is one of only two signal transduction pathways regulating

development that are restricted to all metazoans (Richards and Degnan 2009; Babonis

and Martindale 2017; Richter et al. 2018) and it plays diverse roles in cellular and

developmental processes and in homeostasis (Liu et al. 2002; MacDonald et al. 2009;

Petersen and Reddien 2009; Loh et al. 2016). One highly conserved function of the

cWnt pathway is its role in establishing the first embryonic axis in bilaterians (anterior-

posterior) and cnidarians (oral-aboral) (Wikramanayake et al. 2003; Petersen and

Reddien 2009; Loh et al. 2016; Holstein 2022). We and others have also shown that the

cWnt pathway also plays a critical role in segregating germ layers in cnidarians and

invertebrate bilaterians (Wikramanayake et al. 1998; Logan et al. 1999; Imai et al. 2000;

Wikramanayake et al. 2003; Henry et al. 2008; Darras et al. 2011; McCauley et al. 2015;

Holstein 2022). Considering the conservation of the role of cWnt signaling in these

fundamental processes it is remarkable how divergent the regulation of the cWnt

destruction complex appears to be between early branching metazoans and Bilateria.

Below, we suggest a model for the evolution of the cWnt DC that argues that the cWnt

DC in *Nematostella* is not "divergent" but it resembles the ancestral cWnt DC in the

cnidarian-bilaterian LCA that was functionally sufficient to regulate cWnt signaling in

animals that had relatively simple body plans. But this ancestral DC had promiscuous

interactions that increased the evolvability of the cWnt pathway during its broad

deployment as bilaterians underwent a massive radiation to evolve taxa with complex

body plans.

We know very little about cWnt signaling in ctenophores (Pang et al. 2010;

Walters et al. 2025) and it has also been difficult to study embryonic development in two

other early emerging taxa, poriferans and placozoans, due to the challenges in accessing

their embryos. Hence, our current focus has been on reconstructing the ancestral cWnt

DC in the cnidarian-bilaterian LCA, since we can experimental manipulate

Nematostella development and there is accumulated knowledge on the developmental

roles of the cWnt pathway in cnidarian development. We propose that the ancestral DC

that was present in the LCA of Bilateria and Cnidaria was scaffolded by an APC-like

protein that bound β-catenin and Axin, and an Axin protein that bound GSK-3β and had

two βcatBM-like sites that had low-affinity β-catenin-binding activity. This DC was

capable of regulating β-catenin stability in the cWnt pathway in the cnidarian-bilaterian

LCA. Whether this DC was able to directly target β-catenin for degradation through the

proteasome is not known at this time but regardless of the mechanism, it had to be very

different from how β-catenin is targeted for degradation in the bilaterian cWnt DC

(Roberts et al. 2011; Ranes et al. 2021).

Following the split of the bilaterians from the cnidarian-bilaterian LCA, the

βcatBM-like site at the C-terminus positioned between the GID and DAX domains of

bilaterian Axin gained amino acids that increased its affinity for β-catenin possibly by

gaining the functionally significant histidine residue seen in the bilaterian βcatBM

(Kremer et al. 2010). Likely in parallel, APC-like gained a long carboxy-terminal region

that led to the acquisition of redundant β-catenin and Axin-binding sites. In addition,

some bilaterian APC proteins gained domains that are involved in processes such as

oligomerization and interaction with microtubules (Nelson and Näthke 2013; Abbott

and Näthke 2023) (Fig. 8). The mechanism for domain addition to APC is unknown but

it is likely due to gene fusion since that is considered to be the predominant mechanism

for domain evolution in eukaryotic proteins and it generally occurs at the N- or C-

termini of proteins (Patthy 1999; Liu et al. 2005; Weiner et al. 2006).

We were unable to identify a βcatBM-like sequence in the Axin-RGS domain of

all the bilaterians we examined (listed in Table 1). This sequence was most likely lost

through drift, since with the acquisition of APC domains that interacted with high

affinity with the Axin RGS domain of the βcatBM-like sequence embedded in the RGS

would not have been able to interact with β-catenin if it had an earlier function in

binding β-catenin. The overall consequence of these changes may have been the

evolution of higher-affinity protein-protein interactions within the DC that allowed a

more efficient capture of free β-catenin from the cytoplasm and an increased level of

redundancy for β-catenin sequestration that served to buffer this critical regulatory

module from misregulation (Peterson-Nedry et al. 2008; Roberts et al. 2011) (Fig. 8C).

This idea is consistent with modeling studies which have shown that signaling pathways

evolve increased levels of complexity under standard evolutionary conditions, and this

complexity also provides increased robustness (Soyer and Bonhoeffer 2006). It is also

likely that the increased number of interfaces provided by the bilaterian Axin and APC

proteins provided additional targets for context-dependent regulation of the cWnt DC

during the evolution of complex body plans in the Bilateria. In conclusion, we suggest

that the cnidarian-bilaterian LCA cWnt DC had promiscuous protein interactions that

may have increased the evolvability of the cWnt pathway during its broad deployment

during bilaterian development and evolution. We believe that the approach we have

taken in this study illustrate the power of leveraging biodiversity to gain insight into the

evolution of protein function and protein-protein interactions networks that regulate

evolution and development.

Materials and Methods

Animal care, spawning and culture

Adult *N. vectensis* were spawned and eggs prepared for microinjections as

previously described (Wijesena et al. 2022).

Adult *Lytechinus variegatus* (Lv) were obtained from Duke University Marine

Lab (Beaufort, NC) or Pelagic Corporation (Sugarloaf Key, FL). Spawning of animals

and collection and treatments of gametes was done as previously described (Byrum et al.

2009; Sun et al. 2021).

Cloning full length NvAxin and Reverse Transcriptase-PCR analysis

Full length *N. vectensis* Axin (NvAxin) cDNA was PCR-amplified from egg stage

using primers designed against the NvAxin sequence (NEMVEDRAFT_v1g182113). The

primer sequences were: NvAxin FL forward primer:

CGCGCGGATCCATGAGCGAAGAGTTGTGTAG, and NvAxin FL reverse primer:

CGACCATCGATTGAGTCTACTTTTTCTACTTTGC. Egg, cleavage, blastula, gastrula and

planula stage embryos were collected separately for RNA extraction using Trizol reagent

(cat# 15596026, Invitrogen). Approximately 1 μg of total RNA was used for cDNA

synthesis using SuperScript IV first strand synthesis system (cat# 18091200,

Invitrogen). For PCR, NvAxin cDNA was amplified from different stage embryos. The

primers used to amplify NvAxin for RT-PCR were: NvAxin FP:

ATGAGCGAAGAGTTGTGTAG, and NvAxin RP: CGTGCATCTTGCTCGTCGTG.

The PCR procedures followed the protocol provided with the Q5 high fidelity

DNA polymerase kit (cat# M0491, NEB) with thermal cycling conditions, 98°C for 30

seconds, 35 cycles with (98°C 10 seconds, 62°C 30 seconds, 72°C 20-30 seconds/kb)

and 72°C for 2 minutes. The wildtype NvAxin was inserted into the GFP-pCS2+ vector

by BamH1 and Cla1 restriction enzyme digestion and ligation with T4 DNA ligase.

Making NvAxin domain deletion constructs by site-directed mutagenesis

NvAxinΔRGS, NvAxinΔGID, NvAxinΔDAX constructs were made using the Q5 Site-

directed mutagenesis kit (E0554, NEB). Primers were designed to delete the RGS, GID

and DAX domains separately. The following primer sequences were used: NvAxin

ΔRGS forward primer, CAGGCTCACAGAGACACC, and NvAxin ΔRGS reverse primer,

GAGTTAGCCCATCGCTCG; NvAxin ΔGID forward primer,

TGAGAGTGAATATGGCCAG, and NvAxin ΔGID reverse primer,

GGATTTTTGGCTTCCTTTG; NvAxinΔDAX forward primer,

CAATCGATTCGAATTCAAG, and NvAxinΔDAX reverse primer,

TTGTGTTTCTCCTTTTTCC. To perform the PCR procedure, the NvAxin-GFP pCS2+

plasmid was mixed with pairs of primers, the Q5 hot start high-fidelity 2X master mix

and nuclease free water to perform PCR procedure, followed by kinase, ligase & Dpnl

(KLD) treatment.

Microinjection of morpholino antisense oligonucleotides and mRNAs

For knockdown experiments, morpholino antisense oligonucleotides to Axin or APC-like

(NvAxin MO or NvAPC MO) and a standard control morpholino (Control MO) to a

random nucleotide sequence (5'-CCTCTTACCTCAGTTACAATTTATA-3') were

synthesized at Gene Tools, LLC (Eugene, OR). The NvAxin MO:

TACACAACTCTTCGCTCATCTCACG or NvAPC-like MO:

ATCCTCTTCAAAATCGTCCATGTTG were designed to span the start codon of the Axin

or APC-like transcripts. Each MO was injected at a final concentration of 2 mM in 30-

40% glycerol and dextran rhodamine 0.2 μg/μl (D1824, Invitrogen).

For overexpression experiments in *N. vectensis* or *L. variegatus*, the NvAxin-

GFP, NvAxin ΔRGS-GFP, NvAxin ΔGID-GFP, NvAxin ΔDIX-GFP, SpAxin-GFP, Nvβ-

catenin-mCherry, and Spβ-catenin-mCherry, NvAxin GID-GFP, and a control GFP

mRNA were all cloned into the pCS2+ vector. The pCS2+ vectors containing the

respective cDNAs were linearized with NotI and mRNA was transcribed using the SP6

mMessage mMachine Kit (Ambion® ThermoFisher Scientific). Dominant negative

NvTcf (NvdnTCF) (Röttinger et al. 2012) was used for downregulating cWNT signaling .

This construct was in the Venus-pDEST expression vector. The dnTCF-Venus construct

was linearized with Acc65I and *in vitro* transcribed as previously described (Röttinger et

al. 2012). The mRNAs were mixed in 40% glycerol to a final concentration of 0.5 µg/µl.

GFP mRNA was overexpressed as a control at the same molar ratio as previously

described (Bince and Wikramanayake 2008).

For co-overexpression experiments in *N. vectensis* or *L. variegatus*, *NvAxin-*

GFP, *NvAxin ΔGID-GFP*, *SpAxin-GFP*, *Nvβ-catenin-mCherry*, *Spβ-catenin-mCherry*

mRNAs were injected at a final concentration of 0.5 µg/µl, 0.5 µg/µl, 0.5 µg/µl, 0.2

µg/µl, 0.2 µg/µl respectively in 40% glycerol.

For each experiment, more than 100 embryos were injected per MO or per

mRNA. The experiments were repeated at least three times for biological replication

(Sun et al. 2021; Wijesena et al. 2022).

CRISPR/Cas9 mediated knockouts and microinjections

To knock out *NvAxin* or *NvAPC-lik*e, the synthetic guide RNAs (sgRNA;

Synthego, Inc.) and Cas9 complex (RNP complex) were introduced into *N. vectensis*

embryos by microinjection into zygotes. The *NvAxin* or *NvAPC-like* single guide RNAs

were designed using CRISPRscan and CHOPCHOP as previously described (Salinas-

Saavedra et al. 2018). Each single guide RNA was resuspended to 50 μM and stored at

−80°C. Cas9 was obtained from PNA bio (Cat.# CP01). Cas9 was reconstituted to 5 mM

and was aliquoted for single use and stored at −80°C.

A mixture (2.5 μl) containing sgRNA (12 μM), Cas9 (0.6mM), 40% Glycerol,

dextran rhodamine (0.2 μg/μl) and nuclease free H_2O was prepared and 1 μl of mixture

was loaded into needles for microinjection. The same amount of Cas9 only was injected

as a control. *NvAxin* sgRNA was used to introduce mutations into *NvAxin*, and *NvAPC-*

like sgRNA was used to introduce mutations into *NvAPC-like*, and the mixture of

NvAxin and *NvAPC-like* sgRNAs were used to mutate both genes in single embryos.

Genomic DNA extraction from single embryos and validation of DNA editing by

CRISPR/Cas9

Single *NvAxin* gRNA injected embryos were collected into 0.2 ml PCR tubes and

lysed by adding 20 μl of extraction buffer for 100 μl buffer: 1 μl 1M Tris, pH8.0; 0.3 μl

Tween 20; 0.2 μl 0.5M EDTA; 5 μl 1M KCl; 0.3 μl NP40; 2.5 μl proteinase K (20

mg/ml); nuclease free water). The tubes with lysed embryos were incubated for 3 hr at

55°C in a thermal cycler with periodic vortexing and the reaction was inactivated by

heating the tubes at 98°C for 5min. For the PCR analysis of *NvAxin* gRNA injected

embryos, 5 μl of genomic DNA was used as a template to carry out the PCR reaction as

described above for NvAxin cloning and then the amplicon was gel extracted, purified

and sequenced at Eurofins Scientific. The following primers were used to generate the

NvAxin amplicons from genomic DNA from embryos edited with the *NvAxin* gRNA:

forward primer: ATGGCCAGTCTAAAAAGTCTATCC, and reverse primer:

CTGGTGGTTCAATGTCAAGAACGGTAT.

Quantitative PCR (qPCR)

To evaluate the effects of injected gRNAs, MO or mRNA on lineage-specific gene

expression, quantitative PCR (qPCR) was performed on *N. vectensis*, or *L. variegatus*

cDNA samples synthesized from total RNA that was extracted from gRNA-, morpholino-

or mRNA-injected embryos. The control and treated samples were collected at the

blastula stage; *N. vectensis* embryos were at 24 hpf at 17°C and *L. variegatus* embryos

were at 6 hpf at 25°C. Total RNA was isolated using the Trizol reagent solution (cat#

AM9738, Invitrogen). The cDNA was synthesized from 50 -100 ng total RNA of control

and experimental embryos respectively using the qScript cDNA Synthesis kit (Quanta

Biosciences, Beverly). The PerfeCTa SYBR Green FastMix (Quanta Biosciences,

Beverly) was used for assembling the qPCR reactions. Each experiment was repeated 3

times with separate batches of embryos and each PCR reaction was done in triplicate.

The expression of selected genes was analyzed using the delta delta Ct ($2^{-\Delta\Delta Ct}$)

method and using GADPH as a control as previously described (Sun et al. 2021).

Statistical significance was determined using the Unpaired t test without specific fold

change cut off. The values represent the fold change relative to control. All the *L.*

variegatus primer sequences were obtained from David McClay's laboratory at Duke

University.

Whole mount in situ hybridization (WMISH)

NvAxin expression during early development was determined using whole

mount in situ hybridization (WMISH).   The *NvAxin* in situ hybridization probe was

made using a pair of primers specifically targeting NvAxin: NvAxin-F: 5'

CGAAGGTAGAGCCGTATC 3', and NvAxin-R:

ATTTAGGTGACACTATAGGTCCGTCAGAGCATCACT, and was designed to contain an

Sp6 promoter for *in vitro* RNA synthesis. Embryo fixation and WMISH were performed

as previously described (Salinas-Saavedra et al. 2018). Embryos were fixed in 4%

paraformaldehyde with 0.2% glutaraldehyde in PTw (1XPBS+ 0.1%Tween-20) for 2 min

at room temperature, and then fixed in 4% paraformaldehyde in PTw for 1 hr at 4 °C. In

each experiment, a final concentration of 1 ng/μL anti-sense *NvAxin* probe was used

which was hybridized at 63°C for 2 days. *NvAxin* expression was detected using

NBT/BCIP as a substrate for the alkaline phosphatase conjugated to anti-digoxigenin

fab fragments (1:5000; Roche).

F-actin and nuclear staining

The morphology of *dnTcf-*, *NvAxin-*, *SpAxin-* and control *GFP*-RNA injected

embryos was examined using embryos collected at late gastrula stage (48 hpf at room

temperature) that were fixed and stained for F-actin and nuclear staining. Embryos

were collected and fixed as previously described with modifications (Salinas-Saavedra et

al. 2018). Briefly, embryos were first fixed in fix1 (100 mM HEPES pH 6.9; 0.05 M

EGTA; 5 mM MgSO4; 200 mM NaCl; 1x PBS; 3.7% Formaldehyde; 0.2%

Glutaraldehyde; 0.5% Triton X-100; and autoclaved ultrapure water) for 2-3 minutes

and then fixed in fix2 (100 mM HEPES pH 6.9; 0.05 M EGTA; 5 mM MgSO4; 200 mM

NaCl; 1x PBS; 3.7% Formaldehyde; 0.05% Glutaraldehyde; 0.5% Triton X-100; and

autoclaved ultrapure water) for one hour at room temperature on the rocker. Fixed

embryos were rinsed in PBT (X% Tween 20 in 1 x PBS, pH7.4) for 30 mins x 4 times.

Embryos were blocked with 5% normal sheep in PBT buffer for 1 hour at room

temperature on the rocker then rinsed with PBT for 30 mins x 4 times. For the F-actin

staining, embryos were stained with Phalloidin in PBT for 1 hour and then rinsed with

PBT for 10 min x 2 times. For nuclear staining, embryos were incubated in DAPI in PBT

for 1 hour and then rinsed in 1xPBS (PH 7.4) for 10 min x 3 times. Following these

washes, the embryos were dehydrated using a gradient of isopropanol (50, 75, 90, 100%,

100%). The embryos were then cleared with 1:2 benzyl benzoate: benzyl alcohol

(BB:BA) for 1 min x 2 times and mounted in BB:BA for imaging using scanning confocal

microscopy.

In vitro Transcription and Translation (TNT) and Western blot analysis

All GFP, mCherry, Flag and their fused constructs were in vitro transcribed and

translated using the TnT® Quick Coupled Transcription/Translation System with SP6

RNA polymerase promoter respectively (cat # L2080, Promega). Plasmids (at 1 μg

each) containing the respective cDNAs coding for the various fusion proteins were

incubated with TNT quick master mix at 30°C for 90 minutes. Protein expression was

examined using Western blot analysis with anti-GFP (cat# sc9996), anti-mCherry

antibodies (PA5-34974) or anti-flag (cat# 740001) respectively.

Protein-protein interaction was tested using ChromoTek GFP-Trap® Magnetic

Agarose following instructions to perform immunoprecipitation (IP) assay (cat # gtma,

Proteintech Group, INC., Rosemont, IL). Denatured IP protein samples were examined

on an 8% SDS-PAGE gel followed by Western blot analysis. Western blot was done as

previously described (Sun et al. 2024).

Split luciferase assay

N- and C-terminal fragments of click beetle green (CBG) split luciferase,

including polypeptide linkers, were amplified and blunt-end-ligated into pSPE3-RfA

vectors in a single restriction-ligation reaction. Coding sequences for *N. vectensis* and S.

purpuratus Axin and β-catenin were cloned into pENTR using D-TOPO cloning kit

(Invitrogen K240020) and recombined into CBG-tagged destination vectors via

Gateway cloning (Invitrogen 11791020). Nanoluciferase (NanoLuc) was similarly cloned

and recombined into pSPE3-RfA.

Recombined Axin and β-catenin constructs were linearized, purified, and used for

in vitro mRNA synthesis (Invitrogen AM1348). Equimolar CBG-tagged *Axin* and *β-*

catenin mRNA mixtures, along with NanoLuc mRNA and fluorescent dextran, were

microinjected into *Nematostella* zygotes. At 28 hours post-fertilization(hpf), live

embryos were collected, and luminescence was measured after adding D-luciferin

substrate (Goldbio LUCNA-100). Embryos were then lysed with ONE-Glo EX

Luciferase Assay Reagent (Promega N1610), and further measurements were taken.

NanoDLR Stop & Glo Reagent (Promega N1610) was used to stop the CBG reaction and

initiate NanoLuc luminescence. NanoLuc signal was used to normalize protein levels

and CBG intensities. Average NanoLuc signal from 15-20 minutes after the addition of

substrate was used to determine ratios of protein levels. Average relative intensities

across timepoints were calculated for each construct combination.

Primers:

1021

Name	Sequence
linker-N-luc Fw	CAGATCTCGTACGCGTCCC
N-luc Rev	TTAGCCGTCGTCGTCGAT
Kozak-CBG-C-linker Fw	CAGAAAAAATGAAGGGGTTATGTCAATAACGTTG
C-luc-linker Rev	TCCGCCCCGGGACG
NvAxin Fw	CACCATGAGCGAAGAGTTGTGTAG
NvAxin Rev	TGAGTCTACTTTTTCTACTTTGC
NvBeta-catenin Fw	CACCATGGGGATGCAGCAGGG
NvBeta-catenin-STOP Rev	TCAGAGGTCTGTGTCATACAGAG
SpAxin Fw	CACCATGAGTCTAGAAGTGTATAGG
SpAxin Rev	GTGATCATCGACAGATTC
SpBeta-catenin Fw	CACCATGGAGCAGGCAAGGTAC
SpBeta-catenin-STOP Rev	TTACAGATCGGTATCAAAGAACG
NanoLuc Fw	CACCATGGTCTTCACACTCG
NanoLuc Rev	TTACGCCAGAATGCG

1022

1023

AlphaFold 3 Protein-Protein interaction prediction

Computational methods were used to predict protein-protein interactions among

the components of the β-catenin destruction complex in *S. purpuratus* (sea urchin) and

Nematostella. Protein sequences for the individual components of the β-catenin

destruction complex were used as input in AlphaFold 3. The prediction accuracy was

assessed using the predicted align error (PAE) and confidence scores such as the

predicted TM-score (PTM) and inter-chain predicted TM-score (iPTM), which are

indicators of the reliability of the predicted interactions. A PTM score above 0.5

indicates that the overall predicted fold for the complex may resemble the true

structure. iPTM Values greater than 0.8 represent confident, high-quality predictions,

while values below 0.5 suggest likely failed predictions. ipTM scores between 0.5 and

0.8 fall into an intermediate zone where predictions may be correct or incorrect[21].

AlphaFold 3 allowed for the visualization of possible binding interfaces and the

determination of whether specific regions, such as the β-catenin binding domains, could

facilitate protein-protein interactions, even in cases where canonical domains were not

evident.

1040

1041

1042   References

1043   Abbott J, Näthke IS. 2023. The adenomatous polyposis coli protein 30 years on. *Semin Cell Dev*
1044   *Biol* 150–151:28–34.

1045   Adamska M, Larroux C, Adamski M, Green K, Lovas E, Koop D, Richards GS, Zwafink C,
1046   Degnan BM. 2010. Structure and expression of conserved Wnt pathway components in
1047   the demosponge Amphimedon queenslandica. *Evol Dev* 12:494–518.

1048   Babonis LS, Martindale MQ. 2017. Phylogenetic evidence for the modular evolution of
1049   metazoan signalling pathways. *Philosophical Transactions of the Royal Society B:*
1050   *Biological Sciences* 372:20150477.

1051   Bince JM, Wikramanayake AH. 2008. Functional analysis of Wnt signaling in the early sea
1052   urchin embryo using mRNA microinjection. *Methods Mol Biol* 469:213–222.

1053   Bryant P, Pozzati G, Elofsson A. 2022. Improved prediction of protein-protein interactions using
1054   AlphaFold2. *Nat Commun* 13:1265.

1055   Byrum CA, Xu R, Bince JM, McClay DR, Wikramanayake AH. 2009. Blocking Dishevelled
1056   signaling in the noncanonical Wnt pathway in sea urchins disrupts endoderm formation
1057   and spiculogenesis, but not secondary mesoderm formation. *Dev Dyn* 238:1649–1665.

1058   Chou H-Y, Howng S-L, Cheng T-S, Hsiao Y-L, Lieu A-S, Loh J-K, Hwang S-L, Lin C-C, Hsu
1059   C-M, Wang C, et al. 2006. GSKIP Is Homologous to the Axin GSK3β Interaction
1060   Domain and Functions as a Negative Regulator of GSK3β. *Biochemistry* 45:11379–
1061   11389.

1062   Dajani R, Fraser E, Roe SM, Yeo M, Good VM, Thompson V, Dale TC, Pearl LH. 2003.
1063   Structural basis for recruitment of glycogen synthase kinase 3β to the axin–APC scaffold
1064   complex. *EMBO J* 22:494–501.

1065   Darras S, Gerhart J, Terasaki M, Kirschner M, Lowe CJ. 2011. β-catenin specifies the
1066   endomesoderm and defines the posterior organizer of the hemichordate Saccoglossus
1067   kowalevskii. *Development* 138:959–970.

1068   Dunn CW, Hejnol A, Matus DQ, Pang K, Browne WE, Smith SA, Seaver E, Rouse GW, Obst
1069   M, Edgecombe GD, et al. 2008. Broad phylogenomic sampling improves resolution of
1070   the animal tree of life. *Nature* 452:745–749.

1071   Fagotto F, Jho E, Zeng L, Kurth T, Joos T, Kaufmann C, Costantini F. 1999. Domains of Axin
1072   Involved in Protein–Protein Interactions, Wnt Pathway Inhibition, and Intracellular
1073   Localization. *J Cell Biol* 145:741–756.

1074   Fodde R, Smits R, Clevers H. 2001. APC, Signal transduction and genetic instability in
1075   colorectal cancer. *Nat Rev Cancer* 1:55–67.

Gammons M, Bienz M. 2018. Multiprotein complexes governing Wnt signal transduction. *Curr*
Opin Cell Biol 51:42–49.

Gerhart J. 1999. 1998 warkany lecture: Signaling pathways in development. *Teratology* 60:226–
239.

Ha N-C, Tonozuka T, Stamos JL, Choi H-J, Weis WI. 2004. Mechanism of phosphorylation-
dependent binding of APC to beta-catenin and its role in beta-catenin degradation. *Mol*
Cell 15:511–521.

Hedgepeth CM, Deardorff MA, Rankin K, Klein PS. 1999. Regulation of glycogen synthase
kinase 3beta and downstream Wnt signaling by axin. *Mol Cell Biol* 19:7147–7157.

Henry JQ, Perry KJ, Wever J, Seaver E, Martindale MQ. 2008. β-Catenin is required for the
establishment of vegetal embryonic fates in the nemertean, *Cerebratulus lacteus*.
Developmental Biology 317:368–379.

Holstein TW. 2022. The role of cnidarian developmental biology in unraveling axis formation
and Wnt signaling. *Dev Biol* 487:74–98.

Holzem M, Boutros M, Holstein TW. 2024. The origin and evolution of Wnt signalling. *Nat Rev*
Genet 25:500–512.

Houliston E, Leclère L, Munro C, Copley RR, Momose T. 2022. Past, present and future of
Clytia hemisphaerica as a laboratory jellyfish. In: Goldstein B, Srivastava M, editors.
Current Topics in Developmental Biology. Vol. 147. Emerging Model Systems in
Developmental Biology. Academic Press. p. 121–151. Available from:
https://www.sciencedirect.com/science/article/pii/S0070215321001046

Ikeda S, Kishida S, Yamamoto H, Murai H, Koyama S, Kikuchi A. 1998. Axin, a negative
regulator of the Wnt signaling pathway, forms a complex with GSK-3beta and beta-
catenin and promotes GSK-3beta-dependent phosphorylation of beta-catenin. *EMBO J*
17:1371–1384.

Imai K, Takada N, Satoh N, Satou Y. 2000. β-catenin mediates the specification of endoderm
cells in ascidian embryos. *Development* 127:3009–3020.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates
R, Žídek A, Potapenko A, et al. 2021. Highly accurate protein structure prediction with
AlphaFold. *Nature* 596:583–589.

van Kappel EC, Maurice MM. 2017. Molecular regulation and pharmacological targeting of the
β-catenin destruction complex. *Br J Pharmacol* 174:4575–4588.

Kimelman D, Xu W. 2006. beta-catenin destruction complex: insights and questions from a
structural perspective. *Oncogene* 25:7482–7491.

Kinzler KW, Vogelstein B. 1996. Lessons from hereditary colorectal cancer. *Cell* 87:159–170.

Kishida S, Yamamoto H, Ikeda S, Kishida M, Sakamoto I, Koyama S, Kikuchi A. 1998. Axin, a
Negative Regulator of the Wnt Signaling Pathway, Directly Interacts with Adenomatous
Polyposis Coli and Regulates the Stabilization of β-Catenin. *Journal of Biological*
Chemistry 273:10823–10826.

Kohler EM, Derungs A, Daum G, Behrens J, Schneikert J. 2008. Functional definition of the
mutation cluster region of adenomatous polyposis coli in colorectal tumours. *Hum Mol*
Genet 17:1978–1987.

Kraus Y, Aman A, Technau U, Genikhovich G. 2016. Pre-bilaterian origin of the blastoporal
axial organizer. *Nat Commun* 7:11694.

Kremer SA, Erdeniz N, Peterson-Nedry W, Swanson EA, Wehrli M. 2010. In vivo analysis in
Drosophila reveals differential requirements of contact residues in Axin for interactions
with GSK3beta or beta-catenin. *Dev Biol* 337:110–123.

Kumburegama S, Wijesena N, Xu R, Wikramanayake AH. 2011. Strabismus-mediated primary
archenteron invagination is uncoupled from Wnt/β-catenin-dependent endoderm cell fate
specification in Nematostella vectensis (Anthozoa, Cnidaria): Implications for the
evolution of gastrulation. *EvoDevo* 2:2.

Layden MJ, Rentzsch F, Röttinger E. 2016. The rise of the starlet sea anemone Nematostella
vectensis as a model system to investigate development and regeneration. *Wiley*
Interdiscip Rev Dev Biol 5:408–428.

Lebedeva T, Aman AJ, Graf T, Niedermoser I, Zimmermann B, Kraus Y, Schatka M, Demilly
A, Technau U, Genikhovich G. 2021. Cnidarian-bilaterian comparison reveals the
ancestral regulatory logic of the β-catenin dependent axial patterning. *Nat Commun*
12:4032.

Lee PN, Kumburegama S, Marlow HQ, Martindale MQ, Wikramanayake AH. 2007.
Asymmetric developmental potential along the animal-vegetal axis in the anthozoan
cnidarian, Nematostella vectensis, is mediated by Dishevelled. *Dev Biol* 310:169–186.

Liu C, Li Y, Semenov M, Han C, Baeg GH, Tan Y, Zhang Z, Lin X, He X. 2002. Control of
beta-catenin phosphorylation/degradation by a dual-kinase mechanism. *Cell* 108:837–
847.

Liu J, Xiao Q, Xiao J, Niu C, Li Y, Zhang X, Zhou Z, Shu G, Yin G. 2022. Wnt/β-catenin
signalling: function, biological mechanisms, and therapeutic opportunities. *Sig Transduct*
Target Ther 7:1–23.

Liu M, Walch H, Wu S, Grigoriev A. 2005. Significant expansion of exon-bordering protein
domains during animal proteome evolution. *Nucleic Acids Research* 33:95–105.

Logan CY, Miller JR, Ferkowicz MJ, McClay DR. 1999. Nuclear β-catenin is required to specify
vegetal cell fates in the sea urchin embryo. *Development* 126:345–357.

Loh KM, van Amerongen R, Nusse R. 2016. Generating Cellular Diversity and Spatial Form:
Wnt Signaling and the Evolution of Multicellular Animals. *Dev Cell* 38:643–655.

MacDonald BT, Tamai K, He X. 2009. Wnt/β-Catenin Signaling: Components, Mechanisms,
and Diseases. *Developmental Cell* 17:9–26.

McCauley BS, Akyar E, Saad HR, Hinman VF. 2015. Dose-dependent nuclear β-catenin
response segregates endomesoderm along the sea star primary axis. *Development*
142:207–217.

Mendoza-Topaz C, Mieszczanek J, Bienz M. 2011. The Adenomatous polyposis coli tumour
suppressor is essential for Axin complex assembly and function and opposes Axin's
interaction with Dishevelled. *Open Biol* 1:110013.

Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, Ferrin TE. 2023.
UCSF ChimeraX: Tools for structure building and analysis. *Protein Sci* 32:e4792.

Momose T, Derelle R, Houliston E. 2008. A maternally localised Wnt ligand required for axial
patterning in the cnidarian Clytia hemisphaerica. *Development* 135:2105–2113.

Momose T, Houliston E. 2007. Two oppositely localised frizzled RNAs as axis determinants in a
cnidarian embryo. *PLoS Biol* 5:e70.

Nelson S, Näthke IS. 2013. Interactions and functions of the adenomatous polyposis coli (APC)
protein at a glance. *Journal of Cell Science* 126:873–877.

Pang K, Ryan JF, Mullikin JC, Baxevanis AD, Martindale MQ. 2010. Genomic insights into Wnt
signaling in an early diverging metazoan, the ctenophore Mnemiopsis leidyi. *EvoDevo*
1:10.

Patthy L. 1999. Genome evolution and the evolution of exon-shuffling--a review. *Gene* 238:103–
114.

Petersen CP, Reddien PW. 2009. Wnt signaling and the polarity of the primary body axis. *Cell*
139:1056–1068.

Peterson-Nedry W, Erdeniz N, Kremer S, Yu J, Baig-Lewis S, Wehrli M. 2008. Unexpectedly
robust assembly of the Axin destruction complex regulates Wnt/Wg signaling in
Drosophila as revealed by analysis in vivo. *Dev Biol* 320:226–241.

Pires-daSilva A, Sommer RJ. 2003. The evolution of signalling pathways in animal
development. *Nat Rev Genet* 4:39–49.

Polakis P. 1995. Mutations in the APC gene and their implications for protein structure and
function. *Current Opinion in Genetics & Development* 5:66–71.

1179 Pronobis MI, Deuitch N, Posham V, Mimori-Kiyosue Y, Peifer M. 2017. Reconstituting
1180   regulation of the canonical Wnt pathway by engineering a minimal β-catenin destruction
1181   machine. *Mol Biol Cell* 28:41–53.

1182 Pronobis MI, Rusan NM, Peifer M. 2015. A novel GSK3-regulated APC:Axin interaction
1183   regulates Wnt signaling by driving a catalytic cycle of efficient βcatenin destruction.
1184   *eLife* [Internet]. Available from: https://elifesciences.org/articles/08022

1185 Qiu L, Sun Y, Ning H, Chen G, Zhao W, Gao Y. 2024. The scaffold protein AXIN1: gene
1186   ontology, signal network, and physiological function. *Cell Communication and Signaling*
1187   22:77.

1188 Ranes M, Zaleska M, Sakalas S, Knight R, Guettler S. 2021. Reconstitution of the destruction
1189   complex defines roles of AXIN polymers and APC in β-catenin capture, phosphorylation,
1190   and ubiquitylation. *Mol Cell* 81:3246-3261.e11.

1191 Richards GS, Degnan BM. 2009. The dawn of developmental signaling in the metazoa. *Cold*
1192   *Spring Harb Symp Quant Biol* 74:81–90.

1193 Richter DJ, Fozouni P, Eisen MB, King N. 2018. Gene family innovation, conservation and loss
1194   on the animal stem lineage.Telford MJ, editor. *eLife* 7:e34226.

1195 Richter DJ, King N. 2013. The genomic and cellular foundations of animal origins. *Annu Rev*
1196   *Genet* 47:509–537.

1197 Roberts DM, Pronobis MI, Poulton JS, Waldmann JD, Stephenson EM, Hanna S, Peifer M.
1198   2011. Deconstructing the ßcatenin destruction complex: mechanistic roles for the tumor
1199   suppressor APC in regulating Wnt signaling. *Mol Biol Cell* 22:1845–1863.

1200 Rokas A. 2008. The origins of multicellularity and the early history of the genetic toolkit for
1201   animal development. *Annu Rev Genet* 42:235–251.

1202 Ros-Rocher N, Pérez-Posada A, Leger MM, Ruiz-Trillo I. 2021. The origin of animals: an
1203   ancestral reconstruction of the unicellular-to-multicellular transition. *Open Biol*
1204   11:200359.

1205 Röttinger E, Dahlin P, Martindale MQ. 2012. A Framework for the Establishment of a Cnidarian
1206   Gene Regulatory Network for "Endomesoderm" Specification: The Inputs of ß-
1207   Catenin/TCF Signaling. *PLOS Genetics* 8:e1003164.

1208 Salinas-Saavedra M, Rock AQ, Martindale MQ. 2018. Germ layer-specific regulation of cell
1209   polarity and adhesion gives insight into the evolution of mesoderm. *Elife* 7:e36740.

1210 Schneider SQ, Finnerty JR, Martindale MQ. 2003. Protein evolution: structure-function
1211   relationships of the oncogene beta-catenin in the evolution of multicellular animals. *J Exp*
1212   *Zool B Mol Dev Evol* 295:25–44.

Schneikert J, Behrens J. 2007. The canonical Wnt signalling pathway and its APC partner in
colon cancer development. *Gut* 56:417–425.

Schultz DT, Haddock SHD, Bredeson JV, Green RE, Simakov O, Rokhsar DS. 2023. Ancient
gene linkages support ctenophores as sister to other animals. *Nature* 618:110–117.

Servetnick MD, Steinworth B, Babonis LS, Simmons D, Salinas-Saavedra M, Martindale MQ.
2017. Cas9-mediated excision of Nematostella brachyury disrupts endoderm
development, pharynx formation and oral-aboral patterning. *Development* 144:2951–
2960.

Shor B, Schneidman-Duhovny D. 2024. CombFold: predicting structures of large protein
assemblies using a combinatorial assembly algorithm and AlphaFold2. *Nat Methods*
21:477–487.

Soyer OS, Bonhoeffer S. 2006. Evolution of complexity in signaling pathways. *Proceedings of*
the National Academy of Sciences 103:16337–16342.

Stamos JL, Weis WI. 2013. The β-catenin destruction complex. *Cold Spring Harb Perspect Biol*
5:a007898.

Sun H, Peng CJ, Wang L, Feng H, Wikramanayake AH. 2021. An early global role for Axin is
required for correct patterning of the anterior-posterior axis in the sea urchin embryo.
Development 148:dev191197.

Sun H, Shami Shah A, Chiu D-C, Bonfini A, Buchon N, Baskin JM. 2024. Wnt/β-catenin
signaling within multiple cell types dependent upon kramer regulates Drosophila
intestinal stem cell proliferation. *iScience* 27:110113.

Valenta T, Hausmann G, Basler K. 2012. The many faces and functions of β-catenin. *EMBO J*
31:2714–2736.

Villalobos V, Naik S, Bruinsma M, Dothager RS, Pan M-H, Samrakandi M, Moss B, Elhammali
A, Piwnica-Worms D. 2010. Dual-Color Click Beetle Luciferase Heteroprotein Fragment
Complementation Assays. *Chem Biol* 17:1018–1029.

Walters BM, Guttieres LJ, Goëb M, Marjenberg SJ, Martindale MQ, Wikramanayake AH. 2024.
β-catenin localization in the ctenophore Mnemiopsis leidyi suggests an ancestral role in
cell adhesion and nuclear function. :2024.08.29.610370. Available from:
https://www.biorxiv.org/content/10.1101/2024.08.29.610370v1

Weiner J, Beaussart F, Bornberg-Bauer E. 2006. Domain deletions and substitutions in the
modular protein evolution. *FEBS J* 273:2037–2047.

Wijesena N, Sun H, Kumburegama S, Wikramanayake AH. 2022. Distinct Frizzled receptors
independently mediate endomesoderm specification and primary archenteron
invagination during gastrulation in *Nematostella*. *Developmental Biology* 481:215–225.

Wikramanayake AH, Hong M, Lee PN, Pang K, Byrum CA, Bince JM, Xu R, Martindale MQ.
2003. An ancient role for nuclear β-catenin in the evolution of axial polarity and germ
layer segregation. *Nature* 426:446–450.

Wikramanayake AH, Huang L, Klein WH. 1998. β-Catenin is essential for patterning the
maternally specified animal-vegetal axis in the sea urchin embryo. *Proceedings of the*
National Academy of Sciences 95:9343–9348.

Xing Y, Clements WK, Kimelman D, Xu W. 2003. Crystal structure of a β-catenin/Axin
complex suggests a mechanism for the β-catenin destruction complex. *Genes Dev.*
17:2753–2764.

Yang J, Zhang W, Evans PM, Chen X, He X, Liu C. 2006. Adenomatous polyposis coli (APC)
differentially regulates beta-catenin phosphorylation and ubiquitination in colon cancer
cells. *J Biol Chem* 281:17751–17757.

Acknowledgements and funding sources

This work was partially supported by NASA (80NSSC18K1067) to AHW and MQM and
NSF (1755364) to MQM. We thank Ryan Christopher Vignogna, Lin Luan, and Le Li for
help with AlphaFold analysis.

1290 **Figures**

1291

1292
1293
Figure 1- A phylogenetic overview perspective of the evolution of changes in
protein-protein interaction domains in the Wnt/ β-catenin pathway
destruction complex scaffolding proteins Axin and APC. (A) A simplified
phylogenetic tree showing the evolutionary relationships between Bilateria and non-
Bilateria taxa. (B) Simplified representation of the cWnt pathway in its "off "and "on"
states, showing the activity of the destruction complex scaffolded by the Axin and APC
proteins. (C) Schematic diagram comparing the main protein binding domains of Axin
in bilaterians and non-bilaterians. (D) Schematic diagram of APC protein domain
structures in bilaterians and non-bilaterians. *Hs*, *Homo sapiens*; *Dm*, *Drosophila*
melanogaster; *Sp*, *Strongylocentrotus purpuratus*; *Nv*, *Nematostella vectensis*; *Ta*,
Trichoplax adhaerens; *Aq*, *Amphimedon queenslandica*; *Ml*, *Mnemiopsis leidyi*. The
numbers under the protein structures indicate the amino acid position. The color codes
for the domains are indicated within or below the protein structures.
1307

1308

1309
1310
Figure 2- Overexpression of Axin in *Nematostella* embryos disrupts Wnt/β-
catenin signaling. (A) Live imaging of embryos showed that in GFP and Nvβ-catenin-
mCherry co-expressing embryos β-catenin-mCherry was enriched in the nuclei at one
side of the embryo. However, co-overexpression of NvAxin-GFP (B) or SpAxin-GFP (C)
with Nvβ-catenin-mCherry eliminated the nuclearization of β-catenin-mCherry. (D-F)
Compared to GFP overexpressing control embryos which had a well-formed gut
epithelium (D), NvAxin-GFP overexpressing embryos (E) and sea urchin Axin
overexpressing embryos (SpAxin-GFP) (F) initiated primary invagination but failed to
maintain a normal gut epithelium at the late gastrula stage. Green: phalloidin; Blue:
DAPI. (G) Gene syn-expression domains as defined in Röttinger et al. (2012). Figure on
the left shows a lateral view and the figure on the right shows the oral view. (H, I) qPCR
analyses of NvAxin or SpAxin overexpressing embryos at 24 hpf (hours post
fertilization) showed that all endomesoderm gene markers (orange, and red bars) and
pharynx markers (blue bars) were downregulated, except for lhx6. The values represent
the fold change relative to the GFP control (black bars). The qPCR experiments were
replicated with three separate batches of embryos, with three technical replicates in
each experiment. GAPDH was used as the internal control. Statistical significance was
determined using the Unpaired t test. ****, $p<0.0001$; ***, $p<0.001$; **, $p<0.01$; *,
$p<0.05$; n=3.
1330

1331
1332

1333

Figure 3- CRISPR/Cas9 mediated knockout of *Axin* and *APC-like* induces ectopic tentacles and upregulates endomesoderm gene expression in *Nematostella* embryos. (A) *Nematostella* 4-tentacle polyp, 7 days post-fertilization (7dpf) injected with Cas9 at the zygote stage (a). *NvAxin* guide RNA/Cas9-injected polyps produced ectopic tentacles (b), which is similar to what is seen in *NvAPC-like* gRNA/Cas9 injected polyps (c). Co-injection of *Axin* and *APC-like* gRNA/Cas9 produced a more severe phenotype (d) than *Axin* or *APC-like* gRNA/Cas9 single

injections. Scale bar = 20μm. For each injection experiment, 100-200 embryos were
injected with either gRNA/Cas9 or Cas9 alone. Among these, 70-75% of the
gRNA/Cas9-injected embryos and more than 95% Cas9 only injected embryos exhibited
the same morphology as shown in the representative figures. (B) While Cas9-injected
embryos have normal oral pole *Axin* localization (a, blue staining), in situ hybridization
showed that *NvAxin*/Cas9 injected embryos displayed reduced *Axin* expression (b).
WMISH showed that 81.9% of the control embryos showed strong *NvAxin* expression at
the animal pole; 13.7% showed detectable but weaker asymmetrical *NvAxin* expression,
and 4.4% showed no detectable signal (n=204) (Fig. 2Ba). By contrast, 72.5% of *NvAxin*
gRNA/Cas9-injected embryos showed no detectable *Axin* expression, 24.4% showed
weak *Axin* expression, and only 3.1% showed a signal like that seen in the control
(n=131) (Fig. 2Bb). The embryos with weak *NvAxin* signals may be due to incomplete
excision of *NvAxin* or the generation of mosaic embryos where the gene was only
excised in some nuclei. (C and D) qPCR shows the effects of *Axin* and *APC-like* knock-
out by CRISPR/Cas9 on the expression of endomesoderm and pharynx gene markers.
Fz10, *gata*, *foxA*, *foxB*, *Wnt8*, *lmx*, *Brachyury (Bra)* (orange and red bars) are endo
mesodermal markers, *Chordin*, *lhx6* (blue bars) are markers for the pharynx; *Fz5* and
six3 (green bars) are ectoderm gene makers. qPCR experiments were replicated with
three separate batches of embryos with three technical replicates in each
experiment. GAPDH was used as the internal control. Graphs shown fold change in
mRNA levels normalized to the Cas9 only control (black bars). Statistical significance
was determined using the Unpaired t test. **, p<0.01; *, p<0.05; n=3.

1363
1364

1365

1366

1374

1375

Figure 4- Overexpression of *Nematostella* Axin in sea urchin embryos
blocks nuclearization of β-catenin and cWnt signaling. (A) In the control
where GFP and Spβ-catenin-mCherry were co-expressed, β-catenin was enriched in the
nuclei at the vegetal pole (a). Co-overexpression of NvAxin (b) or SpAxin (c) with Spβ-
catenin-mCherry downregulated the nuclearization of β-catenin. (B) Control embryos
injected with *GFP* mRNA developed normally (a, a'), while embryos developing from
SpAxin or *NvAxin* mRNA-injected zygotes became severely anteriorized (b, b' and c, c').
Scale bar = 10μm. (C and D) qPCR analyses showed that NvAxin or SpAxin
overexpressing embryos downregulated endomesoderm genes and upregulated
neuroectoderm genes. The bar graph shows the fold change of each gene between *Axin*
mRNA-injected and *GFP* mRNA-injected control embryos. *Blimp1*, *Wnt8*, *Hox11/13b*,
GataE, *Alx1*, *Delta* and *GCM* are endomesoderm gene markers; *Foxq2* and *Six3* are
neuroectoderm gene markers. qPCR experiments were replicated with three separate
batches of embryos with three technical replicates in each experiment. GAPDH was
used as the internal control. Statistical significance was determined using the Unpaired t
test. ****, p<0.0001; ***, p<0.001; **, p<0.01; *, p<0.05; n=3. For each injection

experiment, 200-300 embryos were injected with each construct and more than 95% of
embryos had the same morphology as shown in the representative figures.

1414

Figure 5- Sea urchin embryos posteriorized by Axin knock-down can be rescued by full-length NvAxin, NvAxinΔRGS, and NvAxinΔDAX, but not NvAxinΔGID. (A) Schematic representations of full-length Axin and single domain-deleted Axin constructs. (B) Gastrula and pluteus larva developing from Control MO, LvAxin MO, *GFP* mRNA or *NvAxin* mRNA injected zygotes from the sea urchin *Lytechinus variegatus* (a-d'). Control MO-injected embryos develop normally (a,a'), while LvAxin MO-injected embryos become posteriorized (b, b'). *GFP* mRNA-injected zygotes produce normal embryos (c, c'), while *NvAxin* mRNA-injected zygotes develop into anteriorized embryos (d, d'). (C) The posteriorized phenotype induced in LvAxin MO-injected embryos is not rescued by control *GFP* mRNA co-injection (a, a') but is rescued by the overexpression of NvAxin (b, b'), NvAxinΔRGS (c, c'), and NvAxinΔDAX (d, d'). Injection of *NvAxinΔGID* mRNA fails to rescue the posteriorization induced by Axin knockdown (e,e'). Scale bar = 10 µm. The numbers in the panels indicate the number of embryos examined and the number that showed the phenotype.

1429

1430

1431

1432

1433
Figure 6- Overexpression of the NvAxin GID domain upregulated
endomesodermal and pharyngeal gene expression in *Nematostella*
embryos. (A) Comparison of animals developing from *Nematostella* zygotes injected
with *GFP* mRNA (control) or *NvAxin-GID (NvGID)* mRNA at 7 days post fertilization
(7dpf). At 7dpf control animals were at the polyp stage while *NvGID* RNA-injected
embryos developed abnormally. Scale bar =10 µm. (B) qPCR analyses of NvGID
overexpressing embryos at 24 hpf (hours post fertilization) showed that all
endomesoderm gene markers (orange, red and blue bars) were upregulated. Some genes
were unregulated at modest levels while others showed significant increases. The two
apical domain genes were significantly downregulated. The qPCR experiments were
replicated with three separate batches of embryos, with three technical replicates in
each experiment. GAPDH was used as the internal control. Graphs shown fold change
in mRNA levels normalized to the GFP control (black bars). Statistical significance was
determined using the Unpaired t test. ****, $p < 0.0001$; ***, $p < 0.001$; *, $p < 0.05$; n=3. (C)
Western blot analysis of anti-GFP antibody-mediated immunoprecipitation of *in vitro*
synthesized NvGID-GFP and NvGSK-3β-mCherry fused proteins.

1450

1451
1452

Figure 7- *In vitro* analysis of protein-protein interactions indicate direct
interactions between Nvβ-catenin and components of the *Nematostella* β-
catenin destruction complex. (A) *In vitro*-synthesized GFP- or mCherry-fused
proteins were tested for potential interactions by mixing a GFP-fused partner with an
mCherry-fused partner and performing immunoprecipitation (IP) with an anti-GFP
antibody. The labels at the top of the panel show the interacting proteins in each
experiment. The top part of the Western blot analysis shows the protein-protein
interactions between protein pairs. The bottom part of the figure shows the Western blot
of GFP-tagged proteins. Sea urchin (*Strongylocentrotus purpuratus*) β-catenin (Spβ-
catenin) shows a strong interaction with sea urchin Axin (SpAxin) (Lane 5), while Nvβ-
catenin shows a consistent but considerably weaker interaction with NvAxin (Lane 1).
NvAxin also shows a relatively weaker interaction with Spβ-catenin (compare lanes 2
and 5). Unexpectedly, SpAxin shows a weaker interaction with Nvβ-catenin than with
SpAxin (compare lane 4 and lane 5). Sp and Nv Axin bind the SpGSK-3β and NvGSK-3β
respectively (Lanes 3 and 6). (B) NvAPC-like and Nvβ-catenin interact strongly in vitro
(Lane 2) while NvAPC-like shows weaker but consistent interactions with NvAxin and
NvGSK-3β (Lanes 1 and 3). (C) NvAxin-GFP and Nvβ-catenin-mCherry interact in vitro

(Lane 2) as previously shown (Figure 7A, Lane 1), changing the conserved leucine on the
Axin-RGS located βcatBM-like sequence to aspartic acid reduces the in vitro interaction
between NvAxin-GFP and Nvβ-catenin-mCherry (Lane 3) as does the same mutation to
the βcatBM-like sequence towards the C-terminus of NvAxin (Lane 4). Mutating the
leucine to aspartic acid in both NvAxin βcatBM-like sequences completely abolishes the
interaction of NvAxin-GFP with Nvβ-catenin-mCherry (Lane 5).
1476

1477

1478

1479

1480

1481

1482

1483

1484

1485

1486

1487

1488

Figure 8- A model for the co-evolution of the β-catenin destruction complex scaffolding proteins Axin and APC-like/APC. The last common ancestor (LCA) to metazoans most likely had an Axin protein that lacked the canonical β-catenin-binding motif (βcatBM) but had a βcatBM-like motif embedded within the RGS domain with possibly promiscuous β-catenin-binding activity. The APC-like protein in this metazoan LCA had ARM domains with strong similarity to APC proteins in extant species. We do not have any evidence that the Axin-APC-like proteins in the metazoan LCA scaffolded a functional β-catenin destruction complex. There were relatively few changes in the main domains in Axin and APC-like in the LCA to the Porifera and later branching metazoans, but there was the acquisition of a critical leucine in the βcatBM-like motif embedded within the RGS domain that may have imparted a higher β-catenin-binding affinity to this motif. Based on the available evidence, we propose that in the Cnidarian-Bilaterian LCA, the βcatBM-like motif embedded within the RGS domain was duplicated with the duplicated copy inserted between the GID and DAX domains of Axin. The Axin and APC-like proteins in this LCA had several protein-protein interactions not reported in bilaterians and as such, may be weak interactions not previously defined in bilaterians that scaffolded a nascent destruction complex. In the Bilateria, the βcatBM on Axin evolved with the acquisition of additional critical amino acids including a critical histidine and the βcatBM-like motif in the RGS domain was lost, possibly through drift. The Axin protein in the Bilateria did not gain additional major domains, but the APC protein gained multiple domains that allowed it to interact strongly with several components of the destruction complex and over 100 other proteins (Nelson and Nathke, 2013). Here, the domains on the bilaterian APC have been simplified for clarity but the additional domains can be seen in Fig. 1D. Model was created using BioRender.

Supplementary Table 1. The primary protein interaction domains on Axin proteins from bilaterian and non-bilaterian taxa. In bilaterians (black), Axin proteins contain APC- (RGS), GSK-3β- (GID), β-catenin- (β-cat), and Disheveled- (DAX) binding domains. In non-bilaterians (red), the conserved β-catenin binding domain between the GID and DAX domains is absent.

Axin Domains Species Name & Accession #	RGS	GID	β-cat	DAX
Strongylocentrotus purpuratus (XP_003729076)	✔	✔	✔	✔
Saccoglossus kowalevskii (NP_001158475)	✔	✔	✔	✔
Platynereis dumerilii (ANS60441)	✔	✔	✔	✔
Crassostrea gigas (EKC24785.1)	✔	?	✔	✔
Branchiostoma floridae (XP_035695674)	✔	✔	✔	✔
Homo sapiens (NP_003493)	✔	✔	✔	✔
Mus musculus (AAC53285)	✔	✔	✔	✔
Xenopus laevis (NP_001081874)	✔	✔	✔	✔
Danio rerio (BAA92439)	✔	✔	✔	✔
Hofstenia miamia (KJ658749.1)	?	✔	✔	✔
Tribolium castaneum (NP_001280495)	✔	?	✔	✔
Drosophila melanogaster (NP_733336)	✔	✔	✔	✔
Apis mellifera (XP_006564444)	✔	✔	✔	✔
Nematostella vectensis (XP_048576223)	✔	✔	✘	✔
Mnemiopsis leidyi (ML04208a)	✔	✔	✘	✔
Amphimedon queenslandica (NP_001266244)	✔	✔	✘	✔
Trichoplax adhaerens (XP_002109778)	✔	✔	✘	✔

1529
1530
1531
1532 **Supplementary Figure 1- Whole mount RNA in situ hybridization shows the**
1533 **expression of *Axin* in *Nematostella* eggs and embryos.** Axin is ubiquitously
1534 expressed in the egg, and during cleavage and blastula stages (A-C), however Axin is
1535 asymmetrically expressed at the animal pole at early and late gastrula stage (D, E) and
1536 the oral side at planula stage (F). Scale bar = 10 μm.

1554

1555
1556
Supplementary Figure 2-The effect of NvTcf knockdown on primary
invagination and endomesodermal and pharyngeal gene expression in
***Nematostella* embryos.** (A) Compared to GFP overexpressing control embryos (a),
NvdnTCF-overexpressing embryos (b) initiated primary invagination but failed to
maintain a normal archenteron epithelium at the late gastrula stage. (B) qPCR analysis
of NvdnTCF-overexpressing embryos at 24 hpf (hours post fertilization) showed that all
genes expressed in cells derived from animal pole blastomeres (orange, red, and blue
bars) are downregulated. qPCR experiments were replicated with three separate batches
of embryos with three technical replicates in each experiment. GAPDH was used as the
internal control. Statistical significance was determined using the Unpaired t test. ****,
$p<0.0001$; ***, $p<0.001$; **, $p<0.01$; *, $p<0.05$; n=3.
1568

1569

1570
Supplementary Figure 3- Morpholino mediated knockdown of Axin or APC-
like upregulates endomesoderm gene expression in *Nematostella* embryos.
(A,B) Endomesoderm and pharynx gene markers were upregulated in NvAxinMO or
NvAPC-like MO-injected embryos compared to control MO (black bars). The qPCR
experiments were replicated with three separate batches of embryos, with three
technical replicates in each experiment. GAPDH was used as the internal control.
Statistical significance was determined using the Unpaired t test. **, p<0.01; *, p<0.05;
n=3.
1579

1580
1581

1582
1583

Supplementary Figure 4- Confirmation of successful CRISPR/Cas9-mediated gene editing with Sanger sequencing using a single injected embryo from each group. Compared to the Cas9 only injected embryo (A), the sequencing chromatograms showed that injection of NvAxingRNA/Cas9 induced deletions or mutations (cyan box) at the targeted region of the Axin gene (B-E).

Supplementary Figure 5- The protein sequence alignment of Axin GID domains in various metazoan taxa. The top panel shows the alignments of amino acids sequence using Clustal Omega and Jalview. Conserved amino acids are highlighted in different colors, while gaps are represented in white. The bottom panel is the analysis of conservation and consensus. *Ml, Mnemiopsis leidyi*; *Aq, Amphimedon queenslandica*; *Ta, Trichoplax adhaerens*; *Nv, Nematostella vectensis*; *Hm, Hofstenia miamia*; *Dm, Drosophila melanogaster*; *Sp, Strongylocentrotus purpuratus*; *Lv, Lytechinus variegatus*; *Am, Apis mellifera*; *Pd, Platynereis dumerilii*; *Hs, Homo sapiens*; *Mm, Mus musculus*; *Bf, Branchiostoma floridae*; *Sk, Saccoglossus kowalevskii*. Blue boxes highlight the critical residues for the interactions between Axin and GSK-3β.

1624

1625
1626
Supplementary Figure 6-The interaction of Nv- and SpAxin with Nv- or Spβ-
catenin proteins in *Nematostella* embryos or embryonic lysates. Nv/SpAxin
and Nv/Spβ-catenin proteins were fused to click beetle green (CBG) split luciferase
fragments, and luciferase activity was detected in embryo lysates (A) and live embryos
(B) (n ≈ 150-300 per sample). Axin proteins were labeled C-terminally with the N-
terminus of CBG (CBG-N) while β-catenin proteins were labeled N-terminally with the
C-terminus of CBG (CBG-C), each connected by a flexible linker peptide. Proximity
caused by protein-protein interaction results in complementation of CBG split-luc
fragments into functional luciferase and hence stronger luminescence. Plots show
moving averages (lines) and 95% confidence intervals (shaded areas) from three
independent experiments that represent biological and technical replicates. Percentages
are relative intensities compared to a combination of constructs with the strongest
signal (SpAxin + Spβ-catenin) averaged across all replicates and timepoints. $RLU_{corr.}$ −
intensity of luminescence in relative luciferase units normalized to the amount of
expressed protein in each sample (as determined by Nanoluciferase signal). T, T_{corr} −
time of measurement (corrected by the delay necessary to lyse embryos) *** $p < 0.001$,
n.s. − not significant, Student's t-test.
1644

Supplementary Figure 7- AlphaFold-based predictions of protein-protein interactions between Nvβ-catenin/Spβ-catenin and components of the β-catenin destruction complex. The expected aligned error (PAE) plots for models of interactions between (A) SpAxin and Spβ-catenin (Spβ-cat), (B) SpAxin and SpGSK, (C) NvAxin and NvGSK-3β, (D) NvAxin and Nvβ-cat, (E) Nvβ-cat and NvAPC-like, (F) NvAxin and NvAPC-like, (G) NvGSK-3β and NvAPC-like, (H) NvAPC-like_β-catBD and Nvβ-cat, (I) NvAPC-like Axin binding domain (NvAPC-like_AxinBD) and NvAxin RGS. In the PAE plots, the dark green regions indicate low position error (high confidence in the relative positioning of the interacting residues), whereas light green regions indicate higher positional error (lower confidence). The iPTM and PTM scores, which represent confidence in the predicted inter-protein and intra-protein interactions, are also provided for each model on the right side of figure.

Supplementary Figure 8- Prediction of non-bilaterian Axin-β-catenin binding motif (βcatBM) using AlphaFold 3. (A-E) The Predicted Aligned Error (PAE) plots show the predicted interactions between NvAxin_βcatBM or NvAxin_βcatBM2 (duplicated βcatBM in NvAxin) and Nvβ-catenin, TaAxin_βcatBM and Taβ-catenin, AqAxin_βcatBM and Aqβ-catenin, MlAxin_βcatBM and Mlβ-catenin. In the PAE plot, dark green regions indicate low positional error (high confidence in the relative positioning of the interacting residues), while light green regions indicate higher positional error (lower confidence). The iPTM and PTM scores, which represent confidence in the predicted inter-protein and intra-protein interactions, are also provided for each model. (F) Protein sequence alignments of various bilaterian Axin-βcatBM domains with the predicted NvAxin-βcat, TaAxin-βcat, AqAxin-βcat and MlAxin-βcat motifs. Each residue in the alignment is color-coded, while gaps are shown in white. Conservation and consensus analysis are presented at the bottom of the sequence alignment. Sequence alignments were performed using Clustal Omega and Jalview. (G) An in vitro analysis of *Mnemiopsis* Axin and *Mnemiopsis* β-catenin interaction. Co-IP analysis of MlAxin-Flag and Mlβ-catenin-GFP fail to show an interaction between these two proteins in vitro (lane 3). *Ml, Mnemiopsis leidyi*; *Aq, Amphimedon queenslandica*; *Ta, Trichoplax adhaerens*; *Nv, Nematostella vectensis*; *Hm, Hofstenia miamia*; *Tc, Tribolium castaneum*; *Dm, Drosophila melanogaster*; *Am, Apis mellifera*; *Hs, Homo sapiens*; *Mm, Mus musculus*; *Bf, Branchiostoma floridae*; *Pd, Platynereis dumerilii*; *Cg, Crassostrea gigas*; *Sk, Saccoglossus kowalevskii*; *Sp, Strongylocentrotus purpuratus*.

1694

1695
1696
Supplementary Figure 9-The protein sequence alignment of Axin-β-catenin
binding motif (βcatBM) -like sequence in cnidarians and ctenophores. The
top panels show the alignments of amino acids sequence of βcatBM-like in Axin-RGS
(A) or C-terminal (B) generated using Clustal Omega and visualized with Jalview in
cnidarian Axins. (C) Similar alignment of the βcatBM-like in Axin-RGS from three
ctenophore species. Conserved amino acids are highlighted in distinct colors, while gaps
are shown in white. The bottom panels in A, B and C show the analysis of conservation
and consensus derived from the alignment. *Nv, Nematostella vectensis; Ed, Exaiptasia*
diaphana; At, Actinia tenebrosa; Op, Oculina patagonica; Am, Acropora muricata;
Stp, Stylophora pistillata; Of, Orbicella faveolata; Pv, Pocillopora verrucosa; Ml,
Mnemiopsis leidyi; Bo, Beroe ovata; Bm, Bolinopsis microptera.
1708
1709

1710
1711
1712
1713