Evaluation of Tumor-colonizing *Salmonella* Strains using the Chick Chorioallantoic Membrane Model

Khin K. Z. Mon[1], and Linda J. Kenney[1,2*].

[1]Department of Biochemistry and Molecular Biology, [2]Sealy Center for Structural Biology, University of Texas Medical Branch, Galveston, Texas, USA

*Address correspondence to: likenney@utmb.edu

Keywords: Chick Chorioallantoic Membrane, Tumor Colonizing *Salmonella,* CAM tumor, Therapeutic agent, 3D tumor spheroids

Abbreviations: STm, *Salmonella* Typhimurium*;* MOI, multiplicity of infection; ED, embryonic development day; DPI, days post-infection; HPI, hours post-infection; CAM, chorioallantoic membrane; LDH, Lactate Dehydrogenase; H&E, Hematoxylin and Eosin; WT, wildtype;

ABSTRACT

The chick embryo chorioallantoic membrane (CAM) tumor model is a valuable preclinical model for studying the tumor colonizing process of *Salmonella enterica* serovar Typhimurium. It offers advantages such as cost-effectiveness, rapid turnaround, reduced engraftment issues, and ease of observation. In this study, we explored and validated the applicability of the partially immune-deficient CAM tumor model. Herein, we demonstrate that *Salmonella* preferentially colonizes tumors and directly causes tumor cell death. Bacterial migration, tumor colonization, and intra-tumor distribution did not require flagellar-mediated motility. The vast majority of *Salmonella* that colonized the CAM tumor were extracellular. Thus, tumor invasion was independent of both SPI1- and SPI2-encoded type three secretion systems. Surprisingly, the extracellular residence of *Salmonella* on CAM tumors did not require biofilm formation. We evaluated our wild-type parental strain compared to the attenuated clinical strain VNP20009 and discovered a reduced tumor colonization capability of VNP20009. The inability to effectively colonize CAM tumors potentially explains the reduced anti-tumor efficacy of VNP20009. Our work establishes the xenograft CAM model as an informative and predictive screening platform for studying tumor-colonizing *Salmonella*.

IMPORTANCE

Cancer has a major impact on society, as it poses a significant health burden to human populations worldwide. *Salmonella* Typhimurium has demonstrated promise in cancer treatment by exerting direct tumoricidal effects and enhancing host-mediated anti-tumor immunity in xenograft mouse studies. A general understanding of its pathogenesis and the relative ease of genetic manipulation support the development of attenuated strains

for therapeutic use. Alternative *in ovo* models such as the CAM tumor model present a

suitable screening platform to accelerate the development of therapeutic strains. It allows

for rapid evaluation of *Salmonella* strains to assess their efficacy and potential as

oncolytic agents. The present study establishes that the *in ovo* tumor model can be

utilized as a preclinical tool for evaluating oncolytic *Salmonella*, bridging the gap

between *in vitro* and *in vivo* screening.

INTRODUCTION

In the field of bacterial-mediated cancer therapy, *Salmonella enterica* serovar

Typhimurium (STm) has emerged as an ideal therapeutic candidate. This is in part due to

its intrinsic tumor targeting ability (1), its facultative anaerobic properties, which allow

colonization in both hypoxic and oxygenated regions of heterogeneous tumor

microenvironments (2), its invasive capability (3), and its direct or immune-mediated

anti-tumor effects (3–6). STm infection mainly causes self-limiting gastrointestinal

disease in healthy humans but can sometimes manifest as extraintestinal systemic

infections in immune-compromised individuals (7). To promote its safe application in

cancer patients, bacterial pathogenic properties need to be eliminated or attenuated

without compromising the tumor colonizing capability, and thus subsequent anti-tumor

potency. Therefore, tuning the balance between virulence attenuation and tumor-

colonization capability is an important criterion in the design of suitable STm strains.

The chick embryo chorioallantoic membrane (CAM) has long been used as a

robust model for cancer and cardiovascular research, as well as for preclinical evaluation

of cancer drug therapies (8–10). The CAM model is an immunodeficient model that is

receptive to xenografting of various mammalian tumor cells without inducing a host

immune response (11). The well-vascularized embryonic tissue of the CAM supports and

preserves heterogeneous tumor growth with a much shorter experimental timeline

(typically 3-4 days) for tumor development compared to mouse models, which requires

weeks (12). Moreover, the CAM model also allows real-time visualization of tumor

growth (13) as well as the dynamics of bacterial infection via fluorescence

stereomicroscopy (14). These attributes indicate that the CAM tumor model could be

well suited for preliminary evaluation of *Salmonella*-based therapies for cancer treatment.

The engineered STm strain VNP20009 is attenuated by purine auxotrophy (a

deletion in *purI*) and lipid A modification (a deletion in *msbB*) (15). It demonstrated an

excellent safety profile with observable tumor growth retardation in a variety of pre-

clinical xenograft mouse models (16–19). Despite the high success rate in the mouse

studies, no anti-tumor effect was observed in humans. The intravenous infusion of the

VNP20009 strain to melanoma patients in a clinical trial failed and was attributed to a

low bacterial tumor colonization and rapid bacterial clearance by the host immune

response (20). The failure of the VNP20009 strain to colonize tumors highlighted the

xenograft mouse model as a poor predictor of therapeutic success in humans. Hence,

there is a need to explore alternative models to overcome some of the translational

limitations associated with traditional mouse models. We hypothesized that the CAM

tumor model might serve as a preliminary screening platform to study the tumor-bacterial

interactions to evaluate the efficacy of tumor-targeted *Salmonella* strains for therapeutic

purposes.

In the current work, we developed the *in ovo* CAM tumor model and compared it

with an *in vitro* three-dimensional (3D) tumor spheroid model to examine the relevance

of the screening data obtained (21). While 3D tumor spheroids can mimic *in vivo* solid

tumor features such as structural organization, cellular assembly and nutrient gradients

(22), they cannot fully simulate the heterogenous tumor microenvironment established

within a living host. Therefore, the CAM tumor model serves as a better pre-clinical

model, bridging an informational gap between *in vitro* cell-based models and *in vivo*

animal-based systems for screening oncolytic *Salmonella*. With the CAM tumor model,

we demonstrated that STm pathogenic traits such as flagella, virulence factors and

biofilm formation were not essential for tumor colonization. We further discovered that

reduced tumor colonization by VNP20009 was attributed to its growth sensitivity within

the tumor microenvironment, which subsequently impacted its anti-tumor effectiveness.

RESULTS

Validation of the CAM model for studying tumor-colonizing *Salmonella*

A general overview of the CAM tumor STm infection model is depicted in Figure

1. We first determined the infectious dose of STm by infecting tumor bearing chick

embryos with different doses of bacteria: approximately 500 colony-forming units

(CFU), 50 CFU, or 5 CFU. Chick embryo mortality was monitored daily post-bacterial

infection with PBS as the control group. The highest inoculum of ~500 CFU caused the

greatest mortality (38% survival) at one day post-infection (DPI) compared to groups that

received lower dosages of ~50 CFU (66% survival) or ~5 CFU (77% survival)

(Supplementary Figure 1). To minimize the mortality of highly susceptible tumor-bearing

chick embryos as well as to ensure effective tumor colonization, we selected the mid-

range of ~50 CFU as the optimal infectious dose.

The important features of tumor-targeting STm are the higher tumor specificity in

colonization compared to organ colonization, and the ability to penetrate tumor tissue

(23, 24). These criteria were tested first in the CAM tumor model. At 1 DPI, WT STm

accumulated preferentially in tumors over livers of chick embryos at a ratio of 1000:1.

The mean CFU/g in tumors was 1.9×10^8 CFU/g compared to 1.9×10^5 CFU/g in livers

(Figure 2A, note log scale). Next, we examined the intra-tumoral distribution of STm.

Infected whole tumors were harvested, embedded in paraffin and sectioned at different

tissue depths (top, middle and bottom). The tumor sections were stained with antibodies

against bacterial lipopolysaccharide (green), nuclei (blue) and E-cadherin (magenta) and

examined by confocal microscopy. STm was observed at all tumor tissue depths as

evident by lipopolysaccharide staining (LPS = green, second panel Figure 2B and

Supplementary Figure 2). Specifically, bacteria were mostly present in the metastatic

tumor cell regions, where decreased E-cadherin staining indicated a loss of cell-to-cell

adhesion junctions.

STm colonization drove direct tumor apoptosis in the CAM model

We next tested the hypothesis that STm infection directly promotes tumor cell

death. A trypan blue exclusion assay was performed on dissociated tumor cells to test *ex*

vivo tumor cell viability. At 3 DPI, STm infection significantly reduced the percentage of

tumor cell viability (74%) compared to PBS treated tumors (84%) (Figure 2C). Because

the presence of STm was detected throughout the entire tumor (Figure 2B), it raised the

question of whether its localization directly induced tumor cell apoptosis. We utilized the

terminal deoxynucleotidyl transferase-mediated deoxyuridine triphosphate nick end-

labeling (TUNEL) assay to detect *Salmonella*-induced DNA fragmentation in tumor

cells. A marked accumulation of TUNEL-positive cells (red signals) was observed in

STm- infected tumors relative to PBS-treated tumor sections at the top, middle and

bottom levels (Figure 2D). Immunofluorescence images of TUNEL-stained tumor

sections were further quantified using Fiji software (25). The percentage of TUNEL-

positive nuclei in STm infected tumor tissue sections was significantly higher than PBS

tumors (52% vs 33%, respectively), indicating that tumor cells underwent increased

apoptosis following bacterial infection (Figure 2E). Lastly, the histological changes in

Hematoxylin and Eosin (H&E)-stained tumor sections were analyzed. A more extensive

necrotic area in *Salmonella*-infected tumors (22%) was detected compared to the PBS

control group (4%) (Figure 2F, 2G).

Bacterial motility was not required for tumor colonization or intra-tumoral distribution

The prevailing hypothesis regarding a role for flagella in tumor colonization is

that motility is essential for bacteria to effectively target, colonize, and disperse within

the tumor environment (26–28). To test this hypothesis, WT or a *ΔflgK* null strain were

directly inoculated onto CAM tumors. The *flgK* gene encodes a flagellar hook-associated

protein; its deletion renders *Salmonella* non-motile (29). Live *in ovo* images of the

bacterial colonization pattern, as well as CFU enumeration from tumors was comparable

between the WT and the *ΔflgK* null strain (Figure 3A, B). We reasoned that the role of

flagella might be dispensable when bacteria were directly inoculated onto the tumor,

rendering colonization a passive process. Therefore, we also tested a distant inoculation

route by depositing bacteria a short distance away from the tumor (~0.5 cm) on filter

papers to test whether *Salmonella* required flagellar-mediated motility for tumor

colonization. The *ΔflgK* strain showed no defect in CAM tumor colonization, and

colonized tumors to the same extent as the WT (Figure 3A, C). The intra-tumoral

distribution pattern of the non-motile *ΔflgK* strain was also comparable to that of the

motile WT (Figure 2B), indicating that the *ΔflgK* strain was able to disperse and penetrate

all depths of the tumor (Figure 3D). These results provided strong evidence that STm

migration, tumor colonization and intra-tumoral distribution were not dependent on

bacterial motility in the CAM tumor model.

3D tumor spheroids do not recapitulate the CAM tumor model with respect to key

virulence factors encoded on pathogenicity islands 1 and 2

In the host gastrointestinal tract, STm utilizes a Type III secretion system (T3SS1)

encoded by *Salmonella* Pathogenicity Island-1 (SPI-1) to invade host intestinal epithelial

cells and a SPI-2-encoded T3SS2 for intracellular growth and survival within host

macrophages (30–33). We therefore investigated whether bacterial invasion or replication

within tumors was also dependent on SPI-1 and SPI-2. To address this question, we first

employed the *in vitro* 3D tumor spheroid model before assessing their roles *in vivo* in the

CAM tumor model. The use of 3D magnetic cell culture technology produced structurally

relevant tumor spheroids that mimicked some features of *in vivo* solid tumors. The spatial

architecture with regard to oxygen and nutrient gradients, extracellular matrix, and cell-

to-cell interactions is maintained (21, 22). Within tumor spheroids, intracellular invasion

and replication rates of STm strains were examined using a gentamicin protection assay.

The elimination of *hilD* (the SPI-1 master regulator) resulted in a significant invasion

defect with a mean reduction of 1.4 log CFU/well compared to WT (Figure 4A),

suggesting that SPI-1 played a significant role in early bacterial invasion of tumor cells.

At 16 HPI, an intracellular replication rate of both Δ*hilD* (mean log CFU/well = 3.5) and

the SPI-2 master regulator, Δ*ssrB* (mean log CFU/well = 4.5) were significantly reduced

relative to WT (mean log CFU/well = 5.6) (Figure 4A). Consequently, the overall

replication rate (16 HPI/ 2 HPI) was higher for WT (28-fold) compared to either Δ*hilD*

and Δ*ssrB* strains (6-fold and 4-fold, respectively) (Figure 4B). We next examined the

cytotoxic damage induced by bacterial infection of 3D tumor spheroids using a lactate

dehydrogenase (LDH) assay 16 HPI. The amount of LDH release is directly correlated

with cell cytotoxicity levels. Tumor spheroids infected with WT bacteria released higher

amounts of LDH (53% of total maximum LDH release at 100%) compared to SPI-1

(22%) or SPI-2 (33%) deficient strains (Figure 4C). Whole-mount immunofluorescence

staining of 3D tumor spheroids exhibited a specific cellular structural organization.

Proliferating tumor cells were observed in the periphery, with an inner layer of quiescent

cells and a necrotic core region. These were evident by distinct fluorescence intensity

profiles of the cell proliferation marker Ki67, in which the intensity varied from high to

low from the periphery to the core (Figure 4D). From representative images of infected

3D tumor spheroids at 16 HPI, both SPI-1- and SPI-2-deficient strains were observed at

reduced levels compared to the WT (Figure 4D) consistent with the quantification of

CFU (Figure 4A).

We then investigated whether tumor colonization and intracellular invasion or

replication required SPI-1 or SPI-2 in the CAM tumor model. Using an ex vivo

gentamicin assay, we quantified total tumor-associated bacteria (i.e. both extra- and intra-

cellular bacteria) recovered from the CAM tumors. Both SPI-1 (*ΔhilD)* and SPI-2

(*ΔssrB*)-deficient strains were found to colonize (extracellular) and invade (intracellular)

tumor cells to the same extent as WT at 1 DPI (Figure 5A, B). In the mouse tumor model,

Salmonella is found almost exclusively extracellularly, forming biofilms on tumors (34).

We therefore assessed whether *Salmonella* adapted a similar residence lifestyle in our

CAM tumor model. Based on the percentage of the total tumor-associated bacterial

population, approximately 99% of bacteria were extracellular, with ~1% of the bacteria

found inside the CAM tumor cells (WT, *ΔhilD* and *ΔssrB*) (Figure 5C). Hence, this

finding corroborated that the loss of SPI-1 and SPI-2 genes did not impair the tumor-

colonizing ability of STm (Figure 5B), because STm were predominantly extracellular on

the CAM tumors.

To address the question of whether extracellular STm also form biofilms on CAM

tumors, we infected tumors with *Salmonella* WT or a *ΔcsgD* null strain; CsgD is the

master biofilm regulator (35). The extracellular bacterial load recovered indicated that

both strains successfully colonized the tumor to the same extent: a mean log CFU/g of ~7

to 8, at 1 and 3 DPI (Figure 5D). Thus, in contrast to the mouse tumor model (34), the

ability to form biofilms was not required for the extracellular accumulation of STm on

CAM tumors.

How does the clinical strain VNP2009 compare with WT *Salmonella*?

Despite promising results in xenograft mouse models (effective tumor

colonization with tumor growth retardation) the clinical strain VNP20009 did not

perform well (low bacterial tumor colonization) in a human clinical trial. It was therefore

of interest to evaluate colonization by VNP20009 compared to its parental WT strain (15,

36) in both 3D tumor spheroid and CAM tumor models.

The clinical strain VNP20009 replicates poorly within 3D tumor spheroids

We first investigated the performance of the clinical strain VNP20009 within 3D

tumor spheroids. Compared to WT, the VNP20009 strain displayed a modest invasion

defect with a mean reduction of 0.6 log CFU/well (Figure 6A, left columns). However,

the replication defect at 16 HPI was striking, with an overall mean reduction of 1.6 log

CFU/well relative to WT (Figure 6A, right columns). Replication of the WT was 28-fold,

but VNP20009 only increased 3-fold (Figure 6B). These differences indicated a stark

difference in the ability of the attenuated VNP20009 strain to grow within the tumor

spheroid compared to the WT. The possibility that differential growth rates between

strains contributed to tumor invasion was ruled out, as the growth rate of VNP20009 was

not compromised under bacterial culture conditions (Supplementary Figure 3).

Furthermore, bacterial inoculums were prepared under the same SPI-1 inducing condition

for both strains. The percentage of bacteria that were SPI-1 positive (using a P*prgH*-

mCherry reporter) for WT and VNP20009 were similar at 67% and 62%, respectively, of

the total population (Supplementary Figure 4). We next compared the ability of

VNP20009 to induce tumor cell cytotoxicity with the LDH assay after 16 HPI. WT-

infected tumor spheroids released higher amounts of LDH (53%) than VNP20009 (36%)

(Figure 6C).

Representative images of VNP20009-infected tumor spheroids highlighted the

reduction in intracellular bacteria compared to WT (Figure 6D), corroborating the

bacterial enumeration dataset (Figure 6A, B). Furthermore, tumor spheroids infected with

VNP20009 revealed distinct cell morphological defects of the strain with elongated

bacterial cells (Figure 6E), confirming the strain growth defects within the tumor

spheroid (see Discussion).

Reduced CAM tumor colonization of VNP20009 compromises its anti-tumor immunity

We examined the tumor colonization ability of VNP2009 in the CAM tumor model. The effect of VNP20009 administration (~50 CFU) on chick embryo mortality was first evaluated against the WT and PBS control groups. A significant improvement in survival rate and prolonged survival time was noted for tumor-bearing chick embryos infected with VNP20009 compared to the WT (Figure 7A). Chick embryos completely succumbed to WT infection by four days, whereas VNP-infected embryos survived for up to seven days. We next assessed the tumor-colonizing capability of VNP20009. Live *in ovo* images of mCherry-tagged bacterial strains colonizing the CAM tumor highlighted the marked reduction in fluorescence signals of VNP20009 at 1 DPI (Figure 7B) compared to the WT. The image emphasizes the strain attenuation of VNP2009 in tumor colonization, which was also apparent when we examined the intra-tumoral distribution (Supplementary Figure 5). Compared to WT, VNP20009 was remarkably reduced throughout the tumor tissue sections at all depths. This result was further confirmed by an overall reduction in the extracellular bacterial load recovered from VNP20009-infected tumors (Figure 7C). At 1 DPI, the extracellular bacterial load of VNP20009 was only ~1% of WT. A reduction of VNP20009 tumor colonization was still evident at 3 DPI and was ~36% of WT colonization (Figure 7C). When we compared the two strains and their ability to colonize the liver (Figure 7D), VNP2009 exhibited an early defect in dissemination (~10% of WT). However, the attenuated strain was able to colonize chick embryo livers to a similar level as the WT by 3 DPI (Figure 7D). This was in contrast to tumor colonization (Figure 7C), where the rate of VNP2009 colonization remained

weaker than the WT over time. Furthermore, a cell morphological defect was observed,

as elongated chains of VNP20009 were detected in CAM tumor tissue sections at 3 DPI

(Figure 7E). These chains were strikingly similar to our previous observations in 3D

tumor spheroids (Figure 6E). The CAM tumor invasion rate of VNP20009 was also

compromised compared to the WT invasion rate (see intracellular bacterial counts, Figure

7F).

Chick embryos possess only partial immunity at early developmental stages

(before ED 9), but as immunocompetence develops, embryos are capable of mounting an

inflammatory response with cytokine expression from host immune cells (37). Induction

of pro-inflammatory cytokines such as tumor necrosis factor-alpha (TNF-α) and

Interleukin-1 beta (IL-1β) following *Salmonella* infection in tumor-bearing hosts are

indicative of host-mediated tumoricidal effects (38). To investigate whether host immune

factors play an indirect role in driving anti-tumor immunity, CAM tumors were infected

with WT, VNP20009 or PBS at ED11, and cytokine expression in tumor lysates at 2 HPI

was measured. TNF-α levels were similar across uninfected and infected tumors (WT and

VNP20009) at 2 HPI (Figure 7G). In contrast, IL-1β was highly expressed in WT-

infected tumors, while IL-1β levels were similar between uninfected (PBS) and

VNP20009-infected tumors at 2 HPI (Figure 7H). This result suggests that reduced tumor

colonization by VNP20009 failed to trigger activation of host-mediated anti-tumor

immunity in the CAM model.

DISCUSSION

To the best of our knowledge, this is the first study to exploit the CAM tumor

model as an *in vivo* model to study *Salmonella*-mediated anticancer therapy. Our work

established that the chick embryo CAM model reproduces key features of tumor-

targeting STm reported in the mouse model: (1) tumor specificity in colonization (Figure

2A), (2) intra-tumoral distribution (Figure 2B), (3) direct induction of tumor cell death

(Figure 2C-G) and (4) indirect stimulation of anti-tumor immunity to some degree in the

partial immunocompetent host (Figure 7H). To assess whether the CAM model

recapitulated the published mouse studies, *Salmonella* key pathogenic traits were

investigated for their respective role in tumor colonization and invasion.

Flagellar-dependent motility as a driver of bacterial migration, tumor

colonization, and intra-tumoral distribution was not essential, as a non-motile *Salmonella*

(Δ*flgK)* was not impaired for colonization in the CAM model (Figure 3). This finding

agrees with published mouse data indicating that tumor colonization by *Salmonella* is a

passive process (not requiring bacterial motility), where bacterial infection induces a

cytokine storm and hemorrhage, facilitating the deposition of bacteria at the tumor site

(23, 39). In the CAM tumor model, the passive tumor migration process of STm (via a

distant inoculation route, Figure 3A, C) was most likely aided by the active movement

motion of a live chick embryo just beneath the CAM layer.

In our *in vitro* 3D tumor spheroids, both SPI-1 and SPI-2 contributed to STm

intracellular replication within the tumor spheroids (Figure 4). In the mouse model, there

are conflicting reports as to whether SPI-1 or SPI-2 virulence genes are required for

tumor invasion. Although SPI-1-mediated tumor invasion (40) and SPI-2-associated

intra-tumoral replication (41) were reported to be important, in the majority of mouse

tumor studies, virulence functions were not required, as bacteria were extracellular on

tumor sites (23, 34, 42, 43). Consistent with these latter findings, virulence functions

were not required for CAM tumor colonization and invasion (Figure 5A, B). In our

experiments, *Salmonella* strains (WT, Δ*hilD,* Δ*ssrB*) were predominately extracellular on

the CAM tumor (Figure 5C). The inconsistency between *in vitro* and *in ovo* data is likely

attributed to the fact that 3D spheroids consist of a single cancer cell-type, which lacks

the heterogeneity and vascularization of the chick CAM model to mimic the complexity

of the tumor microenvironment within the host and its interaction with bacteria.

In the mouse model, an almost exclusive extracellular residence as biofilms on

solid tumors was reported to be a bacterial protective mechanism against phagocytosis by

host immune cells, specifically neutrophils (34). However, the CAM tumor model did not

reproduce the STm biofilm phenotype, as the Δ*csgD* strain was able to colonize tumors to

similar levels as WT (Figure 5D). Immunological differences in the tumor

microenvironment between the two models (mouse vs chick embryos) might explain the

differences in an adaptation of bacterial lifestyles on solid tumors. In contrast to a

mammalian host, chick embryos have heterophils, the avian equivalent of neutrophils, but

lack myeloperoxidase (37). Previously it was shown that heterophil recruitment to the

inflammation site following lipopolysaccharide stimulation was detectible at ED7 (44),

but fully functional heterophils within developing embryos was only described at ED18,

three days before hatching (45, 46). Our CAM experimental timeline is completed by

ED14 due to the low survival percentage of tumor-bearing chick embryos at an STm

infectious dose of ~50 CFU. Therefore, the lack of functional heterophils within the chick

embryonic immune system in the current study timeline likely influences the non-biofilm

residence of STm on CAM tumors. As with every model, the CAM tumor model has its

limitations. As highlighted above, the lack of a functional host immune system and the

short experimental timeline (inability to assess STm impact on tumor growth or

regression) should be considered during experimental design and data interpretation.

Previous studies examining the anti-tumor efficacy of the highly attenuated STm

strain VNP20009 failed to directly compare the tumor colonizing level of the strain

relative to its parental WT (15, 16, 18, 47–49). Here, we used both the *in vitro* 3D tumor

spheroids and CAM tumor model to provide evidence of reduced tumor colonization,

invasion and replication by VNP20009 compared to WT (Figures 6AB, 7BC, F). In 3D

tumor spheroids, the reduced intracellular replication of VNP20009 has consequences

that limit its ability to kill tumor cells (Figure 6C). In addition, we observed cell

morphology defects of VNP20009 (i.e., elongated bacteria cells) in both models during

infection (Figures 6E, 7E). Although deletion of *msbB* in VNP20009 has been reported to

exhibit severe growth and morphology defects under *in vitro* growth conditions in the

presence of 5% CO_2, acidic pH and high osmolarity (50, 51), our work provides

additional evidence of the inherent growth sensitivity of VNP20009 in the *in vivo* tumor

microenvironment (Figure 7E). We reason that this growth defect is related to cancer

metabolism, which undergoes aerobic glycolysis (also known as the Warburg effect),

resulting in elevated CO_2 levels and lactic acid in the tumor microenvironment (52).

A poor human clinical outcome of VNP20009 was mainly the result of low tumor

colonization that compromised its anti-tumor efficacy (20). In our CAM tumor model, we

showed that IL-1β expression levels in tumor lysates of VNP20009 and PBS controls

were similar (Figure 7I), suggesting that the reduced presence of VNP20009 on tumors

was insufficient to elicit the anti-tumor immune response in the CAM model. These

results highlighted the importance of maximizing the tumor colonizing capability of the

bacterial strain as a primary factor in strengthening its anti-tumor effectiveness.

In summary, the chick embryo CAM model offers a valuable tool for initial

screening and preclinical assessment of tumor-colonizing *Salmonella* strains. Its

accessibility, vascularization, and relevance to tumor biology make it a suitable choice

for studying bacterial interactions with tumors and assessing potential therapeutic strains

for bacterial-mediated cancer therapy.

MATERIALS AND METHODS

3D Spheroid infection model

A549 tumor cell spheroids were formed with the 96-well bioprinting kit (Greiner Bio-

One catalog. 655841) according to the manufacturer's guidelines. Cells were first

magnetized with NanoShuttle-PL, nanoparticles overnight. Magnetized cells were

counted and seeded at the same cell density of ~50,000 cells/well. Using the 96-well

magnetic drive placed at the bottom of the well, magnetized cells were printed into

spheroid for 24 h. The next day, STm strains grown under SPI-1 inducing conditions

were added into each well at an MOI of 100. Bacteria were added for 30 min, high

gentamicin concentrations of 100 μg/ml were added for 1.5 h for invasion (2 HPI) and

changed to low gentamicin of 20 μg/ml for the remainder of the infection process. We

confirmed that all extracellular bacteria were eliminated post-gentamicin treatment by

plating the cell culture medium. At each step of the cell media replacement and wash

process, the 96-well plate was placed on the holding drive to hold the spheroids down

while aspirating the solution to prevent cell loss. At both 2 HPI (invasion) and 16 HPI

(replication), cell media were aspirated, washed with phosphate-buffered saline, PBS

(Roche, 11666789001), and replaced with 1% Triton X-100 (Sigma, T8787-100ml) for

10 min to lyse the cells. Spheroids were homogenized by the vigorous pipetting method.

Lysates were then serially diluted and plated for intracellular bacterial enumeration.

Whole-mount Immunofluorescence staining of 3D tumor spheroids

At the end of a 16 h STm infection, tumor spheroids were fixed in 4% paraformaldehyde

for ~5 h, washed with PBS thrice, and then processed for whole-mount

immunofluorescence staining. Each tumor spheroid was individually picked up using the

magpen with teflon caps (Catalog no: 657850, Grenier, Bio-One), during each step of the

staining process for solution replacement. Antigen retrieval was performed by

transferring spheroids into a citrate buffer (Sigma, C9999-100ml) solution for 20 minutes

in 99°C water bath before allowing the solution to cool to room temperature. The

permeabilization step was performed with 0.3% Triton-X/PBS for 15 min followed by

incubation with blocking buffer, 5% bovine serum albumin in PBS (BSA; Sigma-

Aldrich, Catalog A7030) for 1 h. Spheroids were then incubated with primary antibody,

recombinant anti-Ki67 (Abcam Catalog ab16667) diluted in blocking solution (1:200),

overnight at 4°C in the dark. The next day, spheroids were incubated with a secondary

antibody to Ki67, Alexa Fluor 488 goat anti-rabbit IgG H&L (Abcam Catalog ab150081),

and Hoechst (blue, Invitrogen, H3570) for 1 h. Whole spheroids were then mounted onto

the slide with a prolong gold antifade mountant (Invitrogen Catalog: P36934), coverslip

sealed with nail polish and dried overnight in the dark. Whole-mount spheroids were

imaged with Olympus SpinSR-10 spinning disk confocal at 10x air objective (NA 0.40),

and 100x oil objective (NA 1.5). Acquired microscopy images were processed with Fiji

software (25).

Embryonated chicken CAM tumor model

Specific pathogen-free (SPF), premium fertilized chicken eggs were purchased from

AVS bio and set horizontally in a Rcom Max 50 incubator at 37°C and 60% humidity.

On ED3, CAM of the chick embryo was dropped by inserting a thin needle at the apex of

the egg and removing ~4 ml of albumin with a syringe and 21G needle. A small window

(~1 cm^2) was made on top of the eggshell and resealed with 3M Tegaderm TM

transparent film dressing. On ED7, A549 cell pellets (~2 x 10^6 cells/egg) were mixed

with 50 µl Matrigel (Corning 354234 or 354263) and were grafted into the center of a

sterilized silicone ring placed on top of the CAM layer. Eggshell windows were then re-

sealed with Tegaderm and put back into the incubator to allow tumor growth for four

days. On ED11, the silicone ring was first carefully removed from the CAM without

disrupting the established tumor before the application of the bacterial inoculum directly

onto the tumor with a pipette. Bacterial inoculums were plated each time to confirm that a

similar dose was applied for all strains. For the distant inoculation route experiment,

bacteria were deposited onto a sterile cut-out filter paper, placed a short distance away

from the tumor (~0.5 cm). At different time points, tumors and livers were harvested for

enumeration of viable bacteria by homogenization, serial dilution, and plating methods.

All chick embryo experiments were repeated twice (N = 2) with 5-10 tumor bearing

embryos in each treated group.

TUNEL assay and apoptosis quantification with Fiji

Detection of apoptotic cells on tumor tissue sections was performed with Click-iT Plus

TUNEL assay (Invitrogen, Thermo Fisher Scientific Lot 2559160) according to the

manufacturer's guidelines. Immunofluorescence images were acquired with an Olympus

Super Res Spinning Disk, SpinSR-10 microscope at 100x oil objective (NA 1.5).

Quantification of TUNEL-positive nuclei as well as DAPI nuclei was performed with Fiji

(25) following the optimized and unbiased image processing protocol. Briefly, images

were first split into separate color channels and compressed with z-axis maximum

intensity projection. Background noise reduction was performed by subtracting

background values from each channel. Images were then converted to greyscale with 8-

bit. The Binarization thresholding function available in Fiji (25) was utilized and

thresholding algorithms that capture the best outline of stained cells for each channel

were selected (53). Moments and Li auto-thresholding algorithms were opted for

highlighting TUNEL-positive nuclei and DAPI-positive cells, respectively. Lastly, the

analyze particle function was applied for the count read-outs. The percentage of TUNEL-

stained cells was calculated against the total cell numbers (DAPI-positive cells) within

the same region.

Statistical Analysis

All statistical analysis were performed with GraphPad Prism 10.2.1.

Acknowledgements: We are extremely grateful for the thoughtful comments provided

by our reviewers. We thank Dr. Ruby Yun-Ju Huang from National University of

Singapore for guidance and expertise on the chick chorioallantoic membrane (CAM)

tumor model. This work was supported by Cancer prevention and Research Institute of

Texas (CPRIT) Grant RP2000650 awarded to LJ.K.

Conflict of interests: All authors declared no conflict of interests.

FIGURE LEGENDS

Figure 1: **Experimental scheme of the CAM tumor, STm infection model.**

Figure 2: STm promotes tumor cell death in the CAM model. (A-G) Chick embryos

with pre-established A549 tumors were directly inoculated with *Salmonella* WT or PBS

on ED 11. Tumors and livers were harvested at 1 and 3 DPI. (A) *Salmonella* WT CFU

recovered per gram of tissue from tumors and livers at 1 DPI (tumor to liver ratio

1000:1). (B) Representative images of *Salmonella* WT LPS (green), E-cadherin

(magenta) and Hoechst (blue) staining detected in tumor sections from top, middle and

bottom of whole tumors harvested at 1 DPI. Scale bar = 100 µm (10x objective) and 10

µm (100x objective). The middle panel is the zoomed image of the white box and the

right panel is the zoomed image of the red box at 100x objective. (C) A trypan blue

exclusion assay was performed on *ex vivo* dissociated tumor cell suspensions, indicating

a loss of tumor cell viability in the group infected with WT *Salmonella* at 3 DPI. (D)

Representative images of TUNEL-stained (red) and Hoechst (blue) tumor sections from 3

DPI (top, middle, bottom). Scale bar = 10 μm, 100x objective. (E) Quantitative

comparison of the percentage of TUNEL-positive nuclei between WT-infected tumors

(52%) and PBS-treated tumors (33%) at 3 DPI. (F) Representative images of H&E tumor

sections. Zoomed images of the red box of the necrotic region at 3 DPI. N: necrotic areas.

(G) The percentage of the necrotic area as quantified by an external pathologist from H

and E-stained tumor sections infected with WT (22%) and PBS (4%) at 3 DPI. The

results represent the mean ± SD. For all, the statistical significance was by unpaired t-

test, not significant (ns). P-value * ≤ 0.05, ** ≤ 0.01, *** ≤ 0.001, **** ≤ 0.0001.

Figure 3: Flagellar motility is not required for tumor colonization or intra-tumoral

distribution. (A) Representative live *in ovo* images of mCherry-tagged *Salmonella.* WT

and *ΔflgK* colonizing A549 tumors on CAMs via either direct (bacteria inoculated

directly on tumors) or distant inoculation (bacteria deposited on filter paper, ~0.5 cm

away from tumors) at 1 DPI. (B-C) Total *Salmonella* WT and *ΔflgK* CFU recovered per

gram of tumor tissue at 1 DPI from direct and distant inoculation. The results represent

the mean ± SD. The statistical significance was by unpaired t-test, not significant (ns).

(D) Representative images of *Salmonella ΔflgK* LPS (green), E-cadherin (magenta) and

Hoechst (blue) staining detected in tumor sections from top, middle and bottom levels of

whole tumors harvested at 1 DPI. Scale bar = 100 µm (10x objective) and 10 µm (100x

objective). The second column is the zoomed in details of the white box at 100x

objective.

Figure 4: SPI-1 and SPI-2 are required for intracellular replication of STm within

3D tumor spheroids. (A) Intracellular bacterial enumeration of STm strains (WT, *ΔhilD*

and ΔssrB) in A549 tumor spheroids measured at invasion (mean log CFU/well at 2 HPI:

WT = 4.2, *ΔhilD* = 2.8, *ΔssrB* = 3.9) and replication (mean log CFU/well at 16 HPI: WT

= 5.6, *ΔhilD* = 3.5, *ΔssrB* = 4.5) stages of infection. Dots: biological replicates, each with

a minimum of five technical replicates. The bars represent the mean +/- SD. (B) Fold

replication (16 HPI/2 HPI) of each STm strain (WT = 28-fold, *ΔhilD* = 6-fold, *ΔssrB* = 4-

fold) within A549 tumor spheroids. The bars represent the means of four biological

replicates, a minimum of five technical replicates each. Error bars +/- SD. (C) The

amount of LDH released from A549 tumor spheroids after infection with STm (WT=

53% , *ΔhilD* = 22% and *ΔssrB = 33%)* at 16 HPI. The bars represent the means of three

biological replicates, a minimum of five technical replicates each. Error bars +/- SD. For

all, the statistical significance was determined by one way analysis of variance (ANOVA)

with Sidak's multiple comparison test. Not significant (ns), P-value *≤ 0.05, **≤ 0.01,

≤ 0.001, *≤ 0.0001. (D) Representative images of immunofluorescence-labeling

of 3D tumor spheroids with Ki67 AlexaFluor 488 (green), Hoeschst (blue) and mCherry-

tagged bacterial strains (red) at both 10x and 100x objectives (edge and core of

spheroids). Maximum intensity projections of Z-stack images were taken with an

Olympus Super Resolution Spinning Disk SpinSR-10 microscope. Scale bar = 100 μm

(10x) and 10 μm (100x).

Figure 5: In the CAM tumor model, STm resides in an extracellular, non-biofilm

state. (A) Extracellular CFU recovered after infection with WT *Salmonella*, *ΔhilD* and

ΔssrB per gram of tumor at 1 DPI. Statistical significance by one-way ANOVA with

Sidak's multiple comparison test. (B) Intracellular WT *Salmonella*, *ΔhilD* and *ΔssrB*

CFU recovered per gram of tumor at 1 DPI. Statistical significance by one-way ANOVA

with Sidak's multiple comparison test. (C) Percentage of total tumor-associated STm

(WT, *ΔhilD, ΔssrB)* recovered before and after an ex vivo gentamicin assay from infected

CAM tumors at 1 DPI. EX: extracellular, IN: intracellular. Statistical significance was by

one-way ANOVA with Sidak's multiple comparison test. (D) Extracellular *Salmonella*

WT and *ΔcsgD* CFU recovered per gram of tumor at 1 and 3 DPI. The results represent

the mean ± SD. The statistical significance was by unpaired t-test. For all, not significant

(ns), P-value * ≤ 0.05, ** ≤ 0.01, *** ≤ 0.001, **** ≤ 0.0001.

Figure 6: Replication of VNP20009 within 3D tumor spheroids is impaired. (A)

Intracellular bacterial enumeration from WT and VNP20009 infected tumor spheroids,

measured at invasion (mean log CFU/well at 2 HPI: WT = 4.2, VNP20009 = 3.6) and

replication (mean log CFU/well at 16 HPI: WT = 5.6, VNP20009 = 4) stages of infection.

Dots: biological replicates, each with a minimum of five technical replicates. The bars

represent the mean +/- SD. The statistical significance was determined by unpaired t-test.

(B) Fold replication (16 HPI/2 HPI) of WT (28-fold) and VNP20009 (3-fold) strains

within tumor spheroids. The bars represent the means of four biological replicates, a

minimum of five technical replicates each. Error bars = +/- SD. The statistical

significance was by unpaired t-test. (C) LDH release from tumor spheroids infected with

WT (53%) and VNP20009 (36%) at 16 HPI. The bars represent the means of three

biological replicates, a minimum of five technical replicates each. Error bars = +/- SD.

Statistical significance was by one-way ANOVA with Sidak's multiple comparison test.

For all, non-significant (ns), P-value *≤ 0.05, **≤ 0.01, ***≤ 0.001, ****≤ 0.0001. (D)

Representative immunofluorescence images of 3D tumor spheroids with Ki47

AlexaFluor 488 (green), Hoeschst (blue) and mCherry-tagged bacterial strains (red) at

10x and 100x objectives. Maximum intensity projections of Z-stack images were taken

with an Olympus Super Resolution Spinning Disk SpinSR-10 microscope. (E) Zoomed

300% magnification of white box ROI from 100x tumor spheroids edge of both WT and

VNP20009 to compare and visualize the cell morphology defect of VNP20009, outlined

by dotted lines.

Figure 7: Reduced tumor colonization by VNP20009 impacts its anti-tumor

immunity. (A) Tumor bearing chick embryos infected with VNP20009 (purple line)

showed significant improvement in the percentage survival as well as longer survival

times compared to chick embryos that were infected with WT (red line) at a dose of ~50

CFU. For survival analysis, the Log-rank (Mantel-Cox) test was used. (B) Representative

live, *in ovo* images of mCherry-tagged *Salmonella,* WT and VNP20009 colonizing CAM

tumors at 1 DPI. (C-D) Extracellular *Salmonella* WT and VNP20009 CFU recovered per

gram of tumor and liver at 1 and 3 DPI. The statistical significance was determined by

unpaired t-test. (E) Super Resolution 320x magnified representative images of WT

compared with distinct cell morphology defects of VNP20009 observed within CAM

tumor tissue sections at 3 DPI. LPS (green) and Hoechst (blue). Elongated bacteria cells

are outlined with white and red lines. (F) Intracellular *Salmonella* WT and VNP20009

CFU recovered per gram of tumor at 1 DPI. The statistical significance was determined

by unpaired t-test. (G-H) TNF-α and IL-1β expression level in tumor lysates at 2 HPI,

measured by ELISA, data were expressed as pg per mg of total protein. The results

represent the mean $\pm$ SD. Statistical significance was by one-way ANOVA with Sidak's

multiple comparison test. Not significant (ns), P-value * ≤ 0.05, ** ≤ 0.01, *** ≤ 0.001,

**** ≤ 0.0001. N = 2 with 5-10 tumor-bearing chick embryos per group.

References

1.   Pawelek JM, Low KB, Bermudes D. 1997. Tumor-targeted Salmonella as a novel

anticancer vector. Cancer Res 57:4537–4544.

2.   Leschner S, Weiss S. 2010. Salmonella—allies in the fight against cancer. J Mol

Med 88:763–773.

3.   Avogadri F, Martinoli C, Petrovska L, Chiodoni C, Transidico P, Bronte V, Longhi

R, Colombo MP, Dougan G, Rescigno M. 2005. Cancer Immunotherapy Based on

Killing of Salmonella-Infected Tumor Cells. Cancer Res 65:3920–3927.

4.   Kaimala S, Al-Sbiei A, Cabral-Marques O, Fernandez-Cabezudo MJ, Al-Ramadi

BK. 2018. Attenuated Bacteria as Immunotherapeutic Tools for Cancer Treatment.

Front Oncol 8:136.

5.   Saccheri F, Pozzi C, Avogadri F, Barozzi S, Faretta M, Fusi P, Rescigno M. 2010.

Bacteria-Induced Gap Junctions in Tumors Favor Antigen Cross-Presentation and

Antitumor Immunity. Sci Transl Med 2:44ra57-44ra57.

6.   Zhou S, Gravekamp C, Bermudes D, Liu K. 2018. Tumor-targeting bacteria

engineered to fight cancer. Nat Rev Cancer 18:727–743.

7.   Giannella RA. 1996. Salmonella, p. . *In* Baron, S (ed.), Medical Microbiology, 4th

ed. University of Texas Medical Branch at Galveston, Galveston (TX).

8.   Sys G, Van Bockstal M, Forsyth R, Balke M, Poffyn B, Uyttendaele D, Bracke M,

De Wever O. 2012. Tumor grafts derived from sarcoma patients retain tumor

morphology, viability, and invasion potential and indicate disease outcomes in the

chick chorioallantoic membrane model. Cancer Lett 326:69–78.

9.   Kain KH, Miller JWI, Jones-Paris CR, Thomason RT, Lewis JD, Bader DM,

Barnett JV, Zijlstra A. 2014. The chick embryo as an expanding experimental model

for cancer and cardiovascular research. Dev Dyn Off Publ Am Assoc Anat

243:216–228.

10.   Mitrevska K, Merlos Rodrigo MA, Cernei N, Michalkova H, Splichal Z, Hynek D,

Zitka O, Heger Z, Kopel P, Adam V, Milosavljevic V. 2023. Chick chorioallantoic

membrane (CAM) assay for the evaluation of the antitumor and antimetastatic

activity of platinum-based drugs in association with the impact on the amino acid

metabolism. Mater Today Bio 19:100570.

11. Miebach L, Berner J, Bekeschus S. 2022. In ovo model in cancer research and

tumor immunology. Front Immunol 13.

12. Ishihara M, Hu J, Wong A, Cano-Ruiz C, Wu L. 2019. Chapter Five - Mouse- and

patient-derived CAM xenografts for studying metastatic renal cell carcinoma, p. 59–

80. *In* Tamanoi, F (ed.), The Enzymes. Academic Press.

13. Fischer D, Fluegen G, Garcia P, Ghaffari-Tabrizi-Wizsy N, Gribaldo L, Huang RY-

J, Rasche V, Ribatti D, Rousset X, Pinto MT, Viallet J, Wang Y, Schneider-Stock

R. 2022. The CAM Model—Q&A with Experts. Cancers 15:191.

14. Mon KKZ, Si Z, Chan-Park MB, Kenney LJ. 2022. Polyimidazolium Protects

against an Invasive Clinical Isolate of Salmonella Typhimurium. Antimicrob Agents

Chemother 66:e00597-22.

15. Low KB, Ittensohn M, Luo X, Zheng L-M, King I, Pawelek JM, Bermudes D. 2004.

Construction of VNP20009, p. 47–59. *In* Springer, CJ (ed.), Suicide Gene Therapy:

Methods and Reviews. Humana Press, Totowa, NJ.

16. Clairmont C, Lee KC, Pike J, Ittensohn M, Low KB, Pawelek J, Bermudes D,

Brecher SM, Margitich D, Turnier J, Li Z, Luo X, King I, Zheng LM. 2000.

Biodistribution and Genetic Stability of the Novel Antitumor Agent VNP20009, a

Genetically Modified Strain of Salmonella typhimuvium. J Infect Dis 181:1996–

2002.

17. Low KB, Ittensohn M, Le T, Platt J, Sodi S, Amoss M, Ash O, Carmichael E,

Chakraborty A, Fischer J, Lin SL, Luo X, Miller SI, Zheng L, King I, Pawelek JM,

Bermudes* D. 1999. Lipid A mutant Salmonella with suppressed virulence and

TNFα induction retain tumor-targeting in vivo. 1. Nat Biotechnol 17:37–41.

18. Luo X, Li Z, Lin S, Le T, Ittensohn M, Bermudes D, Runyab JD, Shen S, Chen J,

King IC, Zheng L. 2001. Antitumor Effect of VNP20009, an Attenuated Salmonella

, in Murine Tumor Models. Oncol Res Featur Preclin Clin Cancer Ther 12:501–508.

19. King I, Luo X, Feng M, Ittensohn M, Li Z, Belcourt M, Lin S, Le T, Pike J, Troy K,

Sznol M, Clairmont C, Bermudes D, Zheng L-M. 2000. Tumour therapy using

Salmonella. Emerg Drugs 5:211–219.

20. Toso JF, Gill VJ, Hwu P, Marincola FM, Restifo NP, Schwartzentruber DJ, Sherry

RM, Topalian SL, Yang JC, Stock F, Freezer LJ, Morton KE, Seipp C, Haworth L,

Mavroukakis S, White D, MacDonald S, Mao J, Sznol M, Rosenberg SA. 2002.

Phase I Study of the Intravenous Administration of Attenuated Salmonella

typhimurium to Patients With Metastatic Melanoma. J Clin Oncol Off J Am Soc

Clin Oncol 20:142–152.

21. Tseng H, Gage JA, Raphael RM, Moore RH, Killian TC, Grande-Allen KJ, Souza

GR. 2013. Assembly of a Three-Dimensional Multitype Bronchiole Coculture

Model Using Magnetic Levitation. Tissue Eng Part C Methods 19:665–675.

22. Zanoni M, Piccinini F, Arienti C, Zamagni A, Santi S, Polico R, Bevilacqua A,

Tesei A. 2016. 3D tumor spheroid models for in vitro therapeutic screening: a

systematic approach to enhance the biological relevance of data obtained. Sci Rep

6:19103.

23.   Crull K, Bumann D, Weiss S. 2011. Influence of infection route and virulence

factors on colonization of solid tumors by Salmonella enterica serovar

Typhimurium. FEMS Immunol Med Microbiol 62:75–83.

24.   Wang C-Z, Kazmierczak RA, Eisenstark A. 2016. Strains, Mechanism, and

Perspective: Salmonella-Based Cancer Therapy. Int J Microbiol 2016:5678702.

25.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,

Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein

V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for

biological-image analysis. Nat Methods 9:676–682.

26.   Toley BJ, Forbes NS. 2012. Motility is critical for effective distribution and

accumulation of bacteria in tumor tissue. Integr Biol 4:165–176.

27.   Thornlow DN, Brackett EL, Gigas JM, Van Dessel N, Forbes NS. 2015. Persistent

enhancement of bacterial motility increases tumor penetration. Biotechnol Bioeng

112:2397–2405.

28.   Coutermarsh-Ott SL, Broadway KM, Scharf BE, Allen IC. 2017. Effect of

Salmonella enterica serovar Typhimurium VNP20009 and VNP20009 with restored

chemotaxis on 4T1 mouse mammary carcinoma progression. Oncotarget 8:33601–

33613.

29. Homma M, DeRosier DJ, Macnab RM. 1990. Flagellar hook and hook-associated

proteins of *Salmonella typhimurium* and their relationship to other axial components

of the flagellum. J Mol Biol 213:819–832.

30. Cirillo DM, Valdivia RH, Monack DM, Falkow S. 1998. Macrophage-dependent

induction of the Salmonella pathogenicity island 2 type III secretion system and its

role in intracellular survival. Mol Microbiol 30:175–188.

31. Main-Hester KL, Colpitts KM, Thomas GA, Fang FC, Libby SJ. 2008. Coordinate

regulation of Salmonella pathogenicity island 1 (SPI1) and SPI4 in Salmonella

enterica serovar Typhimurium. Infect Immun 76:1024–1035.

32. Haraga A, Ohlson MB, Miller SI. 2008. Salmonellae interplay with host cells. 1. Nat

Rev Microbiol 6:53–66.

33. Shetty D, Kenney LJ. A pH-sensitive switch activates virulence in Salmonella. eLife

12:e85690.

34. Crull K, Rohde M, Westphal K, Loessner H, Wolf K, Felipe-López A, Hensel M,

Weiss S. 2011. Biofilm formation by Salmonella enterica serovar Typhimurium

colonizing solid tumours. Cell Microbiol 13:1223–1233.

35. Desai SK, Winardhi RS, Periasamy S, Dykas MM, Jie Y, Kenney LJ. 2016. The

horizontally-acquired response regulator SsrB drives a Salmonella lifestyle switch

by relieving biofilm silencing. eLife 5:e10747.

36.   Low KB, Ittensohn M, Luo X, Zheng L-M, King I, Pawelek JM, Bermudes D. 2004.

Construction of VNP20009: a novel, genetically stable antibiotic-sensitive strain of

tumor-targeting Salmonella for parenteral administration in humans. Methods Mol

Med 90:47–60.

37.   Garcia P, Wang Y, Viallet J, Macek Jilkova Z. 2021. The Chicken Embryo Model:

A Novel and Relevant Model for Immune-Based Studies. Front Immunol 12.

38.   Kim J-E, Phan TX, Nguyen VH, Dinh-Vu H-V, Zheng JH, Yun M, Park S-G, Hong

Y, Choy HE, Szardenings M, Hwang W, Park J-A, Park S, Im S-H, Min J-J. 2015.

Salmonella typhimurium Suppresses Tumor Growth via the Pro-Inflammatory

Cytokine Interleukin-1β. Theranostics 5:1328–1342.

39.   Leschner S, Westphal K, Dietrich N, Viegas N, Jablonska J, Lyszkiewicz M,

Lienenklaus S, Falk W, Gekara N, Loessner H, Weiss S. 2009. Tumor Invasion of

Salmonella enterica Serovar Typhimurium Is Accompanied by Strong Hemorrhage

Promoted by TNF-α. PLOS ONE 4:e6692.

40.   Avogadri F, Martinoli C, Petrovska L, Chiodoni C, Transidico P, Bronte V, Longhi

R, Colombo MP, Dougan G, Rescigno M. 2005. Cancer Immunotherapy Based on

Killing of Salmonella-Infected Tumor Cells. Cancer Res 65:3920–3927.

41.   Pawelek JM, Sodi S, Chakraborty AK, Platt JT, Miller S, Holden DW, Hensel M,

Low KB. 2002. Salmonella pathogenicity island-2 and anticancer activity in mice.

10. Cancer Gene Ther 9:813–818.

42.   Loessner H, Endmann A, Leschner S, Westphal K, Rohde M, Miloud T,

Hämmerling G, Neuhaus K, Weiss S. 2007. Remote control of tumour-targeted

Salmonella enterica serovar Typhimurium by the use of l-arabinose as inducer of

bacterial gene expression in vivo. Cell Microbiol 9:1529–1537.

43.   Silva-Valenzuela CA, Desai PT, Molina-Quiroz RC, Pezoa D, Zhang Y, Porwollik

S, Zhao M, Hoffman RM, Contreras I, Santiviago CA, McClelland M. 2016. Solid

tumors provide niche-specific conditions that lead to preferential growth of

Salmonella. Oncotarget 7:35169.

44.   Valdes TI, Kreutzer D, Moussy F. 2002. The chick chorioallantoic membrane as a

novel in vivo model for the testing of biomaterials. J Biomed Mater Res 62:273–

282.

45.   Kogut M, Lowry V, Moyes R, Bowden L, Bowden R, Genovese K, Deloach. 1998.

Lymphokine-augmented activation of avian heterophils. Poult Sci 77:964–971.

46.   Kogut MH, Holtzapple C, Lowry VK, Genovese K, Stanker LH. 1998. Functional

responses of neonatal chicken and turkey heterophils following stimulation by

inflammatory agonists. Am J Vet Res 59:1404–1408.

47.   Li M, Lu M, Lai Y, Zhang X, Li Y, Mao P, Liang Z, Mu Y, Lin Y, Zhao AZ, Zhao

Z, Zhou S, Li F. 2020. Inhibition of acute leukemia with attenuated Salmonella

typhimurium strain VNP20009. Biomed Pharmacother 129:110425.

48.   Liu X, Guo Y, Sun Y, Chen Y, Tan W, Min J-J, Zheng JH. 2022. Comparison of

Anticancer Activities and Biosafety Between Salmonella enterica Serovar

Typhimurium ΔppGpp and VNP20009 in a Murine Cancer Model. Front Microbiol

13.

49. Ahmed SG, Oliva G, Shao M, Wang X, Mekalanos JJ, Brenner GJ. 2022.

Intratumoral injection of schwannoma with attenuated Salmonella typhimurium

induces antitumor immunity and controls tumor growth. Proc Natl Acad Sci

119:e2202719119.

50. Karsten V, Murray SR, Pike J, Troy K, Ittensohn M, Kondradzhyan M, Low KB,

Bermudes D. 2009. msbB deletion confers acute sensitivity to CO_2 in Salmonella

enterica serovar Typhimurium that can be suppressed by a loss-of-function mutation

in zwf. BMC Microbiol 9:170.

51. Murray SR, Bermudes D, de Felipe KS, Low KB. 2001. Extragenic Suppressors of

Growth Defects in msbB Salmonella. J Bacteriol 183:5554–5561.

52. Vander Heiden MG, Cantley LC, Thompson CB. 2009. Understanding the Warburg

Effect: The Metabolic Requirements of Cell Proliferation. Science 324:1029–1033.

53. de Noiron J, Hoareau M, Colin J, Guénal I. 2021. Apoptosis Quantification in

Tissue: Development of a Semi-Automatic Protocol and Assessment of Critical

Steps of Image Processing. 10. Biomolecules 11:1523.

Cell pellets + matrigel mixture

A549 lung cancer cells

ED 0

Set eggs horizontally
(37°C, 60% humidity)

ED 3

Remove 4ml of albumin
(CAM drop)

ED 7

Inoculate A549 cancer cells
(~2 x 10^6 cells/egg)

ED 11

Inoculate *Salmonella typhimurium*
(~50 CFU/egg)

Tumor

Liver

ED 12, ED 14 (1 DPI & 3 DPI)

Harvest tumor & liver
CFU enumeration
Formalin-Fixed Paraffin-Embedded (FFPE)

Live in-ovo Imaging

Created with BioRender.com

(A)

Direct Inoculation Route

WT

1000 µm | 1000 µm | 1000 µm

Δ*flgk*

1000 µm | 1000 µm | 1000 µm

Distant Inoculation Route

WT

1000 µm | 1000 µm | 1000 µm

Δ*flgk*

1000 µm | 1000 µm | 1000 µm

(D)

Δ*flgk* Top

100 µm | 10 µm

Δ*flgk* Middle

100 µm | 10 µm

Δ*flgk* Bottom

100 µm | 10 µm

(B) Direct Inoculation

ns

WT   Δ*flgk*

1 DPI

(C) Distant Inoculation

ns

WT   Δ*flgk*

1 DPI