Evaluation of Tumor-colonizing *Salmonella* Strains using the Chick Chorioallantoic Membrane Model

Khin K. Z. Mon[1], and Linda J. Kenney[1,2*].

[1]Department of Biochemistry and Molecular Biology, [2]Sealy Center for Structural Biology, University of Texas Medical Branch, Galveston, Texas, USA

*Address correspondence to: likenney@utmb.edu

Keywords: Chick Chorioallantoic Membrane, Tumor Colonizing *Salmonella,* CAM tumor, Therapeutic agent, 3D tumor spheroids

Abbreviations: STm, *Salmonella* Typhimurium*;* multiplicity of infection, MOI; h, hour; min, minutes; ED, embryonic development day; dpi, days post-infection; HPI, h post-infection; SD, standard deviation; CAM, chorioallantoic membrane; Lactate Dehydrogenase, LDH; Hematoxylin and Eosin (H and E); WT, wildtype;

ABSTRACT

The chick embryo chorioallantoic membrane (CAM) tumor model is a valuable preclinical model for studying the tumor colonizing process of *Salmonella enterica* serovar Typhimurium. It offers advantages such as cost-effectiveness, rapid turnaround, reduced engraftment issues, and ease of observation. In this study, we explored and validated the applicability of the partially immune-deficient CAM tumor model. Herein, we demonstrate that *Salmonella* preferentially colonizes tumors and directly causes tumor cell death. Bacterial migration, tumor colonization, and intra-tumor distribution did not require flagellar-mediated motility. Both tumor colonization and invasion were independent of known virulence factors, as an extracellular lifestyle of *Salmonella* predominated. Surprisingly, the extracellular residence of *Salmonella* on CAM tumors did not require biofilm formation. We evaluated our wild-type parental strain compared to an attenuated clinical strain VNP20009 and discovered a reduced tumor colonization capability of VNP20009. The inability to effectively colonize tumor cells potentially explains the reduced anti-tumor efficacy of VNP20009. Our work establishes the xenograft CAM model as an informative and predictive screening platform for studying tumor-colonizing *Salmonella*.

IMPORTANCE

Cancer has a major impact on society, as it poses a significant health burden to human populations worldwide. *Salmonella* Typhimurium has demonstrated promise in cancer treatment by exerting direct tumoricidal effects and enhancing host-mediated anti-tumor immunity. A general understanding of its pathogenesis and the relative ease of genetic manipulation support the development of attenuated strains for therapeutic use. The chick

embryo chorioallantoic membrane tumor model represents a suitable screening platform

to accelerate the development of therapeutic strains. It allows for rapid evaluation of

Salmonella strains to assess their efficacy and potential as oncolytic agents. The present

study establishes that the *in ovo* tumor model can be utilized as a preclinical tool for

evaluating oncolytic *Salmonella*, bridging the gap between *in vitro* and *in vivo* screening.

INTRODUCTION

In the field of bacterial-mediated cancer therapy, *Salmonella enterica* serovar

Typhimurium (STm) has emerged as an ideal therapeutic candidate. This is in part due to

its intrinsic tumor targeting ability (1), its facultative anaerobic properties, which allow

colonization in both hypoxic and oxygenated regions of heterogeneous tumor

microenvironments (2), its invasive capability (3), and its direct or immune-mediated

anti-tumor effects (3–6). STm infection mainly causes self-limiting gastrointestinal

disease in healthy humans but can sometimes manifest as extraintestinal systemic

infections in immune-compromised individuals (7). To promote its safe application in

cancer patients, bacterial pathogenic properties need to be eliminated or attenuated

without compromising the tumor colonizing capability, and thus subsequent anti-tumor

potency. Therefore, tuning the balance between virulence attenuation and tumor-

colonization capability is an important criterion in the design of suitable STm strains.

The chick embryo chorioallantoic membrane (CAM) has long been used as a

robust model for cancer and cardiovascular research, as well as for preclinical evaluation

of cancer drug therapies (8–10). The CAM model is an immunodeficient model that is

receptive to xenografting of various mammalian tumor cells without inducing a host

immune response (11). The well-vascularized embryonic tissue of the CAM supports and

preserves heterogeneous tumor growth with a much shorter experimental timeline

(typically 3-4 days) for tumor development compared to mouse models, which requires

weeks (12). Moreover, the CAM model also allows real-time visualization of tumor

growth (13) as well as the dynamics of bacterial infection via fluorescence

stereomicroscopy (14). These attributes highlight the CAM tumor model as a promising

valuable platform to evaluate *Salmonella*-based therapies for cancer treatment.

The engineered STm strain VNP20009 is attenuated by purine auxotrophy (a

deletion in *purI*) and lipid A modification (a deletion in *msbB*) (15). It demonstrated an

excellent safety profile (16) with observable tumor growth retardation in a variety of pre-

clinical xenograft mouse models (17–20). Despite the high success rate in the mouse

studies, no anti-tumor effect was observed in humans. The intravenous infusion of the

VNP20009 strain to melanoma patients in a clinical trial failed and was attributed to a

low bacterial tumor colonization and rapid bacterial clearance by the host immune

response (21). The failure of the VNP20009 strain to colonize tumors highlighted the

xenograft mouse model as a poor predictor of therapeutic success in humans. Hence,

there is a need to explore alternative models to overcome some of the translational

limitations associated with traditional mouse models. We hypothesized that the CAM

tumor model might serve as a preliminary screening platform to study the tumor-bacterial

interactions to evaluate the efficacy of tumor-targeted *Salmonella* strains for therapeutic

purposes.

In the current work, we developed the *in ovo* CAM tumor model and compared it

with an *in vitro* three-dimensional (3D) tumor spheroid model to examine the relevance

of the screening data obtained (22). While 3D tumor spheroids can mimic *in vivo* solid

tumor features such as structural organization, cellular assembly and nutrient gradients

(23), they cannot fully simulate the heterogenous tumor microenvironment established

within a living host. Therefore, the CAM tumor model serves as a better pre-clinical

model, bridging an informational gap between *in vitro* cell-based models and *in vivo*

animal-based systems for screening oncolytic *Salmonella*. With the CAM tumor model,

we demonstrated that STm pathogenic traits such as flagella, virulence factors and

biofilm formation were not essential for tumor colonization. We further discovered that

reduced tumor colonization by VNP20009 was attributed to its growth sensitivity within

the tumor microenvironment, which subsequently impacted its anti-tumor effectiveness.

RESULTS

Validation of the CAM model for studying tumor-colonizing *Salmonella*

A general overview of the CAM tumor STm infection model is depicted in Figure

1. We first determined the infectious dose of STm by infecting tumor bearing chick

embryos with different doses of bacteria: approximately 500 colony-forming units

(CFU), 50 CFU, or 5 CFU. Chick embryo mortality was monitored daily post-bacterial

infection with PBS as the control group. The highest inoculum of ~500 CFU caused the

greatest mortality (38% survival) at one day post-infection (dpi) compared to groups that

received lower dosages of ~50 CFU (66% survival) or ~5 CFU (77% survival)

(Supplementary Figure 1). To minimize the mortality of highly susceptible tumor-bearing

chick embryos as well as to ensure effective tumor colonization, we selected the mid-

range ~50 CFU as the optimal infectious dose.

The important features of tumor-targeting STm are the higher tumor specificity in

colonization compared to organ colonization, and the ability to penetrate tumor tissue

(24, 25). These criteria were tested first in the CAM tumor model. At 1 dpi, WT STm

accumulated preferentially in tumors over livers of chick embryos at a ratio of 1000:1.

The mean CFU/g in tumors was 1.9 x 10^8 CFU/g compared to livers at 1.9 x 10^5 CFU/g

(Figure 2A). Next, we examined the intra-tumoral distribution of STm within tumor

tissue. Infected whole tumors were harvested, embedded in paraffin and sectioned at

different tissue depths (top, middle and bottom). The tumor sections were stained with

antibodies against bacterial lipopolysaccharide (green), nuclei (blue) and e-cadherin

(magenta) and their immunofluorescence was examined by confocal microscopy. STm

was observed at all tumor tissue depths as evident by lipopolysaccharide staining (LPS =

green, second panel Figure 2B and Supplementary Figure 3). Specifically, bacteria were

mostly present in the metastatic tumor cell regions, where loss of cell-to-cell adhesion

junctions was apparent as a loss of e-cadherin staining.

STm colonization drove direct tumor apoptosis in the CAM model

We next tested the hypothesis that STm infection directly promoted tumor cell

death. A trypan blue exclusion assay was performed on dissociated tumor cells at 3 dpi to

test *ex vivo* tumor cell viability. STm infection significantly reduced the percentage of

tumor cell viability (74%) compared to PBS-treated tumors (84%) (Figure 2C). Because

the presence of STm was detected throughout the entire tumor (Figure 2B), it raised the

question of whether its localization directly induced tumor cell apoptosis. We utilized the

terminal deoxynucleotidyl transferase-mediated deoxyuridine triphosphate nick end-

labeling (TUNEL) assay to detect *Salmonella*-induced DNA fragmentation in tumor

cells. A marked accumulation of TUNEL-positive cells (red signals) was observed in

STm- infected tumors relative to PBS-infected tumors sections at the top, middle and

bottom levels (Figure 2D). Immunofluorescence images of TUNEL-stained tumor sections were further quantified using Fiji software. The percentage of TUNEL-positive nuclei in STm infected tumor tissue sections was significantly higher than PBS tumors (52% vs 33%, respectively), indicating that tumor cells underwent increased apoptosis following bacterial infection (Figure 2E). Lastly, the histological changes in Hematoxylin and Eosin (H and E)-stained tumor sections were analyzed. A more extensive necrotic area in *Salmonella*-infected tumors (22%) was detected compared to the PBS control group (4%) (Figure 2F, 2G).

Bacterial motility was not required for tumor colonization or intra-tumoral distribution

The prevailing hypothesis regarding a role for flagella in tumor colonization is that motility is essential for bacteria to effectively target, colonize, and disperse within the tumor environment (26–28). To test this hypothesis, WT or a *ΔflgK* null strain were directly inoculated onto CAM tumors. The *flgK* gene encodes a flagellar hook-associated protein; its deletion renders *Salmonella* non-motile (29). Live *in ovo* images of the bacterial colonization pattern, as well as CFU enumeration from tumors was comparable between the WT and the *ΔflgK* null strain (Figure 3A, B). We reasoned that the role of flagella might be dispensable when bacteria were directly inoculated onto the tumor, rendering colonization a passive process. Therefore, we also tested distant inoculation route by depositing bacteria a short distance away from the tumor (~0.5 cm) on filter papers to test whether *Salmonella* required flagellar-mediated motility to migrate towards the tumor for colonization. Despite the loss in motility, the *ΔflgK* strain was able to migrate, colonize and accumulate at tumors to the same extent as the WT (Figure 3A, C).

The intra-tumoral distribution pattern of the non-motile *ΔflgK* strain was also comparable

to that of the motile WT (Figure 2B), indicating that the *ΔflgK* strain was able to disperse

and penetrate all depths of the tumor (Figure 3D). These results provided strong evidence

that STm migration, tumor colonization and intra-tumoral distribution were not

dependent on bacterial motility in the CAM tumor model.

3D tumor spheroids do not recapitulate the CAM-tumor model with respect to key

virulence factors encoded on pathogenicity islands 1 and 2

During infection, STm utilizes *Salmonella* Pathogenicity Island-1 (SPI-1) to

invade host intestinal epithelial cells and SPI-2 associated genes for intracellular growth

and survival within host macrophages (30–33). It was therefore of interest to investigate

whether bacterial invasion or replication within tumor cells was also dependent on SPI-1

and SPI-2. To address this question, we first employed the *in vitro* 3D tumor spheroids

model before assessing their roles *in vivo* in the CAM tumor model. The use of 3D

magnetic cell culture technology produced structurally relevant tumor spheroids that

mimicked some features of *in vivo* solid tumors. The spatial architecture with regard to

oxygen and nutrient gradients, extracellular matrix, and cell-to-cell interactions is

maintained. Within tumor spheroids, intracellular invasion and replication rates of STm

strains were examined using a gentamicin protection assay. The elimination of *hilD* (the

SPI-1 master regulator) resulted in a significant invasion defect with a mean reduction of

1.4 log CFU/well compared to WT (Figure 4A), suggesting that SPI-1 played a

significant role in early bacterial invasion of tumor cells. At 16 HPI, an intracellular

replication rate of both Δ*hilD* (mean log CFU/well = 3.5) and the SPI-2 master regulator,

Δ*ssrB* (mean log CFU/well = 4.5) were significantly reduced relative to WT (mean log

CFU/well = 5.6) (Figure 4A). Consequently, the overall replication rate (16 HPI/ 2 HPI)

was higher for WT (28-fold) compared to either $\Delta hilD$ and $\Delta ssrB$ strains (6-fold and 4-

fold, respectively) (Figure 4B). We next examined the cytotoxic damage induced by

bacterial infection of 3D tumor spheroids using a lactate dehydrogenase (LDH) assay 16

HPI. The amount of LDH release is directly correlated with cell cytotoxicity levels.

Tumor spheroids infected with WT bacteria released significantly higher amounts of

LDH (53% of total maximized LDH release control at 100%) compared to SPI-1 (22%)

or SPI-2 (33%) deficient strains (Figure 4C). Whole-mount immunofluorescence staining

of 3D tumor spheroids exhibited a specific cellular structural organization. Proliferating

tumor cells were observed in the periphery, with an inner layer of quiescent cells and a

necrotic core region. These were evident by distinct fluorescence intensity profiles of the

cell proliferation marker Ki67, in which the intensity varied from high to low from the

periphery to the core (Figure 4D). From representative images of infected 3D tumor

spheroids at 16 HPI, both SPI-1- and SPI-2-deficient strains were observed at reduced

levels compared to the WT (Figure 4D). These results were consistent with the

quantification of colony-forming units (Figure 4A).

We then investigated whether maximal tumor colonization and intracellular

invasion or replication required SPI-1 or SPI-2 in the CAM tumor model. Using an ex

vivo gentamicin assay, we quantified both extracellular and intracellular bacterial load

recovered from the CAM-tumors. Both SPI-1 (*ΔhilD)* and SPI-2 (*ΔssrB*)-deficient strains

were found to colonize (extracellular) and invade (intracellular) tumor cells to the same

extent as WT at 1 dpi (Figure 5A, B). In the mouse tumor model, *Salmonella* was found

almost exclusively to reside extracellularly, forming biofilms on tumors (34). We

therefore assessed whether *Salmonella* adapted a similar residence lifestyle in our CAM

tumor model. Based on percentage of total tumor-associated bacteria population,

approximately 99% of bacteria were extracellular with ~1% of the bacterial population

found inside the CAM-tumor cells (WT, *ΔhilD* and *ΔssrB*) (Figure 5C). Hence, this

finding corroborated that the loss of SPI-1 and SPI-2 genes did not impair the tumor-

colonizing ability of STm (Figure 5B), because STm were predominantly extracellular on

the CAM tumors.

To address the question of whether extracellular STm also formed biofilms, we

infected CAM tumors with *Salmonella* WT or a *ΔcsgD* null strain; CsgD is the master

biofilm regulator (35). The extracellular bacterial load recovered indicated that both

strains successfully colonized the tumor to the same extent: a mean log CFU/g of ~7 and

8, respectively at 1- and 3 dpi (Figure 5D). Deletion of the *csgD* gene, rendering the

bacteria unable to form biofilms did not reduce the extracellular bacterial load recovered

from tumors. In contrast to the mouse tumor model (34), the extracellular residence of

STm on CAM tumors was not in the form of biofilms (see Discussion).

How does the clinical strain VNP2009 compare with WT *Salmonella*?

There were discrepancies in reports on the performance of the clinical strain

VNP20009 between the xenograft mouse models (effective tumor colonization with

tumor growth retardation) and the human clinical trial (low bacterial tumor colonization).

It was therefore of interest to evaluate VNP20009 compared to its parental WT strain (15,

36) in both 3D tumor spheroid and CAM tumor models.

The clinical strain VNP20009 replicates poorly within 3D tumor spheroids

We first investigated the performance of the clinical strain VNP20009 within 3D

tumor spheroids. Compared to WT, the VNP20009 strain displayed a modest invasion

defect with a mean reduction of 0.6 log CFU/well (Figure 6A, left columns). However,

the replication defect at 16 HPI was striking, with an overall mean reduction of 1.6 log

CFU/well relative to WT (Figure 6A, right columns). The fold replication of the WT was

28-fold, but VNP20009 only increased 3-fold (Figure 6B). These differences indicated a

stark difference in the ability of the attenuated VNP20009 strain to grow within the tumor

spheroid compared to the WT. We next compared the ability of VNP20009 to induce

tumor cell cytotoxicity with the LDH assay after 16 HPI. WT-infected tumor spheroids

released significantly higher amounts of LDH (53%) than VNP20009 (36%) (Figure 6C).

This result implies that the reduced intracellular replication of VNP20009 within tumor

spheroids has consequences that limit its ability to kill tumor cells (Figure 6A, right

columns). The possibility of differential growth rates between strains contributing to

tumor invasion was ruled out, as the growth rate of VNP20009 was not compromised

under bacterial culture conditions (Supplementary Figure 2). Representative images of

VNP20009-infected tumor spheroids highlighted the reduced intracellular bacteria

compared to WT (Figure 6D), corroborating the bacterial enumeration dataset (Figure

6A, B). Furthermore, tumor spheroid infected with VNP20009 revealed distinct cell

morphological defects of the strain with elongated bacteria cells (Figure 6E), confirming

the strain growth defects within the tumor spheroid (see Discussion).

Reduced CAM tumor colonization of VNP20009 compromised its anti-tumor

immunity

We then examined the tumor colonization ability of VNP2009 in the CAM tumor model.

The effect of VNP20009 administration (~50 CFU) on chick embryo mortality was first

evaluated against the WT and PBS control groups. A significant improvement in survival

rate and prolonged survival time was noted for tumor-bearing chick embryos infected

with VNP20009 compared to the WT (Figure 7A). Chick embryos completely

succumbed to WT infection by four days, whereas VNP-infected embryos survived for

up to seven days. We next assessed the tumor-colonizing capability of VNP20009. Live

in ovo images of mCherry-tagged bacterial strains colonizing the CAM tumor highlighted

the marked reduction in fluorescence signals of VNP20009 at 1 dpi (Figure 7B)

compared to the WT. The image emphasizes the strain attenuation of VNP2009 in tumor

colonization, which was also apparent when we examined the intra-tumoral distribution

(Supplementary Figure 4). Compared to WT, VNP20009 was remarkably reduced

throughout the tumor tissue sections at all depths. This result was further confirmed by an

overall reduction in the extracellular bacterial load recovered from VNP20009-infected

tumors (Figure 7C). At 1 dpi, the extracellular bacterial load of VNP20009 was only ~1%

of WT. A reduction of VNP20009 tumor colonization continued at 3 dpi and was ~36%

of WT colonization (Figure 7C). When we compared the two strains and their ability to

colonize the liver (Figure 7D), VNP2009 exhibited an early defect in dissemination

(~10% of WT). However, the attenuated strain was able to colonize chick embryo livers

to a similar level as the WT by 3 dpi (Figure 7D). This was in contrast to tumor

colonization (Figure 7C), where the rate of VNP2009 colonization remained weaker than

the WT over time. Furthermore, cell morphological defect was observed as elongated

chains of VNP20009 were detected in CAM tumor tissue sections at 3 dpi (Figure 7E).

These chains were strikingly similar to our previous observations in 3D tumor spheroids

(Figure 6E). The CAM-tumor invasion rate of VNP20009 was also compromised

compared to WT invasion rate (intracellular bacteria count, Figure 7F).

Chick embryos possess only partial immunity at early developmental stages

(before ED 9), but as immunocompetence develops, embryos are capable of mounting an

inflammatory response with cytokine expression from host immune cells (37). Induction

of pro-inflammatory cytokines such as tumor necrosis factor-alpha (TNF-α) and

Interleukin-1 beta (IL-1β) following *Salmonella* infection in tumor-bearing hosts are

indicative of host-mediated tumoricidal effects (38). To investigate whether host immune

factors play an indirect role in driving anti-tumor immunity, CAM tumors were infected

with WT, VNP20009 or PBS at ED11, and cytokine expression in tumor lysates at 2 HPI

was measured. TNF-α levels were similar across uninfected and infected tumors (WT and

VNP20009) at 2 HPI (Figure 7H). In contrast, IL-1β was highly expressed in WT-

infected tumors, while IL-1β levels were similar between uninfected (PBS) and

VNP20009-infected tumors at 2 HPI (Figure 7I). This result suggests that reduced tumor

colonization by VNP20009 failed to trigger activation of host-mediated anti-tumor

immunity in the CAM model.

DISCUSSION

To the best of our knowledge, this is the first study to exploit the CAM tumor

model as an *in vivo* model to study *Salmonella*-mediated anticancer therapy. Our work

established that the chick embryo CAM model reproduces key features of tumor targeting

STm reported in the mouse model: (1) tumor specificity in colonization (Figure 2A), (2)

intra-tumoral distribution (Figure 2B), (3) direct induction of tumor cell death (Figure

2C-G) and (4) indirect stimulation of anti-tumor immunity to some degree in the partial

immunocompetent host (Figure 7H). To assess whether the CAM model recapitulated the

published mouse studies, *Salmonella* key pathogenic traits were investigated for their

respective role in tumor colonization and invasion.

Flagellar-dependent motility as a driver of bacterial migration, tumor

colonization, and intra-tumoral distribution was not essential, as non-motile *Salmonella*

(Δ*flgK)* was not impaired for colonization in the CAM model (Figure 3). This finding

agrees with published mouse data indicating that tumor colonization by *Salmonella* is a

passive process (not requiring bacterial motility), where bacterial infection induces a

cytokine storm and hemorrhage, facilitating the deposition of bacteria at the tumor site

(24, 39). In the CAM tumor model, the passive tumor migration process of STm (via a

distant inoculation route, Figure 3A, C) was most likely aided by the active movement

motion of a live chick embryo just beneath the CAM layer.

In our *in vitro* 3D tumor spheroids, both SPI-1 and SPI-2 contributed to STm

intracellular replication within the tumor spheroids (Figure 4). In the mouse model, there

are conflicting reports as to whether SPI-1 or SPI-2 virulence genes are required for

tumor invasion. Although SPI-1-mediated tumor invasion (40) and SPI-2-associated

intra-tumoral replication (41) were reported to be important, in the majority of mouse

tumor studies, virulence functions were not required, as bacteria were extracellular on

tumor sites (24, 34, 42, 43). Consistent with these latter findings, virulence functions

were not required for CAM tumor colonization and invasion (Figure 5A, B). In our

experiments, *Salmonella* strains (WT, Δ*hilD,* Δ*ssrB*) were predominately extracellular on

the CAM tumor (Figure 5C). The inconsistency between our *in vitro* and *in ovo* data is

likely attributed to the fact that 3D spheroids consisted of a single cancer cell-type, which

lacks the heterogeneity and vascularization of the chick CAM model to mimic the

complexity of the tumor microenvironment within the host and its interaction with

bacteria.

In the mouse model, an almost exclusive extracellular residence as biofilms on

solid tumors was reported to be a bacterial protective mechanism against phagocytosis by

host immune cells, specifically neutrophils (34). However, the CAM tumor model did not

reproduce the STm biofilm phenotype, as the Δ*csgD* strain was able to colonize tumors to

similar level as WT (Figure 5D). Immunological differences in the tumor

microenvironment between the two models (mouse vs chick embryos) might explain the

differences in an adaptation of bacterial lifestyles on solid tumors. In contrast to a

mammalian host, chick embryos have heterophils, the avian equivalent of neutrophils, but

lacking myeloperoxidase (37). Although heterophil recruitment to the inflammation site

following lipopolysaccharide stimulation was reported at ED7 (44), the full functionality

of heterophils within developing embryos was only described at ED18, three days before

hatching (45, 46). Our CAM experimental timeline is completed by ED14 due to the low

survival percentage of tumor-bearing chick embryos at an STm infectious dose of ~50

CFU (~6% survival at 3 dpi, Figure 7A). Therefore, the lack of functional heterophils

within the chick embryonic immune system in the current study timeline likely influences

the non-biofilm residence of STm on CAM tumors. As with every model, the CAM

tumor model also has its limitations. As highlighted above, the lack of a functional host

immune system and the short experimental timeline (unable to assess STm impact on

tumor growth or regression) should be considered during experimental design and data

interpretation.

Previous studies examining the anti-tumor efficacy of the highly attenuated STm

strain VNP20009 failed to directly compare the tumor colonizing level of the strain

relative to its parental WT (15, 17, 19, 47–49). Here, we used both the *in vitro* 3D tumor

spheroids and CAM tumor model to provide evidence of reduced tumor colonization,

invasion and replication by VNP20009 compared to WT (Figures 6AB, 7BC, F). In

addition, we observed cell morphology defects of VNP20009 (i.e., elongated bacteria

cells) in both models during infection (Figures 6E, 7E). Although deletion of *msbB* in

VNP20009 has been reported to exhibit severe growth and morphology defects under *in*

vitro growth conditions in the presence of 5% CO_2, acidic pH and high osmolarity (50,

51), our work provides additional evidence of the inherent growth sensitivity of

VNP20009 in the *in vivo* tumor microenvironment (Figure 7E). We reason that this

growth defect is related to cancer metabolism, which undergoes aerobic glycolysis (also

known as Warburg effect), resulting in elevated CO_2 levels and lactic acid in the tumor

microenvironment (52).

A poor human clinical outcome of VNP20009 was mainly the result of low tumor

colonization that subsequently compromised its anti-tumor efficacy (21). In our CAM-

tumor model, we showed that IL-1β expression levels in tumor lysates of VNP20009 and

PBS controls were similar (Figure 7I), suggesting that the reduced presence of

VNP20009 on tumors was insufficient to elicit the anti-tumor immune response in the

CAM model. These results highlighted the importance of maximizing the tumor

colonizing capability of the bacteria strain as a primary factor in strengthening its anti-tumor effectiveness.

In summary, the chick embryo CAM model offers a valuable tool for initial screening and preclinical assessment of tumor-colonizing *Salmonella* strains. Its accessibility, vascularization, and relevance to tumor biology make it a suitable choice for studying bacterial interactions with tumors and assessing potential therapeutic strains for bacterial-mediated cancer therapy.

MATERIALS AND METHODS

Cell culture and bacterial strains growth conditions

A549 human lung carcinoma cells (ATCC CCL-185) were purchased from the American Type Culture Collection (ATCC). The cell line was cultured in DMEM (Gibco 11965-092) supplemented with 10% heat-inactivated fetal bovine serum (GenDEPOT, F0900-050) and 1% penicillin/streptomycin (GenDEPOT, CA005-010), at 37°C with 5% CO_2. The bacterial strains and plasmid used in this study are listed in Supplementary Table 1. Bacterial strains (transformed with reporter plasmid) were grown in Luria-Bertani (LB) medium in presence of 100 µg/ml ampicillin (Sigma, A0166) at 37 °C at 225 rpm overnight. Sub-cultures were grown to the late-log phase to induce SPI-1 associated genes and diluted to the desired dosage for inoculum preparation in both 3D tumor spheroids and CAM tumor models.

3D Spheroid infection model

A549 tumor cell spheroids were formed with the 96-well bioprinting kit (Greiner Bio-One catalog. 655841) according to the manufacturer's guidelines. Cells were first magnetized with NanoShuttle-PL, nanoparticles overnight. Magnetized cells were

counted and seeded in same cell density of ~50,000 cells/well. Using the 96-well

magnetic drive placed at the bottom of the well, magnetized cells were printed into

spheroid for 24 hr. The next day, STm strains grown under the SPI-1 inducing condition

were added into each well at MOI of 100. Bacteria were added for 30 min, high

gentamicin concentration of 100 ug/ml were added for 1.5 h for invasion (2 HPI) and

changed to low gentamicin of 20 ug/ml for remainder of infection process. We confirmed

that all extracellular bacteria were eliminated post-gentamicin treatment by plating the

cell culture medium. At each step of the cell media replacement and wash process, the

96-well plate was placed on the holding drive to hold the spheroids down while aspirating

the solution to prevent cell loss. At both 2 HPI (invasion) and 16 HPI (replication), cell

media were aspirated, washed with phosphate-buffered saline, PBS (Roche,

11666789001), and replaced with 1% Triton X-100 (Sigma, T8787-100ml) for 10 min to

lyse the cells. Spheroids were homogenized by the vigorous pipetting method. Lysates

were then serially diluted and plated for intracellular bacterial enumeration.

Whole-mount Immunofluorescence staining of 3D tumor spheroids

At the end of 16 hr STm infection, tumor spheroids were fixed in 4% paraformaldehyde

for ~5 h, washed with PBS thrice, and then processed for whole-mount

immunofluorescence staining. Each tumor spheroids were individually picked up using

the magpen with teflon caps (Catalog no: 657850, Grenier, Bio-One), during each step of

the staining process for solution replacement. Antigen retrieval was performed by

transferring spheroids into a citrate buffer (Sigma, C9999-100ml) solution for 20 minutes

at 99°C water bath before allowing the solution to cool to room temperature. The

permeabilization step was performed with 0.3% Triton-X/PBS for 15 min followed by

incubation with blocking buffer, 5% bovine serum albumin in PBS (BSA; Sigma-

Aldrich, Catalog A7030) for 1 hr. Spheroids were then incubated with primary antibody,

recombinant anti-Ki67 (Abcam Catalog ab16667) diluted in blocking solution (1:200),

overnight at 4°C in the dark. The next day, spheroids were incubated with a secondary

antibody to Ki67, Alexa Fluor 488 goat anti-rabbit IgG H&L (Abcam Catalog ab150081),

and Hoechst (blue, Invitrogen, H3570) for 1 hr. Whole spheroids were then mounted onto

the slide with a prolong gold antifade mountant (Invitrogen Catalog: P36934), coverslip

sealed with nail polish and dried overnight in the dark. Whole-mount spheroids were

imaged with Olympus SpinSR-10 spinning disk confocal at 10x air objective (NA 0.40),

and 100x oil objective (NA 1.5). Acquired microscopy images were processed with Fiji

software.

Lactate Dehydrogenase Assay

The cell culture supernatants from A549 tumor spheroids infected with STm strains were

collected at 16 hr post-infection timepoint and Lactate Dehydrogenase (LDH) assay was

performed with CytoTox96 non-radioactive cytotoxicity assay according to the

manufacturer's guideline (Promega, Catalog no: G1780). Maximum releasable LDH

activity (100%) in the cells was achieved by treating the uninfected spheroids with lysis

buffer for 45 min at room temperature and maximum LDH release control measurement

was used to calculate the percentage of cytotoxicity in STm-infected cells. The

colorimetric absorbance of the samples was measured with a plate reader at 490 nm (Bio-

rad iMark microplate absorbance reader #1681130).

Embryonated chicken chorioallantoic membrane (CAM) infection model

Specific pathogen-free (SPF), premium fertilized chicken eggs were purchased from

AVS bio and set horizontally in a Rcom Max 50 incubator at 37 °C and 60% humidity.

On embryonic development day three (ED3), the chorioallantoic membrane (CAM) of

the chick embryo was dropped by inserting a thin needle at the apex of the egg and

removing ~4ml of albumin with syringe and 21G needle. A small window was made on

top of the eggshell and resealed with 3M Tegaderm TM transparent film dressing. On

ED7, A549 cell pellets (~2 x 10^6 cells/egg) were mixed with 50 µl Matrigel (Corning

354234 or 354263) and were grafted into the center of a sterilized silicone ring placed on

top of the CAM layer. Eggshell windows were then re-sealed with Tegaderm and put

back into the incubator to allow tumor growth for four days. On ED11, the silicone ring

was first carefully removed from the CAM without disrupting the established tumor

before the application of the bacterial inoculum (~50 CFU) directly onto the tumor with a

pipette. Bacterial inoculums were plated each time to confirm that a similar rate of dose

amount was applied for all strains. For distant inoculation route experiment, bacteria were

deposited onto a sterile cut-out filter paper, placed a short distance away from the tumor

(~0.5 cm). At different time points of post-bacterial infection, tumors, and livers were

harvested for enumeration of viable bacteria by homogenization, serial dilution, and

plating methods.

CAM tumor dissociation for tumor cell viability

The whole tumors harvested were placed in the sterile petri dish and minced with scissors

in DPBS++ solution (Corning 21-030-CV). Minced tissue fragments with the solution

were transferred into 15 ml centrifuge tubes. DPBS++ solution was removed and

replaced with 2 ml (1mg/ml) of collagenase (Millipore Sigma SCR103). Tubes were

turned over several times and incubated at 37 °C for a minimum of 3 h. At the end of

incubation, 5 ml of DMEM media (without antibiotic) were added to each tube and

resuspended thoroughly by repeated pipetting from 20 to 40 times for complete tissue cell

dissociation. Tissue fragments with media solution were filtered through a 40 µm nylon

mesh cell strainer (Falcon 352340). Filtered cell suspensions were then centrifuged at 500

x g for 5 minutes. Supernatants were removed and the dissociated cell pellet was

resuspended with fresh DMEM media. The total viable tumor cell count was obtained

with the trypan blue staining method.

Ex vivo **gentamicin protection assay**

To evaluate the percentage of intracellular bacteria in the tumor tissue, an *ex vivo*

gentamicin protection assay was performed. The whole tumor tissue was homogenized in

1 ml of cold PBS. A 0.5 ml aliquot of tumor homogenate suspension was serially diluted

and plated to determine the extracellular bacterial load in the tumor. The remaining 0.5

ml aliquot was treated with 200 µg/ml of gentamicin for 90 minutes at 37 °C to kill

extracellular bacteria. Tissue suspensions were then centrifuged at 5000 rpm for 10 min,

washed twice with PBS, and resuspended in 1% Triton X-100 PBS for 15 min at 37 °C to

lyse the cells. The lysate was then serially diluted and plated for the intracellular bacteria

enumeration.

Histology and immunofluorescence staining

Tumors harvested at 1 and 3 dpi were fixed in 10% formalin (Sigma, H7501128) before

samples were sent to the UTMB histopathology core facilities for paraffin embedding,

tissue sections as well as Hemolysin and eosin (H&E) staining. H&E-stained tumor

sections were sent to the Research Histology Core Laboratory (RHCL) at MD Anderson

for external histopathologist assessment. The sample thickness was 5 µm at each depth:

top, middle, and bottom cuts of the tumor samples. Tissues were deparaffinized and

rehydrated in a series of xylene (Sigma, 534056-500ML) and ethanol (Fisher Scientific,

BP2818500) washes. Deparaffinized and rehydrated tumor sections were boiled in

antigen retrieval solution, citrate buffer for 35 min at 99°C water bath before allowing the

solution to cool to room temperature. The permeabilization step was performed with

0.3% Triton-X/PBS for 15 min followed by incubation with blocking buffer, 5% bovine

serum albumin in PBS for 1 hr. Tumor sections were then stained and prepared for

immunofluorescence with lipopolysaccharides, LPS (Abcam, ab35156) 488 (green) for

Salmonella, E-cadherin (BD Biosciences, 610182) 647 (far-red), and Hoechst (blue) for

the nucleus. Images were acquired with an Olympus Super Res Spinning Disk, SpinSR-

10 microscope at 100x oil objective (NA 1.5) and super resolution imaging with optical

zoom, 320x of 100x region of interest (ROI) for VNP20009 elongated cells observation.

Acquired microscopy images were processed with Fiji software.

TUNEL assay and apoptosis quantification with Fiji

Detection of apoptotic cells on tumor tissue sections was performed with Click-iT Plus

TUNEL assay (Invitrogen, Thermo Fisher Scientific Lot 2559160) according to the

manufacturer's guidelines. Immunofluorescence images were acquired with an Olympus

Super Res Spinning Disk, SpinSR-10 microscope at 100x oil objective (NA 1.5).

Quantification of TUNEL-positive nuclei as well as DAPI nuclei was performed with Fiji

(53) following the optimized and unbiased image processing protocol. Briefly, images

were first split into separate color channels and compressed with z-axis maximum

intensity projection. Background noise reduction is performed by subtracting background

values from each channel. Images are then converted to greyscale with 8-bit. Binarization

thresholding function available in Fiji was utilized and thresholding algorithms that

capture the best outline of stained cells for each channel were selected. Moments and Li

thresholding methods were opted for highlighting TUNEL-positive nuclei and DAPI-

positive cells, respectively. Lastly, the analyze particle's function was applied for the

count read-outs. The percentage of TUNEL-stained cells was calculated against the total

cell numbers (DAPI-positive cells) within the same region.

ELISA

Tumor tissues were collected at 2 hr post-infection (HPI). Tumor tissues were

homogenized in Radioimmunoprecipitation assay (RIPA) buffer (Cell Signaling

Technology, 9806S) containing protease inhibitor cocktail (Roche, 11836153001).

Protein lysates were collected from supernatant after a round of centrifugation at 13,000 x

g for 20 minutes, 4°C and were stored immediately at -80 °C until analysis. Total protein

concentration from tumor lysates was measured with Pierce BCA assay

(ThermoScientific 23225) following the manufacture's instruction. Tumor lysates were

normalized for total protein content and analyzed by chicken-specific enzymatic-linked

immunosorbent assay (ELISA) kits for expression levels of TNF-alpha (MyBioSource

MBS:260419) and IL-1beta (MyBioSource MBS:2024496) according to the

manufacturer's guidelines.

Statistical Analysis

All statistical analysis were performed with GraphPad Prism 10.2.1.

Acknowledgements: This work was supported by Cancer prevention and Research

Institute of Texas (CPRIT) Grant RP2000650 awarded to LJ.K.

Conflict of interests: All authors declared no conflict of interests.

Figure 1: Experimental scheme of the CAM tumor, STm infection model. ED0: Specific

pathogen-free (SPF), premium fertilized chicken eggs were purchased from AVS bio and

set horizontally in an Rcom Max 50 incubator at 37 °C and 60% humidity. ED3: The

chorioallantoic membrane (CAM) of the chick embryo was dropped by inserting a thin

needle at the apex of the egg and removing ~4ml of albumin with syringe and 21G

needle. A small window was made on top of the eggshell and resealed with transparent

Tegaderm. ED7: A549 cell pellets (~2 x 10^6 cells/egg) with 50 µl Matrigel mixture was

grafted into the center of a sterilized silicone ring placed on top of the CAM layer with a

pipette. ED11:

the silicone ring was first carefully removed from the CAM without disrupting the

established tumor before direct inoculation of the STm strains with a pipette (~50

CFU/egg) onto established tumor. ED12 and ED14: End timepoint. Tumors and chick

embryos' livers were harvested for analysis. ED: embryonic day.

Figure 2: STm promotes tumor cell death in the CAM model. (A-G) Chick embryos

with pre-established A549 tumor cells were directly inoculated with *Salmonella* WT or

PBS on ED 11. Tumors and livers were harvested at 1- and 3-dpi. (A) *Salmonella* WT

CFU recovered per gram of tissue from both tumors and livers at 1 dpi (tumor to liver

ratio 1000:1). (B) Representative images of *Salmonella* WT LPS (green), E-cadherin

(magenta) and Hoechst (blue) staining detected in tumor sections from top, middle and

bottom cut-depth levels of whole tumors harvested at 1 dpi. Scale bar = 100 µm (10x

objective) and 10 µm (100x objective). The middle panel is the zoomed image of the

white box and the right panel is the zoomed image of the red box at 100x objective. (C) A

trypan blue exclusion assay was performed on *ex vivo* dissociated tumor cells suspension,

indicating a loss of tumor cell viability in the group infected with WT *Salmonella* at 3

dpi. (D) Representative images of TUNEL-stained (red) and Hoechst (blue) tumor

sections from 3 dpi (top, middle, bottom). Scale bar = 10 μm, 100x objective. (E)

Quantitative comparison of the percentage of TUNEL-positive nuclei between WT-

infected tumors (52%) and PBS uninfected tumors (33%) at 3 dpi. (F) Representative

images of H and E-stained tumor sections. Zoomed images of the red box of the necrotic

region at 3 dpi. N: necrotic areas. (G) The percentage of the necrotic area as quantified by

an external pathologist from H and E-stained tumor sections infected with WT (22%) and

PBS (4%) at 3 dpi. The results represent the mean ± SD. For all, the statistical

significance was by unpaired t-test, not significant (ns) = P-value * ≤ 0.05, ** ≤ 0.01, ***

≤ 0.001, **** ≤ 0.0001.

Figure 3: Flagellar motility is not required for tumor colonization or intra-tumoral distribution. (A) Representative live *in ovo* images of mCherry-tagged *Salmonella.* WT and *ΔflgK* colonizing A549 tumors on CAMs via either direct (bacteria inoculated directly on tumors) or distant inoculation (bacteria deposited on filter paper, a short distance away from tumors) at 1 dpi. (B-C) Total *Salmonella* WT and *ΔflgK* CFU recovered per gram of tumor tissue at 1 dpi from direct and distant inoculation. The results represent the mean ± SD. The statistical significance was by unpaired t-test, non-significant (ns). (D) Representative images of *Salmonella ΔflgK* LPS (green), E-cadherin (magenta) and Hoechst (blue) staining detected in tumor sections from top, middle and bottom levels of whole tumors harvested at 1 dpi. Scale bar = 100 µm (10x objective) and

10 µm (100x objective). The second column is the zoomed in details of the white box at

100x objective.

Figure 4: SPI-1 and SPI-2 are required for intracellular replication of STm within

3D tumor spheroids. (A) Intracellular bacteria enumeration of STm strains (WT, *ΔhilD*

and ΔssrB) in A549 tumor spheroids measured at invasion (mean log CFU/well at 2 HPI:

WT = 4.2, *ΔhilD* = 2.8, *ΔssrB* = 3.9) and replication (mean log CFU/well at 16 HPI: WT

= 5.6, *ΔhilD* = 3.5, *ΔssrB* = 4.5) stages of infection. Dots: biological replicates, each with

a minimum of five technical replicates. The bars represent the mean +/- standard

deviation (SD). (B) Fold replication (16 HPI/2 HPI) of each STm strain (WT = 28-fold,

ΔhilD = 6-fold, *ΔssrB* = 4-fold) within A549 tumor spheroids. The bars represent the

means of four biological replicates, a minimum of five technical replicates each. Error

bars +/- SD. (C) The amount of LDH released from A549 tumor spheroids after infection

with STm (WT= 53% , *ΔhilD* = 22% *and ΔssrB* = 33%*)* at 16 HPI. The bars represent the

means of three biological replicates, a minimum of five technical replicates each. Error

bars +/- SD. For all, the statistical significance was determined by one way analysis of

variance (ANOVA) with Sidak's multiple comparison test. Not significant (ns) = P-value

*≤ 0.05, **≤ 0.01, ***≤ 0.001, ****≤ 0.0001. (D) Representative images of

immunofluorescence-labeling of 3D tumor spheroids with Ki67 AlexaFluor 488 (green),

Hoeschst (blue) and mCherry-tagged bacterial strains (red) at both 10x and 100x

objectives (edge and core of spheroids). Maximum intensity projections of Z-stack

images were taken with an Olympus Super Resolution Spinning Disk SpinSR-10

microscope. Scale bar = 100 μm (10x) and 10 μm (100x).

Figure 5: In the CAM tumor model, STm resides in an extracellular, non-biofilm

state. (A) Extracellular CFU recovered after infection with WT *Salmonella*, *ΔhilD* and

ΔssrB per gram of tumor at 1 dpi. Statistical significance by one-way ANOVA with

Sidak's multiple comparison test. (B) Intracellular WT *Salmonella*, *ΔhilD* and *ΔssrB*

CFU recovered per gram of tumor at 1dpi. Statistical significance by one-way ANOVA

with Sidak's multiple comparison test. (C) Percentage of total tumor-associated STm

(WT, *ΔhilD, ΔssrB)* recovered before and after ex vivo gentamicin assay from infected

CAM tumors at 1dpi. EX: extracellular, IN: intracellular. Statistical significance was by

one-way ANOVA with Sidak's multiple comparison test. (D) Extracellular *Salmonella*

WT and *ΔcsgD* CFU recovered per gram of tumor at 1- and 3-dpi. The results represent

627   the mean ± SD. The statistical significance was by unpaired t-test. For all, not significant

628   (ns) = P-value * ≤ 0.05, ** ≤ 0.01, *** ≤ 0.001, **** ≤ 0.0001.

Figure 6: Replication of VNP20009 within 3D tumor spheroids is impaired. (A)
Intracellular bacteria enumeration from WT and VNP20009 infected tumor spheroids,
measured at invasion (mean log CFU/well at 2 HPI: WT = 4.2, VNP20009 = 3.6) and
replication (mean log CFU/well at 16 HPI: WT = 5.6, VNP20009 = 4) stages of infection.
Dots: biological replicates, each with a minimum of five technical replicates. The bars
represent the mean +/- standard deviation (SD). The statistical significance was
determined by unpaired t-test. (B) Fold replication (16 HPI/2 HPI) of WT (28-fold) and
VNP20009 (3-fold) strains within tumor spheroids. The bars represent the means of four
biological replicates, a minimum of five technical replicates each. Error bars = +/- SD.
The statistical significance was by unpaired t-test. (C) LDH release from tumor spheroids
infected with WT (53%) and VNP20009 (36%) at 16 HPI. The bars represent the means

of three biological replicates, a minimum of five technical replicates each. Error bars =

+/- SD. Statistical significance was by one-way ANOVA with Sidak's multiple

comparison test. For all, non-significant (ns) = P-value *≤ 0.05, **≤ 0.01, ***≤ 0.001,

****≤ 0.0001. (D) Representative immunofluorescence images of 3D tumor spheroids

with Ki47 AlexaFluor 488 (green), Hoeschst (blue) and mCherry-tagged bacteria strains

(red) at 10x and 100x objectives. Maximum intensity projections of Z-stack images were

taken with an Olympus Super Resolution Spinning Disk SpinSR-10 microscope. (E)

Zoomed 300% magnification of white box ROI from 100x tumor spheroids edge of both

WT and VNP20009 to compare and visualize the cell morphology defect of VNP20009,

outlined by dotted lines.

Figure 7: Reduced tumor colonization by VNP20009 impacts its tumoricidal potential and anti-tumor immunity. (A) CAM tumors infected with VNP20009 (purple line) showed significant improvement in percentage survival and longer survival times compared to WT-infected CAM tumors (red line) at a dose of ~50 CFU. For survival analysis, the Log-rank (Mantel-Cox) test was used. (B) Representative live, *in ovo* images of mCherry-tagged *Salmonella,* WT and VNP20009 colonizing CAM tumors at 1 dpi. (C-D) Extracellular *Salmonella* WT and VNP20009 CFU recovered per gram of tumor and liver at 1- and 3-dpi. The statistical significance was determined by unpaired t-test. (E) Super Resolution 320x magnified representative images of WT compared with distinct cell morphology defects of VNP20009 observed within CAM tumor tissue sections at 3 dpi. LPS (green) and Hoechst (blue). Elongated bacteria cells are outlined

with white and red lines. (F) Intracellular bacterial loads recovered from WT and

VNP20009 CFU per gram of tumor at 1dpi. The statistical significance was determined

by unpaired t-test. (G-H) TNF-α and IL-1β expression level in tumor lysates at 2 HPI,

measured by ELISA, data were expressed as pg per mg of total protein. The results

represent the mean ± SD. Statistical significance was by one-way ANOVA with Sidak's

multiple comparison test. Non-significant (ns) = P-value * ≤ 0.05, ** ≤ 0.01, *** ≤ 0.001,

**** ≤ 0.0001.

Strain or plasmid	Description	Source or reference
Wildtype	*Salmonella enterica* serovar Typhimurium 14028s, pFPV::mCherry	Lab strain collection
Δ*flgK*	*flgK::cm,* pFPV::mCherry	Lab strain collection
Δ*hilD*	*hilD* deletion, pFPV::mCherry	Lab strain collection
Δ*ssrB*	*ssrB::kan,* pFPV::mCherry	Desai et al. (2016)
Δ*csgD*	*csgD::chl,* pFPV::mCherry	Desai et al. (2016)
VNP20009	*Salmonella enterica* subsp. *enterica* strain YS1646, pFPV::mCherry	ATCC BAA3199
pFPV::mCherry plasmid	mCherry cloned between XbaI and SphI sites in pFPV	Addgene plasmid # 20956 http://n2t.net/addgene:20956; RRID:Addgene_20956

699 **Supplementary Table 1: List of bacterial strains and plasmid used in the study.**

710
Supplementary Figure 1: Optimal infective dose of STm for CAM tumor model was

approximately 50 CFU. Dose response survival analysis was performed by infecting the

tumor bearing chick embryos at different infectious doses of STm (~500 CFU, ~50 CFU,

~5 CFU) together with PBS uninfected group. Mortality of chick embryos were

monitored and recorded daily following STm infection. Death of chick embryo was

identified through a small window opening on the top of an eggshell (covered with

transparent Tegaderm) characterized by loss of blood vessels and non-movement of chick

inside the egg.

723
Supplementary Figure 2: VNP20009 was not defective in growth under lab growth

conditions. Similar growth rates were observed among all *Salmonella typhimurium*

strains. STm strains deleted in *flgK, hilD, ssrB, csgD,* VNP20009, and the wild type

14028s were grown for 10 hours in Luria-Bertani broth at 37 °C, 250 rpm and the

absorbance at 600 nm was measured every 2 hours.

Supplementary Figure 3: Representative images of STm WT LPS (green), E-cadherin

(magenta), and Hoechst (blue) staining were detected in tumor sections from top, middle,

and bottom depth levels of whole tumors harvested at 3 dpi. Scale bar = 100 µm (10x

objective) and 10 µm (100x objective). The second panel is the zoomed image of the

white box at 100x objective.

Supplementary Figure 4: Representative images of STm VNP20009 LPS (green), E-

cadherin (magenta), and Hoechst (blue) staining detected in tumor sections from top,

middle, and bottom depth levels of whole tumors harvested at (A) 1 dpi and (B) 3 dpi.

Scale bar = 100 µm (10x objective) and 10 µm (100x objective). The second panel is the

zoomed image of the white box at 100x objective.

References

1. Pawelek JM, Low KB, Bermudes D. 1997. Tumor-targeted Salmonella as a novel

anticancer vector. Cancer Res 57:4537–4544.

2. Leschner S, Weiss S. 2010. Salmonella—allies in the fight against cancer. J Mol

Med 88:763–773.

3. Avogadri F, Martinoli C, Petrovska L, Chiodoni C, Transidico P, Bronte V, Longhi

R, Colombo MP, Dougan G, Rescigno M. 2005. Cancer Immunotherapy Based on

Killing of Salmonella-Infected Tumor Cells. Cancer Research 65:3920–3927.

4. Kaimala S, Al-Sbiei A, Cabral-Marques O, Fernandez-Cabezudo MJ, Al-Ramadi

BK. 2018. Attenuated Bacteria as Immunotherapeutic Tools for Cancer Treatment.

Front Oncol 8:136.

5. Saccheri F, Pozzi C, Avogadri F, Barozzi S, Faretta M, Fusi P, Rescigno M. 2010.

Bacteria-Induced Gap Junctions in Tumors Favor Antigen Cross-Presentation and

Antitumor Immunity. Science Translational Medicine 2:44ra57-44ra57.

6. Zhou S, Gravekamp C, Bermudes D, Liu K. 2018. Tumor-targeting bacteria

engineered to fight cancer. Nat Rev Cancer 18:727–743.

7. Giannella RA. 1996. Salmonella, p. . *In* Baron, S (ed.), Medical Microbiology, 4th

ed. University of Texas Medical Branch at Galveston, Galveston (TX).

8. Sys G, Van Bockstal M, Forsyth R, Balke M, Poffyn B, Uyttendaele D, Bracke M,

De Wever O. 2012. Tumor grafts derived from sarcoma patients retain tumor

morphology, viability, and invasion potential and indicate disease outcomes in the

chick chorioallantoic membrane model. Cancer Letters 326:69–78.

9.   Kain KH, Miller JWI, Jones-Paris CR, Thomason RT, Lewis JD, Bader DM,

Barnett JV, Zijlstra A. 2014. The chick embryo as an expanding experimental model

for cancer and cardiovascular research. Dev Dyn 243:216–228.

10. Mitrevska K, Merlos Rodrigo MA, Cernei N, Michalkova H, Splichal Z, Hynek D,

Zitka O, Heger Z, Kopel P, Adam V, Milosavljevic V. 2023. Chick chorioallantoic

membrane (CAM) assay for the evaluation of the antitumor and antimetastatic

activity of platinum-based drugs in association with the impact on the amino acid

metabolism. Mater Today Bio 19:100570.

11. Miebach L, Berner J, Bekeschus S. 2022. In ovo model in cancer research and

tumor immunology. Front Immunol 13.

12. Ishihara M, Hu J, Wong A, Cano-Ruiz C, Wu L. 2019. Chapter Five - Mouse- and

patient-derived CAM xenografts for studying metastatic renal cell carcinoma, p. 59–

80. *In* Tamanoi, F (ed.), The Enzymes. Academic Press.

13. Fischer D, Fluegen G, Garcia P, Ghaffari-Tabrizi-Wizsy N, Gribaldo L, Huang RY-

J, Rasche V, Ribatti D, Rousset X, Pinto MT, Viallet J, Wang Y, Schneider-Stock

R. 2022. The CAM Model—Q&A with Experts. Cancers (Basel) 15:191.

14. Mon KKZ, Si Z, Chan-Park MB, Kenney LJ. 2022. Polyimidazolium Protects

against an Invasive Clinical Isolate of Salmonella Typhimurium. Antimicrobial

Agents and Chemotherapy 66:e00597-22.

15. Low KB, Ittensohn M, Luo X, Zheng L-M, King I, Pawelek JM, Bermudes D. 2004.

Construction of VNP20009, p. 47–59. *In* Springer, CJ (ed.), Suicide Gene Therapy:

Methods and Reviews. Humana Press, Totowa, NJ.

16. Lee KC, Zheng L-M, Luo X, Clairmont C, Fischer J, Margitich D, Turnier J,

Almassian B, Bermudes D, King I. 2000. Comparative Evaluation of the Acute

Toxic Effects in Monkeys, Pigs and Mice of a Genetically Engineered Salmonella

Strain (VNP20009) Being Developed as an Antitumor Agent. Int J Toxicol 19:19–

25.

17. Clairmont C, Lee KC, Pike J, Ittensohn M, Low KB, Pawelek J, Bermudes D,

Brecher SM, Margitich D, Turnier J, Li Z, Luo X, King I, Zheng LM. 2000.

Biodistribution and Genetic Stability of the Novel Antitumor Agent VNP20009, a

Genetically Modified Strain of Salmonella typhimuvium. The Journal of Infectious

Diseases 181:1996–2002.

18. Low KB, Ittensohn M, Le T, Platt J, Sodi S, Amoss M, Ash O, Carmichael E,

Chakraborty A, Fischer J, Lin SL, Luo X, Miller SI, Zheng L, King I, Pawelek JM,

Bermudes* D. 1999. Lipid A mutant Salmonella with suppressed virulence and

TNFα induction retain tumor-targeting in vivo. 1. Nat Biotechnol 17:37–41.

19. Luo X, Li Z, Lin S, Le T, Ittensohn M, Bermudes D, Runyab JD, Shen S, Chen J,

King IC, Zheng L. 2001. Antitumor Effect of VNP20009, an Attenuated Salmonella

, in Murine Tumor Models. Oncology Research Featuring Preclinical and Clinical

Cancer Therapeutics 12:501–508.

20.   King I, Luo X, Feng M, Ittensohn M, Li Z, Belcourt M, Lin S, Le T, Pike J, Troy K,

Sznol M, Clairmont C, Bermudes D, Zheng L-M. 2000. Tumour therapy using

Salmonella. Emerging Drugs 5:211–219.

21.   Toso JF, Gill VJ, Hwu P, Marincola FM, Restifo NP, Schwartzentruber DJ, Sherry

RM, Topalian SL, Yang JC, Stock F, Freezer LJ, Morton KE, Seipp C, Haworth L,

Mavroukakis S, White D, MacDonald S, Mao J, Sznol M, Rosenberg SA. 2002.

Phase I Study of the Intravenous Administration of Attenuated Salmonella

typhimurium to Patients With Metastatic Melanoma. J Clin Oncol 20:142–152.

22.   Tseng H, Gage JA, Raphael RM, Moore RH, Killian TC, Grande-Allen KJ, Souza

GR. 2013. Assembly of a Three-Dimensional Multitype Bronchiole Coculture

Model Using Magnetic Levitation. Tissue Engineering Part C: Methods 19:665–

675.

23.   Zanoni M, Piccinini F, Arienti C, Zamagni A, Santi S, Polico R, Bevilacqua A,

Tesei A. 2016. 3D tumor spheroid models for in vitro therapeutic screening: a

systematic approach to enhance the biological relevance of data obtained. Sci Rep

6:19103.

24.   Crull K, Bumann D, Weiss S. 2011. Influence of infection route and virulence

factors on colonization of solid tumors by Salmonella enterica serovar

Typhimurium. FEMS Immunology & Medical Microbiology 62:75–83.

25. Wang C-Z, Kazmierczak RA, Eisenstark A. 2016. Strains, Mechanism, and

Perspective: Salmonella-Based Cancer Therapy. International Journal of

Microbiology 2016:5678702.

26. Toley BJ, Forbes NS. 2012. Motility is critical for effective distribution and

accumulation of bacteria in tumor tissue. Integrative Biology 4:165–176.

27. Thornlow DN, Brackett EL, Gigas JM, Van Dessel N, Forbes NS. 2015. Persistent

enhancement of bacterial motility increases tumor penetration. Biotechnology and

Bioengineering 112:2397–2405.

28. Coutermarsh-Ott SL, Broadway KM, Scharf BE, Allen IC. 2017. Effect of

Salmonella enterica serovar Typhimurium VNP20009 and VNP20009 with restored

chemotaxis on 4T1 mouse mammary carcinoma progression. Oncotarget 8:33601–

33613.

29. Homma M, DeRosier DJ, Macnab RM. 1990. Flagellar hook and hook-associated

proteins of *Salmonella typhimurium* and their relationship to other axial components

of the flagellum. Journal of Molecular Biology 213:819–832.

30. Cirillo DM, Valdivia RH, Monack DM, Falkow S. 1998. Macrophage-dependent

induction of the Salmonella pathogenicity island 2 type III secretion system and its

role in intracellular survival. Molecular Microbiology 30:175–188.

31. Main-Hester KL, Colpitts KM, Thomas GA, Fang FC, Libby SJ. 2008. Coordinate

regulation of Salmonella pathogenicity island 1 (SPI1) and SPI4 in Salmonella

enterica serovar Typhimurium. Infect Immun 76:1024–1035.

32.   Haraga A, Ohlson MB, Miller SI. 2008. Salmonellae interplay with host cells. 1. Nat

Rev Microbiol 6:53–66.

33.   Shetty D, Kenney LJ. A pH-sensitive switch activates virulence in Salmonella. eLife

12:e85690.

34.   Crull K, Rohde M, Westphal K, Loessner H, Wolf K, Felipe-López A, Hensel M,

Weiss S. 2011. Biofilm formation by Salmonella enterica serovar Typhimurium

colonizing solid tumours. Cellular Microbiology 13:1223–1233.

35.   Desai SK, Winardhi RS, Periasamy S, Dykas MM, Jie Y, Kenney LJ. 2016. The

horizontally-acquired response regulator SsrB drives a Salmonella lifestyle switch

by relieving biofilm silencing. eLife 5:e10747.

36.   Low KB, Ittensohn M, Luo X, Zheng L-M, King I, Pawelek JM, Bermudes D. 2004.

Construction of VNP20009: a novel, genetically stable antibiotic-sensitive strain of

tumor-targeting Salmonella for parenteral administration in humans. Methods Mol

Med 90:47–60.

37.   Garcia P, Wang Y, Viallet J, Macek Jilkova Z. 2021. The Chicken Embryo Model:

A Novel and Relevant Model for Immune-Based Studies. Frontiers in Immunology

12.

38.   Kim J-E, Phan TX, Nguyen VH, Dinh-Vu H-V, Zheng JH, Yun M, Park S-G, Hong

Y, Choy HE, Szardenings M, Hwang W, Park J-A, Park S, Im S-H, Min J-J. 2015.

Salmonella typhimurium Suppresses Tumor Growth via the Pro-Inflammatory

Cytokine Interleukin-1β. Theranostics 5:1328–1342.

39. Leschner S, Westphal K, Dietrich N, Viegas N, Jablonska J, Lyszkiewicz M,

Lienenklaus S, Falk W, Gekara N, Loessner H, Weiss S. 2009. Tumor Invasion of

Salmonella enterica Serovar Typhimurium Is Accompanied by Strong Hemorrhage

Promoted by TNF-α. PLOS ONE 4:e6692.

40. Avogadri F, Martinoli C, Petrovska L, Chiodoni C, Transidico P, Bronte V, Longhi

R, Colombo MP, Dougan G, Rescigno M. 2005. Cancer Immunotherapy Based on

Killing of Salmonella-Infected Tumor Cells. Cancer Research 65:3920–3927.

41. Pawelek JM, Sodi S, Chakraborty AK, Platt JT, Miller S, Holden DW, Hensel M,

Low KB. 2002. Salmonella pathogenicity island-2 and anticancer activity in mice.

10. Cancer Gene Ther 9:813–818.

42. Loessner H, Endmann A, Leschner S, Westphal K, Rohde M, Miloud T,

Hämmerling G, Neuhaus K, Weiss S. 2007. Remote control of tumour-targeted

Salmonella enterica serovar Typhimurium by the use of l-arabinose as inducer of

bacterial gene expression in vivo. Cellular Microbiology 9:1529–1537.

43. Silva-Valenzuela CA, Desai PT, Molina-Quiroz RC, Pezoa D, Zhang Y, Porwollik

S, Zhao M, Hoffman RM, Contreras I, Santiviago CA, McClelland M. 2016. Solid

tumors provide niche-specific conditions that lead to preferential growth of

Salmonella. Oncotarget 7:35169.

44. Valdes TI, Kreutzer D, Moussy F. 2002. The chick chorioallantoic membrane as a

novel in vivo model for the testing of biomaterials. J Biomed Mater Res 62:273–

282.

45.   Kogut M, Lowry V, Moyes R, Bowden L, Bowden R, Genovese K, Deloach. 1998.

Lymphokine-augmented activation of avian heterophils. Poultry Science 77:964–

971.

46.   Kogut MH, Holtzapple C, Lowry VK, Genovese K, Stanker LH. 1998. Functional

responses of neonatal chicken and turkey heterophils following stimulation by

inflammatory agonists. Am J Vet Res 59:1404–1408.

47.   Li M, Lu M, Lai Y, Zhang X, Li Y, Mao P, Liang Z, Mu Y, Lin Y, Zhao AZ, Zhao

Z, Zhou S, Li F. 2020. Inhibition of acute leukemia with attenuated Salmonella

typhimurium strain VNP20009. Biomedicine & Pharmacotherapy 129:110425.

48.   Liu X, Guo Y, Sun Y, Chen Y, Tan W, Min J-J, Zheng JH. 2022. Comparison of

Anticancer Activities and Biosafety Between Salmonella enterica Serovar

Typhimurium ΔppGpp and VNP20009 in a Murine Cancer Model. Front Microbiol

13.

49.   Ahmed SG, Oliva G, Shao M, Wang X, Mekalanos JJ, Brenner GJ. 2022.

Intratumoral injection of schwannoma with attenuated Salmonella typhimurium

induces antitumor immunity and controls tumor growth. Proceedings of the National

Academy of Sciences 119:e2202719119.

50.   Karsten V, Murray SR, Pike J, Troy K, Ittensohn M, Kondradzhyan M, Low KB,

Bermudes D. 2009. msbB deletion confers acute sensitivity to CO2 in Salmonella

enterica serovar Typhimurium that can be suppressed by a loss-of-function mutation

in zwf. BMC Microbiol 9:170.

51.   Murray SR, Bermudes D, de Felipe KS, Low KB. 2001. Extragenic Suppressors of

914   Growth Defects in msbB Salmonella. J Bacteriol 183:5554–5561.

915   52.   Vander Heiden MG, Cantley LC, Thompson CB. 2009. Understanding the Warburg

916   Effect: The Metabolic Requirements of Cell Proliferation. Science 324:1029–1033.

917   53.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,

918   Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein

919   V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for

920   biological-image analysis. Nat Methods 9:676–682.