Spatial Expression of Long Non-Coding RNAs in Human Brains of Alzheimer's Disease

Bernard Ng[1, #], Denis R. Avey[1, 5, #], Katia de Paiva Lopes[1], Masashi Fujita[2], Ricardo A. Vialle[1], Himanshu Vyas[1], Nicola A. Kearns[1, 5], Shinya Tasaki[1], Artemis Iatrou[1,3], Sashini De Tissera[1], Tien-Hao Chang[1], Jishu Xu[1], Chunjiang Yu[1], Faraz Sultan[1], Vilas Menon[2], Chris Gaiteri[1,4], Philip L. De Jager[2], David A. Bennett[1, 5], Yanling Wang[1, 5, *]

1. Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL, 60612, USA
2. Center for Translational and Computational Neuroimmunology, Department of Neurology, Columbia University Irving Medical Center, New York, NY, USA
3. Current affiliation: Department of Psychiatry, McLean Hospital, Harvard Medical School, Belmont, MA, 02478, USA
4. Department of Psychiatry, Upstate Medical University, Syracuse, NY, 13210, USA
5. Department of Neurological Sciences, Rush University Medical Center, Chicago, IL, 60612, USA

#Contributed equally
*Correspondence to yanling_wang@rush.edu

Abstract

Long non-coding RNAs (lncRNAs) are critical regulators of physiological and pathological processes, with their dysregulation increasingly implicated in aging and Alzheimer's disease (AD). Using spatial transcriptomics, we analyzed 78 postmortem brain sections from 21 ROSMAP participants to map the spatial expression of lncRNAs in the dorsolateral prefrontal cortex of aged human brains. Compared to mRNAs, lncRNAs exhibited greater subregion-specific expression, with enrichment in antisense and lincRNA biotypes. Network analysis identified 193 gene modules across eight subregions, including lncRNA-enriched modules involved in critical biological processes. We also identified AD differentially expressed (DE) lncRNAs, which showed greater subregion specificity than AD DE mRNAs. Gene set enrichment analysis highlighted the involvement of these AD DE lncRNAs in epigenetic regulation and chromatin remodeling, including enrichment for HDAC target genes such as *OIP5-AS1*. Statistical modeling suggested that interactions between *OIP5-AS1* and HDAC proteins, particularly HDAC11, were associated with tau tangles in excitatory neurons and plaque burden in microglia. This study provides a comprehensive resource of lncRNA spatial expression in the aged human brain and uncovers potential functional roles of lncRNAs in AD pathogenesis.

Introduction

One of the most remarkable discoveries of the 21st century is that a significant portion of the genome is transcribed into RNAs with little or no protein-coding potential[1,2]. While approximately 90% of the human genome is transcribed, only about 2% is annotated as protein-coding genes. This paradigm-shifting discovery has spurred extensive research into the regulatory roles of non-coding RNAs (ncRNAs) in gene expression[2,3], redefining our understanding of the genome and challenging the traditional view that only protein-coding genes are functionally significant.

Among ncRNAs, long non-coding RNAs (lncRNAs) represent a prominent class of transcripts longer than 200 nucleotides, predominantly transcribed by RNA Polymerase II. LncRNAs regulate gene expression through diverse mechanisms, including epigenetic modification, chromatin remodeling, and post-transcriptional processes[4,5]. Compared to coding RNAs, lncRNAs exhibit greater tissue and cell-type specificity[6,7] and are particularly abundant in the human brain, where they play critical roles in neuronal differentiation, synaptogenesis, and neuroplasticity[1,8]. Dysregulated lncRNA expression has been implicated in numerous neurological disorders, including Alzheimer's disease (AD). In AD, aberrant lncRNA expression has been linked to key pathological processes, such as amyloid-beta metabolism, tau pathology, neuroinflammation, and neuronal death[9–13]. However, the spatial and cellular contexts in which lncRNAs might influence AD-related processes remain largely unexplored.

In this study, we present the spatial expression profiles of lncRNAs in the dorsolateral prefrontal cortex (DLPFC) of 21 participants from Religious Orders Study and Memory and Aging Project (ROSMAP). We identify subregion-specific lncRNAs and examine their overlap with cell-type-specific lncRNAs. Additionally, we map lncRNAs and mRNAs that are differentially expressed (DE) in AD across distinct subregions, comparing our spatial transcriptomics (ST) results with findings from bulk RNA-seq and single-nucleus RNA-seq (snRNA-seq) analyses. Finally, we infer potential interactions between *OIP5-AS1* and histone deacetylase (HDAC) proteins, particularly HDAC11, providing new insights into the functional roles of lncRNAs in AD pathogenesis.

Results

Spatial expression of lncRNAs in aged human brains

Using our recently acquired ST data[14], we analyzed lncRNA expression from 78 DLPFC sections of 21 female individuals (AD = 13; no cognitive impairment (NCI) = 8). In total, we detected 7,634 lncRNAs and 16,769 protein-coding mRNAs across 8 spatial spot clusters corresponding to subregions: layer 1 (L1), layers 2–3 (L2–3), layers 3–5 (L3–5), layer 5 (L5), layer 6 (L6), white matter (WM), meninges, and

blood vessels (Fig. 1A). Consistent with prior studies[6,7], lncRNAs exhibited lower expression levels than mRNAs (Fig. 1B), with most lncRNAs detected in fewer than 0.01% of ST spots (Fig. 1C). For downstream analysis, we retained 1,578 lncRNAs and 13,808 mRNAs with adequate expression (counts per million [CPM] >1 in >80% of brain sections). Among the detected lncRNAs, the most prevalent biotypes were lincRNAs and antisense transcripts (Fig. 1D). Notably, lncRNAs demonstrated greater subregion specificity, with only 39.9% of lncRNAs detected across all eight subregions, compared to 89.3% of mRNAs (Fig. 1E).

By contrasting the expression of spatial spots in one subregion against all others, we identified 79 subregion-specific lncRNAs (Fig. 2A–2C; Table S1). To investigate whether subregion-specific lncRNAs also exhibit cell-type specificity, we cross-referenced a published snRNA-seq dataset derived from the DLPFC of 424 ROSMAP participants[15]. While we observed limited overlap between subregion-specific and cell-type-specific lncRNAs (Fig. 2D; Table S1), some exceptions emerged. For example, *AL031056.1* was enriched in L1 (Fig. 2A) and astrocytes (Fig. 2D), *LINC00507* in L2–3 (Fig. 2A) and excitatory neurons (Fig. 2D), *AC021613.1* in L6 (Fig. 2A, 2C) and excitatory neurons (Fig. 2D), and *AC092958.1*, *TYMSOS*, and *LINC00844* in WM (Fig. 2A, 2B) and oligodendrocytes (Fig. 2C, 2D). However, the majority of subregion-specific lncRNAs were not confined to specific cell types (Fig. 2D).

We next examined whether cell-type-specific lncRNAs exhibited subregion specificity. Although general trends emerged, such as astrocyte-specific lncRNAs being more abundant in L1, excitatory/inhibitory neuron-specific lncRNAs in cortical layers, and oligodendrocyte-specific lncRNAs in WM, most cell-type-specific lncRNAs were expressed across multiple subregions (Fig. 2E; Table S1). These observations likely reflect the heterogeneity of cell types within each subregion and the shared developmental origins of cell types that span cortical layers. Nonetheless, the identification of both subregion-specific and cell-type-specific lncRNAs suggests that some lncRNAs function within precise spatial regions to support localized processes, while others serve the specific needs of individual cell types.

lncRNA-enriched gene modules in aged human brains

To investigate the molecular systems within each subregion, we derived a pseudo-bulk matrix for each subregion and performed co-expression network analysis using the Speakeasy algorithm[16] (Fig. 3A). Both lncRNAs and mRNAs were included to capture their potential interactions across biological processes. This analysis generated 193 modules, each containing at least 30 genes (Table S2). Module preservation analysis revealed that cortical layer modules were more preserved than those derived from WM, blood vessels, and meninges (Fig. 3B; Table S3). Within individual subregions, 15–60% of

the modules were lncRNA-enriched (Fig. 3C). For example, the L2-3_m14 module, comprising 1,261

genes, was enriched with 210 lncRNAs (Bonferroni p < 0.05), including 4 reported AD-differential

lncRNAs[9,10,17], 3 AD-GWAS genes[18], and 23 AD-differential lncRNA genes based on our ST analysis

(Fig. 3D; Tables S2, S6). Gene Ontology (GO) analysis revealed that L2-3_m14 was enriched for

biological processes, such as metabolism, cell cycle regulation, ncRNA processing, and positive

regulation of telomerase (Fig. 3E, Table S4). Similarly, the WM_m11 module, containing 1,148 genes,

was enriched with 156 lncRNAs (Bonferroni p < 0.05), including 2 previously reported AD-differential

lncRNAs[9,10,17], 4 AD-GWAS genes[18], and 29 AD-differential lncRNAs based on our ST analysis (Fig.

3F; Tables S2, S6). GO analysis indicated that WM_m11 was associated with processes such as

carbohydrate derivative metabolism, lipid metabolic modification, and nervous system development

(Fig. 3G, Table S4). In addition, we conducted module-trait association analysis and detected a small

fraction of modules associated with AD traits (nominal p < 0.05), including lncRNA-enriched modules

(Table S5).

AD DE lncRNAs and cross-modality comparisons

To identify DE lncRNAs in AD compared to NCI, we aggregated spot-level expression values within

each brain section to generate a pseudo-bulk matrix for each subregion. These estimates were

normalized using TMM-voom[19], and linear mixed models (LMM) were applied to contrast AD against

NCI while adjusting for age, RNA Integrity Number (RIN), postmortem interval (PMI). Batch effects were

included as fixed effects, and within-subject correlations were modeled as random effects. After

applying false discovery rate (FDR) correction across genes and subregions, we identified 275 AD DE

lncRNAs (Table S6). Notably, a greater proportion of these lncRNAs exhibited negative associations

with AD compared to mRNAs (Fig. 4A, 4B). Biotype analysis revealed enrichment of bidirectional

promoter lncRNAs in L3–5 and lincRNAs in L6 and WM (Fig. 4C).

Next, we compared our ST-derived AD DE lncRNAs and mRNAs with those identified from bulk RNA-

seq data of the DLPFC in 1,049 individuals[20] (Table S6). Although the overlap of AD DE genes between

ST and bulk RNA-seq was modest (Fig. 4D), t-values for all lncRNAs and mRNAs exhibited moderate

correlations between the two modalities (R = 0.3 for lncRNAs; R = 0.39 for mRNAs; Fig. 4E).

Concordant findings included several downregulated lncRNAs, such as *LY86-AS1*, *FAM106A*,

LINC01007, and *MIR7-3HG*, as well as upregulated lncRNAs, including *NEAT1*, *LINC00632*, *FGD5-*

AS1, and *LINC01094* in AD brains (Fig. 4E).

Consistent with our ST findings, snRNA-seq analysis also showed that DE lncRNAs were more

frequently negatively associated with AD than DE mRNAs (Fig. S1A– S1B; Table S6). Additionally, AD

DE lncRNAs were enriched for lincRNAs, particularly in inhibitory neurons and oligodendrocyte
precursor cells (OPCs) (Fig. S1C). These DE lncRNAs also displayed greater cell-type specificity in
oligodendrocytes, inhibitory neurons, and excitatory neurons compared to DE mRNAs (Fig. S1D).
However, overlaps between subregion-specific and cell-type-specific AD DE lncRNAs were minimal
(Fig. S1E).

Pairwise t-value correlations revealed moderate alignment between L3–5 and inhibitory neurons, as
well as between blood vessels and microglia/oligodendrocytes (Fig. S1F). While a larger proportion of
DE mRNAs were shared across subregions and cell types (Fig. S1E), t-value correlations between
subregions and cell types remained weak (Fig. S1G), likely reflecting differences in resolution and
sensitivity between ST and snRNA-seq platforms.

Spatial expression of AD DE lncRNAs and their functional implications

When comparing AD DE lncRNAs and mRNAs, we found that DE lncRNAs were generally more
subregion-specific than DE mRNAs, with 115 of 275 DE lncRNAs showing subregion-specific
differential expression in the cortex (Fig. 5A; Table S6). A higher proportion of DE lncRNAs were
specifically expressed in the meninges, L2–3, L3–5, L5, and L6 (Fig. 5A). Notably, these subregion-
specific DE lncRNAs, regardless of whether they were up- or downregulated, did not appear to be cell-
type-specific (Fig. 5B). The remaining 160 of 275 AD DE lncRNAs were differentially expressed across
multiple subregions, with the majority (~93%) being novel lncRNAs (Fig. 5C). Among the 11 DE
lncRNAs that were previously reported[9,10,17], 8 were also-replicated in bulk RNA-seq datasets (Fig. 5C,
bottom row; Table S6).

Some AD DE lncRNAs displayed high expression in one subregion but showed predominant AD effects
in other subregions. For example, *LINC00844* was highly expressed in WM/oligodendrocytes (Fig. 2A,
2B) but exhibited its strongest AD effects in L2-3 (Fig. 5B). Similarly, *LINC00507* showed high
expression in L2-3 but demonstrated predominant AD effects in L3-6 (Fig. 5C). On the other hand, the
X-inactive-specific transcript (*XIST*), which was broadly expressed across all subregions (Fig. 2A, Fig.
5D), showed its AD effects primarily in L1 and L2–3 (Fig. 5C, 5E). The detection of DE *XIST* prompted
an investigation into other X-linked genes, revealing that *MECP2* was upregulated in L5, while *PGK1*
was downregulated in L1, L2–3, and blood vessels in AD brains (Fig. 5E).

To further explore functional implications, we applied gene set enrichment analysis (GSEA) on t-values
for all genes across subregions, focusing on gene sets containing AD DE lncRNAs. This analysis
revealed enrichment for processes such as cell motility (*MALAT1*, *SPATA13*), apoptosis (*ARMCX5-*
GPRASP2), chromatin organization (*XIST*, *KCNQ1OT1*), and regulation of defense responses

(*NEAT1*) (Fig. 5F; Table S7–8). Notably, we also observed enrichment for HDAC targets, which

included three AD DE lncRNAs (*FGD5-AS1*, *NNT-AS1*, *OIP5-AS1*) (Fig. 5F; Table S8). Consistent with

this finding, several HDACs (*HDAC1*, *HDAC3*, *HDAC6*, *HDAC7*, and *HDAC10*) were differentially

expressed in one or more subregions of AD brains (Fig. 5G). These results support growing evidence

implicating HDACs in AD pathogenesis[21,22] and suggest a potential functional link between specific

lncRNAs and HDAC activity.

Inferred interaction between *OIP5-AS1* and HDAC11 is associated with AD traits

HDAC inhibition has been extensively studied as a potential therapeutic approach for AD[21,23–25].

However, the lack of selectivity in current HDAC inhibitors often results in unintended off-target

effects[26,27]. Our GSEA identified specific lncRNAs, including *FGD5-AS1*, *NNT-AS1*, and *OIP5-AS1*, that

may interact with HDACs, suggesting a potential strategy to selectively modulate HDAC activity in AD

treatment. Among these lncRNAs, *OIP5-AS1* (Cyrano) has been implicated in a wide range of biological

processes, including development[28–30], cancer progression[31–34], and cerebral ischemia-reperfusion

injury[35].

To validate *OIP5-AS1* expression, we performed RNAscope and immunohistochemistry (IHC) on

postmortem DLPFC sections from ROSMAP participants (n=8). Consistent with our ST findings, *OIP5-*

AS1 was broadly expressed across cortical layers, particularly in L2-3 (Fig. S2A, S2B). At the cellular

level, *OIP5-AS1* expression was detected in MAP2+ neurons and IBA1+ microglia (Fig. S2A, S2C).

These observations prompted further investigation into whether genetic variation influences *OIP5-AS1*

expression and whether its interactions with HDACs are associated with clinical and pathological AD

traits. Using the published snRNA-seq dataset[15], we first curated expression quantitative trait loci

(eQTL) statistics of SNPs within 1 Mb of *OIP5-AS1's* transcription start site (TSS) across six major brain

cell types. Multivariate fine-mapping[36] identified rs1655558 as a potential causal variant influencing

OIP5-AS1 expression in multiple cell types, including excitatory neurons, astrocytes, and

oligodendrocytes (Fig. 6A). We then examined the interaction between *OIP5-AS1* expression in six cell

types and HDAC protein abundance in the DLPFC using a published proteomics dataset [37]. These

interactions were tested for associations with AD-related traits: cognitive decline, neuritic plaques, and

tau tangles (Fig. 6B). Significant interaction effects were declared at an α of 0.05, corrected for the

number of cell types and HDACs tested. Our analysis revealed that interactions between *OIP5-AS1*

and HDACs were more strongly associated with tau tangles and neuritic plaques than with cognitive

decline (Fig. 6C, left). Notably, some interaction effects exhibited p-values an order of magnitude lower

than the main effects of HDACs alone (circled dots in Fig. 6D, 6G), suggesting that *OIP5-AS1* plays a

critical modulatory role in HDAC-mediated AD pathology. Comparable trends were observed when using HDAC mRNA expression instead of protein abundance, although interaction effects were weaker (Fig. 6C, right; Fig. S3), indicating that *OIP5-AS1* primarily interacts with HDAC proteins rather than their transcripts in influencing AD pathophysiology.

The interaction between *OIP5-AS1* and HDAC11 was particularly striking. In excitatory neurons, this interaction was associated with tau pathology (Fig. 6D), with the strongest effects observed in the hippocampus and medial temporal lobe (Fig. 6E). In microglia, the interaction was associated with neuritic plaques (Fig. 6G), with similarly pronounced effects in the same brain regions (Fig. 6H). Stratifying 424 ROSMAP participants[15] into tertiles based on *OIP5-AS1* expression levels in excitatory neurons or microglia revealed that HDAC11 protein abundance was positively associated with tau tangles only when *OIP5-AS1* expression was high in excitatory neurons (Fig. 6F). Conversely, HDAC11 protein abundance was positively associated with plaques only when *OIP5-AS1* expression was low in microglia (Fig. 6I). These findings suggest that *OIP5-AS1* interacts with HDACs, particularly HDAC11, in a cell-type-specific manner, modulating their effects on AD pathology.

Discussion

Recent advancements in sequencing technologies have revealed that a significant portion of the mammalian genome is transcribed into lncRNAs[1]. While the brain expresses more tissue-specific lncRNAs than most other tissues[38,39], their spatial and cellular contexts in the human brain remain poorly understood. In this study, we mapped the spatial expression of lncRNAs in DLPFC of aged human brains and found that lncRNAs exhibit lower overall expression levels and more restricted spatial patterns compared to mRNAs (Fig. 1). Using a published snRNA-seq dataset[15], we compared subregion- and cell-type-specific lncRNAs and observed limited overlap, with some alignment between white matter and oligodendrocytes (Fig. 2). Additionally, we identified numerous subregion-specific lncRNA-enriched gene modules associated with diverse biological processes (Fig. 3), suggesting that lncRNAs may influence brain function by interacting with mRNAs within shared molecular systems.

Out of 275 DE lncRNAs, 11 DE lncRNAs were previously reported in AD studies (Fig. 5C). For example, we observed higher expression of *MALAT1*, *NEAT1*, *BACE1-AS*, and *BDNF-AS* in AD individuals, consistent with their known roles in AD pathogenesis. *BACE1-AS* promotes amyloid production[40–42], while *BDNF-AS* modulates synaptic plasticity through interactions with its mRNA counterpart[43–45]. *MALAT1*, a major organizer of nuclear speckles, suppresses neuronal apoptosis and inflammation by modulating *miR-125b*[46,47]. Similarly, *NEAT1* regulates tau phosphorylation and Aβ metabolism through interactions with *miR-124*, *miR-107*, and histone acetyltransferases[48,49]. In addition, we identified

emerging immune- and aging-associated lncRNAs, such as *LY86-AS1*[50], *FAM106A*[51], and *MIR7-3HG*[52].

Increased *XIST* levels have been previously reported in the brains of female AD patients and linked to X Chromosome Interaction (XCI), a process contributing to female-specific vulnerability in AD[53,54]. We detected an upregulation of *XIST* in AD, primarily in the upper cortical layers (Fig. 5C-5E). Since our study included only females, this upregulation suggests disrupted XCI in AD, potentially leading to silencing of some X-linked genes and/or overexpression of escapee genes. Consistently, we observed altered expression of X-linked genes, such as *MECP2*, *PGK1*, *MAOA*, and *HPRT1* (Fig. 5D, 5E; Table S6), which are critical for neuronal function, epigenetic regulation, and cellular metabolism[55–59].

Gene set enrichment analysis revealed that AD DE lncRNAs are involved in diverse regulatory processes, including epigenetic regulation, chromatin remodeling, and RNA metabolism. One key finding was the enrichment of HDAC target genes, including *FGD5-AS1*, *NNT-AS1*, and *OIP5-AS1*. Among these, *OIP5-AS1* stands out for its diverse regulatory roles[60]. *OIP5-AS1* has been shown to regulate cell proliferation, apoptosis, migration, and invasion in cancer through miRNA sponging and interactions with chromatin modifiers, such as HDACs[31–34,61]. In the brain, *OIP5-AS1* is upregulated in M2 microglia-derived exosomes, where it inhibits neuronal pyroptosis via ubiquitination of its binding protein TXNIP[35]. Additionally, it acts as a central player in a non-coding RNA network by degrading mature *miR-7*, thereby modulating neuronal activity[62].

Using statistical modeling, we inferred that the interactions between *OIP5-AS1* and HDAC proteins, particularly HDAC11, are associated with tau tangles in excitatory neurons and plaque burden in microglia (Fig. 6D-I). These findings indicate that *OIP5-AS1* may contribute to AD pathology through epigenetic regulation in a cell-type-specific and context-dependent manner. Notably, the interaction between *OIP5-AS1* and HDAC11 in excitatory neurons was strongly associated with tau tangle burden in the hippocampus and medial temporal lobe (Fig. 6D, 6E). Given that tau pathology typically originates in these regions and progresses to other cortical areas, our findings suggest that the interaction between *OIP5-AS1* and HDAC11 may play a central role in tau progression and early hippocampal vulnerability.

Limitations

This study has several inherent limitations. First, our spatial transcriptomics (ST) samples were collected exclusively from non-Latino white females, limiting the generalizability of our findings to younger individuals, males, or other racial and ancestral groups. Second, the 10x Visium platform relies on polyadenylated (poly(A)) RNA capture, which excludes non-polyadenylated lncRNAs as well as

those with alternative 3' end processing or shorter poly(A) tails. Third, the platform's resolution of 55 µm constrains our ability to resolve cell-type-specific lncRNA expression within the spatial context. Fourth, the relatively small sample size of the study reduces statistical power, likely leading to an underestimation of the total number of AD-associated DE lncRNAs. Finally, while we provide insights into potential functional roles of lncRNAs, including *OIP5-AS1*, experimental validation is necessary to confirm their mechanistic contributions to AD pathogenesis.

Conclusion and Future Directions

This study provides ST profiles of lncRNAs in DLPFC of aged human brains. We mapped subregion-specific lncRNA changes and identified hundreds of AD-associated DE lncRNAs, encompassing previously known AD-associated lncRNAs, emerging immune-related lncRNAs, and novel lncRNAs. These findings lay the groundwork for future investigations using higher-resolution spatial profiling platforms, advanced lncRNA enrichment techniques, and co-detection of proteins in the same tissue sections, enabling a more precise delineation of cell-type-specific lncRNA changes in response to distinct AD pathologies. Additionally, we explored the potential interaction between *OIP5-AS1* and HDACs in mediating AD-associated phenotypes. Emerging RNA-targeting technologies, such as antisense oligonucleotides (ASOs)[63] and CRISPR-Cas13d[64,65], present exciting opportunities for functional studies on *OIP5-AS1* and other AD-associated lncRNAs. Notably, a key limitation of HDAC inhibitors in clinical trials has been their lack of specificity. *OIP5-AS1* may offer a promising avenue for selectively modulating HDAC activity and stratifying patients for precision medicine.

Acknowledgments

We are grateful to those who agreed to donate their brains for research. We thank all the employees at RADC for their support and assistance. This study was supported by NIA grants R01AG074082 and R01AG079223 (to Y.W.), P30AG10161, P30AG72975, R01AG015819, R01AG017917, and U01AG61356 (to D. A. B.), and R01AG061798 (to C.G.).

Author contributions

B.N., D.R.A., K.P.L., S.T., R.A.V., T.C., and A.I. performed data analysis. D.R.A. and S.D.T. conducted ST experiments. J.X. performed ST data processing. M.F., V.M., and P.L.D. provided snRNA-seq data. H.V. and N.A.K. conducted RNAScope and IHC experiments. Y.W., B.N., D.R.A., and K.P.L. wrote the manuscript. D.A.B. runs the parent ROSMAP study and provides critical input. Y.W. conceived and directed this study, overseeing research activities and leading manuscript preparation. All authors reviewed and approved the final manuscript.

Lead Contact

Further information and requests may be directed to the corresponding author Yanling Wang
(Yanling_Wang@rush.edu)

Materials Availability

This study did not generate new unique reagents.

Data Availability

Subregion- and cell-type-enriched lncRNAs, ST gene expression modules and AD-differential genes are
provided in the Supplementary Information. Raw and normalized count matrices of the ST data is available at
Synapse under accession code syn53141470.

Online Methods

Tissue collection for Spatial Transcriptomics

The human brain tissues were obtained through the ROSMAP (Religious Orders Study and Rush
Memory and Aging Project) cohort studies at the Rush Alzheimer's Disease Center, Rush University
Medical Center (RUMC), Chicago, IL[66]. Both studies were approved by the Institutional Review Board
of RUMC. Written informed consent was obtained as was a repository consent and an Anatomic Gift
Act, in compliance with national ethical guidelines. We selected 13 female individuals clinically
diagnosed as AD and 8 controls with no cognitive impairment (NCI) based on criteria as previously
reported[67–69]. The brain tissue of those individuals had RIN> 5 and minimal freezing artifacts.   Frozen
brain blocks of the right-hemisphere DLPFC were cut (~ 1 cm^3), transferred on dry ice, and stored at -
80C.

Frozen brain blocks were embedded in OCT and cryosectioned coronally to a thickness of 10 µm
using a Leica CM1950 cryostat set at -17C. Sectioning was done in sets of three (middle section for
ST and two adjacent sections for IF), with 10-50 µm between each set. Sections were attached to
Visium slides and stored at -80C in a slide mailer for up to one week before proceeding with spatial
transcriptomics. ST-adjacent sections were attached to pre-labeled Leica superfrost plus slides and
stored at -80C in slide mailers for up to two weeks before proceeding with immunohistochemistry.

Spatial Transcriptomics

Spatial Transcriptomics was performed according to the Visium Spatial Gene Expression User Guide
(10X Genomics; CG000239). Briefly, cryosectioned tissue on Visium slides was transferred from -80C

to the lab on dry ice and quickly thawed for 1 min on a heat block at 37C, then immediately

transferred to methanol pre-equilibrated to -20C and fixed at -20C for 30 min. Sections were then

stained with hematoxylin and eosin, and brightfield images were acquired with a Nikon Ti2 with NIS-

Elements AR 5.11.01 64-bit software. After imaging, tissues were immediately permeabilized in 0.1%

pepsin diluted in 0.1 M HCl for 18 min at 37C. Reverse transcription, second-strand synthesis, cDNA

amplification (15 cycles), and library construction were performed according to the manufacturer's

instructions. For the first four individuals (#2, 8, 12, and 18), protocol version 'Rev A' was followed,

and the sample index PCR was 12 cycles. For the remaining samples, protocol version 'Rev D' was

followed, and the sample index PCR was 13-14 cycles. Library concentration was quantified by Qubit

1x dsDNA HS using a SpectraMax M3 plate reader. Library concentration and size distribution were

assessed by Fragment Analyzer (Agilent). Sequencing was performed in three batches on an Illumina

NovaSeq, targeting a minimum of 50,000 raw reads per spot.

Spatial transcriptomics sequencing data processing

Sequencing data were pre-processed with the Space Ranger pipeline (10x Genomics), which

mapped the barcodes against the human genome (Gencode 27), generated FASTQ files, and

generated a count matrix for each section. The individual files were then merged into a Seurat object

(Seurat v4.2.0) and were processed in R. We calculated per-spot quality metrics and spots with low-

measured genes (< 500) were removed. The Seurat::SCTransform function was used to normalize

and scale the UMI counts based on regularized negative binomial regression. Principal component

analysis (PCA) was performed on the top 3000 variable genes. Using the top 50 PCs and accounting

for technical and biological covariates such as age, RIN, library batch, and donor, we integrated the

datasets using the Harmony package (v0.1.0). We explored various tools for data integration and

chose Harmony for its high performance, usability, and scalability. We only used Harmony integration

to identify clusters corresponding to brain regions. All downstream analyses (differential gene

expression, cell proportion estimates, and ligand-receptor analysis) were performed on pre-

integration count data. After visualizing the standard deviation attributed to each Harmony embedding

as an elbow plot, we decided to use the top 10 Harmony embeddings. We applied Seurat::runUMAP

to compute UMAP (uniform manifold approximation and projection) dimension reduction and

Seurat::findNeighbors and Seurat::findClusters to identify clusters based on a shared nearest

neighbor (SNN) clustering algorithm (Louvain clustering with resolution set to 0.3; all other

parameters default). This resulted in 10 clusters, which were annotated based on their canonical

markers and spatialLIBD (v7.5) to the closest anatomical cortical layer (L1, L2-3, L3-5, L5, L6), white

matter (WM), meninges, and specific cellular subpopulations, namely cells with high expression of

hemoglobin genes ('Blood Vessel') and SST/NPY ('Interneuron'). The last identified cluster, which

carried a glial signature, had very few spots and so was not included in downstream analyses.

Seurat::FindMarkers was used to identify cluster-enriched (e.g. subregion-specific) lncRNAs reported

in Fig. 2A and Table S1. For Fig. 2A, a log2FC cutoff of 0.1 was employed resulting in 50 lncRNAs.

Co-expression network analysis in spatial transcriptomics

The modules of co-expressed genes were derived using the Speakeasy (SE) algorithm v1, Matlab

version[16]. We created pseudo-bulk gene expression by summing the counts per donor, then splitting

by UMAP cluster. Lowly expressed genes (less than 1 count per million, CPM) were removed, and

the variancePartition R function was applied to capture the major sources of variation, as previously

described[14]. The input for SE consisted of the residuals after normalization using tmm.voom and

adjustment for library batch, RIN, age, and PMI. We built the modules separately for each UMAP

cluster, using the results from 100 initializations of SE and Spearman correlation. This resulted in 193

modules, each with at least 30 genes (L1 = 30, L2-3 = 21, L3-5 = 20, L5 = 20, L6 = 20, Meninges =

33, WM = 20, and Blood Vessels = 29). The module assignments are provided in Table S2.

Functional Enrichment Analysis (FEA)

Each module was annotated using the gprofiler software (v2_0.2.1) via the g:GOSt R function [70]. This

tool maps genes to functional information sources and detects statistically significantly enriched

terms. These sources include the GO database, pathways from KEGG, Reactome, and

WikiPathways, miRNA targets from miRTarBase, regulatory motif matches from TRANSFAC, tissue

specificity from the Human Protein Atlas, protein complexes from CORUM, and human disease

phenotypes from the Human Phenotype Ontology. For multiple testing correction, we used the default

g:SCS method unless otherwise specified. The FEA results are available in Table S4.

Module Preservation (MP) Analysis

To assess whether the modules were preserved across distinct UMAP clusters, we used the MP

function from the WGCNA R package (v1.70-3)[71]. Pairwise comparisons were performed for all 193

modules. We then estimated the mean and variance of the preservation statistic under the null

hypothesis that there is no relationship between module assignments in the reference and test

datasets. Fifteen distinct preservation statistics were calculated (e.g., module density, mean

expression, connectivity), each standardized by its mean and variance to define a Z statistic for each

preservation measure. Under these assumptions, each Z statistic follows a normal distribution if the

module is not preserved. To compare preservation across modules, we calculated a composite

preservation measure using the following equation:

$$Zsummary = \frac{Zdensity + Zconnectivity}{2}$$

384   Modules with Z < 2 were considered not preserved, those with Z between 2 and 10 were considered

385   preserved, and modules with Z > 10 were considered highly preserved. Zsummary stats are available

386   in Table S3.

Statistical analysis to identify AD-differential genes

For each brain section of a subject, we summed the UMI counts of all spots belonging to a subregion

to generate a pseudobulk estimate for each gene. We only kept expressed genes with counts per

million (CPM) > 1 in at least 80% of the brain sections, resulting in 1,578 lncRNAs and 13,808

mRNAs. With the pseudobulk estimates, we estimated the library sizes of the brain sections using the

trimmed mean of M-values (TMM) of edgeR[72] (edgeR_3.36.0) and transformed the pseudobulk

estimates into $\log_2$ CPM using voom-limma (limma_3.50.1). For each subregion, we excluded brain

sections with <50 spots. To contrast the normalized pseudobulk expression between AD and controls,

we applied linear mixed models (LMM), accounting for age, PMI, RIN, and batch as fixed effects and

modeled within-subject correlation with a random effect. We declared significance at an α of 0.05 with

false discovery rate (FDR) correction across all genes and all subregions.

For interpretation, we applied GSEA[73] with AD-differential t-values of each subregion as scores, and

GO terms and canonical pathways as genesets. To characterize lncRNAs with respect to brain cell

types, we used the pseudobulk expression data from a published snRNAseq dataset of 424

ROSMAP individuals[15]. For each of the major cell types, we contrasted pseudobulk expression of AD

subjects vs. controls using multiple regression, accounting for age, PMI, RIN, and batch. For further

comparison, we repeated this AD-differential expression analysis with bulk DLPFC RNA-seq data

from 1049 individuals[20].

Follow-up Analysis on *OIP5-AS1*

We used the eQTL statistics from the published snRNAseq dataset[15]. We first assembled the eQTL

statistics from all SNPs within 1Mb from the transcription start site of *OIP5-AS1*. This was done for six

major brain cell types (astrocytes, excitatory neurons, inhibitory neurons, microglia, oligodendrocytes,

and OPC) that showed adequate expression across subjects. We then applied mvSuSiE[36], which

aggregates the statistics across SNPs and cell types to estimate the posterior inclusion probability

that a given SNP casually affects expression of *OIP5-AS1*.

In addition, we further tested the interaction effects of *OIP5-AS1* and HDACs on 3 categories of AD traits: cognitive decline, neuritic plaques, and tau tangles. For cognitive decline, scores from tests on episodic memory (ep), semantic memory (se), working memory (wo), presentation speed (ps), and presentation order (po) were examined with their aggregate (global) being the trait of focus. For neuritic plaques, measurements were taken from medial frontal (mf), medial temporal (mt), anterior gyrus (ag), entorhinal cortex(ec), and hippocampus (hip) with their aggregate as the trait of focus. For tau tangles, measurements were taken from mf, ag, ec, hip, inferior temporal (it), cingulate gyrus (cg), and mesial temporal (mes_t) with their aggregate as the trait of focus. For each of the six major brain cell types[15] and each HDAC protein detected in a bulk DLPFC dataset (HDACs 2, 4, 5, 6, 8, 11)[37], we applied multiple regression to model an AD trait by the main effect of *OIP5-AS1* expression, the main effect of a HDAC protein, and their interaction effect, while accounting for age, PMI, RIN, and batch. To aid interpretation, we further divided the subjects into tertiles based on *OIP5-AS1* expression and examined how each AD trait relates to HDAC abundance for each tertile. The same analysis was repeated with HDAC protein abundance replaced by HDAC expression level from the snRNA-seq dataset.

RNAscope and Immunofluorescence co-staining and analysis

Human FFPE DLPFC sections were stained with neuron (MAP2; Sigma #M2320-100UL) and microglia (IBA1; Wako #019-19741) markers followed by RNAscope with Hs-OIP5-AS1-O1-C1 probe (ACD #1559738-C1) using a Bond-Rx research stainer (Leica Biosystems) using an Opal 6-plex detection kit (Akoya Biosciences #NEL871001KT). Briefly, sections were baked for 30 min at 60 °C prior to dewaxing for 30 s at 72 °C. Sections were antigen retrieved using HIER with a pH 6.0 citrate buffer solution (Leica Biosystems #AR9961) for 20 min at 100 °C followed by 10 min peroxide block. MAP2 (1:100) was incubated for 30 min followed by 10 min Opal polymer HRP Ms+Rb and 10 min Opal signal development with Opal 520 diluted at 1:100. Sections were antigen retrieved for 20 min in ph 6.0 buffer again and the process is repeated with IBA1 (1:500) paired with Opal 690 (1:150). The sections were then stained with RNAscope probe against *OIP5-AS1* using RNAScope LS Multiplex Fluorescent Reagent Kit (ACD #322800) according to the manufacturer with the following modifications. The sections were antigen retrieved using HIER with a pH 6.0 buffer for 20 min at 100 °C, washed, and treated with ACD Protease III for 15 min at 40 °C prior to probe hybridization for 2 h at 40 °C. Opal 620 (1:500) was used for signal development. Slides were counter stained with 10X spectral DAPI for 5 min before mounting. Imaging was performed using Phenocycler-Fusion 2.0 (Akoya Biosciences) at 40X. Images were spectrally umixed using inForm (Akoya Biosciences) and composite TIFs exported to QuPath (v0.3.2) for quantitative analysis. Regions of interest that

encompassed cortical layers were identified and selected as ROIs based on tissue cytoarchitecture

patterns from nuclei and neuron stains. Cells were defined in each ROI using QuPath Cell Detection

tool with the following parameters, requested pixel size = 0.5µm, background radius = 8µm, median

filter radius =0µm, Sigma value for Gaussian filter = 1.5µm, minimum nucleus area = 10µm2 ,

maximum nucleus area =400µm2 , and intensity threshold = 10. Cell parameters include a cell

expansion of 5 µm from the cell nucleus and general parameters include smoothing of the detected

nucleus/cell boundaries. To quantify OIP5-AS1, subcellular spot parameters were set to detection

threshold = 0.08, split by intensity and shape to true, expected spot size = 0.05µm2, min spot size =

0.01µm2, max spot size =1µm2 and include clusters set to true. For cell type specific expression,

cells were thresholded on their MAP2 and IBA1 mean cell intensity by section and cells of ambiguous

identity were not included in the analysis.

References

1.   Mattick, J. S. *et al.* Long non-coding RNAs: definitions, functions, challenges and recommendations. *Nat. Rev. Mol. Cell Biol.* **24**, 430–447 (2023).

2.   Statello, L., Guo, C.-J., Chen, L.-L. & Huarte, M. Gene regulation by long non-coding RNAs and its biological functions. *Nat. Rev. Mol. Cell Biol.* **22**, 96–118 (2020).

3.   Wang, K. C. & Chang, H. Y. Molecular mechanisms of long noncoding RNAs. *Mol. Cell* **43**, 904–914 (2011).

4.   Guttman, M. *et al.* Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature* **458**, 223–227 (2009).

5.   Kung, J. T. Y., Colognori, D. & Lee, J. T. Long noncoding RNAs: past, present, and future. *Genetics* **193**, 651–669 (2013).

6.   Djebali, S. *et al.* Landscape of transcription in human cells. *Nature* **489**, 101–108 (2012).

7.   Cabili, M. N. *et al.* Integrative annotation of human large intergenic noncoding RNAs reveals global properties and specific subclasses. *Genes Dev.* **25**, 1915–1927 (2011).

8.   Derrien, T. *et al.* The GENCODE v7 catalog of human long noncoding RNAs: analysis of their gene structure, evolution, and expression. *Genome Res.* **22**, 1775–1789 (2012).

9.   Wu, X., Xia, P., Yang, L., Lu, C. & Lu, Z. The roles of long non-coding RNAs in Alzheimer's disease diagnosis, treatment, and their involvement in Alzheimer's disease immune responses. *Noncoding RNA Res.* **9**, 659–666 (2024).

10. Policarpo, R. & d'Ydewalle, C. Missing lnc(RNAs) in Alzheimer's Disease? *Genes* **13**, (2021).

11. Lan, Z., Chen, Y., Jin, J., Xu, Y. & Zhu, X. Long non-coding RNA: Insight into mechanisms of Alzheimer's disease. *Front. Mol. Neurosci.* **14**, 821002 (2021).

12. Huang, Z., Zhao, J., Wang, W., Zhou, J. & Zhang, J. Depletion of LncRNA NEAT1 rescues mitochondrial dysfunction through NEDD4L-dependent PINK1 degradation in animal models of Alzheimer's disease. *Front. Cell. Neurosci.* **14**, 28 (2020).

13. Riva, P., Ratti, A. & Venturin, M. The Long Non-Coding RNAs in Neurodegenerative Diseases: Novel Mechanisms of Pathogenesis. *Curr. Alzheimer Res.* **13**, 1219–1231 (2016).

14. Avey, D. R. *et al.* Uncovering plaque-Glia niches in human Alzheimer's Disease brains using spatial transcriptomics. *bioRxivorg* (2024) doi:10.1101/2024.09.05.611566.

15. Fujita, M. *et al.* Cell subtype-specific effects of genetic variation in the Alzheimer's disease brain. *Nat. Genet.* **56**, 605–614 (2024).

16. Gaiteri, C. *et al.* Identifying robust communities and multi-community nodes by combining top-down and bottom-up approaches to clustering. *Sci. Rep.* **5**, 1–14 (2015).

17. Li, D. *et al.* Insights into lncRNAs in Alzheimer's disease mechanisms. *RNA Biol.* **18**, 1037–1047 (2021).

18. Bellenguez, C. *et al.* New insights into the genetic etiology of Alzheimer's disease and related dementias. *Nat. Genet.* **54**, 412–436 (2022).

19. Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: Precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biol.* **15**, R29 (2014).

20. Mostafavi, S. *et al.* A molecular network of the aging human brain provides insights into the pathology and cognitive decline of Alzheimer's disease. *Nat. Neurosci.* **21**, 811–819 (2018).

21. Shukla, S. & Tekwani, B. L. Histone deacetylases inhibitors in neurodegenerative diseases, neuroprotection and neuronal differentiation. *Front. Pharmacol.* **11**, 537 (2020).

22. Govindarajan, N. *et al.* Reducing HDAC6 ameliorates cognitive deficits in a mouse model for

Alzheimer's disease. *EMBO Mol. Med.* **5**, 52–63 (2013).

23. Onishi, T. *et al.* A novel orally active HDAC6 inhibitor T-518 shows a therapeutic potential for
Alzheimer's disease and tauopathy in mice. *Sci. Rep.* **11**, 15423 (2021).

24. Yang, S.-S., Zhang, R., Wang, G. & Zhang, Y.-F. The development prospection of HDAC
inhibitors as a potential therapeutic direction in Alzheimer's disease. *Transl. Neurodegener.* **6**, 19
(2017).

25. Zhang, M. *et al.* Histone deacetylase inhibitors VPA and WT161 ameliorate the pathological
features and cognitive impairments of the APP/PS1 Alzheimer's disease mouse model by
regulating the expression of APP secretases. *Alzheimers. Res. Ther.* **16**, 15 (2024).

26. Kumar, V., Kundu, S., Singh, A. & Singh, S. Understanding the role of histone deacetylase and
their inhibitors in neurodegenerative disorders: Current targets and future perspective. *Curr.*
Neuropharmacol. **20**, 158–178 (2022).

27. Ru, J. *et al.* Technologies of targeting histone deacetylase in drug discovery: Current progress
and emerging prospects. *Eur. J. Med. Chem.* **261**, 115800 (2023).

28. Yang, J.-H. *et al.* LncRNA OIP5-AS1-directed miR-7 degradation promotes MYMX production
during human myogenesis. *Nucleic Acids Res.* **50**, 7115–7133 (2022).

29. Zhuang, A. *et al.* Loss of the long non-coding RNA OIP5-AS1 exacerbates heart failure in a sex-
specific manner. *iScience* **24**, 102537 (2021).

30. Hunkler, H. J. *et al.* The long non-coding RNA Cyrano is dispensable for pluripotency of Murine
and human pluripotent stem cells. *Stem Cell Reports* **15**, 13–21 (2020).

31. Kotake, Y., Matsunaga, N., Wakasaki, T. & Okada, R. OIP5-AS1 promotes proliferation of non-
small-cell lung cancer and head and neck squamous cell carcinoma cells. *Cancer Genomics*
Proteomics **18**, 543–548 (2021).

32. Wu, L., Liu, Y., Guo, C. & Shao, Y. LncRNA OIP5-AS1 promotes the malignancy of pancreatic
ductal adenocarcinoma via regulating miR-429/FOXD1/ERK pathway. *Cancer Cell Int.* **20**, 296
(2020).

33. Arunkumar, G. *et al.* LncRNA OIP5-AS1 is overexpressed in undifferentiated oral tumors and
integrated analysis identifies as a downstream effector of stemness-associated transcription
factors. *Sci. Rep.* **8**, 7018 (2018).

34. Gong, M. *et al.* OIP5 is a novel prognostic biomarker in clear cell renal cell cancer correlating
with immune infiltrates. *Front. Immunol.* **13**, 805552 (2022).

35. Li, Z. *et al.* Exosomal OIP5-AS1 attenuates cerebral ischemia-reperfusion injury by negatively
regulating TXNIP protein stability and inhibiting neuronal pyroptosis. *Int. Immunopharmacol.* **127**,
111310 (2024).

36. Zou, Y., Carbonetto, P., Xie, D., Wang, G. & Stephens, M. Fast and flexible joint fine-mapping of
multiple traits via the Sum of Single Effects model. *bioRxivorg* (2024)
doi:10.1101/2023.04.14.536893.

37. Johnson, E. C. B. *et al.* Large-scale deep multi-layer analysis of Alzheimer's disease brain
reveals strong proteomic disease-related changes not observed at the RNA level. *Nat. Neurosci.*
25, 213–225 (2022).

38. Jiang, C. *et al.* Identifying and functionally characterizing tissue-specific and ubiquitously
expressed human lncRNAs. *Oncotarget* **7**, 7120–7133 (2016).

39. de Goede, O. M. *et al.* Population-scale tissue transcriptomics maps long non-coding RNAs to
complex disease. *Cell* **184**, 2633–2648.e19 (2021).

40. Li, F. *et al.* The effect of BACE1-AS on β-amyloid generation by regulating BACE1 mRNA

expression. *BMC Mol. Biol.* **20**, 23 (2019).

41. Faghihi, M. A. *et al.* Expression of a noncoding RNA is elevated in Alzheimer's disease and
drives rapid feed-forward regulation of beta-secretase. *Nat. Med.* **14**, 723–730 (2008).

42. Zhang, W., Zhao, H., Wu, Q., Xu, W. & Xia, M. Knockdown of BACE1-AS by siRNA improves
memory and learning behaviors in Alzheimer's disease animal model. *Exp. Ther. Med.* **16**, 2080–
2086 (2018).

43. Modarresi, F. *et al.* Inhibition of natural antisense transcripts in vivo results in gene-specific
transcriptional upregulation. *Nat. Biotechnol.* **30**, 453–459 (2012).

44. Jiao, S.-S. *et al.* Brain-derived neurotrophic factor protects against tau-related neurodegeneration
of Alzheimer's disease. *Transl. Psychiatry* **6**, e907 (2016).

45. Ding, Y., Luan, W., Shen, X., Wang, Z. & Cao, Y. LncRNA BDNF-AS as ceRNA regulates the
miR-9-5p/BACE1 pathway affecting neurotoxicity in Alzheimer's disease. *Arch. Gerontol. Geriatr.*
99, 104614 (2022).

46. Ma, P. *et al.* Long non-coding RNA MALAT1 inhibits neuron apoptosis and neuroinflammation
while stimulates neurite outgrowth and its correlation with MiR-125b mediates PTGS2, CDK5 and
FOXQ1 in Alzheimer's disease. *Curr. Alzheimer Res.* **16**, 596–612 (2019).

47. Bernard, D. *et al.* A long nuclear-retained non-coding RNA regulates synaptogenesis by
modulating gene expression. *EMBO J.* **29**, 3082–3093 (2010).

48. Dong, L.-X. *et al.* LncRNA NEAT1 promotes Alzheimer's disease by down regulating micro-27a-
3p. *Am. J. Transl. Res.* **13**, 8885–8896 (2021).

49. Ke, S. *et al.* Long noncoding RNA NEAT1 aggravates Aβ-induced neuronal damage by targeting
miR-107 in Alzheimer's disease. *Yonsei Med. J.* **60**, 640–650 (2019).

50. Bútová, R. *et al.* LncRNAs LY86-AS1 and VIM-AS1 distinguish plasma cell leukemia patients
from multiple myeloma patients. *Biomedicines* **9**, 1637 (2021).

51. Kurniansyah, N. *et al.* An integrated multi-omics analysis of sleep-disordered breathing traits
implicates P2XR4 purinergic signaling. *Commun. Biol.* **6**, 125 (2023).

52. Tan, Y. *et al.* lncRNA-associated ceRNA network revealing the potential regulatory roles of
ferroptosis and immune infiltration in Alzheimer's disease. *Front. Aging Neurosci.* **15**, 1105690
(2023).

53. Bajic, V. P. *et al.* The X files: 'the mystery of X chromosome instability in Alzheimer's disease'.
Front. Genet. **10**, 1368 (2019).

54. Garcia, A. X., Xu, J., Cheng, F., Ruppin, E. & Schäffer, A. A. Altered gene expression in
excitatory neurons is associated with Alzheimer's disease and its higher incidence in women.
Alzheimers Dement. (N. Y.) **9**, e12373 (2023).

55. Sun, X. *et al.* The MAOA gene influences the neural response to psychosocial stress in the
human brain. *Front. Behav. Neurosci.* **14**, 65 (2020).

56. Ziermans, T. *et al.* Working memory brain activity and capacity link MAOA polymorphism to
aggressive behavior during development. *Transl. Psychiatry* **2**, e85 (2012).

57. Chua, G. N. L. *et al.* Differential dynamics specify MeCP2 function at nucleosomes and
methylated DNA. *Nat. Struct. Mol. Biol.* 1–9 (2024).

58. Chu, S.-L. *et al.* Phosphoglycerate kinase 1 acts as a cargo adaptor to promote EGFR transport
to the lysosome. *Nat. Commun.* **15**, 1021 (2024).

59. Vinokurov, A. Y. *et al.* HPRT1 deficiency induces alteration of mitochondrial energy metabolism
in the brain. *Mol. Neurobiol.* **60**, 3147–3157 (2023).

60. Wooten, S. & Smith, K. N. Long non-coding RNA OIP5-AS1 (Cyrano): A context-specific

regulator of normal and disease processes. *Clin. Transl. Med.* **12**, e706 (2022).

61.　Mei, J. *et al.* OIP5-AS1 modulates epigenetic regulator HDAC7 to enhance non-small cell lung
cancer metastasis via miR-140-5p. *Oncol. Lett.* **20**, 7 (2020).

62.　Kleaveland, B., Shi, C. Y., Stefano, J. & Bartel, D. P. A network of noncoding regulatory RNAs
acts in the mammalian brain. *Cell* **174**, 350–362.e17 (2018).

63.　Mummery, C. J. *et al.* Tau-targeting antisense oligonucleotide MAPTRx in mild Alzheimer's
disease: a phase 1b, randomized, placebo-controlled trial. *Nat. Med.* **29**, 1437–1447 (2023).

64.　Kuo, H.-C., Prupes, J., Chou, C.-W. & Finkelstein, I. J. Massively parallel profiling of RNA-
targeting CRISPR-Cas13d. *Nat Commun* **15**, (2024).

65.　Wessels, H.-H. *et al.* Massively parallel Cas13 screens reveal principles for guide RNA design.
Nat. Biotechnol. **38**, 722–727 (2020).

66.　Bennett, D. A. *et al.* Religious Orders Study and Rush Memory and Aging Project. *J. Alzheimers.*
Dis. **64**, S161–S189 (2018).

67.　Bennett, D. A. *et al.* Natural history of mild cognitive impairment in older persons. *Neurology* **59**,
198–205 (2002).

68.　Bennett, D. A. *et al.* Neuropathology of older persons without cognitive impairment from two
community-based studies. *Neurology* **66**, 1837–1844 (2006).

69.　Bennett, D. A. *et al.* Decision rules guiding the clinical diagnosis of Alzheimer's disease in two
community-based cohort studies compared to standard practice in a clinic-based cohort study.
Neuroepidemiology **27**, 169–176 (2006).

70.　Kolberg, L. *et al.* g:Profiler-interoperable web service for functional enrichment analysis and gene
identifier mapping (2023 update). *Nucleic Acids Res.* **51**, W207–W212 (2023).

71.　Langfelder, P., Luo, R., Oldham, M. C. & Horvath, S. Is my network module preserved and
reproducible? *PLoS Comput. Biol.* **7**, e1001057 (2011).

72.　Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

73.　Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for
interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences*
102, 15545–15550 (2005).

Main Figures

645
646
647 **Figure 1. Spatial expression of lncRNAs. (A)** Study overview. **(B)** Comparison of the expression distributions
648 of lncRNAs and protein-coding genes (mRNAs). **(C)** Comparison of percentage of ST spots with non-zero
649 expression for lncRNAs and mRNAs. **(D)** Proportion of detected lncRNAs (aggregated across subregions)
650 belonging to each biotype. **(E)** Upset plots showing the numbers of lncRNAs (left) and mRNAs (right) detected
651 in the top 20 most common intersections of subregions.
652

Figure 2. Subregion- and cell-type-enriched lncRNAs. (A) Heatmap showing scaled expression (z-score) of the top 50 subregion-specific lncRNAs. The last column to the right shows the percentage of spots with non-zero expression for each lncRNA (*exemplar lncRNAs shown in panel B). **(B)** UMAP and expression feature plots for 4 representative sections, showing 5 examples of lncRNAs with enriched expression in specific cortical layers or white matter (WM). **(C)** Violin plots showing expression distributions among 6 major brain cell types for the 5

lncRNAs shown in (B). **(D)** Scaled snRNA-seq expression of the same 50 subregion-specific lncRNAs shown in (A) for the 6 most abundant brain cell types (*exemplar lncRNAs shown in panel B; n.d.: not detected). **(E)** Scaled expression by cell type (top) and ST subregion (bottom) for the top 10 cell-type-specific lncRNAs from each of the 6 major brain cell types.

716
717
Figure 3. Subregion-specific gene modules.
(A) Schematic of gene expression module generation and downstream analyses. **(B)** Heatmap showing the
percentage of preserved modules between each pair of subregions. **(C)** Stacked barplot shows the number of
modules in each subregion. Red colors for the lncRNA-enriched modules. **(D)** Network connectivity map for L2-
3 m14, illustrating high connectivity of lncRNAs (correlation > 0.9). **(E)** Enriched Gene Ontology (GO) terms (FDR
at 0.05) from functional enrichment analysis of an example lncRNA-enriched module (L2-3 m14). **(F)** Network
connectivity map for WM module 11, illustrating high connectivity of lncRNAs (correlation > 0.95). **(G)** Enriched
GO terms (FDR at 0.05) from functional enrichment analysis for the WM module 11. **(E, G)** The GO biological
process terms were collapsed, keeping only the top GO parent functions to avoid redundancy. The y-axis and
bar size represent the significance in -log₁₀(Adj. p-Value). GO biological process (BP), cellular component (CC),
and GO molecular function (MF).
729

Figure 4. Differential lncRNAs and mRNAs in AD and cross-modality comparison with bulk RNA-seq. **(A)** Histogram showing the distribution of AD-differential t-values for mRNAs and lncRNAs. **(B)** Total numbers of AD-differential lncRNAs and mRNAs for each subregion. **(C)** Heatmap showing the enrichment of AD-differential lncRNAs for specific biotypes. **(D)** Overlaps in AD-differential lncRNAs and mRNAs between ST and bulk RNA-seq. **(E)** Scatterplots of ST (y-axis) vs. bulk RNA-seq (x-axis) AD-differential t-values for lncRNAs (left) and mRNAs (right). Correlation value (R) is shown at the top left of each plot.

Figure 5. Characterization of AD-differential lncRNAs detected by ST. (A) The percentage of DEGs in each ST subregion, split by lncRNA/mRNA, up/down in AD, and whether they are subregion-specific (ss) or not (nss). Fisher's exact test was applied to assess differences in subregion-specificity between lncRNA and mRNA with p-values indicated. **(B)** (top) Heatmap showing t-values of AD-differential lncRNAs passing FDR correction specifically for one ST subregion, i.e. subregion-specific (*p<5e-4 in designated subregion and p>0.001 in all others). (bottom) AD-differential t-values for 6 brain cell types. An additional row below shows the percentage of ST spots with non-zero expression for each lncRNA. **(C)** (left) Heatmap showing t-values of AD-differential lncRNAs passing FDR correction in two or more ST subregions, i.e. not subregion-specific (*p<5e-4). An additional row below shows the percentage of ST spots with non-zero expression for each lncRNA. (right) AD-differential t-values for a subset of 13 lncRNAs with previously reported AD association, including t-values from DLPFC bulk RNA-seq below. **(D)** XIST spatial expression plots for two control (Ctrl) and two AD sections. **(E)** Boxplots of subregion-specific pseudobulk expression from ST data for XIST, MECP2, and PGK1 (X-linked genes), all of which show altered expression in AD in one or more subregions. **(F)** Sankey diagram and heatmap showing a subset of enriched genesets and AD differential lncRNAs belonging to those genesets. **(G)** A subset of HDAC genes display AD-differential expression in specific ST subregions (*p<5e-4).

809
810
Figure 6. Genetic analysis of *OIP5-AS1* and its interaction with HDAC. (A) Colocalization analysis of *OIP5-*
AS1 eQTLs across major brain cell types. Applying multivariate colocalization found rs1655558 to be a
potential driver of *OIP5-AS1* expression in astrocytes, oligodendrocytes, and excitatory neurons (Ast:
Astrocyte, Exc: excitatory, Inh: Inhibitory, Mic: Microglia, Oli: Oligodendrocyte, OPC: Oligodendrocyte
progenitor cell, PIP: posterior inclusion probability; cartoon cell type images created in BioRender.com). **(B)**
Summary of *OIP5-AS1*xHDAC interaction analysis. snRNAseq[14] and proteomic[37] data from human DLPFC
were used to associate the statistical interaction between *OIP5-AS1* expression level (in specific cell types)
and protein abundance of each of six HDACs with AD phenotypes, including cognitive decline (cogdx), amyloid
plaques, and tau tangles. Cognition was assessed over multiple domains and pathologies were measured in

multiple brain regions. **(C)** Each point in the violin plot represents the –log10(p) of the interaction effect of *OIP5-AS1* and HDAC for a given combination of cell type, HDAC, and AD phenotype. Higher associations were observed for plaques and tangles. **(D)** Scatterplots showing t-values of the HDAC main effect (x-axis) and the *OIP5-AS1xHDAC* interaction for tangles measured in multiple brain regions**.** Dots are colored but not circled if interaction p<0.05 and an order of magnitude lower than the main effect p-value. Circled dots are interactions passing multiple comparison correction for the number of HDACs and cell types. *OIP5-AS1xHDAC11* interaction shows the strongest association with tangles. **(E)** Heatmap of *OIP5-AS1xHDAC11* interaction p-values across cell types and tangles in different brain regions shown (ag: anterior gyrus, ec: entorhinal cortex, hip: hippocampus, mf: medial frontal, mt: medial temporal, cg: cingulate gyrus, it: inferior temporal, mes_t: mesial temporal). **(F)** An illustration of interaction effect. Individuals were stratified into three groups based on the expression of *OIP5-AS1* in excitatory neurons. For each group, HDAC11 protein abundance was plotted against tangle level in the hippocampus. **(G-I)** are plaque burden counterparts of (D-F).

Supplementary Figures

Figure S1. Differential lncRNAs and mRNAs in AD and cross-modality comparison with snRNA-seq. (**A**) Histogram showing the distribution of AD-differential t-values for mRNAs and lncRNAs. (**B**) Total numbers of AD-differential lncRNAs and mRNAs for each subregion. (**C**) Heatmap showing the enrichment of AD-differential lncRNAs for specific biotypes. (**D**) The percentage of DEGs in each snRNA cell type, split by lncRNA/mRNA, up/down in AD, and whether they are cell-type-specific (cs) or not (ncs). Fisher's exact test was applied to assess differences in cell-type-specificity between lncRNA and mRNA with p-values indicated. (**E**) Overlaps in AD-differential lncRNAs and mRNAs between ST and snRNA-seq. (**F-G**) Heatmap showing correlation (R) between subregions and cell types based on their AD differential t-values (*p<0.05) for lncRNAs (F) or mRNAs (G).

882
Figure S2. Validation of *OIP5-AS1* expression in human DLPFC by RNAscope.
(A) Representative RNAscope image of human DLPFC showing transcripts of *OIP5-AS1 a*nd
immunostaining for MAP2 and IBA1. Scale bars = 100 µm (overview), 20 µm (insets). (B) Boxplots of
mean *OIP5-AS1* spot quantification in cortical layers, range n =103-1788 cells per layer from 8
individuals. (C) Boxplots of mean *OIP5-AS1* spot quantification by cell type, derived from a minimum
of 200 MAP2+ neurons and 50 IBA1+ microglia for each of 8 individuals.

Figure S3. Association of lncRNA *OIP5-AS1* with AD phenotypes using cell-type-specific expression of HDAC RNA. (**A**) RNA expression of *OIP5-AS1* and HDAC genes in designated snRNA-seq cell types and bulk DLPFC RNA-seq. (**B-D**) Scatterplots showing t-values of the HDAC main effect (x-axis) and the *OIP5-AS1*xHDAC interaction for cognition (**B**), plaque burden (**C**), and tangles (**D**) measured in multiple brain regions. Dots are colored but not circled if interaction p<0.05 and an order of magnitude lower than the main effect p-value. The circled dot in (**B**), Inh:*HDAC7*, is the only interaction passing multiple comparison correction for the number of HDACs and cell types.

Supplementary Tables

926
927
Table S1. Subregion- and cell-type-enriched lncRNAs
Table S2. ST gene expression module assignments
Table S3. ST gene expression module preservation
Table S4. ST gene expression module FEA
Table S5. ST gene expression module AD trait associations
Table S6. AD-differential lncRNAs (ST, bulk, snRNA)
Table S7. AD-differential GSEA of ST genes
Table S8. AD-differential genesets including lncRNAs
936
937

943