Spatial Expression of Long Non-Coding RNAs in Human Brains of Alzheimer's Disease

Bernard Ng[1, #], Denis R. Avey[1, 5, #], Katia de Paiva Lopes[1], Masashi Fujita[2], Shinya Tasaki[1], Ricardo

A. Vialle[1], Artemis Iatrou[1,3], Sashini De Tissera[1], Tien-Hao Chang[1], Jishu Xu[1], Himanshu Vyas[1],

Nicola A. Kearns[1, 5], Vilas Menon[2], Chris Gaiteri[1,4], Philip L. De Jager[2], David A. Bennett[1, 5], Yanling

Wang[1, 5, *]

1. Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL, 60612, USA

2. Center for Translational and Computational Neuroimmunology, Department of Neurology, Columbia

University Irving Medical Center, New York, NY, USA

3. Current affiliation: Department of Psychiatry, McLean Hospital, Harvard Medical School, Belmont,

MA, 02478, USA

4. Department of Psychiatry, Upstate Medical University, Syracuse, NY, 13210, USA

5. Department of Neurological Sciences, Rush University Medical Center, Chicago, IL, 60612, USA

[#]Contributed equally

*Correspondence to yanling_wang@rush.edu

Abstract

Long non-coding RNAs (lncRNAs) are highly versatile in their modes of action and play critical roles in

both normal physiological and disease processes. The dysregulation of lncRNA expression has been

implicated in aging and many neurological disorders, including Alzheimer's disease (AD). Here, we

report a spatial expression atlas of 7,634 lncRNA genes in aged human brains, covering 258,987

microdomains from 78 postmortem brain sections of 21 ROSMAP participants. We detected a greater

proportion of lncRNAs expressed specifically in distinct cortical subregions than mRNAs, and most

belong to antisense and lincRNA biotypes. We generated 193 gene modules from gene networks of 8

subregions and identified lncRNA-enriched gene modules implicated in multiple AD-relevant biological

processes. By cross-referencing published snRNA-seq data, we detected a partially overlapping but

independent pattern between sub-region and cell-type specificity of lncRNA expression. We mapped

spatially differentially expressed (SDE) lncRNAs and mRNAs in AD brains and observed that SDE

lncRNAs encompass more subregion-specific transcripts. Gene set enrichment analysis indicates that

AD SDE lncRNAs are involved in epigenetic regulation, chromatin remodeling, RNA metabolism,

synaptic signaling, and apoptosis. Particularly relevant to AD therapeutic potentials is an enrichment

for HDAC target genes, including *OIP5-AS1*, a lncRNA extensively studied in cancer and, most recently,

in AD. We then applied multivariate fine-mapping to the expression quantitative trait loci (eQTLs)

proximal to the *OIP5-AS1* loci and identified rs1655558 as a potential genetic driver of *OIP5-AS1*

expression. Using statistical modeling, we inferred that the interaction between *OIP5-AS1* and HDAC

proteins, especially HDAC11, is associated with tau tangles in excitatory neurons and plaque burden

in microglia. Our study represents a valuable resource of lncRNA spatial expression in the aged human

brain, shedding mechanistic insights into their functional roles in AD and neurodegeneration.

Introduction

One remarkable yet unexpected discovery of the eukaryotic transcriptome in the 21st century is that a

large portion of the genome is actively transcribed into RNAs with little or no protein-coding potential[1,2].

Specifically, about 90% of the human genome is transcribed, but only about 2% is annotated as protein-

coding genes. Since this discovery, there has been a surge of publications highlighting the regulatory

roles of non-coding RNAs (ncRNAs) in controlling gene expression[2,3], debunking the long-held belief

that functional genetic information is only contained in protein-coding genes and RNAs are mere

intermediaries along the central dogma between DNA and proteins.

Long non-coding RNAs (lncRNAs) belong to a class of ncRNAs that are longer than 200 nucleotides

and are generated mainly by Polymerase II, with little or no protein-coding potential[4,5]. Accumulated

studies show that lncRNAs affect gene expression through various regulatory mechanisms, including

epigenetic modification, chromatin modeling, and post-transcriptional and translational regulations.

Compared with coding RNAs, lncRNAs exhibit substantially greater tissue- and cell-type-specificity[6,7].

Notably, the number of lncRNAs often increases with the complexity of organisms. In the human

genome, approximately 40% of lncRNAs are explicitly expressed in the brain, and many of those

lncRNAs are involved in various biological processes, including neuronal differentiation,

synaptogenesis, and neuroplasticity[1,8].

The dysregulation of lncRNA expression has been implicated in many neurological disorders, including

brain aging and Alzheimer's disease (AD). In AD, aberrant expression of lncRNAs has been found in

the brains of AD animal models and AD patients. Those dysregulated lncRNAs have been implicated

in various aspects of AD pathogenesis, such as amyloid-beta metabolism, tau protein pathology,

neuron inflammation, and cell death[9–13]. Despite these insights, our understanding of the spatial

expression patterns of lncRNAs in the human brain remains limited, hindering our understanding of

their functional roles in health and disease contexts.

Here, we report a spatial expression atlas of 7,634 lncRNA genes in aged human brains, covering

258,987 microdomains from 78 brain sections across 21 ROSMAP participants with and without AD

dementia. We identify spatially resolved lncRNA genes and lncRNA-enriched gene modules. We cross-
reference ROSMAP snRNA-seq data[14] to explore the patterns between sub-region and cell-type
specificity of lncRNA expression. We map spatially differentially expressed (SDE) lncRNAs and mRNAs
in AD brains and compare our ST findings with those from bulk RNA-seq and snRNA-seq. Finally, we
infer the potential functions of SDE lncRNAs and uncover a potential interaction between *OIP5-AS1*
and HDACs in mediating AD pathogenesis. Our study represents a valuable resource of lncRNA spatial
expression in the aged human brain, shedding mechanistic insights into their functional roles in AD and
neurodegeneration.

Results

Spatial expression of lncRNAs in aged human brains

Leveraging our recently acquired spatial transcriptomic data[15], we analyzed lncRNA expression in the
dorsolateral prefrontal cortex (DLPFC) from 78 brain sections across 21 female individuals (AD=13;
NCI=8). In total, we detected 7,634 lncRNAs and 16,769 protein-coding mRNAs across all spot clusters,
including Layer (L)1, L2-3, L3-5, L5, and L6, white matter (WM), meninges, and blood vessels (Fig 1A).
As expected, most lncRNAs present lower expressions than mRNAs (Fig. 1B). Out of the total ~260K
spots, we only detected lncRNAs in <0.01% of spots (Fig. 1C), and those lncRNAs are mostly long
intergenic non-coding RNAs (lincRNA) or antisense (Fig. 1D). We then assessed the subregion-
specificity of lncRNAs. We found a greater proportion of lncRNAs expressed in a single subregion (or
distinct subsets of subregions) than mRNAs (Fig. 1E).

Contrasting the expression of ST spots in one subregion against all other subregions, we identified 79
subregion-specific lncRNAs (Fig. 2A-B, Table S1). To examine whether subregion-specific lncRNA
aligns with cell-type-specific expression and vice versa, we cross-referenced a published snRNA-seq
dataset derived from the DLPFC of 424 ROSMAP individuals[14]. We found a subset of cell-type specific
lncRNAs (Table S1) aligns with the ST subregion-specific lncRNAs (Fig. 2C; Table S1). For example,
LINC00844 is highly expressed in WM (Fig 2A-B) and oligodendrocytes (Fig. 2C-D). In general,
astrocyte-specific lncRNAs are more abundantly expressed in L1, excitatory/inhibitory neuron-specific
lncRNAs in cortical layers (especially L2-3), and oligodendrocyte-specific lncRNAs in WM (Fig. 2E).
However, the majority of the subregion-specific lncRNAs are not cell-type-specific and vice versa (Fig.
2C, 2E), indicating a partially overlapping but independent relationship between sub-region and cell-
type specificity of lncRNA expression.

Spatially resolved lncRNA-enriched gene modules in aged human brains

We reason that lncRNAs may work with mRNAs in the same gene networks. To this end, we derived pseudo-bulk matrices from each subregion as inputs and then performed co-expression network analysis using the Speakeasy algorithm[16]. We generated 193 modules, with at least 30 genes per module (Table S2). We then conducted module preservation (MP) analysis and revealed that cortical layer modules are more preserved than those derived from WM, blood vessels, or meninges (Fig. 3B; Table S3). Within each subregion, 15~60% of modules are lncRNA-enriched (Fig. 3C). For instance, the cortical module 14 (L2-3_m14) contains 1,261 genes, enriched with 210 lncRNAs (Bonf. $p < 0.05$), including 4 known AD-differential lncRNAs[9,10,17], 3 AD-GWAS genes[18], and 23 AD-differential lncRNA genes based on our ST analysis (Fig. 3D). Gene Ontology (GO) analysis indicates that the L2-3_m14 module is involved in many biological processes (BP), including metabolic and cell cycle processes, ncRNA processing, positive regulation of telomerase, and meiosis I (Fig. 3E, Table S4). In addition, the WM module 11 (WM_m11) contains 1,148 genes, enriched with 156 lncRNAs (Bonf. $p < 0.05$), 2 of which are known AD-differential lncRNAs[9,10,17], 4 are AD-GWAS genes[18] and 29 AD differential based on our ST analysis (Fig. 3F). Gene Ontology (GO) analysis indicates that WM_m11 is involved in carbohydrate derivative metabolism, lipid metabolic modification, and nervous system development (Fig. 3G, Table S4). This result suggests that lncRNAs may exert their roles through interactions with mRNAs within shared regulatory networks.

Spatially differentially expressed (SDE) lncRNAs in postmortem AD brains

To identify spatially differentially expressed (SDE) lncRNAs in postmortem AD brains, we first summed the expression values of all spots within a brain section for each subregion to generate a pseudobulk matrix. We then normalized the pseudobulk estimates across brain sections using TMM-voom[19] and applied linear mixed models (LMM) to contrast AD against NCI individuals. We accounted for age, RNA Integrity Number (RIN), postmortem interval (PMI), and batch as fixed effects and within-subject correlation as a random effect. We declared significance at an α of 0.05 with false discovery rate (FDR) correction for all genes (lncRNA and mRNA) and tested subregions. We found 275 SDE lncRNAs (Table S5), with a greater proportion displaying negative AD association than mRNAs (Fig. 4A-B). Breaking down the SDE lncRNAs by biotypes, we observed enrichment for bidirectional promoter lncRNA in L3-5 and lincRNA in L6 and WM (Fig. 4C). We then compared SDE lncRNAs and mRNAs from our ST analysis with those identified from bulk DLPFC RNA-seq [20] and snRNA-seq on 1,049 and 424 ROSMAP individuals[14], respectively. Despite only a mild overlap in AD DE genes (Fig. 4D), we observed strong correlations of t-values between ST and bulk RNA-seq results for all lncRNAs (R=0.3)

and mRNAs (R=0.39; Fig. 4E), with the remarkable concordance for *LY86-AS1*, *FAM106A*, *LINC01007* and *MIR7-3HG* (downregulated in AD), as well as *NEAT1*, *LINC00632*, *FGD5-AS1* and *LINC01094* (upregulated in AD; Fig. 4E). Similarly, we also detected minimal overlapping between subregion-specific and cell-type-specific AD DE lncRNAs and mRNAs (Fig. 4F). When correlating pairwise t-values across subregions and cell types, we noticed a few exceptions, such as the moderate correlations between L3-5 and inhibitory neurons (Fig. 4H). In addition, we also detected a DE lncRNA alignment between blood vessels and microglia or oligodendrocytes (Fig. 4G), indicating shared lncRNA functions in neurovascular and neuroinflammatory processes. Thus, we detected both consistent and divergent results of AD DE lncRNAs between ST data and other data sets, highlighting the need to explore lncRNA expressions in a range of data layers to capture their potential context-specific roles or responses.

Among AD SDE lncRNAs detected in our study, a greater proportion is subregion-specific (Fig. 5A). We then cross-referenced the snRNA-seq data[14] to examine whether subregion-specific lncRNA expression arises from distinct cell types. However, in most cases, we did not observe a specific cell type driving expression of these subregion-specific lncRNAs (Fig. 5B). On the other side, we detected novel and known SDE lncRNAs[9,10,17] expressed in multiple subregions (Fig. 5C). For previously reported lncRNAs[9,10,17], we also observed an alignment with our bulk RNA-seq analysis (Fig. 5C, bottom panel). Interestingly, some AD SDE lncRNAs show high expression in one subregion (Fig. 2), but their AD effects are predominant in other subregions. One example is *LINC00844*, which displays higher expression in WM/oligodendrocytes (Fig. 2), but its AD effects are predominantly in L2-3 (Fig. 5B). Another is *LINC00507*, which is highly expressed in L2-3, but its AD effects are predominantly in L3-6 (Fig. 5C). In addition, the X-linked gene *XIST* is among SDE lncRNAs, which prompted us to examine the expression of other X-linked genes. Indeed, we found that *MECP2* and *MAOA* are upregulated in AD, while *PGK1* and *HPRT1* are downregulated in AD (Fig. 5D-5F). Applying geneset enrichment analysis (GSEA) to t-values of all genes for each subregion and then focusing on genesets containing AD SDE lncRNAs, we found enrichments for cell motility (*MALAT1*, *SPATA13*), apoptotic process (*ARMCX5-GPRASP2*), chromatin organization (*XIST*, *KCNQ1OT1*) and regulation of defense response (*NEAT1*) (Fig. 5G). Notably, we also detected an enrichment of the "HDAC3 targets dn" geneset that contains 3 AD SDE lncRNAs (*FGD5-AS1*, *NNT-AS1*, and *OIP5-AS1*) (Fig. 5G). In addition, we found that *HDAC1*, *HDAC3*, *HDAC6*, *HDAC7*, and *HDAC10* are spatially differentially expressed in AD brains (Fig. 5H). This finding echoes the accumulated studies showing HDACs' involvement in the AD pathological process[21,22], suggesting a possible functional link between specific lncRNAs and HDAC activity.

Statistically inferred interaction between *OIP5-AS1* and HDAC11 is associated with AD phenotypes

HDAC inhibition has been actively investigated as a potential therapeutic approach in AD[21,23–25]. However, HDAC inhibitors also face challenges due to a lack of selectivity, which often results in unintended off-target effects[26,27]. Our GESA results (Fig. 5D) indicate that HDACs could interact with specific lncRNAs (*FGD5-AS1*, *NNT-AS1*, and *OIP5-AS1*), providing a possible solution for selectively modulating HDAC activity in AD treatment. Among those three lncRNAs, *OIP5-AS1* (*Cyrano*) has been implicated in many developmental processes[28–30], in the onset and progression of various cancers [31,34], and recently in cerebral ischemia-reperfusion injury[35]. Thus, we decided to follow up on *OIP5-AS1* to examine 1) whether genetic variation affects OIP5 AS1 expression and 2) whether OIP5 AS 1 may interact with HDACs and, if so, to what extent the interaction correlates with AD clinical and pathological traits.

We curated a set of expression quantitative trait loci (eQTLs) within 1Mb from the transcription starting site (TSS) of *OIP5-AS1* for each brain cell type (Fig. 6A) from the published snRNA-seq dataset[14] and applied multivariate fine-mapping [36]. Based on the posterior probability of each SNP being causal, we found rs1655558 a potential causal variant influencing OIP5-AS1 expression in multiple cell types, including excitatory neurons, astrocytes, and oligodendrocytes (Fig. 6A). We then investigated the interactions between *OIP5-AS1* expression levels in six major brain cell types and HDAC protein abundance in the DLPFC by incorporating published proteomics data [37]. We associated the interaction between cell-type-specific *OIP5-AS1* expression levels and HDAC protein abundance with three categories of AD traits, i.e., cognitive decline, amyloid plaques, and tau tangles (Fig. 6B). We discovered interaction effects for tau tangles and amyloid plaques, but not cognitive decline (Fig. 6C). Importantly, some interactions show stronger associations with AD traits than the HDAC main effect (more than an order of magnitude lower in p-value, non-circled colored dots in Fig. 6D, 6G), suggesting the interaction with *OIP5-AS1* an essential element for HDACs to impact tangles and amyloid plaques. We repeated this analysis with cell-type-specific HDAC mRNA expression and observed a similar trend in which some interactions show stronger AD associations than main effects, but association strength is lower than its protein counterpart (Fig. 6C; Fig. S1). This finding indicates that *OIP5-AS1* may interact with HDAC proteins rather than mRNAs in affecting AD pathophysiology.

To narrow down which HDAC protein(s) interact with *OIP5 AS1* and in which cell type(s), we adjusted the number of cell types and HDACs in our statistical modeling and found that *OIP5-AS1*x*HDAC11* interaction but not the main effect of HDAC11, is associated with tau pathology in excitatory neurons (Fig. 6D). Notably, the interaction effect is most strongly associated with tangle measurements of

hippocampus and medial temporal lobe (Fig 6E). Given that tau pathology typically progresses from
the hippocampus and medial temporal lobe to other cortical areas, our result indicates that the *OIP5-*
*AS1*xHDAC11 interaction may play a central role in tau progression and early hippocampal
vulnerability. Indeed, when we stratified the 424 ROSMAP individuals in the snRNA-seq data[14] into
three groups based on *OIP5-AS1* expression in excitatory neurons, we found that tau tangles are only
positively associated with HDAC11 protein level within the *OIP5-AS1*-high group (Fig. 6F).

The OIP5-AS1xHDAC11 interaction effect was also detected for amyloid plaques but in microglia, not
excitatory neurons (Fig. 6G). Similarly, the OIP5-AS1xHDAC11 interaction is strongly associated with
plaque measurements in the hippocampus and mesial temporal lobe (Fig. 6H). Stratifying the 424
ROSMAP individuals into three groups based on *OIP5-AS1* expression in microglia, we observed that
amyloid plaques are only positively associated with HDAC11 protein abundance within the OIP5 *AS1-*
low group (Fig. 6I). Overall, our analysis infers that *OIP5-AS1* interacts with HDACs, especially
HDAC11, in a cell-type-specific manner in contributing to AD pathophysiology.

Discussion

Recent advancements in sequencing technologies have revealed that a large fraction of the
mammalian genome is transcribed into lncRNAs[1]. Despite studies showing that brains express more
tissue-specific lncRNAs than most other tissue types[38,39], little is known about the spatial and cellular
context of lncRNAs in the human brain. We mapped the spatial expression of 7,634 lncRNAs in aged
human brains. Overall, we detected lower expression levels and more restricted spatial expression
patterns of lncRNAs than mRNAs (Fig. 1). By cross-referencing published snRNA-seq data, we detect
that some lncRNAs exhibit both sub-regional and cell-type specificity while others are uniquely
restricted to a particular brain subregion or specific cell type (Fig. 2). This partially overlapping but
independent pattern between sub-region and cell-type specificity of lncRNA expression indicates that
lncRNAs may have dual roles, operating within precise spatial regions to support localized functions
while also being tailored to the needs of individual cell types. Indeed, we detected many subregion-
specific lncRNA-enriched gene modules (Fig. 3), supporting their nuanced roles in complex regulatory
networks that involve cross-talks between different cell types.

Comparing the spatial expression of AD SDE lncRNAs to mRNAs, we also observed that lncRNAs are
generally more subregion-specific than mRNAs, with 115 out of 275 AD SDE lncRNAs showing
differential expression specifically in one subregion of the cortex. At least 13 of the AD SDE lncRNAs
detected in our study have been previously reported in AD studies (Fig. 5C, Table S5). For instance,
we detected higher expression of *MALAT1*, *NEAT1*, *BACE1-AS*, and *BDNF-AS* in AD individuals, which

is in line with extensive literature supporting their roles in AD pathogenesis through different pathways. *BACE1-AS* is reported to promote amyloid production[40–42], whereas *BDNF-AS* modulates synaptic plasticity by interacting with their mRNA counterparts[43–45]. As the major organizers of nuclear speckles and paraspeckles, *MALAT1* suppresses neuronal apoptosis and inflammation by modulating *miR-125b* expression[46,47], whereas *NEAT1* regulates tau phosphorylation and Aβ metabolism through interacting with *miR-124* and *miR-107* and histone acetyltransferases[48,49]. Furthermore, our analysis detected emerging immune- and aging-associated lncRNAs, such as *LY86-AS1*[50]*, FAM106A*[51], and *MIR7-3HG*[52].

Increased *XIST* levels have been previously reported in the brains of female AD patients linked to X Chromosome Interaction (XCI), an important factor contributing to female-specific vulnerability in AD[53,54]. We detected an upregulation of *XIST* in AD, primarily in the upper cortical layers (Fig. 5C, 5D). Given that we conducted our study in only females, this upregulation of *XIST* in AD females indicates that the XCI process is disrupted in AD, which may lead to silencing of some X-linked genes or/and overexpression of other genes that escape XCI. Indeed, we detected altered expression of X-linked genes such as *MAOA, MECP2, PGK1,* and *HPRT1*(Fig. 5D, 5E)*,* which have been reported to be essential for neuronal function, epigenetic regulations, and cellular metabolism [55–59].

Our gene set enrichment analysis shows that AD SDE lncRNAs are involved in diverse gene regulation processes, including epigenetic regulation, chromatin remodeling, RNA metabolism, synaptic signaling, and cell apoptosis. One of the mechanistic insights from this analysis is the enrichment of HDAC targets, including *FGD5-AS1*, *NNT-AS1*, and *OIP5-AS1*. HDAC pathways have been widely implicated in AD pathogenesis. Among the HDAC targets, *OIP5-AS1* is particularly interesting due to its diverse regulatory roles in cell biology[60]. First, *OIP5-AS1* has been shown to play a crucial role in cancer by regulating cell proliferation, apoptosis, migration, and invasion, primarily through its ability to sponge miRNAs and interact with chromatin modifiers such as HDACs[31–34,61]. Second, *OIP5-AS1* is upregulated in M2 microglia-derived exosomes, which inhibits neuronal pyroptosis through ubiquitination of its binding protein TXNIP[35]. Third, *OIP5-AS1* acts as a central player of a non-coding RNA network to destroy mature miR-7 and regulate neuronal activity[62]. Using statistical modeling, we inferred that the interaction between *OIP5-AS1* and HDAC proteins, especially HDAC11, is associated with tau tangles in excitatory neurons and plaque burden in microglia (Fig. 6D-I). These results indicate that *OIP5-AS1* may be involved in epigenetic regulations in AD through its interaction with HDACs in a cell-type-specific and context-dependent manner.

Limitations

This study has several data-intrinsic limitations: 1) Our ST samples were collected from aged control and AD female individuals. Therefore, whether our results can generalize to younger age groups or males remains to be determined; 2) Since the 10X Visium platform captures RNA through a polyadenylated (poly(A)) approach, our study misses non-polyadenylated lncRNAs, as well as any lncRNAs with alternative 3' end processing or shorter poly(A) tails; 3) the Visium platform is at the 55-μm resolution, limiting us from analyzing cell-type specific lncRNA expression in the spatial context; 4) the sample size of this study is relatively small, so we are likely underpowered to detect all AD DE lncRNAs; and 5) experiments are needed to validate the functional roles of lncRNAs, such as *OIP5-AS1* in AD pathogenesis.

Conclusion and Future Directions

We report a spatial expression atlas of 7,634 lncRNA genes in aged human brains, demonstrating their relatively restricted spatial expressions. We chart subregion-specific lncRNA changes and identify hundreds of AD DE lncRNAs, including known AD-associated lncRNAs, emerging immune-associated lncRNAs, and novel lncRNAs. These findings open exciting possibilities for future studies with high-resolution spatial profiling platforms, specific enrichment techniques for lncRNAs of interest, and co-detection of protein(s) in the same section, enabling finer delineation of cell-type-specific lncRNA changes in response to different AD pathologies. Furthermore, we explore the potential interaction effects of *OIPS-AS1* and HDACs in mediating AD-associated phenotypes. With various RNA-targeting technologies, such as antisense oligonucleotides (ASOs)[63] and CRISPR-Cas13d[64,65], further functional studies on *OIPS-AS1* and other AD-associated lncRNAs will shed new insights into their potential roles in AD. Because a key limitation of HDAC inhibitors in clinical trials is their non-selective action, *OIP5-AS1* might offer a possible solution for selectively modulating HDAC activity and stratifying patients for precision medicine.

Acknowledgments

We are grateful to those who agreed to donate their brains for research. We thank all the employees at RADC for their support and assistance. This study was supported by NIA grants R01AG074082 and R01AG079223 (to Y.W.), P30AG10161, P30AG72975, R01AG015819, R01AG017917, and U01AG61356 (to D. A. B.), and R01AG061798 (to C.G.).

Author contributions

B.N, D.R.A. K.L. S.T. R.V., and A.I. performed data analysis. S.D.T. and D.R.A. conducted ST experiments. J.X. performed ST data processing. M. F., V. M., and P.L.D. provided snRNA-seq data.

B.N., D.R.A., K.L., and Y.W. wrote the manuscript. D.A.B runs the parent ROSMAP study. Y.W.
conceived and supervised this study. All authors reviewed and approved the final manuscript.

Lead Contact

Further information and requests may be directed to the corresponding author Yanling Wang
(Yanling_Wang@rush.edu)

Materials Availability

This study did not generate new unique reagents.
303

Online Methods

Tissue collection for Spatial Transcriptomics

The human brain tissues were obtained through the ROSMAP (Religious Orders Study and Memory
and Aging Project) cohort studies at the Rush Alzheimer's Disease Center, Rush University Medical
Center, Chicago, IL. Written informed consent was given by the donors for brain autopsy and for the
use of material and clinical data for research purposes, in compliance with national ethical guidelines.
We selected 13 individuals clinically diagnosed as AD with high pathology and 8 controls with no
cognitive impairment (NCI) and no or very little pathology. All individuals are female and matched for
age. We chose all female individuals because there were insufficient males in the ROSMAP cohort
matching our selection criteria (e.g., RIN > 5, age-matched, minimal freezing artifacts in frozen
DLPFC). Frozen brain blocks of the right-hemisphere DLPFC from 21 individuals were cut (~ 1 cm^3),
transferred on dry ice, and stored at -80C.
316

Frozen brain blocks were embedded in OCT and cryosectioned coronally to a thickness of 10 μm
using a Leica CM1950 cryostat set at -17C. Sectioning was done in sets of three (middle section for
ST and two adjacent sections for IF), with 10-50 μm between each set. Sections were attached to
Visium slides and stored at -80C in a slide mailer for up to one week before proceeding with spatial
transcriptomics. ST-adjacent sections were attached to pre-labeled Leica superfrost plus slides and
stored at -80C in slide mailers for up to two weeks before proceeding with immunohistochemistry.
323

Spatial Transcriptomics

Spatial Transcriptomics was performed according to the Visium Spatial Gene Expression User Guide
(10X Genomics; CG000239). Briefly, cryosectioned tissue on Visium slides was transferred from -80C
to the lab on dry ice and quickly thawed for 1 min on a heat block at 37C, then immediately
transferred to methanol pre-equilibrated to -20C and fixed at -20C for 30 min. Sections were then
stained with hematoxylin and eosin, and brightfield images were acquired with a Nikon Ti2 with NIS-
Elements AR 5.11.01 64-bit software. After imaging, tissues were immediately permeabilized in 0.1%
pepsin diluted in 0.1 M HCl for 18 min at 37C. Reverse transcription, second-strand synthesis, cDNA
amplification (15 cycles), and library construction were performed according to the manufacturer's
instructions. For the first four individuals (#2, 8, 12, and 18), protocol version 'Rev A' was followed,
and the sample index PCR was 12 cycles. For the remaining samples, protocol version 'Rev D' was

followed, and the sample index PCR was 13-14 cycles. Library concentration was quantified by Qubit 1x dsDNA HS using a SpectraMax M3 plate reader. Library concentration and size distribution were assessed by Fragment Analyzer (Agilent). Sequencing was performed in three batches on an Illumina NovaSeq, targeting a minimum of 50,000 raw reads per spot.

Spatial transcriptomics sequencing data processing

Sequencing data were pre-processed with Space Ranger pipeline (10x Genomics), which mapped the barcodes against the human genome (Gencode 27), generated FASTQ files, and generated a count matrix for each section. The individual files were then merged in a Seurat object (Seurat v4.2.0) and were processed in R. We calculated per-spot quality metrics and spots with low-measured genes (< 500) were removed. The Seurat::SCTransform function was used to normalize and scale the UMI counts based on regularized negative binomial regression. Principal component analysis (PCA) was performed on the top 3000 variable genes. Using the top 50 PCs (Supplementary Fig. 1Fig. S1D) and accounting for technical and biological covariates such as age, RNA integrity number (RIN), library batch, and donor, we integrated the datasets using the Harmony package (v0.1.0). We explored various tools for data integration and chose Harmony for its high performance, usability, and scalability. We only used Harmony integration to identify clusters corresponding to brain regions. All downstream analyses (differential gene expression, cell proportion estimates, and ligand-receptor analysis) were performed on pre-integration count data. After visualizing the standard deviation attributed to each Harmony embedding as an elbow plot (Supplementary Fig. 1Fig. S1E), we decided to use the top 10 Harmony embeddings. We applied Seurat::runUMAP to compute UMAP (uniform manifold approximation and projection) dimension reduction and Seurat::findNeighbors and Seurat::findClusters to identify clusters based on a shared nearest neighbor (SNN) clustering algorithm (Louvain clustering with resolution set to 0.3; all other parameters default). This resulted in 11 clusters, which were annotated based on their canonical markers and spatialLIBD (v7.5) to the closest anatomical cortical layer (L1, L2-3, L3-5, L5, L6), white matter (WM), meninges, and specific cellular subpopulations, namely cells with high expression of hemoglobin genes ('Blood Vessel') and SST/NPY ('Interneuron'). The last identified cluster, which carried a glial signature, had very few spots and so was not included in downstream analyses.

Co-expression network analysis in spatial transcriptomics

The modules of co-expressed genes were derived using the Speakeasy (SE) algorithm v1, Matlab version[16] . We created pseudo-bulk gene expression by summing the counts per donor, then splitting by UMAP cluster. Lowly expressed genes (less than 1 count per million, CPM) were removed, and the variancePartition R function was applied to capture the major sources of variation, as previously described [15]. The input for SE consisted of the residuals after normalization using tmm.voom and adjustment for library batch, RIN, age, and PMI. We built the modules separately for each UMAP cluster, using the results from 100 initializations of SE and Spearman correlation. This resulted in 193 modules, each with at least 30 genes (L1 = 30, L2-3 = 21, L3-5 = 20, L5 = 20, L6 = 20, Meninges = 33, WM = 20, and blood vessels = 29). The module assignments are provided in Table S2.

Functional Enrichment Analysis (FEA)

Each module was annotated using the gprofiler software (v2_0.2.1) via the g:GOSt R function [66]. This
tool maps genes to functional information sources and detects statistically significantly enriched
terms. These sources include the GO database, pathways from KEGG, Reactome, and
WikiPathways, miRNA targets from miRTarBase, regulatory motif matches from TRANSFAC, tissue
specificity from the Human Protein Atlas, protein complexes from CORUM, and human disease
phenotypes from the Human Phenotype Ontology. For multiple testing correction, we used the default
g:SCS method unless otherwise specified. The FEA results are available in Table S4.
384
Module Preservation (MP) Analysis
To assess whether the modules were preserved across distinct UMAP clusters, we used the MP
function from the WGCNA R package (v1.70-3)[67]. Pairwise comparisons were performed for all 193
modules. We then estimated the mean and variance of the preservation statistic under the null
hypothesis that there is no relationship between module assignments in the reference and test
datasets. Fifteen distinct preservation statistics were calculated (e.g., module density, mean
expression, connectivity), each standardized by its mean and variance to define a Z statistic for each
preservation measure. Under these assumptions, each Z statistic follows a normal distribution if the
module is not preserved. To compare preservation across modules, we calculated a composite
preservation measure using the following equation:
395
396
$$Zsummary = \frac{Zdensity + Zconnectivity}{2}$$
397
398   Modules with Z < 2 were considered not preserved, those with Z between 2 and 10 were considered
399   preserved, and modules with Z > 10 were considered highly preserved. Zsummary stats are available
400   in Table S3.
401
402   **Statistical analysis to identify AD-differential genes**
403
For each brain section of a subject, we summed the UMI counts of all spots belonging to a subregion
to generate a pseudobulk estimate for each gene. We only kept expressed genes with counts per
million (CPM) > 1 in at least 80% of the brain sections. With the pseudobulk estimates, we estimated
the library sizes of the brain sections using the trimmed mean of M-values (TMM) of edgeR[68]
(edgeR_3.36.0) and transformed the pseudobulk estimates into log_2 CPM using voom-limma
(limma_3.50.1). For each subregion, we excluded brain sections with <50 spots. To contrast the
normalized pseudobulk expression between AD and controls, we applied linear mixed models (LMM),
accounting for age, PMI, RIN, and batch as fixed effects and modeled within-subject correlation with a
random effect. We declared significance at an α of 0.05 with false discovery rate (FDR) correction
across all genes and all subregions.
414
415   For interpretation, we applied GSEA[69] with AD differential t-values of each subregion as scores, and
416   GO terms and canonical pathways as genesets. To characterize lncRNAs with respect to brain cell
417   types, we used the pseudobulk expression data from a published snRNAseq dataset of 424
418   ROSMAP individuals [14]. For each of the major cell types, we contrasted pseudobulk expression of AD
419   subjects vs. controls using multiple regression, accounting for age, PMI, RIN, and batch. For further

comparison, we repeated this AD differential expression analysis with bulk DLPFC RNA-seq data from 1049 individuals[20].

References

1. Mattick, J. S. *et al.* Long non-coding RNAs: definitions, functions, challenges and recommendations. *Nat. Rev. Mol. Cell Biol.* **24**, 430–447 (2023).
2. Statello, L., Guo, C.-J., Chen, L.-L. & Huarte, M. Gene regulation by long non-coding RNAs and its biological functions. *Nat. Rev. Mol. Cell Biol.* **22**, 96–118 (2020).
3. Wang, K. C. & Chang, H. Y. Molecular mechanisms of long noncoding RNAs. *Mol. Cell* **43**, 904–914 (2011).
4. Guttman, M. *et al.* Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature* **458**, 223–227 (2009).
5. Kung, J. T. Y., Colognori, D. & Lee, J. T. Long noncoding RNAs: past, present, and future. *Genetics* **193**, 651–669 (2013).
6. Djebali, S. *et al.* Landscape of transcription in human cells. *Nature* **489**, 101–108 (2012).
7. Cabili, M. N. *et al.* Integrative annotation of human large intergenic noncoding RNAs reveals global properties and specific subclasses. *Genes Dev.* **25**, 1915–1927 (2011).
8. Derrien, T. *et al.* The GENCODE v7 catalog of human long noncoding RNAs: analysis of their gene structure, evolution, and expression. *Genome Res.* **22**, 1775–1789 (2012).
9. Wu, X., Xia, P., Yang, L., Lu, C. & Lu, Z. The roles of long non-coding RNAs in Alzheimer's disease diagnosis, treatment, and their involvement in Alzheimer's disease immune responses. *Noncoding RNA Res.* **9**, 659–666 (2024).
10. Policarpo, R. & d'Ydewalle, C. Missing lnc(RNAs) in Alzheimer's Disease? *Genes* **13**, (2021).
11. Lan, Z., Chen, Y., Jin, J., Xu, Y. & Zhu, X. Long non-coding RNA: Insight into mechanisms of Alzheimer's disease. *Front. Mol. Neurosci.* **14**, 821002 (2021).
12. Huang, Z., Zhao, J., Wang, W., Zhou, J. & Zhang, J. Depletion of LncRNA NEAT1 rescues mitochondrial dysfunction through NEDD4L-dependent PINK1 degradation in animal models of Alzheimer's disease. *Front. Cell. Neurosci.* **14**, 28 (2020).
13. Riva, P., Ratti, A. & Venturin, M. The Long Non-Coding RNAs in Neurodegenerative Diseases: Novel Mechanisms of Pathogenesis. *Curr. Alzheimer Res.* **13**, 1219–1231 (2016).
14. Fujita, M. *et al.* Cell subtype-specific effects of genetic variation in the Alzheimer's disease brain. *Nat. Genet.* **56**, 605–614 (2024).
15. Avey, D. R. *et al.* Uncovering plaque-Glia niches in human Alzheimer's Disease brains using spatial transcriptomics. *bioRxivorg* (2024) doi:10.1101/2024.09.05.611566.
16. Gaiteri, C. *et al.* Identifying robust communities and multi-community nodes by combining top-down and bottom-up approaches to clustering. *Sci. Rep.* **5**, 1–14 (2015).
17. Li, D. *et al.* Insights into lncRNAs in Alzheimer's disease mechanisms. *RNA Biol.* **18**, 1037–1047 (2021).
18. Bellenguez, C. *et al.* New insights into the genetic etiology of Alzheimer's disease and related dementias. *Nat. Genet.* **54**, 412–436 (2022).
19. Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: Precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biol.* **15**, R29 (2014).
20. Mostafavi, S. *et al.* A molecular network of the aging human brain provides insights into the pathology and cognitive decline of Alzheimer's disease. *Nat. Neurosci.* **21**, 811–819 (2018).

21. Shukla, S. & Tekwani, B. L. Histone deacetylases inhibitors in neurodegenerative diseases, neuroprotection and neuronal differentiation. *Front. Pharmacol.* **11**, 537 (2020).

22. Govindarajan, N. *et al.* Reducing HDAC6 ameliorates cognitive deficits in a mouse model for Alzheimer's disease. *EMBO Mol. Med.* **5**, 52–63 (2013).

23. Onishi, T. *et al.* A novel orally active HDAC6 inhibitor T-518 shows a therapeutic potential for Alzheimer's disease and tauopathy in mice. *Sci. Rep.* **11**, 15423 (2021).

24. Yang, S.-S., Zhang, R., Wang, G. & Zhang, Y.-F. The development prospection of HDAC inhibitors as a potential therapeutic direction in Alzheimer's disease. *Transl. Neurodegener.* **6**, 19 (2017).

25. Zhang, M. *et al.* Histone deacetylase inhibitors VPA and WT161 ameliorate the pathological features and cognitive impairments of the APP/PS1 Alzheimer's disease mouse model by regulating the expression of APP secretases. *Alzheimers. Res. Ther.* **16**, 15 (2024).

26. Kumar, V., Kundu, S., Singh, A. & Singh, S. Understanding the role of histone deacetylase and their inhibitors in neurodegenerative disorders: Current targets and future perspective. *Curr. Neuropharmacol.* **20**, 158–178 (2022).

27. Ru, J. *et al.* Technologies of targeting histone deacetylase in drug discovery: Current progress and emerging prospects. *Eur. J. Med. Chem.* **261**, 115800 (2023).

28. Yang, J.-H. *et al.* LncRNA OIP5-AS1-directed miR-7 degradation promotes MYMX production during human myogenesis. *Nucleic Acids Res.* **50**, 7115–7133 (2022).

29. Zhuang, A. *et al.* Loss of the long non-coding RNA OIP5-AS1 exacerbates heart failure in a sex-specific manner. *iScience* **24**, 102537 (2021).

30. Hunkler, H. J. *et al.* The long non-coding RNA Cyrano is dispensable for pluripotency of Murine and human pluripotent stem cells. *Stem Cell Reports* **15**, 13–21 (2020).

31. Kotake, Y., Matsunaga, N., Wakasaki, T. & Okada, R. OIP5-AS1 promotes proliferation of non-small-cell lung cancer and head and neck squamous cell carcinoma cells. *Cancer Genomics Proteomics* **18**, 543–548 (2021).

32. Wu, L., Liu, Y., Guo, C. & Shao, Y. LncRNA OIP5-AS1 promotes the malignancy of pancreatic ductal adenocarcinoma via regulating miR-429/FOXD1/ERK pathway. *Cancer Cell Int.* **20**, 296 (2020).

33. Arunkumar, G. *et al.* LncRNA OIP5-AS1 is overexpressed in undifferentiated oral tumors and integrated analysis identifies as a downstream effector of stemness-associated transcription factors. *Sci. Rep.* **8**, 7018 (2018).

34. Gong, M. *et al.* OIP5 is a novel prognostic biomarker in clear cell renal cell cancer correlating with immune infiltrates. *Front. Immunol.* **13**, 805552 (2022).

35. Li, Z. *et al.* Exosomal OIP5-AS1 attenuates cerebral ischemia-reperfusion injury by negatively regulating TXNIP protein stability and inhibiting neuronal pyroptosis. *Int. Immunopharmacol.* **127**, 111310 (2024).

36. Zou, Y., Carbonetto, P., Xie, D., Wang, G. & Stephens, M. Fast and flexible joint fine-mapping of multiple traits via the Sum of Single Effects model. *bioRxivorg* (2024) doi:10.1101/2023.04.14.536893.

37. Johnson, E. C. B. *et al.* Large-scale deep multi-layer analysis of Alzheimer's disease brain reveals strong proteomic disease-related changes not observed at the RNA level. *Nat. Neurosci.* **25**, 213–225 (2022).

38. Jiang, C. *et al.* Identifying and functionally characterizing tissue-specific and ubiquitously expressed human lncRNAs. *Oncotarget* **7**, 7120–7133 (2016).

39. de Goede, O. M. *et al.* Population-scale tissue transcriptomics maps long non-coding RNAs to complex disease. *Cell* **184**, 2633–2648.e19 (2021).

40. Li, F. *et al.* The effect of BACE1-AS on β-amyloid generation by regulating BACE1 mRNA expression. *BMC Mol. Biol.* **20**, 23 (2019).

41. Faghihi, M. A. *et al.* Expression of a noncoding RNA is elevated in Alzheimer's disease and drives rapid feed-forward regulation of beta-secretase. *Nat. Med.* **14**, 723–730 (2008).

42. Zhang, W., Zhao, H., Wu, Q., Xu, W. & Xia, M. Knockdown of BACE1-AS by siRNA improves memory and learning behaviors in Alzheimer's disease animal model. *Exp. Ther. Med.* **16**, 2080–2086 (2018).

43. Modarresi, F. *et al.* Inhibition of natural antisense transcripts in vivo results in gene-specific transcriptional upregulation. *Nat. Biotechnol.* **30**, 453–459 (2012).

44. Jiao, S.-S. *et al.* Brain-derived neurotrophic factor protects against tau-related neurodegeneration of Alzheimer's disease. *Transl. Psychiatry* **6**, e907 (2016).

45. Ding, Y., Luan, W., Shen, X., Wang, Z. & Cao, Y. LncRNA BDNF-AS as ceRNA regulates the miR-9-5p/BACE1 pathway affecting neurotoxicity in Alzheimer's disease. *Arch. Gerontol. Geriatr.* **99**, 104614 (2022).

46. Ma, P. *et al.* Long non-coding RNA MALAT1 inhibits neuron apoptosis and neuroinflammation while stimulates neurite outgrowth and its correlation with MiR-125b mediates PTGS2, CDK5 and FOXQ1 in Alzheimer's disease. *Curr. Alzheimer Res.* **16**, 596–612 (2019).

47. Bernard, D. *et al.* A long nuclear-retained non-coding RNA regulates synaptogenesis by modulating gene expression. *EMBO J.* **29**, 3082–3093 (2010).

48. Dong, L.-X. *et al.* LncRNA NEAT1 promotes Alzheimer's disease by down regulating micro-27a-3p. *Am. J. Transl. Res.* **13**, 8885–8896 (2021).

49. Ke, S. *et al.* Long noncoding RNA NEAT1 aggravates Aβ-induced neuronal damage by targeting miR-107 in Alzheimer's disease. *Yonsei Med. J.* **60**, 640–650 (2019).

50. Bútová, R. *et al.* LncRNAs LY86-AS1 and VIM-AS1 distinguish plasma cell leukemia patients from multiple myeloma patients. *Biomedicines* **9**, 1637 (2021).

51. Kurniansyah, N. *et al.* An integrated multi-omics analysis of sleep-disordered breathing traits implicates P2XR4 purinergic signaling. *Commun. Biol.* **6**, 125 (2023).

52. Tan, Y. *et al.* lncRNA-associated ceRNA network revealing the potential regulatory roles of ferroptosis and immune infiltration in Alzheimer's disease. *Front. Aging Neurosci.* **15**, 1105690 (2023).

53. Bajic, V. P. *et al.* The X files: 'the mystery of X chromosome instability in Alzheimer's disease'. *Front. Genet.* **10**, 1368 (2019).

54. Garcia, A. X., Xu, J., Cheng, F., Ruppin, E. & Schäffer, A. A. Altered gene expression in excitatory neurons is associated with Alzheimer's disease and its higher incidence in women. *Alzheimers Dement. (N. Y.)* **9**, e12373 (2023).

55. Sun, X. *et al.* The MAOA gene influences the neural response to psychosocial stress in the human brain. *Front. Behav. Neurosci.* **14**, 65 (2020).

56. Ziermans, T. *et al.* Working memory brain activity and capacity link MAOA polymorphism to aggressive behavior during development. *Transl. Psychiatry* **2**, e85 (2012).

57. Chua, G. N. L. *et al.* Differential dynamics specify MeCP2 function at nucleosomes and methylated DNA. *Nat. Struct. Mol. Biol.* 1–9 (2024).

58. Chu, S.-L. *et al.* Phosphoglycerate kinase 1 acts as a cargo adaptor to promote EGFR transport to the lysosome. *Nat. Commun.* **15**, 1021 (2024).

59. Vinokurov, A. Y. *et al.* HPRT1 deficiency induces alteration of mitochondrial energy metabolism
in the brain. *Mol. Neurobiol.* **60**, 3147–3157 (2023).

60. Wooten, S. & Smith, K. N. Long non-coding RNA OIP5-AS1 (Cyrano): A context-specific
regulator of normal and disease processes. *Clin. Transl. Med.* **12**, e706 (2022).

61. Mei, J. *et al.* OIP5-AS1 modulates epigenetic regulator HDAC7 to enhance non-small cell lung
cancer metastasis via miR-140-5p. *Oncol. Lett.* **20**, 7 (2020).

62. Kleaveland, B., Shi, C. Y., Stefano, J. & Bartel, D. P. A network of noncoding regulatory RNAs
acts in the mammalian brain. *Cell* **174**, 350–362.e17 (2018).

63. Mummery, C. J. *et al.* Tau-targeting antisense oligonucleotide MAPTRx in mild Alzheimer's
disease: a phase 1b, randomized, placebo-controlled trial. *Nat. Med.* **29**, 1437–1447 (2023).

64. Kuo, H.-C., Prupes, J., Chou, C.-W. & Finkelstein, I. J. Massively parallel profiling of RNA-
targeting CRISPR-Cas13d. *Nat Commun* **15**, (2024).

65. Wessels, H.-H. *et al.* Massively parallel Cas13 screens reveal principles for guide RNA design.
Nat. Biotechnol. **38**, 722–727 (2020).

66. Kolberg, L. *et al.* g:Profiler-interoperable web service for functional enrichment analysis and gene
identifier mapping (2023 update). *Nucleic Acids Res.* **51**, W207–W212 (2023).

67. Langfelder, P., Luo, R., Oldham, M. C. & Horvath, S. Is my network module preserved and
reproducible? *PLoS Comput. Biol.* **7**, e1001057 (2011).

68. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

69. Subramanian, A. *et al.* Gene set enrichment analysis: A knowledge-based approach for
interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences*
102, 15545–15550 (2005).

Main Figures

Figure 1. ST reveals the spatial expression of lncRNAs. (A) Study overview. **(B)** Comparison of the expression distributions of lncRNAs and protein-coding genes (mRNAs). **(C)** Comparison of percentage of ST spots with non-zero expression for lncRNAs and mRNAs. **(D)** Proportion of detected lncRNAs (aggregated across subregions) belonging to each biotype. **(E)** Upset plots showing the numbers of lncRNAs (left) and mRNAs (right) detected in the top 20 most common intersections of subregions.

Figure 2. Identification of subregion- and cell-type-enriched lncRNAs. (A) Heatmap showing scaled expression (z-score) of the top 50 subregion-specific lncRNAs. An additional column to the right shows the percentage of spots with non-zero expression for each lncRNA (*exemplar lncRNAs shown in panel B). **(B)** UMAP and expression feature plots for 4 representative sections, showing 5 examples of lncRNAs with enriched expression in specific cortical layers or white matter (WM). **(C)** Scaled snRNA-seq expression of the same 50 subregion-specific lncRNAs shown in (A) for the 6 most abundant brain cell types (*exemplar lncRNAs shown in panel B; n.d.: not detected). **(D)** Violin plots showing expression distributions among 6 major brain cell types for the 5 lncRNAs shown in (B). **(E)** Scaled expression by cell type (top) and ST subregion (bottom) for the top 10 cell-type-specific lncRNAs from each of the 6 major brain cell types.

Figure 3. Subregion-specific gene modules from ST data highlight a central role for lncRNAs.
(A) Schematic of gene expression module generation and downstream analyses. **(B)** Heatmap showing the percentage of preserved modules between each pairwise comparison of UMAP clusters. **(C)** Stacked barplot shows the number of modules in each subregion. Red colors for the lncRNA-enriched modules. **(D)** Network connectivity map for L2-3 m14, illustrating high connectivity of lncRNAs (correlation > 0.9). **(E)** Enriched Gene Ontology (GO) terms (FDR at 0.05) from functional enrichment analysis of an example lncRNA-enriched module (L2-3 m14). **(F)** Network connectivity map for WM module 11, illustrating high connectivity of lncRNAs (correlation > 0.95). **(G) Enriched** GO terms (FDR at 0.05) from functional enrichment analysis for the WM module 11. **(E, G)** The GO biological process terms were collapsed, keeping only the top GO parent functions to avoid redundancy. The y-axis and bar size represent the significance in -log$_{10}$(Adj. p-Value). GO biological process (BP), cellular component (CC), and GO molecular function (MF).

632
633
634

Figure 4. Contrasting AD-differential lncRNAs and mRNAs across data modalities
(A) Histogram showing the distribution of AD-differential t-values for mRNAs and lncRNAs. **(B)** Total numbers of
AD-differential lncRNAs and mRNAs for each subregion. **(C)** Heatmap showing the enrichment of AD-differential
lncRNAs for specific biotypes. **(D)** Overlapsin AD-differential lncRNAs and mRNAs between ST and bulk RNA-
seq. **(E)** Scatterplots of ST (y-axis) vs. bulk RNA-seq (x-axis) AD-differential t-values for lncRNAs (left) and
mRNAs (right). Correlation (R) value is shown at the top left of each plot. **(F)** Overlaps in AD-differential lncRNAs
and mRNAs between ST and snRNA-seq. **(G)** Heatmap showing correlation (R) between subregions-cell-types
based on their AD differential t-values(*p<0.05). **(H)** Scatterplot of ST L3-5 (y-axis) vs. snRNA-seq inhibitory
neuron (x-axis) AD-differential t-values (restricted tolncRNAs).

Figure 5. Characterization of AD-differential lncRNAs by ST. (A) The percentage of DEGs in each ST subregion, split by lncRNA/mRNA, up/down in AD, and whether they are subregion-specific (ss) or not (nss). Fisher's exact test was applied to assess differences in subregion-specificity between lncRNA and mRNA with p-values indicated. **(B)** (top) Heatmap showing t-values of AD-differential lncRNAs passing FDR correction specifically for one ST subregion, i.e. subregion-specific (*p<5e-4 in designated subregion and p>0.001 in all others). (bottom) AD-differential t-values for 6 brain cell types. An additional row below shows the percentage of ST spots with non-zero expression for each lncRNA. **(C)** (left) Heatmap showing t-values of AD-differential lncRNAs passing FDR correction in two or more ST subregions, i.e. not subregion-specific (*p<5e-4). An additional row below shows the percentage of ST spots with non-zero expression for each lncRNA. (right) AD-differential t-values for a subset of 13 lncRNAs with previously reported AD association, including t-values from DLPFC bulk RNA-seq below. **(D)** Boxplots of subregion-specific pseudobulk expression from ST data for XIST, MECP2, and PGK1 (X-linked genes), all of which show altered expression in AD in one or more subregions. **(E)** Heatmap showing a subset of X-linked genes with AD-differential expression in one or more subregions (*p<5e-4). **(F)** XIST spatial expression plots for two control (Ctrl) and two AD sections. **(G)** Sankey diagram and heatmap showing a subset of enriched genesets and AD differential lncRNAs belonging to those genesets. **(H)** A subset of HDAC genes display AD-differential expression in specific ST subregions (*p<5e-4).

661
662
Figure 6. Genetic analysis of OIP5-AS1 and its interaction with HDAC. (A) Colocalization analysis of OIP5-
AS1 eQTLs across major brain cell types. Applying mvSuSiE found rs1655558 to be a potential driver of OIP5-
AS1 expression in astrocytes, oligodendrocytes, and excitatory neurons (Ast: Astrocyte, Exc: excitatory, Inh:
Inhibitory, Mic: Microglia, Oli: Oligodendrocyte, OPC: Oligodendrocyte progenitor cell, PIP: posterior inclusion
probability; cartoon cell type images created in BioRender.com). **(B)** Summary of OIP5-AS1xHDAC interaction
analysis. snRNAseq (Masashi et al., 2024) and proteomic (Wingo et al., 2024) data from human DLPFC were
used to associate the statistical interaction between OIP5-AS1 expression level (in specific cell types) and
protein abundance of each of six HDACs with AD phenotypes, including cognitive decline (cogdx), amyloid
plaques, and tau tangles. Cognition was assessed over multiple domains and pathologies were measured in
multiple brain regions. **(C)** Each point in the violin plot represents the −log10(p) of the interaction effect of
OIP5-AS1 and HDAC for a given combination of cell type, HDAC, and AD phenotype. Higher associations
were observed for plaques and tangles. **(D)** Scatterplots showing t-values of the HDAC main effect (x-axis) and
the OIP5-AS1xHDAC interaction for tangles measured in multiple brain regions. Dots are colored but not
circled if interaction p<0.05 and an order of magnitude lower than the main effect p-value. Circled dots are
interactions passing multiple comparison correction for the number of HDACs and cell types. OIP5-
AS1xHDAC11 interaction shows the strongest association with tangles. (E) Heatmap of OIP5-AS1xHDAC11
interaction p-values across cell types and tangles in different brain regions shown (ag: anterior gyrus, ec:

entorhinal cortex, hip: hippocampus, mf: medial frontal, mt: medial temporal, cg: cingulate gyrus, it: inferior temporal, mes_t: mesial temporal). **(F)** An exemplary visualization of interaction effect. Individuals were stratified into three groups based on the expression of OIP5-AS1 in excitatory neurons. For each group, HDAC11 protein abundance was plotted against tangle level in the hippocampus. (G-I) are plaque burden counterparts of (D-F).

Supplementary Tables.

Table S1. Subregion- and cell-type-enriched lncRNAs

Table S2. ST gene expression module assignments

Table S3. ST gene expression module preservation

Table S4. ST gene expression module FEA

Table S5. Differentially expressed lncRNAs in AD (ST, bulk, snRNA)

Table S6. GSEA of ST genes

Supplementary Figures.

Figure S1. Association of lncRNA OIP5-AS1 with AD phenotypes using HDAC