Genome-wide consensus transcriptional signatures identify synaptic pruning

linking Alzheimer's disease and epilepsy

Huihong Li[1,2,3,4], Zhiran Xie[4], Yuxuan Tian[4], Yaxi Yang[4], Bingying Lin[1,2,3,5], Si

Chen[1,2,3,5], Jianwei Li[1,2,3,5], Jie Wu[1,2,3], Mingjie Chen[6], Xueke Liu[6], Yushan Sun[6],

Zihan Deng[1,2,3,5], Mengqi Hong[1,2,3,5], Bing Huang[1,2,3,5], Naili Wei[1], Xiaoyu Ji[1,2,3,5*]

[1]Department of Neurosurgery, The First Affiliated Hospital of Shantou University

Medical College, Shantou, Guangdong 515041, China

[2]Brain Function and Disease Laboratory, Shantou University Medical College,

Shantou, Guangdong 515041, China

[3]Department of Pharmacology, Shantou University Medical College, Shantou,

Guangdong 515041, China

[4]Program in Computational Biology and Biomedical Informatics, Yale University,

New Haven, Connecticut, United States of America

[5]Pain and Related Diseases Research Laboratory, Shantou University Medical College,

Shantou, Guangdong 515041, China

[6]Department of Biotechnology and Food Engineering, Guangdong Technion – Israel

Institute of Technology, Shantou, Guangdong 515063 , China

[*]**Corresponding authors**

Contact: Dr. Xiaoyu Ji, xyji@stu.edu.cn

Abstract

Alzheimer's disease (AD) and epilepsy (EP) share a complex bidirectional

relationship, yet the molecular mechanisms underlying their comorbidity remain

insufficiently explored. To identify potential consensus transcriptional programs

across animal models and human patients with AD and EP, we conducted

comprehensive genome-wide transcriptomic analyses of multiple datasets. Our

investigation included mouse models of temporal lobe epilepsy (pilocarpine- and

kainic acid-induced; n = 280), mouse models of AD (7 transgenic models expressing

human tau or amyloid pathology; n = 257), and performed cross-species validation in

human cohorts (EP: n = 182; AD: n = 301). Our unsupervised gene co-expression

analysis revealed a highly conserved immune-related module across all models and

patient cohorts. The hub consensus signatures of this module were centered around a

microglial synaptic pruning pathway involving *TYROBP*, *TREM2*, and *C1Q*

complement components. Gene regulatory network analysis identified *TYROBP* as the

key upstream hub signature. These hub consensus signatures showed consistent

upregulation in both AD and EP human bulk and single-nucleus datasets, preserved

their regulatory relationship across species, and demonstrated strong diagnostic value.

Computational modeling further demonstrates the asymmetric sensitivity of synaptic

pruning to neural network dynamics. Loss of inhibitory synapses has a

disproportionately greater impact on the excitation/inhibition (E/I) balance, shifting it

toward hyperexcitation and increasing neural network synchronization. Our findings

support a mechanistic hypothesis that identifies microglial complement-mediated

synaptic pruning as a conserved central pathway linking neurodegeneration to

epileptogenesis and suggested that targeting this pathway may offer therapeutic

benefits for AD and EP comorbidity.

Keywords: Alzheimer's disease, epilepsy, synaptic pruning, computational

neuroscience, modeling, transcriptome, *TYROBP, TREM2, C1Q*

Graphic abstract

52
53

Highlights

- Genome wide transcriptomic analysis across epilepsy (EP) and Alzheimer's disease (AD) models and patients identified a conserved immune module.

- *TYROBP-TREM2-C1Q* microglial synaptic pruning pathway was identified as central consensus signature across model and patient datasets.

- Consensus transcriptional signatures showed strong diagnostic value in both AD and EP patient cohorts.

- Computational modeling demonstrates the asymmetric effects of synaptic pruning, whereby loss of inhibitory synapses disproportionately disrupts the E/I balance, leading to hyperexcitability and increased network synchronization.

1. Background

Alzheimer's disease (AD) and epilepsy (EP) are two common neurological diseases with a complex bidirectional relationship that contributes to their respective pathogenesis [1–3]. AD, characterized by progressive cognitive decline and neurodegeneration, is the leading cause of dementia worldwide, affecting an estimated 55 million individuals. This number is projected to triple by 2050 as life expectancy increases [4]. EP, characterized by recurrent unprovoked seizures, affects over 50 million individuals worldwide with a particularly high incidence among elderly populations, especially those with neurodegenerative conditions such as AD [5,6].

Epidemiological studies demonstrate a strong association between AD and EP. AD patients show increased susceptibility to EP, with 10–22% experiencing unprovoked seizures, while 22–54% demonstrating subclinical epileptiform activity during disease progression [7,8]. These seizures, predominantly occurring in the temporal lobe, correlate with accelerated cognitive decline and enhanced AD pathology [9]. Conversely, EP patients exhibit a two-fold higher risk of developing dementia compared to controls, with late-onset EP patients showing a three-fold increased risk [10,11]. This elevated risk is particularly evident in temporal lobe epilepsy (TLE), where numerous pathological features overlap with AD [9].

The pathological overlap between AD and EP extends across multiple biological

levels [12]. At the molecular and cellular levels, both conditions feature amyloid

deposition, tau accumulation, and hippocampal sclerosis—particularly evident TLE

[13–15]. Chronic seizures increase neuronal activity and metabolic demands, potentially

accelerating the production and aggregation of amyloid-β (Aβ) and

hyperphosphorylated tau protein [16,17]. Conversely, the progressive accumulation of

these pathological proteins in AD appears to lower seizure thresholds, promoting

hyperexcitability and epileptiform activity [18,19]. At the network level, both disorders

demonstrate profound disruption of the excitation/inhibition (E/I) balance, with a shift

toward hyperexcitability driving both cognitive impairment and seizure susceptibility.

This network dysfunction is accompanied by impaired synaptic plasticity, altered

synchronization patterns, and progressive circuit reorganization [20–22].

Despite these established clinical and pathological connections, the precise molecular

mechanisms governing the AD-EP relationship remain insufficiently explored. While

previous studies have identified various potential contributors—including

dysregulated protein homeostasis, mitochondrial dysfunction, and chronic

inflammatory responses—most investigations have examined these conditions in

isolation rather than specifically addressing their comorbidity [23–25]. Current research

approaches have been limited by several factors: 1) inadequate integration of

transcriptomic data across multiple disease models and human cohorts; 2) insufficient

attention to shared cellular processes that bridge molecular alterations with network

dysfunction; and 3) limited application of computational modeling to connect

molecular findings with clinically relevant physiological outcomes.

The challenge of treating EP in AD patients further underscores the need for deeper

mechanistic insights. Clinicians face complex decisions regarding seizure treatment in

AD patients, including whether to treat subclinical epileptiform activity and which

antiseizure medications minimize cognitive side effects [26]. Emerging therapeutic

approaches require more precise understanding of the shared pathological

mechanisms. Systems biology and computational neuroscience approaches offer

powerful tools to integrate insights from both animal models and human patients,

potentially revealing actionable therapeutic targets that address both conditions

simultaneously.

Emerging evidence suggests that AD and EP may share underlying molecular

mechanisms. In this study, we investigated the hypothesis that consensus

transcriptional programs exist across animal models and human patients with AD and

EP. To test this, 1) we performed a genome-wide transcriptomic analysis of large-scale

datasets from well-established mouse models of TLE (pilocarpine- and kainic

acid-induced; n = 280) and AD (multiple transgenic models expressing human tau

[TAU, rTg4510] or amyloid pathology [J20, TAS10, TPM, HO-TASTPM,

HET-TASTPM]; n = 257), aiming to identify robust, consensus transcriptional

modules and signatures across distinct disease modeling paradigms, 2) cross-species

validation of the diagnostic value of consensus transcriptional signatures and

assessment of module preservation in human patient cohorts (EP cohorts: n = 182; AD

cohorts: n = 301), 3) developed computational models to link molecular insights to

network mechanisms of epileptogenesis. Our multi-cohort, cross-species analysis

indicated strong, consensus transcriptional signatures centered on the microglial

synaptic pruning pathway, which includes *TYROBP*, *TREM2*, and *C1Q* complement

component. Computational modeling further suggests that neural network dynamics

exhibit differential sensitivity to synaptic pruning, with inhibitory synapse loss having

a more pronounced effect than excitatory synapse pruning. This asymmetric

sensitivity drives network hyperexcitability and increased synchronization, potentially

explaining the mechanistic link between AD and seizure susceptibility.

2. Materials and Methods

2.1 Data Collection

We conducted a comprehensive search to identify relevant transcriptomic datasets

from both animal models and patient cohorts. The selection process involved

screening based on sample size adequacy (n > 40) and research design to ensure

statistical robustness and minimize bias. Our final selection comprised several

large-scale, methodologically sound datasets from both EP and AD research that

focused on genome-wide expression patterns in well-characterized disease models

and appropriate controls. We ultimately selected four large-scale datasets from animal

models, along with three EP and two AD patient cohorts.

The EP model datasets included two of the most used TLE modeling approaches. The

first dataset (PRJEB18790) contained 200 hippocampi of pilocarpine-induced (PO)

model mice and wild-type (WT) controls [27]. The second dataset (GSE73878)

comprised 80 hippocampal samples from the kainic acid-induced (KA) model and

WT control. For the AD model datasets, we selected two datasets that systematically

examined genome-wide expression alterations in amyloid and tau pathology

progression. The Matarin dataset (GSE64398) contained 113 hippocampal samples

from five mouse models and their controls: TAU mice, TAS10 mice, TPM mice, and

TASTPM mice (both heterozygous and homozygous variants) [28]. The Castanho

dataset (GSE125957) included J20 and rTg4510 models with 144 hippocampal and

entorhinal cortex samples and matched controls [29]. To validate disease co-expression

modules across species, we constructed modules using 129 hippocampal samples

from TLE patients (GSE63808) and incorporated 30 consensus AD modules from the

AMP-AD consortium's meta-analysis of 2,114 samples across seven brain regions [30,31].

To evaluate the diagnostic value of transcriptional signatures, we analyzed 47

hippocampal samples from TLE patients and controls (E-MTAB-3123), along with

253 samples from various brain regions of AD patients and controls (GSE48350). In

addition, we reanalyzed single-nucleus RNA-sequencing (snRNAseq) datasets from

TLE patients (cells = 66305) and AD cohorts (cells = 50985) to characterize

cell-type-specific expression [32,33].

2.2 Quality control and preprocessing

All raw RNA sequences underwent quality control and trimming to obtain clean

sequences, using FastQC (0.12.1) and Fastp (0.22.0) with stringent parameters

(--qualified_quality_phred 30 --n_base_limit 5 --detect_adapter_for_pe --cut_tail

--trim_front1 15 --dedup --trim_poly_x --trim_poly_g) to ensure high data quality [34].

Ribodetector (0.2.7) was used to remove potential rRNA contamination [35].

Low-quality regions in the reads (due to platform-specific artifacts) were also

trimmed using Fastp. Trimmed sequences were then aligned to the GRCm39.111

reference mouse genome using HISAT2 (2.2.1) and followed by quantification using

featureCounts (2.0.6) [36,37]. Detailed information on mapping metrics is provided in

Table S1. Principal Component Analysis (PCA) and hierarchical clustering were

performed to identify potential outlier samples. All datasets underwent appropriate

normalization: RNAseq data were processed with variance-stabilizing transformation

using DESeq2 (1.46.0), while microarray data were subjected to log2 transformation

and variance-stabilizing normalization using the "limma" (3.62.1) package [38,39]. Batch

effects were removed using limma:removeBatchEffect(). We assessed dataset

comparability by calculating general network properties such as average gene

expression and overall network connectivity, with results indicating comparable

across all datasets.

2.3 Gene Co-expression Networks Construction

To construct and compare gene co-expression networks (GCNs) between EP and AD

model datasets, we implemented the standard WGCNA comparative methodology [40,41],

which effectively identifies conserved gene modules across datasets while mitigating

batch effects. Using the WGCNA package (1.73), we constructed signed

co-expression networks in an unsupervised, unbiased manner based on genome-wide

expression profiles. Network construction employed the minimum soft-thresholding

power ($\beta = 12$) that achieved scale-free topology ($R^2 \geq 0.85$) across all model datasets.

We generated adjacency matrices using the signed network option and transformed

them into topological overlap matrices (TOM) to quantify gene connectivity profile

similarities. Module identification proceeded through average linkage hierarchical

clustering of TOM dissimilarity (1-TOM), followed by dynamic tree cutting

(parameters: deepSplit = 1, minClusterSize = 27). For each module, we calculated

module eigengenes (MEs) from the first principal component to summarize module

expression profiles. We then identified disease-associated modules by computing

Pearson correlation coefficients between MEs and genotype information, with

significance thresholds of $P < 0.05$ and |correlation coefficient| > 0.35. Additionally,

we determined module membership (MM) and genotype significance (GS) by calculating Pearson correlations between individual genes and their respective MEs or genotypes. Gene Ontology (GO) enrichment analyses for module annotation were conducted using clusterProfiler (4.10.1) [42].

2.4 Module Preservation Analysis

To evaluate the preservation of EP model modules across AD model datasets, we implemented dual analytical approaches as described by Langfelder et al. [43]. First, we calculated Zsummary preservation statistics using the modulePreservation function (networkType = "signed", nPermutations = 100). This composite metric comprehensively assesses preservation by integrating module size, density, and connectivity parameters, quantifying how effectively EP module structures are maintained in AD datasets. We interpreted the preservation evidence according to established thresholds: strong (Zsummary > 10), moderate (2 < Zsummary < 10), or absent (Zsummary < 2). We complemented this analysis with the medianRank preservation statistic, which evaluates relative module preservation with reduced sensitivity to module size variations, where lower ranks indicate stronger preservation. Additionally, we performed cross-tabulation analysis to quantify module overlap between EP and AD model datasets, as well as between model and patient datasets. This approach generated contingency tables detailing the number of genes shared between reference and test dataset modules. We determined overlap significance

(adjusted $P < 0.05$) using Fisher's exact test with Benjamini-Hochberg (BH)

correction for multiple testing.

2.5 Consensus Network and Signatures Identification

After confirming significant preservation of key modules across datasets, we

constructed consensus modules to identify gene co-expression patterns consistently

present in both conditions. This consensus approach involved calculating the

component-wise minimum values of the topological overlap matrices from all datasets,

thereby capturing shared network topology while filtering dataset-specific variations.

We applied hierarchical clustering, using the same parameters as those applied to

individual networks, to the consensus dissimilarity matrix (1-consensusTOM) to

identify robust consensus modules [41]. Hub genes act as key drivers within a module

and were identified using stringent criteria to minimize false positives: genes with

MM > 0.9 and ranking in the top 5% for intramodular connectivity within each

module. Consensus hub signatures were determined by intersecting the hub genes of

key modules in both EP and AD datasets, requiring an overlap count of at least three

across datasets. The expression and diagnostic value of these consensus hub

signatures were further explored in the AD and EP patient datasets (GSE48350 and

E-MTAB-3123).

To explore interactions and potential regulatory relationships among consensus hub

signatures, we constructed a protein-protein interaction (PPI) network using the

STRING database (12.0) (https://string-db.org/cgi/input.pl) with a confidence score

threshold of 0.4 and a false discovery rate (FDR) of 5% [44]. We also performed

regulatory network analysis with the GENIE3 (1.28.0) algorithm, which applies a

random forest approach to infer potential regulatory relationships from gene

expression data. GENIE3 ranks regulatory interactions by weight, with higher weights

indicating stronger regulatory links [45]. Gene expression profiles of the consensus hub

signatures from all datasets were used to infer plausible regulatory interactions. To

reduce false positives, only the top 30% plausible regulatory links were retained to

construct gene regulatory networks.

2.6 Machine Learning

To evaluate the diagnostic potential of consensus hub signatures, we conducted binary

classification (disease vs. control) across datasets using these signatures as predictive

features. Prior to analysis, gene expression values underwent standardization to ensure

comparable scaling across genes. We implemented classification models in Python

3.8.20 using scikit-learn, XGBoost, and CatBoost libraries. Seven supervised learning

algorithms were evaluated: Logistic Regression (with increased maximum iterations

of 1000), Support Vector Machine with RBF kernel, Random Forest, Naive Bayes,

XGBoost, AdaBoost, and CatBoost. To ensure robust performance estimation and

prevent overfitting, we employed 5-fold stratified cross-validation, with each fold

maintaining proportional representation of disease and control samples. Performance

was assessed through multiple metrics including Accuracy, Precision, Recall, F1 score,

and Area Under the Receiver Operating Characteristic Curve (AUC). The AUC was

calculated by pooling predicted probabilities from all cross-validation splits to

generate Receiver Operating Characteristic (ROC) curves.

2.7 Cell Composition Deconvolution and Single-nucleus RNA-seq Analysis

In silico deconvolution can estimate the cellular composition of a tissue sample from

its bulk gene expression profile, with CIBERSORT particularly suited for cellular

composition deconvolution of brain transcriptomes [46,47]. Cell composition

deconvolution was performed using CIBERSORT (1.04) via the BrainDeconvShiny

platform, with cell-type signatures specific to *Mus musculus* from immunopurified

mouse brain tissue [46,47]. Associations between identified cell types and modules

were determined by calculating Pearson correlation coefficients.

snRNA-seq datasets from EP and AD cohorts were reanalyzed to investigate

cell-type-specific transcriptional changes of consensus transcriptional signatures. For

EP cohort, we utilized snRNA-seq dataset (GSE190453) profiling temporal lobe

tissue from TLE patients and non-epileptic controls (n = 8) [32]. For AD cohort, we

analyzed snRNA-seq data from the Religious Orders Study and Rush Memory and

Aging Project (ROSMAP) cohort, encompassing 48 prefrontal cortex (PFC) samples

from AD patients and non-pathology controls (n =48) [33]. Data preprocessing was

performed using the Seurat package (5.0.1) according to the original studies [48]. Nuclei

were filtered based on standard quality control criteria: nuclei with fewer than 200

detected genes, excessive gene counts (thresholds adjusted per sample), or

mitochondrial transcript percentages exceeding 20% were excluded. Mitochondrial

content was calculated using the PercentageFeatureSet() function. After filtering,

SCTransform normalization was applied to each sample, regressing out total RNA

counts and mitochondrial content to correct for technical variability. Batch correction

across samples was performed via Seurat's integration workflow, anchoring on the top

3,000 variable features. Dimensionality reduction was conducted through PCA, with

the first 30 components used for clustering and UMAP visualization. Cell-type

annotation followed the definitions established in the original studies and calculate

scores for each nucleus using the AddModuleScore() function, and cell-type identity

was assigned based on the highest-scoring module. Cluster annotations were further

refined by majority voting based on the predominant cell-type identity within each

cluster. Differential expressions of consensus transcriptional signature between

control and disease groups were evaluated using two-sided Wilcoxon rank-sum tests.

2.8 Computational modeling

We implemented a spiking neural network using the Izhikevich neuron model to

simulate hippocampal CA3 circuit dynamics and explore the contribution of synaptic

pruning to pathological changes in AD and EP [49–51]. The network consisted of 1000

standard Izhikevich neurons (800 excitatory, 200 inhibitory) and simulated for

1000ms using the Brian2 package [52]. Network architecture included random sparse

connectivity (0.1 probability) with conductance-based synapses modeling AMPA

(excitatory) and GABA-a (inhibitory) currents across four connection types: E→E,

E→I, I→E, and I→I [53]. Synaptic weights were drawn from a standard normal

distribution and normalized to [0,1]. We simulated synaptic pruning by selectively

removing excitatory or inhibitory synapses with the lowest weights (0-50%, with 5%

increments), repeating each simulation at least 50 times.

To quantify network dynamics, we calculated the excitation/inhibition (E/I) ratio and

event synchrony (ES) [54]. The E/I ratio compared total excitatory currents to the

absolute magnitude of inhibitory currents, while ES measured spike timing

coordination between excitatory neuron pairs within 5-ms windows. Both metrics

were normalized to baseline unpruned values to reflect pruning-induced changes.

Statistical analysis included Pearson correlation coefficients and linear regression to

quantify the effect sizes of excitatory and inhibitory pruning on network dynamics.

Complete details of the computational model parameters, equations, and analysis

methods can be found in **Text S1.**

3. Results

3.1 Identification of Consensus Gene Co-Expression Network and Hub

Signatures

To identify conserved molecular signatures underlying EP and AD, we employed an

unbiased, genome-wide approach to construct GCNs across four independent datasets:

two EP models (pilocarpine-induced [EP-PO] and kainic acid-induced [EP-KA]) and

two AD datasets (AD-Castanho: [rTg4510, J20]; AD-Matarin: [TAU, TAS10, TPM,

HO-TASTPM, HET-TASTPM]). After removing only genes with negligible or

undetectable expression, we retained all expressed genes (ranging from 12,558 to

17,813) to avoid selection bias.

Network construction using consistent soft-thresholding power ($\beta = 12$) identified

common and distinct GCNs modules across all model datasets (**Fig 1A, S1**): EP-PO

revealed 19 modules (97- 1203 genes), EP-KA contained 14 modules (36-1534 genes),

AD-Castanho yielded 28 modules (55-820 genes), and AD-Matarin showed 16

modules (33-1354 genes). Correlation analysis between module eigengenes and

disease status identified disease-associated modules in each model (**Table S2**). In the

EP-PO dataset, 12 modules showed significant association with the EP genotype ($|r| >$

0.35, $P < 0.05$). Similarly, EP-KA had 2 disease-associated modules, AD-Castanho

contained 12 modules correlated with tau pathology, with 4 modules correlated with

general AD pathology, and AD-Matarin revealed 5 modules associated with tau

pathology and 2 modules associated with amyloid pathology, with 3 modules

correlated with general AD pathology (**Fig 1A, Table S2**).

Functional enrichment analysis of disease-specific modules revealed several distinct

biological processes unique to each condition (**Table S2**). EP-specific modules were

enriched in term such as "synapse organization" (PO-lightcyan), "modulation of

chemical synaptic transmission" (PO-yellow) and "regulation of calcium ion import"

(KA-midnightblue) while AD-specific modules showed enrichment in term such as

"establishment of protein localization to organelle" (Matarin-blue), "dendrite

development "(Castanho-saddlebrown) reflecting the unique pathophysiological

mechanisms in each disease (**Table S2**). Despite disease-specific differences,

cross-condition preservation analysis revealed remarkable conservation of an immune

module across all four models (**Fig. 1A**). Using Z-summary preservation statistics and

cross-tabulation approaches, we identified a large immune-related module (turquoise)

with exceptionally highest preservation (Z-summary > 50) across all pairwise

comparisons (**Fig. 1A, Table S2**). This module demonstrated significant overlap

among all four models (adjusted P < 1E-300) and was negatively correlated neuron

proportion and positively correlated with disease progression genotype in each model

(**Table S2**): EP-PO (r = 0.80, P = 6.13E-42), EP-KA (r = 0.52, P = 9.31E-07),

AD-Castanho (r =0.46, P = 8.63E-08), and AD-Matarin (r = 0.56, P = 9.14E-11).

The strong preservation pattern of this immune module across both diseases prompted

us to construct consensus GCNs by integrating network topology information from all

datasets (**Fig. S2**). This approach identified a robust immune-related and

disease-associated consensus module containing 1617 genes common to all models

(**Table S2-3**). To identify key drivers within this consensus immune module, we

applied stringent criteria based on intramodular connectivity and module membership

across all four datasets, which identified 19 consensus hub signatures consistently

present in the immune modules across all models (**Fig. 1B**). These hub signatures

included microglial activation markers (*TREM2, TYROBP, CD68*), *C1Q* complement

components (*C1QA, C1QB, C1QC*), and inflammatory mediators (*TLR2, FCGR2B,*

CYBA). Additional immune-related genes comprised lysosomal enzymes (*CTSH,*

CTSZ, GUSB), immune signaling regulators (*CD84, CD52, SLC11A1, UNC93B1*),

genes involved in phagocytosis and motility (*MYO1F, PROS1*), and the immune

checkpoint receptor *LAG3*.

To investigate the biological significance and potential regulatory relationships among

these hub signatures, we employed a multi-layered network analysis approach

combining protein-protein interaction data from STRING with gene regulatory

networks inferred by GENIE3 (**Fig. 1B**). Network centrality analysis identified

TYROBP as the most upstream hub signature with highest total degree

(regulator/target ratio = 1.33, total degree = 24.19), followed by *CD68*, *TREM2*, *C1Q*

complement components and *SLC11A1* (**Fig. 1B, Table S2**). GO enrichment

analysis of the hub signature network identified "synaptic pruning" as the most

significantly enriched biological process with strongest signal (FDR = 4.02E-07,

signal = 2.91), which include *C1Q* complement components and microglial receptors

(*TREM2*) (**Fig. 1C**). Genes involved in "Synaptic pruning" also participate in other

enriched processes such as "Complement-mediated synapse pruning" (*C1QA*, *TREM2*)

and "Microglial cell activation" (*TYROBP*, *C1QA*, *TREM2, SLC11A1, TLR2*) (**Fig. 1C,**

Table S2).

Taken together, our genome wide transcriptomic analysis revealed a highly conserved

immune module in EP and AD model, with a consensus regulatory network centered

around *TYROBP, TREM2, CD68, C1Q* components that coordinate microglia

activation and synaptic pruning. This consensus molecular pathway represents a

potential central mechanism linking both neurological conditions.

3.2 Consensus GCNs Significantly Overlap with GCNs in Human Patients

To validate the translational relevance of our identified consensus GCNs, we

compared them with disease-associated GCNs from human patient datasets. We

employed a comprehensive cross-species approach to identify overlaps between our

consensus modules and published AD-associated co-expression modules from 2,114

samples across seven brain regions, as well as our newly constructed EP patient

modules from 127 TLE patient hippocampal samples (**Fig. 2A, S3, Table S3**).

Our analysis revealed significant overlaps between the consensus immune module and

multiple human AD modules across different brain regions. The strongest associations

were observed with modules from cerebellum (CBE), dorsolateral prefrontal cortex

(DLPFC), and superior temporal gyrus (STG) regions. The consensus module

demonstrated particularly strong overlap with the " CBE-turquoise" module (n = 502

genes, 25.39% of the patient module, P < 1E-300), the "DLPFC-blue" module (n =

502 genes, 28.67% of the patient module, P < 1E-300) and "STG-blue"(n = 389 genes,

33.22% of the human module, P < 1E-300) (**Fig. 2A, Table S2**). Similarly, substantial

overlaps were detected with modules from inferior frontal gyrus (IFG), superior

temporal gyrus (STG), temporal cortex (TCX), and frontal pole (FP) but the overlap

percentage varied from 2% to 33% (**Table S2**).

For EP patient dataset, our consensus module showed significant overlap with two EP

patient's co-expression modules (yellow: n = 153 genes, 39.64% of the patient

module, P < 7.07E-88; blue: n = 279 genes, 22.85% of the patient module, P <

1.06E-84) (**Fig. 2A, Table S2**). Notably, the top hub genes identified in our animal

model analysis—*TYROBP*, *TREM2*, and the *C1Q* components—were also identified

as central hub genes in the corresponding EP patients's yellow module, with *TYROBP*

showed the top 2 intramodular connectivity (**Table S3**).

To further validate our hub consensus signatures findings, we examined whether the

19 previously identified hub signatures remained clustered within the same modules

in human disease networks (**Fig. 2B**). The co-occurrence analysis revealed strong

co-expression patterns, with particularly strong co-expression observed among *C1Q*

complement components, *TREM2*, *TYROBP*, *CYBA*, *SLC11A1*. These genes showed

the highest co-occurrence frequency (dark blue in the heatmap), suggesting they

function as a coordinated unit across species and disease conditions. Collectively,

these cross-species analyses demonstrate that the identified immune modules and hub

signatures represent conserved biological entities relevant to both AD and EP

pathogenesis in humans.

3.3 Consensus Hub Signatures Displayed Strong Diagnostic Value

To validate the clinical relevance of the identified consensus hub transcriptional

signatures in AD and EP animal models, we analyzed the expression of 19 hub

consensus signatures in bulk and single nucleus datasets from AD and EP patients

(GSE48350, E-MTAB-3123, GSE190453 and ROSMAP PFC) compared to their

controls.

Firstly, we localized the expression of these signatures in patient cell types, which

demonstrated predominantly expression in microglia (**Fig. 3A-D**). Some signatures

such as *CTSH*, *MYO1F*, and *GUSB* showed broad expression across various cell types.

Among all the microglia-specific signatures, expression of *TYROBP*, *TREM2*, and the

C1Q complement genes were consistently detectable and exhibited high average

expression across EP and AD cohorts (**Fig. 3B, D**). We then performed differential

expression analysis on microglia and bulk datasets (**Fig. 3E-H**). We observed that

almost all signatures were significantly upregulated in the EP cohort, and most

signatures were also significantly upregulated in the AD cohort, with particularly high

expression in microglia of late pathology (**Fig. 3G**). Notably, *TYROBP*, *TREM2*, and

the *C1Q* complement genes—identified as central consensus signatures implicated in

microglial synaptic pruning—were consistently upregulated in both disorders (**Fig.**

3E-H).

Using these gene expression profiles from bulk datasets (GSE48350, E-MTAB-3123),

we developed diagnostic models with multiple machine learning algorithms to explore

the diagnostic value of these 19 hub consensus signatures. These models generally

achieved excellent discrimination in the EP dataset (AUC > 0.9) and good

performance in the AD cohort (AUC > 0.7) (**Fig. S4**). The superior classification

accuracy in EP likely reflects the more substantial transcriptional alterations observed

in these patients. These findings demonstrate that these consensus signatures also

dysregulated in human patients and possess strong diagnostic utility. This is especially

true for genes involved in microglia activation and synaptic pruning (e.g., *TYROBP*,

TREM2, and the *C1Q* complement genes).

3.4 Asymmetric Effects of Synaptic Pruning on Neural Network Dynamics

To better understand how synaptic pruning influences neural network dynamics and

its potential role in the comorbidity of AD and EP, we developed a neural network

model simulating the hippocampal CA3 circuit. In this model, we systematically

varied the extent of synaptic pruning (0% to 50%) at either excitatory or inhibitory

synapses. We then quantified changes in the E/I ratio and synchronization levels,

conducting at least 50 simulation runs per condition (**Fig. 4**).

Our findings indicate that pruning either excitatory or inhibitory synapses alters the

E/I ratio, but with differential effects. Inhibitory synaptic pruning significantly

elevated the E/I ratio, while excitatory pruning reduced it. These bidirectional trends

were consistent across all simulations and in experiments where excitatory or

inhibitory pruning was varied independently (**Fig. 4A**). Correlation analysis revealed

a negative association between excitatory pruning and the E/I ratio ($r = -0.474$) and a

stronger positive correlation between inhibitory pruning and the E/I ratio ($r = 0.684$)

(**Fig. 4B**). Linear regression further confirmed this differential sensitivity, with

excitatory pruning reducing the E/I ratio ($\beta = -1.038$), whereas inhibitory pruning

increased more substantially ($\beta = 1.454$), indicating that E/I balance is more sensitive

to changes in inhibitory connectivity (**Fig. 4B**).

Notably, network synchronization exhibited an even more pronounced differential

sensitivity to pruning type. Inhibitory pruning displayed a moderate positive

correlation with event synchronization (r = 0.346), while excitatory pruning showed

only a negligible negative correlation (r = −0.039) (**Fig. 4B**). Linear regression

analysis further confirmed this asymmetric sensitivity, with inhibitory pruning

exerting a substantially stronger effect on synchronization (β = 0.358) compared to

excitatory pruning (β = −0.041) (**Fig. 4B**). Additionally, the E/I ratio itself was

positively correlated with synchronization (r = 0.491), suggesting that the influence

of synaptic pruning on network synchronization may be indirectly mediated through

its effects on E/I balance (**Fig. 4C**).

These simulations demonstrate that while both types of synaptic pruning affect

network properties, inhibitory pruning exhibits a heightened potency in disrupting

both E/I balance and synchronization dynamics. This differential sensitivity to

inhibitory pruning may represent a key mechanism by which synaptic loss in

neurodegeneration conditions could precipitate network hyperexcitability and

hypersynchrony characteristic of epileptiform activity.

4. Discussion

While Alzheimer's disease (AD) and epilepsy (EP) share a complex bidirectional

relationship, the molecular pathways linking AD-related neurodegeneration to

increased EP susceptibility remain insufficiently explored. To address this gap, we

employed robust genome-wide, unsupervised bioinformatic and computational

approaches to analyze transcriptional data from multiple large-scale animal models

and human patient datasets. This comprehensive analysis identified consensus

molecular pathways, cellular processes, and neural network dynamics associated with

this comorbidity across diverse experimental paradigms.

Conserved Neuroinflammation Pathways in Epilepsy and Alzheimer's Disease

Our genome-wide analysis provides compelling evidence supporting the hypothesis

that neuroinflammatory pathways form a common molecular substrate between AD

and EP [55]. We identified remarkable preservation of the immune-related turquoise

module across both disease models and human patient datasets **(Fig. 1-2)**. Functional

enrichment analysis of the preserved module revealed significant enrichment in

neuroinflammatory processes, such as "inflammatory response", "cytokine

production", "blood vessel morphogenesis", "glial cell activation", and "complement

and coagulation cascades" (**Table S2**).

Neuroinflammation has been extensively documented in both diseases and appears to

serve a complex dual role in disease progression [55–57]. While activated microglia

facilitate Aβ plaque clearance and reactive astrocytes provide metabolic support and

ionic homeostasis that may attenuate neural damage and seizures, chronic activation

or dysfunction of these glial populations can promote disease progression through

persistent neuroinflammation [58–60]. Specifically, activated microglia and astrocytes

secrete proinflammatory cytokines and chemokines, including IL-1β, TNF-α, and

IL-6, which can enhance neuronal hyperexcitability, increase seizure susceptibility,

exacerbate Aβ and tau aggregation, and accelerate neurodegeneration and cognitive

decline [61].

Additional inflammatory components, including complement cascade activation and

blood-brain barrier (BBB) dysfunction, are implicated in both pathologies [62–65].

Complement activation contributes to synaptic loss in AD and is elevated in TLE

patient tissue [64,65]. BBB disruption precedes neuronal hyperactivity in EP, promoting

astrocyte dysfunction and seizures. In AD, BBB disruption facilitates peripheral

inflammatory cell infiltration, thereby amplifying neuroinflammation [62,63].

These results establish neuroinflammation as a key mechanistic link between AD and

EP [55]. AD-associated inflammation can enhance neuronal hyperexcitability and

seizure susceptibility. Conversely, EP-induced inflammation may accelerate Aβ and

tau accumulation, contributing to cognitive decline. Our results support previous

studies and suggest that neuroinflammation is highly conserved across both diseases

and species.

Synaptic Pruning was Identified as a Core Pathway in this Comorbidity

Our systematic in-depth analysis of the conserved immune module revealed a central

synaptic pruning pathway involving *TYROBP*, *TREM2*, and *C1Q* complement genes

(**Fig. 1**). These genes demonstrated consistent co-dysregulated in microglia and bulk

expression data and demonstrated strong diagnostic value across both AD and EP

patient, (**Fig. 2-3**). Collectively, these findings suggest that this microglia-mediated

synaptic pruning may represent a core mechanistic pathway underlying the

comorbidity of AD and EP.

This pathway has been extensively documented in AD pathology and experimental

studies, demonstrating its involvement in both tau and Aβ pathology [66–69].

Dysregulation or mutations in *TYROBP* and *TREM2* are established risk factors for

AD due to their effects on microglial function [70–72]. Under physiological conditions,

microglia maintain synaptic homeostasis through a regulated process in which

complement proteins, including *C1Q*, tag unnecessary synapses for elimination [73].

Microglial receptors, particularly *TREM2* and *TYROBP*, recognize these tags and

initiate phagocytic clearance, a process critical for neural circuit refinement and

maintenance in the adult brain [66]. In AD pathology, these receptors enable microglia

to recognize protein deposits, triggering their transition to a disease-associated

microglial (DAM) state characterized by enhanced phagocytic activity and elevated

expression of *TYROBP*, *TREM2*, *APOE*, and complement proteins including *C1Q* [66].

While *TYROBP* or *TREM2* deficiency disrupts DAM activation and impairs critical

microglial functions such as plaque clearance, potentially accelerating AD pathology,

overactivation of this pathway can also contribute to neurodegeneration and cognitive

decline through excessive *C1Q*-mediated synaptic pruning [67–69].

Synaptic pruning can exert effects on both AD and EP disease progression through

two primary mechanisms: synaptic transmission and synaptic plasticity. Regarding

synaptic plasticity, research demonstrated that *C1Q* inhibition in aged experimental

models can restore both long-term potentiation (LTP) and long-term depression (LTD)

[67–69]. In non-aged models, increased *C1Q* expression also impairs the LTP process [74].

This is particularly relevant for AD, where impaired synaptic plasticity is a feature of

cognitive decline. In EP, compromised synaptic plasticity may contribute to the

formation and maintenance of epileptogenic circuits. Regarding synaptic transmission,

elevated *C1Q* expression or Aβ accumulation in the CA1 region leads to decreased

miniature and evoked excitatory postsynaptic currents [75,76]. In AD, this reduction in

synaptic transmission contributes to cognitive decline by disrupting normal neural

communication. For EP, this alteration affects both excitatory and inhibitory neurons

[74], potentially contributing to the imbalance between excitation and inhibition that

characterizes epileptic activity. Notably, these effects can be reversed through *C1Q*

downregulation [77,78]. The relationship between these mechanisms appears

bidirectional: excessive synaptic pruning leads to neuronal damage and reduced LTP,

which in turn promotes disease progression in both conditions, creating a harmful

feedback loop where synaptic dysfunction promotes pathological processes.

Notably, interventions targeting the synaptic pruning pathway, whether through the

knockdown of regulatory genes like *TYROBP* and *TREM2* or direct inhibition of *C1Q*

accumulation, have successfully restored both electrophysiological and cognitive

functions in AD models. This restoration occurred even in the presence of significant

amyloid and tau pathology [66–69]. Our results establish a regulatory network that is

consistent with previous observations and further extends its role in the comorbidity

of AD and EP (**Fig. 1**).

While this pathway has been thoroughly investigated in AD pathology, to the best of

our knowledge, no previous studies have linked it to the comorbidity between EP and

AD. Additionally, our understanding of how synaptic pruning relates to EP remains

limited, particularly regarding its effects on global neural network dynamics [79]. To

address this gap, we conducted an in-silico experiment to shed light on this issue.

Asymmetric Effects of Synaptic Pruning Promote Network Synchronization

Computational modeling has become an important tool for understanding AD and EP

pathology, with models ranging from molecular-level simulations of ion dynamics to

large-scale network models [80,81]. To explore how synaptic pruning contributes to the

comorbidity of AD and EP, we developed a computational model of the hippocampal

CA3 circuit building upon Liu's biologically plausible EP modeling framework [53].

This model allowed us to quantify how pruning of excitatory and inhibitory synapses

affects E/I balance (E/I ratio) and the temporal coordination of neural firing

(synchronization).

Our computational model reveals a differential sensitivity of neural networks to

synaptic pruning, with E/I balance being substantially more responsive to inhibitory

synapse loss than excitatory synapse elimination (**Fig. 4**). Quantitative analysis by

linear regression demonstrated that excitatory synapse elimination decreased the E/I

ratio (β = -1.038), while inhibitory synapse removal led to a significantly larger

increase in the E/I ratio (β = 1.454) (**Fig. 4B**). This heightened sensitivity to

inhibitory pruning can be explained by the architectural role of inhibitory interneurons

within neural circuits. These neurons often establish synaptic connections with

multiple excitatory neurons and act as crucial regulators of circuit coordination and

synchronization. Therefore, when inhibitory control mechanisms are impaired, the

resulting dysregulation cascades throughout more widespread circuits [82].

Consequently, a reduction in inhibitory synapses produces a disproportionately larger

perturbation in neural networks than an equivalent reduction in excitatory synapses.

Even more striking was the asymmetric sensitivity of network synchronization to

different types of synaptic pruning. Our findings reveal that inhibitory pruning

exhibited a moderate positive correlation with synchronization ($r = 0.346$), whereas

excitatory pruning had a negligible correlation ($r = -0.039$) (**Fig. 4B**). This

differential sensitivity suggests that synchronization dynamics are primarily

vulnerable to disruptions in inhibitory circuitry. Notably, the E/I ratio itself

demonstrated a strong correlation with synchronization levels ($r = 0.491$) (**Fig. 4C**).

While it seems intuitive that enhancing network excitability can promote synchrony,

our analysis highlights the importance of considering the specific location and

functional role of pruned synapses in neural circuits. For example, in a circuit that

exerts an inhibitory effect on the broader network, pruning inhibitory synapses that

inhibits this circuit paradoxically reduces network synchronization. This explains the

indirect, nonlinear association of synaptic pruning with network synchronization.

Although establishing a direct causal relationship between synaptic pruning and

AD-EP comorbidity remains challenging, our simulation results suggest a mechanistic

pathway whereby neuronal or synaptic loss in neurodegenerative diseases disrupts E/I

balance with heightened sensitivity to inhibitory synapse loss, thereby promoting EP

susceptibility (**Fig. 4**). This mechanism may extend beyond AD, as EP is a

well-documented comorbidity of multiple neurodegenerative conditions including

Huntington's and Parkinson's disease [83]. Our simulations demonstrate that even with a

proportional loss of both synapse types (or even with greater excitatory synapse loss),

the effect of inhibitory synapse loss can overpower that of excitatory synapse loss.

This occurs due to differential sensitivity, leading to more extensive disruption of

network E/I balance and promoting network synchronization (**Fig. 4**). Similar to

TYROBP and *TREM2* being established risk factors for AD [71,72], Mendelian

randomization studies have also indicated that *C1Q* a is a risk factor for both AD and

EP (**Fig. S5**) [84]. Notably, in AD, synaptic loss often precedes cognitive decline and EP,

suggesting it may be an initiating event rather than a consequence [1].

Our results emphasize the critical role of inhibitory synaptic pruning in promoting

network synchronization due to the neural network's heightened sensitivity to

inhibitory synapse loss. At the molecular level, *C1Q* functions as an "eat me" signal,

tagging synapses for microglial elimination [79]. While this tagging process itself is not

synapse-type specific, interestingly, several recent studies have demonstrated that

microglia may preferentially prune inhibitory synapses [79,85]. Dejanovic et al.

demonstrated that the *TREM2-C1Q* microglial synaptic pruning pathway engulfs

inhibitory synapses more frequently than excitatory synapses in AD models, while

Fan et al. observed similar preferential pruning of inhibitory synapses during recurrent

seizures that exacerbate EP model pathology [85,86]. Studies have shown that *C1Q*

blockade reduces both the frequency and severity of seizures in traumatic brain injury

models [87]. However, it is also important to note that *C1Q* knockout also affects

synaptic elimination during development, resulting in enhanced connectivity and

seizure activity [88,89]. Recently, Aloi et al. also reported that overactivation of microglia

reduced Aβ pathology but also caused hyperexcitability and seizures via synaptic

pruning [90].

These findings collectively suggest that the network's differential sensitivity to

inhibitory synapse loss, coupled with the apparent preferential pruning of inhibitory

synapses by microglia, may represent a critical mechanism underlying AD-EP

comorbidity. The management of microglial *C1Q*-mediated synaptic pruning during

AD pathology and aging warrants further investigation, particularly regarding the

regulation of this potential preferential pruning of inhibitory synapses.

Limitations

Our study presents two key limitations. First, while we identified a conserved

microglial synaptic pruning pathway across multiple datasets, our reliance on bulk

expression data limits our ability to fully characterize microglial and neuron

subtype-specific contributions. Despite our computational deconvolution efforts and

cell type expression analyses, future research should employ spatial transcriptomics to

more precisely map the cell subtype-specific dynamics in AD-EP comorbidity,

particularly focusing on microglial activation patterns and their synaptic targeting

behavior.

Second, our computational model is informative about synaptic pruning's impact on

network dynamics. However, it needs to simplify several aspects of neural circuits,

including network topology and dendritic complexity. Future modeling cam

incorporate more detailed cellular mechanisms and circuit architectures to better

capture the molecular progression of AD-EP comorbidity and the asymmetric effects

of inhibitory versus excitatory synaptic pruning on network synchronization.

Conclusion

Our study involved a genome-wide analysis of large-scale transcriptomic datasets

from multiple AD and EP animal models, as well as human patient cohorts, revealing

a consensus molecular signature converging on microglial synaptic pruning. We

identified a highly preserved immune-related co-expression module centered on the

TYROBP-TREM2-C1Q complement pathway that is consistently upregulated across

species and disease conditions. To facilitate module reuse and validation by the

research community, we also developed and provided a web application

(https://github.com/HuihongLi/OverlapModule) that enables researchers to test the

overlap preservation of their own modules across our module. Importantly, our

computational modeling demonstrates that neural networks exhibit differential

sensitivity to synaptic pruning, with inhibitory synapse loss exerting

disproportionately powerful effects on E/I balance and network synchronization

compared to excitatory synapse loss.

These findings illuminate a mechanistic link between neuroinflammation and network

dysfunction in AD-EP comorbidity. The complement-mediated synaptic pruning

pathway can provide a unifying framework that explains the bidirectional relationship

between these conditions observed clinically. The asymmetric sensitivity of neural

networks to inhibitory pruning—coupled with evidence that microglia preferentially

target inhibitory synapses—creates a pathological loop that drives hyperexcitability

and hypersynchrony. These findings suggest that targeting microglia–complement

signaling may provide disease-modifying benefits by preserving inhibitory circuit

integrity and preventing the onset of network hyperexcitability. Future research

utilizing spatial transcriptomics and selective manipulation of complement pathways

in specific microglial subpopulations will be crucial to validate this mechanism and

develop targeted therapeutic strategies for patients suffering from AD, EP, or their

comorbidity.

Abbreviations

EP   Epilepsy

AD   Alzheimer's disease

E/I   Excitation/Inhibition

TLE   Temporal lobe epilepsy

759	Aβ	Amyloid-β
760	PO	Pilocarpine-induced
761	KA	Kainic acid-induced
762	GCNs	Gene co-expression networks
763	WT	Wild-type
764	PCA	Principal Component Analysis
765	BH	Benjamini-Hochberg
766	GO	Gene ontology
767	TOM	Topological overlap matrices
768	MEs	Module eigengenes
769	MM	Module membership
770	GS	Gene-genotype significance
771	PPI	Protein-protein interaction
772	FDR	False discovery rate
773	AUC	Area Under the Receiver Operating Characteristic Curve
774	ROC	Receiver Operating Characteristic
775	CBE	Cerebellum
776	DLPFC	Dorsolateral prefrontal cortex
777	STG	Superior temporal gyrus
778	IFG	Inferior frontal gyrus
779	TCX	Temporal cortex

780	FP	Frontal pole
781	SVM	Support vector machine
782	ES	Event synchrony
783	BBB	Blood-brain barrier
784	DAM	Disease-associated microglial
785	LTP	Long-term potentiation
786	LTD	Long-term depression
787	BP	Biological processes
788	ROSMAP	Rush Memory and Aging Project
789	snRNAseq	Single nucleus RNA-sequencing

Figures and legends

Figure 1. Identification of conserved immune-related GCNs and hub signatures

(A) Chord diagram showing module preservation across EP and AD models. Each

segment represents a gene module identified in different datasets. Purple connections

indicate statistically significant module overlap (Fisher's exact test, adjusted $P <$

1E-300), with color intensity corresponding to significance level (-log10(padj)). Inner

tracks display module correlations with disease status (red: positive correlation; blue:

negative correlation) and neuron proportion.

(B) Integrated PPI and regulatory network of consensus hub signatures identified

across all models. Nodes represent hub genes consistently present in the preserved

immune module, with node size proportional to total weighted degree. Solid lines

indicate validated protein-protein interactions (STRING), dotted lines indicate

computationally inferred gene regulation (GENIE3), with edge width reflecting

interaction strength. Node color indicates regulator/target ratio, with red showing

stronger regulatory potential.

(C) GO enrichment analysis of consensus hub signatures. Bar length corresponds to

enrichment signal strength, color indicates significance level (FDR), and size reflects

gene count within each category. Bars represent significantly enriched biological

processes, with "Synaptic pruning" emerging as the most significant enriched process

with highest signal.

Figure 2. Comparison of consensus modules with patient modules.

(A) The overlap between the consensus immune module and disease-associated GCNs

from human patients. Peripheral nodes represent modules from different brain regions

in AD and EP patient datasets. Line thickness indicates the number of overlapping

genes between each patient module and the consensus module, with thicker lines

showing stronger connections. Line color intensity corresponds to statistical

significance (-log10(padj)), with red-orange hues indicating the highest significance

levels. The significant associations (adjust $P < 0.05$) were observed with modules

across brain regions in AD patients, and with yellow and blue modules in EP patients.

(B) Co-occurrence heatmap of the 19 identified consensus hub signatures across

human disease modules. Color intensity represents the frequency of gene

co-expression (darker blue indicates higher co-occurrence counts).

Figure 3. Expression of consensus hub signatures in EP and AD patient cohorts.

(A-D) UMAP visualization and dot plot showing cell type-specific expression of

consensus hub genes in EP patient samples **(A-B)** and AD patient samples **(C-D)**.

Microglia show the highest expression of most hub genes, with *TYROBP*, *TREM2*,

and *C1Q* complement genes exhibiting particularly strong expression.

(E-H) Differential expression analysis of consensus hub genes in microglia and bulk

tissue from EP patients (**E-F**) and AD patients (**G-H**) compared to controls. Bar

heights represent fold changes with statistical significance indicated (*p < 0.05, **p <

0.01, ***p < 0.001).

Figure 4. Asymmetric Effects of synaptic pruning on neural network dynamics.

(**A**) Simulation results of excitatory pruning ratio (0-0.5), inhibitory pruning ratio

(0-0.5), E/I ratio, and synchronization. X-axis: Excitatory pruning ratio; Y-axis:

Inhibitory pruning ratio; Z-axis: Average E/I ratio. Colors indicate average

synchronization levels from dark purple (close to 1, indicating no change in

synchrony) to yellow (indicating higher synchrony).

(B) The distribution of E/I ratio and synchronization values across different pruning

levels. Each violin represents the distribution of values at a given pruning ratio (0–0.5

in 0.05 increments). Embedded boxplots display median and interquartile ranges.

Overlaid dots represent individual simulation outcomes. Regression results (slope,

p-value, and Pearson's r) are displayed in the upper right of each panel, indicating that

inhibitory pruning has a stronger positive correlation with both E/I ratio and

synchronization, while excitatory pruning shows weaker or negative correlations.

(C) The relationship between E/I ratio and synchronization across all simulation

conditions. A positive correlation is observed, suggesting that changes in E/I balance

mediate alterations in network synchrony. Regression statistics are displayed in the

corner.

Legends of supplementary files

Text S1: Computational modeling details

Table S1: Summary of Quality Control and Mapping Statistics

Table S2: Statistics of Module Preservation and Functional Enrichment

Table S3. Module gene information

Figure S1. Construction of co-expression modules in model datasets.

(A) Scale independence and mean connectivity plots for different soft-thresholding

powers across all four datasets (EP-PO, EP-KA, AD-Matarin, AD-Castanho). A

power of 12 was selected as the minimum threshold that achieved scale-free topology

($R^2 \geq 0.85$) across all datasets. (B-E) Sample dendrograms and trait heatmaps showing

hierarchical clustering of genes and resulting modules for (B) AD-Matarin, (C)

EP-KA, (D) AD-Castanho, and (E) EP-PO datasets. Upper dendrograms show gene

clustering, while lower heatmaps display module-trait relationships, with red

indicating positive correlations and blue indicating negative correlations with disease

status.

Figure S2. Dendrogram of the consensus network

Hierarchical clustering dendrogram of the consensus network constructed by

integrating topological overlap matrices from all four datasets (EP-PO, EP-KA,

AD-Matarin, AD-Castanho). The color band beneath shows consensus module

assignments, while the heatmap displays module-trait associations across multiple

model and disease conditions and age variables. This consensus approach identified

robust co-expression patterns conserved across both AD and EP models.

Figure S3. Construction of co-expression module in EP patient datasets.

Dendrogram of gene modules identified in the EP patient dataset using hierarchical

clustering and dynamic tree cutting based on TOM. (parameters: Soft-thresholding

power of 7 ensuring a scale-free topology ($R^2 \geq 0.85$), deepSplit = 1, minClusterSize =

28).

Figure S4. ROC curves of diagnostic models using shared hub gene expression.

ROC curves for classifiers trained on (A) AD patient dataset (GSE48350) and (B) EP

patient dataset (E-MTAB-3123). Multiple machine learning classifiers, including

Logistic Regression, Support Vector Machine (SVM), Random Forest, Naive Bayes,

XGBoost, AdaBoost, and CatBoost, were evaluated using 5-fold stratified

cross-validation. Models were trained using standardized expression values of all

consensus hub genes.

Figure S5: Forest plot of Mendelian randomization result from DMRdb.

Forest plot showing hazard ratios (HR) and 95% confidence intervals (CI) for

associations between Alzheimer's disease subtypes, epilepsy, and *C1QA*

http://www.inbirg.com/DMRdb/. Red dots represent point estimates for hazard ratios,

and horizontal lines indicate 95% confidence intervals. p-values are shown for each

association. Values greater than 1.0 indicate increased risk for the outcome.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets analyzed during the current study are available in the National Center for

Biotechnology Information (NCBI), European Nucleotide Archive (ENA), and

Synapse under the following accession numbers: PRJEB18790, GSE125957,

GSE73878, GSE64398, GSE63808, E-MTAB-3123, GSE48350, GSE190453,

syn51758062.1.

Competing Interests

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Funding

This research was supported by the Young Scientists Fund of the National Natural

Science Foundation of China (Grant Nos. 82201629) and the Guangdong Provincial

Science and Technology Special Fund (Grant Nos. 210803136872196)

Authors' contribution

HL conceptualized the study, conducted the investigation, developed the methodology,

performed formal analysis, and contributed to writing the original draft and

reviewing/editing the manuscript. ZX, YT, YY, and BL contributed to the

investigation, performed formal analysis, and participated in writing the original draft

and reviewing/editing the manuscript. SC JL contributed to writing the original draft

and reviewing/editing the manuscript. JW was involved in conceptualization and

contributed to writing the original draft and reviewing/editing the manuscript. MC,

XL, YS, ZD, and MH contributed to the investigation and participated in

reviewing/editing the manuscript. BH and NW conceptualized the study and

contributed to writing the original draft and reviewing/editing the manuscript. XJ

provided supervision, acquired funding, contributed to conceptualization, and

participated in writing the original draft and reviewing/editing the manuscript. All

authors read and approved the final manuscript.

Acknowledgments

The results published here are in whole or in part based on data obtained from the **AD**

Knowledge Portal

Reference

1. Vossel, K. A. *et al.* Seizures and Epileptiform Activity in the Early Stages of

Alzheimer Disease. *JAMA Neurol.* **70**, 1158–1166 (2013).

2.Palop, J. J. & Mucke, L. Epilepsy and Cognitive Impairments in Alzheimer Disease.

Arch. Neurol. **66**, 435–440 (2009).

3. Fang, Y. *et al.* Alzheimer Disease and Epilepsy: A Mendelian Randomization Study. *Neurology* **101**, (2023).

4. Prince, M. *et al.* World Alzheimer Report 2015. The Global Impact of Dementia: An analysis of prevalence, incidence, cost and trends. (Alzheimer's Disease International, 2015).

5. Ngugi, A. K., Bottomley, C., Kleinschmidt, I., Sander, J. W. & Newton, C. R. Estimation of the burden of active and life-time epilepsy: A meta-analytic approach. *Epilepsia* **51**, 883–890 (2010).

6. Mendez, M. F. & Lim, G. T. H. Seizures in Elderly Patients with Dementia. *Drugs Aging* **20**, 791–803 (2003).

7. Leitner, D. *et al.* Similar brain proteomic signatures in Alzheimer's disease and epilepsy. *Acta Neuropathol. (Berl.)* **147**, 27 (2024).

8. Pandis, D. & Scarmeas, N. Seizures in Alzheimer Disease: Clinical and Epidemiological Data: Seizures in Alzheimer Disease. *Epilepsy Curr.* **12**, 184–187 (2012).

9. Tombini, M. *et al.* Temporal Lobe Epilepsy and Alzheimer's Disease: From Preclinical to Clinical Evidence of a Strong Association. *J. Alzheimers Dis. Rep.* **5**, 243–261 (2021).

10. Johnson, E. L. *et al.* Association Between Midlife Risk Factors and Late-Onset Epilepsy: Results From the Atherosclerosis Risk in Communities Study. *JAMA Neurol.* **75**, 1375–1382 (2018).

11. Stefanidou, M. *et al.* Bi-directional association between epilepsy and dementia.

Neurology **95**, e3241–e3247 (2020).

12. Noebels, J. A perfect storm: Converging paths of epilepsy and Alzheimer's

dementia intersect in the hippocampal formation. *Epilepsia* **52**, 39–46 (2011).

13. Gourmaud, S. *et al.* Alzheimer-like amyloid and tau alterations associated with

cognitive deficit in temporal lobe epilepsy. *Brain* **143**, 191–209 (2020).

14. Tai, X. Y. *et al.* Hyperphosphorylated tau in patients with refractory epilepsy

correlates with cognitive decline: a study of temporal lobe resections. *Brain* **139**,

2441–2455 (2016).

15. Nelson, P. T. *et al.* Hippocampal sclerosis in advanced age: clinical and

pathological features. *Brain* **134**, 1506–1518 (2011).

16. Watts, M. E., Pocock, R. & Claudianos, C. Brain energy and oxygen metabolism:

Emerging role in normal function and disease. *Front. Mol. Neurosci.* **11**, 396508

(2018).

17. Zou, Y. *et al.* Epileptic seizures induced by pentylenetetrazole kindling accelerate

Alzheimer-like neuropathology in 5×FAD mice. *Front. Pharmacol.* **15**, 1500105

(2024).

18. Born, H. A. Seizures in Alzheimer's disease. *Neuroscience* **286**, 251–263 (2015).

19. Toniolo, S., Sen, A. & Husain, M. Modulation of Brain Hyperexcitability:

Potential New Therapeutic Approaches in Alzheimer's Disease. *Int. J. Mol. Sci.*

21, 9318 (2020).

20. Palop, J. J. *et al.* Aberrant Excitatory Neuronal Activity and Compensatory

Remodeling of Inhibitory Hippocampal Circuits in Mouse Models of Alzheimer's

Disease. *Neuron* **55**, 697–711 (2007).

21. Toniolo, S., Sen, A. & Husain, M. Modulation of Brain Hyperexcitability:

Potential New Therapeutic Approaches in Alzheimer's Disease. *Int. J. Mol. Sci.*

2020 Vol 21 Page 9318 **21**, 9318 (2020).

22. Targa Dias Anastacio, H., Matosin, N. & Ooi, L. Neuronal hyperexcitability in

Alzheimer's disease: what are the drivers behind this aberrant phenotype? *Transl.*

Psychiatry **12**, 1–14 (2022).

23. Liew, Y. *et al.* Neuroinflammation: A Common Pathway in Alzheimer's Disease

and Epilepsy. *J. Alzheimer's Dis.* **94**, S253–S265 (2023).

24. Kalyvas, A.-C., Dimitriou, M., Ioannidis, P., Grigoriadis, N. & Afrantou, T.

Alzheimer's Disease and Epilepsy: Exploring Shared Pathways and Promising

Biomarkers for Future Treatments. *J. Clin. Med.* **13**, 3879 (2024).

25. Giorgi, F. S., Saccaro, L. F., Busceti, C. L., Biagioni, F. & Fornai, F. Epilepsy and

Alzheimer's Disease: Potential mechanisms for an association. *Brain Res. Bull.*

160, 107–120 (2020).

26. Bosco, F. *et al.* Antiseizure Medications in Alzheimer's Disease from Preclinical

to Clinical Evidence. *Int. J. Mol. Sci.* **24**, 12639 (2023).

27. Srivastava, P. K. *et al.* Genome-wide analysis of differential RNA editing in

epilepsy. *Genome Res.* **27**, 440–450 (2017).

28. Matarin, M. *et al.* A Genome-wide Gene-Expression Analysis and Database in

Transgenic Mice during Development of Amyloid or Tau Pathology. *Cell Rep.* **10**,

633–644 (2015).

29. Castanho, I. *et al.* Transcriptional Signatures of Tau and Amyloid Neuropathology.

Cell Rep. **30**, 2040-2054.e5 (2020).

30. Johnson, M. R. *et al.* Systems genetics identifies Sestrin 3 as a regulator of a

proconvulsant gene network in human epileptic hippocampus. *Nat. Commun.* **6**,

6031 (2015).

31. Wan, Y.-W. *et al.* Meta-Analysis of the Alzheimer's Disease Human Brain

Transcriptome and Functional Dissection in Mouse Models. *Cell Rep.* **32**, (2020).

32. Chen, Z.-P. *et al.* Lipid-accumulated reactive astrocytes promote disease

progression in epilepsy. *Nat. Neurosci.* **26**, 542–554 (2023).

33. Mathys, H. *et al.* Single-cell transcriptomic analysis of Alzheimer's disease.

Nature **570**, 332–337 (2019).

34. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

35. Deng, Z.-L., Münch, P. C., Mreches, R. & McHardy, A. C. Rapid and accurate

identification of ribosomal RNA sequences via deep learning. *Nucleic Acids Res.*

50, e60 (2022).

36. Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based

genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat.*

Biotechnol. **37**, 907–915 (2019).

37. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. *Bioinformatics* **30**,

923–930 (2014).

38. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

39. Ritchie, M. E. *et al.* limma powers differential expression analyses for

RNA-sequencing and microarray studies. *Nucleic Acids Res.* **43**, e47 (2015).

40. Miller, J. A., Horvath, S. & Geschwind, D. H. Divergence of human and mouse

brain transcriptome highlights Alzheimer disease pathways. *Proc. Natl. Acad. Sci.*

107, 12698–12703 (2010).

41. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation

network analysis. *BMC Bioinformatics* **9**, 559 (2008).

42. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics

data. *The Innovation* **2**, (2021).

43. Langfelder, P., Luo, R., Oldham, M. C. & Horvath, S. Is My Network Module

Preserved and Reproducible? *PLOS Comput. Biol.* **7**, e1001057 (2011).

44. Szklarczyk, D. *et al.* The STRING database in 2025: protein networks with

directionality of regulation. *Nucleic Acids Res.* **53**, D730–D737 (2025).

45. Huynh-Thu, V. A., Irrthum, A., Wehenkel, L. & Geurts, P. Inferring Regulatory

Networks from Expression Data Using Tree-Based Methods. *PLOS ONE* **5**,

e12776 (2010).

46. Newman, A. M. *et al.* Robust enumeration of cell subsets from tissue expression profiles. *Nat. Methods* **12**, 453–457 (2015).

47. Sutton, G. J. *et al.* Comprehensive evaluation of deconvolution methods for human brain gene expression. *Nat. Commun.* **13**, 1358 (2022).

48. Dictionary learning for integrative, multimodal and scalable single-cell analysis | Nature Biotechnology. https://www.nature.com/articles/s41587-023-01767-y.

49. Izhikevich, E. M. Simple model of spiking neurons. *IEEE Trans. Neural Netw.* **14**, 1569–1572 (2003).

50. Sammons, R. P. *et al.* Structure and function of the hippocampal CA3 module. *Proc. Natl. Acad. Sci.* **121**, e2312281120 (2024).

51. Vyleta, N. P., Borges-Merjane, C. & Jonas, P. Plasticity-dependent, full detonation at hippocampal mossy fiber–CA3 pyramidal neuron synapses. *eLife* **5**, e17977 (2016).

52. Stimberg, M., Brette, R. & Goodman, D. F. Brian 2, an intuitive and efficient neural simulator. *eLife* **8**, e47314 (2019).

53. Liu, Y., Grigorovsky, V. & Bardakjian, B. Excitation and Inhibition Balance Underlying Epileptiform Activity. *IEEE Trans. Biomed. Eng.* **67**, 2473–2481 (2020).

54. Penn, Y., Segal, M. & Moses, E. Network synchronization in hippocampal neurons. *Proc. Natl. Acad. Sci.* **113**, 3341–3346 (2016).

55. Liew, Y. *et al.* Neuroinflammation: A Common Pathway in Alzheimer's Disease

and Epilepsy. *J. Alzheimers Dis.* **94**, S253–S265 (2023).

56. Vezzani, A., French, J., Bartfai, T. & Baram, T. Z. The role of inflammation in

epilepsy. *Nat. Rev. Neurol.* **7**, 31–40 (2011).

57. Heneka, M. T., Golenbock, D. T. & Latz, E. Innate immunity in Alzheimer's

disease. *Nat. Immunol.* **16**, 229–236 (2015).

58. Pekny, M. & Pekna, M. Astrocyte Reactivity and Reactive Astrogliosis: Costs and

Benefits. *Physiol. Rev.* **94**, 1077–1098 (2014).

59. Chen, Z.-P. *et al.* Lipid-accumulated reactive astrocytes promote disease

progression in epilepsy. *Nat. Neurosci.* **26**, 542–554 (2023).

60. Onyango, I. G., Jauregui, G. V., Čarná, M., Bennett, J. P. & Stokin, G. B.

Neuroinflammation in Alzheimer's Disease. *Biomedicines* **9**, 524 (2021).

61. Vezzani, A. & Viviani, B. Neuromodulatory properties of inflammatory cytokines

and their impact on neuronal excitability. *Neuropharmacology* **96**, 70–82 (2015).

62. van Vliet, E. A., Aronica, E. & Gorter, J. A. Blood–brain barrier dysfunction,

seizures and epilepsy. *Semin. Cell Dev. Biol.* **38**, 26–34 (2015).

63. Zlokovic, B. V. Neurovascular pathways to neurodegeneration in Alzheimer's

disease and other disorders. *Nat. Rev. Neurosci.* **12**, 723–738 (2011).

64. Kopczynska, M. *et al.* Complement system biomarkers in epilepsy. *Seizure* **60**,

1–7 (2018).

65. Veerhuis, R., Nielsen, H. M. & Tenner, A. J. Complement in the brain. *Mol.*

Immunol. **48**, 1592–1603 (2011).

66. Haure-Mirande, J.-V., Audrain, M., Ehrlich, M. E. & Gandy, S. Microglial TYROBP/DAP12 in Alzheimer's disease: Transduction of physiological and pathological signals across TREM2. *Mol. Neurodegener.* **17**, 55 (2022).

67. Haure-Mirande, J.-V. *et al.* Integrative approach to sporadic Alzheimer's disease: deficiency of TYROBP in cerebral Aβ amyloidosis mouse normalizes clinical phenotype and complement subnetwork molecular pathology without reducing Aβ burden. *Mol. Psychiatry* **24**, 431–446 (2019).

68. Haure-Mirande, J.-V. *et al.* Deficiency of TYROBP, an adapter protein for TREM2 and CR3 receptors, is neuroprotective in a mouse model of early Alzheimer's pathology. *Acta Neuropathol. (Berl.)* **134**, 769–788 (2017).

69. Audrain, M. *et al.* Integrative approach to sporadic Alzheimer's disease: deficiency of TYROBP in a tauopathy mouse model reduces C1q and normalizes clinical phenotype while increasing spread and state of phosphorylation of tau. *Mol. Psychiatry* **24**, 1383–1397 (2019).

70. Roussos, P. *et al.* The triggering receptor expressed on myeloid cells 2 (*TREM2*) is associated with enhanced inflammation, neuropathological lesions and increased risk for Alzheimer's dementia. *Alzheimers Dement.* **11**, 1163–1170 (2015).

71. Ma, J., Jiang, T., Tan, L. & Yu, J.-T. TYROBP in Alzheimer's Disease. *Mol. Neurobiol.* **51**, 820–826 (2015).

72. Jonsson, T. *et al.* Variant of TREM2 Associated with the Risk of Alzheimer's

Disease. *N. Engl. J. Med.* **368**, 107–116 (2013).

73. Stevens, B. *et al.* The Classical Complement Cascade Mediates CNS Synapse

Elimination. *Cell* **131**, 1164–1178 (2007).

74. Wu, X. *et al.* Complement C1q drives microglia-dependent synaptic loss and

cognitive impairments in a mouse model of lipopolysaccharide-induced

neuroinflammation. *Neuropharmacology* **237**, 109646 (2023).

75. Neniskyte, U. & Gross, C. T. Errant gardeners: glial-cell-dependent synaptic

pruning and neurodevelopmental disorders. *Nat. Rev. Neurosci.* **18**, 658–670

(2017).

76. Hu, Y. *et al.* Wogonin mitigates microglia-mediated synaptic over-pruning and

cognitive impairment following epilepsy. *Phytomedicine* **135**, 156222 (2024).

77. Bie, B., Wu, J., Foss, J. F. & Naguib, M. Activation of mGluR1 Mediates

C1q-Dependent Microglial Phagocytosis of Glutamatergic Synapses in

Alzheimer's Rodent Models. *Mol. Neurobiol.* **56**, 5568–5585 (2019).

78. Wu, J., Bie, B., Foss, J. F. & Naguib, M. Amyloid Fibril–Induced Astrocytic

Glutamate Transporter Disruption Contributes to Complement C1q-Mediated

Microglial Pruning of Glutamatergic Synapses. *Mol. Neurobiol.* **57**, 2290–2300

(2020).

79. Andoh, M., Ikegaya, Y. & Koyama, R. Synaptic Pruning by Microglia in Epilepsy.

J. Clin. Med. **8**, 2170 (2019).

80. Yuan, P. *et al.* PLD3 affects axonal spheroids and network defects in Alzheimer's

disease. *Nature* **612**, 328–337 (2022).

81. Stefanescu, R. A., Shivakeshavan, R. G. & Talathi, S. S. Computational models of

epilepsy. *Seizure* **21**, 748–759 (2012).

82. Luo, L. Architectures of neuronal circuits. *Science* **373**, eabg7285 (2021).

83. Neri, S., Mastroianni, G., Gardella, E., Aguglia, U. & Rubboli, G. Epilepsy in

neurodegenerative diseases. *Epileptic. Disord.* **24**, 249–273 (2022).

84. Zheng, X. *et al.* DMRdb: a disease-centric Mendelian randomization database for

systematically assessing causal relationships of diseases with genes, proteins,

CpG sites, metabolites and other diseases. *Nucleic Acids Res.* **53**, D1363–D1371

(2025).

85. Dejanovic, B. *et al.* Complement C1q-dependent excitatory and inhibitory

synapse elimination by astrocytes and microglia in Alzheimer's disease mouse

models. *Nat. Aging* **2**, 837–850 (2022).

86. Fan, J. *et al.* Preferential pruning of inhibitory synapses by microglia contributes

to alteration of the balance between excitatory and inhibitory synapses in the

hippocampus in temporal lobe epilepsy. *CNS Neurosci. Ther.* **29**, 2884–2900

(2023).

87. Holden, S. S. *et al.* Complement factor C1q mediates sleep spindle loss and

epileptic spikes after mild brain injury. *Science* **373**, eabj2685 (2021).

88. Chu, Y. *et al.* Enhanced synaptic connectivity and epilepsy in C1q knockout mice.

Proc. Natl. Acad. Sci. **107**, 7975–7980 (2010).

89. Ma, Y., Ramachandran, A., Ford, N., Parada, I. & Prince, D. A. Remodeling of

dendrites and spines in the C1q knockout model of genetic epilepsy. *Epilepsia* **54**,

1232–1239 (2013).

90. Aloi, M. S. *et al.* Microglia specific deletion of miR-155 in Alzheimer's disease

mouse models reduces amyloid-β pathology but causes hyperexcitability and

seizures. *J. Neuroinflammation* **20**, 60 (2023).

1165

Figure 1

Figure 2

Figure 3

Figure 4