Shared Transcriptomic Signatures Reveal Synaptic Pruning as a Link Between Alzheimer's Disease and Epilepsy

Huihong Li[1,2,3,4], Zhiran Xie[4], Yuxuan Tian[4], Bingying Lin[1,2,3,5], Jianwei Li[1,2,3,5], Jie Wu[1,2,3], Mingjie Chen[6], Xueke Liu[6], Yushan Sun[6], Si Chen[1,2,3,5], Zihan Deng[1,2,3,5], Mengqi Hong[1,2,3,5], Bing Huang[1,2,3,5], Naili Wei[1], Xiaoyu Ji[1,2,3,5*]

[1]Department of Neurosurgery, The First Affiliated Hospital of Shantou University Medical College, Shantou, Guangdong 515041, China

[2]Brain Function and Disease Laboratory, Shantou University Medical College, Shantou, Guangdong 515041, China

[3]Department of Pharmacology, Shantou University Medical College, Shantou, Guangdong, 51504, China

[4]Program in Computational Biology and Biomedical Informatics, Yale University, New Haven, Connecticut, United States of America

[5]Pain and Related Diseases Research Laboratory, Shantou University Medical College, Shantou, Guangdong 515041, China

[6]Department of Biotechnology and Food Engineering, Guangdong Technion – Israel Institute of Technology, Shantou, China 515063

*Corresponding authors

Contact: Dr. Xiaoyu Ji, xyji@stu.edu.cn

Abstract

Alzheimer's disease (AD) and epilepsy (EP) share a complex, bidirectional relationship, yet the molecular basis of their comorbidity remains unclear. To address this, we analyzed publicly available large-scale transcriptomic datasets from pilocarpine-induced EP mouse models (n = 200), two AD mouse models expressing human tau (rTg4510) or amyloid precursor protein (J20) (n = 141), and transcriptomic profiles from AD and EP patient cohorts. Differential expression and weighted gene co-expression network analyses revealed a highly conserved immune-related module enriched in microglia-specific genes, centered around *Tyrobp*, *Trem2*, and *C1q* complement component genes (*C1qa*, *C1qb*, *C1qc*) involved in the synaptic pruning pathway. Gene regulatory network analysis showed that *Tyrobp* regulates these *C1q* components gene through *Trem2*. These hub genes were markedly upregulated in both models and human patient datasets, preserved their regulatory relationship across species, and displayed strong diagnostic value. Cell composition deconvolution and single-cell transcriptomics confirmed their high, cell-type-specific expression in microglia from both animal model and human patient datasets. Moreover, computational modeling indicated that excessive synaptic pruning, particularly of inhibitory synapses, alters the excitation/inhibition balance toward excitability that increases neural network synchrony. Collectively, our findings reveal how microglial complement-driven synapse elimination may promote both seizure activity and cognitive decline. We suggest that microglia play a central role in the comorbidity of

43 EP and AD, and further exploration of this synaptic pruning pathway in the context of

44 EP may provide promising insights into the treatment of AD and EP comorbidity.

46 **Keywords:** Alzheimer's disease, epilepsy, synaptic pruning, computational

47 neuroscience, modeling, transcriptome, *Tyrobp, Trem2, C1q*

49 **Graphic abstract**

Highlights

- Transcriptomic analysis across epilepsy (EP) and Alzheimer's disease (AD) models and patient datasets revealed a conserved immune gene module.

- Shared transcriptomic signatures identified in animal models show strong diagnostic value in both AD and EP patient datasets.

- *Tyrobp-Trem2-C1q* microglial synaptic pruning pathway was identified as a central shared molecular mechanism across model and patient datasets.

- Computational modeling revealed that excessive synaptic pruning, particularly of inhibitory synapses, alters E/I balance toward excitability and increases synchrony in neural network.

1. Background

Alzheimer's disease (AD) and epilepsy (EP) are two common neurological diseases with a complex bidirectional relationship that contributes to their respective pathogenesis (1–3). AD, characterized by progressive cognitive decline and neurodegeneration is the leading cause of dementia globally, affecting an estimated 55 million people. This number is projectedto triple by 2050 as life expectancy increases (4). EP, characterized by recurrent unprovoked seizures, affects more than 50 million people worldwide with a particularly high incidence among elderly populations, especially those with neurodegenerative conditions such as AD (5,6).

Epidemiological studies have revealed a complex interplay between AD and EP. AD patients show increased susceptibility to EP, with 10–22% experiencing unprovoked seizures, while 22–54% demonstrating subclinical epileptiform activity during disease progression (7,8). These seizures, predominantly occurring in the temporal lobe, correlate with accelerated cognitive decline and enhanced AD pathology (9). Conversely, EP patients exhibit a two-fold higher risk of developing dementia compared to controls, with late-onset EP patients showing a three-fold increased risk (10,11). This elevated risk is particularly evident in temporal lobe epilepsy (TLE), where numerous pathological features overlap with AD (9).

Both diseases show overlapping pathological characteristics, especially in TLE,

including amyloid deposition, tau pathology, and hippocampal sclerosis (12,13).

Chronic seizures increase neuronal activity and metabolic demands, potentially

accelerating the production and aggregation of amyloid-β (aβ) and

hyperphosphorylated tau protein (14,15). Conversely, the progressive accumulation of

these pathological proteins in AD may lower seizure thresholds, promoting

hyperexcitability and epileptiform activity (16,17). Furthermore, neuroinflammation,

synaptic loss, and microglial activation, features of both AD and TLE, may

synergistically accelerate neuronal damage and network dysfunction (18,19),

potentially explaining the high comorbidity rates observed clinically.

The molecular mechanisms underlying this comorbidity remain incompletely

understood, though mounting evidence suggests shared molecular pathways driving

neuronal hyperexcitability and protein aggregation in both disorders (20). Recent

findings implicate dysregulated protein homeostasis, mitochondrial dysfunction, and

chronic inflammatory responses as central contributors (7,19,21). Additionally,

alterations in glutamatergic and GABAergic signaling, alongside disruptions in

synaptic plasticity pathways, appear to converge in both conditions, particularly

within the hippocampal region (22–24).

The treatment of EP in AD patients has been considered a complex clinical challenge

in terms of the decision to treat seizures and the choice of antiseizure medications (25).

Emerging therapeutic approaches, including targeted immunotherapies and novel

small molecules, show promise but require a deeper understanding of the shared

pathological mechanisms. Therefore, systems biology and computational modeling

approaches are urgently needed to integrate insights from animal models and patients.

Such approaches could reveal common molecular pathways, cellular processes, and

neural network dynamics alterations underlying this comorbidity, which is crucial for

developing more effective, targeted therapeutic strategies.

To address this gap, we hypothesize that conserved molecular pathways exist in AD

and EP animal models and human patients. To test this, we (1) analyzed and

integrated large-scale transcriptomic data from pilocarpine-induced EP mouse models

(n = 200), two AD mouse models expressing human tau (rTg4510, TG) or amyloid

precursor protein (J20) (n = 141) to identify shared transcriptional signatures and

pathways; (2) validated these findings in AD and EP human patient datasets; and (3)

developed computational models to link molecular insights to network mechanisms of

epileptogenesis. Our analysis identified a conserved synaptic pruning pathway

involving *Tyrobp*, *Trem2*, and *C1q* complement genes. Computational modeling

further suggested that preferential synaptic elimination of inhibitory synapses

contributes to neuronal network excitability and the cognitive decline associated with this comorbidity.

2. Materials and Methods

2.1 Data Collection

A comprehensive database search was conducted to retrieve transcriptomic data from AD and EP models as well as human patients. Databases including Gene Expression Omnibus (GEO) and European Nucleotide Archive (ENA), were searched using the keywords "epilepsy" and "Alzheimer's disease". We downloaded and reanalyzed two extensive bulk RNA-seq datasets from EP and AD mouse models. The EP model dataset was obtained from the ENA (PRJEB18790) and includes bulk RNA-seq data from 200 hippocampi of pilocarpine-induced EP model mice and wild-type (WT) control. The AD dataset was downloaded from GEO (GSE125957) and includes bulk RNA-seq data from 141 samples of the entorhinal cortex and hippocampus of AD model mice that expressed either human tau (rTg4510, TG) or amyloid precursor protein (J20), with their corresponding healthy controls at various ages (range from 2 to 12 months).

To assess the clinical relevance of insights derived from animal models at the gene, pathway, and cellular levels, we also retrieved and analyzed several large-scale

transcriptome datasets from AD and EP patients. For cross-species validation of

disease co-expression modules, we used 129 hippocampal samples from TLE patients

(GSE63808) and 97 middle temporal gyrus samples from AD patients (GSE132903).

To evaluate the diagnostic value of shared transcriptional signatures, we analyzed 47

hippocampal samples from EP patients and controls (E-MTAB-3123), along with 253

different brain region samples from AD patients and controls (GSE48350). Finally,

single-cell RNA sequencing (scRNA-Seq) data from 85,780 immune cells of six

drug-refractory EP patients (GSE201048) and single-nucleus RNA sequencing

(snRNA-Seq) data from 174,420 immune cells from 217 AD patients and 226 controls

(GSE227223) were employed to identify the cell types and subtypes expressing these

shared transcriptional signatures.

2.2 Quality control and preprocessing

All downloaded raw RNA sequences from AD and EP model datasets underwent

quality control and trimming to obtain clean sequences, using FastQC (0.12.1) and

Fastp (0.22.0) with stringent parameters (--qualified_quality_phred 30 --n_base_limit

5 --detect_adapter_for_pe --cut_tail --trim_front1 15 --dedup --trim_poly_x

--trim_poly_g) to ensure high data quality (26,27). Ribodetector (0.2.7) was used to

remove potential rRNA contamination (28). Low-quality regions in the sequences,

attributed to instability in the sequencing platform, were also trimmed using Fastp.

Trimmed sequences were then aligned to the GRCm39.111 reference mouse genome

using HISAT2 (2.2.1) (29) and followed by quantification using featureCounts (2.0.6)

(30).

For the AD model dataset, samples were excluded from subsequent analysis based on

the criteria outlined in the original publication, resulting in 121 high-quality RNA-seq

samples. For the EP model dataset, duplicate and low-quality samples were initially

removed. Then a variance-stabilizing transformation was applied to the count matrix

before performing Principal Component Analysis (PCA) to identify potential outlier

samples. PCA was then performed to identify outliers, with Mahalanobis distances

calculated from PCA scores. A chi-squared distribution threshold of 0.975 was used to

detect significant outliers, resulting in 11 samples subsequently excluded from

analysis. Detailed information on quality control and mapping metrics is provided in

Table S1. Following quality control, mapping, and exclusion of low-quality samples,

no significant differences in read depth were observed between model groups and

controls.

2.3 Differential Expression Analysis

Differential expression analysis was perform using DESeq2 (1.46.0) (31). Before

analysis, non-expressed and lowly expressed genes were filtered out. For the EP

model dataset, only genes with at least 20 counts in a minimum of 90 samples (~90

samples per group) were retained, while for the AD dataset, genes with at least 20

counts in at least 6 samples (6–8 samples per group) were kept. The DESeq function

was then used to estimate size factors and dispersions and to perform statistical testing

for differential expression (31).

For the EP model dataset, the design formula was specified as ~Condition (EP/WT),

and results were extracted by comparing the "EP" condition to the "WT" condition.

For the AD dataset, we aimed to detect progressive differentially expressed genes

(DEGs), following a previous approach (32), using DESeq's likelihood-ratio test to fit

the statistical model: *Gene expression = Genotype + Age + Genotype × Age*

(Interaction). p-values were adjusted for multiple testing using the

Benjamini-Hochberg (BH) correction.

DEGs were identified based on an adjusted p-value < 0.05 and |log2 fold change| > 1.

These thresholds were applied to the EP model dataset and to the final time point of

the AD dataset. Shared DEGs between the EP and AD model datasets were identified

by taking the intersection of their DEG results. Kyoto Encyclopedia of Genes and

Genomes (KEGG) pathway and gene ontology (GO) analyses for shared DEGs were

performed using WebGestalt (2024), with weighted set cover redundancy removal

(33).

2.4 Weighted Gene Co-expression Network Construction

To construct weighted gene co-expression networks and compare results from EP and

AD model datasets, we employed a standard meta-WGCNA comparison method

(34–36). All datasets underwent variance-stabilizing transformation using DESeq2

(1.46.0) to normalize the variance across different samples. We then calculated the

median absolute deviation (MAD) and retained the top 85% of genes with the highest

MAD in each dataset, resulting in 10,997 genes common to all datasets for

comparability in further analyses. Outliers were identified through hierarchical

clustering, excluding two samples from the rTg4510 dataset and three from the J20

dataset. To assess dataset comparability, general network properties such as average

gene expression and overall network connectivity were calculated.

We constructed signed co-expression networks using the WGCNA R package (1.73)

(35), selecting the lowest soft-thresholding power to achieve a scale-free topology fit

index (R^2) of at least 0.85 across datasets. Adjacency matrices were generated using

the signed network option and converted to topological overlap matrices (TOM) to

quantify similarity in gene connectivity profiles. Modules were identified by average

linkage hierarchical clustering of TOM dissimilarity (1-TOM), followed by dynamic

tree cutting with parameters deepSplit = 2 and minClusterSize = 24. This procedure

was also applied to patient microarray datasets (GSE63808 and GSE132903), using

variance-stabilizing normalization with the "limma" (3.62.1) package (37).

To match modules between EP and AD model datasets, we color-mapped the AD modules to the best matching EP modules according to the hypergeometric test (34). Module profiles were summarized using module eigengenes (MEs), derived from the first principal component of each module. To identify modules significantly associated with disease genotypes (EP or interaction of TG/J20), we calculated Pearson correlation coefficients between MEs and genotype information, with significance defined as $p < 0.05$ and an absolute correlation coefficient > 0.65 (38). Module membership (MM) and gene-genotype significance (GS) were also determined by calculating Pearson correlations between individual genes and MEs or genotypes, respectively. KEGG pathway and GO analyses for module annotation were conducted using WebGestalt (2024) with weighted set cover redundancy removal (33).

2.5 Module Preservation Analysis

To assess module preservation of the EP model module in AD model datasets, we used two approaches (39): Zsummary preservation statistics and cross-tabulation-based preservation statistics.

We employed the modulePreservation function from the WGCNA R package, with parameters set to networkType = "signed", nPermutations = 100, to calculate composite Z-summary preservation statistics. This metric evaluates preservation

based on module size, network density, and connectivity, offering a comprehensive

assessment of how well the structure of EP modules is maintained in AD datasets. The

Z-summary thresholds were interpreted as follows: Z-summary > 10 indicates strong

evidence of preservation, 2 < Z-summary < 10 suggests moderate evidence, and

Z-summary < 2 indicates no evidence of preservation (39). To complement the

Z-summary statistic, we also used the medianRank preservation statistic, which

assesses relative preservation across modules with less dependency on module size. A

lower median rank denotes stronger observed preservation (39).

Additionally, we used a cross-tabulation method to examine module overlap between

EP and AD model datasets, as well as between model and patient datasets (GSE63808

and GSE132903). This method generates a contingency table reporting the

overlapping gene number between each module in the reference and test datasets (39).

The statistical significance of overlaps was assessed using Fisher's exact test, with

p-values adjusted using the BH correction. A module was reported as significantly

overlapping if the overlap count was > 30 and the adjusted p-value was < 0.05.

2.6 Hub Gene Identification and Regulatory Network Analysis

Hub genes act as key drivers within a module and were identified using stringent

criteria to minimize false positives: genes ranking in the top 10% for intramodular

connectivity within each module, MM > 0.9, and GS > 0.4. Shared hub genes were

determined by intersecting the hub genes of key modules in both EP and AD datasets,

along with the DEGs. The expression and diagnostic value of these shared hub genes

were further explored in the AD and EP patient datasets (GSE48350 and

E-MTAB-3123).

To further explore interactions and potential regulatory relationships among shared

hub genes in the EP and AD model datasets, we constructed a protein-protein

interaction (PPI) network using the STRING database (12.0)

(https://string-db.org/cgi/input.pl) with a confidence score threshold of 0.4 and a false

discovery rate (FDR) of 5% (40). We then performed regulatory network analysis

with the GENIE3 (1.28.0) algorithm, which applies a random forest approach to infer

potential regulatory relationships from gene expression data. GENIE3 ranks

regulatory interactions by weight, with higher weights indicating stronger regulatory

links (41). Gene expression profiles of the key module from the EP and TG model

datasets were used to infer plausible regulatory interactions among shared hub genes.

To reduce false positives, a weight threshold of 0.01 was applied to filter out weaker

regulatory links. Additionally, only the top 30 plausible regulatory links between

shared hub genes were retained to construct gene regulatory networks. Finally, the

shared regulatory network of hub genes was constructed by intersecting the regulatory

networks inferred from the EP and TG model datasets.

2.7 Machine Learning

To validate the diagnostic value of the transcriptomic signatures identified in animal models, multiple machine-learning models were used to train diagnostic models based on these signatures in EP and AD patient datasets (GSE48350 and E-MTAB-3123) to predict disease status based on gene expression profiles. The main goal was to distinguish patients with AD or EP from healthy controls. First, the gene expression data was standardized by applying the StandardScaler function in the sklearn.preprocessing module in Python. Standardization ensures that all features/gene signatures have a mean of 0 and a standard deviation of 1, making them comparable across scales. Several machine learning algorithms were used in the classification process, including logistic regression, support vector machine (SVM), random forest, naive Bayes, XGBoost, AdaBoost, and CatBoost. To evaluate the performance of each model, a 5-fold stratified cross-validation method was used. Different metrics were used to measure the performance of the models, including accuracy, precision, recall, F1 score, and AUC. The ROC curve and AUC score for each classifier were plotted to visually examine the trade-off between sensitivity and specificity.

2.8 Cell Composition Deconvolution

In silico deconvolution can estimate the cellular composition of a tissue sample from its gene expression profile, with CIBERSORT particularly suited for cellular

composition deconvolution of brain transcriptomes (42,43). The raw count expression

matrix was converted to FPKM values using edgeR (4.4.1) (44), followed by cell

composition deconvolution using CIBERSORT (1.04) via the BrainDeconvShiny

platform, with cell-type signatures specific to *Mus musculus* from immunopurified

mouse brain tissue (43,45).　For statistical analysis, the Mann-Whitney test ($p < 0.05$)

was applied to compare the EP dataset with the control group, while the

Kruskal-Wallis test was used for the AD datasets, followed by pairwise comparisons

across age and genotype groups. The BH method was used to control the FDR ($q <$

0.05). Associations between identified cell types and shared hub genes were

determined by calculating Pearson correlation coefficients.

2.9 Single-cell and Single-nucleus RNA Sequencing Data Analysis

To localize and validate the expression of transcriptomic signatures identified in

animal models at the cellular level in AD and EP patients, count matrices of

sc/snRNA-seq (GSE201048 and GSE227223) were downloaded and processed

according to the methods outlined in the original publications (46–48). Quality control

was performed by filtering cells with high mitochondrial content (>20% for the EP

dataset and >5% for the AD dataset) and selecting cells based on cellular complexity

(300–5,000 genes per cell for the EP dataset and >500 genes for the AD dataset).

Subsequent data processing was conducted using the standard Seurat pipeline,

including scaling, normalization, variable gene identification, and clustering. PCA

was performed on the 2,000 most variable genes, with the first 20 principal

components (PCs) used for t-SNE and clustering in the EP dataset, and the first 30

PCs used for the AD dataset. Cluster identities were determined by identifying DEGs

with the FindMarkers function, followed by annotation based on the original

publications.

2.10 Computational modelling

To explore the contribution of the identified key pathways to pathological changes in

AD and EP at the network level, we implemented a spiking neural network using the

Izhikevich neuron model to simulate hippocampal CA3 circuit dynamics with

simulation spans of 1000ms, employing the Brian2 package (49). The network

consists of 1000 Izhikevich neurons (800 excitatory and 200 inhibitory), with

dynamics governed by (50):

$$\frac{dv}{dt} = 0.04v^2 + 5v + 140 - u + I$$
$$\frac{du}{dt} = a(bv - u)$$

where v is the membrane potential, u is the recovery variable. When v≥30□mV, the

following reset conditions are applied:$v = c;\ u = u + d$. Parameters a, b, c and d

are set to standard regular spiking for excitatory neurons and fast-spiking for

inhibitory neurons following the Izhikevich framework (50). I is external input

incorporating both suprathreshold Poisson noise at 1 Hz and presynaptic inputs (50).

Following Liu's EP network modeling, synaptic connectivity is sparse, with a 0.1

probability of random connections between neurons, and interactions are modeled

using conductance-based synapses to allow for voltage-dependent temporal and

spatial summation (50). Synaptic interactions simulate AMPA (excitatory) and

GABA-a (inhibitory) currents for four connection types: excitatory-excitatory (E→E),

excitatory-inhibitory (E→I), inhibitory-excitatory (I→E), and inhibitory-inhibitory

(I→I) (50). Synaptic currents are calculated as (51):

$$I_j = w S_{i \to j}(t)\left(E_i - v_j\right)$$

we introduce w to leverage the synaptic weight, while $S_{i \to j}(t))$ is the synaptic

conductance evolving as $\frac{dS_{i \to j}(t)}{dt} = -\frac{S_{i \to j}(t)}{\tau(i)}$ and E_i is the synaptic reversal potential

(50). Synaptic conductance is updated upon presynaptic spikes by adding a peak

conductance $S_{i \to j}(max)$, while we calibrate unitary synapse to induce EPSPs for

E→E/E→I and IPSPs for I→E/I→I of 1.0, 2.5, −2.0, −1.0 mV with time

constants: $\tau = \{3, 3, 5, 5\}$ ms and reversal potentials: $E = \{0, 0, -75, -75\}$ mV

(52–54). Each of four types of synaptic weights (w) are drawn independently from a

standard normal distribution, normalized to [0, 1], and set to 0 during synaptic

pruning. We simulated synaptic pruning by selectively removing excitatory (E→E and

E→I) or inhibitory (I→E and I→I) synapses, starting with those having the lowest

weights based on the synaptic weight distribution (ranging from 0% to 50%, with 5%

increments). Each pruning simulation was repeated at least 50 times.

The excitation/inhibition (E/I) ratio quantifies the balance between synaptic drives in

the network, indicating firing rate stability, while event synchrony measures temporal

coordination of firing patterns, implying network oscillation patterns. Both large E/I

ratio and excessive synchrony are associated with EP and AD (51).

To analyze the network, we calculated the E/I ratio and event synchrony to measure

the effects of synaptic pruning on firing rate stability and temporal coordination. The

E/I ratio compares total excitatory currents (from E→E and E→I synapses) to the

absolute magnitude of inhibitory currents (from I→E and I→I synapses), averaged

over the simulation period T (51):

$$E/IRatio = \frac{\sum_{i=1}^{N} I_i^E(t)}{\left| \sum_{i=1}^{N} I_i^I(t) \right|}$$

where:

$I_i^E(t)$ represents excitatory synaptic currents onto neuron i at time t

$I_i^I(t)$ represents inhibitory synaptic currents onto neuron i at time t

N is the total number of neurons

$\bar{x}$ represents the mean over time T

Event synchrony (ES) quantifies spike timing coordination between excitatory neuron

pairs by calculating the fraction of their spikes occurring within 5-ms windows. For

each excitatory neuron pair (i, j), synchronous events are counted (with exact

coincident spikes weighted by 0.5 and near-coincident spikes within the time window

weighted by 1.0), then normalized by the geometric mean of both neurons' total spike

counts (55). During total simulation time T:

$$ES_{ij} = \frac{\sum_{s_i \in S_i} \sum_{s_j \in S_j} C(s_i, s_j)}{\sqrt{N_i N_j}}$$

where:

Si is the set of all spike times from neuron i during the simulation.

Sj is the set of all spike times from neuron j during the simulation.

Ni and Nj are the total number of spikes from neuron i and j respectively over the

entire simulation period.

C(si, sj) counts synchronized events for each pair of spikes:

$$C(s_i, s_j) = \begin{cases} 1 & if \; 0 < |s_i - s_j| < 5ms \\ 0.5 & if \; s_i = s_j \\ 0 & otherwise \end{cases}$$

412 The network-wide event synchrony is computed as:

$$\text{Network ES} = \frac{1}{P} \sum_{i,j} ES_{ij}$$

413 Where $P = \frac{Ne \, (Ne-1)}{2}$ is the total number of unique excitatory neuron pairs, Ne is the

414 number of excitatory neurons.

Both E/I ratio and ES values are normalized to baseline unpruned values to highlight

pruning-induced changes in network dynamics. For statistical analysis, we conducted

a correlation analysis by calculating Pearson correlation coefficients and linear

regression analysis to quantify the effect sizes of excitatory and inhibitory pruning on

both E/I ratio and synchronization.

3. Results

3.1 Identification and Functional Enrichment of Shared DEGs

3.1.1 Differential Expression Analysis

To reveal the shared molecular signatures underlying EP and AD models, we

performed differential expression analysis using DESeq2. In the EP dataset, we

identified 378 DEGs, with 351 upregulated and 27 downregulated genes (**Fig. 1A**).

The TG dataset yielded 426 progressive DEGs, including 377 upregulated and 49

downregulated genes (**Fig. 1B**). In contrast, the J20 dataset contained only one

upregulated progressive DEG, Cst7 (**Fig. 1C**). Notably, 101 genes were differentially

expressed in both the EP and TG datasets, representing 26.72% and 23.71% of the

DEGs in each dataset, respectively (**Fig. 1A–B**). This substantial overlap suggests

shared molecular pathways between EP and TG models. Detailed DEGs information

is available in **Table S2**.

3.1.2 Functional Enrichment Analysis of Shared DEGs

To investigate the biological relevance of these shared DEGs between EP and TG

models, we conducted a functional enrichment analysis. KEGG pathway analysis

identified several most enriched pathways, including "complement and coagulation

cascades," "phagosome," and "cytokine-cytokine receptor interaction" (**Fig. 1D**). GO

enrichment analysis highlighted key biological processes, such as "defense response,"

"immune response," and "regulation of cytokine production" (**Fig. 1E**). Additionally,

enriched molecular functions included "signaling receptor binding" and "immune

receptor activity," with cellular components such as the "extracellular matrix" and

"side of membrane" also significantly represented. These findings indicate that the

shared DEGs are primarily involved in immune processes, reflecting shared molecular

pathways in both EP and TG models. Detailed GO enrichment analysis results can be

found in **Table S2**.

3.2 Construction of Weighted Gene Co-Expression Network

WGCNA was particularly suited for this study due to its ability to identify modules of

highly correlated genes and reveal underlying biological pathways conserved across

different conditions. By applying WGCNA to both EP and AD datasets, we aimed to

identify conserved gene modules that may represent common molecular mechanisms

between EP and AD.

After filtering out outlier samples, low-expression, and low-variance genes in each

model dataset, we retained 10,997 genes common to all model datasets for further

analysis (**Fig. S1A-C**). For network construction, we selected a soft-thresholding

power of 9 to ensure that networks in all model datasets exhibited scale-free topology

with a fit index > 0.85 (**Fig. S1D**). This threshold was consistently applied across

analyses to maintain network comparability. To assess comparability between EP and

AD model networks, we examined correlations of ranked mean gene expression and

ranked gene connectivity between dataset pairs. For EP and J20 model datasets, high

correlations were observed for both expression ($r = 0.98$, $p < 1e-200$) and connectivity

($r = 0.23$, $p < 5.7e-132$). Similarly, EP and TG model datasets showed significant

correlations (expression: $r = 0.96$, $p = 1e-200$; connectivity: $r = 0.43$, $p = 1e-200$) (**Fig.**

S1E), indicating comparability across model datasets (34).

We used the WGCNA package to construct gene co-expression networks for the EP,

TG, and J20 model datasets. The analysis identified diverse gene modules across the

three models: the EP dataset revealed 33 modules (28–790 genes per module) (**Fig.**

2A), TG had 29 modules (34–1,537 genes) (**Fig. 2B**), and J20 contained 32 modules

(27–613 genes) (**Fig. 2C**). Detailed module information can be found in **Table S3**.

Associations between module eigengenes (MEs) and model genotypes were assessed.

The EP model dataset identified seven modules significantly associated with the EP

genotype: three with positive correlations (green, steelblue, turquoise; $0.73 \leq r \leq 0.79$,

$p < 3e\text{-}31$) and four with negative associations (lightyellow, red, lightcyan,

paleturquoise; $-0.92 \leq r \leq -0.69$, $p < 8e\text{-}27$). Notably, five of these modules contained

DEGs linked to the EP genotype (turquoise: n = 198, green: n = 86, steelblue: n = 12,

lightcyan: n = 12, red: n = 5) (**Fig. 2A, D**). In the TG model dataset, seven modules

showed significant associations with the TG genotype: two with positive correlations

(turquoise, yellow; $0.72 \leq r \leq 0.87$, $p < 3e\text{-}10$) and five with negative associations

(brown, saddlebrown, darkorange2, darkmagenta, royalblue; $-0.83 \leq r \leq -0.67$, $p <$

$1e\text{-}08$). Nine modules contained DEGs associated with the TG genotype (turquoise: n

= 268, yellow: n = 14, green: n = 7, darkorange2: n = 4, blue: n = 2, orangered4: n = 2,

royalblue: n = 2, darkmagenta: n = 1, tan: n = 1) (**Fig. 2B, E**). Consistent with the

DEGs analysis that showed the J20 model dataset had fewer DEGs, none of the

modules exhibited significant associations with the J20 genotype ($r < 0.4$) (**Fig. 2C,**

F).

3.3 Module Preservation Analysis

Following module construction, we assessed the preservation of EP model modules in

the AD model datasets to reveal conserve molecular pathways between EP and AD

models. Our analysis revealed varying degrees of module preservation between EP

and AD model datasets (**Fig. 3A-B**). In the TG model, the steelblue module had low

preservation scores (Z-summary.pres = 0.89. Similarly, in the J20 model, the steelblue

and yellow modules showed weak preservation (Z-summary.pres < 4), suggesting

these may represent unique biological processes specific to the EP model dataset. In

contrast, several EP model modules, including green, blue, and brown, exhibited high

preservation scores and significant overlap with AD modules in both models,

suggesting they represent conserved biological processes. Preservation statistics are

shown in **Table S3**.

Notably, the turquoise EP model module had the highest preservation scores

(Z-summary.pres = 83.01) and showed strong overlap with the TG model turquoise

module, displaying the highest overlap counts and the lowest adjusted p-values (**Fig.**

3A, C). Both turquoise modules were positively correlated with disease genotype (EP:

r = 0.79, p = 5e-41; TG: r = 0.87, p = 4e-18) and contained most DEGs in each model

(**Fig. 2A, C**). Other EP modules, such as lightcyan, showed significant correlations

but did not display preservation. GO enrichment analysis of this model turquoise

module revealed enrichment in immune-related terms (**Table S3**). We further assessed

the preservation of the mouse turquoise module in EP and AD patient dataset

(GSE63808 and GSE132903), finding that the turquoise immune modules from EP

and TG models significantly overlapped with modules constructed from EP and AD

patient datasets, respectively (**Fig. 3E-F, S2**). This suggests the turquoise module

represents a conserved immune molecular signature across species and disease states,

suggesting that immune processes play a preserved and significant role in these

modules across conditions.

3.4 Shared Hub Gene Identification and Regulatory Network Analysis

To identify hub genes that may play a crucial role in both EP and AD, we intersected

the hub genes of the turquoise module from EP and TG model datasets with the

previously identified 101 DEGs. This analysis identified 19 shared hub genes

common to both conditions (two shared hub genes not shown in the figure due to their

ranking: *Apobec* and *Cxcl16*) (**Fig. 4A-B**). We then constructed a PPI network to

investigate the biological significance of these 19 genes and the results showed a PPI

network consisting of 14 genes **(Fig. 4C, Table S4)**. This PPI network revealed strong

connectivity among *C1q* complex component genes (*C1qa*, *C1qb*, *C1qc*) and other

immune-related genes, including *Trem2*, *Tyrobp*, and *Adgre1*. GO biological process

enrichment analysis highlighted the most significant processes involved in "Synaptic

pruning" (**Fig. 4D, Table S4**). Genes involved in "Synaptic pruning" (*C1qc*, *C1qb*,

C1qa, *Trem2*) also participate in other enriched processes, such as

"Complement-mediated synapse pruning" (*C1qa*, *Trem2*) and "Microglial cell

activation" (*Tyrobp*, *C1qa*, *Trem2*) (**Table S4**).

While WGCNA reveals gene co-occurrence patterns, it does not identify regulatory

mechanisms. To explain potential regulatory relationships among these 19 genes, we

applied the GENIE3 algorithm to gene expression profiles of the turquoise module in

both EP and TG conditions (**Table S5**). By intersecting predicted regulatory networks

from both conditions, we identified a shared regulatory pathway: *Tyrobp* regulates

Trem2, which in turn regulates the complement component genes *C1q* via *C1qa* (**Fig.**

4E). These genes, particularly *Tyrobp*, *Trem2*, and *C1q* complex components gene

(*C1qa*, *C1qb*, *C1qc*), play central roles in co-expression modules of both EP and TG

model and are involved in multiple enriched terms, especially synaptic pruning.

Notably, these genes are also progressive DEGs in the TG model (**Fig. 4F**). The

successful reconstruction of a regulatory network involving *Tyrobp*, *Trem2*, and *C1q*

component genes suggests they may represent a conserved molecular pathway

underlying both conditions.

3.5 Validation of the Diagnostic Value of Shared Hub Genes

To further validate the clinical relevance of the identified transcriptional signatures in

AD and EP animal models, we analyzed the expression of 19 shared hub genes in

datasets from AD and EP patients (GSE48350 and E-MTAB-3123) compared to

controls. Thirteen of these 19 genes were expressed in both datasets, with 10

exhibiting significantly higher expression levels ($p < 0.05$) compared to controls (**Fig.**

5A-B). Notably, genes such as *Tyrobp*, *Trem2*, and *C1q* complex component genes

demonstrated pronounced significance in both datasets (**Fig. 5A-B**). To assess their

diagnostic potential, we developed diagnostic models leveraging the expression

profiles of these genes in the hippocampus (**Fig. S2**). Using multiple machine learning

methods, these models effectively differentiated disease from control groups, with

particularly high performance in the EP dataset (AUC > 0.9) and satisfactory

performance in the AD patient dataset (AUC > 0.7) (**Fig. S3**). These findings confirm

the diagnostic value of the transcriptional signatures identified in animal models in

human AD and EP patients.

Given that these genes were shared hub genes of co-expression modules in EP and

AD models, likely reflecting conserved expression pathways and progressive changes,

we examined their co-expression patterns in AD and EP patient datasets (GSE48350

and E-MTAB-3123) by calculating Pearson correlation coefficients. *Tyrobp*, *Trem2*,

and the *C1q* component genes demonstrated strong correlations in both AD and EP

patient datasets (r > 0.85), whereas co-expression patterns among other genes were

less consistent across datasets (**Fig.5C-D**). Collectively, these findings suggested that

synaptic pruning pathway genes such as *Tyrobp*, *Trem2*, and the *C1q* component are

not only conserved between AD and EP animal models and human patients but also

likely play critical roles in the underlying pathology of these conditions.

3.6 Cellular Composition Alteration and Genes Localization in Cell Type

To investigate cellular composition changes and their associations with the key hub

genes (*Tyrobp*, *Trem2*, *C1q* complex genes) in TG and EP conditions, we employed

CIBERSORT for cell composition deconvolution of bulk RNA-seq data. Our analysis

revealed significant cellular composition alterations. Firstly, we observed a notable

reduction in neuronal composition in both EP and TG conditions compared to controls

(p < 0.05) (**Fig. 6A, C**). Secondly, there was a marked increase in microglial

composition in both models (p < 0.05), consistent with the enrichment of

immune-related processes identified in previous analyses (**Fig. 6A, C**). While

astrocyte and endothelial cell alterations were also detected in EP and TG models (at 8

months, p < 0.05), these changes were less pronounced than the neuronal decrease and

microglial increase observed in the TG model (**Fig. 6A, C**). No significant differences

were found in oligodendrocyte composition in either condition.

We examined the relationship between these cellular composition changes and the key

hub genes such as *Tyrobp*, *Trem2*, and *C1q* complex genes by conducting Pearson

correlation analyses (**Fig. 6B, D**). Notably, all key hub genes showed strong negative

correlations (r < -0.7) with neuronal composition in both EP and TG conditions,

suggesting an inverse relationship between their expression levels and neuronal

proportion. Conversely, strong positive correlations were observed between these

genes and microglial and endothelial cell composition (r > 0.7). Correlations with

astrocytes and oligodendrocytes were more variable and showed less consistency

across genes and conditions. Although no cellular composition changes were observed

in the J20 model (**Fig. S4A**), we found that genes such as *Tyrobp*, *Trem2*, and *C1q*

complex components, were strongly associated with microglial composition,

indicating that these genes demonstrate a strong link with microglial cell types (**Fig.**

S4B). We further localized the expression of these genes to cell types using the whole

mouse brain transcriptome cell type atlas from the Allen Brain Atlas (56). The results

showed that the majority (15 of 19 genes) were predominantly expressed in immune

cells, particularly microglia and macrophages (**Fig. S5**).

To further explore the expression of these genes in human patient cell types,

especially immune cells, we explored the expression of these genes in two large-scale

sc/sn RNAseq datasets of immune cells from AD and EP patients (GSE201048 and

GSE2272230) (**Fig. S6**). We found that 12 of the shared hub genes were detectable in

both datasets. Among these, *Tyrobp* and *Trem2*, along with the *C1q* complex

component genes, displayed notably high and predominant expression in microglia

compared to other shared hub genes (**Fig. S6**). Collectively, these findings confirmed

the high and predominant expression of *Tyrobp*, *Trem2*, and *C1q* components genes in

the microglia of both AD and EP patients. This consistency between animal models

and human patients suggests that microglia and these synaptic pruning genes play

crucial roles in the progression of both AD and EP pathology and may represent a

potential conserved regulatory pathway.

3.7 Computational modeling of synaptic pruning on neural network

To better understand how synaptic pruning influences neural network dynamics and

its potential role in the comorbidity of AD and EP, we developed a neural network

model simulating the hippocampal CA3 circuit. In this model, we systematically

varied the extent of synaptic pruning (0% to 50%) at either excitatory or inhibitory

synapses. We then quantified changes in the excitatory/inhibitory (E/I) ratio and

synchronization levels, conducting at least 50 simulation runs per condition (**Fig. 7**).

Our findings indicate that pruning of either excitatory or inhibitory synapses alters the

E/I ratio, with distinct effects. Inhibitory synaptic pruning significantly elevated the

E/I ratio, while excitatory pruning reduced it. These trends were consistent across all

simulations and in experiments where excitatory or inhibitory pruning was varied

independently (**Fig. 7A**). Correlation analysis showed a negative association between

excitatory pruning and the E/I ratio (r = −0.474) and a positive correlation between

inhibitory pruning and the E/I ratio (r = 0.684) (**Fig. 7B**). Linear regression further

confirmed that excitatory pruning reduced the E/I ratio (β = -1.038), whereas

inhibitory pruning increased it significantly (β = 1.454), with the effect of inhibitory

pruning surpassing that of excitatory pruning.

Interestingly, inhibitory pruning displayed a moderate positive correlation with event

synchronization (r = 0.346), while excitatory pruning showed only a weak negative

correlation (r = −0.039) (**Fig. 7B**). Linear regression analysis reflected this pattern,

with inhibitory pruning exerting a stronger effect on synchronization (β = 0.358) than

excitatory pruning ($\beta = -0.041$). Additionally, the E/I ratio itself was positively

correlated with synchronization ($r = 0.491$) (**Fig. 7B**), suggesting that the influence of

synaptic pruning on network synchronization may be nonlinear and indirectly

mediated by changes in the E/I balance.

These simulations suggest that excessive synapses pruning alters the E/I ratio, and

inhibitory pruning overpowering the effects of excitatory pruning and substantially

enhancing synchronization.

4. Discussion

Alzheimer's disease (AD) and epilepsy (EP) share a complex bidirectional

relationship, though the molecular pathways connecting AD-related

neurodegeneration and increased EP susceptibility remain unclear. To address this

issue, we used rigorous bioinformatic and computational methods to integrate and

analyze transcriptional data from multiple large-scale publicly available animal

models and human patient datasets. We identified consensus molecular pathways,

cellular processes, and neural network dynamics associated with this comorbidity

across multiple datasets.

Neuroinflammation Pathways Conserved in Epilepsy and Alzheimer's Disease

Our findings provide strong evidence supporting the hypothesis that

neuroinflammatory pathways are common between AD and EP, particularly those

directly triggered by tau pathology (7,19). Differential expression analysis

demonstrated significant gene overlap between EP and TG models (**Fig. 1A-B**).

WGCNA revealed preservation of the immune-related turquoise module across both

disease models as well as human patient datasets (**Fig. 2-3**). These results align with

previous analyses of this dataset that identified significant conservation of this module

across multiple AD patient cohorts (32). We extended these findings by demonstrating

the module's conservation and overlap in EP models and patient datasets, suggesting it

represents a fundamental molecular pathway shared between AD and EP (**Fig. 3,5**).

Functional enrichment analysis of shared DEGs and the preserved module revealed

significant enrichment in neuroinflammatory processes, including inflammatory

response, cytokine production, blood vessel morphogenesis, glial cell activation, and

complement and coagulation cascades (**Fig. 1D-E, Table S3**). Additionally, we

observed increased populations of neuroinflammation-associated cell types, including

astrocytes, microglia, and endothelial cells, in both TG and EP animal models (**Fig.**

6).

Neuroinflammation has been extensively documented in both conditions and appears

to serve a complex dual role in disease progression (57,58). While activated microglia

facilitate Aβ plaque clearance and reactive astrocytes provide metabolic support and

ionic homeostasis that may attenuate neural damage and seizures, chronic activation or dysfunction of these glial populations can promote disease progression through persistent neuroinflammation (59–62). Specifically, activated microglia and astrocytes secrete proinflammatory cytokines and chemokines, including IL-1β, TNF-α, and IL-6, which can enhance neuronal hyperexcitability, increase seizure susceptibility, exacerbate Aβ and tau aggregation, and accelerate neurodegeneration and cognitive decline (59–62) .

Additional inflammatory components, including complement cascade activation and blood-brain barrier (BBB) dysfunction, are implicated in both pathologies (63–66). Complement activation contributes to synaptic loss in AD and is elevated in TLE patient tissue (65,66). BBB disruption precedes neuronal hyperactivity in EP, promoting astrocyte dysfunction and seizures, while in AD it facilitates peripheral inflammatory cell infiltration, amplifying neuroinflammation (63,64).

These findings support the emerging view of neuroinflammation as a mechanistic bridge between AD and EP (19). AD-associated inflammation can enhance neuronal hyperexcitability and seizure susceptibility. Conversely, EP-induced inflammation may accelerate Aβ and tau accumulation, contributing to cognitive decline. Our results support previous studies and suggest that neuroinflammation is highly conserved between these diseases and between species. Future studies should focus on

revealing the precise causal relationship between these neuroinflammatory

components and AD-EP comorbidity.

Synaptic Pruning was Identified as a Core Pathway in this Comorbidity

A comprehensive analysis of this conserved module revealed a central synaptic

pruning pathway involving *Tyrobp*, *Trem2*, and *C1q* complement genes (**Fig. 4**).

These genes demonstrated consistent upregulation across both AD and EP patient

datasets, with preserved regulatory relationships in both animal models and human

patients (**Fig. 5**). Moreover, we confirmed their specific expression in microglial cells

across both animal and human patient samples (**Fig. S5-6**). Collectively, these

findings suggest that microglia-mediated synaptic pruning represents a core

mechanistic pathway underlying the comorbidity of AD and EP.

This pathway has been extensively characterized in AD pathology (67). Dysregulation

or mutations in *Tyrobp* and *Trem2* have established risk factors for AD through their

effects on microglial function (68,69). Under physiological conditions, microglia

maintain synaptic homeostasis through a regulated process in which complement

proteins, including *C1q*, tag unnecessary synapses for elimination (70). Microglial

receptors, particularly *Trem2* and *Tyrobp*, recognize these tags and initiate phagocytic

clearance, a process critical for neural circuit refinement and maintenance in the adult

brain (71). In AD pathology, these receptors enable microglia to recognize protein

deposits, triggering their transition to a disease-associated microglial (DAM) state

characterized by enhanced phagocytic activity and elevated expression of *Tyrobp*,

Trem2, *Apoe*, and complement proteins including *C1q* (67). While *Tyrobp* or *Trem2*

deficiency disrupts DAM activation and impairs critical microglial functions such as

plaque clearance, potentially accelerating AD pathology, this pathway overactivation

can also contribute to neurodegeneration and cognitive decline through excessive

C1q-mediated synaptic pruning (67,72–74).

Synaptic pruning can exert effects on both AD and EP disease progression through

two primary mechanisms: synaptic transmission and synaptic plasticity. Regarding

synaptic transmission, elevated *C1q* expression or Aβ accumulation in the CA1 region

leads to decreased miniature and evoked excitatory postsynaptic currents (75,76). In

AD, this reduction in synaptic transmission contributes to cognitive decline by

disrupting normal neural communication. For EP, this alteration affects both

excitatory and inhibitory neurons (77), potentially contributing to the imbalance

between excitation and inhibition that characterizes epileptic activity. Notably, these

effects can be reversed through *C1q* downregulation (78,79).

The pathway's impact on synaptic plasticity is equally significant. Research

demonstrates that C1q inhibition in aged experimental models can restore both

long-term potentiation (LTP) and long-term depression (LTD) (72,74,80). In non-aged

models, increased *C1q* expression also impairs the LTP process (77). This is

particularly relevant for AD, where impaired synaptic plasticity is a feature of

cognitive decline. In EP, compromised synaptic plasticity may contribute to the

formation and maintenance of epileptogenic circuits. The relationship between these

mechanisms appears bidirectional: excessive synaptic pruning leads to neuronal

damage and reduced LTP, which in turn promotes disease progression in both

conditions, creating a harmful feedback loop where synaptic dysfunction promotes

pathological processes.

Notably, interventions targeting this synaptic pruning pathway, either through

knockdown of regulatory genes such as *Tyrobp* and *Trem2* or direct inhibition of *C1q*

accumulation—have successfully restored electrophysiological level and cognitive

function in AD models, even in the presence of significant amyloid and tau pathology

(67,72–74). Our results establish a regulatory network that is consistent with previous

observations and further extends its role in the comorbidity of AD and EP (**Fig. 4**).

While this pathway is extensively studied in the context of AD, to the best of our

knowledge, no prior research has established a connection between this pathway and

the comorbidity of EP and AD. Our understanding of the relationship between

synaptic pruning and EP remains nascent, especially how to interpret its contribution

to the network dynamics (81). To address this gap, we conducted an in-silico

experiment to shed light on this issue.

Synaptic Pruning Alter E/I Balance to Promote Network Synchronization

Computational modeling has become an important tool for understanding AD and EP

pathology, models ranging from molecular-level simulations of ion dynamics to

large-scale network models (82,83). Disruption of excitatory/inhibitory (E/I) balance

and neural synchronization represent a key point of convergence between AD and EP

pathology, supported by a wealth of experimental evidence. In EP, patch-clamp

recordings show reduced GABAergic inhibition and enhanced glutamatergic

transmission, leading to network hyperexcitability (84,85). In AD, studies have shown

that AD mouse models exhibit spontaneous epileptiform activity, while human studies

using magnetoencephalography have demonstrated disruption of synchronization in

multiple frequency bands (16,86).

To explore how synaptic pruning contributes to the comorbidity of AD and EP, we

developed a computational model of the hippocampal CA3 circuit on top of Liu's

biologically plausible EP modeling. We calibrate fixed synaptic strength by adjusting

conductance-based synapses to produce realistic postsynaptic potential based on

animal experiments. (16,86), while leveraging synaptic connection pruning across

simulation trials on independently standard normal synaptic weights to transition the

network into abnormal states. This model allowed us to quantify how pruning of

excitatory and inhibitory synapses affects E/I balance (E/I ratio) and the temporal

coordination of neural firing (synchronization) (85).

Our computational model suggests that synaptic pruning can regulate the E/I balance

in neural networks, with inhibitory synapse pruning having a more pronounced effect.

Quantitative analysis by linear regression showed that excitatory synapse elimination

decreased the E/I ratio (β = -1.038), while inhibitory synapse removal led to a larger

increase in the E/I ratio (β = 1.454) (**Fig. 7**). Our finding that inhibitory pruning has a

stronger (enhancing) effect on network dynamics can be explained by the function of

inhibitory interneurons within neural circuits. These neurons often establish synaptic

connections with multiple excitatory neurons and act as regulators of circuit

coordination and synchronization (87). Therefore, when inhibitory control

mechanisms are impaired, the resulting disruption of regulation will trigger

dysregulation that extends to more widespread circuits. Moreover, endogenous

homeostatic mechanisms such as synaptic scaling appear to have a limited capacity to

compensate for the loss of inhibitory synapses (88). Ultimately, a reduction in

inhibitory synapses may produce a larger perturbation in neural networks than an

equivalent reduction in excitatory synapses.

Interestingly, our findings reveal a weaker correlation of synaptic pruning on network

synchronization compared to its impact on the E/I ratio. Specifically, inhibitory

pruning exhibited a moderate positive correlation with synchronization (r = 0.346), whereas excitatory pruning had a negligible correlation (r = –0.039). Notably, the E/I ratio itself demonstrated a strong correlation with synchronization levels (r = 0.491). While it seems intuitive that enhancing network excitability can enhance synchrony, the results of our analysis predict the importance of considering the specific location and functional role of pruned synapses in neural circuits (87). For instance, in a circuit that has an inhibitory effect on the broader network, pruning inhibitory synapses that inhibit this circuit will paradoxically reduce network synchronization. This explains the indirect, nonlinear association of synaptic pruning with network synchronization (**Fig. 7B**).

.

Although it is still difficult to establish a causal relationship between synaptic pruning and this comorbidity, our simulation results suggest a potential relationship, that is, the loss of neurons or synapses in neurodegenerative diseases disrupt the E/I balance of the neural network, may thereby promote epilepsy susceptibility (**Fig. 7**). In fact, epilepsy is well known as a comorbidity of multiple neurodegenerative diseases such as Huntington's and Parkinson's disease (89–91). Our simulations show that in the case of equal loss of excitatory or inhibitory synapses (even in the case of loss more excitatory synapses than inhibitory synapses) (**Fig. 7A**), the effect of inhibitory loss can overpower the effect of excitatory loss, causing a more extensive network E/I balance disruption and promoting network synchronization (**Fig. 7B**). Similar to the

established risk factors for AD for *Tyrobp* and *Trem2*, Mendelian randomization

studies also found that *C1qa* is a risk factor for both AD and EP (**Fig. S7**) (92). In fact,

in AD, synaptic loss often precedes cognitive decline and EP, which suggests that it

may be an initiating event rather than a consequence (93).

Our results highlight the role of inhibitory synaptic pruning in promoting network

synchronization. Notably, several recent studies have shown that microglial synaptic

pruning appear to have inhibitory pruning tendencies (94). Dejanovic B et al. had

shown that the *Trem2-C1q* microglial synaptic pruning pathway preferentially engulfs

inhibitory synapses in AD model, while Fan J et al. demonstrated similar preferential

pruning of inhibitory synapses during recurrent seizures that exacerbate EP model

pathology (94,95). At the molecular level, *C1q* acts as an "eat me" signal, tagging

synapses for microglia to eliminate (81). While this tagging process is not

synapse-type specific, microglia demonstrate selective pruning of inhibitory synapses

(81,94). Studies have shown that *C1q* blockade reduces both the frequency and

severity of seizures in traumatic brain injury models (96). However, it is also

important to note that *C1q* knockout also affects synaptic elimination during

development, resulting in enhanced connectivity and seizure activity (97,98). These

findings suggest the importance of managing microglial *C1q*-mediated synaptic

pruning during AD pathology and aging, and further exploration and regulation of this

potential preferential pruning of inhibitory synapses is warranted,

Conclusion

In this study, we integrated large-scale transcriptomic datasets from multiple AD and

EP animal models, as well as human patient cohorts, to identify a shared molecular

pathway that converges on microglial synaptic pruning. Specifically, we found a

highly preserved immune-related co-expression module centered on the

Tyrobp-Trem2-C1q complement pathway. Differential gene expression and gene

regulatory network analysis confirmed that this pathway is consistently upregulated in

both animal models and human patients. Our computer simulations further suggest a

mechanistic basis for how excessive synapses pruning (possibly preferential pruning

of inhibitory pruning) may tip the excitation-inhibition balance toward

hyperexcitation, leading to the comorbidity of AD and EP.

Together, these findings highlight a plausible microglial-driven,

complement-mediated synaptic pruning mechanism at the intersection of AD and EP.

In linking the dysregulated immune response to excitatory/inhibitory imbalances, our

work provides a unifying framework that may explain the strong bidirectional

comorbidity observed clinically. More broadly, these results underscore the potential

of targeting microglia–complement signaling for therapeutic interventions aimed at

ameliorating seizures and slowing cognitive decline in patients with AD and EP.

Future studies integrating spatial transcriptomics, advanced single-cell profiling, and

in vivo manipulation of complement pathways are warranted to elucidate how

microglial subtypes selectively prune synapses in AD and EP, and whether modulating

this process can indeed confer disease-modifying benefits.

Abbreviations

EP   Epilepsy

AD   Alzheimer's disease

E/I   Excitation/Inhibition

TLE   Temporal lobe epilepsy

Aβ   Amyloid-β

GEO   Gene Expression Omnibus

ENA   European Nucleotide Archive

WT   Wild-type

scRNA-Seq   Single-cell RNA sequencing

snRNA-Seq   Single-nucleus RNA sequencing

PCA   Principal Component Analysis

DEGs   Differentially expressed genes

BH   Benjamini-Hochberg

KEGG   Kyoto Encyclopedia of Genes and Genomes

GO   Gene ontology

MAD   Median absolute deviation

899	TOM	Topological overlap matrices
900	MEs	Module eigengenes
901	MM	Module membership
902	GS	Gene-genotype significance
903	PPI	Protein-protein interaction
904	FDR	False discovery rate
905	SVM	Support vector machine
906	PCs	Principal components
907	ES	Event synchrony
908	BBB	Blood-brain barrier
909	DAM	Disease-associated microglial
910	LTP	Long-term potentiation
911	LTD	Long-term depression
912	BP	Biological processes
913	CC	Cellular components
914	MF	Molecular functions
915	HR	Hazard ratios
916	CI	Confidence intervals
917	NCBI	National Center for Biotechnology Information
918		
919		

 Figures and legends

 Figure 1: Differential Expression and Functional Enrichment Analyses in EP and

 AD Model Datasets.

(A) Volcano plot shows the DEGs in the EP model dataset. The x-axis represents the

log2 fold change, and the y-axis shows the -log10 adjusted p-value. Upregulated

genes are highlighted in red, and downregulated genes are in blue. The dashed lines

mark the thresholds for significance (adjusted p-value < 0.05 and |log2 fold change| >

1). **(B)** Differential expression analysis of the TG model dataset. Scatter plot showing

progressive DEGs identified across different ages (4, 6, and 8 months) compared to

their control and initial months. DEGs are color-coded based on their expression level

(log2 fold change) compared to the control and initial month, with dot size to show

significance (adjusted p-value). The shaded regions correspond to the different age

groups, with the numbers indicating the count of EP-DEGs (numerator) and total

DEGs (denominator) with a log2 fold change below 1 that marks the threshold for

non-significant genes. **(C)** The single upregulated progressive DEG, Cst7, detected in

the J20 model dataset across different ages (6, 8, 10, and 12 months). The plot shows

the expression level of *Cst7* over time (x-axis: Age in months; y-axis: Normalized

counts), with different genotypes (J20 and WT) represented by different colors.

Dashed lines represent mean paths for each age. (LRTstatistic = 43.70,

log2foldchange[6-12months] = 2.73, FDR = 2.91E-5). **(D)** KEGG pathway

enrichment analysis of shared DEGs between the EP and TG datasets. The bubble plot

indicates the significantly enriched pathways, with bubble size representing the gene

count and color indicating the significance level (-log10 adjusted p-value). **(E)** GO

enrichment analysis of shared DEGs between the EP and TG model datasets. The dot

plot shows the enriched biological processes (BP), cellular components (CC), and

molecular functions (MF). Dot size corresponds to gene count, while color represents

the significance level (-log10 adjusted p-value). Immune-related functions were

prominently enriched.

Figure 2: Co-Expression Network Construction and Model-Trait Association.

(A-C) Hierarchical clustering dendrograms are presented for the EP (**A**), TG (**B**), and

J20 (**C**) model datasets, with gene modules identified through dynamic tree cutting.

The colors in the heatmap below each dendrogram correspond to distinct gene

modules, where the clustering of similar colors between the EP and other model

datasets (TG and J20) reflects the conservation of modules across conditions. The GS

is shown with darker colors indicating stronger correlations: red represents positive

correlations, and blue represents negative correlations with disease genotypes (EP for

panel A, TG for panel B, and J20 for panel C). Overlapping red areas in the

module-trait relationships suggest that these modules may be positively correlated

with both EP and AD model, as seen in the turquoise module in the TG model dataset.

The DEG row shows the distribution of DEGs within the modules: red indicates

DEGs, while grey indicates non-DEGs. **(D-F)** Heatmaps of the module-trait

relationships in EP (**D**), TG (**E**), and J20 (**F**) model datasets. The heatmaps show the

Pearson correlation coefficients between MEs and genotypes. The color scale

represents the strength and direction of the correlation, with red indicating positive

and blue indicating negative correlations. Modules are considered as significant

association with genotype using threshold (p-value < 0.05, |r| > 0.65).

Figure 3: Module Preservation Analysis Across EP and AD Datasets.

(A, B) Zsummary preservation statistics of EP model modules in the TG and J20

model datasets, respectively. Each point represents an EP model module, with the

y-axis displaying the Zsummary preservation statistic. A Zsummary > 10 indicates

strong evidence of preservation, 2 < Zsummary < 10 indicates moderate evidence, and

Zsummary < 2 suggests no evidence of preservation. The gray horizontal line at Z =

10 and the black horizontal line at Z = 2 serve as reference points in the plots. The

module size is indicated by the x-axis, and the color represents the module color. **(C,**

D) Bubble plots show the cross-tabulation of EP model modules with TG (**C**) and J20

(**D**) model modules; the cross-tabulation of EP patient (GSE63808) and EP model

datasets (**E**); cross-tabulation of AD (GSE132903) and TG mouse model datasets (**F**).

The color scale represents the adjusted p-value for the overlap significance, while the

size of the dots corresponds to the number of overlapping genes. Larger, more

significant dots indicate stronger overlaps between EP and AD modules such as the

turquoise module. Only significantly overlapping with overlap count > 30 and

adjusted p-value < 0.05 was colored, while those with insignificant overlap are

marked in gray.

Figure 4: Identification and Network Analysis of Shared Hub Genes in EP and

AD Datasets.

(**A**) Network plot illustrating the top 50 genes within the conserved turquoise module

across both the EP and TG model datasets. Each node represents a gene, and edges

signify gene co-expression connections. Grey edges correspond to connections in the

EP dataset, while blue edges represent those in the TG dataset. The color of the nodes

indicates their level of connectivity, with deeper color, indicating higher connectivity

within the network such as *Capg*, *Msn*, and *Npc2*. 17 red nodes are shared hub genes

that exhibit high connectivity in both networks and are identified as DEGs under both

conditions. Two shared hub genes are not shown in the figure due to their ranking in

both modules: *Apobec* and *Cxcl16*. (**B**) Venn diagram depicting the overlap between

hub genes identified in the EP and AD datasets and shared DEGs. 19 shared hub

genes were identified as key drivers in both conditions. (**C**) PPI network of the 14

shared hub genes constructed from 19 shared hub genes using the STRING database.

The size and color of the nodes are colored by their connectivity in the PPI network.

(**D**) The bubble plot showed the top GO biological process enrichment terms based on

p-values for the 19 shared hub genes. (**E**) The shared gene regulatory network was

inferred using the GENIE3 algorithm, showing the regulatory relationships among the

shared hub genes. (**F**) Progressive upregulation of key 5 shared hub genes involved in microglial synaptic pruning across different ages in the TG mouse model. The plot shows the expression level of these genes over time (x-axis: Age in months; y-axis: Normalized counts), with different genotypes represented by different colors (WT: black, TG: teal). Dashed lines represent mean paths for each age. LRT statistics of these genes can be found in **Table S2**.

Figure 5. Expression analysis and correlation patterns of shared hub genes in AD and EP patient datasets.

(**A, B**) Violin plots showing the differential expression of shared hub genes between control and disease groups in (**A**) EP patient dataset (E-MTAB-3123) and (**B**) AD patient dataset (GSE48350). Red violins represent control groups and blue violins represent disease groups. The width of each violin indicates the density of data points, and the internal box plots show the median and interquartile ranges. The Wilcoxon p-value is shown for each comparison. (**C, D**) Correlation heatmap showing the relationships between shared hub genes and age in the EP patient dataset (E-MTAB-3123) and AD patient dataset (GSE48350). The color intensity and box size represent the strength of Pearson correlation coefficients (r), with red indicating positive correlations and blue indicating negative correlations.

Figure 6: Cellular Composition Deconvolution and Association with Shared Hub

Genes in EP and TG Datasets.

(**A, C**) The plot shows the estimated proportions of different cell types (astrocytes,

neurons, oligodendrocytes, microglia, and endothelial) in control and EP model

conditions (**A**), and TG model condition (**C**). The x-axis represents different

conditions or months, and the y-axis shows the proportions of specific cell types in

each condition, with different genotypes represented by different colors. The

Mann-Whitney test was performed to determine significant cell composition

difference ($p < 0.05$) in EP model dataset. The Kruskal-Wallis test was performed to

determine the significance ($q < 0.05$) with the BH method was used to adjust the

p-value in pairwise comparisons among different groups in the TG model dataset. (**B,**

D) Correlation heatmap showing the relationships between key shared hub genes and

cell composition in the EP model dataset. The color intensity represents the strength

of Pearson correlation coefficients (r), with red indicating positive correlations and

blue indicating negative correlations. The significance of correlations with shared hub

gene is shown by red arrows, with arrow color proportional to correlation strength.

1040

1041 **Figure 7. Effects of excitatory and inhibitory synaptic pruning on neural**

1042 **network dynamics.**

1043 (**A**) Four-dimensional plot showing the simulation results of excitatory pruning ratio

1044 (0-0.5), inhibitory pruning ratio (0-0.5), E/I ratio, and synchronization. The surface

1045 visualizes the variation of the average E/I ratio with different pruning condition.

Colors indicate average synchronization levels from dark purple (close to 1, indicating

no change in synchrony) to yellow (indicating higher synchrony). (**B**) Correlation plot

showing the relationships among synapse pruning, E/I ratio and synchronization. The

color intensity represents the strength of Pearson correlation coefficients (r), with red

indicating positive correlations and blue indicating negative correlations.

Legends of supplementary files

Table S1: Summary of Quality Control and Mapping Statistics for Bulk RNA-seq Samples.

Detailed summary of the QC and mapping statistics for all RNA-seq samples

analyzed in this study. For each sample, the total number of raw reads, the number of

reads passing quality filtering through Fastp, and the reads retained after rRNA

contamination removal by ribodetector are reported. Each sample is categorized by

the experimental group, including relevant age and condition tags, and further

classified into either the AD or EP dataset. Notes regarding any specific issues or

annotations on individual samples are also provided.

Table S2: Differential Expression and Functional Enrichment Analysis of Shared DEGs.

Summary of differential expression and functional enrichment analyses across three

datasets: EP, TG, and J20 mouse models.

Table S3: Module Construction, Preservation, and Functional Enrichment of Gene Co-Expression Networks in Mouse Models.

Summary of WGCNA on EP, TG, and J20 mouse models. The table is divided into multiple sections, including the gene modules' membership identified in each dataset and their correlations with model-specific genotypes. Additionally, module preservation analyses between the EP and AD datasets are included. GO enrichment analyses of the turquoise modules are also shown.

Table S4: Protein-Protein Interaction Network and GO Biological Process Enrichment Analysis of Shared Hub Genes.

Analysis results of the PPI network constructed for 19 shared hub genes identified across EP and TG models. The table is divided into three sections. The PPI GO sheet summarizes the results of the GO biological process enrichment analysis. The PPI Degree sheet lists each gene and its degree of connectivity within the network. The PPI Interaction sheet details the specific interactions between these genes.

Table S5: Regulatory Network Prediction Using GENIE3 for Turquoise Module in Mice Models.

Results of regulatory network inference were conducted using the GENIE3 machine

learning algorithm on the gene expression profiles of the turquoise module from the

EP) and TG models. The table is divided into two sections: GENIE3TG and

GENIE3EP, representing the predicted regulatory relationships within the TG and EP

datasets, respectively. Each section lists the regulatory gene, the target gene, and the

corresponding weight of the predicted regulatory interaction.

Table S6: Cell Composition Deconvolution of Bulk RNA-seq Data.

Results of cell composition deconvolution analysis conducted using CIBERSORT

from BrainDeconvShiny on bulk RNA-seq data from the EP and AD models. The

table includes the estimated proportions of major brain cell types, including neurons,

astrocytes, oligodendrocytes, microglia, and endothelial for each sample, alongside

the sample identifiers, group classification, age, and condition type (e.g., rTg4510,

Wild Type).

Figure S1. Construction of co-expression module in model datasets.

(**A-C**) Hierarchical clustering dendrograms and trait heatmaps for the EP, rTg4510,

and J20 datasets. (**A**) EP dataset, (**B**) rTg4510 dataset, and (**C**) J20 dataset.

Hierarchical clustering was performed on variance-stabilized gene expression data to

identify potential outliers. The top 85% of genes with the highest Median Absolute

Deviation (MAD) were retained, and outliers were detected and removed before

network construction. Two samples were removed from the rTg4510 dataset, and

three samples were removed from the J20 dataset. The heatmaps below each

dendrogram display the trait data associated with each sample, with red indicating

outliers or specific traits of interest. **(D-F)** Soft-thresholding power Selection and

Comparability Assessment, **(D)** Selection of the soft-thresholding power for network

construction. The left panel shows the scale-free topology fit index (y-axis) as a

function of the soft-thresholding power (x-axis) for EP, TG, and J20 datasets. The

right panel depicts the mean connectivity (y-axis) against soft-thresholding power,

indicating that a power of 9 was chosen for all datasets to achieve scale-free topology

with $R^2 > 0.85$. **(F)** Comparability assessment of the EP, TG, and J20 datasets. Scatter

plots display the correlations between ranked mean gene expression (top) and ranked

gene connectivity (bottom) between EP and TG datasets (left) and between EP and

J20 datasets (right). Positive correlations and significant P values suggest strong

comparability between datasets.

Figure S2. Construction of co-expression module in patient datasets.

(**A**) Scale independence and mean connectivity plots across soft-thresholding powers

for the EP (GSE63808) and AD (GSE132903) patient datasets, with a selected

soft-thresholding power of 9 ensuring a scale-free topology ($R^2 \geq 0.85$). (**B-C**)

Dendrogram of gene modules identified in the EP and AD dataset using hierarchical

clustering and dynamic tree cutting based on topological overlap matrices (TOM).

Figure S3: ROC curve of Diagnostic model.

(**A**) in EP patient dataset (E-MTAB-3123) and (**B**) in AD patient dataset (GSE48350). ROC curves of various machine learning classifiers in classifying disease cases from control participants based on selected gene expression data. Remarkably, most of the models have an AUC between 0.85 and 0.92 in EP patient dataset, with the best performance recorded with XGBoost, reaching the highest AUC score in this EP dataset (AUC = 0.92). Although the AD model diagnostic efficiency is not as high as that of the EP dataset, which may reflect the complex disease background of AD, these diagnostic models still show high diagnostic efficiency (AUC > 0.7).

Figure S4: Cellular Composition Deconvolution and Association with Shared Hub Genes in J20 Datasets.

(**A**) The plot shows the cell composition level over time in J20 dataset (x-axis: Age in months; y-axis: Normalized counts), with different genotypes represented by different colors (WT: black, J20: teal). Dashed lines represent mean paths for each age. The Kruskal-Wallis test was performed to determine the significance with the BH method was used to adjust the p-value in pairwise comparisons among different groups, with a significance level of q < 0.05. (**B**) Correlation heatmap showing the relationships between key shared hub genes and cell composition in the J20 model dataset. The color intensity represents the strength of Pearson correlation coefficients (r), with red indicating positive correlations and blue indicating negative correlations.

1151

Figure S5: Expression Localization of Shared Hub Genes in Mouse Brain Cell

Types.

This figure shows the expression patterns of 19 shared hub genes across various cell

types in the mouse brain, based on the Whole Mouse Brain Transcriptomic Cell Type

Atlas (https://assets.nemoarchive.org/dat-qg7n1b0) from the Allen Brain Cell Atlas.

The top left panel illustrates the localization and color of various cell types, such as

astrocytes, microglia, macrophages, endothelial cells, and oligodendrocytes in the

brain atlas. Each circular panel highlights the specific expression profile of one gene,

with darker shading representing higher expression levels. Most of these genes (15

out of 19) are predominantly expressed in immune cells, specifically microglia and

macrophages.

1163

Figure S6: sc/nRNAseq Analysis of Shared Hub Gene Expression in Immune

Cells of Patients.

(**A, D**) Dot plot displaying the expression of shared hub genes across different

identified cell types in the EP patient dataset. The size of the dots represents the

proportion of cells expressing each gene, and the color intensity indicates the average

expression level. (**A**) EP patient dataset (GSE227223), (**D**) AD patient dataset

(GSE201048). (**B, E**) t-SNE plot showing the clustering of cells in the EP and AD

dataset based on their gene expression profiles, with distinct colors representing

1172 different cell types. (**B**) EP dataset, (**E**) AD dataset. (**C, F**) Feature plots showing the

1173 expression patterns of individual hub genes within the EP dataset, over on the t-SNE

1174 map. Darker purple areas represent higher expression levels. *TYROBP*, *TREM2*,

1175 *C1QA*, *C1QB*, and *C1QC* exhibit high expression predominantly in microglia clusters.

1176 (C) EP dataset, (F) AD dataset.

1177

Figure S7: Forest plot of Mendelian randomization result from DMRdb.

Forest plot showing hazard ratios (HR) and 95% confidence intervals (CI) for

associations between Alzheimer's disease subtypes, epilepsy, and *C1QA*

http://www.inbirg.com/DMRdb/. Red dots represent point estimates for hazard ratios,

and horizontal lines indicate 95% confidence intervals. p-values are shown for each

association. Values greater than 1.0 indicate increased risk for the outcome.

1184

1185 **Declarations**

1186

1187 **Ethics approval and consent to participate**

1188 Not applicable.

1189

1190 **Consent for publication**

1191 Not applicable.

1192

Availability of data and materials

The datasets analysed during the current study are available in the National Center for

Biotechnology Information (NCBI) or European Nucleotide Archive (ENA) under the

following accession numbers: PRJEB18790, GSE125957, GSE48350, E-MTAB-3123

GSE63808, GSE132903, GSE201048, and GSE227223.

Competing Interests

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Funding

This research was supported by the Young Scientists Fund of the National Natural

Science Foundation of China (Grant Nos. 82201629) and the Guangdong Provincial

Science and Technology Special Fund (Grant Nos. 210803136872196)

Authors' contribution

HL conceptualized the study, conducted the investigation, developed the methodology,

performed formal analysis, and contributed to writing the original draft and

reviewing/editing the manuscript. ZX, YT, and BL contributed to the investigation,

performed formal analysis, and participated in writing the original draft and

reviewing/editing the manuscript. JL contributed to writing the original draft and

reviewing/editing the manuscript. JW was involved in conceptualization and

contributed to writing the original draft and reviewing/editing the manuscript. MC,

XL, YS, SC, ZD, and MH contributed to the investigation and participated in

reviewing/editing the manuscript. BH and NW conceptualized the study and

contributed to writing the original draft and reviewing/editing the manuscript. XJ

provided supervision, acquired funding, contributed to conceptualization, and

participated in writing the original draft and reviewing/editing the manuscript. All

authors read and approved the final manuscript.

Acknowledgments

Not applicable

References

1. Giorgi FS, Saccaro LF, Busceti CL, Biagioni F, Fornai F. Epilepsy and Alzheimer's Disease: Potential mechanisms for an association. Vol. 160, Brain Research Bulletin. 2020.
2. Kamondi A, Grigg-Damberger M, Löscher W, Tanila H, Horvath AA. Epilepsy and epileptiform activity in late-onset Alzheimer disease: clinical and pathophysiological advances, gaps and conundrums. Vol. 20, Nature Reviews Neurology. 2024.
3. Fang Y, Si X, Wang J, Wang Z, Chen Y, Liu Y, et al. Alzheimer Disease and Epilepsy: A Mendelian Randomization Study. Neurology. 2023;101(4).
4. Wimo A, Jönsson L, Bond J, Prince M, Winblad B. The worldwide economic impact of dementia 2010. Alzheimer's and Dementia. 2013;9(1).
5. Ngugi AK, Bottomley C, Kleinschmidt I, Sander JW, Newton CR. Estimation of the burden of active and life-time epilepsy: A meta-analytic approach. Epilepsia. 2010;51(5).
6. Mendez MF, Lim GTH. Seizures in elderly patients with dementia: Epidemiology and management. Drugs Aging. 2003;20(11).
7. Leitner D, Pires G, Kavanagh T, Kanshin E, Askenazi M, Ueberheide B, et al. Similar brain proteomic signatures in Alzheimer's disease and epilepsy. Acta Neuropathol. 2024;147(1).

8.   Pandis D, Scarmeas N. Seizures in Alzheimer Disease: Clinical and Epidemiological Data.
https://doi.org/105698/1535-7511-125184 [Internet]. 2012;12(5):184–7. Available from:
https://journals.sagepub.com/doi/full/10.5698/1535-7511-12.5.184
9.   Tombini M, Assenza G, Ricci L, Lanzone J, Boscarino M, Vico C, et al. Temporal Lobe
Epilepsy and Alzheimer's Disease: From Preclinical to Clinical Evidence of a Strong
Association. Vol. 5, Journal of Alzheimer's Disease Reports. 2021.
10.   Johnson EL, Krauss GL, Lee AK, Schneider ALC, Dearborn JL, Kucharska-Newton AM,
et al. Association between Midlife Risk Factors and Late-Onset Epilepsy: Results from the
Atherosclerosis Risk in Communities Study. JAMA Neurol. 2018;75(11).
11.   Stefanidou M, Beiser AS, Himali JJ, Peng TJ, Devinsky O, Seshadri S, et al. Bi-directional
association between epilepsy and dementia: The Framingham Heart Study. Neurology.
2020;95(24).
12.   Gourmaud S, Shou H, Irwin DJ, Sansalone K, Jacobs LM, Lucas TH, et al. Alzheimer-like
amyloid and tau alterations associated with cognitive deficit in temporal lobe epilepsy.
Brain. 2020;143(1).
13.   Tai XY, Koepp M, Duncan JS, Fox N, Thompson P, Baxendale S, et al.
Hyperphosphorylated tau in patients with refractory epilepsy correlates with cognitive
decline: A study of temporal lobe resections. Brain. 2016;139(9).
14.   Zou Y, Wang C, Li H, Zhong M, Lin J, Hu Y, et al. Epileptic seizures induced by
pentylenetetrazole kindling accelerate Alzheimer-like neuropathology in 5×FAD mice.
Front Pharmacol. 2024 Oct 31;15:1500105.
15.   Watts ME, Pocock R, Claudianos C. Brain energy and oxygen metabolism: Emerging role
in normal function and disease. Front Mol Neurosci [Internet]. 2018 Jun 22 [cited 2025 Jan
3];11:396508. Available from: www.frontiersin.org
16.   Born HA. Seizures in Alzheimer's disease. Neuroscience. 2015 Feb 12;286:251–63.
17.   Toniolo S, Sen A, Husain M. Modulation of Brain Hyperexcitability: Potential New
Therapeutic Approaches in Alzheimer's Disease. International Journal of Molecular
Sciences 2020, Vol 21, Page 9318 [Internet]. 2020 Dec 7 [cited 2025 Jan 3];21(23):9318.
Available from: https://www.mdpi.com/1422-0067/21/23/9318/htm
18.   Kalyvas AC, Dimitriou M, Ioannidis P, Grigoriadis N, Afrantou T. Alzheimer's Disease
and Epilepsy: Exploring Shared Pathways and Promising Biomarkers for Future
Treatments. Journal of Clinical Medicine 2024, Vol 13, Page 3879 [Internet]. 2024 Jul 1
[cited   2025   Jan   3];13(13):3879.   Available   from:
https://www.mdpi.com/2077-0383/13/13/3879/htm
19.   Liew Y, Retinasamy T, Arulsamy A, Ali I, Jones NC, O'Brien TJ, et al. Neuroinflammation:
A Common Pathway in Alzheimer's Disease and Epilepsy. Journal of Alzheimer's Disease.
2023 Jan 1;94(s1):S253–65.
20.   Targa Dias Anastacio H, Matosin N, Ooi L. Neuronal hyperexcitability in Alzheimer's
disease: what are the drivers behind this aberrant phenotype? Translational Psychiatry
2022 12:1 [Internet]. 2022 Jun 22 [cited 2025 Jan 3];12(1):1–14. Available from:
https://www.nature.com/articles/s41398-022-02024-7
21.   Bele T, Ingale S. Network-based investigation to identify the common gene-disease

linkage between Alzheimer's disease, Parkinson's disease, and epilepsy. Network Biology
[Internet]. [cited 2025 Jan 3];2024(4):254–65. Available from: www.iaees.orgArticle

22. Mao R, Hu M, Liu X, Ye L, Xu B, Sun M, et al. Impairments of GABAergic transmission
in hippocampus mediate increased susceptibility of epilepsy in the early stage of
Alzheimer's disease. Cell Communication and Signaling [Internet]. 2024 Dec 1 [cited
2025 Jan 3];22(1):1–16. Available from:
https://biosignaling.biomedcentral.com/articles/10.1186/s12964-024-01528-7

23. Barker-Haliski M, Steve White H. Glutamatergic Mechanisms Associated with Seizures
and Epilepsy. Cold Spring Harb Perspect Med [Internet]. 2015 Aug 1 [cited 2025 Jan
3];5(8):a022863. Available from:
http://perspectivesinmedicine.cshlp.org/content/5/8/a022863.full

24. Tang Y, Yan Y, Mao J, Ni J, Qing H. The hippocampus associated GABAergic neural
network impairment in early-stage of Alzheimer's disease. Ageing Res Rev. 2023 Apr
1;86:101865.

25. Bosco F, Guarnieri L, Rania V, Palma E, Citraro R, Corasaniti MT, et al. Antiseizure
Medications in Alzheimer's Disease from Preclinical to Clinical Evidence. International
Journal of Molecular Sciences 2023, Vol 24, Page 12639 [Internet]. 2023 Aug 10 [cited
2025 Jan 3];24(16):12639. Available from:
https://www.mdpi.com/1422-0067/24/16/12639/htm

26. Andrews S. FastQC. Babraham Bioinformatics. 2010;

27. Chen S, Zhou Y, Chen Y, Gu J. Fastp: An ultra-fast all-in-one FASTQ preprocessor. In:
Bioinformatics. 2018.

28. Deng ZL, Münch PC, Mreches R, McHardy AC. Rapid and accurate identification of
ribosomal RNA sequences via deep learning. Nucleic Acids Res. 2022;50(10).

29. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol. 2019;37(8).

30. Liao Y, Smyth GK, Shi W. FeatureCounts: An efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 2014;30(7).

31. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15(12).

32. Castanho I, Murray TK, Hannon E, Jeffries A, Walker E, Laing E, et al. Transcriptional
Signatures of Tau and Amyloid Neuropathology. Cell Rep. 2020;30(6).

33. Elizarraras JM, Liao Y, Shi Z, Zhu Q, Pico AR, Zhang B. WebGestalt 2024: faster gene set
analysis and new support for metabolomics and multi-omics. Nucleic Acids Res [Internet].
2024 Jul 5 [cited 2025 Feb 9];52(W1):W415–21. Available from:
https://dx.doi.org/10.1093/nar/gkae456

34. Miller JA, Horvath S, Geschwind DH. Divergence of human and mouse brain
transcriptome highlights Alzheimer disease pathways. Proc Natl Acad Sci U S A.
2010;107(28).

35. Langfelder P, Horvath S. WGCNA: An R package for weighted correlation network
analysis. BMC Bioinformatics. 2008;9.

36. Zhang B, Horvath S. A general framework for weighted gene co-expression network

analysis. Stat Appl Genet Mol Biol. 2005;4(1).

37. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. Limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res. 2015;43(7).

38. Schober P, Schwarte LA. Correlation coefficients: Appropriate use and interpretation. Anesth Analg. 2018;126(5).

39. Langfelder P, Luo R, Oldham MC, Horvath S. Is my network module preserved and reproducible? PLoS Comput Biol. 2011;7(1).

40. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. Nucleic Acids Res. 2023;51(1 D).

41. Huynh-Thu VA, Irrthum A, Wehenkel L, Geurts P. Inferring regulatory networks from expression data using tree-based methods. PLoS One. 2010;5(9).

42. Newman AM, Liu CL, Green MR, Gentles AJ, Feng W, Xu Y, et al. Robust enumeration of cell subsets from tissue expression profiles. Nat Methods. 2015;12(5).

43. Sutton GJ, Poppe D, Simmons RK, Walsh K, Nawaz U, Lister R, et al. Comprehensive evaluation of deconvolution methods for human brain gene expression. Nat Commun. 2022;13(1).

44. Robinson MD, McCarthy DJ, Smyth GK. edgeR: A Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics. 2009;26(1).

45. Zhang Y, Chen K, Sloan SA, Bennett ML, Scholze AR, O'Keeffe S, et al. An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. Journal of Neuroscience. 2014;34(36).

46. Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, et al. Dictionary learning for integrative, multimodal and scalable single-cell analysis. Nat Biotechnol. 2024;42(2).

47. Kumar P, Lim A, Hazirah SN, Chua CJH, Ngoh A, Poh SL, et al. Single-cell transcriptomics and surface epitope detection in human brain epileptic lesions identifies pro-inflammatory signaling. Nat Neurosci. 2022;25(7).

48. Sun N, Victor MB, Park YP, Xiong X, Scannail AN, Leary N, et al. Human microglial state dynamics in Alzheimer's disease progression. Cell. 2023;186(20).

49. Stimberg M, Brette R, Goodman DFM. Brian 2, an intuitive and efficient neural simulator. Elife. 2019 Aug 1;8.

50. Izhikevich EM. Simple model of spiking neurons. IEEE Trans Neural Netw. 2003 Nov;14(6):1569–72.

51. Tzavellas NP, Tsamis KI, Katsenos AP, Davri AS, Simos Y V., Nikas IP, et al. Firing Alterations of Neurons in Alzheimer's Disease: Are They Merely a Consequence of Pathogenesis or a Pivotal Component of Disease Progression? Vol. 13, Cells. 2024.

52. Sammons RP, Vezir M, Moreno-Velasquez L, Cano G, Orlando M, Sievers M, et al. Structure and function of the hippocampal CA3 module. Proc Natl Acad Sci U S A. 2024;121(6).

53. Vyleta NP, Borges-Merjane C, Jonas P. Plasticity-dependent, full detonation at

hippocampal   mossy   fiber-CA3   pyramidal   neuron   synapses.   Elife. 2016;5(OCTOBER2016).

54.   Liu Y, Grigorovsky V, Bardakjian B. Excitation and Inhibition Balance Underlying Epileptiform Activity. IEEE Trans Biomed Eng. 2020;67(9).

55.   Penn Y, Segal M, Moses E. Network synchronization in hippocampal neurons. Proc Natl Acad Sci U S A. 2016;113(12).

56.   Yao Z, van Velthoven CTJ, Kunst M, Zhang M, McMillen D, Lee C, et al. A high-resolution transcriptomic and spatial atlas of cell types in the whole mouse brain. Nature. 2023;624(7991).

57.   Vezzani A, French J, Bartfai T, Baram TZ. The role of inflammation in epilepsy. Nature Reviews   Neurology   2010   7:1   [Internet].   2010;7(1):31–40.   Available   from: https://www.nature.com/articles/nrneurol.2010.178

58.   Heneka MT, Golenbock DT, Latz E. Innate immunity in Alzheimer's disease. Nature Immunology   2015   16:3   [Internet].   2015;16(3):229–36.   Available   from: https://www.nature.com/articles/ni.3102

59.   Pekny M, Pekna M. Astrocyte reactivity and reactive astrogliosis: Costs and benefits. Physiol Rev. 2014;94(4).

60.   Chen ZP, Wang S, Zhao X, Fang W, Wang Z, Ye H, et al. Lipid-accumulated reactive astrocytes promote disease progression in epilepsy. Nat Neurosci. 2023;26(4).

61.   Onyango IG, Jauregui G V, Čarná M, Bennett JP, Stokin GB. Neuroinflammation in Alzheimer's disease. Vol. 9, Biomedicines. 2021.

62.   Vezzani A, Viviani B. Neuromodulatory properties of inflammatory cytokines and their impact on neuronal excitability. Neuropharmacology. 2015;96(PA):70–82.

63.   van Vliet EA, Aronica E, Gorter JA. Blood-brain barrier dysfunction, seizures and epilepsy. Vol. 38, Seminars in Cell and Developmental Biology. 2015.

64.   Zlokovic B V. Neurovascular pathways to neurodegeneration in Alzheimer's disease and other disorders. Vol. 12, Nature Reviews Neuroscience. 2011.

65.   Kopczynska M, Zelek WM, Vespa S, Touchard S, Wardle M, Loveless S, et al. Complement system biomarkers in epilepsy. Seizure. 2018;60.

66.   Veerhuis R, Nielsen HM, Tenner AJ. Complement in the brain. Vol. 48, Molecular Immunology. 2011.

67.   Haure-Mirande JV, Audrain M, Ehrlich ME, Gandy S. Microglial TYROBP/DAP12 in Alzheimer's disease: Transduction of physiological and pathological signals across TREM2. Vol. 17, Molecular Neurodegeneration. 2022.

68.   Pottier C, Ravenscroft TA, Brown PH, Finch NCA, Baker M, Parsons M, et al. TYROBP genetic variants in early-onset Alzheimer's disease. Neurobiol Aging. 2016 Dec 1;48:222.e9-222.e15.

69.   Neumann H, Daly MJ. Variant TREM2 as Risk Factor for Alz hei mer's Disease. 2013;

70.   Stevens B, Allen NJ, Vazquez LE, Howell GR, Christopherson KS, Nouri N, et al. The Classical Complement Cascade Mediates CNS Synapse Elimination. Cell. 2007;131(6).

71.   Paolicelli RC, Ferretti MT. Function and dysfunction of microglia during brain development: Consequences for synapses and neural circuits. Vol. 9, Frontiers in Synaptic

1411   Neuroscience. 2017.

1412   72.   Audrain M, Haure-Mirande JV, Wang M, Kim SH, Fanutza T, Chakrabarty P, et al.
1413   Integrative approach to sporadic Alzheimer's disease: deficiency of TYROBP in a
1414   tauopathy mouse model reduces C1q and normalizes clinical phenotype while increasing
1415   spread and state of phosphorylation of tau. Mol Psychiatry. 2019;24(9).

1416   73.   Haure-Mirande JV, Audrain M, Fanutza T, Kim SH, Klein WL, Glabe C, et al. Deficiency
1417   of TYROBP, an adapter protein for TREM2 and CR3 receptors, is neuroprotective in a
1418   mouse model of early Alzheimer's pathology. Acta Neuropathol [Internet]. 2017 Nov 1
1419   [cited   2024   Oct   1];134(5):769–88.   Available   from:
1420   https://link.springer.com/article/10.1007/s00401-017-1737-3

1421   74.   Haure-Mirande JV, Wang M, Audrain M, Fanutza T, Kim SH, Heja S, et al. Integrative
1422   approach to sporadic Alzheimer's disease: deficiency of TYROBP in cerebral Aβ
1423   amyloidosis mouse normalizes clinical phenotype and complement subnetwork molecular
1424   pathology without reducing Aβ burden. Molecular Psychiatry 2018 24:3 [Internet]. 2018
1425   Oct   3   [cited   2025   Feb   9];24(3):431–46.   Available   from:
1426   https://www.nature.com/articles/s41380-018-0255-6

1427   75.   Neniskyte U, Gross CT. Errant gardeners: glial-cell-dependent synaptic pruning and
1428   neurodevelopmental disorders. Nature Reviews Neuroscience 2017 18:11 [Internet]. 2017
1429   Sep   21   [cited   2025   Feb   9];18(11):658–70.   Available   from:
1430   https://www.nature.com/articles/nrn.2017.110

1431   76.   Hu Y, Qi H, Yang J, Wang F, Peng X, Chen X, et al. Wogonin mitigates
1432   microglia-mediated synaptic over-pruning and cognitive impairment following epilepsy.
1433   Phytomedicine. 2024 Dec 1;135:156222.

1434   77.   Wu X, Gao Y, Shi C, Tong J, Ma D, Shen J, et al. Complement C1q drives
1435   microglia-dependent synaptic loss and cognitive impairments in a mouse model of
1436   lipopolysaccharide-induced neuroinflammation. Neuropharmacology. 2023;237.

1437   78.   Bie B, Wu J, Foss JF, Naguib M. Activation of mGluR1 Mediates C1q-Dependent
1438   Microglial Phagocytosis of Glutamatergic Synapses in Alzheimer's Rodent Models. Mol
1439   Neurobiol [Internet]. 2019 Aug 15 [cited 2025 Feb 9];56(8):5568–85. Available from:
1440   https://link.springer.com/article/10.1007/s12035-019-1467-8

1441   79.   Wu J, Bie B, Foss JF, Naguib M. Amyloid Fibril–Induced Astrocytic Glutamate
1442   Transporter Disruption Contributes to Complement C1q-Mediated Microglial Pruning of
1443   Glutamatergic Synapses. Mol Neurobiol [Internet]. 2020 May 1 [cited 2025 Feb
1444   9];57(5):2290–300.   Available   from:
1445   https://link.springer.com/article/10.1007/s12035-020-01885-7

1446   80.   Iweka CA, Seigneur E, Hernandez AL, Paredes SH, Cabrera M, Blacher E, et al. Myeloid
1447   deficiency of the intrinsic clock protein BMAL1 accelerates cognitive aging by disrupting
1448   microglial synaptic pruning. J Neuroinflammation [Internet]. 2023 Dec 1 [cited 2025 Feb
1449   9];20(1):1–16.   Available   from:
1450   https://link.springer.com/articles/10.1186/s12974-023-02727-8

1451   81.   Andoh M, Ikegaya Y, Koyama R. Synaptic pruning by microglia in epilepsy. Vol. 8,
1452   Journal of Clinical Medicine. 2019.

1453   82.   Yuan P, Zhang M, Tong L, Morse TM, McDougal RA, Ding H, et al. PLD3 affects axonal
1454   spheroids and network defects in Alzheimer's disease. Nature 2022 612:7939 [Internet].
1455   2022   Nov   30   [cited   2025   Jan   3];612(7939):328–37.   Available   from:
1456   https://www.nature.com/articles/s41586-022-05491-6

1457   83.   Stefanescu RA, Shivakeshavan RG, Talathi SS. Computational models of epilepsy. Seizure.
1458   2012 Dec 1;21(10):748–59.

1459   84.   Fritschy JM. Epilepsy, E/I balance and GABAA receptor plasticity. Front Mol Neurosci
1460   [Internet].   2008   Mar   28   [cited   2025   Jan   3];1(MAR):201.   Available   from:
1461   www.frontiersin.org

1462   85.   Ren X, Brodovskaya A, Hudson JL, Kapur J. Connectivity and Neuronal Synchrony
1463   during Seizures. Journal of Neuroscience [Internet]. 2021 Sep 8 [cited 2025 Jan
1464   3];41(36):7623–35. Available from: https://www.jneurosci.org/content/41/36/7623

1465   86.   van Nifterick AM, de Haan W, Stam CJ, Hillebrand A, Scheltens P, van Kesteren RE, et al.
1466   Functional network disruption in cognitively unimpaired autosomal dominant Alzheimer's
1467   disease: a magnetoencephalography study. Brain Commun [Internet]. 2024 Oct 30 [cited
1468   2025 Jan 3];6(6). Available from: https://dx.doi.org/10.1093/braincomms/fcae423

1469   87.   Luo L. Architectures of neuronal circuits. Science (1979) [Internet]. 2021 Sep 3 [cited
1470   2025   Jan   3];373(6559).   Available   from:
1471   https://www.science.org/doi/10.1126/science.abg7285

1472   88.   Dzyubenko E, Fleischer M, Manrique-Castano D, Borbor M, Kleinschnitz C, Faissner A,
1473   et al. Inhibitory control in neuronal networks relies on the extracellular matrix integrity.
1474   Cellular and Molecular Life Sciences [Internet]. 2021 Jul 1 [cited 2025 Jan
1475   3];78(14):5647–63.   Available   from:
1476   https://link.springer.com/article/10.1007/s00018-021-03861-3

1477   89.   Neri S, Mastroianni G, Gardella E, Aguglia U, Rubboli G. Epilepsy in neurodegenerative
1478   diseases. Epileptic Disorders [Internet]. 2022 Apr 1 [cited 2025 Feb 15];24(2):249–73.
1479   Available from: https://onlinelibrary.wiley.com/doi/full/10.1684/epd.2021.1406

1480   90.   Cano A, Fonseca E, Ettcheto M, Sánchez-López E, de Rojas I, Alonso-Lana S, et al.
1481   Epilepsy in Neurodegenerative Diseases: Related Drugs and Molecular Pathways.
1482   Pharmaceuticals 2021, Vol 14, Page 1057 [Internet]. 2021 Oct 18 [cited 2025 Feb
1483   15];14(10):1057. Available from: https://www.mdpi.com/1424-8247/14/10/1057/htm

1484   91.   Paudel YN, Angelopoulou E, Piperi C, Othman I, Shaikh MF. Revisiting the impact of
1485   neurodegenerative proteins in epilepsy: Focus on alpha-synuclein, beta-amyloid, and Tau.
1486   Vol. 9, Biology. 2020.

1487   92.   Zheng X, Tian Z, Che X, Zhang X, Xiang Y, Ge Z, et al. DMRdb: a disease-centric
1488   Mendelian randomization database for systematically assessing causal relationships of
1489   diseases with genes, proteins, CpG sites, metabolites and other diseases. Nucleic Acids Res
1490   [Internet]. 2025 Jan 6 [cited 2025 Feb 15];53(D1):D1363–71. Available from:
1491   https://dx.doi.org/10.1093/nar/gkae853

1492   93.   Chin J, Scharfman HE. Shared cognitive and behavioral impairments in epilepsy and
1493   Alzheimer's disease and potential underlying mechanisms. Epilepsy & Behavior. 2013
1494   Mar 1;26(3):343–51.

1495   94.   Dejanovic B, Wu T, Tsai MC, Graykowski D, Gandham VD, Rose CM, et al. Complement
1496   C1q-dependent excitatory and inhibitory synapse elimination by astrocytes and microglia
1497   in Alzheimer's disease mouse models. Nat Aging. 2022;2(9).

1498   95.   Fan J, Dong X, Tang Y, Wang X, Lin D, Gong L, et al. Preferential pruning of inhibitory
1499   synapses by microglia contributes to alteration of the balance between excitatory and
1500   inhibitory synapses in the hippocampus in temporal lobe epilepsy. CNS Neurosci Ther
1501   [Internet]. 2023 Oct 1 [cited 2025 Jan 3];29(10):2884–900. Available from:
1502   https://onlinelibrary.wiley.com/doi/full/10.1111/cns.14224

1503   96.   Holden SS, Grandi FC, Aboubakr O, Higashikubo B, Cho FS, Chang AH, et al.
1504   Complement factor C1q mediates sleep spindle loss and epileptic spikes after mild brain
1505   injury. Science (1979). 2021;373(6560).

1506   97.   Chu Y, Jin X, Parada I, Pesic A, Stevens B, Barres B, et al. Enhanced synaptic connectivity
1507   and epilepsy in C1q knockout mice. Proc Natl Acad Sci U S A. 2010;107(17).

1508   98.   Ma Y, Ramachandran A, Ford N, Parada I, Prince DA. Remodeling of dendrites and spines
1509   in the C1q knockout model of genetic epilepsy. Epilepsia. 2013;54(7).

1510

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Cell Composition by Type

Figure 6

Figure 7