Genome-wide consensus transcriptional signatures identify synaptic pruning

linking Alzheimer's disease and epilepsy

Huihong Li[1,2,3,4], Zhiran Xie[4], Yuxuan Tian[4], Ruoyin Zhou[4], Yaxi Yang[4], Bingying

Lin[1,2,3,5], Si Chen[1,2,3,5], Jie Wu[1,2,3], Zihan Deng[1,2,3,5], Jianwei Li[1,2,3,5], Mingjie Chen[6],

Xueke Liu[6], Yushan Sun[6], Bing Huang[1,2,3,5], Naili Wei[1], Xiaoyu Ji[1,2,3,5*]

[1]Department of Neurosurgery, The First Affiliated Hospital of Shantou University

Medical College, Shantou, Guangdong 515041, China

[2]Brain Function and Disease Laboratory, Shantou University Medical College,

Shantou, Guangdong 515041, China

[3]Department of Pharmacology, Shantou University Medical College, Shantou,

Guangdong 515041, China

[4]Program in Computational Biology and Biomedical Informatics, Yale University,

New Haven, Connecticut, United States of America

[5]Pain and Related Diseases Research Laboratory, Shantou University Medical College,

Shantou, Guangdong 515041, China

[6]Department of Biotechnology and Food Engineering, Guangdong Technion – Israel

Institute of Technology, Shantou, Guangdong 515063, China

[*]**Corresponding authors**

Contact: Dr. Xiaoyu Ji, xyji@stu.edu.cn

Abstract

Alzheimer's disease (AD) and epilepsy (EP) share a complex bidirectional relationship, yet the molecular mechanisms underlying their comorbidity remain insufficiently explored. To identify potential transcriptional programs across animal models and human patients with AD and EP, we conducted a comprehensive genome-wide transcriptomic analysis. Our investigation included mouse models of temporal lobe epilepsy (pilocarpine- and kainic acid-induced; n = 280), AD transgenic models (7 transgenic models expressing human tau or amyloid pathology; n = 257), and performed cross-species validation in human cohorts (EP: n = 182; AD: n = 301). We identified a highly conserved immune-related module across all models and patient cohorts. The hub consensus signatures of this module were centered around a microglial synaptic pruning pathway involving *TYROBP*, *TREM2*, and *C1Q* complement components. Gene regulatory network analysis identified *TYROBP* as the key upstream hub signature. These signatures showed consistent up-regulation in both conditions and strong diagnostic potential. Differential expression analyses revealed their predominant expression in specific microglial subpopulations associated with complement-mediated synaptic pruning and immune activation. Neural circuit modeling further demonstrates the asymmetric sensitivity of synaptic pruning to network dynamics. Loss of inhibitory synapses has a disproportionately significant impact on neural network excitation/inhibition balance and synchronization. Our findings support microglial complement-mediated synaptic pruning as a conserved

43   central pathway linking neurodegeneration to epileptogenesis, suggesting a promising

44   therapeutic target for AD and EP comorbidity.

46   **Keywords:** Alzheimer's disease, epilepsy, synaptic pruning, computational

47   neuroscience, modeling, transcriptome, *TYROBP, TREM2, C1Q*

49 **Graphic abstract**

50
51

Highlights

- Genome-wide transcriptomic analysis across epilepsy (EP) and Alzheimer's disease (AD) models and patients identified a conserved immune module.

- The *TYROBP-TREM2-C1Q* microglial synaptic pruning pathway was identified as a central consensus signature across model and patient cohorts.

- Consensus signatures possess potential diagnostic utility in AD and EP patients, with predominant expression in specific microglial subpopulations

- Neural circuit modeling demonstrates the asymmetric effects of synaptic pruning, whereby loss of inhibitory synapses disproportionately disrupts the E/I balance, leading to increased network synchronization.

1. Introduction

Alzheimer's disease (AD) and epilepsy (EP) are two common neurological diseases with a complex bidirectional relationship that contributes to their respective pathogenesis (1–3). AD, characterized by progressive cognitive decline and neurodegeneration, is the leading cause of dementia worldwide, affecting an estimated 55 million individuals. This number is projected to triple by 2050 as life expectancy increases (4). EP, characterized by recurrent unprovoked seizures, affects over 50 million individuals worldwide, with an exceptionally high incidence among elderly populations, especially those with neurodegenerative conditions such as AD (5,6).

Epidemiological studies demonstrate a strong association between AD and EP. AD patients exhibit increased seizure susceptibility, with 10–22% experiencing unprovoked seizures, while 22–54% demonstrate subclinical epileptiform activity during disease progression (7,8). These seizures, predominantly occurring in the temporal lobe, correlate with accelerated cognitive decline and exacerbated AD pathology (9). Conversely, EP patients demonstrate a two- to three-fold increased risk of developing dementia (10,11). This elevated risk is particularly evident in temporal lobe epilepsy (TLE), where numerous pathological features overlap with AD (9).

The pathological overlap between AD and EP manifests across multiple biological scales (12). Both conditions feature amyloid deposition, tau accumulation, and

hippocampal sclerosis at the molecular and cellular levels, particularly in TLE

(13–15). Chronic seizures increase neuronal activity and metabolic demands,

potentially accelerating the production and aggregation of amyloid-β (Aβ) and

hyperphosphorylated tau protein (16,17). Conversely, the progressive accumulation of

these pathological proteins in AD appears to lower seizure thresholds, promoting

hyperexcitability and epileptiform activity (18,19). At the network level, both

disorders demonstrate profound disruption of the excitation/inhibition (E/I) balance,

with a shift toward hyperexcitability driving both cognitive impairment and seizure

susceptibility. This network dysfunction is accompanied by impaired synaptic

plasticity, altered synchronization patterns, and progressive circuit reorganization

(19–21).

Despite these established clinical and pathological connections, the precise molecular

mechanisms governing the AD-EP relationship remain insufficiently explored. While

previous studies have identified potential contributors—including dysregulated

protein homeostasis, mitochondrial dysfunction, and chronic inflammatory

responses—most investigations have examined these conditions in isolation rather

than explicitly addressing their comorbidity (22–24). Several factors have limited

current research approaches: 1) inadequate integration of transcriptomic data across

multiple disease models and human cohorts; 2) insufficient attention to shared cellular

processes that bridge molecular alterations with network dysfunction; and 3) limited

application of computational modeling to connect molecular findings with clinically

relevant physiological outcomes.

The challenge of treating EP in AD patients further highlights the need for deeper

mechanistic insights. Clinicians face complex decisions regarding seizure treatment in

AD patients, including whether to treat subclinical epileptiform activity and which

antiseizure medications minimize cognitive side effects (25). Emerging therapeutic

approaches require a more precise understanding of the shared pathological

mechanisms. Systems biology and computational neuroscience offer powerful tools to

integrate insights from animal models and human patients, potentially revealing

actionable therapeutic targets that simultaneously address both conditions.

This study investigated the molecular mechanisms underlying the comorbidity

between AD and EP through genome-wide transcriptional analyses. Our approach

included: 1) comprehensive transcriptomic analysis of mouse models of TLE (n = 280)

and AD (n = 257) to identify consensus signatures across disease paradigms, 2)

cross-species validation in human cohorts (EP: n = 182; AD: n = 301) to assess

clinical relevance, and 3) computational modeling to connect molecular findings with

network-level mechanisms. Through these integrated analyses, we aimed to identify

conserved transcriptional programs that might explain the mechanistic link between

neurodegeneration and epileptogenesis, potentially revealing novel therapeutic targets

for both conditions.

2. Materials and Methods

A workflow overview of the datasets and analyses can be found in the figure below.

2.1 Data Collection

We retrieved several large-scale, well-characterized datasets from EP and AD research

that focused on genome-wide expression patterns in well-characterized disease

models and appropriate controls. The EP model datasets included two of the most

used TLE modeling approaches. The first dataset (PRJEB18790) contained 200

hippocampi of pilocarpine-induced (PO) model mice and wild-type (WT) controls

(26). The second dataset (GSE73878) comprised 80 hippocampal samples from the

kainic acid-induced (KA) model and WT control. For the AD model datasets, we

selected two datasets that systematically examined genome-wide expression

alterations in amyloid and tau pathology progression. The Matarin dataset (GSE64398)

contained 113 hippocampal samples from five mouse models and their controls: TAU

mice, TAS10 mice, TPM mice, and TASTPM mice (both heterozygous and

homozygous variants) (27). The Castanho dataset (GSE125957) included J20 and

rTg4510 models with 144 hippocampal and entorhinal cortex samples and matched

controls (28). To validate disease co-expression modules across species, we

constructed modules using 129 hippocampal samples from TLE patients (GSE63808)

and incorporated 30 consensus AD modules from the AMP-AD consortium's

meta-analysis of 2,114 samples across seven brain regions (29,30). To evaluate the

diagnostic value of transcriptional signatures, we analyzed 47 hippocampal samples

from TLE patients and controls (E-MTAB-3123) and 253 samples from AD patients

and controls (GSE48350). In addition, we reanalyzed single-nucleus RNA-sequencing

(snRNA-seq) datasets to characterize cell-type and subtype-specific expression of

identified signatures. Details about data collection, quality control, preprocessing,

parameters, and additional methods can be found in **Text S1**.

2.2 Gene Co-expression Networks Construction

To construct and compare gene co-expression networks (GCNs) between EP and AD model datasets, we implemented the standard meta-WGCNA comparative methodology (31,32), effectively identifying conserved gene modules while mitigating batch effects between datasets. Using the WGCNA package (1.73), we constructed signed GCNs based on genome-wide expression profiles. Network construction employed the minimum soft-thresholding power that achieved scale-free topology ($R^2 \geq 0.85$) across all model datasets. Using the signed network option, we generated adjacency matrices and transformed them into topological overlap matrices (TOM) to quantify gene connectivity profile similarities. Module identification proceeded through average linkage hierarchical clustering of TOM dissimilarity (1-TOM), followed by dynamic tree cutting (parameters: deepSplit = 1, minClusterSize = 27). We also constructed consensus modules to identify GCNs consistently in all conditions. This consensus approach involved calculating the component-wise minimum values of the topological overlap matrices from all datasets. We calculated module eigengenes (MEs) from the first principal component to summarize module expression profiles for each module. We then identified disease-associated modules by computing Pearson correlation coefficients between MEs and genotype information, with significance thresholds of $P < 0.05$ and |correlation coefficient| > 0.35. Additionally, we determined module membership (MM) and genotype significance (GS) by calculating Pearson correlations between

individual genes and their respective MEs or genotypes. Gene Ontology (GO)

enrichment analyses for module annotation were conducted using clusterProfiler

(4.10.1) (33). Hub genes act as key drivers within a module and were identified using

stringent criteria to minimize false positives: genes with MM > 0.9 and ranking in the

top 5% for intramodular connectivity within each module. Consensus hub signatures

were obtained by intersecting the hub genes of key modules across different datasets.

2.3 Module Preservation Analysis

To evaluate the preservation of EP model modules across AD model datasets, we

implemented dual analytical approaches as described by Langfelder et al. (34). First,

we calculated Zsummary preservation statistics using the modulePreservation

function. This composite metric comprehensively assesses preservation by integrating

module size, density, and connectivity parameters, quantifying how effectively EP

module structures are maintained in AD datasets. We interpreted the preservation

evidence according to established thresholds: strong (Zsummary > 10), moderate (2 <

Zsummary < 10), or absent (Zsummary < 2). Additionally, we performed

cross-tabulation analysis to quantify module overlap between EP and AD model

datasets, as well as between model and patient datasets. This approach generated

contingency tables detailing the genes shared between the reference and test dataset

modules. We determined overlap significance using Fisher's exact test with

Benjamini-Hochberg (BH) correction for multiple testing.

2.4 Protein-Protein Interaction and Gene Regulatory Network Analysis

To explore interactions and potential regulatory relationships among consensus hub

signatures, we constructed a protein-protein interaction (PPI) network using the

STRING database (12.0) with a confidence score threshold of 0.4 and a false

discovery rate (FDR) of 5% (35). We also performed regulatory network analysis with

the GENIE3 (1.28.0) algorithm, which applies a random forest approach to infer

potential regulatory relationships from gene expression data. GENIE3 ranks

regulatory interactions by weight, with higher weights indicating stronger regulatory

links (36). Gene expression profiles of the consensus hub signatures from all datasets

were used to infer plausible regulatory interactions. Only the top 30% plausible

regulatory links were retained to construct gene regulatory networks to reduce false

positives.

2.5 Single-nucleus RNA-seq Analysis

snRNA-seq datasets from EP and AD cohorts were reanalyzed to investigate

cell-type-specific transcriptional changes of consensus transcriptional signatures. For

the EP cohort, we utilized the snRNA-seq dataset (GSE190453) profiling temporal

lobe tissue from TLE patients and non-epileptic controls (cells = 66305) (37). For the

AD cohort (syn51758062.1), we analyzed snRNA-seq data from the Religious Orders

Study and Rush Memory and Aging Project (ROSMAP) cohort, encompassing 48

prefrontal cortex (PFC) samples from AD patients and non-pathology controls (cells =

50985) (38). Data preprocessing was performed using the Seurat package (5.0.1)

according to the original studies (39). Dimensionality reduction was conducted

through PCA, with the first 30 components used for clustering and UMAP

visualization. Cell-type annotation followed the definitions established in the original

studies and calculated scores for each nucleus using the AddModuleScore() function,

and cell-type identity was assigned based on the highest-scoring module. Differential

expressions of consensus transcriptional signature between control and disease groups

were evaluated using two-sided Wilcoxon rank-sum tests. To demonstrate the

expression of our signatures in microglial subtypes, we utilized the comprehensive

single-cell microglial dataset (GSE204702), which covers 215,680 microglia from

different neurological diseases and brain regions (40). We performed differential

expression analysis for their 12 subclasses and analyzed the expression patterns of our

consensus signatures across the identified microglial subtypes.

2.6 Neural Circuit Modeling

We implemented a spiking neural network using the Izhikevich neuron model to

simulate neural circuit dynamics and explore the contribution of synaptic pruning to

pathological changes in AD and EP. The network consisted of 1000 standard

Izhikevich neurons (800 excitatory, 200 inhibitory) and was simulated for 1000ms

using the Brian2 package (41). Network architecture included random sparse

connectivity (0.1 probability) with conductance-based synapses modeling AMPA

(excitatory) and GABA-a (inhibitory) currents across four connection types: E→E,

E→I, I→E, and I→I (42). Synaptic weights were drawn from a standard normal

distribution and normalized to [0,1]. We simulated synaptic pruning by selectively

removing excitatory or inhibitory synapses with the lowest weights (0-50%, with 5%

increments), repeating each simulation at least 50 times. To quantify network

dynamics, we calculated the excitation/inhibition (E/I) ratio and event synchrony (ES)

(43). The E/I ratio compared total excitatory currents to the absolute magnitude of

inhibitory currents, while ES measured spike timing coordination between excitatory

neuron pairs within 5-ms windows. Both metrics were normalized to baseline

unpruned values to reflect pruning-induced changes. Statistical analysis included

Pearson correlation coefficients and linear regression to quantify excitatory and

inhibitory pruning effect sizes on network dynamics. Complete details of the

computational model parameters, equations, and analysis methods can be found in

Text S2.

3. Results

3.1 Identification of Consensus Gene Co-Expression Network and Hub

Signatures

To identify conserved molecular signatures underlying EP and AD, we employed an

unbiased, genome-wide approach to construct GCNs across four independent model

datasets: two EP models (pilocarpine-induced [EP-PO] and kainic acid-induced

[EP-KA]) and two AD model datasets (AD-Castanho: [rTg4510, J20]; AD-Matarin:

[TAU, TAS10, TPM, HO-TASTPM, HET-TASTPM]). After removing only genes

with negligible or undetectable expressions, we retained all expressed genes (ranging

from 12,558 to 17,813) to avoid selection bias. We identified common and distinct

GCN modules across all model datasets (**Fig. 1A, S1**): EP-PO revealed 19 modules

(97- 1203 genes), EP-KA contained 14 modules (36-1534 genes), AD-Castanho

yielded 28 modules (55-820 genes), and AD-Matarin showed 16 modules (33-1354

genes). Correlation analysis between module eigengenes and disease status identified

disease-associated modules in each model (**Table S2**). In the EP-PO dataset, 12

modules were significantly associated with the EP genotype ($|r| > 0.35$, $P < 0.05$).

Similarly, EP-KA had two disease-associated modules, AD-Castanho contained 12

modules correlated with tau pathology, with four modules correlated with general AD

pathology, and AD-Matarin revealed five modules associated with tau pathology and

two modules related to amyloid pathology, with three modules correlated with general

AD pathology (**Fig. 1A, Table S2**).

Despite model and disease-specific differences, cross-condition preservation analysis

revealed remarkable conservation of an immune module across all datasets (**Fig. 1A**).

Using Z-summary preservation statistics and cross-tabulation approaches, we

identified a large immune-related module (turquoise) with exceptionally highest

preservation (mean Z-summary = 70 & mean median rank = 3 & Fisher's adjusted *P*

< 1E-300) across all pairwise comparisons (**Fig. 1A, Table S2**). This module was

negatively correlated with deconvoluted neuron proportion and positively correlated

with disease progression genotype in each dataset ($|r| > 0.46$, $P < 9.31E\text{-}07$) (**Table S2,**

Text S1). The strong preservation pattern of this immune module across both diseases

prompted us to construct consensus GCNs by integrating network topology

information from all datasets (**Fig. S2**). We identified a robust immune-related and

disease-associated consensus module containing 1617 genes common to all models

with 19 consensus hub signatures (**Fig. 1B, Table S2-3**).

Among these 19 consensus hub signatures, PPI and gene regulatory network centrality

analysis identified *TYROBP* as the most upstream hub signature with the highest total

degree (regulator/target ratio = 1.33, total degree = 24.19), followed by *CD68*,

TREM2, *C1Q* complement components, and *SLC11A1* (**Fig. 1B, Table S2**). GO

enrichment analysis of the hub signature network identified "synaptic pruning" (*C1Q*

complement components, *TREM2*) as the most significantly enriched biological

process with the strongest signal (**Fig. 1C, Table S2**).

Taken together, our genome-wide transcriptomic analysis revealed a highly conserved

immune module across EP and AD models, with a consensus regulatory network

centered around *TYROBP*, *TREM2*, *CD68*, and *C1Q* components that coordinate

microglia activation and synaptic pruning. This consensus molecular pathway

represents a potential central mechanism linking both neurological conditions.

3.2 Consensus GCNs Significantly Overlap with GCNs in Human Patients

To validate the translational relevance of our identified consensus GCN, we compared

it with disease-associated GCNs from patient cohorts. We employed a cross-species

approach to identify overlaps between our consensus modules and AD-associated

GCN modules from 2,114 samples across seven brain regions, as well as our newly

constructed EP patient modules from 127 EP patient hippocampal samples (**Fig. 2A,**

S3, Table S3).

Our analysis revealed significant overlaps between the consensus immune module and

multiple human AD modules across different brain regions. The strongest associations

were observed with modules from the cerebellum (CBE), dorsolateral prefrontal

cortex (DLPFC), and superior temporal gyrus (STG) regions. The consensus module

demonstrated a powerful overlap with the " CBE-turquoise" module (n = 502, 25.39%

of genes, P < 1E-300), the "DLPFC-blue" module (n = 502, 28.67% of genes, P <

1E-300) and "STG-blue" (n = 389, 33.22% of genes, P < 1E-300) (**Fig. 2A, Table S2**).

Similarly, substantial overlaps were detected with modules from the inferior frontal

gyrus (IFG), superior temporal gyrus (STG), temporal cortex (TCX), and frontal pole

(FP), with the overlap percentage varying from 2% to 33% (**Table S2**). For the EP

patient cohort, our consensus module showed significant overlap with two EP patients'

GCN modules (yellow: n = 153, 39.64% of genes, $P < 7.07E-88$; blue: n = 279, 22.85%

of genes, $P < 1.06E-84$) (**Fig. 2A, Table S2**). Notably, the top hub genes identified in

our animal model analysis—*TYROBP*, *TREM2*, and the *C1Q* components—were also

identified as central hub genes (top 2 intramodular connectivity) in the EP patient's

yellow module (**Table S3**). To further validate our hub consensus signatures findings,

we examined whether the 19 previously identified hub signatures remained clustered

within the same modules in human disease networks (**Fig. 2B**). The co-occurrence

analysis revealed strong co-expression patterns among these signatures, with

particularly strong co-occurrence observed among *C1Q* complement components,

TREM2, *TYROBP*, *CYBA*, *and SLC11A1*. These genes showed the highest

co-occurrence frequency (dark blue in the heatmap), suggesting they function as a

coordinated unit across species and disease conditions.

Collectively, these cross-species preservation analyses demonstrate that the identified

immune modules and hub signatures represent conserved biological entities relevant

to both AD and EP pathogenesis in humans.

3.3 Consensus Hub Signatures Showed Microglial Specificity Upregulation

To validate the clinical relevance of our identified consensus hub signatures, we

investigated their expression patterns in bulk and single-nucleus human cohorts and

explored their distribution across microglial subtypes.

Expression profiling across patient cell types demonstrated that these signatures were

predominantly localized to microglia (**Fig. 3A-D**), though some genes, including

CTSH, *MYO1F*, and *GUSB*, exhibited broader expression across multiple cell types.

In general, nearly all signatures showed significant upregulation in the EP cohort,

while most were also upregulated in AD patient cohorts, with particularly elevated

expression in microglia from late-stage pathology (**Fig. 3E-H**). Furthermore,

diagnostic models developed by various machine learning algorithms based on bulk

expression profiles of 19 consensus features achieved excellent discrimination

between patients and controls (EP: AUC > 0.9, AD: AUC > 0.7). (**Fig. S4, Text S1**).

Notably, *TYROBP*, *TREM2*, and *C1Q* complement genes—central consensus

signatures involved in microglial synaptic pruning—demonstrated consistent

upregulation in both disorders (**Fig. 3E-H**).

To understand the specific microglial contributions, we examined the expression

patterns of our consensus signatures across microglial subtypes using a

comprehensive single-cell RNA-sequencing dataset from Tuddenham et al.,

encompassing 215,680 microglial cells from 74 donors. Distinct expression patterns

of consensus hub signatures across 12 defined microglial subtypes were found (**Fig.**

3I). Key signatures, including *TYROBP*, *TREM2*, *CD68*, and *C1Q* complement genes,

showed prominent upregulation in clusters 10, 4, and 9, while being downregulated in

clusters 12, 7, 6, and 1. According to GO functional characterization, these

high-expressing clusters represent distinct microglial states: cluster 10 is enriched for

genes involved in antigen presentation and complement-mediated synaptic pruning,

while clusters 4 and 9 share genes associated with oxidative phosphorylation and

immune response pathways (**Fig. 3I**). In contrast, clusters showing downregulation of

our consensus signatures (clusters 1, 6, and 7) represent alternative microglial states

characterized by metabolic and transcriptional regulatory functions.

These findings demonstrate that consensus hub signatures, particularly those involved

in microglial activation and synaptic pruning, possess potential diagnostic utility and

are expressed explicitly in functionally distinct microglial subpopulations.

Immune-activated microglial subtypes, especially cluster 10, are primary mediators of

synaptic pruning, potentially contributing to the comorbidity of AD and EP.

3.4 Asymmetric Effects of Synaptic Pruning on Neural Network Dynamics

To better understand how synaptic pruning influences neural network dynamics and

its potential role in the comorbidity of AD and EP, we developed a spiking neural

network model simulating the neural circuit. In this model, we systematically varied

the extent of synaptic pruning (0% to 50%) at excitatory or inhibitory synapses and

quantified changes in the E/I ratio and synchronization levels (**Fig. 4**).

Our findings indicate that pruning excitatory or inhibitory synapses alters the E/I ratio,

but with differential effects. Inhibitory synaptic pruning significantly elevated the E/I

ratio, while excitatory pruning reduced it. These bidirectional trends were consistent

across all simulations. In experiments where excitatory or inhibitory pruning was

varied independently (**Fig. 4A**). Correlation analysis revealed a negative association

between excitatory pruning and the E/I ratio ($r = -0.474$) and a stronger positive

correlation between inhibitory pruning and the E/I ratio ($r = 0.684$) (Fig. 4B). Linear

regression further confirmed this differential sensitivity, with excitatory pruning

reducing the E/I ratio ($\beta = -1.038$). In contrast, inhibitory pruning increased more

substantially ($\beta = 1.454$), indicating that E/I balance is more sensitive to changes in

inhibitory connectivity (**Fig. 4B**).

Network synchronization exhibited an even more pronounced differential sensitivity

to pruning type. Inhibitory pruning positively correlated with event synchronization (r

$= 0.346$). In contrast, excitatory pruning showed only a negligible negative

correlation ($r = -0.039$) (**Fig. 4B**). Linear regression analysis further confirmed this

asymmetric sensitivity, with inhibitory pruning exerting a substantially more

substantial effect on synchronization ($\beta = 0.358$) compared to excitatory pruning ($\beta =$

-0.041) (**Fig. 4B**). Additionally, the E/I ratio itself was positively correlated with

synchronization ($r = 0.491$), suggesting that the influence of synaptic pruning on

network synchronization may be indirectly mediated through its effects on E/I balance

(**Fig. 4C**).

These simulations demonstrate that while both types of synaptic pruning affect

network properties, inhibitory pruning exhibits a heightened potency in disrupting

both E/I balance and synchronization dynamics. This differential sensitivity to

inhibitory pruning may represent a key mechanism by which synapse loss in

neurodegenerative diseases may trigger the hypersynchrony characteristic of network

hyperexcitability and epileptiform activity.

4. Discussion

While Alzheimer's disease (AD) and epilepsy (EP) share a complex bidirectional

relationship, the molecular mechanisms underlying their comorbidity remain poorly

understood. Our genome-wide transcriptional analysis of multiple animal models

and patient cohorts revealed several key molecular insights into this relationship,

particularly regarding shared neuroinflammatory pathways and synaptic pruning

mechanisms.

Conserved Neuroinflammation Pathways in Epilepsy and Alzheimer's Disease

Our genome-wide analysis proves that neuroinflammatory pathways form a common

molecular substrate between AD and EP (22). We identified remarkable preservation

of the immune-related turquoise module across disease models and human patient

cohorts **(Fig. 1-2**). Functional enrichment analysis of the preserved module revealed

significant enrichment in neuroinflammatory processes, such as "inflammatory

response", "cytokine production", "blood vessel morphogenesis", "glial cell

activation", and "complement and coagulation cascades" (**Table S2**).

Neuroinflammation has been extensively documented in both diseases and serves a

complex dual role in disease progression (22,44,45). While activated microglia

facilitate Aβ plaque clearance and reactive astrocytes provide metabolic support and

ionic homeostasis that may attenuate neural damage and seizures, chronic activation

or dysfunction of these glial populations can promote disease progression through

persistent neuroinflammation (37,46,47). Specifically, activated microglia and

astrocytes secrete proinflammatory cytokines and chemokines, including IL-1β,

TNF-α, and IL-6, which can enhance neuronal hyperexcitability, increase seizure

susceptibility, exacerbate Aβ and tau aggregation, and accelerate neurodegeneration

and cognitive decline (48). Additional inflammatory components, including

complement cascade activation and blood-brain barrier (BBB) dysfunction, further

link these pathologies. Complement activation contributes to synaptic loss in AD and

is elevated in TLE patient tissue. BBB disruption promotes both seizure susceptibility

and neuroinflammation (49–52).

Our results supported previous findings that identified neuroinflammation as a key

mechanistic link between AD and EP. AD-associated inflammation enhances seizure

susceptibility, and EP-induced inflammation may accelerate Aβ and tau aggregation

as well as cognitive decline.

Synaptic Pruning as Core Signatures in this Comorbidity

Our analysis of the conserved immune module revealed a central synaptic pruning

pathway involving *TYROBP*, *TREM2*, and *C1Q* complement genes (**Fig. 1**). These

genes showed consistent co-dysregulation in microglia and demonstrated diagnostic

value across both conditions (**Figs. 2-3**). The expression profile notably differentiated

between activated and alternative metabolic state microglial populations (**Fig. 3I**).

Collectively, these findings suggest that this microglia-mediated synaptic pruning may

represent a core mechanistic pathway underlying the comorbidity of AD and EP.

This pathway is well-documented in AD pathology, with *TYROBP* and *TREM2*

established as risk factors due to their effects on microglial function (53–55). Under

physiological conditions, microglia maintain synaptic homeostasis through

complement-mediated pruning, where *C1Q* tags unnecessary synapses for elimination

(56). *TREM2* and *TYROBP* recognize these tags and initiate phagocytic clearance,

which is critical for neural circuit maintenance (57). However, in AD pathology, these

receptors trigger microglia's transition to a disease-associated state (DAM),

characterized by enhanced phagocytic activity and elevated expression of *TYROBP*,

TREM2, and *C1Q* (57). While *TYROBP/TREM2* deficiency impairs plaque clearance,

pathway overactivation can also promote neurodegeneration through excessive

synaptic pruning (58–60).

The impact of synaptic pruning on AD and EP progression can operate through two

primary mechanisms: synaptic transmission and plasticity. *C1Q* inhibition in aged

models can restore both long-term potentiation (LTP) and depression (LTD), while

increased *C1Q* expression impairs LTP in non-aged models (58–61). This is

particularly relevant as impaired synaptic plasticity characterizes cognitive decline in

AD and may contribute to epileptogenic circuit formation in EP. Elevated *C1Q*

expression leads to decreased synaptic transmission in the CA1 region, affecting both

excitatory and inhibitory neurons (67–69), which can create a harmful feedback loop

where synaptic dysfunction promotes disease progression in both conditions (62,63).

Notably, interventions targeting the synaptic pruning pathway, whether through the

knockdown of regulatory genes like *TYROBP/TREM2* or direct inhibition of *C1Q*

accumulation, have successfully restored both electrophysiological and cognitive

functions in AD models. This restoration occurred even in the presence of significant

Aβ and tau pathology (57–60). Our results establish a regulatory network that is

consistent with previous observations and further extends its role in the comorbidity

of AD and EP (**Fig. 1B**). Interestingly, microglial clusters 1 and 6 showed

downregulation of consensus signatures including *TYROBP*, *TREM2*, and *C1Q* genes

(**Fig. 3I**). These clusters have been previously reported to correlate positively with

amyloid-beta and tau tangle pathology, while correlating negatively with the slope of

cognitive decline (40). Conversely, clusters 4, 9, and 10 upregulated these genes

(*TYROBP*, *TREM2*, *C1Q*) and were enriched for genes that generally correlated

negatively with amyloid-beta and tau tangle pathology, while correlating positively

with the slope of cognitive decline (**Fig. 3I**) (40). These results may reflect the

complex and stage-dependent role of microglia in the pathogenesis of AD. While

microglia are designed to reduce the burden of AD pathology, their excessive

activation may paradoxically contribute to neurodegeneration and cognitive decline.

While this pathway has been extensively studied in AD, its role in AD-EP

comorbidity remains unexplored. To address this gap and understand how synaptic

pruning affects global neural network dynamics in AD-EP comorbidity (64), we

conducted an in silico experiment.

Asymmetric Effects of Synaptic Pruning Promote Network Synchronization

Computational modeling has improved our understanding of AD and EP pathologies.

It can predict quantitative mechanisms behind them, with models ranging from

molecular-level ion dynamics simulations to large-scale network models (65,66). We

developed a neural circuit model building upon Liu's biologically plausible EP

modeling framework (42). This model allowed us to quantify how pruning of

excitatory and inhibitory synapses affects E/I balance (E/I ratio) and the temporal

coordination of neural firing (synchronization).

Our model reveals a differential sensitivity of neural networks to synaptic pruning,

with E/I balance being substantially more responsive to inhibitory synapse loss than

excitatory synapse elimination (**Fig. 4**). Excitatory synapse elimination decreased the

E/I ratio but had negligible effects on network synchronization. In contrast, inhibitory

synapse pruning resulted in a significant increase in the E/I ratio and network

synchronization (**Fig. 4**). This heightened sensitivity to inhibitory pruning can be

explained by the architectural role of inhibitory interneurons within neural circuits.

These neurons often establish synaptic connections with multiple excitatory neurons

and act as crucial circuit coordination and synchronization regulators. Therefore,

when inhibitory control mechanisms are impaired, dysregulation cascades throughout

more widespread circuits (67). Consequently, a reduction in inhibitory synapses

produces a disproportionately more significant perturbation in neural networks than

an equivalent reduction in excitatory synapses. This result implies a plausible

mechanistic pathway whereby neuronal or synapse loss in neurodegenerative diseases

disrupts E/I balance, enhancing sensitivity to inhibitory synapse loss and thereby

promoting EP susceptibility. (**Fig. 4**). This mechanism may extend beyond AD, as EP

is a well-documented comorbidity of multiple neurodegenerative conditions,

including Huntington's, Parkinson's disease, and multiple sclerosis (68). Notably, in

AD, synaptic loss often precedes cognitive decline and EP, suggesting it may be an

initiating event rather than a consequence (1).

Our results emphasize the critical role of inhibitory synaptic pruning in promoting

network synchronization due to the neural network's heightened sensitivity to

inhibitory synapse loss. At the molecular level, *C1Q* functions as an "eat me" signal,

tagging synapses for microglial elimination (64). While this tagging process is not

synapse-type specific, interestingly, several recent studies have demonstrated that

microglia may preferentially prune inhibitory synapses (64,69). Dejanovic et al.

demonstrated that the *TREM2-C1Q* microglial synaptic pruning pathway engulfs

inhibitory synapses more frequently than excitatory synapses in AD models. At the

same time, Fan et al. observed similar preferential pruning of inhibitory synapses

during recurrent seizures that exacerbate EP model pathology (69,70). Studies have

shown that *C1Q* blockade reduces both the frequency and severity of seizures in

traumatic brain injury models (71). However, it is also important to note that *C1Q*

knockout also affects synaptic elimination during development, enhancing

connectivity and seizure activity (72,73). Recently, Aloi et al. reported that

overactivation of microglia reduced Aβ pathology and caused hyperexcitability and

seizures via synaptic pruning (74).

These findings collectively suggest that the network's differential sensitivity to

inhibitory synapse loss and the apparent preferential pruning of inhibitory synapses by

microglia may represent a critical mechanism underlying AD-EP comorbidity.

Managing microglial *C1Q*-mediated synaptic pruning during AD pathology and aging

warrants further investigation, particularly regarding regulating this potential

preferential pruning of inhibitory synapses.

Conclusion

Our genome-wide transcriptional analysis of AD and EP animal models and human

cohorts revealed a consensus molecular signature converging on microglial synaptic

pruning, with a highly preserved immune module centered on the

TYROBP-TREM2-C1Q complement pathway consistently upregulated across species

and conditions. To support community validation, we developed a web application

(https://github.com/HuihongLi/OverlapModule) enabling researchers to test our

module overlap preservation with theirs. Neural circuit modeling demonstrated that

inhibitory synapse loss through complement-mediated pruning exerts

disproportionately powerful effects on E/I balance and network synchronization

compared to excitatory loss, creating a pathological loop of hyperexcitability and

hypersynchrony that mechanistically links neuroinflammation to network dysfunction

in AD-EP comorbidity. These findings provide a unifying framework explaining the

bidirectional relationship between AD and EP, suggesting that targeting

microglial-complement signaling may offer disease-modifying benefits.

Abbreviations

EP   Epilepsy

AD   Alzheimer's disease

E/I   Excitation/Inhibition

TLE   Temporal lobe epilepsy

Aβ   Amyloid-β

PO   Pilocarpine-induced

KA   Kainic acid-induced

GCNs   Gene co-expression networks

WT   Wild-type

PCA   Principal Component Analysis

BH   Benjamini-Hochberg

GO   Gene Ontology

TOM   Topological overlap matrices

MEs   Module eigengenes

MM   Module membership

GS   Gene-genotype significance

PPI   Protein-protein interaction

598	FDR	False discovery rate
599	AUC	Area Under the Receiver Operating Characteristic Curve
600	ROC	Receiver Operating Characteristic
601	CBE	Cerebellum
602	DLPFC	Dorsolateral prefrontal cortex
603	STG	Superior temporal gyrus
604	IFG	Inferior frontal gyrus
605	TCX	Temporal cortex
606	FP	Frontal pole
607	SVM	Support vector machine
608	ES	Event synchrony
609	BBB	Blood-brain barrier
610	DAM	Disease-associated microglia
611	LTP	Long-term potentiation
612	LTD	Long-term depression
613	ROSMAP	Rush Memory and Aging Project
614	snRNA-seq	Single nucleus RNA-sequencing

Figures and legends

Figure 1. Identification of conserved immune-related GCNs and hub signatures.

(A) Chord diagram showing module preservation across EP and AD models. Each

segment represents a gene module identified in different datasets. Purple connections

indicate statistically significant module overlap (Fisher's exact test, adjusted $P <$

1E-300), with color intensity corresponding to significance level (-log10(padj)). Inner

tracks display module correlations with disease status (red: positive correlation; blue:

negative correlation) and neuron proportion.

(B) Integrated PPI and regulatory network of consensus hub signatures identified

across all models. Nodes represent hub genes consistently present in the preserved

immune module, with node size proportional to total weighted degree. Solid lines

indicate validated protein-protein interactions (STRING), dotted lines indicate

computationally inferred gene regulation (GENIE3), with edge width reflecting

interaction strength. Node color indicates regulator/target ratio, with red showing

stronger regulatory potential.

(C) GO enrichment analysis of consensus hub signatures. Bar length corresponds to

enrichment signal strength, color indicates significance level (FDR), and size reflects

gene count within each category. Bars represent significantly enriched biological

processes, with "Synaptic pruning" emerging as the most enriched process with the

highest signal.

Figure 2. Comparison of consensus modules with patient modules.

(A) The overlap between the consensus immune module and disease-associated GCNs

from human patients. Peripheral nodes represent modules from different brain regions

in AD and EP patient cohorts. Line thickness indicates the number of overlapping

genes between each patient module and the consensus module, with thicker lines

showing stronger connections. Line color intensity corresponds to statistical

significance (-log10(adjusted *P*)), with red-orange hues indicating the highest

significance levels. The significant associations (adjusted *P* < 0.05) were observed

with modules across brain regions in AD patients, and with yellow and blue modules

in EP patients.

(B) Co-occurrence heatmap of the 19 identified consensus hub signatures across

human disease modules. Color intensity represents the frequency of gene

co-occurrence (darker blue indicates higher co-occurrence counts).

Figure 3. Expression of consensus hub signatures in EP and AD patient cohorts.

(A-D) UMAP visualization and dot plot showing cell type-specific expression of

consensus hub genes in EP patient samples **(A-B)** and AD patient samples **(C-D)**.

Microglia show the highest expression of most hub genes, with *TYROBP*, *TREM2*,

and *C1Q* complement genes exhibiting powerful expression.

(E-H) Differential expression analysis of consensus hub genes in microglia and bulk

tissue from EP patients (**E-F**) and AD patients (**G-H**) compared to controls. Bar

heights represent fold changes with statistical significance indicated (*p < 0.05, **p <

0.01, ***p < 0.001).

(I) Heatmap showing the expression patterns (average log fold change) of 19

consensus gene signatures across 12 distinct microglial clusters derived from

Tuddenham's data. The color scale represents average log fold change (logFC) or

statistical significance for GO enrichment analysis. The dendrogram shows

hierarchical clustering of both genes and microglial subtypes.

Figure 4. Neural circuit modeling of synaptic pruning

(**A**) Simulation results of excitatory pruning ratio (0-0.5), inhibitory pruning ratio

(0-0.5), E/I ratio, and synchronization. X-axis: Excitatory pruning ratio; Y-axis:

Inhibitory pruning ratio; Z-axis: Average E/I ratio. Colors indicate average

synchronization levels from dark purple (close to 1, indicating no change in

synchrony) to yellow (indicating higher synchrony).

(**B**) The distribution of E/I ratio and synchronization values across different pruning

levels. Each violin represents the distribution of values at a given pruning ratio (0–0.5

in 0.05 increments). Embedded boxplots display median and interquartile ranges.

Overlaid dots represent individual simulation outcomes. Regression results (slope,

p-value, and Pearson's r) are shown in the upper right of each panel, indicating that

inhibitory pruning has a stronger positive correlation with both the E/I ratio and

synchronization. In contrast, excitatory pruning shows weaker correlations.

(**C**) The E/I ratio and synchronization relationship across all simulation conditions. A

positive correlation is observed, suggesting that changes in E/I balance mediate

alterations in network synchrony. Regression statistics are displayed in the corner.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

The datasets analyzed during the current study are available in the National Center for

Biotechnology Information (NCBI), European Nucleotide Archive (ENA), and

Synapse under the following accession numbers: PRJEB18790, GSE125957,

GSE73878, GSE64398, GSE63808, E-MTAB-3123, GSE48350, GSE190453,

syn51758062.1, GSE204702.

Competing Interests

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Funding

This research was supported by the Young Scientists Fund of the National Natural Science Foundation of China (Grant Nos. 82201629) and the Guangdong Provincial Science and Technology Special Fund (Grant Nos. 210803136872196)

Authors' contribution

HL conceptualized the study, conducted the investigation, developed the methodology, performed formal analysis, and contributed to writing the original draft and reviewing/editing the manuscript. ZX, YT, RZ, YY, and BL contributed to the investigation, performed formal analysis, and participated in writing the original draft and reviewing/editing the manuscript. SC, ZD, and JL contributed to writing the original draft and reviewing/editing the manuscript. JW was involved in conceptualization and contributed to writing the original draft and reviewing/editing the manuscript. MC, XL, and YS contributed to the investigation and participated in reviewing/editing the manuscript. BH and NW conceptualized the study and contributed to writing the original draft and reviewing/editing the manuscript. XJ provided supervision, acquired funding, contributed to conceptualization, and participated in writing the original draft and reviewing/editing the manuscript. All authors read and approved of the final manuscript.

Acknowledgments

The results published here are in whole or in part based on data obtained from the **AD**

Knowledge Portal.

References

1. Vossel KA, Beagle AJ, Rabinovici GD, Shu H, Lee SE, Naasan G, et al. Seizures and Epileptiform Activity in the Early Stages of Alzheimer Disease. JAMA Neurol. 2013 Sep 1;70(9):1158–66.

2. Palop JJ, Mucke L. Epilepsy and Cognitive Impairments in Alzheimer Disease. Arch Neurol. 2009 Apr 1;66(4):435–40.

3. Fang Y, Si X, Wang J, Wang Z, Chen Y, Liu Y, et al. Alzheimer Disease and Epilepsy. Neurology. 2023 Jul 25;101(4):e399–409.

4. Prince M, Wimo A, Guerchet M, Ali GC, Wu YT, Prina M. World Alzheimer Report 2015. The Global Impact of Dementia: An analysis of prevalence, incidence, cost and trends. Alzheimer's Disease International; 2015.

5. Ngugi AK, Bottomley C, Kleinschmidt I, Sander JW, Newton CR. Estimation of the burden of active and life-time epilepsy: A meta-analytic approach. Epilepsia. 2010;51(5):883–90.

6. Mendez MF, Lim GTH. Seizures in Elderly Patients with Dementia. Drugs Aging. 2003 Sep 1;20(11):791–803.

7. Pandis D, Scarmeas N. Seizures in Alzheimer Disease: Clinical and Epidemiological Data: Seizures in Alzheimer Disease. Epilepsy Curr. 2012 Sep 1;12(5):184–7.

8. Leitner D, Pires G, Kavanagh T, Kanshin E, Askenazi M, Ueberheide B, et al. Similar brain proteomic signatures in Alzheimer's disease and epilepsy. Acta Neuropathol (Berl). 2024 Jan 30;147(1):27.

9. Tombini M, Assenza G, Ricci L, Lanzone J, Boscarino M, Vico C, et al. Temporal Lobe Epilepsy and Alzheimer's Disease: From Preclinical to Clinical Evidence of a Strong Association. J Alzheimers Dis Rep. 2021 Mar 11;5(1):243–61.

10. Johnson EL, Krauss GL, Lee AK, Schneider ALC, Dearborn JL, Kucharska-Newton AM, et al. Association Between Midlife Risk Factors and Late-Onset Epilepsy: Results From the Atherosclerosis Risk in Communities Study. JAMA Neurol. 2018 Nov 1;75(11):1375–82.

11. Stefanidou M, Beiser AS, Himali JJ, Peng TJ, Devinsky O, Seshadri S, et al.
Bi-directional association between epilepsy and dementia. Neurology. 2020 Dec
15;95(24):e3241–7.

12. Noebels J. A perfect storm: Converging paths of epilepsy and Alzheimer's
dementia intersect in the hippocampal formation. Epilepsia. 2011;52(s1):39–46.

13. Gourmaud S, Shou H, Irwin DJ, Sansalone K, Jacobs LM, Lucas TH, et al.
Alzheimer-like amyloid and tau alterations associated with cognitive deficit in
temporal lobe epilepsy. Brain. 2020 Jan 1;143(1):191–209.

14. Tai XY, Koepp M, Duncan JS, Fox N, Thompson P, Baxendale S, et al.
Hyperphosphorylated tau in patients with refractory epilepsy correlates with
cognitive decline: a study of temporal lobe resections. Brain. 2016 Sep
1;139(9):2441–55.

15. Nelson PT, Schmitt FA, Lin Y, Abner EL, Jicha GA, Patel E, et al. Hippocampal
sclerosis in advanced age: clinical and pathological features. Brain. 2011 May
1;134(5):1506–18.

16. Zou Y, Wang C, Li H, Zhong M, Lin J, Hu Y, et al. Epileptic seizures induced by
pentylenetetrazole kindling accelerate Alzheimer-like neuropathology in 5×FAD
mice. Front Pharmacol. 2024 Oct 31;15:1500105.

17. Watts ME, Pocock R, Claudianos C. Brain Energy and Oxygen Metabolism:
Emerging Role in Normal Function and Disease. Front Mol Neurosci. 2018 Jun
22;11:216.

18. Born HA. Seizures in Alzheimer's disease. Neuroscience. 2015 Feb
12;286:251–63.

19. Toniolo S, Sen A, Husain M. Modulation of Brain Hyperexcitability: Potential
New Therapeutic Approaches in Alzheimer's Disease. Int J Mol Sci. 2020
Jan;21(23):9318.

20. Palop JJ, Chin J, Roberson ED, Wang J, Thwin MT, Bien-Ly N, et al. Aberrant
Excitatory Neuronal Activity and Compensatory Remodeling of Inhibitory
Hippocampal Circuits in Mouse Models of Alzheimer's Disease. Neuron. 2007
Sep 6;55(5):697–711.

21. Targa Dias Anastacio H, Matosin N, Ooi L. Neuronal hyperexcitability in
Alzheimer's disease: what are the drivers behind this aberrant phenotype? Transl
Psychiatry. 2022 Jun 22;12(1):1–14.

22. Liew Y, Retinasamy T, Arulsamy A, Ali I, Jones NC, O'Brien TJ, et al.

Neuroinflammation: A Common Pathway in Alzheimer's Disease and Epilepsy. J
Alzheimer's Dis. 2023 Jul 25;94(s1):S253–65.

23. Kalyvas AC, Dimitriou M, Ioannidis P, Grigoriadis N, Afrantou T. Alzheimer's
Disease and Epilepsy: Exploring Shared Pathways and Promising Biomarkers for
Future Treatments. J Clin Med. 2024 Jan;13(13):3879.

24. Giorgi FS, Saccaro LF, Busceti CL, Biagioni F, Fornai F. Epilepsy and
Alzheimer's Disease: Potential mechanisms for an association. Brain Res Bull.
2020 Jul 1;160:107–20.

25. Bosco F, Guarnieri L, Rania V, Palma E, Citraro R, Corasaniti MT, et al.
Antiseizure Medications in Alzheimer's Disease from Preclinical to Clinical
Evidence. Int J Mol Sci. 2023 Jan;24(16):12639.

26. Srivastava PK, Bagnati M, Delahaye-Duriez A, Ko JH, Rotival M, Langley SR, et
al. Genome-wide analysis of differential RNA editing in epilepsy. Genome Res.
2017 Mar 1;27(3):440–50.

27. Matarin M, Salih DA, Yasvoina M, Cummings DM, Guelfi S, Liu W, et al. A
Genome-wide Gene-Expression Analysis and Database in Transgenic Mice during
Development of Amyloid or Tau Pathology. Cell Rep. 2015 Feb 3;10(4):633–44.

28. Castanho I, Murray TK, Hannon E, Jeffries A, Walker E, Laing E, et al.
Transcriptional Signatures of Tau and Amyloid Neuropathology. Cell Rep. 2020
Feb 11;30(6):2040-2054.e5.

29. Johnson MR, Behmoaras J, Bottolo L, Krishnan ML, Pernhorst K, Santoscoy
PLM, et al. Systems genetics identifies Sestrin 3 as a regulator of a proconvulsant
gene network in human epileptic hippocampus. Nat Commun. 2015 Jan
23;6(1):6031.

30. Wan YW, Al-Ouran R, Mangleburg CG, Perumal TM, Lee TV, Allison K, et al.
Meta-analysis of the Alzheimer's disease human brain transcriptome and
functional dissection in mouse models. Cell Rep. 2020;32(2).

31. Miller JA, Horvath S, Geschwind DH. Divergence of human and mouse brain
transcriptome highlights Alzheimer disease pathways. Proc Natl Acad Sci. 2010
Jul 13;107(28):12698–703.

32. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation
network analysis. BMC Bioinformatics. 2008 Dec 29;9(1):559.

33. Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal
enrichment tool for interpreting omics data. The innovation. 2021;2(3).

34. Langfelder P, Luo R, Oldham MC, Horvath S. Is My Network Module Preserved
and Reproducible? PLOS Comput Biol. 2011 Jan 20;7(1):e1001057.

35. Szklarczyk D, Nastou K, Koutrouli M, Kirsch R, Mehryary F, Hachilif R, et al.
The STRING database in 2025: protein networks with directionality of regulation.
Nucleic Acids Res. 2025 Jan 6;53(D1):D730–7.

36. Huynh-Thu VA, Irrthum A, Wehenkel L, Geurts P. Inferring Regulatory Networks
from Expression Data Using Tree-Based Methods. PLOS ONE. 2010 Sep
28;5(9):e12776.

37. Chen ZP, Wang S, Zhao X, Fang W, Wang Z, Ye H, et al. Lipid-accumulated
reactive astrocytes promote disease progression in epilepsy. Nat Neurosci. 2023
Apr;26(4):542–54.

38. Mathys H, Davila-Velderrain J, Peng Z, Gao F, Mohammadi S, Young JZ, et al.
Single-cell transcriptomic analysis of Alzheimer's disease. Nature. 2019
Jun;570(7761):332–7.

39. Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, et al.
Dictionary learning for integrative, multimodal and scalable single-cell analysis.
Nat Biotechnol. 2024;42(2):293–304.

40. Tuddenham JF, Taga M, Haage V, Marshe VS, Roostaei T, White C, et al. A
cross-disease resource of living human microglia identifies disease-enriched
subsets and tool compounds recapitulating microglial states. Nat Neurosci. 2024
Dec;27(12):2521–37.

41. Stimberg M, Brette R, Goodman DF. Brian 2, an intuitive and efficient neural
simulator. elife. 2019;8:e47314.

42. Liu Y, Grigorovsky V, Bardakjian B. Excitation and Inhibition Balance
Underlying Epileptiform Activity. IEEE Trans Biomed Eng. 2020
Sep;67(9):2473–81.

43. Penn Y, Segal M, Moses E. Network synchronization in hippocampal neurons.
Proc Natl Acad Sci. 2016 Mar 22;113(12):3341–6.

44. Vezzani A, French J, Bartfai T, Baram TZ. The role of inflammation in epilepsy.
Nat Rev Neurol. 2011 Jan;7(1):31–40.

45. Heneka MT, Golenbock DT, Latz E. Innate immunity in Alzheimer's disease. Nat
Immunol. 2015 Mar;16(3):229–36.

46. Pekny M, Pekna M. Astrocyte Reactivity and Reactive Astrogliosis: Costs and

Benefits. Physiol Rev. 2014 Oct;94(4):1077–98.

47. Onyango IG, Jauregui GV, Čarná M, Bennett JP, Stokin GB. Neuroinflammation

in Alzheimer's Disease. Biomedicines. 2021 May;9(5):524.

48. Vezzani A, Viviani B. Neuromodulatory properties of inflammatory cytokines and
their impact on neuronal excitability. Neuropharmacology. 2015 Sep 1;96:70–82.

49. van Vliet EA, Aronica E, Gorter JA. Blood–brain barrier dysfunction, seizures
and epilepsy. Semin Cell Dev Biol. 2015 Feb 1;38:26–34.

50. Zlokovic BV. Neurovascular pathways to neurodegeneration in Alzheimer's
disease and other disorders. Nat Rev Neurosci. 2011 Dec;12(12):723–38.

51. Kopczynska M, Zelek WM, Vespa S, Touchard S, Wardle M, Loveless S, et al.
Complement system biomarkers in epilepsy. Seizure. 2018 Aug 1;60:1–7.

52. Veerhuis R, Nielsen HM, Tenner AJ. Complement in the brain. Mol Immunol.
2011 Aug 1;48(14):1592–603.

53. Roussos P, Katsel P, Fam P, Tan W, Purohit DP, Haroutunian V. The triggering
receptor expressed on myeloid cells 2 (*TREM2*) is associated with enhanced
inflammation, neuropathological lesions and increased risk for Alzheimer's
dementia. Alzheimers Dement. 2015 Oct 1;11(10):1163–70.

54. Ma J, Jiang T, Tan L, Yu JT. TYROBP in Alzheimer's Disease. Mol Neurobiol.
2015 Apr 1;51(2):820–6.

55. Jonsson T, Stefansson H, Steinberg S, Jonsdottir I, Jonsson PV, Snaedal J, et al.
Variant of TREM2 Associated with the Risk of Alzheimer's Disease. N Engl J
Med. 2013 Jan 10;368(2):107–16.

56. Stevens B, Allen NJ, Vazquez LE, Howell GR, Christopherson KS, Nouri N, et al.
The Classical Complement Cascade Mediates CNS Synapse Elimination. Cell.
2007 Dec 14;131(6):1164–78.

57. Haure-Mirande JV, Audrain M, Ehrlich ME, Gandy S. Microglial
TYROBP/DAP12 in Alzheimer's disease: Transduction of physiological and
pathological signals across TREM2. Mol Neurodegener. 2022 Aug 24;17(1):55.

58. Haure-Mirande JV, Audrain M, Fanutza T, Kim SH, Klein WL, Glabe C, et al.
Deficiency of TYROBP, an adapter protein for TREM2 and CR3 receptors, is
neuroprotective in a mouse model of early Alzheimer's pathology. Acta

Neuropathol (Berl). 2017 Nov 1;134(5):769–88.

59. Audrain M, Haure-Mirande JV, Wang M, Kim SH, Fanutza T, Chakrabarty P, et al. Integrative approach to sporadic Alzheimer's disease: deficiency of TYROBP in a tauopathy mouse model reduces C1q and normalizes clinical phenotype while increasing spread and state of phosphorylation of tau. Mol Psychiatry. 2019 Sep;24(9):1383–97.

60. Haure-Mirande JV, Wang M, Audrain M, Fanutza T, Kim SH, Heja S, et al. Integrative approach to sporadic Alzheimer's disease: deficiency of TYROBP in cerebral Aβ amyloidosis mouse normalizes clinical phenotype and complement subnetwork molecular pathology without reducing Aβ burden. Mol Psychiatry. 2019 Mar;24(3):431–46.

61. Hu Y, Qi H, Yang J, Wang F, Peng X, Chen X, et al. Wogonin mitigates microglia-mediated synaptic over-pruning and cognitive impairment following epilepsy. Phytomedicine. 2024 Dec 1;135:156222.

62. Bie B, Wu J, Foss JF, Naguib M. Activation of mGluR1 Mediates C1q-Dependent Microglial Phagocytosis of Glutamatergic Synapses in Alzheimer's Rodent Models. Mol Neurobiol. 2019 Aug 1;56(8):5568–85.

63. Wu J, Bie B, Foss JF, Naguib M. Amyloid Fibril–Induced Astrocytic Glutamate Transporter Disruption Contributes to Complement C1q-Mediated Microglial Pruning of Glutamatergic Synapses. Mol Neurobiol. 2020 May 1;57(5):2290–300.

64. Andoh M, Ikegaya Y, Koyama R. Synaptic Pruning by Microglia in Epilepsy. J Clin Med. 2019 Dec;8(12):2170.

65. Yuan P, Zhang M, Tong L, Morse TM, McDougal RA, Ding H, et al. PLD3 affects axonal spheroids and network defects in Alzheimer's disease. Nature. 2022 Dec;612(7939):328–37.

66. Stefanescu RA, Shivakeshavan RG, Talathi SS. Computational models of epilepsy. Seizure. 2012 Dec 1;21(10):748–59.

67. Luo L. Architectures of neuronal circuits. Science. 2021 Sep 3;373(6559):eabg7285.

68. Neri S, Mastroianni G, Gardella E, Aguglia U, Rubboli G. Epilepsy in neurodegenerative diseases. Epileptic Disord. 2022;24(2):249–73.

69. Dejanovic B, Wu T, Tsai MC, Graykowski D, Gandham VD, Rose CM, et al. Complement C1q-dependent excitatory and inhibitory synapse elimination by astrocytes and microglia in Alzheimer's disease mouse models. Nat Aging. 2022

Sep;2(9):837–50.

70. Fan J, Dong X, Tang Y, Wang X, Lin D, Gong L, et al. Preferential pruning of
inhibitory synapses by microglia contributes to alteration of the balance between
excitatory and inhibitory synapses in the hippocampus in temporal lobe epilepsy.
CNS Neurosci Ther. 2023;29(10):2884–900.

71. Holden SS, Grandi FC, Aboubakr O, Higashikubo B, Cho FS, Chang AH, et al.
Complement factor C1q mediates sleep spindle loss and epileptic spikes after
mild brain injury. Science. 2021 Sep 10;373(6560):eabj2685.

72. Chu Y, Jin X, Parada I, Pesic A, Stevens B, Barres B, et al. Enhanced synaptic
connectivity and epilepsy in C1q knockout mice. Proc Natl Acad Sci. 2010 Apr
27;107(17):7975–80.

73. Ma Y, Ramachandran A, Ford N, Parada I, Prince DA. Remodeling of dendrites
and spines in the C1q knockout model of genetic epilepsy. Epilepsia.
2013;54(7):1232–9.

74. Aloi MS, Prater KE, Sánchez REA, Beck A, Pathan JL, Davidson S, et al.
Microglia specific deletion of miR-155 in Alzheimer's disease mouse models
reduces amyloid-β pathology but causes hyperexcitability and seizures. J
Neuroinflammation. 2023 Mar 7;20(1):60.

Figure 1

Figure 2

Figure 3

Figure 4