Shared Transcriptomic Signatures Reveal Synaptic Pruning as a Link Between Alzheimer's Disease and Epilepsy

Huihong Li[1,2,3,4], Zhiran Xie[4], Yuxuan Tian[4], Bingying Lin[1,2,3,5], Jianwei Li[1,2,3,5], Jie Wu[1,2,3], Mingjie Chen[6], Xueke Liu[6], Yushan Sun[6], Si Chen[1,2,3,5], Zihan Deng[1,2,3,5], Mengqi Hong[1,2,3,5], Bing Huang[1,2,3,5], Naili Wei[1], Xiaoyu Ji[1,2,3,5*]

[1]Department of Neurosurgery, The First Affiliated Hospital of Shantou University Medical College, Shantou, Guangdong 515041, China

[2]Brain Function and Disease Laboratory, Shantou University Medical College, Shantou, Guangdong 515041, China

[3]Department of Pharmacology, Shantou University Medical College, Shantou, Guangdong 51504, China

[4]Program in Computational Biology and Biomedical Informatics, Yale University, New Haven, Connecticut, United States of America

[5]Pain and Related Diseases Research Laboratory, Shantou University Medical College, Shantou, Guangdong 515041, China

[6]Department of Biotechnology and Food Engineering, Guangdong Technion – Israel Institute of Technology, Shantou, Guangdong 515063，China

[*]**Corresponding authors**

Contact: Dr. Xiaoyu Ji, xyji@stu.edu.cn

Abstract

Alzheimer's disease (AD) and epilepsy (EP) exhibit a complex, bidirectional relationship, however, the molecular mechanisms underlying their comorbidity remain poorly understood. To address this gap, we analyzed large-scale transcriptomic datasets from pilocarpine-induced EP mouse models (n = 200), two AD mouse models expressing human tau (rTg4510) or amyloid precursor protein (J20) (n = 141), and transcriptomic profiles from AD and EP patient cohorts. Differential expression and weighted gene co-expression network analyses identified a highly conserved immune-related module enriched in microglia-specific genes, particularly *Tyrobp*, *Trem2*, and *C1q* complement component genes (*C1qa*, *C1qb*, *C1qc*) implicated in the synaptic pruning pathway. Gene regulatory network analysis showed that *Tyrobp* regulates these *C1q* components gene through *Trem2*. These hub genes were markedly upregulated in both models and human patient datasets, preserved their regulatory relationship across species, and displayed strong diagnostic value. Cell composition deconvolution and single-cell transcriptomics confirmed their high, cell-type-specific expression in microglia from both animal model and human patient datasets. Moreover, computational modeling indicated that excessive synaptic pruning, particularly of inhibitory synapses, alters the excitation/inhibition (E/I) balance toward excitability that increases neural network synchrony. Collectively, our findings reveal how microglial complement-driven synapse elimination may promote both seizure activity and cognitive decline. We suggest that microglia play a central role in

the comorbidity of EP and AD, and further exploration of this synaptic pruning

pathway in the context of EP may provide promising insights into the treatment of AD

and EP comorbidity.

Keywords: Alzheimer's disease, epilepsy, synaptic pruning, computational

neuroscience, modeling, transcriptome, *Tyrobp, Trem2, C1q*

Graphic abstract

Highlights

- Transcriptomic analysis across epilepsy (EP) and Alzheimer's disease (AD) models and patient datasets revealed a conserved immune gene module.

- Shared transcriptomic signatures identified in animal models show strong diagnostic value in both AD and EP patient datasets.

- *Tyrobp-Trem2-C1q* microglial synaptic pruning pathway was identified as a central shared molecular mechanism across model and patient datasets.

- Computational modeling revealed that excessive synaptic pruning-particularly of inhibitory synapses-alters E/I balance toward excitability and increases synchrony in neural network.

1. Background

Alzheimer's disease (AD) and epilepsy (EP) are two common neurological diseases
with a complex bidirectional relationship that contributes to their respective
pathogenesis [1–3]. AD, characterized by progressive cognitive decline and
neurodegeneration is the leading cause of dementia worldwide, affecting an estimated
55 million individuals. This number is projected to triple by 2050 as life expectancy
increases [4]. EP, characterized by recurrent unprovoked seizures, affects over 50
million individuals worldwide with a particularly high incidence among elderly
populations, especially those with neurodegenerative conditions such as AD [5,6].

Epidemiological studies demonstrate strong association between AD and EP. AD
patients show increased susceptibility to EP, with 10–22% experiencing unprovoked
seizures, while 22–54% demonstrating subclinical epileptiform activity during disease
progression [7,8]. These seizures, predominantly occurring in the temporal lobe,
correlate with accelerated cognitive decline and enhanced AD pathology [9]. Conversely,
EP patients exhibit a two-fold higher risk of developing dementia compared to
controls, with late-onset EP patients showing a three-fold increased risk [10,11]. This
elevated risk is particularly evident in temporal lobe epilepsy (TLE), where numerous
pathological features overlap with AD [9].

Both conditions share overlapping pathological features, particularly in TLE,

including amyloid deposition, tau accumulation, and hippocampal sclerosis [12,13].

Chronic seizures increase neuronal activity and metabolic demands, potentially

accelerating the production and aggregation of amyloid-β (aβ) and

hyperphosphorylated tau protein [14,15]. Conversely, the progressive accumulation of

these pathological proteins in AD may lower seizure thresholds, promoting

hyperexcitability and epileptiform activity [16,17]. Furthermore, neuroinflammation,

synaptic loss, and microglial activation, features of both AD and TLE, may

synergistically accelerate neuronal damage and network dysfunction [18,19], potentially

explaining the high comorbidity rates observed clinically.

The molecular mechanisms underlying this comorbidity remain incompletely

understood, though mounting evidence suggests shared molecular pathways driving

neuronal hyperexcitability and protein aggregation in both disorders [20]. Recent

findings implicate dysregulated protein homeostasis, mitochondrial dysfunction, and

chronic inflammatory responses as central contributors [7,18]. Additionally, alterations in

glutamatergic and GABAergic signaling, alongside disruptions in synaptic plasticity

pathways, appear to converge in both conditions, particularly within the hippocampal

region [21].

Treating epilepsy in AD patients presents a complex clinical challenge, particularly

regarding the decision to treat seizures and the selection of appropriate antiseizure

medications [22]. Emerging therapeutic approaches, including targeted immunotherapies

and novel small molecules, show promise but require a deeper understanding of the

shared pathological mechanisms. Therefore, systems biology and computational

modeling approaches are urgently needed to integrate insights from animal models

and patients. Such approaches could reveal common molecular pathways, cellular

processes, and neural network dynamics alterations underlying this comorbidity,

which is crucial for developing more effective, targeted therapeutic strategies.

We hypothesize that conserved molecular mechanism exist in AD and EP animal

models and human patients. To address this, we (1) analyzed and integrated

large-scale transcriptomic data from pilocarpine-induced EP mouse models (n = 200),

two AD mouse models expressing human tau (rTg4510) or amyloid precursor protein

(J20) (n = 141) to identify shared transcriptional signatures and pathways; (2)

validated these findings in AD and EP human patient datasets; and (3) developed

computational models to link molecular insights to network mechanisms of

epileptogenesis. Our analysis identified a conserved synaptic pruning pathway

involving *Tyrobp*, *Trem2*, and *C1q* complement genes. Computational modeling

further suggested that preferential synaptic elimination of inhibitory synapses

contributes to neuronal network excitability and the cognitive decline associated with

this comorbidity.

2. Materials and Methods

2.1 Data Collection

A comprehensive database search was performed to retrieve transcriptomic data from

both AD and EP models and human patients. Databases including Gene Expression

Omnibus (GEO) and European Nucleotide Archive (ENA), were searched using the

keywords "epilepsy" and "Alzheimer's disease". We downloaded and reanalyzed two

large-scale bulk RNA-seq datasets from EP and AD mouse models. The EP model

dataset was obtained from the ENA (PRJEB18790) and includes bulk RNA-seq data

from 200 hippocampi of pilocarpine-induced EP model mice and wild-type (WT)

control. The AD dataset was downloaded from GEO (GSE125957) and includes bulk

RNA-seq data from 141 samples of the entorhinal cortex and hippocampus of AD

model mice that expressed either human tau (rTg4510, TG) or amyloid precursor

protein (J20), with their corresponding healthy controls at various ages (range from 2

to 12 months).

We also analyzed multiple large-scale transcriptome datasets from AD and EP patients

to validate findings from animal models at the gene, pathway, and cellular levels. For

cross-species validation of disease co-expression modules, we used 129 hippocampal

samples from TLE patients (GSE63808) and 97 middle temporal gyrus samples from

AD patients (GSE132903). To evaluate the diagnostic value of shared transcriptional

signatures, we analyzed 47 hippocampal samples from EP patients and controls

(E-MTAB-3123), along with 253 different brain region samples from AD patients and

controls (GSE48350). Finally, single-cell RNA sequencing (scRNA-Seq) data from

85,780 immune cells of six drug-refractory EP patients (GSE201048) and

single-nucleus RNA sequencing (snRNA-Seq) data from 174,420 immune cells from

217 AD patients and 226 controls (GSE227223) were employed to identify the cell

types and subtypes expressing these shared transcriptional signatures.

2.2 Quality control and preprocessing

All downloaded raw RNA sequences from AD and EP model datasets underwent

quality control and trimming to obtain clean sequences, using FastQC (0.12.1) and

Fastp (0.22.0) with stringent parameters (--qualified_quality_phred 30 --n_base_limit

5 --detect_adapter_for_pe --cut_tail --trim_front1 15 --dedup --trim_poly_x

--trim_poly_g) to ensure high data quality [23,24]. Ribodetector (0.2.7) was used to

remove potential rRNA contamination [25]. Low-quality regions in the sequences,

attributed to instability in the sequencing platform, were also trimmed using Fastp.

Trimmed sequences were then aligned to the GRCm39.111 reference mouse genome

using HISAT2 (2.2.1) [26] and followed by quantification using featureCounts (2.0.6) [27].

For the AD model dataset, samples were excluded from subsequent analysis based on

the criteria outlined in the original publication, resulting in 121 high-quality RNA-seq

samples. For the EP model dataset, duplicate and low-quality samples were initially

removed. Then a variance-stabilizing transformation was applied to the count matrix

before performing Principal Component Analysis (PCA) to identify potential outlier

samples. PCA was then performed to identify outliers, with Mahalanobis distances

calculated from PCA scores. A chi-squared distribution threshold of 0.975 was used to

detect significant outliers, resulting in 11 samples subsequently excluded from

analysis. Detailed information on quality control and mapping metrics is provided in

Table S1. Following quality control, mapping, and exclusion of low-quality samples,

no significant differences in read depth were observed between model groups and

controls.

2.3 Differential Expression Analysis

Differential expression analysis was perform using DESeq2 (1.46.0) [28]. Before

analysis, non-expressed and lowly expressed genes were filtered out to ensure that

genes had at least 20 counts per sample in each design group. The DESeq function

was then used to estimate size factors and dispersions and to perform statistical testing

for differential expression [28]. For the EP model dataset, results were extracted by

comparing the "EP" condition to the "WT" condition. For the AD dataset, we aimed to

detect progressive differentially expressed genes (DEGs), as describe previously [29],

using DESeq's likelihood-ratio test to fit the statistical model: *Gene expression =*

Genotype + Age + Genotype×Age (Interaction). p-values were adjusted for multiple

testing using the Benjamini-Hochberg (BH) correction.

DEGs were identified based on an adjusted p-value < 0.05 and |log2 fold change| > 1.

These thresholds were applied to the EP model dataset and to the final time point of

the AD dataset. Shared DEGs between the EP and AD model datasets were identified

by taking the intersection of their DEG results. Kyoto Encyclopedia of Genes and

Genomes (KEGG) pathway and gene ontology (GO) analyses for shared DEGs were

performed using WebGestalt (2024), with weighted set cover redundancy removal [30].

2.4 Weighted Gene Co-expression Network Construction

To construct weighted gene co-expression networks and compare results from EP and

AD model datasets, we employed a standard meta-WGCNA comparison method [31–33].

This method is particularly suitable for comparative analysis across datasets to

identify conserved gene modules, thus resolving batch effects between datasets. All

datasets underwent variance-stabilizing transformation using DESeq2 (1.46.0) to

normalize the variance across different samples. We then calculated the median

absolute deviation (MAD) and retained the top 85% of genes with the highest MAD

in each dataset, resulting in 10,997 genes common to all datasets for comparability in

further analyses. Outliers were identified through hierarchical clustering, excluding

two samples from the TG dataset and three from the J20 dataset. To assess dataset

comparability, general network properties such as average gene expression and

overall network connectivity were calculated.

We constructed signed co-expression networks using the WGCNA R package (1.73)

[32], selecting the lowest soft-thresholding power ($\beta = 9$) to achieve a scale-free

topology fit index (R^2) of at least 0.85 across datasets. Adjacency matrices were

generated using the signed network option and converted to topological overlap

matrices (TOM) to quantify similarity in gene connectivity profiles. Modules were

identified by average linkage hierarchical clustering of TOM dissimilarity (1-TOM),

followed by dynamic tree cutting with parameters deepSplit = 2 and minClusterSize =

24. This procedure was also applied to patient microarray datasets (GSE63808 and

GSE132903), using variance-stabilizing normalization with the "limma" (3.62.1)

package [34].

To match modules between EP and AD model datasets, we color-mapped the AD

modules to the best matching EP modules according to the hypergeometric test [31].

Module profiles were summarized using module eigengenes (MEs), derived from the

first principal component of each module. To identify modules significantly

associated with disease genotypes (EP or interaction of TG/J20), we calculated

Pearson correlation coefficients between MEs and genotype information, with significance defined as $p < 0.05$ and an absolute correlation coefficient > 0.65 [35]. Module membership (MM) and gene-genotype significance (GS) were also determined by calculating Pearson correlations between individual genes and MEs or genotypes, respectively. KEGG pathway and GO analyses for module annotation were conducted using WebGestalt (2024) with weighted set cover redundancy removal [30].

2.5 Module Preservation Analysis

To assess module preservation of the EP model module in AD model datasets, we used two approaches [36]: Zsummary preservation statistics and cross-tabulation-based preservation statistics.

We employed the modulePreservation function from the WGCNA R package, with parameters set to networkType = "signed", nPermutations = 100, to calculate composite Z-summary preservation statistics. This metric evaluates preservation based on module size, network density, and connectivity, offering a comprehensive assessment of how well the structure of EP modules is maintained in AD datasets. The Z-summary thresholds were interpreted as follows: Z-summary > 10 indicates strong evidence of preservation, $2 <$ Z-summary < 10 suggests moderate evidence, and Z-summary < 2 indicates no evidence of preservation [36]. To complement the Z-summary statistic, we also used the medianRank preservation statistic, which

assesses relative preservation across modules with less dependency on module size. A

lower median rank denotes stronger observed preservation [36].

Additionally, we used a cross-tabulation method to examine module overlap between

EP and AD model datasets, as well as between model and patient datasets (GSE63808

and GSE132903). This method generates a contingency table reporting the

overlapping gene number between each module in the reference and test datasets [36].

The statistical significance of overlaps was assessed using Fisher's exact test, with

p-values adjusted using the BH correction. A module was reported as significantly

overlapping if the overlap count was > 30 and the adjusted p-value was < 0.05.

2.6 Hub Gene Identification and Regulatory Network Analysis

Hub genes act as key drivers within a module and were identified using stringent

criteria to minimize false positives: genes ranking in the top 10% for intramodular

connectivity within each module, MM > 0.9, and GS > 0.4. Shared hub genes were

determined by intersecting the hub genes of key modules in both EP and AD datasets,

along with the DEGs. The expression and diagnostic value of these shared hub genes

were further explored in the AD and EP patient datasets (GSE48350 and

E-MTAB-3123).

To further explore interactions and potential regulatory relationships among shared

hub genes in the EP and AD model datasets, we constructed a protein-protein

interaction (PPI) network using the STRING database (12.0)

(https://string-db.org/cgi/input.pl) with a confidence score threshold of 0.4 and a false

discovery rate (FDR) of 5% [37]. We then performed regulatory network analysis with

the GENIE3 (1.28.0) algorithm, which applies a random forest approach to infer

potential regulatory relationships from gene expression data. GENIE3 ranks

regulatory interactions by weight, with higher weights indicating stronger regulatory

links [38]. Gene expression profiles of the key module from the EP and TG model

datasets were used to infer plausible regulatory interactions among shared hub genes.

To reduce false positives, a weight threshold of 0.01 was applied to filter out weaker

regulatory links. Additionally, only the top 30 plausible regulatory links between

shared hub genes were retained to construct gene regulatory networks. Finally, the

shared regulatory network of hub genes was constructed by intersecting the regulatory

networks inferred from the EP and TG model datasets.

2.7 Machine Learning

To validate the diagnostic value of the transcriptomic signatures identified in animal

models, multiple machine-learning models were used to train diagnostic models based

on these signatures in EP and AD patient datasets (GSE48350 and E-MTAB-3123) to

predict disease status based on gene expression profiles. The main goal was to

distinguish patients with AD or EP from healthy controls. First, the gene expression

data was standardized by applying the StandardScaler function in the

sklearn.preprocessing module in Python. Standardization ensures that all

features/gene signatures have a mean of 0 and a standard deviation of 1, making them

comparable across scales. Several machine learning algorithms were used in the

classification process, including logistic regression, support vector machine (SVM),

random forest, naive Bayes, XGBoost, AdaBoost, and CatBoost. To evaluate the

performance of each model, a 5-fold stratified cross-validation method was used.

Different metrics were used to measure the performance of the models, including

accuracy, precision, recall, F1 score, and AUC. The ROC curve and AUC score for

each classifier were plotted to visually examine the trade-off between sensitivity and

specificity.

2.8 Cell Composition Deconvolution

In silico deconvolution can estimate the cellular composition of a tissue sample from

its gene expression profile, with CIBERSORT particularly suited for cellular

composition deconvolution of brain transcriptomes [39,40]. The raw count expression

matrix was converted to FPKM values using edgeR (4.4.1) [41], followed by cell

composition deconvolution using CIBERSORT (1.04) via the BrainDeconvShiny

platform, with cell-type signatures specific to *Mus musculus* from immunopurified

mouse brain tissue [40,42]. For statistical analysis, the Mann-Whitney test ($p < 0.05$)

was applied to compare the EP dataset with the control group, while the

Kruskal-Wallis test was used for the AD datasets, followed by pairwise comparisons across age and genotype groups. The BH method was used to control the FDR (q < 0.05). Associations between identified cell types and shared hub genes were determined by calculating Pearson correlation coefficients.

2.9 Single-cell and Single-nucleus RNA Sequencing Data Analysis

To localize and validate the expression of transcriptomic signatures identified in animal models at the cellular level in AD and EP patients, count matrices of sc/snRNA-seq (GSE201048 and GSE227223) were downloaded and processed according to the methods outlined in the original publications [43–45]. Quality control was performed to filter cells with high mitochondrial content (>20% for the EP dataset and >5% for the AD dataset) and select cells based on cellular complexity (300–5,000 genes per cell for the EP dataset and >500 genes for the AD dataset). Subsequent data processing was conducted using the standard Seurat pipeline, including scaling, normalization, variable gene identification, and clustering. PCA was performed on the 2,000 most variable genes, with the first 20 principal components (PCs) used for t-SNE and clustering in the EP dataset, and the first 30 PCs used for the AD dataset. Cluster identities were determined by identifying DEGs with the FindMarkers function, followed by annotation based on the original publications.

2.10 Computational modelling

To explore the contribution of the identified key pathways to pathological changes in AD and EP at the network level, we implemented a spiking neural network using the Izhikevich neuron model to simulate hippocampal CA3 circuit dynamics with simulation spans of 1000ms, employing the Brian2 package [46]. The network consists of 1000 Izhikevich neurons (800 excitatory and 200 inhibitory), with dynamics governed by [47]:

$$\frac{dv}{dt} = 0.04v^2 + 5v + 140 - u + I$$
$$\frac{du}{dt} = a(bv - u)$$

where v is the membrane potential, u is the recovery variable. When v≥30□mV, the following reset conditions are applied:$v = c$; $u = u + d$. Parameters a, b, c and d are set to standard regular spiking for excitatory neurons and fast-spiking for inhibitory neurons following the Izhikevich framework [47]. I is external input incorporating both suprathreshold Poisson noise at 1 Hz and presynaptic inputs [47].

Following Liu's EP network modeling, synaptic connectivity is sparse, with a 0.1 probability of random connections between neurons, and interactions are modeled using conductance-based synapses to allow for voltage-dependent temporal and spatial summation [47]. Synaptic interactions simulate AMPA (excitatory) and GABA-a (inhibitory) currents for four connection types: excitatory-excitatory (E→E), excitatory-inhibitory (E→I), inhibitory-excitatory (I→E), and inhibitory-inhibitory

363 (I→I) [47]. Synaptic currents are calculated as [48]:

$$I_j = wS_{i \to j}(t)(E_i - v_j)$$

we introduce w to leverage the synaptic weight, while $S_{i \to j}(t)$) is the synaptic

conductance evolving as $\frac{dS_{i \to j}(t)}{dt} = -\frac{S_{i \to j}(t)}{\tau(i)}$ and E_i is the synaptic reversal potential

[47]. Synaptic conductance is updated upon presynaptic spikes by adding a peak

conductance $S_{i \to j}(max)$, while we calibrate unitary synapse to induce EPSPs for

E→E/E→I and IPSPs for I→E/I→I of 1.0, 2.5, −2.0, −1.0 mV with time

constants:$\tau = \{3, 3, 5, 5\}$ ms and reversal potentials:$E = \{0, 0, -75, -75\}$ mV [49–51].

Each of four types of synaptic weights (w) is drawn independently from a standard

normal distribution, normalized to [0, 1], and set to 0 during synaptic pruning. We

simulated synaptic pruning by selectively removing excitatory (E→E and E→I) or

inhibitory (I→E and I→I) synapses, starting with those having the lowest weights

based on the synaptic weight distribution (ranging from 0% to 50%, with 5%

increments). Each pruning simulation was repeated at least 50 times.

The excitation/inhibition (E/I) ratio quantifies the balance between synaptic drives in

the network, indicating firing rate stability, while event synchrony measures temporal

coordination of firing patterns, implying network oscillation patterns. Both large E/I

ratio and excessive synchrony are associated with EP and AD [48].

To analyze the network, we calculated the E/I ratio and event synchrony to measure

the effects of synaptic pruning on firing rate stability and temporal coordination. The

E/I ratio compares total excitatory currents (from E→E and E→I synapses) to the

absolute magnitude of inhibitory currents (from I→E and I→I synapses), averaged

over the simulation period T [48]:

$$E/IRatio = \frac{\sum_{i=1}^{N} I_i^E(t)}{\left| \sum_{i=1}^{N} I_i^I(t) \right|}$$

where:

$I_i^E(t)$ represents excitatory synaptic currents onto neuron i at time t

$I_i^I(t)$ represents inhibitory synaptic currents onto neuron i at time t

N is the total number of neurons

$\bar{x}$ represents the mean over time T

Event synchrony (ES) quantifies spike timing coordination between excitatory neuron

pairs by calculating the fraction of their spikes occurring within 5-ms windows. For

each excitatory neuron pair (i, j), synchronous events are counted (with exact

coincident spikes weighted by 0.5 and near-coincident spikes within the time window

weighted by 1.0), then normalized by the geometric mean of both neurons' total spike

counts [52]. During total simulation time T:

$$ES_{ij} = \frac{\sum_{s_i \in S_i} \sum_{s_j \in S_j} C(s_i, s_j)}{\sqrt{N_i N_j}}$$

where:

Si is the set of all spike times from neuron i during the simulation.

Sj is the set of all spike times from neuron j during the simulation.

Ni and Nj are the total number of spikes from neuron i and j respectively over the

entire simulation period.

C(si, sj) counts synchronized events for each pair of spikes:

$$C(s_i, s_j) = \begin{cases} 1 & if\ 0 < |s_i - s_j| < 5ms \\ 0.5 & if\ s_i = s_j \\ 0 & otherwise \end{cases}$$

407 The network-wide event synchrony is computed as:

$$Network\ ES = \frac{1}{P} \sum_{i,j} ES_{ij}$$

Where $P = \frac{Ne\ (Ne-1)}{2}$ is the total number of unique excitatory neuron pairs, Ne is the

number of excitatory neurons.

Both E/I ratio and ES values are normalized to baseline unpruned values to highlight

pruning-induced changes in network dynamics. After normalization, the E/I ratio and

ES value fluctuate around 1. A value greater than 1 means a higher E/I ratio or ES

value than after synaptic pruning, and a value less than 1 means a smaller E/I ratio or

ES value after synaptic pruning. For statistical analysis, we conducted a correlation

analysis by calculating Pearson correlation coefficients and linear regression analysis

to quantify the effect sizes of excitatory and inhibitory pruning on E/I ratio and

synchronization.

3. Results

3.1 Identification and Functional Enrichment of Shared DEGs

3.1.1 Differential Expression Analysis

To reveal the shared molecular signatures underlying EP and AD models, we performed differential expression analysis using DESeq2. In the EP dataset, we identified 378 DEGs, with 351 upregulated and 27 downregulated genes (**Fig. 1A**). The TG dataset yielded 426 progressive DEGs, including 377 upregulated and 49 downregulated genes (**Fig. 1B**). In contrast, the J20 dataset contained only one upregulated progressive DEG, Cst7 (**Fig. 1C**). Notably, 101 genes were differentially expressed in both the EP and TG datasets, representing 26.72% and 23.71% of the DEGs in each dataset, respectively (**Fig. 1A–B**). This substantial overlap suggests shared molecular pathways between EP and TG models. Detailed DEGs information is available in **Table S2**.

3.1.2 Functional Enrichment Analysis of Shared DEGs

To investigate the biological relevance of these shared DEGs between EP and TG models, we conducted a functional enrichment analysis. KEGG pathway analysis identified several most enriched pathways, including "complement and coagulation

cascades," "phagosome," and "cytokine-cytokine receptor interaction" (**Fig. 1D**). GO

enrichment analysis highlighted key biological processes, such as "defense response,"

"immune response," and "regulation of cytokine production" (**Fig. 1E**). Additionally,

enriched molecular functions included "signaling receptor binding" and "immune

receptor activity," with cellular components such as the "extracellular matrix" and

"side of membrane" also significantly represented. These findings indicate that the

shared DEGs are primarily involved in immune processes, reflecting shared molecular

pathways in both EP and TG models. Detailed GO enrichment analysis results can be

found in **Table S2**.

3.2 Construction of Weighted Gene Co-Expression Network

WGCNA was particularly suited for this study due to its ability to identify modules of

highly correlated genes and reveal underlying biological pathways conserved across

different conditions. By applying WGCNA to both EP and AD datasets, we aimed to

identify conserved gene modules that may represent common molecular mechanisms

between EP and AD.

After filtering out outlier samples, low-expression, and low-variance genes in each

model dataset, we retained 10,997 genes common to all model datasets for further

analysis (**Fig. S1A-C**). For network construction, we selected a soft-thresholding

power of 9 to ensure that networks in all model datasets exhibited scale-free topology

with a fit index > 0.85 (**Fig. S1D**). This threshold was consistently applied across

analyses to maintain network comparability. To assess comparability between EP and

AD model networks, we examined correlations of ranked mean gene expression and

ranked gene connectivity between dataset pairs. For EP and J20 model datasets, high

correlations were observed for both expression ($r = 0.98$, $p < 1e-200$) and connectivity

($r = 0.23$, $p < 5.7e-132$). Similarly, EP and TG model datasets showed significant

correlations (expression: $r = 0.96$, $p = 1e-200$; connectivity: $r = 0.43$, $p = 1e-200$) (**Fig.**

S1E), indicating comparability across model datasets [31].

We used the WGCNA package to construct gene co-expression networks for the EP,

TG, and J20 model datasets. The analysis identified diverse gene modules across the

three models: the EP dataset revealed 33 modules (28–790 genes per module) (**Fig.**

2A), TG had 29 modules (34–1,537 genes) (**Fig. 2B**), and J20 contained 32 modules

(27–613 genes) (**Fig. 2C**). Detailed module information can be found in **Table S3**.

Associations between module eigengenes (MEs) and model genotypes were assessed.

The EP model dataset identified seven modules significantly associated with the EP

genotype: three with positive correlations (green, steelblue, turquoise; $0.73 \le r \le 0.79$,

$p < 3e-31$) and four with negative associations (lightyellow, red, lightcyan,

paleturquoise; $-0.92 \le r \le -0.69$, $p < 8e-27$). Notably, five of these modules contained

DEGs linked to the EP genotype (turquoise: $n = 198$, green: $n = 86$, steelblue: $n = 12$,

lightcyan: $n = 12$, red: $n = 5$) (**Fig. 2A, D**). In the TG model dataset, seven modules

showed significant associations with the TG genotype: two with positive correlations

(turquoise, yellow; $0.72 \leq r \leq 0.87$, $p < 3e\text{-}10$) and five with negative associations

(brown, saddlebrown, darkorange2, darkmagenta, royalblue; $-0.83 \leq r \leq -0.67$, $p <$

$1e\text{-}08$). Nine modules contained DEGs associated with the TG genotype (turquoise: n

= 268, yellow: n = 14, green: n = 7, darkorange2: n = 4, blue: n = 2, orangered4: n = 2,

royalblue: n = 2, darkmagenta: n = 1, tan: n = 1) (**Fig. 2B, E**). Consistent with the

DEGs analysis that showed the J20 model dataset had fewer DEGs, none of the

modules exhibited significant associations with the J20 genotype ($r < 0.4$) (**Fig. 2C,**

F).

3.3 Module Preservation Analysis

Following module construction, we assessed the preservation of EP model modules in

the AD model datasets to reveal conserve molecular pathways between EP and AD

models. Our analysis revealed varying degrees of module preservation between EP

and AD model datasets (**Fig. 3A-B**). In the TG model, the steelblue module had low

preservation scores (Z-summary.pres = 0.89. Similarly, in the J20 model, the steelblue

and yellow modules showed weak preservation (Z-summary.pres < 4), suggesting

these may represent unique biological processes specific to the EP model dataset. In

contrast, several EP model modules, including green, blue, and brown, exhibited high

preservation scores and significant overlap with AD modules in both models,

suggesting they represent conserved biological processes. Preservation statistics are

shown in **Table S3**.

Notably, the turquoise EP model module had the highest preservation scores

(Z-summary.pres = 83.01) and showed strong overlap with the TG model turquoise

module, displaying the highest overlap counts and the lowest adjusted p-values (**Fig.**

3A, C). Both turquoise modules were positively correlated with disease genotype (EP:

r = 0.79, p = 5e-41; TG: r = 0.87, p = 4e-18) and contained most DEGs in each model

(**Fig. 2A, C**). Other EP modules, such as lightcyan, showed significant correlations

but did not display preservation. GO enrichment analysis of this model turquoise

module revealed enrichment in immune-related terms (**Table S3**). We further assessed

the preservation of the mouse turquoise module in EP and AD patient dataset

(GSE63808 and GSE132903), finding that the turquoise immune modules from EP

and TG models significantly overlapped with modules constructed from EP and AD

patient datasets, respectively (**Fig. 3E-F, S2**). This suggests the turquoise module

represents a conserved immune molecular signature across species and disease states,

suggesting that immune processes play a preserved and significant role in these

modules across conditions.

3.4 Shared Hub Gene Identification and Regulatory Network Analysis

To identify hub genes that may play a crucial role in both EP and AD, we intersected

the hub genes of the turquoise module from EP and TG model datasets with the

previously identified 101 DEGs. This analysis identified 19 shared hub genes

common to both conditions (two shared hub genes not shown in the figure due to their

ranking: *Apobec* and *Cxcl16*) (**Fig. 4A-B**). We then constructed a PPI network to

investigate the biological significance of these 19 genes and the results showed a PPI

network consisting of 14 genes (**Fig. 4C, Table S4**). This PPI network revealed strong

connectivity among *C1q* complex component genes (*C1qa*, *C1qb*, *C1qc*) and other

immune-related genes, including *Trem2*, *Tyrobp*, and *Adgre1*. GO biological process

enrichment analysis highlighted the most significant processes involved in "Synaptic

pruning" (**Fig. 4D, Table S4**). Genes involved in "Synaptic pruning" (*C1qc*, *C1qb*,

C1qa, *Trem2*) also participate in other enriched processes, such as

"Complement-mediated synapse pruning" (*C1qa*, *Trem2*) and "Microglial cell

activation" (*Tyrobp*, *C1qa*, *Trem2*) (**Table S4**).

While WGCNA reveals gene co-occurrence patterns, it does not identify regulatory

mechanisms. To explain potential regulatory relationships among these 19 genes, we

applied the GENIE3 algorithm to gene expression profiles of the turquoise module in

both EP and TG conditions (**Table S5**). By intersecting predicted regulatory networks

from both conditions, we identified a shared regulatory pathway: *Tyrobp* regulates

Trem2, which in turn regulates the complement component genes *C1q* via *C1qa* (**Fig.**

4E). These genes, particularly *Tyrobp*, *Trem2*, and *C1q* complex components gene

(*C1qa*, *C1qb*, *C1qc*), play central roles in co-expression modules of both EP and TG

model and are involved in multiple enriched terms, especially synaptic pruning.

Notably, these genes are also progressive DEGs in the TG model (**Fig. 4F**). The

successful reconstruction of a regulatory network involving *Tyrobp*, *Trem2*, and *C1q*

component genes suggests they may represent a conserved molecular pathway

underlying both conditions.

3.5 Validation of the Diagnostic Value of Shared Hub Genes

To further validate the clinical relevance of the identified transcriptional signatures in

AD and EP animal models, we analyzed the expression of 19 shared hub genes in

datasets from AD and EP patients (GSE48350 and E-MTAB-3123) compared to

controls. Thirteen of these 19 genes were expressed in both datasets, with 10

exhibiting significantly higher expression levels ($p < 0.05$) compared to controls (**Fig.**

5A-B). Notably, genes such as *Tyrobp*, *Trem2*, and *C1q* complex component genes

demonstrated pronounced significance in both datasets (**Fig. 5A-B**). To assess their

diagnostic potential, we developed diagnostic models leveraging the expression

profiles of these genes in the hippocampus (**Fig. S2**). Using multiple machine learning

methods, these models effectively differentiated disease from control groups, with

particularly high performance in the EP dataset (AUC > 0.9) and satisfactory

performance in the AD patient dataset (AUC > 0.7) (**Fig. S3**). These findings confirm

the diagnostic value of the transcriptional signatures identified in animal models in

human AD and EP patients.

Given that these genes were shared hub genes of co-expression modules in EP and

AD models, likely reflecting conserved expression pathways and progressive changes,

we examined their co-expression patterns in AD and EP patient datasets (GSE48350

and E-MTAB-3123) by calculating Pearson correlation coefficients. *Tyrobp*, *Trem2*,

and the *C1q* component genes demonstrated strong correlations in both AD and EP

patient datasets ($r > 0.85$), whereas co-expression patterns among other genes were

less consistent across datasets (**Fig.5C-D**). Collectively, these findings suggested that

synaptic pruning pathway genes such as *Tyrobp*, *Trem2*, and the *C1q* component are

not only conserved between AD and EP animal models and human patients but also

likely play critical roles in the underlying pathology of these conditions.

3.6 Cellular Composition Alteration and Genes Localization in Cell Type

To investigate cellular composition changes and their associations with the key hub

genes (*Tyrobp*, *Trem2*, *C1q* complex genes) in TG and EP conditions, we employed

CIBERSORT for cell composition deconvolution of bulk RNA-seq data. Our analysis

revealed significant cellular composition alterations. Firstly, we observed a notable

reduction in neuronal composition in both EP and TG conditions compared to controls

($p < 0.05$) (**Fig. 6A, C**). Secondly, there was a marked increase in microglial

composition in both models ($p < 0.05$), consistent with the enrichment of

immune-related processes identified in previous analyses (**Fig. 6A, C**). While

astrocyte and endothelial cell alterations were also detected in EP and TG models (at 8

months, $p < 0.05$), these changes were less pronounced than the neuronal decrease and

microglial increase observed in the TG model (**Fig. 6A, C**). No significant differences

were found in oligodendrocyte composition in either condition.

We examined the relationship between these cellular composition changes and the key

hub genes such as *Tyrobp*, *Trem2*, and *C1q* complex genes by conducting Pearson

correlation analyses (**Fig. 6B, D**). Notably, all key hub genes showed strong negative

correlations ($r < -0.7$) with neuronal composition in both EP and TG conditions,

suggesting an inverse relationship between their expression levels and neuronal

proportion. Conversely, strong positive correlations were observed between these

genes and microglial and endothelial cell composition ($r > 0.7$). Correlations with

astrocytes and oligodendrocytes were more variable and showed less consistency

across genes and conditions. Although no cellular composition changes were observed

in the J20 model (**Fig. S4A**), we found that genes such as *Tyrobp*, *Trem2*, and *C1q*

complex components, were strongly associated with microglial composition,

indicating that these genes demonstrate a strong link with microglial cell types (**Fig.**

S4B). We further localized the expression of these genes to cell types using the whole

mouse brain transcriptome cell type atlas from the Allen Brain Atlas [53]. The results

showed that the majority (15 of 19 genes) were predominantly expressed in immune

cells, particularly microglia and macrophages (**Fig. S5**).

To further explore the expression of these genes in human patient cell types, especially immune cells, we explored the expression of these genes in two large-scale sc/sn RNAseq datasets of immune cells from AD and EP patients (GSE201048 and GSE2272230) (**Fig. S6**). We found that 12 of the shared hub genes were detectable in both datasets. Among these, *Tyrobp* and *Trem2*, along with the *C1q* complex component genes, displayed notably high and predominant expression in microglia compared to other shared hub genes (**Fig. S6**). Collectively, these findings confirmed the high and predominant expression of *Tyrobp*, *Trem2*, and *C1q* components genes in the microglia of both AD and EP patients. This consistency between animal models and human patients suggests that microglia and these synaptic pruning genes play crucial roles in the progression of both AD and EP pathology and may represent a potential conserved regulatory pathway.

3.7 Computational modeling of synaptic pruning on neural network

To better understand how synaptic pruning influences neural network dynamics and its potential role in the comorbidity of AD and EP, we developed a neural network model simulating the hippocampal CA3 circuit. In this model, we systematically varied the extent of synaptic pruning (0% to 50%) at either excitatory or inhibitory synapses. We then quantified changes in the E/I ratio and synchronization levels, conducting at least 50 simulation runs per condition (**Fig. 7**).

630 Our findings indicate that pruning of either excitatory or inhibitory synapses alters the

631 E/I ratio, with distinct effects. Inhibitory synaptic pruning significantly elevated the

632 E/I ratio, while excitatory pruning reduced it. These trends were consistent across all

633 simulations and in experiments where excitatory or inhibitory pruning was varied

634 independently (**Fig. 7A**). Correlation analysis showed a negative association between

635 excitatory pruning and the E/I ratio ($r = -0.474$) and a positive correlation between

636 inhibitory pruning and the E/I ratio ($r = 0.684$) (**Fig. 7B**). Linear regression further

637 confirmed that excitatory pruning reduced the E/I ratio ($\beta = -1.038$), whereas

638 inhibitory pruning increased it significantly ($\beta = 1.454$), with the effect of inhibitory

639 pruning surpassing that of excitatory pruning.

641 Interestingly, inhibitory pruning displayed a moderate positive correlation with event

642 synchronization ($r = 0.346$), while excitatory pruning showed only a weak negative

643 correlation ($r = -0.039$) (**Fig. 7B**). Linear regression analysis reflected this pattern,

644 with inhibitory pruning exerting a stronger effect on synchronization ($\beta = 0.358$) than

645 excitatory pruning ($\beta = -0.041$). Additionally, the E/I ratio itself was positively

646 correlated with synchronization ($r = 0.491$) (**Fig. 7B**), suggesting that the influence of

647 synaptic pruning on network synchronization may be nonlinear and indirectly

648 mediated by changes in the E/I balance.

These simulations suggest that excessive synapses pruning alters the E/I ratio, and

inhibitory pruning overpowering the effects of excitatory pruning and substantially

enhancing synchronization.

4. Discussion

While Alzheimer's disease (AD) and epilepsy (EP) share a complex bidirectional

relationship, the molecular pathways linking AD-related neurodegeneration to

increased EP susceptibility remain poorly understood. To address this issue, we used

rigorous bioinformatic and computational methods to integrate and analyze

transcriptional data from multiple large-scale publicly available animal models and

human patient datasets. We identified consensus molecular pathways, cellular

processes, and neural network dynamics associated with this comorbidity across

multiple datasets.

Conserved Neuroinflammation Pathways in Epilepsy and Alzheimer's Disease

Our findings provide strong evidence supporting the hypothesis that

neuroinflammatory pathways are common between AD and EP, particularly those

directly triggered by tau pathology [7,18]. Consistent with previous results, we identified

no significant transcriptional changes in the J20 model due to the inherent

characteristics of the model rather than technical limitations (**Fig. 1C**), such as slower

and later accumulation of pathology and variable neuronal cell loss [29]. However,

differential expression analysis demonstrated significant gene overlap between EP and

TG models (**Fig. 1A-B**). WGCNA revealed preservation of the immune-related

turquoise module across both disease models as well as human patient datasets (**Fig.

2-3**). These results align with previous analyses of this dataset that identified

significant conservation of this module across multiple AD patient cohorts [29].

Although highly conserved in the EP and TG datasets, the J20 dataset did not show

strong preservation of this module (**Fig. 3B**), which further reinforces that this module

is the result of a transcriptional state change that is conserved in EP and TG. We

extended these findings by demonstrating the module's conservation and overlap in

EP models and patient datasets, suggesting it represents a fundamental molecular

pathway shared between AD and EP (**Fig. 3,5**).

Functional enrichment analysis of shared DEGs and the preserved module revealed

significant enrichment in neuroinflammatory processes, including inflammatory

response, cytokine production, blood vessel morphogenesis, glial cell activation, and

complement and coagulation cascades (**Fig. 1D-E, Table S3**). Additionally, we

observed increased populations of neuroinflammation-associated cell types, including

astrocytes, microglia, and endothelial cells, in both TG and EP animal models (**Fig.

6**).

Neuroinflammation has been extensively documented in both conditions and appears

to serve a complex dual role in disease progression [54,55]. While activated microglia

facilitate Aβ plaque clearance and reactive astrocytes provide metabolic support and

ionic homeostasis that may attenuate neural damage and seizures, chronic activation

or dysfunction of these glial populations can promote disease progression through

persistent neuroinflammation [56–59]. Specifically, activated microglia and astrocytes

secrete proinflammatory cytokines and chemokines, including IL-1β, TNF-α, and

IL-6, which can enhance neuronal hyperexcitability, increase seizure susceptibility,

exacerbate Aβ and tau aggregation, and accelerate neurodegeneration and cognitive

decline [56–59].

Additional inflammatory components, including complement cascade activation and

blood-brain barrier (BBB) dysfunction, are implicated in both pathologies [60–63].

Complement activation contributes to synaptic loss in AD and is elevated in TLE

patient tissue [62,63]. BBB disruption precedes neuronal hyperactivity in EP, promoting

astrocyte dysfunction and seizures, while in AD it facilitates peripheral inflammatory

cell infiltration, amplifying neuroinflammation [60,61].

These results establish neuroinflammation as a key mechanistic link between AD and

EP [18]. AD-associated inflammation can enhance neuronal hyperexcitability and

seizure susceptibility. Conversely, EP-induced inflammation may accelerate Aβ and

tau accumulation, contributing to cognitive decline. Our results support previous

studies and suggest that neuroinflammation is highly conserved between these

diseases and between species. Future studies should focus on revealing the precise

causal relationship between these neuroinflammatory components and AD-EP

comorbidity.

Synaptic Pruning was Identified as a Core Pathway in this Comorbidity

A comprehensive analysis of this conserved module revealed a central synaptic

pruning pathway involving *Tyrobp*, *Trem2*, and *C1q* complement genes (**Fig. 4**).

These genes demonstrated consistent upregulation across both AD and EP patient

datasets, with preserved regulatory relationships in both animal models and human

patients (**Fig. 5**). Moreover, we confirmed their specific expression in microglial cells

across both animal and human patient samples (**Fig. S5-6**). Collectively, these

findings suggest that microglia-mediated synaptic pruning represents a core

mechanistic pathway underlying the comorbidity of AD and EP.

This pathway has been extensively documented in AD pathology, with its role

extending to both tau and Aβ pathology [64]. Dysregulation or mutations in *Tyrobp* and

Trem2 are established risk factors for AD due to their effects on microglial function

[65,66]. Under physiological conditions, microglia maintain synaptic homeostasis

through a regulated process in which complement proteins, including *C1q*, tag

unnecessary synapses for elimination [67]. Microglial receptors, particularly *Trem2* and

Tyrobp, recognize these tags and initiate phagocytic clearance, a process critical for

neural circuit refinement and maintenance in the adult brain [68]. In AD pathology, these

receptors enable microglia to recognize protein deposits, triggering their transition to

a disease-associated microglial (DAM) state characterized by enhanced phagocytic

activity and elevated expression of *Tyrobp*, *Trem2*, *Apoe*, and complement proteins

including *C1q* [64]. While *Tyrobp* or *Trem2* deficiency disrupts DAM activation and

impairs critical microglial functions such as plaque clearance, potentially accelerating

AD pathology, this pathway overactivation can also contribute to neurodegeneration

and cognitive decline through excessive *C1q*-mediated synaptic pruning [64,69–71].

Synaptic pruning can exert effects on both AD and EP disease progression through

two primary mechanisms: synaptic transmission and synaptic plasticity. Regarding

synaptic transmission, elevated *C1q* expression or Aβ accumulation in the CA1 region

leads to decreased miniature and evoked excitatory postsynaptic currents [72,73]. In AD,

this reduction in synaptic transmission contributes to cognitive decline by disrupting

normal neural communication. For EP, this alteration affects both excitatory and

inhibitory neurons [74], potentially contributing to the imbalance between excitation and

inhibition that characterizes epileptic activity. Notably, these effects can be reversed

through *C1q* downregulation [75,76].

The pathway's impact on synaptic plasticity is equally significant. Research

demonstrates that C1q inhibition in aged experimental models can restore both

long-term potentiation (LTP) and long-term depression (LTD) [69,71,77]. In non-aged

models, increased *C1q* expression also impairs the LTP process [74]. This is particularly

relevant for AD, where impaired synaptic plasticity is a feature of cognitive decline.

In EP, compromised synaptic plasticity may contribute to the formation and

maintenance of epileptogenic circuits. The relationship between these mechanisms

appears bidirectional: excessive synaptic pruning leads to neuronal damage and

reduced LTP, which in turn promotes disease progression in both conditions, creating

a harmful feedback loop where synaptic dysfunction promotes pathological processes.

Notably, interventions targeting this synaptic pruning pathway, whether through

knockdown of regulatory genes such as *Tyrobp* and *Trem2* or direct inhibition of *C1q*

accumulation, have successfully restored electrophysiological function and cognitive

function in AD models, even in the presence of significant amyloid and tau pathology

[64,69–71] . Our results establish a regulatory network that is consistent with previous

observations and further extends its role in the comorbidity of AD and EP (**Fig. 4**).

While this pathway is extensively studied in the context of AD, to the best of our

knowledge, no prior research has established a connection between this pathway and

the comorbidity of EP and AD. Our understanding of the relationship between

synaptic pruning and EP remains nascent, especially how to interpret its contribution

to the network dynamics [78]. To address this gap, we conducted an in-silico experiment

to shed light on this issue.

Synaptic Pruning Alter E/I Balance to Promote Network Synchronization

Computational modeling has become an important tool for understanding AD and EP

pathology, models ranging from molecular-level simulations of ion dynamics to

large-scale network models [79,80]. Disruption of excitatory/inhibitory (E/I) balance and

neural synchronization represent a key point of convergence between AD and EP

pathology, supported by a wealth of experimental evidence. In EP, patch-clamp

recordings demonstrate both reduced GABAergic inhibition and enhanced

glutamatergic transmission, resulting in network hyperexcitability [81,82]. In AD, studies

have shown that AD mouse models exhibit spontaneous epileptiform activity, while

human studies using magnetoencephalography have demonstrated disruption of

synchronization in multiple frequency bands [16,83].

To explore how synaptic pruning contributes to the comorbidity of AD and EP, we

developed a computational model of the hippocampal CA3 circuit on top of Liu's

biologically plausible EP modeling. We calibrate fixed synaptic strength by adjusting

conductance-based synapses to produce realistic postsynaptic potential based on

animal experiments. [16,83], while leveraging synaptic connection pruning across

simulation trials on independently standard normal synaptic weights to transition the

network into abnormal states. This model allowed us to quantify how pruning of

excitatory and inhibitory synapses affects E/I balance (E/I ratio) and the temporal

coordination of neural firing (synchronization) [82].

Our computational model suggests that synaptic pruning can regulate the E/I balance

in neural networks, with inhibitory synapse pruning having a more pronounced effect.

Quantitative analysis by linear regression also suggests that excitatory synapse

elimination decreased the E/I ratio (β = -1.038), while inhibitory synapse removal led

to a larger increase in the E/I ratio (β = 1.454) (**Fig. 7**). Our finding that inhibitory

pruning has a stronger (enhancing) effect on network dynamics can be explained by

the function of inhibitory interneurons within neural circuits. These neurons often

establish synaptic connections with multiple excitatory neurons and act as regulators

of circuit coordination and synchronization [84]. Therefore, when inhibitory control

mechanisms are impaired, the resulting disruption of regulation will trigger

dysregulation that extends to more widespread circuits. Moreover, endogenous

homeostatic mechanisms such as synaptic scaling appear to have a limited capacity to

compensate for the loss of inhibitory synapses [85]. Ultimately, a reduction in inhibitory

synapses may produce a larger perturbation in neural networks than an equivalent

reduction in excitatory synapses.

Interestingly, our findings reveal a weaker correlation of synaptic pruning on network

synchronization compared to its impact on the E/I ratio. Specifically, inhibitory

pruning exhibited a moderate positive correlation with synchronization (r = 0.346),

whereas excitatory pruning had a negligible correlation (r = −0.039). Notably, the E/I

ratio itself demonstrated a strong correlation with synchronization levels (r = 0.491).

While it seems intuitive that enhancing network excitability can enhance synchrony,

the results of our analysis predict the importance of considering the specific location

and functional role of pruned synapses in neural circuits [84]. For instance, in a circuit

that has an inhibitory effect on the broader network, pruning inhibitory synapses that

inhibit this circuit will paradoxically reduce network synchronization. This explains

the indirect, nonlinear association of synaptic pruning with network synchronization

(**Fig. 7B**).

.

Although it is still difficult to establish a causal relationship between synaptic pruning

and this comorbidity, our simulation results suggest a potential relationship, that is,

the loss of neurons or synapses in neurodegenerative diseases disrupt the E/I balance

of the neural network, may thereby promote epilepsy susceptibility (**Fig. 7**). In fact,

epilepsy is well known as a comorbidity of multiple neurodegenerative diseases such

as Huntington's and Parkinson's disease [86–88]. Our simulations show that in the case of

equal loss of excitatory or inhibitory synapses (even in the case of loss more

excitatory synapses than inhibitory synapses) (**Fig. 7A**), the effect of inhibitory

synapses loss can overpower the effect of excitatory synapses loss, causing a more

extensive network E/I balance disruption and promoting network synchronization

(**Fig. 7B**). Similar to *Tyrobp* and *Trem2* being known risk factors for AD, Mendelian

randomization studies have also indicated that *C1qa* is a risk factor for both AD and

EP (**Fig. S7**) [89]. In fact, in AD, synaptic loss often precedes cognitive decline and EP,

which suggests that it may be an initiating event rather than a consequence [90].

Our results highlight the role of inhibitory synaptic pruning in promoting network

synchronization. Notably, several recent studies have shown that microglial synaptic

pruning appear to have inhibitory pruning preference [91]. Dejanovic B et al. had shown

that the *Trem2-C1q* microglial synaptic pruning pathway engulfs inhibitory synapses

than excitatory synapses in AD model, while Fan J et al. demonstrated similar

preferential pruning of inhibitory synapses during recurrent seizures that exacerbate

EP model pathology [91,92]. At the molecular level, *C1q* acts as an "eat me" signal,

tagging synapses for microglia to eliminate [78]. While this tagging process is not

synapse-type specific, microglia demonstrate selective pruning of inhibitory synapses

[78,91]. Studies have shown that *C1q* blockade reduces both the frequency and severity

of seizures in traumatic brain injury models [93]. However, it is also important to note

that *C1q* knockout also affects synaptic elimination during development, resulting in

enhanced connectivity and seizure activity [94,95]. These findings suggest the

importance of managing microglial *C1q*-mediated synaptic pruning during AD

pathology and aging, and further exploration and regulation of this potential

preferential pruning of inhibitory synapses is warranted,

Our study has two limitations: while our analysis mined a conserved microglial

synaptic pruning pathway across multiple datasets, we primarily relied on bulk

RNAseq data that cannot fully capture cell subtype specificity. Although we attempted

to address this limitation through computational deconvolution and validation of

single-cell datasets, future studies should employ spatial transcriptomic approaches to

advance understanding of the cell subtype-specific contribution to AD-EP comorbidity,

for example, by tracking changes in microglial activation patterns and synaptic action

in AD and EP. In addition, while our computational model provides insights into the

contribution of synaptic pruning to network dynamics, it simplifies many aspects of

real neural circuits, such as topology, cell types, and dendritic details within neural

circuits. Future models could incorporate more detailed cellular mechanisms to better

reflect the progression of AD and EP comorbidity at the molecular level.

Conclusion

In this study, we integrated large-scale transcriptomic datasets from multiple AD and

EP animal models, as well as human patient cohorts, to identify a shared molecular

pathway that converges on microglial synaptic pruning. Specifically, we found a

highly   preserved   immune-related   co-expression   module   centered   on   the

Tyrobp-Trem2-C1q complement pathway. Differential gene expression and gene

regulatory network analysis confirmed that this pathway is consistently upregulated in

both animal models and human patients. Our computational modeling further suggests

a mechanistic basis for how excessive synapses pruning tip the E/I balance toward

hyperexcitation, leading to the comorbidity of AD and EP.

These findings highlight a microglial-driven, complement-mediated synaptic pruning

mechanism common to AD and EP. In linking the dysregulated immune response to

E/I imbalances, our work provides a unifying framework that may explain the strong

bidirectional comorbidity observed clinically. More broadly, these results highlight

the potential of targeting microglia–complement signaling for therapeutic

interventions aimed at ameliorating seizures and slowing cognitive decline in patients

with AD and EP. Future studies integrating spatial transcriptomics, advanced

single-cell profiling, and in vivo manipulation of complement pathways are warranted

to elucidate how microglial subtypes selectively prune synapses in AD and EP, and

whether modulating this process can indeed confer disease-modifying benefits.

Abbreviations

EP   Epilepsy

AD   Alzheimer's disease

E/I   Excitation/Inhibition

TLE   Temporal lobe epilepsy

902	Aβ	Amyloid-β
903	GEO	Gene Expression Omnibus
904	ENA	European Nucleotide Archive
905	WT	Wild-type
906	scRNA-Seq	Single-cell RNA sequencing
907	snRNA-Seq	Single-nucleus RNA sequencing
908	PCA	Principal Component Analysis
909	DEGs	Differentially expressed genes
910	BH	Benjamini-Hochberg
911	KEGG	Kyoto Encyclopedia of Genes and Genomes
912	GO	Gene ontology
913	MAD	Median absolute deviation
914	TOM	Topological overlap matrices
915	MEs	Module eigengenes
916	MM	Module membership
917	GS	Gene-genotype significance
918	PPI	Protein-protein interaction
919	FDR	False discovery rate
920	SVM	Support vector machine
921	PCs	Principal components
922	ES	Event synchrony

923	BBB	Blood-brain barrier
924	DAM	Disease-associated microglial
925	LTP	Long-term potentiation
926	LTD	Long-term depression
927	BP	Biological processes
928	CC	Cellular components
929	MF	Molecular functions
930	HR	Hazard ratios
931	CI	Confidence intervals
932	NCBI	National Center for Biotechnology Information

Figures and legends

Figure 1: Differential Expression and Functional Enrichment Analyses in EP and AD Model Datasets.

(A) Volcano plot shows the DEGs in the EP model dataset. The x-axis represents the log2 fold change, and the y-axis shows the -log10 adjusted p-value. The dashed lines mark the thresholds for significance (adjusted p-value < 0.05 and |log2 fold change| > 1). **(B)** Differential expression analysis of the TG model dataset. Scatter plot showing progressive DEGs identified across different ages (4, 6, and 8 months) compared to their control and initial months. The shaded regions correspond to the different age groups, with the numbers indicating the count of EP-DEGs (numerator) and total

DEGs (denominator) with a log2 fold change below 1 that marks the threshold for

non-significant genes. **(C)** The plot shows the expression level of single upregulated

progressive DEG *Cst7* in the J20 model over time (x-axis: Age in months; y-axis:

Normalized counts), with different genotypes (J20 and WT) represented by different

colors. Dashed lines represent mean paths for each age. (LRTstatistic = 43.70,

log2foldchange[6-12months] = 2.73, FDR = 2.91E-5). **(D)** KEGG pathway

enrichment analysis of shared DEGs between the EP and TG datasets. The bubble plot

indicates the significantly enriched pathways, with bubble size representing the gene

count and color indicating the significance level (-log10 adjusted p-value). **(E)** GO

enrichment analysis of shared DEGs between the EP and TG model datasets. The dot

plot shows the enriched biological processes (BP), cellular components (CC), and

molecular functions (MF). Dot size corresponds to gene count, while color represents

the significance level (-log10 adjusted p-value). Immune-related functions were

prominently enriched.

Figure 2: Co-Expression Network Construction and Model-Trait Association.

(A-C) Hierarchical clustering dendrograms are presented for the EP (**A**), TG (**B**), and

J20 (**C**) model datasets, with gene modules identified through dynamic tree cutting.

The colors in the heatmap below each dendrogram correspond to distinct gene

modules, where the clustering of similar colors between the EP and other model

datasets (TG and J20) reflects the conservation of modules across conditions. The GS

is shown with darker colors indicating stronger correlations: red represents positive

correlations, and blue represents negative correlations with disease genotypes (EP for

panel A, TG for panel B, and J20 for panel C). Overlapping red areas in the

module-trait relationships suggest that these modules may be positively correlated

with both EP and AD model, as seen in the turquoise module in the TG model dataset.

The DEG row shows the distribution of DEGs within the modules: red indicates

DEGs, while grey indicates non-DEGs. **(D-F)** Heatmaps of the module-trait

relationships in EP (**D**), TG (**E**), and J20 (**F**) model datasets. The heatmaps show the

Pearson correlation coefficients between MEs and genotypes. The color scale

represents the strength and direction of the correlation, with red indicating positive

and blue indicating negative correlations. Modules are considered as significant

association with genotype using threshold (p-value < 0.05, $|r| > 0.65$).

Figure 3: Module Preservation Analysis Across EP and AD Datasets.

(A, B) Zsummary preservation statistics of EP model modules in the TG and J20

model datasets, respectively. Each point represents an EP model module, with the

y-axis displaying the Zsummary preservation statistic. A Zsummary > 10 indicates

strong evidence of preservation, $2 <$ Zsummary < 10 indicates moderate evidence, and

Zsummary < 2 suggests no evidence of preservation. The gray horizontal line at Z =

10 and the black horizontal line at Z = 2 serve as reference points in the plots. The

module size is indicated by the x-axis, and the color represents the module color. **(C-F)**

Bubble plots show the cross-tabulation of EP model modules with TG (**C**) and J20 (**D**)

model modules; the cross-tabulation of EP patient (GSE63808) and EP model datasets

(**E**); cross-tabulation of AD (GSE132903) and TG mouse model datasets (**F**). The

color scale represents the adjusted p-value for the overlap significance, while the size

of the dots corresponds to the number of overlapping genes. Larger, more significant

dots indicate stronger overlaps between EP and AD modules such as the turquoise

module. Only significantly overlapping with overlap count > 30 and adjusted p-value

< 0.05 was colored, while those with insignificant overlap are marked in gray.

Figure 4: Identification and Network Analysis of Shared Hub Genes in EP and

AD Datasets.

(**A**) Network plot illustrating the top 50 genes within the conserved turquoise module

across both the EP and TG model datasets. Each node represents a gene, and edges

signify gene co-expression connections (Grey for EP, blue for TG). The color of the

nodes indicates gene connectivity. 17 red nodes are shared hub genes that exhibit high

connectivity in both networks and are identified as DEGs under both conditions. Two

shared hub genes are not shown in the figure due to their ranking in both modules:

Apobec and *Cxcl16*. (**B**) Venn diagram depicting the overlap between hub genes

identified in the EP and AD datasets and shared DEGs. 19 shared hub genes were

identified as key drivers in both conditions. (**C**) PPI network of the 14 shared hub

genes constructed from 19 shared hub genes using the STRING database. The size

and color of the nodes are colored by their connectivity in the PPI network. (**D**) The

bubble plot showed the top GO biological process enrichment terms based on

p-values for the 19 shared hub genes. (**E**) The shared gene regulatory network was

inferred using the GENIE3 algorithm, showing the regulatory relationships among the

shared hub genes. (**F**) Progressive upregulation of key 5 shared hub genes involved in

microglial synaptic pruning across different ages in the TG mouse model. The plot

shows the expression level of these genes over time (x-axis: Age in months; y-axis:

Normalized counts), with different genotypes represented by different colors (WT:

black, TG: teal). Dashed lines represent mean paths for each age. LRT statistics of

these genes can be found in **Table S2**.

1017

Figure 5. Expression analysis and correlation patterns of shared hub genes in AD

and EP patient datasets.

(**A, B**) Violin plots showing the differential expression of shared hub genes between

control and disease groups in (**A**) EP patient dataset (E-MTAB-3123) and (**B**) AD

patient dataset (GSE48350). Red violins represent control groups and blue violins

represent disease groups. The width of each violin indicates the density of data points,

and the internal box plots show the median and interquartile ranges. The Wilcoxon

p-value is shown for each comparison. (**C, D**) Correlation heatmap showing the

relationships between shared hub genes and age in the EP patient dataset

(E-MTAB-3123) and AD patient dataset (GSE48350). The color intensity and box

size represent the strength of Pearson correlation coefficients (r), with red indicating

positive correlations and blue indicating negative correlations.

1030

Figure 6: Cellular Composition Deconvolution and Association with Shared Hub

Genes in EP and TG Datasets.

(**A, C**) The plot shows the estimated proportions of different cell types (astrocytes,

neurons, oligodendrocytes, microglia, and endothelial) in control and EP model

conditions (**A**), and TG model condition (**C**). The x-axis represents different

conditions or months, and the y-axis shows the proportions of specific cell types in

each condition, with different genotypes represented by different colors. The

Mann-Whitney test was performed to determine significant cell composition

difference ($p < 0.05$) in EP model dataset. The Kruskal-Wallis test was performed to

determine the significance ($q < 0.05$) with the BH method was used to adjust the

p-value in pairwise comparisons among different groups in the TG model dataset. (**B,**

D) Correlation heatmap showing the relationships between key shared hub genes and

cell composition in the EP model dataset. The color intensity represents the strength

of Pearson correlation coefficients (r), with red indicating positive correlations and

blue indicating negative correlations. The significance of correlations with shared hub

gene is shown by red arrows, with arrow color proportional to correlation strength.

1047

Figure 7. Effects of excitatory and inhibitory synaptic pruning on neural

network dynamics.

(**A**) Four-dimensional plot showing the simulation results of excitatory pruning ratio

(0-0.5), inhibitory pruning ratio (0-0.5), E/I ratio, and synchronization. The surface

visualizes the variation of the average E/I ratio with different pruning condition.

Colors indicate average synchronization levels from dark purple (close to 1, indicating

no change in synchrony) to yellow (indicating higher synchrony). (**B**) Correlation plot

showing the relationships among synapse pruning, E/I ratio and synchronization. The

color intensity represents the strength of Pearson correlation coefficients (r), with red

indicating positive correlations and blue indicating negative correlations.

1058

1059 **Legends of supplementary files**

1060

Table S1: Summary of Quality Control and Mapping Statistics for Bulk

RNA-seq Samples.

Detailed summary of the QC and mapping statistics for all RNA-seq samples

analyzed in this study. For each sample, the total number of raw reads, the number of

reads passing quality filtering through Fastp, and the reads retained after rRNA

contamination removal by ribodetector are reported. Each sample is categorized by

the experimental group, including relevant age and condition tags, and further

classified into either the AD or EP dataset. Notes regarding any specific issues or

annotations on individual samples are also provided.

1070

Table S2: Differential Expression and Functional Enrichment Analysis of Shared DEGs.

Summary of differential expression and functional enrichment analyses across three datasets: EP, TG, and J20 mouse models.

1075

Table S3: Module Construction, Preservation, and Functional Enrichment of Gene Co-Expression Networks in Mouse Models.

Summary of WGCNA on EP, TG, and J20 mouse models. The table is divided into multiple sections, including the gene modules' membership identified in each dataset and their correlations with model-specific genotypes. Additionally, module preservation analyses between the EP and AD datasets are included. GO enrichment analyses of the turquoise modules are also shown.

1083

Table S4: Protein-Protein Interaction Network and GO Biological Process Enrichment Analysis of Shared Hub Genes.

Analysis results of the PPI network constructed for 19 shared hub genes identified across EP and TG models. The table is divided into three sections. The PPI GO sheet summarizes the results of the GO biological process enrichment analysis. The PPI Degree sheet lists each gene and its degree of connectivity within the network. The PPI Interaction sheet details the specific interactions between these genes.

1091

Table S5: Regulatory Network Prediction Using GENIE3 for Turquoise Module in Mice Models.

Results of regulatory network inference were conducted using the GENIE3 machine learning algorithm on the gene expression profiles of the turquoise module from the EP and TG models. The table is divided into two sections: GENIE3TG and GENIE3EP, representing the predicted regulatory relationships within the TG and EP datasets, respectively. Each section lists the regulatory gene, the target gene, and the corresponding weight of the predicted regulatory interaction.

1100

Table S6: Cell Composition Deconvolution of Bulk RNA-seq Data.

Results of cell composition deconvolution analysis conducted using CIBERSORT from BrainDeconvShiny on bulk RNA-seq data from the EP and AD models. The table includes the estimated proportions of major brain cell types, including neurons, astrocytes, oligodendrocytes, microglia, and endothelial for each sample, alongside the sample identifiers, group classification, age, and condition type (e.g., rTg4510, Wild Type).

1108

Figure S1. Construction of co-expression module in model datasets.

(**A-C**) Hierarchical clustering dendrograms and trait heatmaps for the EP, rTg4510, and J20 datasets. (**A**) EP dataset, (**B**) rTg4510 dataset, and (**C**) J20 dataset.

Hierarchical clustering was performed on variance-stabilized gene expression data to identify potential outliers. The top 85% of genes with the highest Median Absolute Deviation (MAD) were retained, and outliers were detected and removed before network construction. Two samples were removed from the rTg4510 dataset, and three samples were removed from the J20 dataset. The heatmaps below each dendrogram display the trait data associated with each sample, with red indicating outliers or specific traits of interest. **(D-E)** Soft-thresholding power Selection and Comparability Assessment, **(D)** Selection of the soft-thresholding power for network construction. The left panel shows the scale-free topology fit index (y-axis) as a function of the soft-thresholding power (x-axis) for EP, TG, and J20 datasets. The right panel depicts the mean connectivity (y-axis) against soft-thresholding power, indicating that a power of 9 was chosen for all datasets to achieve scale-free topology with $R^2 > 0.85$. **(E)** Comparability assessment of the EP, TG, and J20 datasets. Scatter plots display the correlations between ranked mean gene expression (top) and ranked gene connectivity (bottom) between EP and TG datasets (left) and between EP and J20 datasets (right). Positive correlations and significant P values suggest strong comparability between datasets.

Figure S2. Construction of co-expression module in patient datasets.

(**A**) Scale independence and mean connectivity plots across soft-thresholding powers for the EP (GSE63808) and AD (GSE132903) patient datasets, with a selected

1133   soft-thresholding power of 9 ensuring a scale-free topology ($R^2 \geq 0.85$). (**B-C**)

1134   Dendrogram of gene modules identified in the EP and AD dataset using hierarchical

1135   clustering and dynamic tree cutting based on topological overlap matrices (TOM).

1136

Figure S3: ROC curve of Diagnostic model.

(**A**) in EP patient dataset (E-MTAB-3123) and (**B**) in AD patient dataset (GSE48350).

ROC curves of various machine learning classifiers in classifying disease cases from

control participants based on selected gene expression data. Remarkably, most of the

models have an AUC between 0.85 and 0.92 in EP patient dataset, with the best

performance recorded with XGBoost, reaching the highest AUC score in this EP

dataset (AUC = 0.92). Although the AD model diagnostic efficiency is not as high as

that of the EP dataset, which may reflect the complex disease background of AD,

these diagnostic models still show high diagnostic efficiency (AUC > 0.7).

1146

Figure S4: Cellular Composition Deconvolution and Association with Shared

Hub Genes in J20 Datasets.

(**A**) The plot shows the cell composition level over time in J20 dataset (x-axis: Age in

months; y-axis: Normalized counts), with different genotypes represented by different

colors (WT: black, J20: teal). Dashed lines represent mean paths for each age. The

Kruskal-Wallis test was performed to determine the significance with the BH method

was used to adjust the p-value in pairwise comparisons among different groups, with a

significance level of q < 0.05. (**B**) Correlation heatmap showing the relationships between key shared hub genes and cell composition in the J20 model dataset. The color intensity represents the strength of Pearson correlation coefficients (r), with red indicating positive correlations and blue indicating negative correlations.

Figure S5: Expression Localization of Shared Hub Genes in Mouse Brain Cell Types.

This figure shows the expression patterns of 19 shared hub genes across various cell types in the mouse brain, based on the Whole Mouse Brain Transcriptomic Cell Type Atlas (https://assets.nemoarchive.org/dat-qg7n1b0) from the Allen Brain Cell Atlas. The top left panel illustrates the localization and color of various cell types, such as astrocytes, microglia, macrophages, endothelial cells, and oligodendrocytes in the brain atlas. Each circular panel highlights the specific expression profile of one gene, with darker shading representing higher expression levels. Most of these genes (15 out of 19) are predominantly expressed in immune cells, specifically microglia and macrophages.

Figure S6: sc/nRNAseq Analysis of Shared Hub Gene Expression in Immune Cells of Patients.

(**A, D**) Dot plot displaying the expression of shared hub genes across different identified cell types in the EP patient dataset. The size of the dots represents the

proportion of cells expressing each gene, and the color intensity indicates the average

expression level. (**A**) EP patient dataset (GSE227223), (**D**) AD patient dataset

(GSE201048). (**B, E**) t-SNE plot showing the clustering of cells in the EP and AD

dataset based on their gene expression profiles, with distinct colors representing

different cell types. (**B**) EP dataset, (**E**) AD dataset. (**C, F**) Feature plots showing the

expression patterns of individual hub genes within the EP dataset, over on the t-SNE

map. Darker purple areas represent higher expression levels. *TYROBP*, *TREM2*,

C1QA, *C1QB*, and *C1QC* exhibit high expression predominantly in microglia clusters.

(C) EP dataset, (F) AD dataset.

1184

Figure S7: Forest plot of Mendelian randomization result from DMRdb.

Forest plot showing hazard ratios (HR) and 95% confidence intervals (CI) for

associations between Alzheimer's disease subtypes, epilepsy, and *C1QA*

http://www.inbirg.com/DMRdb/. Red dots represent point estimates for hazard ratios,

and horizontal lines indicate 95% confidence intervals. p-values are shown for each

association. Values greater than 1.0 indicate increased risk for the outcome.

1191

1192 **Declarations**

1193

1194 **Ethics approval and consent to participate**

1195 Not applicable.

1196

Consent for publication

Not applicable.

1199

Availability of data and materials

The datasets analyzed during the current study are available in the National Center for

Biotechnology Information (NCBI) or European Nucleotide Archive (ENA) under the

following accession numbers: PRJEB18790, GSE125957, GSE48350, E-MTAB-3123,

GSE63808, GSE132903, GSE201048, and GSE227223.

1205

Competing Interests

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

1209

Funding

This research was supported by the Young Scientists Fund of the National Natural

Science Foundation of China (Grant Nos. 82201629) and the Guangdong Provincial

Science and Technology Special Fund (Grant Nos. 210803136872196)

1214

Authors' contribution

HL conceptualized the study, conducted the investigation, developed the methodology,

performed formal analysis, and contributed to writing the original draft and reviewing/editing the manuscript. ZX, YT, and BL contributed to the investigation, performed formal analysis, and participated in writing the original draft and reviewing/editing the manuscript. JL contributed to writing the original draft and reviewing/editing the manuscript. JW was involved in conceptualization and contributed to writing the original draft and reviewing/editing the manuscript. MC, XL, YS, SC, ZD, and MH contributed to the investigation and participated in reviewing/editing the manuscript. BH and NW conceptualized the study and contributed to writing the original draft and reviewing/editing the manuscript. XJ provided supervision, acquired funding, contributed to conceptualization, and participated in writing the original draft and reviewing/editing the manuscript. All authors read and approved the final manuscript.

Acknowledgments

Not applicable

References

1. Giorgi FS, Saccaro LF, Busceti CL, Biagioni F, Fornai F. Epilepsy and Alzheimer's Disease: Potential mechanisms for an association. Brain Research Bulletin. 2020 Jul 1;160:107-20.
2. Kamondi A, Grigg-Damberger M, Löscher W, Tanila H, Horvath AA.

1240		Epilepsy and epileptiform activity in late-onset Alzheimer disease:
1241		Clinical and pathophysiological advances, gaps and conundrums. Nature
1242		Reviews Neurology. 2024 Mar;20(3):162-82.
1243	3.	Fang Y, Si X, Wang J, Wang Z, Chen Y, Liu Y, Yan Y, Tian J, Zhang B,
1244		Pu J. Alzheimer disease and epilepsy: A Mendelian randomization study.
1245		Neurology. 2023 Jul 25;101(4):e399-409.
1246	4.	Wimo A, Jönsson L, Bond J, Prince M, Winblad B, International AD.
1247		The worldwide economic impact of dementia 2010. Alzheimer's &
1248		dementia. 2013 Jan 1;9(1):1-1.
1249	5.	Ngugi AK, Bottomley C, Kleinschmidt I, Sander JW, Newton CR.
1250		Estimation of the burden of active and life‐time epilepsy: a
1251		meta‐analytic approach. Epilepsia. 2010 May;51(5):883-90.
1252	6.	Mendez MF, Lim GT. Seizures in elderly patients with dementia:
1253		epidemiology and management. Drugs & aging. 2003 Sep;20:791-803.
1254	7.	Leitner D, Pires G, Kavanagh T, Kanshin E, Askenazi M, Ueberheide B,
1255		Devinsky O, Wisniewski T, Drummond E. Similar brain proteomic
1256		signatures in Alzheimer's disease and epilepsy. Acta Neuropathologica.
1257		2024 Jun;147(1):27.
1258	8.	Pandis D, Scarmeas N. Seizures in alzheimer disease: clinical and
1259		epidemiological data: Seizures in alzheimer disease. Epilepsy currents.
1260		2012 Sep;12(5):184-7.
1261	9.	Tombini M, Assenza G, Ricci L, Lanzone J, Boscarino M, Vico C,
1262		Magliozzi A, Di Lazzaro V. Temporal lobe epilepsy and Alzheimer's
1263		disease: from preclinical to clinical evidence of a strong association.
1264		Journal of Alzheimer's disease reports. 2021 Jan 1;5(1):243-61.
1265	10.	Johnson, E. L., Krauss, G. L., Lee, A. K., Schneider, A. L., Dearborn, J.
1266		L., Kucharska-Newton, A. M., ... & Gottesman, R. F. (2018).
1267		Association between midlife risk factors and late-onset epilepsy: results
1268		from the atherosclerosis risk in communities study. *JAMA*
1269		*neurology*, 75(11), 1375-1382.
1270	11.	Stefanidou M, Beiser AS, Himali JJ, Peng TJ, Devinsky O, Seshadri S,
1271		Friedman D. Bi-directional association between epilepsy and dementia:
1272		The Framingham Heart Study. Neurology. 2020 Dec 15;95(24):e3241-7.
1273	12.	Gourmaud S, Shou H, Irwin DJ, Sansalone K, Jacobs LM, Lucas TH,
1274		Marsh ED, Davis KA, Jensen FE, Talos DM. Alzheimer-like amyloid
1275		and tau alterations associated with cognitive deficit in temporal lobe
1276		epilepsy. Brain. 2020 Jan 1;143(1):191-209.
1277	13.	Tai XY, Koepp M, Duncan JS, Fox N, Thompson P, Baxendale S, Liu JY,
1278		Reeves C, Michalak Z, Thom M. Hyperphosphorylated tau in patients
1279		with refractory epilepsy correlates with cognitive decline: a study of
1280		temporal lobe resections. Brain. 2016 Sep 1;139(9):2441-55.
1281	14.	Zou Y, Wang C, Li H, Zhong M, Lin J, Hu Y, Chen Z, Gan CL. Epileptic

seizures induced by pentylenetetrazole kindling accelerate Alzheimer-like neuropathology in 5× FAD mice. Frontiers in Pharmacology. 2024 Oct 31;15:1500105.

15. Watts ME, Pocock R, Claudianos C. Brain energy and oxygen metabolism: emerging role in normal function and disease. Frontiers in molecular neuroscience. 2018 Jun 22;11:216.

16. Born HA. Seizures in Alzheimer's disease. Neuroscience. 2015 Feb 12;286:251-63.

17. Toniolo S, Sen A, Husain M. Modulation of brain hyperexcitability: potential new therapeutic approaches in Alzheimer's disease. International journal of molecular sciences. 2020 Dec 7;21(23):9318.

18. Liew Y, Retinasamy T, Arulsamy A, Ali I, Jones NC, O'Brien TJ, Shaikh MF. Neuroinflammation: A common pathway in Alzheimer's disease and epilepsy. Journal of Alzheimer's Disease. 2023 Jan 1;94(s1):S253-65.

19. Kalyvas AC, Dimitriou M, Ioannidis P, Grigoriadis N, Afrantou T. Alzheimer's Disease and Epilepsy: Exploring Shared Pathways and Promising Biomarkers for Future Treatments. Journal of Clinical Medicine. 2024 Jan;13(13):3879.

20. Targa Dias Anastacio H, Matosin N, Ooi L. Neuronal hyperexcitability in Alzheimer's disease: what are the drivers behind this aberrant phenotype?. Translational psychiatry. 2022 Jun 22;12(1):257.

21. Tang Y, Yan Y, Mao J, Ni J, Qing H. The hippocampus associated GABAergic neural network impairment in early-stage of Alzheimer's disease. Ageing Research Reviews. 2023 Apr 1;86:101865.

22. Bosco F, Guarnieri L, Rania V, Palma E, Citraro R, Corasaniti MT, Leo A, De Sarro G. Antiseizure medications in Alzheimer's disease from preclinical to clinical evidence. International journal of molecular sciences. 2023 Aug 10;24(16):12639.

23. Andrews S, Krueger F, Segonds-Pichon A, Biggins L, Kruger C, Wingett S. FastQC. Babraham Bioinformatics. Cambridge, UK. 2010.

24. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 2018 Sep 1;34(17):i884-90.

25. Deng ZL, Münch PC, Mreches R, McHardy AC. Rapid and accurate identification of ribosomal RNA sequences via deep learning. Nucleic acids research. 2022 Jun 10;50(10):e60-.

26. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nature biotechnology. 2019 Aug;37(8):907-15.

27. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics. 2014 Apr 1;30(7):923-30.

28. Love MI, Huber W, Anders S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. Genome biology. 2014 Dec;15:1-21.

29. Castanho I, Murray TK, Hannon E, Jeffries A, Walker E, Laing E, Baulf H, Harvey J, Bradshaw L, Randall A, Moore K. Transcriptional signatures of tau and amyloid neuropathology. Cell Reports. 2020 Feb 11;30(6):2040-54.

30. Elizarraras JM, Liao Y, Shi Z, Zhu Q, Pico AR, Zhang B. WebGestalt 2024: faster gene set analysis and new support for metabolomics and multi-omics. Nucleic Acids Research. 2024 May 29:gkae456.

31. Miller JA, Horvath S, Geschwind DH. Divergence of human and mouse brain transcriptome highlights Alzheimer disease pathways. Proceedings of the National Academy of Sciences. 2010 Jul 13;107(28):12698-703.

32. Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis. BMC bioinformatics. 2008 Dec;9:1-3.

33. Zhang B, Horvath S. A general framework for weighted gene co-expression network analysis. Statistical applications in genetics and molecular biology. 2005 Aug 12;4(1).

34. Ritchie ME, Phipson B, Wu DI, Hu Y, Law CW, Shi W, Smyth GK. limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic acids research. 2015 Apr 20;43(7):e47-.

35. Schober P, Boer C, Schwarte LA. Correlation coefficients: appropriate use and interpretation. Anesthesia & analgesia. 2018 May 1;126(5):1763-8.

36. Langfelder P, Luo R, Oldham MC, Horvath S. Is my network module preserved and reproducible?. PLoS computational biology. 2011 Jan 20;7(1):e1001057.

37. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, Gable AL, Fang T, Doncheva NT, Pyysalo S, Bork P. The STRING database in 2023: protein–protein association networks and functional enrichment analyses for any sequenced genome of interest. Nucleic acids research. 2023 Jan 6;51(D1):D638-46.

38. Huynh-Thu VA, Irrthum A, Wehenkel L, Geurts P. Inferring regulatory networks from expression data using tree-based methods. PloS one. 2010 Sep 28;5(9):e12776.

39. Newman AM, Liu CL, Green MR, Gentles AJ, Feng W, Xu Y, Hoang CD, Diehn M, Alizadeh AA. Robust enumeration of cell subsets from tissue expression profiles. Nature methods. 2015 May;12(5):453-7.

40. Sutton GJ, Poppe D, Simmons RK, Walsh K, Nawaz U, Lister R, Gagnon-Bartsch JA, Voineagu I. Comprehensive evaluation of deconvolution methods for human brain gene expression. Nature Communications. 2022 Mar 15;13(1):1358.

41. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor

package for differential expression analysis of digital gene expression data. bioinformatics. 2010 Jan 1;26(1):139-40.

42. Zhang Y, Chen K, Sloan SA, Bennett ML, Scholze AR, O'Keeffe S, Phatnani HP, Guarnieri P, Caneda C, Ruderisch N, Deng S. An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. Journal of neuroscience. 2014 Sep 3;34(36):11929-47.

43. Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, Srivastava A, Molla G, Madad S, Fernandez-Granda C, Satija R. Dictionary learning for integrative, multimodal and scalable single-cell analysis. Nature biotechnology. 2024 Feb;42(2):293-304.

44. Kumar P, Lim A, Hazirah SN, Chua CJ, Ngoh A, Poh SL, Yeo TH, Lim J, Ling S, Sutamam NB, Petretto E. Single-cell transcriptomics and surface epitope detection in human brain epileptic lesions identifies pro-inflammatory signaling. Nature Neuroscience. 2022 Jul;25(7):956-66.

45. Sun N, Victor MB, Park YP, Xiong X, Scannail AN, Leary N, Prosper S, Viswanathan S, Luna X, Boix CA, James BT. Human microglial state dynamics in Alzheimer's disease progression. Cell. 2023 Sep 28;186(20):4386-403.

46. Stimberg M, Brette R, Goodman DF. Brian 2, an intuitive and efficient neural simulator. elife. 2019 Aug 20;8:e47314.

47. Izhikevich EM. Simple model of spiking neurons. IEEE Transactions on neural networks. 2003 Nov;14(6):1569-72.

48. Tzavellas NP, Tsamis KI, Katsenos AP, Davri AS, Simos YV, Nikas IP, Bellos S, Lekkas P, Kanellos FS, Konitsiotis S, Labrakakis C. Firing Alterations of Neurons in Alzheimer's Disease: Are They Merely a Consequence of Pathogenesis or a Pivotal Component of Disease Progression?. Cells. 2024 Feb 29;13(5):434.

49. Sammons RP, Vezir M, Moreno-Velasquez L, Cano G, Orlando M, Sievers M, Grasso E, Metodieva VD, Kempter R, Schmidt H, Schmitz D. Structure and function of the hippocampal CA3 module. Proceedings of the National Academy of Sciences. 2024 Feb 6;121(6):e2312281120.

50. Vyleta NP, Borges-Merjane C, Jonas P. Plasticity-dependent, full detonation at hippocampal mossy fiber–CA3 pyramidal neuron synapses. Elife. 2016 Oct 25;5:e17977.

51. Liu Y, Grigorovsky V, Bardakjian B. Excitation and inhibition balance underlying epileptiform activity. IEEE Transactions on Biomedical Engineering. 2020 Jan 1;67(9):2473-81.

52. Penn Y, Segal M, Moses E. Network synchronization in hippocampal neurons. Proceedings of the National Academy of Sciences. 2016 Mar 22;113(12):3341-6.

53. Yao Z, van Velthoven CT, Kunst M, Zhang M, McMillen D, Lee C, Jung
W, Goldy J, Abdelhak A, Aitken M, Baker K. A high-resolution
transcriptomic and spatial atlas of cell types in the whole mouse brain.
Nature. 2023 Dec 14;624(7991):317-32.
54. Vezzani A, French J, Bartfai T, Baram TZ. The role of inflammation in
epilepsy. Nature reviews neurology. 2011 Jan;7(1):31-40.
55. Heneka MT, Golenbock DT, Latz E. Innate immunity in Alzheimer's
disease. Nature immunology. 2015 Mar;16(3):229-36.
56. Pekny M, Pekna M. Astrocyte reactivity and reactive astrogliosis: costs
and benefits. Physiological reviews. 2014 Oct;94(4):1077-98.
57. Chen ZP, Wang S, Zhao X, Fang W, Wang Z, Ye H, Wang MJ, Ke L,
Huang T, Lv P, Jiang X. Lipid-accumulated reactive astrocytes promote
disease progression in epilepsy. Nature neuroscience. 2023
Apr;26(4):542-54.
58. Onyango IG, Jauregui GV, Čarná M, Bennett Jr JP, Stokin GB.
Neuroinflammation in Alzheimer's disease. Biomedicines. 2021 May
7;9(5):524.
59. Vezzani A, Viviani B. Neuromodulatory properties of inflammatory
cytokines and their impact on neuronal excitability. Neuropharmacology.
2015 Sep 1;96:70-82.
60. Van Vliet EA, Aronica E, Gorter JA. Blood–brain barrier dysfunction,
seizures and epilepsy. InSeminars in cell & Developmental biology 2015
Feb 1 (Vol. 38, pp. 26-34). Academic Press.
61. Zlokovic BV. Neurovascular pathways to neurodegeneration in
Alzheimer's disease and other disorders. Nature Reviews Neuroscience.
2011 Dec;12(12):723-38.
62. Kopczynska M, Zelek WM, Vespa S, Touchard S, Wardle M, Loveless S,
Thomas RH, Hamandi K, Morgan BP. Complement system biomarkers
in epilepsy. Seizure. 2018 Aug 1;60:1-7.
63. Veerhuis R, Nielsen HM, Tenner AJ. Complement in the brain.
Molecular immunology. 2011 Aug 1;48(14):1592-603.
64. Haure-Mirande JV, Audrain M, Ehrlich ME, Gandy S. Microglial
TYROBP/DAP12 in Alzheimer's disease: Transduction of physiological
and pathological signals across TREM2. Molecular neurodegeneration.
2022 Aug 24;17(1):55.
65. Pottier C, Ravenscroft TA, Brown PH, Finch NA, Baker M, Parsons M,
Asmann YW, Ren Y, Christopher E, Levitch D, van Blitterswijk M.
TYROBP genetic variants in early-onset Alzheimer's disease.
Neurobiology of aging. 2016 Dec 1;48:222-e9.
66. Neumann H, Daly MJ. Variant TREM2 as risk factor for Alzheimer's
disease. N Engl J Med. 2013 Jan 10;368(2):182-4.
67. Stevens B, Allen NJ, Vazquez LE, Howell GR, Christopherson KS,

Nouri N, Micheva KD, Mehalow AK, Huberman AD, Stafford B, Sher A. The classical complement cascade mediates CNS synapse elimination. Cell. 2007 Dec 14;131(6):1164-78.

68. Paolicelli RC, Ferretti MT. Function and dysfunction of microglia during brain development: consequences for synapses and neural circuits. Frontiers in Synaptic Neuroscience. 2017 May 10;9:9.

69. Audrain M, Haure-Mirande JV, Wang M, Kim SH, Fanutza T, Chakrabarty P, Fraser P, St George-Hyslop PH, Golde TE, Blitzer RD, Schadt EE. Integrative approach to sporadic Alzheimer's disease: deficiency of TYROBP in a tauopathy mouse model reduces C1q and normalizes clinical phenotype while increasing spread and state of phosphorylation of tau. Molecular psychiatry. 2019 Sep;24(9):1383-97.

70. Haure-Mirande JV, Audrain M, Fanutza T, Kim SH, Klein WL, Glabe C, Readhead B, Dudley JT, Blitzer RD, Wang M, Zhang B. Deficiency of TYROBP, an adapter protein for TREM2 and CR3 receptors, is neuroprotective in a mouse model of early Alzheimer's pathology. Acta neuropathologica. 2017 Nov;134:769-88.

71. Haure-Mirande JV, Wang M, Audrain M, Fanutza T, Kim SH, Heja S, Readhead B, Dudley JT, Blitzer RD, Schadt EE, Zhang B. Integrative approach to sporadic Alzheimer's disease: deficiency of TYROBP in cerebral Aβ amyloidosis mouse normalizes clinical phenotype and complement subnetwork molecular pathology without reducing Aβ burden. Molecular psychiatry. 2019 Mar;24(3):431-46.

72. Neniskyte U, Gross CT. Errant gardeners: glial-cell-dependent synaptic pruning and neurodevelopmental disorders. Nature Reviews Neuroscience. 2017 Nov;18(11):658-70.

73. Hu Y, Qi H, Yang J, Wang F, Peng X, Chen X, Zhu X. Wogonin mitigates microglia-mediated synaptic over-pruning and cognitive impairment following epilepsy. Phytomedicine. 2024 Dec 1;135:156222.

74. Wu X, Gao Y, Shi C, Tong J, Ma D, Shen J, Yang J, Ji M. Complement C1q drives microglia-dependent synaptic loss and cognitive impairments in a mouse model of lipopolysaccharide-induced neuroinflammation. Neuropharmacology. 2023 Oct 1;237:109646.

75. Bie B, Wu J, Foss JF, Naguib M. Activation of mGluR1 mediates C1q-dependent microglial phagocytosis of glutamatergic synapses in Alzheimer's rodent models. Molecular neurobiology. 2019 Aug 15;56:5568-85.

76. Wu J, Bie B, Foss JF, Naguib M. Amyloid fibril–induced astrocytic glutamate transporter disruption contributes to complement C1q-mediated microglial pruning of glutamatergic synapses. Molecular Neurobiology. 2020 May;57:2290-300.

77. Iweka CA, Seigneur E, Hernandez AL, Paredes SH, Cabrera M, Blacher

E, Pasternak CT, Longo FM, de Lecea L, Andreasson KI. Myeloid deficiency of the intrinsic clock protein BMAL1 accelerates cognitive aging by disrupting microglial synaptic pruning. Journal of Neuroinflammation. 2023 Feb 24;20(1):48.

78. Andoh M, Ikegaya Y, Koyama R. Synaptic pruning by microglia in epilepsy. Journal of clinical medicine. 2019 Dec 9;8(12):2170.

79. Yuan P, Zhang M, Tong L, Morse TM, McDougal RA, Ding H, Chan D, Cai Y, Grutzendler J. PLD3 affects axonal spheroids and network defects in Alzheimer's disease. Nature. 2022 Dec 8;612(7939):328-37.

80. Stefanescu RA, Shivakeshavan RG, Talathi SS. Computational models of epilepsy. Seizure. 2012 Dec 1;21(10):748-59.

81. Fritschy JM. Epilepsy, E/I balance and GABAA receptor plasticity. Frontiers in molecular neuroscience. 2008 Mar 28;1:201.

82. Ren X, Brodovskaya A, Hudson JL, Kapur J. Connectivity and neuronal synchrony during seizures. Journal of Neuroscience. 2021 Sep 8;41(36):7623-35.

83. van Nifterick AM, de Haan W, Stam CJ, Hillebrand A, Scheltens P, van Kesteren RE, Gouw AA. Functional network disruption in cognitively unimpaired autosomal dominant Alzheimer's disease: a magnetoencephalography study. Brain Communications. 2024;6(6):fcae423.

84. Luo L. Architectures of neuronal circuits. Science. 2021 Sep 3;373(6559):eabg7285.

85. Dzyubenko E, Fleischer M, Manrique-Castano D, Borbor M, Kleinschnitz C, Faissner A, Hermann DM. Inhibitory control in neuronal networks relies on the extracellular matrix integrity. Cellular and Molecular Life Sciences. 2021 Jul;78:5647-63.

86. Neri S, Mastroianni G, Gardella E, Aguglia U, Rubboli G. Epilepsy in neurodegenerative diseases. Epileptic Disorders. 2022 Apr;24(2):249-73.

87. Cano A, Fonseca E, Ettcheto M, Sánchez-López E, de Rojas I, Alonso-Lana S, Morato X, Souto EB, Toledo M, Boada M, Marquie M. Epilepsy in neurodegenerative diseases: related drugs and molecular pathways. Pharmaceuticals. 2021 Oct 18;14(10):1057.

88. Paudel YN, Angelopoulou E, Piperi C, Othman I, Shaikh MF. Revisiting the impact of neurodegenerative proteins in epilepsy: focus on alpha-synuclein, beta-amyloid, and tau. Biology. 2020 Jun 12;9(6):122.

89. Zheng X, Tian Z, Che X, Zhang X, Xiang Y, Ge Z, Zhai Z, Ma Q, Pan J. DMRdb: a disease-centric Mendelian randomization database for systematically assessing causal relationships of diseases with genes, proteins, CpG sites, metabolites and other diseases. Nucleic Acids Research. 2025 Jan 6;53(D1):D1363-71.

90. Chin J, Scharfman HE. Shared cognitive and behavioral impairments in

epilepsy and Alzheimer's disease and potential underlying mechanisms.
Epilepsy & Behavior. 2013 Mar 1;26(3):343-51.

91. Dejanovic B, Wu T, Tsai MC, Graykowski D, Gandham VD, Rose CM,
Bakalarski CE, Ngu H, Wang Y, Pandey S, Rezzonico MG. Complement
C1q-dependent excitatory and inhibitory synapse elimination by
astrocytes and microglia in Alzheimer's disease mouse models. Nature
Aging. 2022 Sep;2(9):837-50.

92. Fan J, Dong X, Tang Y, Wang X, Lin D, Gong L, Chen C, Jiang J, Shen
W, Xu A, Zhang X. Preferential pruning of inhibitory synapses by
microglia contributes to alteration of the balance between excitatory and
inhibitory synapses in the hippocampus in temporal lobe epilepsy. CNS
Neuroscience & Therapeutics. 2023 Oct;29(10):2884-900.

93. Holden SS, Grandi FC, Aboubakr O, Higashikubo B, Cho FS, Chang
AH, Forero AO, Morningstar AR, Mathur V, Kuhn LJ, Suri P.
Complement factor C1q mediates sleep spindle loss and epileptic spikes
after mild brain injury. Science. 2021 Sep 10;373(6560):eabj2685.

94. Chu Y, Jin X, Parada I, Pesic A, Stevens B, Barres B, Prince DA.
Enhanced synaptic connectivity and epilepsy in C1q knockout mice.
Proceedings of the National Academy of Sciences. 2010 Apr
27;107(17):7975-80.

95. Ma Y, Ramachandran A, Ford N, Parada I, Prince DA. Remodeling of
dendrites and spines in the C1q knockout model of genetic epilepsy.
Epilepsia. 2013 Jul;54(7):1232-9.

1557

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

B

Figure 7