Epistasis and cryptic QTL identified using modified bulk segregant analysis of copper resistance in budding yeast

Cassandra Buzby[1,2], Yevgeniy Plavskin[1,2], Federica M.O. Sartori[1,2,3], Qiange Tong[1,2],

Janessa K. Vail[1,2], Mark L. Siegal[1,2]*

[1] Department of Biology, New York University, New York, NY, USA

[2] Center for Genomics and Systems Biology, New York University, New York, NY, USA

[3] Current affiliation: Department of Oncological Sciences, Mount Sinai, New York, NY, USA

* Corresponding author

E-mail: mark.siegal@nyu.edu (MLS)

Abstract

The contributions of genetic interactions to natural trait variation are challenging to estimate experimentally, as current approaches for detecting epistasis are often underpowered. Powerful mapping approaches such as bulk segregant analysis, wherein individuals with extreme phenotypes are pooled for genotyping, obscure epistasis by averaging over genotype combinations. To accurately characterize and quantify epistasis underlying natural trait variation, we have engineered strains of the budding yeast *Saccharomyces cerevisiae* to enable crosses where one parent's chromosome is fixed while the rest of the chromosomes segregate. These crosses allow us to use bulk segregant analysis to identify quantitative trait loci (QTL) whose effects depend on alleles on the fixed parental chromosome, indicating a genetic interaction with that chromosome. Our method, which we term epic-QTL (for *epi*static-with-*c*hromosome *QTL*) analysis, can thus identify interaction loci with high statistical power. Here we perform epic-QTL analysis of copper resistance with chromosome I or VIII fixed in a cross between divergent naturally derived strains. We find seven loci that interact significantly with chromosome VIII and none that interact with chromosome I, the smallest of the 16 budding yeast chromosomes. Each of the seven interactions alters the magnitude, rather than the direction, of an additive QTL effect. We also show that fixation of one source of variation — in this case chromosome VIII, which contains the large-effect QTL mapping to *CUP1* — increases power to detect the contributions of other loci to trait differences.

Author Summary

Most traits of interest that vary in populations are determined by multiple genetic factors, as well as by environmental variation and random chance. These influences may combine in complicated ways, for example when the effect of one genetic variant depends on genetic variants elsewhere in the genome. Such dependencies are difficult to identify and characterize because testing many possible combinations of influences reduces statistical power. We address this challenge by combining bulk segregant analysis, a robust method for detecting effects of individual genetic variants averaged across genetic backgrounds, with chromosome fixation in budding yeast. This approach allows us to detect gene-variant effects that depend on the fixed chromosome with statistical power comparable to that achieved when detecting background-independent effects. Applying the approach to copper sulfate resistance, we identify and characterize interaction effects and, by removing one source of variation (a single yeast chromosome), improve detection of background-independent effects as well.

Introduction

Predicting phenotype from genotype is a central goal of genetics research, yet even with great advances in sequencing technology and experimental scale, accurate characterization of how genetic differences translate into trait differences remains elusive. Many genetic influences are subtle, and most traits of interest that vary in populations are determined by multiple genetic factors, as well as by environmental variation and random chance [1–4]. Moreover, these influences may combine in

complicated ways, for example in the form of gene-by-gene (epistatic) or gene-by-

environment interactions [5–7].

Epistasis is particularly challenging to detect in standard quantitative-genetic

analyses because the number of possible interaction effects vastly exceeds the number

of single-locus (additive) effects, leading to very low statistical power to detect the

interactions [5]. Standard analyses fit additive effects before interaction effects, which

can lead to the mistaken inference that epistasis is less salient than it actually is [8,9]. If

interacting alleles are rare in a mapping population, under-detection of epistasis is even

more severe. Failure to detect epistasis in turn compromises accurate prediction of an

individual's phenotype from genotype, even when an additive model appears to fit well

[8–10].

The distribution of types of epistatic effects influences both detection of loci with

effects on traits and evolutionary trajectories [11–13]. Magnitude epistasis changes the

size of an allele's effect without changing its direction (beneficial vs. detrimental),

whereas sign epistasis changes the direction. Sign epistasis gives rise to peaks and

valleys in fitness landscapes and can alter the order of mutations that increase fitness

[11]. Models of distribution and patterns of these effects have been postulated, with two

extremes being global epistasis and idiosyncratic epistasis. Global epistasis indicates

underlying predictability of the effects of individual alleles, perhaps based on a latent

phenotype such as protein stability or physiological health [14,15]. Idiosyncratic

epistasis, by contrast, refers to unpredictable allele effects that ultimately depend on

details of particular alleles [16,17]. Data on the relative prevalences of magnitude and

sign epistasis, which are currently lacking, would inform which model of epistasis

patterns is more suitable, which in turn will inform models of the evolution of complex

traits.

An ideal method for identifying and characterizing epistatic contributions to trait

variation would have equal (and high) statistical power to detect additive and interaction

effects. Experimental designs with extremely large sample sizes are now common [18–

22], but the most powerful of these favor (or exclusively detect) additive effects. One

such experimental design is bulk segregant analysis (BSA). In a typical BSA

experiment, many progeny are produced from a cross between divergent parental

genotypes, progeny with extreme values of a trait of interest are pooled, and high-

coverage genome sequencing is used to detect loci at which the allele frequency in the

selected bulk deviates significantly from the allele frequency in an unselected (or

oppositely selected) bulk [18,23]. Although BSA benefits from sampling many

individuals — over 10 million in some cases [18] — it cannot detect genetic interaction

effects because pooled sequencing results in an estimate of the effect of any one locus

that is the average across all genotypes at other loci.

Genotyping and phenotyping individuals rather than pools avoids the averaging

issue, but suffers from the problem that potential pairwise interactions far outnumber

additive effects of single loci, resulting in greatly reduced statistical power. Two recent

attempts to detect epistasis using very large sample sizes illustrate this challenge. One

study, which genotyped ~200,000 diploid progeny of a yeast cross, identified numerous

interaction effects, but found a strong positive correlation between the additive effect of

a locus and the number of pairwise interactions it participates in [22]. This correlation

could reflect the true distribution of epistatic effects or an influence of additive-effect size

on detectability of interactions [22]. Another study genotyped ~100,000 haploid

segregants but to maximize power tested only interactions between pairs of additive-

effect QTL [21]. This study found evidence of epistatic effects, but found the additive-

plus-pairwise model to have only modest predictive power for the phenotypes of

reconstructed genotypes, suggesting that effects remain undetected [21]. Another

recent study did not genotype large numbers of segregants, but instead engineered

naturally occurring gene variants into four divergent yeast strains to detect genetic

background-dependent effects on growth. This study found a substantial fraction of

variants to have background-dependent effects, but was limited to variants in and

around 103 genes previously shown to be interaction hubs in genome-wide double-

mutant screens [24,25]. The results of these studies suggest that unbiased approaches

are needed that equalize the power to detect additive and epistatic effects.

126   To identify and characterize genetic interactions with comparable statistical

power to that of additive effects, we combine chromosome substitution with BSA in the

genetically tractable budding yeast, *Saccharomyces cerevisiae*. We call this method

epic-QTL (for *epi*static-with-*c*hromosome *QTL*) analysis. To perform epic-QTL, we cross

two divergent natural isolates of *S. cerevisiae*, but before generating segregating F2

progeny from the resulting F1 diploid, we first eliminate one chromosome from one

parent, fixing the remaining homologous chromosome. The result is that all surviving F2

haploids contain identical copies of one chromosome, but have unique combinations of

the two parents' alleles on all other chromosomes. We also perform the reciprocal

cross, in which the other parent's chromosome is fixed. We then grow the haploid

progeny in selective or non-selective conditions. Here, we use the presence or absence

of copper (II) sulfate ($CuSO_4$) as a selective condition, with the presence of copper and

subsequent oxidative stress selecting for individuals with alleles that promote survival in

this environment. Standard BSA would compare the allele frequencies in bulks grown in

the presence or absence of copper to identify additive QTL for the trait of copper

resistance. In epic-QTL, there are four bulks rather than two. By comparing the QTL

identified in the crosses with the reciprocal fixed chromosomes, we can identify loci

whose effects differ between genetic backgrounds. The loci that are dependent on

parental-chromosome background for their effect are thus epistatic with that fixed

chromosome and are detected with statistical power usually only afforded to additive

effects. In practice, this comparison is done by including a parental-chromosome (or

genetic-background) term in a logistic-regression model that is fit to the data from all

four bulks simultaneously. An additional benefit of chromosome substitution is an

increase in power for detection of additive effects by removing a source of variance in a

population. With only one cross, the fixation of a chromosome containing a major QTL

allows the remaining segregating sites to exert greater influence over membership in

the selected bulk.

Using epic-QTL we can determine where QTL that interact with each fixed

chromosome lie relative to where additive QTL lie, and how prevalent interactions are

for a given trait, in this case copper sulfate resistance. We focus on copper sulfate for

three reasons: 1) its extensive use in prior *S. cerevisiae* studies, allowing comparison

with previous results; 2) the potential for unexplained heritability, in that copper

resistance shows a large gap in broad and narrow sense heritability compared to other

traits [4]; and 3) its ecological and physiological relevance [26–28]. In addition, one

major interaction hub identified by Matsui *et al.* was *CUP1* on chromosome VIII, which is

a major driver of copper resistance, and thus a perfect candidate for chromosome

fixation [22]. We fixed chromosomes I or VIII in a cross between strains derived from an

oak tree or from a wine barrel, which differ on average at 1 in 200 base pairs [29]. The

variants in these strains reflect natural genetic diversity, offering insights into genetic

interactions that might occur within populations. We demonstrate that this method can

be used to remove sources of trait variation to reveal additive QTL otherwise

undetected at this experimental scale, as well as to identify interaction QTL with high

statistical power. We find that these interactions often occur within significant additive

QTL peaks, providing evidence of a lack of pervasive sign epistasis.

Results

Implementation of epic-QTL for copper resistance

Our method to identify QTL interacting with a fixed chromosome, or epic-QTL,

combines chromosome substitution with BSA in a cross of two parental yeast strains.

For the parental strains we used divergent natural isolates of *S. cerevisiae*: NCYC3631

and NCYC3591 (hereafter "Oak" and "Wine", respectively). Before crossing the two

strains, we engineered one strain of each parent to contain the inducible *GAL1*

promoter (*pGAL1*) adjacent to the centromere of one chromosome. Inducing a high

level of transcription through the centromere destabilizes the kinetochore, resulting in

missegregation of the chromosome during mitosis so that a strain with only the

homologous chromosome remaining can be isolated [30]. By this method, we fixed

chromosome I in each background (Fig 1A) and, separately, chromosome VIII in each

background (Fig 1B). We then mated the Wine *pGAL1* strain with the Oak non-*pGAL1*

strain (Fig 1C), and vice versa, to produce diploids with reciprocal conditional

chromosomes. Each diploid strain was then plated on galactose to induce *pGAL1*. After

re-plating colonies from the galactose-containing media to ensure clonality, individual

colonies were checked by Sanger sequencing at biallelic regions to confirm loss of the

conditional chromosome. Each of these aneuploid (2N–1) strains was then sporulated

to produce an F2 population of haploid segregants fixed for a single parent's

chromosome (Fig 1C), which could then be used for BSA (Fig 1D).

Fig 1. Insertion of *GAL1* promoter allows for fixed chromosomes. (A–B) Matings to

produce reciprocal diploids with single conditional chromosomes for chromosome I (A)

and chromosome VIII (B). (C) Induction of the *GAL1* promoter adjacent to the

centromere destabilizes the chromosome's kinetochore, leading to loss of that

chromosome in diploids and fixation of the homologous chromosome in F2 haploid

progeny. One representative cross of the four (Oak I) is shown. (D) BSA experiment for

interactions with chromosome I, allowing detection of additive effects by comparison of

selected (blue) and unselected (white) bulks and of interactions with chromosome I

background (orange or purple). One representative haploid segregant is shown beneath

each bulk's flask.

Before proceeding to BSA, we demonstrated that the Oak and Wine parental

strains, which have been shown to differ phenotypically in multiple traits [31,32], differ in

the phenotype of copper sulfate resistance. We grew populations of haploids of each

parental strain in an array of doses of $CuSO_4$ in rich media (YPD), and compared the

cell counts of each population to a control of the same seed culture. The Wine haploid

strain showed higher resistance to $CuSO_4$, surviving at almost double the percentage at

which the Oak haploid strain survived in mid-range doses of $CuSO_4$ (S1 Fig).

Due to chromosome fixation, the relative intensity of selection varied at the same

doses of $CuSO_4$ between crosses. Specifically, as shown below, Wine chromosome VIII

has a strongly advantageous allele, so the same dose of $CuSO_4$ produces more

survivors in the Wine VIII cross than in the Oak VIII cross. We therefore tested multiple

doses of $CuSO_4$ to determine the dose that produced 1–10% survival of the population.

This range of selection optimizes BSA by balancing detectable effects and retention of

causative genetic variation in the surviving population [33]. Unselected bulks from each

cross were seeded in YPD at 10% the cell number at which selected bulks were seeded

so that their final cell densities were approximately the same as those in the selected

bulks.

Known drivers of resistance are identified as additive QTL

Previous studies have identified additive QTL for the trait of copper sulfate

resistance in a variety of *S. cerevisiae* strains [2,4,18,34]. In addition, multiple studies

have identified mechanistic drivers of copper resistance, including certain alleles of the

FRE1 gene and copy number amplification of the *CUP1* locus [27,35,36]. Because there

is overlap across strains for identification of some QTL [37], one form of validation of

epic-QTL would be for previously known resistance loci to be detected by epic-QTL as

additive effects (that is, effects that do not depend on the parental origin of the fixed

chromosome).

In epic-QTL, as in standard BSA, allele frequency differences between bulks at a

given genomic site indicate a contribution of variation at that site to bulk membership.

For the reciprocal chromosome I crosses and, separately, for the reciprocal

chromosome VIII crosses, we sequenced the four bulks (selected vs. unselected for

each Oak or Wine fixed chromosome) at high depth of coverage (average ~70X per site

per bulk) to estimate allele frequencies. We partitioned the effects of selection, parental-

chromosome background (hereafter "background"), and their interaction using a logistic

regression model as described in *Materials and Methods*. The selection term's

coefficient reflects the additive effect on $CuSO_4$ survival regardless of parental-

chromosome background, whereas the interaction term reflects epistasis with the fixed

chromosome. To visualize significant effects across the genome, we use the z-score of

each coefficient. The z-score is advantageous over the parametric *p*-value to which it

corresponds because its sign indicates the direction of effect of the fixed chromosome,

and because we use permutations of it to determine a threshold of false discovery rate,

detailed in *Materials and Methods*. To compare significance regardless of effect

direction, the majority of plots display absolute values of z-scores. Signed z-scores for

each coefficient and position are included in supplemental data (S1 Table).

The additive (selection) peaks on chromosomes VIII and XII are very close to the

known copper-resistance genes, *CUP1* and *FRE1*, respectively (Fig 2). These genes

are well documented to play critical roles in copper detoxification mechanisms in yeast.

CUP1 encodes a metallothionein involved in sequestering copper ions, thereby

mitigating copper toxicity within the cell [38]. *FRE1* encodes a ferric reductase enzyme

that contributes to iron uptake and detoxification of copper [39,40]. In both fixed-

chromosome backgrounds, we identify *FRE1* as a major driver of resistance, with peak

maxima <6 kb away from the coding region (Fig 2). The crosses for the two fixed-

chromosome backgrounds are independent, so the agreement between the two crosses

further validates the method. In the strains fixed for chromosome I, the peak on

chromosome VIII is upstream of the *CUP1* coding sequence. *CUP1* has two paralogs,

CUP1-1 and *CUP1-2*, within a larger tandemly duplicated region in the reference

genome of the laboratory strain S288C. We found that the region surrounding these

paralogs had poor mapping quality, likely due to repetitive sequences and potential

CUP1 copy variation, which is a known mechanism of resistance [38]. These regions

were therefore excluded from our variant calling, and so the peak upstream of the

coding sequence is as close as can be to this known driver of copper resistance.

Fig 2. Known drivers of copper resistance identified as additive QTL. Traces

represent absolute z-scores of additive (selection) coefficients on chromosomes VIII

and XII, as indicated. Dashed line corresponds to the experiment with chromosome I

fixed and solid line corresponds to the experiment with chromosome VIII fixed.

Horizontal lines indicate the 5% false discovery rate (FDR) threshold for each

experiment. Turquoise vertical lines represent the locations of CUP1 and FRE1, both of

which fall under large-effect additive peaks as expected.

Fixing Large Effect QTL Increases Power for Detection of

Additive Effects

The resolution and reproducibility of epic-QTL are further supported by additive

QTL that do not map to known copper-resistance loci. Between the experiment with

fixed chromosome I and the experiment with fixed chromosome VIII, we identified many

overlapping selection-QTL peaks with locations of highest absolute z-scores within 1.5

kb of one another, indicating not only that there is high resolution and reproducibility but

also that the underlying mechanisms of resistance do not strongly differ in the presence

of one or the other fixed chromosome (Fig 3, S2 Table).

Fig 3. Consistent additive peaks identified in each fixed parental-chromosome

experiment. Additive effects for the experiment with chromosome I fixed (dashed lines)

and the experiment with chromosome VIII fixed (solid lines), shown as the absolute

value of the selection coefficient's z-score, with 5% FDR thresholds shown as horizontal

lines of the same type. The only trace for chromosome I is solid, and the only trace for

chromosome VIII is dashed, because of the fixed chromosome in each experiment.

Peaks for each experiment are denoted as points on the traces as well as vertical lines

below (fixed chromosome I as light blue, top strip, and fixed chromosome VIII as dark

blue, bottom strip). In these strips, regions of significant z-scores are shown in gray,

with intensities proportional to absolute z-score.

Having established that our method identifies additive QTL, we next investigated

the result of chromosome fixation on additive effects. Removal of a major-effect QTL is

expected to increase the power to detect QTL of lesser effect [41], so we investigated

whether epic-QTL has an added benefit of increasing power when a major-effect QTL

exists on a fixed chromosome. We found that indeed the majority of peaks in the fixed

chromosome VIII experiment had z-scores of greater magnitude than the corresponding

peaks in the fixed chromosome I experiment, and that several additional peaks were

detected that were not called when chromosome I was fixed (Fig 3). On chromosome

XII in particular, only a single peak is obvious in the fixed chromosome I experiment,

whereas in the fixed chromosome VIII experiment, two peaks can clearly be

differentiated (Fig 3). Taken together, our results for additive QTL demonstrate that

epic-QTL has high power, resolution and reproducibility; that copper resistance in the

Oak X Wine cross progeny shares at least some underlying mechanisms with other *S.*

cerevisiae strains; and that removal of a major source of variance can increase power to

detect other additive QTL.

Magnitude epistasis is identified with chromosome VIII but not chromosome I

Reciprocal chromosome fixation effectively converts effects that are epistatic with

that chromosome into additive effects that can be detected with high power by BSA. In

the context of the epic-QTL logistic-regression model, these epistatic effects are

identified as significant selection-by-background interactions.

We identify seven interactions with chromosome VIII: one each on chromosomes

VII, X, XII, XIV, and XV, and two on chromosome XI (Fig 4, S3 Table). The z-score of

each interaction peak is less extreme than that of its corresponding additive effect, but

not always in the same direction as the additive effect, indicating that there is no

consistent interaction effect (such as the Oak chromosome always amplifying Oak

alleles' selection effects). We did not identify any interactions at a 5% FDR with

chromosome I (Fig 4A), despite observing >10 additive QTL. As chromosome I is the

smallest of the *S cerevisiae* chromosomes, it is plausible that there would not be any

interactions with the entire chromosome. This result provides evidence that epic-QTL

does not generate spurious interactions, and that interactions are specific and not

widespread across the genome.

Fig 4. Interaction effects indicate magnitude epistasis. (A) Fixed chromosome I

(top) and fixed chromosome VIII (bottom) absolute z-scores for $CuSO_4$ experiments,

with trace colors indicating additive (black) and interaction (red) effects.Vertical red lines

indicate where an interaction peak has been called, with solid lines indicating the

highest peak on a chromosome and dashed lines indicating a second significant peak.

(B) Allele frequencies of the Wine allele at the maximum point of each interaction QTL

per chromosome (indicated by solid lines in 4A and triangles in 4C), in each population.

Oak (O) and Wine (W) fixed chromosome VIII backgrounds are on the horizontal axis,

and the allele frequency of the Wine allele is on the vertical axis. Line crossing of

selected (blue) and non-selected (gray) would indicate sign epistasis. (C) Comparisons

of z-scores at interaction peaks (red triangles) with z-scores at corresponding additive

peaks (black triangles).

The presence of an interaction can indicate different forms of epistasis. One

possibility is sign epistasis, which in the case of epic-QTL would mean that changing the

parent of the fixed chromosome changes the direction of the QTL effect at the

interacting locus. An alternative possibility is magnitude epistasis, in which the QTL

effect size at the interacting locus changes but the direction of the effect remains the

same. In experimental designs such as standard BSA, where effects at any locus are

averaged across a population, sign epistasis obscures phenotypic effect as the effects

of allele pairs cancel each other out. Perfect sign epistasis, in which effect magnitude is

equal but effect direction is opposite, is therefore impossible to detect by BSA without

fixation of one allele in the pair to isolate the effect. In contrast, magnitude epistasis still

produces an additive effect, although the effect size will be the average of that of each

allele combination. To characterize our interaction QTL as exhibiting sign or magnitude

epistasis with the fixed chromosome, we: 1) compared the allele frequencies of the

highest points of the interaction peak (Fig 4C), and 2) scanned for loci above the 5%

FDR threshold where the interaction z-score was more extreme than its corresponding

additive z-score. Sign epistasis would require line-crossing of allele frequencies

between parental backgrounds in the selected bulks, which we do not observe at

interaction peaks (Fig 4B). We also do not find any significant interaction z-scores that

are greater in magnitude than their selection effect (Fig 4C), which would indicate that

the interaction obscures the additive effect through averaging. We therefore conclude

that epistasis with chromosome VIII is magnitude and not sign epistasis.

Prior studies have identified interaction QTL for copper resistance in *S.*

cerevisiae either by testing for interactions between additive QTL [4] or by barcoding

combinations of alleles in natural variants [21,22]. We compared interactions identified

with chromosome VIII in our study with those identified in these other studies, which

utilized strains other than the Oak and Wine strains used here. We find that there is little

overlap between identified loci (S2 Fig). There are, however, consistent additive effects

at specific loci such as *CUP1*, and indeed most interactions with chromosome VIII from

the other studies did involve the *CUP1* locus. These results suggest that interactions

are between specific alleles not necessarily present in all crosses, which could be

corroborated with additional epic-QTL experiments with different parental strains.

Further evidence that interaction effects are idiosyncratic is that not every

additive locus interacts with the fixed chromosome, and there is no correlation in the

direction or magnitude of effect of the interaction compared to additive effect (Fig 4C).

The majority of interaction peaks (4 of 7) were within 6 kb of the additive peak maxima,

indicating that the interactions with chromosome VIII are likely with the same loci that

produce additive effects. Of the remaining interaction peaks, two (the one on

chromosome VII and the lesser one on XI) overlap with what appear as shoulders of

additive peaks, suggesting that multiple QTL occur within that peak, and that the second

largest is interacting with chromosome VIII. The peak on chromosome XV might also

correspond more with the shoulder of the additive peak. The interaction peak and

corresponding additive peak shoulder on chromosome VII are close to the *CUP2* gene,

which induces transcription of *CUP1* [35]. *CUP2* could thus be a candidate for allele

substitution to confirm the interaction with *CUP1* on chromosome VIII. Overall, the co-

occurrences of interaction peaks with additive peaks indicate that interactions occur

primarily with loci that have an additive effect, and the lack of interactions at every

additive peak, or even the most significant additive peaks, suggests that interactions are

specific and idiosyncratic.

Discussion

Identifying epistatic contributions to trait variation presents challenges caused by

the combinatorial nature of interactions and their potential to obscure effects of

individual alleles. Methods to detect pairwise interaction effects suffer from low

statistical power due to the number of pairwise comparisons across the genome to test.

High power can be achieved by BSA experiments, but averaging over combinations of

alleles precludes analysis of interactions. Because of these challenges, approaches to

predicting the effect of genotype often minimize the contribution of epistasis to

phenotypic variance, by first (or only) fitting additive effects [9]. However, although the

effects of epistasis might be statistically underestimated, specific examples of epistasis

have been observed across species and traits such as in tomato size and fruit number

[42,43], corn kernel coloration [44], and mouse coat color (39, reviewed by 12). In non-

natural variants such as deletion mutations or engineered hypomorphs, interactions

have also been quantified; using high-throughput approaches, Costanzo *et.al.* found

~900,000 interactions among 23 million double mutants representing nearly every pair

of genes in the yeast genome [24], and Hale *et al.* found that over one quarter of 1721

CRISPR-interference perturbations showed growth effects that depend on genetic

background [46]. We developed epic-QTL with the aim of bringing the power of a high-

throughput, DNA sequencing-based approach to the identification of interactions

between natural genetic variants. Specifically, we circumvented the statistical problem

of pairwise testing of natural variants by reciprocally fixing a single chromosome from

each of two genetic backgrounds in a cross between divergent yeast strains. With the

high power afforded by BSA, we observe magnitude epistasis with additive QTL but no

evidence of cryptic sign epistasis.

We first verified our method by comparing additive QTL identified by epic-QTL,

both between experiments with chromosome I or chromosome VIII fixed and between

our study and those conducted on copper resistance in other strains. Known drivers of

copper resistance, *CUP1* and *FRE1*, were identified as additive QTL in all experiments

where the loci were allowed to segregate. Additive peaks colocalized between the

experiment with chromosome I fixed and the experiment with chromosome VIII fixed,

but on average those identified in the experiment with fixed chromosome VIII were

stronger, very likely due to the fixation of a major QTL. In comparison with other studies

of copper sulfate resistance, several additive QTL were found in common, but many

were not, suggesting strain-specific differences in which alleles are present that impact

resistance.

The key strength of epic-QTL is identification of interactions with fixed

chromosomes at high statistical power. We next compared interactions found by epic-

QTL to those identified for copper resistance using different strains, different methods

and smaller sample sizes [2,19,21,22], finding that although there is some overlap in

regions containing additive QTL, the majority of interactions are not common between

studies. This finding speaks to the importance of more high-powered studies of

interactions and potentially also suggests that interactions are idiosyncratic in that they

are dependent on the specific alleles found in divergent strains. Because we detected

more additive than interaction QTL, not every detected additive QTL in our study

contained interaction QTL. Moreover, interactions did not necessarily cluster only under

the highest additive peaks. This finding further supports that the interactions are

idiosyncratic, rather than globally caused by nonlinear transformation of the value of a

latent trait with additive genetic architecture [14]. One key overlap with another study to

note, however, is *CUP1* as a hub for interactions on chromosome VIII [22]. That

interactions occur with chromosome VIII but not chromosome I, and that interactions

occur primarily with additive QTL, suggests that our interactions might be with *CUP1*.

The logical next step would be to fix the chromosomes found to interact with fixed

chromosome VIII, allowing chromosome VIII to segregate, and thereby to identify the

interaction partner(s) on this chromosome.

The interaction QTL identified here provide evidence of three compelling trends

that imply a middle ground between completely idiosyncratic epistasis and highly

predictable global epistasis: 1) interaction peaks commonly colocalized with additive

peaks, 2) magnitude epistasis was observed whereas sign epistasis was notably

absent, and 3) interaction peaks differ between studies and are not driven by strength of

an additive peak. Sign epistasis has the potential to completely obscure additive effects,

and thus the lack of sign epistasis implies that interactions occur primarily with loci that

already have an effect. Precise colocalization of additive and interaction maxima

suggest that the additive QTL interacts with the fixed chromosome. These trends

corroborate the conclusion by Bloom *et al.* [4] that interactions occur primarily between

additive QTL, and they validate the approach of reducing the search space by only

looking at interactions between additive QTL as done by Nguyen Ba *et al.* [21].

Sign but not magnitude epistasis can influence available evolutionary trajectories

by creating valleys between peaks on a fitness landscape [11,13]. Examples of specific

reciprocal sign epistasis, for example mutually exclusive adaptive mutations in the

HXT6/7 and *MTH1* genes during experimental evolution under glucose limitation, have

been shown to cause fitness valleys [47]. Sign epistasis involving mutations in specific

interaction hubs has also been observed [25], but the general prevalence of sign

epistasis is yet to be determined. epic-QTL offers an unbiased, powerful method for

detecting sign epistasis. The lack of sign epistasis detected here therefore indicates that

it is relatively rare, at least for the trait, fixed chromosomes, and parental strains we

studied.

It will be important to see whether the trends we observed generalize to other

fixed chromosomes and other complex traits. A complete set of fixed-chromosome

strains would yield an additional benefit of very rapid assessment of the extent to which

epistasis impacts variation in a trait, in that a completely additive genetic architecture

would mean that the sum of chromosome effects should equal the difference between

the parental strains. Such a test only requires chromosome-substituted strains, not

segregants from them. When this kind of test was applied in mice and rats to a large

number of anatomical and physiological traits, it revealed much evidence for epistasis

[48].

475 A widespread lack of sign epistasis would greatly aid in identifying interactions in

476 studies where experimental manipulations of genotypes are not possible. Genome-wide

477 association studies in particular face challenges with respect to the number of subjects

478 relative to the number of possible interactions between markers, and so a variety of

479 methods have been proposed to avoid exhaustively searching all possible pairs of

480 polymorphisms [49]. Prioritizing interactions between additive loci rather than every

481 possible combination could therefore be a principled approach to increase the power to

482 detect interaction effects. Characterizing interaction effects in humans should in turn

483 allow for greater precision in predicting response to medication and susceptibility to

484 disease, as well as facilitate further understanding of the molecular mechanisms of

485 complex traits. In all, our characterization of epistasis not only enhances our

486 understanding of genetic interactions in model organisms but also has implications for

487 improvement of applications in personalized medicine and trait prediction across diverse

488 species.

Materials and Methods

Conditional Chromosome Strain Construction

492 Two divergent haploid yeast strains, one oak tree-derived (NCYC3631, MATα)

493 and one wine barrel-derived (NCYC3591, MATa) [29], were each engineered for

494 chromosome removal from diploids and later selection of haploid progeny. To facilitate

495 crosses, strains were modified by homologous recombination at *URA3*, *HO*, and *MATα*

496 using standard lithium-acetate-based transformation. *URA3* prototrophy was restored

497 (and *KanMX* removed) in wine-derived strains by insertion of *URA3* sequence amplified

from the wine-strain parent (BC187). The positive-selection marker *hph*, which confers

resistance to hygromycin, was removed from the *HO* locus of each parental strain to

allow the marker's use elsewhere. To remove *hph*, we conducted Delitto Perfetto

excision [50] using a cassette that allows positive selection for nourseothricin resistance

(*NatR*) and negative selection against HSV-*TK*, which encodes the thymidine kinase

from herpes simplex virus and thereby confers sensitivity to the pyrimidine analogue 5-

fluorodeoxyuridine (FuDR) [51]. Importantly, two identical sequences (*cyc* terminators)

flank this cassette and thereby lead to spontaneous excision of the cassette by

homology-based repair [50]. Transformants were selected for presence of *NatR* and

checked for loss of *hph*. We then counter-selected using FuDR to select for removal of

the cassette and resulting loss of both *NatR* and HSV-*TK*.

To select for haploids of one mating type in bulks to be sequenced, we

engineered the *MATα* locus to contain the HSV-*TK* negative-selection marker for

removal of any cell that is not a mating-type a haploid. *MATα* contains a bidirectional

promoter driving the expression of the *α1* and *α2* genes, both of which must be active

for the cell to sporulate [52]. The negative-selection marker was inserted upstream of *α1*

and its promoter, and the *STE3* promoter (*pSTE3*), which is activated by the

transcriptional co-activator encoded by *α1*, was placed upstream of *α2* [53]. Thus, the

selection cassette was placed in a way that maintains expression of both *α1* and *α2* in

mating-type α haploids and in diploids. As a result, the cells retain the ability to mate

with mating-type a cells as haploids and to sporulate as diploids, but both diploids and

haploid mating-type α haploids are sensitive to FuDR. For this transformation we used

the positive-selection marker *hph*. We transformed a cassette containing MATα

homology and the positive- and negative-selection markers into the Oak α strain, and

then replaced *hph* with *NatR* using lithium acetate homology-directed transformation,

allowing use of *hph* in later transformations.

For chromosome elimination by driving transcription through the centromere, we

constructed an inducible-promoter cassette by using overlapping PCR to combine the

HSV-*TK* and *hph* markers from Alexander *et al.* [51], the *GAL1* promoter (*pGAL1*), and

regions homologous to the yeast genome adjacent to the centromere. The centromere-

region primers matched the primers used by Reid *et al.* [30] or matched sequences very

close nearby. These cassettes were transformed into either chromosome I or

chromosome VIII in either the Oak α or Wine *a* strains. All strains are detailed in

supplemental data (S4 Table). These haploid strains are referred to as the background

(Wine a or Oak α) plus [I] or [VIII] to indicate the chromosome that can be eliminated.

Wine a[I] and Oak α were mated to produce diploids, and Wine a and Oak α[I]

were mated to produce diploids capable of producing the reciprocal chromosome

fixation (Fig 1A). Diploids were selected by ability to grow in both the absence of uracil

and the presence of nourseothricin, and ploidy was confirmed by flow cytometry using

DNA staining with propidium iodide on the Cytek Aurora. To fix chromosome I, diploids

were then plated onto medium containing 2% D-Galactose (and lacking glucose) to

induce the *GAL1* promoter's transcription through the centromere of the conditional

chromosome. Although we did not use markers to select for the loss of a chromosome,

plating on galactose and restreaking from single colonies was effective for yielding

chromosome loss. Colonies from the restreaked plate were checked by Sanger

sequencing to identify a loss of the conditional chromosome (i.e. by loss of

heterozygosity on that chromosome), then grown in liquid media and frozen as clonal

aneuploid (2N–1) strains, which we term Oak I or Wine I to indicate the parent of origin

of the remaining chromosome I in these otherwise hybrid strains. Populations of ~10^7

clonal aneuploids were sporulated using 2% potassium acetate for ~10 days at room

temperature (25 °C) to produce approximately $4X10^7$ segregants with identical

chromosome I of either Oak or Wine background but segregating parental genotypes at

all other loci. To select for haploids above and beyond the MATα selection, spores were

vortexed for 10 minutes in 50% diethyl ether to kill unsporulated diploids [54]. Surviving

spores were rinsed twice in sterile water, then germinated by incubation in 1 mg/mL

Zymolyase (Z100T) for 12 minutes before culturing in each experiment. The process

was completed identically for the reciprocal fixations of chromosome VIII.

Concentration Determination for Selection

We tested approximate concentrations of copper for selection by growing cells in

media of different concentrations for 24 hours (matching the experiment time) and then

counting the cells either by automatic cell counter (ThermoFisher Countess™ II) or by

manual counting on a hemocytometer. From there, we chose smaller ranges of doses

that would provide between 1% and 10% survival. After conducting experiments, cell

counts in some experiments were checked by hemocytometer to confirm that

experimental treatments were achieving the intended amount of selection.

Bulk Segregant Analysis

Pools of segregants with reciprocal fixed chromosomes (Oak I and Wine I, or

Oak VIII and Wine VIII) were grown in parallel for 24 hours in either YPD or YPD +

copper sulfate at doses determined in each chromosome-fixed bulk population to

produce between 1% and 10% survival. The control cultures grown in YPD were

seeded at 10% of the starting density for the selected group. Each of the four bulk

categories was grown in duplicate. After 24 hours, each of these bulks was harvested

by spinning down cells separately and extracting DNA either by phenol-chloroform-

isoamyl alcohol extraction and ethanol precipitation [55], or by Qiagen DNeasy Blood

and Tissue Extraction Kit 69506.

Sequencing

After DNA was harvested, single-end libraries for DNA sequencing were

prepared according to Baym *et al.*'s protocol [56] using Illumina kit FC-131-2001 and

KAPA HiFi amplification kit KK2612. Fragments were selected to be 200–600 bp by

AMPure XP bead cleanup (Fisher NC9959336), and pooled based on concentrations

determined by LifeTechnologies Qubit 3.0 and Roche 480 LightCycler qPCR. Libraries

were then sequenced on an Illumina NovaSeq 6000, with an average per-bulk depth of

~70X. Reads were aligned to the S288C reference genome R64 using BWA-MEM [57],

sorted using PicardTools [58], and then underwent quality control using Genome

Analysis Toolkit [59] command *BQSR*. Variants were called using GATK

HaplotypeCaller [60] as described by Mansfield and Grumet [61].

epic-QTL Analysis

VCFs were converted to tables of reads to analyze in R using GATK's

VariantsToTable function. We first removed all loci with >2 alternate alleles to control for

mapping quality issues, then assigned each alternate allele to the corresponding parent

based on SNP location from whole genome sequencing of the parent strains [62]. To

increase our signal-to-noise ratio and take advantage of the inherent correlation of

nearby genomic positions in a single F2 cross, we smoothed the read counts in rolling

windows of 200 SNPs using a Gaussian weighted mean.

To partition the effects of fixed-chromosome parent of origin, selection, and their

interaction, we performed logistic regression on these counts for each smoothed

position using the following formula:

$\log(\textit{allele odds}) = \beta 0 + \beta 1 \cdot \textit{selection} + \beta 2 \cdot \textit{background} + \beta 3 \cdot (\textit{selection} \times \textit{background}) + \epsilon$

where *selection* is the presence or absence of copper sulfate in the culture media,

background is the parent of origin of the fixed chromosome, and the

selection×background term captures the dependence on background of the selection

effect, i.e. epistasis. We then used the z-scores of each effect to identify additive

(*selection*) and epistatic (*selection×background*) QTL. Absent any effects of the fixed

chromosome's parent of origin on segregation of the other chromosomes or on growth

in the absence of copper sulfate, we do not expect to see a significant background

effect. For this reason, we set the contrasts of the selection variable to 0 (copper

absent) and 1 (copper present), and for the background variable to –0.5 (Wine) and 0.5

(Oak). Outputs of this analysis are available in supplemental data (S1 Table).

To establish our statistical threshold for interactions, we conducted permutations

of allele frequency ratios by shuffling positions within each parent and replicate in the

non-selected bulks. This permutation approach is based on the assumption that, in the

absence of selection, there would be no loci associated with the trait. For the

permutations, we excluded chromosomes I and VIII from all experiments to remove the

effect of the fixed chromosomes, and removed chromosomes III and V, which contained

engineered regions not identical in the two parents. We then assigned bulks randomly

to these permuted values according to the structure of the original dataset (the same

number of replicates in each group) and calculated our selection, background, and

interaction effects for each. These effects drawn from the bulks of two different parents

form the null distribution of z-scores of interaction effects for each experiment. All

analyses employ a false discovery rate threshold of 5%.

Peaks of absolute z-score (QTL locations) were called by identifying the

maximum points within chromosome segments bounded by chromosome ends or local

minima, defined as positions where the local slope of the absolute z-score trace shifted

from negative to positive. The local slope was calculated within sliding windows of 700

SNPs by linear regression within the window. The highest point in that segment, if

above the 5% FDR, was then called as the peak. Additive and interaction peak locations

are available in supplemental data (S2 and S3 Tables).

Acknowledgements

633 This work was supported in part through the NYU IT High Performance Computing

634 resources, services, and staff expertise. We acknowledge the Zegar Family Foundation

635 for their generous support. We thank the NYU Center for Genomics and System Biology

636 Genomics Core for their assistance and resources. We thank members and alumni of

637 the Siegal lab for input on the paper and discussion about the project.

References

1. Tsouris A, Fournier T, Friedrich A, Hou J, Dunham MJ, Schacherer J. Species-wide survey

of the expressivity and complexity spectrum of traits in yeast. PLOS Genet. 2024;20:

e1011119. doi:10.1371/journal.pgen.1011119

2. Bloom JS, Ehrenreich IM, Loo WT, Lite T-LV, Kruglyak L. Finding the sources of missing

heritability in a yeast cross. Nature. 2013;494: 234–237. doi:10.1038/nature11867

3. Mackay TFC, Stone EA, Ayroles JF. The genetics of quantitative traits: challenges and

prospects. Nat Rev Genet. 2009;10: 565–577. doi:10.1038/nrg2612

4. Bloom JS, Kotenko I, Sadhu MJ, Treusch S, Albert FW, Kruglyak L. Genetic interactions

contribute less than additive effects to quantitative trait variation in yeast. Nat Commun.

2015;6: 8712. doi:10.1038/ncomms9712

5. Falconer D, Mackay TFC. Introduction to Quantitative Genetics. IV. Longman Group Ltd;

1996.

6. Lynch M, Walsh B. Genetics and Analysis of Quantitative Traits. Sinauer Associates, Inc;

7. Boye C, Nirmalan S, Ranjbaran A, Luca F. Genotype × environment interactions in gene

regulation and complex traits. Nat Genet. 2024;56: 1057–1068. doi:10.1038/s41588-024-

01776-w

8.   Morgante F, Huang W, Maltecca C, Mackay TFC. Effect of genetic architecture on the

prediction accuracy of quantitative traits in samples of unrelated individuals. Heredity.

2018;120: 500–514. doi:10.1038/s41437-017-0043-0

9.   Ober U, Huang W, Magwire M, Schlather M, Simianer H, Mackay TFC. Accounting for

Genetic Architecture Improves Sequence Based Genomic Prediction for a Drosophila

Fitness Trait. 2015. doi:doi.org/10.1371/journal.pone.0126880

10. Mackay TFC, Anholt RRH. Pleiotropy, epistasis and the genetic architecture of quantitative

traits. Nat Rev Genet. 2024;25: 639–657. doi:10.1038/s41576-024-00711-3

11. Weinreich DM, Watson RA, Chao L. Perspective: Sign epistasis and genetic constraint on

evolutionary trajectories. Evol Int J Org Evol. 2005;59: 1165–1174.

12. Phillips PC. Epistasis—the essential role of gene interactions in the structure and evolution

of genetic systems. Nat Rev Genet. 2008;9: 855–867. doi:10.1038/nrg2452

13. Phillips PC, Otto SP, Whitlock MC. Beyond the average: the evolutionary importance of

gene interactions and variability of epistatic effects. Epistasis and the Evolutionary

Process. Oxford; 2000.

14. Otwinowski J, Plotkin JB. Inferring fitness landscapes by regression produces biased

estimates of epistasis. Proc Natl Acad Sci. 2014;111: E2301–E2309.

doi:10.1073/pnas.1400849111

15. Kryazhimskiy S, Rice DP, Jerison ER, Desai MM. Microbial evolution. Global epistasis

makes adaptation predictable despite sequence-level stochasticity. Science. 2014;344:

1519–1522. doi:10.1126/science.1250939

16. Johnson MS, Reddy G, Desai MM. Epistasis and evolution: recent advances and an

outlook for prediction. BMC Biol. 2023;21: 120. doi:10.1186/s12915-023-01585-3

17. Reddy G, Desai MM. Global epistasis emerges from a generic model of a complex trait.

Barkai N, Petrov DA, Crona K, Krug J, editors. eLife. 2021;10: e64740.

681 doi:10.7554/eLife.64740

682 18. Ehrenreich IM, Torabi N, Jia Y, Kent J, Martis S, Shapiro JA, et al. Dissection of genetically

683  complex traits with extremely large pools of yeast segregants. Nature. 2010;464: 1039–

684  1042. doi:10.1038/nature08923

685 19. Ehrenreich IM, Bloom J, Torabi N, Wang X, Jia Y, Kruglyak L. Genetic architecture of

686  highly complex chemical resistance traits across four yeast strains. PLoS Genet. 2012;8:

687  e1002570. doi:10.1371/journal.pgen.1002570

688 20. Nadeau JH, Singer JB, Matin A, Lander ES. Analysing complex genetic traits with

689  chromosome substitution strains. Nat Genet. 2000;24: 221–225. doi:10.1038/73427

690 21. Nguyen Ba AN, Lawrence KR, Rego-Costa A, Gopalakrishnan S, Temko D, Michor F, et

691  al. Barcoded bulk QTL mapping reveals highly polygenic and epistatic architecture of

692  complex traits in yeast. Verstrepen KJ, Barkai N, Verstrepen KJ, editors. eLife. 2022;11:

693  e73983. doi:10.7554/eLife.73983

694 22. Matsui T, Mullis MN, Roy KR, Hale JJ, Schell R, Levy SF, et al. The interplay of additivity,

695  dominance, and epistasis on fitness in a diploid yeast cross. Nat Commun. 2022;13: 1463.

696  doi:10.1038/s41467-022-29111-z

697 23. Michelmore RW, Paran I, Kesseli RV. Identification of markers linked to disease-resistance

698  genes by bulked segregant analysis: a rapid method to detect markers in specific genomic

699  regions by using segregating populations. Proc Natl Acad Sci U S A. 1991;88: 9828–9832.

700 24. Costanzo M, VanderSluis B, Koch EN, Baryshnikova A, Pons C, Tan G, et al. A global

701  interaction network maps a wiring diagram of cellular function. Science. 2016;353:

702  aaf1420. doi:10.1126/science.aaf1420

703 25. Ang RML, Chen S-AA, Kern AF, Xie Y, Fraser HB. Widespread epistasis among beneficial

704  genetic variants revealed by high-throughput genome editing. Cell Genomics. 2023;3:

705  100260. doi:10.1016/j.xgen.2023.100260

706 26. Rehman M, Liu L, Wang Q, Saleem MH, Bashir S, Ullah S, et al. Copper environmental

toxicology, recent advances, and future outlook: a review. Environ Sci Pollut Res. 2019;26:

18003–18016. doi:10.1007/s11356-019-05073-6

27. De Freitas J, Wintz H, Hyoun Kim J, Poynton H, Fox T, Vulpe C. Yeast, a model organism

for iron and copper metabolism studies. Biometals. 2003;16: 185–197.

doi:10.1023/A:1020771000746

28. Valentine JS, Gralla EB. Delivering Copper Inside Yeast and Human Cells. Science.

1997;278: 817–818. doi:10.1126/science.278.5339.817

29. Cubillos FA, Louis EJ, Liti G. Generation of a large set of genetically tractable haploid and

diploid Saccharomyces strains. FEMS Yeast Res. 2009;9: 1217–1225. doi:10.1111/j.1567-

1364.2009.00583.x

30. Reid RJD, Sunjevaric I, Voth WP, Ciccone S, Du W, Olsen AE, et al. Chromosome-Scale

Genetic Mapping Using a Set of 16 Conditionally Stable Saccharomyces cerevisiae

Chromosomes. Genetics. 2008;180: 1799–1808. doi:10.1534/genetics.108.087999

31. Clowers KJ, Heilberger J, Piotrowski JS, Will JL, Gasch AP. Ecological and Genetic

Barriers Differentiate Natural Populations of Saccharomyces cerevisiae. Mol Biol Evol.

2015;32: 2317–2327. doi:10.1093/molbev/msv112

32. Gerke JP, Chen CTL, Cohen BA. Natural Isolates of Saccharomyces cerevisiae Display

Complex Genetic Variation in Sporulation Efficiency. Genetics. 2006;174: 985–997.

doi:10.1534/genetics.106.058453

33. Magwene PM, Willis JH, Kelly JK. The Statistics of Bulk Segregant Analysis Using Next

Generation Sequencing. PLOS Comput Biol. 2011;7: e1002255.

doi:10.1371/journal.pcbi.1002255

34. She R, Jarosz DF. Mapping Causal Variants with Single-Nucleotide Resolution Reveals

Biochemical Drivers of Phenotypic Change. Cell. 2018;172: 478-490.e15.

doi:10.1016/j.cell.2017.12.015

35. Welch J, Fogel S, Buchman C, Karin M. The CUP2 gene product regulates the expression

of the CUP1 gene, coding for yeast metallothionein. EMBO J. 1989;8: 255–260.

36. Jungmann J, Reins HA, Lee J, Romeo A, Hassett R, Kosman D, et al. MAC1, a nuclear

regulatory protein related to Cu-dependent transcription factors is involved in Cu/Fe

utilization and stress resistance in yeast. EMBO J. 1993;12: 5051–5056.

doi:10.1002/j.1460-2075.1993.tb06198.x

37. Taylor MB, Ehrenreich IM. Higher-order genetic interactions and their contribution to

complex traits. Trends Genet. 2015;31: 34–40. doi:10.1016/j.tig.2014.09.001

38. Adamo GM, Lotti M, Tamás MJ, Brocca S. Amplification of the CUP1 gene is associated

with evolution of copper tolerance in Saccharomyces cerevisiae. Microbiol Read Engl.

2012;158: 2325–2335. doi:10.1099/mic.0.058024-0

39. Hassett R, Kosman DJ. Evidence for Cu(II) Reduction as a Component of Copper Uptake

by *Saccharomyces cerevisiae*(∗). J Biol Chem. 1995;270: 128–134.

doi:10.1074/jbc.270.1.128

40. Georgatsou E, Alexandraki D. Regulated expression of the Saccharomyces cerevisiae

Fre1p/Fre2p Fe/Cu reductase related genes. Yeast Chichester Engl. 1999;15: 573–584.

doi:10.1002/(SICI)1097-0061(199905)15:7<573::AID-YEA404>3.0.CO;2-7

41. Sinha H, David L, Pascon RC, Clauder-Münster S, Krishnakumar S, Nguyen M, et al.

Sequential Elimination of Major-Effect Contributors Identifies Additional Quantitative Trait

Loci Conditioning High-Temperature Growth in Yeast. Genetics. 2008;180: 1661–1670.

doi:10.1534/genetics.108.092932

42. Eshed Y, Zamir D. Less-than-additive epistatic interactions of quantitative trait loci in

tomato. Genetics. 1996;143: 1807–1817. doi:10.1093/genetics/143.4.1807

43. Aguirre L, Hendelman A, Hutton SF, McCandlish DM, Lippman ZB. Idiosyncratic and dose-

dependent epistasis drives variation in tomato fruit size. Science. 2023;382: 315–320.

doi:10.1126/science.adi5222

44. Ford RH. Inheritance of Kernel Color in Corn: Explanations & Investigations. Am Biol

Teach. 2000;62: 181–188. doi:10.2307/4450870

45.   Bennett DC, Lamoreux ML. The color loci of mice--a genetic century. Pigment Cell Res.

2003;16: 333–344. doi:10.1034/j.1600-0749.2003.00067.x

46.   Hale JJ, Matsui T, Goldstein I, Mullis MN, Roy KR, Ville CN, et al. Genome-scale analysis

of interactions between genetic perturbations and natural variation. Nat Commun. 2024;15:

4234. doi:10.1038/s41467-024-48626-1

47.   Kvitek DJ, Sherlock G. Reciprocal Sign Epistasis between Frequently Experimentally

Evolved Adaptive Mutations Causes a Rugged Fitness Landscape. PLOS Genet. 2011;7:

e1002056. doi:10.1371/journal.pgen.1002056

48.   Shao H, Burrage LC, Sinasac DS, Hill AE, Ernest SR, O'Brien W, et al. Genetic

architecture of complex traits: Large phenotypic effects and pervasive epistasis. Proc Natl

Acad Sci. 2008;105: 19910–19914. doi:10.1073/pnas.0810388105

49.   Van Steen K, Moore JH. How to increase our belief in discovered statistical interactions via

large-scale association studies? Hum Genet. 2019;138: 293–305. doi:10.1007/s00439-

019-01987-w

50.   Storici F, Resnick MA. Delitto PerfettoTargeted Mutagenesis in Yeast with

Oligonucleotides. In: Setlow JK, editor. Genetic Engineering. Boston, MA: Springer US;

2003. pp. 189–207. doi:10.1007/978-1-4615-0073-5_9

51.   Alexander WG, Doering DT, Hittinger CT. High-Efficiency Genome Editing and Allele

Replacement in Prototrophic and Wild Strains of Saccharomyces. Genetics. 2014;198:

859–866. doi:10.1534/genetics.114.170118

52.   Haber JE. Mating-Type Genes and *MAT* Switching in *Saccharomyces cerevisiae*.

Genetics. 2012;191: 33–64. doi:10.1534/genetics.111.134577

53.   Sprague GF, Jensen R, Herskowitz I. Control of yeast cell type by the mating type locus:

positive regulation of the alpha-specific STE3 gene by the MAT alpha 1 product. Cell.

1983;32: 409–415. doi:10.1016/0092-8674(83)90460-9

54.   Dawes IW, Hardie ID. Selective killing of vegetative cells in sporulated yeast cultures by

exposure to diethyl ether. Mol Gen Genet MGG. 1974;131: 281–289.

doi:10.1007/BF00264859

55.   Green MR, Sambrook J. Isolation and Quantification of DNA. Cold Spring Harb Protoc.

2018;2018. doi:10.1101/pdb.top093336

56.   Baym M, Kryazhimskiy S, Lieberman TD, Chung H, Desai MM, Kishony R. Inexpensive

Multiplexed Library Preparation for Megabase-Sized Genomes. PLOS ONE. 2015;10:

e0128036. doi:10.1371/journal.pone.0128036

57.   Li H, Durbin R. Fast and accurate long-read alignment with Burrows–Wheeler transform.

Bioinformatics. 2010;26: 589–595. doi:10.1093/bioinformatics/btp698

58.   Picard Tools - By Broad Institute. [cited 26 Sep 2024]. Available:

http://broadinstitute.github.io/picard/

59.   Van Der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A, et

al. From FastQ Data to High-Confidence Variant Calls: The Genome Analysis Toolkit Best

Practices Pipeline. Curr Protoc Bioinforma. 2013;43. doi:10.1002/0471250953.bi1110s43

60.   Poplin R, Ruano-Rubio V, DePristo MA, Fennell TJ, Carneiro MO, Van Der Auwera GA, et

al. Scaling accurate genetic variant discovery to tens of thousands of samples. 2017.

doi:10.1101/201178

61.   Mansfeld BN, Grumet R. QTLseqr: An R Package for Bulk Segregant Analysis with Next-

Generation Sequencing. Plant Genome. 2018;11. doi:10.3835/plantgenome2018.01.0006

62.   Ziv N, Shuster BM, Siegal ML, Gresham D. Resolving the Complex Genetic Basis of

Phenotypic Variation and Variability of Cellular Growth. Genetics. 2017;206: 1645–1657.

doi:10.1534/genetics.116.195180

Supporting information captions

S1 Fig. Survival percentage by strain in CuSO4. Boxplots of survival percentages of

Oak (NCYC3631) and Wine (NCYC3591) haploid strains at different concentrations of

copper sulfate are shown. Replicate experiments carried out on different dates are

represented by different shapes.

S2 Fig. Comparison with copper sulfate interaction peaks from previous studies.

Interactions with chromosome VIII identified in our study (shown in black trace of the

absolute z-score by genome position) are compared with interaction QTL involving

chromosome VIII identified in other copper-resistance studies, which used both different

experimental designs and different strains. Beneath the trace from this study, NB

(magenta) indicates interaction QTL as identified by Nguyen Ba et al [21], M (turquoise)

represents interaction hotspots between strains from Matsui et al. [22], and B (dark

blue) represents interaction QTL as identified by Bloom et al [4]. Red vertical lines

indicate significant peaks called in our study. *CUP1* is indicated by the vertical light blue

dotted line on Chromosome VIII.

S1 Table. Logistic regression output. Direct output from logistic regression on

sequencing data for each divergent SNP that passes quality filters (see github for code).

Each row (numbered in column A) records the effect term in the model (Background,

Bulk, Interaction, or intercept (column B, "label"); the chromosome fixed in the

experiment (column C, "chr"); the chromosome containing the SNP (column D,

"CHROM"); the chromosomal position of the SNP (column E, "POS"); the z-score of that

SNP in that experiment and for the specified effect (column F, "zscore"); and the

threshold value for a 5% FDR for the specified effect and experiment (column G, "q5").

S2 Table. Additive-effect peaks. Peaks called for the Bulk effect. Each row records

the chromosome fixed in the experiment (column A, "CSS"), the location of the peak

SNP as its chromosome (column B, "CHROM") and chromosomal position (column C,

"POS"), and the z-score of the peak SNP (column D, "zscore").

S3 Table. Interaction peaks. Peaks called for the Interaction effect. Columns are as in

S2 Table.

S4 Table. Yeast strains constructed and used in experiments. Backgrounds,

genotypes, and aliases of all strains used, including those representing intermediate

steps to construct strains for epic-QTL experiments. MLS.CB refers to strains made in

this study; NCYC refers to strains from the NCYC collection [29].

Fig 2

Fig 3

Fig 1

Fig 4