1

Specialized axon initial segment enables low firing threshold and rapid action potential output in fast-spiking interneurons of the human neocortex

Specialized axon initial segment in human neurons

Emoke Bakos[¶1,2], Ádám Tiszlavicz[¶1], Viktor Szegedi[1,2], Abdennour Douida[1], Szabina Furdan[1], Daphne K. Welter[3], Jonathan M. Landry[3], Balazs Bende[4], Gabor Hutoczki[5], Pal Barzo[6], Gabor Tamas[7], Vladimir Benes[3], Attila Szucs[8], Karri Lamsa*[1,2]

ORCID Identifiers:

Emoke Bakos:	https://orcid.org/0009-0001-4790-4352
Adam Tiszlavicz:	https://orcid.org/0009-0008-4830-0594
Viktor Szegedi:	https://orcid.org/0000-0003-4191-379X
Szabina Furdan:	https://orcid.org/0000-0002-8878-0966
Daphne K. Welter:	https://orcid.org/0000-0002-2430-2391
Jonathan M. Landry:	https://orcid.org/0000-0003-2262-9099
Gabor Hutoczki:	https://orcid.org/ 0000-0001-7048-0757
Pal Barzo:	https://orcid.org/0000-0001-8717-748X
Gabor Tamas:	https://orcid.org/0000-0002-7905-6001
Vladimir Benes:	https://orcid.org/0000-0002-0352-2547
Attila Szucs:	https://orcid.org/0000-0001-9733-4135
Karri Lamsa:	https://orcid.org/0000-0002-4609-1337

Affiliations

[1]Hungarian Center of Excellence for Molecular Medicine Research Group for Human

neuron physiology and therapy, Szeged, Hungary

[2]Department of Physiology, Anatomy and Neuroscience, University of Szeged,

Szeged, Hungary

[3]European Molecular Biology Laboratory, Heidelberg, Germany

[4]Hungarian Centre of Excellence for Molecular Medicine Research Group for

Translational Medicine Development, Szeged, Hungary

[5]Department of Neurosurgery, University of Debrecen Clinical Centre, Debrecen

Hungary

[6]Department of Neurosurgery, University of Szeged, Szeged, Hungary

[7]ELKH-SZTE Research Group for Cortical Microcircuits, Department of Physiology,

Anatomy and Neuroscience, University of Szeged, Szeged, Hungary

[8]Neuronal Cell Biology Research Group, Eötvös Loránd University, Budapest,

Hungary

*Corresponding author

Email: klamsa@bio.u-szeged.hu; karri.lamsa@hcemm.eu (KL)

¶These authors contributed equally to this work.

Abstract

The mammalian brain exhibits notable interspecies variation. Microanatomical and

molecular differences in homologous neurons, those with similar locations and developmental

origins across species, are best characterized in the neocortical mantle, the center of complex

brain functions; however, the purpose of these differences remains unclear. We performed

whole-cell microelectrode recordings along with microanatomical and molecular analyses of

human fast-spiking parvalbumin (pvalb)-expressing interneurons in neocortical tissue resected

during brain surgery, comparing them with similar data obtained from the mouse neocortex.

The action potential (AP) firing threshold was lower in human neurons than in mouse neurons.

This was due to a deficiency in low-voltage–activated inhibitory Kv1.1 and Kv1.2 potassium

channels in the axon initial segment (AIS), a specialized axonal region that determines AP

threshold and initiation, in human cells. In contrast, Kv1 ion channels were prominent in mouse

neurons. The AIS was also elongated in humans. Computational simulations of fast-spiking

interneurons revealed that the human-type AIS lowers the AP threshold and shortens the time

lag for AP initiation. We found that the low membrane AP firing threshold in pvalb neurons is

closely linked to slow membrane potential kinetics in the soma. Thus, the human AIS supports

fast in–fast out circuit function in human pvalb neurons, compensating for electrically slow

somatic membrane responses. When formulating therapeutic strategies that involve fast-

spiking neurons, it is crucial to take into account the molecular and functional species

differences.

Short blurb: Fast-spiking neurons in the human neocortex feature structural and molecular

specializations in the axon initial segment that lower firing thresholds and minimize input–

output delay.

Introduction

All organs in the body evolved specialized functions to optimally support animal life and vitality. The mammalian brain exhibits interspecies differences not only in volume and cortical convolution but also in nerve cell microstructure and molecular composition [1, 2]. Although the brain comprises homologous neurons, sharing similar locations and developmental origins across species, these cells show species-specific differentiation at the microscale level [3-6]. The purpose of this neuronal specialization across species remains poorly understood [7, 8], although these adaptations likely optimize neural circuit function [9]. Neuron types and their species-specific features are best characterized in the neocortical mantle, the center of complex high-order brain functions [1, 5, 6, 10-19]. Neocortical function has been intensively studied in relation to brain cognitive and behavioral processes in humans and animal models, the latter aiding in elucidating the roles of neuron types in these processes [20-22]. However, studies using resected human brain tissue have revealed various "human-specific" transcriptomic and microstructural features in neocortical neurons [7, 10, 11, 23-31]. Many of these differences likely contribute to species-specific neuronal functions; however, the structure–function relationship remains unclear [12, 16, 17, 19, 30, 32-34].

Distinctive differences in homologous neurons between the rodent and human neocortex include electrical resistivity and the membrane potential level required to trigger action potentials (APs) [25]. Human neocortical excitatory pyramidal neurons exhibit lower input resistance (Rin) and require greater membrane depolarization (i.e., a higher AP firing threshold) to trigger APs compared with rodent neurons [24, 29, 35]. In contrast, inhibitory interneurons in the human neocortex display higher input resistance and a lower AP firing

threshold relative to their rodent counterparts [26, 27]. Prior studies suggest that these interspecies differences also apply to fast-spiking interneurons [28, 31, 36], the principal inhibitory neuron population in the neocortex, characterized by the expression of *parvalbumin* (*PVALB*), the gene encoding pvalb protein [23]. In humans, pvalb neurons exhibit elevated input resistance; therefore, they generate membrane potential (Vm) changes with reduced electric input [28]. For the same reason, Vm changes, such as excitatory synaptic potentials that lead to APs, occur more slowly [28, 31, 36-39]. The delayed Vm response in human fast-spiking neurons, compared with that in their rodent counterparts, appears counterintuitive, as the rapid transformation of excitatory input to AP output (i.e., input–output transformation) is a hallmark of pvalb interneurons, also known as "fast in–fast out" inhibitory circuits [38, 40]. Pvalb interneurons play a crucial role in complex brain functions. They rapidly inhibit other neurons and orchestrate temporally coordinated network activities that underlie cognitive processes. Their ability to rapidly and reliably inhibit other neurons allows them to precisely control local circuits and shape brain network activity. We hypothesize that a low AP threshold coexists with slow membrane potential kinetics in human pvalb neurons to accelerate their operation in these processes.

In the present study, whole-cell electrophysiological recordings in human and mouse *ex vivo* neocortical slices revealed an association between slow somatic Vm changes and a lowered AP generation threshold in pvalb/*PVALB* interneurons. The reduced AP firing threshold in human cells was accompanied by structural and molecular specialization of the axon initial segment (AIS), a speciated axonal structure that initiates APs [41, 42], which is elongated and deficient in inhibitory Kv1 potassium channels in human neurons. Simulations of fast-spiking interneurons with a human-type AIS showed that a reduced AP threshold shortens the latency of AP initiation, enabling human pvalb neurons to function as fast in–fast out circuits despite slower somatic membrane responses [43].

Results

Pvalb interneurons in humans exhibit a lower AP firing threshold

We measured AP firing thresholds in 91 pvalb-expressing fast-spiking interneurons

[43] from human neocortical tissue resected during deep-brain surgery in 67 patients (Table

S1). The cells were studied using whole-cell clamp recordings in layer 2/3 of acute brain slices

sectioned from the frontal (n = 39 cells), temporal (n = 25 cells), and parietal, occipital, or other

neocortical regions (n = 27 cells) in male (n = 30) and female (n = 34) patients aged 11–85

years. The neurons were filled with biocytin during recordings and visualized *post hoc* using

streptavidin–fluorophore conjugates. All cells were immunopositive for pvalb (n = 79; Fig

1A1–3) or expressed *PVALB* based on mRNA analysis using patch-sequencing following

nucleus harvesting (n = 12; Table S1). For comparison, we also investigated pvalb-expressing

interneurons in the mouse neocortex (n = 95).

Fig. 1. Lowered AP firing threshold in fast-spiking pvalb interneurons in the human versus

mouse neocortex.

(A) AP firing threshold was examined in layer 2/3 interneurons showing a positive immunoreaction for

pvalb. (A1) Excitatory Vm steps (incremental amplitude steps under whole cell current–clamp

conditions at −70 mV) revealed the AP threshold. (A2) Human neocortex pvalb neurons were visualized

postrecording; the soma and dendrites (black) and axon (red) were filled with biocytin and are shown

in three merged 50-µm-thick sections. (A3) Confocal images illustrate pvalb immunoreactivity (Alexa

488) in the cell shown in (*A2*) and a biocytin-filled axon (bcyt, Cy3). Arrows mark colocalized signals.

(B, C) AP phase plot defining the firing threshold. (B1) Four superimposed APs evoked as in *A1* in

human pvalb cells. (B2) Phase plots showing the first derivative (dV/dt) against Vm (four superimposed

curves). (B3) Close-up of the initial AP rise; dotted vertical lines indicate threshold Vm where (dV/dt)

reaches 10 mV/ms (horizontal dotted line). (C1) Mouse pvalb cell APs (four superimposed), with

corresponding phase plots (C2) and the firing threshold definition (C3). (D) Human pvalb neurons s

how a lower AP threshold compared with mouse neurons (Mann–Whitney U test). (D1) AP threshold

histograms for human (blue, 91 cells) and mouse (gray, 95 cells) neurons; bin size: 1 mV. Red lines

denote median (solid) and quartiles (dotted). (D2) Cumulative histograms of AP threshold in human

(blue) and mouse (gray) pvalb cells. (E) In human pvalb neurons, AP threshold was not influenced by

(E1) neocortical region [n = 39 cells, frontal cortex (FC); n = 25 cells, temporal cortex (TC)], (E2) sex

(threshold calculated as the average per patient; 34 females, 30 males; Mann–Whitney U test), or (E3)

age (average per patient; n = 62; Spearman's correlation). Regression line is shown.

We applied positive Vm steps (square-pulse current steps of increasing amplitude, 500 ms at 1

Hz) to cells current-clamped at −70 mV to identify the AP threshold (Fig 1B, C) (defined as

the Vm at which the AP initial rise speed reached 10 mV/ms; hereafter, referred to as the firing

threshold; Fig 1A1) [31, 36, 44]. Human pvalb neurons had a lower AP threshold [median =

−38.9 mV, interquartile range (IQR) = −42.8 to −34.2, n = 91 cells] compared with mouse

neurons (median = −32.5 mV, IQR = −35.9 to −28.3, n = 95 cells, p < 0.001; Fig 1D1, 2). They

also exhibited lower rheobase current (human cells: 87.34 pA, IQR = 16.8–140.0; mouse cells:

247.65 pA, IQR = 53.95–418.4; p < 0.001), a moderately lower AP initial rise slope (human

cells: 30.9 ms^{-1}, IQR = 23.82–35.50; mouse cells: 33.2 ms^{-1}, IQR = 28.90–39.01; p = 0.0011),

and slower AP maximum fall speed (human cells: −122.83 mV/ms, IQR = −165.0 to −82.7;

mouse cells: −153.7 mV/ms, IQR = −213.9 to −113.7; p = 0.001). AP maximum rise speed was

similar across species (human cells: 246.94 mV/ms, IQR = 170.52–335.95; mouse cells: 255.33

mV/ms, 202.10–302.77; p = 1.00). Human neurons also had wider AP half-width (p < 0.001)

and shorter latency to AP generation at rheobase compared with mouse neurons (p < 0.001),

but no differences were observed in AP peak values (p = 1.00) or afterhyperpolarization (AHP)

amplitudes (p = 0.779). These AP properties and interspecies comparisons are summarized in

Table 1 (analyzed via PERMANOVA and Mann–Whitney U test with Bonferroni correction).

Table 1. Features of human and mouse APs from Fig 1.

	Human n = 91			Mouse n = 95			
	Median	1st quartile	3rd quartile	Median	1st quartile	3rd quartile	P-value
AP Thr (mV)	−38,9	−42,8	−34,20258	−2,46562	−35,9	−28,3	**<0.001**
Initial rise slope (ms^{-1})	30,9	23,82813	35,5	33,2	28,9	39,01236	**0.037**
AP peak Vm value (mV)	15,6	9,08	25	15,85557	8,17617	23,54049	1
AP half width (ms)	0,496	0,407	0,676	0,36222	0,32	0,416	**<0.001**
AP rheobase (pA)	87,36667	47,53333	140	247,65	142,2	418,4	**<0.001**
AP latency (ms)	19,618	12,6	33,718	50,5	17,94	79,87667	**<0.001**
Spike number by step	3,83333	1,83333	11,75	5	2,5	8	1
AP AHP ampl (mV)	−19,66087	−16,1987	−23,5826	−18,6386	−15,89205	−21,07239	0.7794
Max rise speed (mV/ms)	246,93813	170,51688	335,94748	255,3305	202,10315	320,77388	1
Max fall speed (mV/ms)	−122,83367	−164,99846	−82,73656	−153,73228	−213,90914	−113,678	**0.001**

AP Thr: action potential firing threshold; initial rise slope: maximum slope of the tangent line in the AP

phase plot above the threshold (10 mV/ms); AP half-width: measured at 50% of the maximal amplitude

from the AP firing threshold to the peak; AP rheobase: amplitude of depolarizing current required to

trigger an AP at the AP Thr; AP latency: average time from rheobase current onset to AP onset (defined

at AP Thr); spike number: number of APs evoked by the rheobase current step; AP AHP ampl.:

amplitude of afterhyperpolarization, measured from AP Thr to AP negative peak; max rise speed:

maximum AP rise speed, measured from the AP phase plot; max fall speed: maximum AP fall speed,

measured from the AP phase plot). Bolded p-values indicate species differences (PERMANOVA and

Mann–Whitney U test with Bonferroni correction).

We confirmed that AP thresholds in human neurons were similar between frontal (−39.0 mV,

IQR = −42.2 to −30.6, n = 39 cells) and temporal cortices (−38.5 mV, IQR = −45.1 to −36.0, n

= 25 cells; p = 0.248, Mann–Whitney U test; Fig 1E1). Average AP threshold per subject

showed no correlation with demographic factors. Across patients, AP firing thresholds were

similar in males (−38.8 mV, IQR = −43.2 to −34.4, n = 30 patients) and females (−39.0 mV,

IQR = −44.2 to −33.1, n = 34 patients; p = 0.956, Mann–Whitney U test; Fig 1E2) and unrelated

to age (r = −0.129, n = 62 patients; p = 0.315, Spearman's correlation; Fig 1E3). These results

show that human pvalb neurons exhibit a lower AP firing threshold in response to somatic

excitation compared with mouse pvalb neurons.

Low AP threshold is associated with electrically slow somata in

pvalb interneurons

Compared with mouse pvalb neurons, human pvalb neurons exhibited higher electrical

resistance, i.e., input resistance (Rin) in their somata, with no difference in capacitance (Cm).

We measured average Rin and Cm using subthreshold Vm steps of +5 to +10 mV (elicited by

square-pulse currents at −70 mV; Fig 2A1). Median Rin was 187.9 MΩ (IQR = 103.8–279.8)

in human cells (n = 60) versus 125.7 MΩ (IQR = 89.5–167.5) in mouse cells (n = 50; p < 0.001,

Mann–Whitney U test; Fig 2A2). Median Cm in the soma was similar across species (humans:

32.4 pF, IQR = 24.3–53.7, n = 60; mice: 35.1 pF, IQR = 26.3–50.6, n = 50; p = 0.728, Mann–

Whitney U test).

Fig. 2. Low AP firing threshold in pvalb interneurons is associated with an electrically slow soma.

(A) Human pvalb cells exhibited higher Rin compared with mouse cells at Vm between −70 mV and

the AP threshold. (A1) Cell Rin was measured using positive current steps inducing 5–10-mV

subthreshold depolarizations. Voltage shifts (ΔVm) and currents steps (ΔI) used to calculate average

Rin are shown in red. (A2) Histograms and box plots show Rin in human (blue) and mouse (gray) pvalb

neurons; bin size: 25 MΩ; box plots present the median, quartiles, and 5th and 95th percentiles (Mann–

Whitney U test). (B) Tau was longer in human than in mouse pvalb cells for Vm shifts in the soma.

(B1) Tau was measured from the ascending phase in +10-mV steps delivered at −70 mV (human: blue;

mouse: gray). Left: Four superimposed +10-mV steps in human and mouse cells. Right: Ascending

phase shows species differences in tau (human average: 8.9 ms; mouse average: 3.4 ms). (B2)

Histograms and box plots illustrate tau differences in human (blue) and mouse (gray) pvalb neurons;

bin size: 0.5 ms (Mann–Whitney U test). (C) Soma tau and Rin were strongly associated in mouse (C1)

but not human (C2) pvalb neurons (Spearman's correlation). (D) Low AP threshold was linked with

slow tau in mouse (D1) and human (D2) pvalb neurons. (D3) Combined data from both species revealed

a strong association between slower tau and lower AP threshold (Spearman's correlation).

The time course of Vm changes (tau), measured from the ascending phase of +10 mV steps,

was slower in human pvalb neurons (median tau of the Vm step: 6.65 ms, IQR = 4.22–8.17, n

= 60) than in mouse pvalb neurons (median: 4.17 ms, IQR = 3.27–6.12, n = 50; $p < 0.001$,

Mann–Whitney U test; Fig 2B1, 2). In mice, tau and Rin showed a significant positive

correlation (r = 0.613, n = 50, $p < 0.001$; Fig 2C1), whereas no such correlation was found in

human pvalb neurons (r = 0.224, n = 60, p = 0.085) (Fig 2C2). However, in both species, lower

AP thresholds were associated with slower tau (human cells: r = −0.274, n = 60, p = 0.034;

mouse cells: r = −0.535, n = 50, $p < 0.001$; Fig 2D1, 2). Combined data from both species

revealed a strong negative correlation between AP threshold and slow tau (r = −0.514, n = 110;

$p < 0.001$, Spearman's correlation; Fig 2D3).

Collectively, these results demonstrate that pvalb neurons with slower Vm kinetics to

depolarization in the soma tend to have lower AP firing thresholds. This association was

particularly strong when analyzing human and mouse pvalb neurons together.

AIS is elongated in human pvalb neurons and exhibits an axonal position associated with AP threshold

To explore whether AIS morphology contributes to the lower AP threshold in human pvalb neurons, we visualized the AIS, a primary site of AP initiation, as its length and position can affect AP threshold [45]. We used immunohistochemistry against beta-IV-spectrin, a cytoskeletal protein enriched in the AIS [42, 46, 47], analyzing its immunofluorescence in three-dimensional (3D) confocal microscopy image stacks, with AIS length and distance from the soma quantified using a custom script. All cells were located in layer 2/3 of the neocortex and confirmed as pvalb-positive (Fig 3A1).

Fig 3. AIS is elongated in human pvalb neurons and axonally located according to AP threshold.

(A) AIS in pvalb neurons was longer in humans than in mice. AIS analysis was performed in triple-immunostained neurons from the human and mouse neocortex. (A1) AIS was measured in 3D fluorescence confocal images of layer 2/3 pvalb-positive neurons. Left: AIS visualized via beta-IV-

spectrin immunofluorescence. Arrows indicate AIS initiation and termination sites (Alexa 488) in a collapsed z-stack image of a human pvalb neuron. Right: Merged beta-IV-spectrin and pvalb (Cy5) immunoreactions. Horizontal dotted curves mark (i) the axon from the soma (s) and the proximal (ii) and (iii) distal AIS ends. (A2) Histograms and box plots summarize AIS parameters from the upper schematic: a = distance from the soma to the AIS initiation site, b = AIS length, and c = distance from the soma to the AIS end. Top: AIS initiation distance from the soma did not differ between species. Middle: AIS was longer in human neurons than in mouse neurons. Bottom: AIS termination site was more distal in human neurons than in mouse neurons. Histogram bin: 1 μm, with kernel density; box plots show the median, quartiles, and 5th and 95th percentiles. (A3) Cumulative AIS data show p-values (Student's *t*-test). (B, C) AIS measurements in electrophysiologically investigated pvalb neurons revealed that a low AP threshold is associated with a more distal AIS termination site. (B1) AIS visualized via beta-IV-spectrin immunopositivity (Cy3) in human and mouse cells filled with biocytin (bcyt) and labeled with streptavidin–Alexa 488 conjugates. Immunofluorescence images show AIS features defined in *A1*. (B2) Electrophysiologically studied human pvalb neurons displayed a longer AIS (*b*) with a more distal termination site (*c*) compared with mouse neurons, whereas the AIS start distance (*a*) from the soma did not differ between species (Mann–Whitney U test). Combined AP threshold data from human (blue) and mouse (gray) pvalb neurons revealed associations with AIS initiation site (C1) and termination site (C3) distance from the soma but no association with AIS length (C2) (Spearman's correlation).

Both human and mouse pvalb neurons exhibited cell-to-cell variation in AIS length and positioning along the axon (Fig 3A2). The median AIS start point was similar between species (humans: 4.12 μm, IQR = 2.95–5.28, n = 100 cells in 17 patients; mice: 3.77 μm, IQR = 2.69–4.68, n = 78 cells in 15 mice; p = 0.265), but the AIS was longer in human cells (median: 16.68 μm, IQR = 14.08–19.78) than in mouse cells (median: 13.99 μm, IQR = 11.15–16.55; p < 0.001). Consequently, the AIS termination site was further from the soma in human neurons

(21.01 μm, IQR = 17.60–24.66, n = 100 cells from 17 samples) than in mouse neurons (17.47

μm, IQR = 14.41–20.66 μm, n = 78 cells from 15 mice; p < 0.001, Mann–Whitney U test; Fig

3A2, 3).

We also investigated whether these morphological differences relate to the lower AP threshold

observed in humans. In electrophysiologically studied and biocytin-filled neurons visualized

via streptavidin–fluorophore conjugates, with confirmed immunopositivity for pvalb, we

observed the AIS based on beta-IV-spectrin positivity and measured its length and distance

from the soma along the axon, comparing these metrics against the AP firing threshold. Of the

28 human and 22 mouse cells recorded in the whole-cell clamp, intact somata and full-length

AIS were successfully recovered (indicated by beta-IV-spectrin immunofluorescence) in 13

human and 14 mouse neurons. In particular, only neurons with biocytin-filled axons extending

beyond the distal end of the beta-IV-spectrin–labeled AIS were included, ensuring complete

AIS capture for measurement (Fig 3B1). The distance from the AIS's initial point to the soma

in human pvalb neurons was 6.10 μm (IQR = 2.55–8.50, n = 13) compared with 4.85 μm (IQR

= 3.80–6.03, n = 14) in mouse neurons (p = 0.437; Fig 3B2). AIS length was greater in human

cells (17.60 μm, IQR = 14.65–20.20 μm) than in mouse cells (14.0 μm, IQR = 12.10–15.33; p

= 0.004), and the AIS terminal site was farther from the soma in human neurons (23.60 μm,

IQR = 20.85–25.75) than in mouse neurons (18.55 μm, IQR = 17.68–20.38; p < 0.001, Mann–

Whitney U test). All AIS properties are presented in Fig 3B2. None of the AIS parameters were

correlated with the AP threshold in either species. The correlation between AIS proximal

distance and AP threshold was 0.536 (p = 0.054) in human cells (n = 13) and −0.297 (p = 0.93)

in mouse cells (n = 14). AIS length was not significantly correlated with AP threshold in

humans (r = 0.203, p = 0.493) or mice (r = 0.079, p = 0.773), nor was distance from the AIS

termination site to the soma correlated with AP threshold in either species (humans: r = −0.239,

p = 0.414; mice: r = 0.024, p = 0.988, Spearman's correlation). However, when pooling data

across species, the AP threshold was significantly correlated with AIS start distance (r = −0.478, p = 0.012) and AIS termination distance (r = −0.383, p = 0.0485) but not AIS length (r = −0.1340, p = 0.516, Spearman's correlation; Fig 3C1–3).

These findings indicate that although AIS length and axonal location differ between human and mouse pvalb neurons, only the AIS position shows a consistent relationship with AP threshold when comparing across species. Specifically, greater distal AIS placement is associated with lower AP threshold.

Axon diameter does not account for interspecies differences in AP threshold

We examined axon diameter in biocytin-filled pvalb interneurons to assess its potential correlation with the AP firing threshold, particularly across species, as axon caliber can influence axon excitability [48]. Axonal diameters were measured using high-magnification fluorescence confocal microscopy images of biocytin-filled neurons at five standardized positions along the axon: two pre-AIS sites and three sites within the AIS (Fig 4A1–3). Reported data represent the mean values per cell for the pre-AIS and within-AIS portions of the axon. No significant differences were observed in axon diameter between human (n = 12) and mouse (n = 14) pvalb neurons in either in the pre-AIS [1.07 μm (IQR = 0.82–1.19) vs. 0.88 μm (IQR = 0.71–1.14); p = 0.939] or within-AIS portion [0.83 μm (IQR = 0.70–0.96) vs. 0.69 μm (IQR = 0.64–0.74); p = 0.260; one-way ANOVA on ranks; Fig 4B]. Additionally, in both AIS portions, axon diameter was not correlated with AP threshold in humans (pre-AIS: r = 0.245, p = 0.429; within-AIS: r = 0.392, p = 0.197; n = 12) or mice (pre-AIS: r = 0.091, p = 0.766; within-AIS: r = 0.056, p = 0.852; n = 12). Pooled data across species also showed no association (pre-AIS: r = 0.137, p = 0.518; within-AIS: r = −0.036, p = 0.866, Spearman's

323 correlation; Fig 4C1, 2). Collectively, these results indicate that axon diameter does not

324 contribute to the observed interspecies differences in AP threshold in pvalb neurons.

Fig 4. Axon diameter fails to explain interspecies differences in pvalb interneuron AP firing

threshold.

(A) Axon cross-diameter was measured in 3D confocal images of biocytin-filled pvalb interneurons.

(A1) Schematic showing axon maximum width at five points. "Pre-AIS" measurements were taken at

the axon's start (0 μm from the soma) and 2 μm from the soma. "AIS" measurements were taken at the

AIS start (shown as *0*) and at *2* and *5* μm from the AIS initiation site. (A2) Collapsed confocal image

showing biocytin-filled human pvalb neurons (Alexa 488), with the AIS visualized via beta-IV-spectrin

immunoreactivity (Cy3). (A3) Enlarged image of the axon highlighting five measurement sites. (B)

Axon thickness was similar between human (blue) and mouse (gray) pvalb neurons. Horizontal bars

indicate medians and quartiles. (C) Axon diameter was not correlated with AP threshold in either the

pre-AIS region (C1) or the AIS (C2). Data from both species were pooled (one-way ANOVA on ranks

with Dunn's pairwise *post hoc* test).

Nav1.6 channels are similarly distributed in the AIS of human and mouse pvalb neurons

We examined immunoreactivity for the low-voltage-gated sodium channel Nav1.6 in

the soma and axon of pvalb neurons (Fig 5A–H), as Nav1.6 density and spatial distribution in

the AIS can regulate AP threshold [49, 50]. Nav1.6 may concentrate at the AIS's distal end to

lower the AP firing threshold [48, 49, 51, 52], or it may be evenly distributed. In layer 2/3 of

the human and mouse neocortex, we assessed Nav1.6 immunofluorescence (Fig 5A1, 2)

intensity using confocal microscopy images of the soma (Fig 5B1, 2) and 3D image stacks

along the axon (Fig 5C1, 2) to evaluate species-specific differences in Nav1.6 channel

distribution. In individual human cells (n = 21; Fig 5D1) and mouse cells (n = 26; Fig 5D2),

24 radial lines were drawn from within the soma to the extracellular space to measure Nav1.6

(Cy3) and pvalb (Cy5) immunofluorescence. Because pvalb is intracellular, its onset along

each line defined the extracellular membrane site (Fig 5D1, 2). We also measured

immunofluorescence intensity for Nav1.6, pvalb, and beta-IV-spectrin in the same cells along

the axon, annotated in 3D image stacks, from the soma to beyond the AIS termination site (Fig

5E1, 2).

Fig 5. AIS displays robust and uniform immunofluorescence for Nav1.6 channels in human and

mouse pvalb neurons.

(A) Triple immunofluorescence for pvalb (Cy5), beta-IV-spectrin (Alexa 488), and Nav1.6 (Cy3)

revealed sodium channel localization in the AIS of human (A1) and mouse (A2) pvalb neurons. Arrows

mark AIS initiation and termination sites; s indicates the soma. Images are collapsed z-stacks from layer

2/3 of the neocortex. (B) Nav1.6 immunofluorescence in the somatic cytoplasm and membrane was

analyzed alongside pvalb in human (B1) and mouse (B2) cells. Immunofluorescence intensity was

measured using a 24-line radial pattern (yellow lines) extending from the extracellular space to the soma

center. (C) Nav1.6 intensity was measured along the axon from the soma (s) past the AIS termination

site in 3D confocal images. Measurement lines in human (C1) and mouse (C2) pvalb neurons overlie

beta-IV-spectrin–defined AIS (masked in the images by the lines). Images are collapsed from z-stacks.

(D) Immunofluorescence intensity [ordinate, arbitrary unit (a.u.)] was measured for pvalb (top, black

line) and Nav1.6 (bottom, red line) along radial lines spanning the extracellular space (negative abscissa

values) to the soma (positive values). Vertical dashed line denotes the membrane location (0-point),

defined as ± 0.5 µm from the first pvalb signal. (D1) Left: Mean fluorescence intensity (F, a.u.) across

24 radial lines in one human cell, spanning -3 to $+1.5$ µm. Schematic illustrates measurement geometry.

Right: Medians and quartiles of intensity values minus the extracellular reference ($F-F_{ex.c}$) in 21 human

cells. (D2) Identical measurements as in *D1* for one mouse cell (left) and 26 mouse cells (medians and

quartiles; right). (E) Fluorescence intensity along 3D pixel lines for pvalb (black), beta-IV-spectrin

(green), and Nav1.6 (red). (E1) Left: Measurement in one human neuron (a.u.); 0 on abscissa = axon

origin. Upper schematic illustrates the measurement line along the axon. Right: Medians and quartiles

of referenced intensity values ($F-F_{ex.c}$) in 21 human cells. Upper schematic indicates the defined AIS

(determined via beta-IV-spectrin immunofluorescence) and pre-AIS and post-AIS axonal regions. AIS

length (green background) was divided into 10 bins per cell. Statistical analysis showed no difference

in Nav1.6 intensity between proximal and distal AIS halves (Mann–Whitney U test). (E2) Same

measurements shown in *E1* for one mouse cell (left) and 26 mouse cells (right; medians and quartiles).

(F, G) Summary of line analyses in human and mouse pvalb neurons; medians and quartiles of $F-F_{ex.c}$

across five cellular domains showed strong Nav1.6 immunoreactivity in the AIS. (H) $F-F_{ex.c}$ values

across domains did not differ between human (blue) and mouse (gray) pvalb cells (PERMANOVA with

Bonferroni *post hoc* test).

Reference-corrected Nav1.6 immunofluorescence (F–$F_{ex.c.}$, defined as the fluorescence intensity with the extracellular average subtracted from other radial or axon pixels) was similar across species in the soma membrane (humans: 7.06, IQR 4.72–15.12, n =21; mice: 8.52, IQR = 3.30–12.50, n = 26; p = 1.00) and the cytoplasm (humans: 23.66, IQR = 16.37–31.64, n = 21; mice: 12.39, IQR = 5.88–20.25, n = 26; p = 0.067).

Nav1.6 immunofluorescence was similarly weak in human and mouse pre-AIS axons (humans: 6.55, IQR = 4.03–20.69, n =21; mice: 6.25, IQR = −0.87–16.24, n =26; p = 1.00) and post-AIS regions (humans: 0.95, IQR = −3.41 to 7.56; mice: 4.59, IQR = −2.49 to 13.13; p = 1.00). Nav1.6 intensity was strong in the AIS but not significantly different between humans (73.18, IQR = 49.32–94.05) and mice (48.40, IQR = 28.84–67.81; p = 0.063, PERMANOVA and Mann–Whitney U test with Bonferroni correction; Fig 5E1, 2). In both species, Nav1.6 immunofluorescence was uniformly distributed within the AIS, with no difference between the first and second halves (humans: p = 0.182; mice: p = 0.833, Mann–Whitney U test; Fig 5E1, 2). Nav1.6 F–$F_{ex.c}$ values are shown in Fig 5F for humans and Fig 5G for mice, with is no species differences observed (Fig 5H).

Human pvalb neurons exhibit low Kv1 channel levels in the AIS relative to mouse neurons

We investigated the immunofluorescence of Kv1.1 and Kv1.2 potassium channels, key inhibitory mechanisms modulating AP threshold in the AIS [53, 54]. Kv1 channels are clustered in the AIS and soma of rodent pvalb neurons [55], where they increase the AP firing threshold [53, 56] (hereafter, Kv1 channels are referred to as inhibitory channels). First, we analyzed Kv1.1 immunofluorescence in the soma and AIS of human (Fig 6A1) and mouse (Fig 6A2) pvalb neurons using triple immunolabeling. Representative cells and analysis are shown

in Fig 6B and C, including those in the soma (Fig 6B1, 2) and along the axon (Fig 6C1, 2). F–$F_{ex.c.}$ analysis showed absent Kv1.1 signal at the human soma membrane (−1.23, IQR = −4.77 to 5.25, n = 31) as well as its presence in mice (5.42, IQR = 0.62–10.58, n = 31; p = 0.041). Kv1.1 signal in the cytoplasm was weak in both species and did not differ significantly (p = 0.817; Fig 6D1, 2).

Fig 6. AIS is deficient in Kv1.1 potassium channels in humans compared with mice.

(A) Immunofluorescence staining revealed a species-specific difference in Kv1.1 expression in the AIS of pvalb neurons. (A1) Triple labeling for beta-IV-spectrin (Alexa 488), Kv1.1 (Cy3), and pvalb (Cy5) in the human neocortex layer 2/3. Open arrows denote the AIS visualized using beta-IV-spectrin in a pvalb+ neuron; solid arrows mark the AIS of a pvalb− neuron; s denotes the pvalb+ soma. The pvalb+ neuron lacked a Kv1.1 signal, whereas the pvalb− AIS exhibited robust Kv1.1 labeling. (A2) Same triple labeling shown in *A1* in the mouse cortex; both pvalb+ (open arrows) and pvalb− (solid arrows) neurons showed robust AIS Kv1.1 expression. (B) Kv1.1 fluorescence intensity in somatic membranes analyzed with pvalb labeling in human (B1) and mouse (B2) neurons. Yellow radial lines indicate the 24-line measurement pattern. (C) Axonal Kv1.1 intensity was measured from the soma (s) in 3D confocal images in human (C1) and mouse (C2) neurons. Measurement lines are shown above the beta-IV-spectrin–labeled AIS in collapsed z-stacks. (D) Fluorescence intensity for pvalb (top, black line) and Kv1.1 (bottom, red line) along radial lines in the soma. Vertical dashed line marks pvalb signal onset (0-point). (D1) Left: Mean intensity (F, a.u.) across 24 radial lines in one human cell, spanning −3 to +1.5 μm. Schematic shows radial lines. Right: Medians and quartiles of F–$F_{ex.c}$ in 31 human cells for pvalb (gray) and Kv1.1. (red). (D2) Same measurements as in *D1* for one mouse cell (left) and 31 mouse cells (right; medians and quartiles). (E) Line analysis of axons for pvalb (black), beta-IV-spectrin (green), and Kv1.1 (red). (E1) Left: Intensity profile of one human cell, as illustrated in upper schematic (0 = the axon origin). Right: Medians and quartiles of F–$F_{ex.}$ intensity values in 31 human cells. Upper schematic shows AIS defined by beta-IV-spectrin labeling. AIS length (green background) was divided into 10 bins per cell. Statistical analysis revealed that Kv1.1 intensity does not differ between the

proximal and distal AIS halves (Mann-Whitney U test). (E2) Identical measurements described in *E1*

for one mouse neuron (left) and 31 mouse neurons (right; medians and quartiles). Statistical analysis

indicated that Kv1.1. is evenly distributed along the AIS (Mann-Whitney U test). (F, G) Summary of

line analyses in human and mouse pvalb neurons; medians and quartiles of $F-F_{ex.c}$ for Kv1.1 across

cellular domains. (H) $F-F_{ex.c}$ values in human (blue) and mouse (gray) pvalb cells across cellular

domains: mouse cells showed stronger Kv1.1 signals in the somatic membrane and AIS compared with

human cells (PERMANOVA test with Bonferroni *post hoc* test).

Axonal analysis showed robust Kv1.1 signal in the AIS of mouse cells (47.56, IQR = 13.98–

74.95, n = 31), consistent with previous reports [53]. In contrast, human AIS Kv1.1 signal was

low or absent (8.54, IQR 5.00–24.43), despite strong signal in nearby pvalb-negative neurons

(Fig 6A1). AIS $F–F_{ex.c.}$ differed between species (p = 0.001; Fig 6E1, 2). Pre-AIS (humans:

−1.06, IQR = −6.71 to 8.77; mice: 9.56, IQR = −5.49 to 20.32; p = 0.287) and post-AIS

(humans: 2.55, IQR = −3.26 to 10.88; mice: 4.58, IQR = −17.97 to 29.20; p = 1.00) regions

showed no significant differences (PERMANOVA and Mann–Whitney U test with Bonferroni

correction.). Kv1.1 intensity showed no difference between AIS halves (humans: p = 0.318;

mice: p = 0.195, Mann–Whitney U test; Fig 6E1, 2). Kv1.1 $F–F_{ex.c.}$ intensity values for each

domain are summarized in Fig 6F and G, and species comparisons are shown in Fig 6H.

We also analyzed Kv1.2 immunofluorescence (Fig 7A1, 2), comparing $F–F_{ex.c.}$ in the soma

(Fig 7B1, 2) and AIS (Fig 7C1, 2) of human and mouse pvalb neurons [57]. A low but

moderately higher Kv1.2 signal was observed at the human soma membrane (5.74, IQR = 2.08–

10.42, n = 60) compared with the mouse membrane (3.42, IQR = −2.66 to 5.30, n = 33; p =

0.013; Fig 7D1, 2). Cytoplasmic signal was also higher in humans (4.47, IQR = −3.73 to 16.39)

than in mice (−6.59, IQR = −12.07 to 4.16; p = 0.002).

Fig 7. Kv1.2 potassium channel expression is low in the AIS of human and mouse pvalb neurons.

(A) Triple immunofluorescence for beta-IV-spectrin (Alexa 488), Kv1.2 (Cy3), and pvalb (Cy5)

revealed weak Kv1.2 immunoreactivity in the AIS of human (A1) and mouse (A2) pvalb neurons. Open

arrows indicate AIS identified via beta-IV-spectrin in a pvalb+ neurons, which show negligible Kv1.2

signal; solid arrow denotes a Kv1.2-positive dendrite and somatic membrane of another neuron; s marks

the pvalb+ soma. (B) Radial intensity measurements of Kv1.2 and pvalb in human (B1) and mouse (B2)

somata using 24-line radial patterns (yellow lines). (C) Line measurements of Kv1.2 intensity along the

axon from the soma (s) in human (C1) and mouse (C2) pvalb neurons. Images are collapsed z-stacks;

line measurement is shown over the axon. (D) Immunofluorescence intensity profiles for pvalb (top,

black line) and Kv1.2 (bottom, red line) along the radial lines in the soma. (D1) Left: Mean fluorescence

intensity (F, a.u.) in 24 pooled lines from one human cell, spanning −3 to +1.5 μm from the 0-point.

Right: Medians and quartiles of $F-F_{ex.c}$ values for pvalb (gray) and Kv1.2. (red) in 60 human cells.

(D2) Identical measurements as in *D1* for one mouse cell (left) and 33 mouse cells (right; medians and

quartiles). (E) Axonal line analysis for pvalb (black), beta-IV-spectrin (green), and Kv1.2 (red). E1)

Left: Line intensity values along the axon in one human cell (illustrated in the upper schematic), with 0

denoting the axon origin. Right: Median and quartiles of $F-F_{ex.}$ values across 60 human cells. AIS

length (green background) was divided into 10 bins. Statistical analysis showed similarly negligible

Kv1.2 intensity in the proximal and distal AIS halves (Mann-Whitney U test). (E2) Identical

measurement as in *E1* for one mouse neuron (left) and 33 mouse neurons (right; medians and quartiles),

showing uniformly low Kv1.2 signal across the AIS (Mann-Whitney U test). (F, G) Summary of line

analyses in human and mouse pvalb cells; medians and quartiles of $F-F_{ex.c}$ values across cellular

domains. (H) $F-F_{ex.c}$ values in human (blue) and mouse (gray) pvalb cells: human cells showed stronger

Kv1.2 signals in the somatic membrane compared with mouse cells (PERMANOVA test with

Bonferroni *post hoc* test).

In the AIS, no difference in Kv1.2 $F–F_{ex.c.}$ was detected between human (2.95, IQR = −3.23 to 9.05, n = 60) and mouse (17.04, IQR = −5.30 to 31.76, n = 30) cells (p = 0.272; Fig 7E1, 2), nor in pre-AIS (p = 0.478) or post-AIS (p = 0.187) sites (PERMANOVA and Mann–Whitney U test with Bonferroni correction). Kv1.2 immunofluorescence intensity showed no difference between AIS halves (humans: p = 0.397; mice: p = 0.984, Mann–Whitney U test; Fig 7E1, 2). Kv1.2 data are summarized in Fig 7F, G and H.

Pvalb neurons in humans lack *KCNA1* mRNA encoding Kv1.1 but possess *KCNA2* mRNA encoding Kv1.2

We used patch-sequencing to measure the mRNA levels of low-threshold ion channels that regulate the AP firing threshold, namely Kv1.1 and Kv1.2 [44, 51, 53-55, 58], in electrophysiologically studied fast-spiking pvalb neurons in human and mouse neocortex layer 2/3. Whole-cell recordings were followed by nucleus extraction and transcriptomic analysis [59, 60]. We detected mRNA for 6,784 genes (IQR = 4872–7399) in humans (n = 21) and 3,617 genes (IQR = 3173–4305) in mice (n = 16; Fig 8A). Transcriptomes were uploaded to the Allen Institute neuron type identification server, which employs an algorithm categorizing neocortex layer 2/3 neuron types by gene expression (https://knowledge.brain-map.org/mapmycells/process/), enabling identification of pvalb neurons, even in cells with undetectable *PVALB* mRNA expression. Using this method, we identified 21 and 16 human and mouse pvalb neurons, respectively. These cells were analyzed for expression of *KCNA1*, *KCNA2*, *KCNC1*, *KCNC2*, *KCND1*, *KCNQ1*, *KCNQ2*, *KCNQ3*, *KCNQ4*, *KCNQ5*, *KCNK2*, *KCNK4*, and *KCK10*, i.e., genes encoding major potassium channels in axons that regulate AP threshold [54, 55]. Additionally, we analyzed *SCN8A* expression and that of the housekeeping genes *ACTB*, *GAPDH*, and *SLC8A1* (Fig 8A).

Fig. 8. Human pvalb neurons lack *KCNA1* mRNA and express low *KCNA2* mRNA levels.

(A) Patch-sequencing analysis (illustrated in the upper schematic) revealed that human pvalb neurons

do not express *KCNA1*, in contrast to its expression in mouse cells. (A1) Heatmap showing transcript

levels (TPM) of genes encoding low-threshold potassium channels in individual human (n = 21) and

mouse (n = 16) pvalb neurons, as well as the housekeeping genes *GAPDH* and *ACTB*. *KCNA1*

expression differed quantitatively (TPM values) and categorically (positive or negative gene

expression) in human and mouse cells. Expression of *KCNA2*, *KCNC1*, *KCNC2*, *KCNK2*, *KCNK4*, and

KCNK10 was observed in both species (*61*), whereas *KCND1* expression was nearly absent in both

(*62*). Human and mouse cells expressed similar levels of *KCNQ1*, *KCNQ2*, and *KCQN4*, whereas

KCNQ3 and *KCNQ5* expression levels were higher in humans. Additionally, *SCN8A* (encoding Nav1.6

channels) and *SLC6A1* (encoding GABA transporter type-1) were highly expressed in both species.

Significant differences in TPM values are marked in red (genes, ordinate). Bottom of the heatmap shows

the pvalb cell subtype classification based on Allen Institute transcriptomic data. (B) Allen Institute

database analysis of layer 2/3 pvalb neuron subtypes in the human and mouse neocortex confirmed

KCNA1 mRNA deficiency in human cells. Heatmap shows average TPM values (scale on the right)

across seven human and eight mouse pvalb neuron subtypes (abscissa). *KCNA1* mRNA was absent in

all human subtypes but robustly expressed in mouse cells. Statistical analyses (one-way MANOVA

with Bonferroni *post hoc* test and chi-squared test with Benjamini–Hochberg adjustment) confirmed

significant differences in both expression level and frequency (p = 0.0011 and p < 0.001, respectively).

KCNQ3 and *KCNQ5* expression levels were higher in humans. Significant differences in TPM values

are marked in red (genes, ordinate). Abscissa shows pvalb cell subtype names.

All 21 human pvalb neurons lacked detectable *KCNA1* mRNA expression, whereas 7 of 16

mouse neurons showed robust expression. The mRNA level heatmap in Fig 8A shows the gene

expression (transcripts per million [(TPM]) of individual pvalb neurons in humans and mice.

Comparing *KCNA1* mRNA expression between human (n = 21) and mouse cells (n = 16)

revealed a significant difference (p = 0.0016, one-way MANOVA with a *post hoc* Bonferroni

test). Furthermore, categorical comparisons (positive or negative mRNA expression)

confirmed that *KCNA1* expression differed significantly between species (p < 0.001, chi-

squared test following Benjamini–Hochberg adjustment at the 5% significance level; Fig 8A).

Both species displayed *KCNA2* mRNA expression without any species-specific difference (p

= 0.970). We also tested whether pooled mRNA levels (TPM) of *KCNA1* and *KCNA2* were

associated with AP threshold in patch-sequenced neurons. No association was observed for

either gene in humans (r = 0.041, n = 20, p = 0.861) or mice (r = 0.236, n = 15, p = 0.388).

When neurons from both species were pooled, the results remained nonsignificant (r = 0.175,

n = 35; p = 0.312, Spearman's correlation test).

Both species exhibited similarly strong *KCNC1* and *KCNC2* mRNA expression, which

respectively encode the Kv.3.1 and Kv3.2 channels characteristics of pvalb interneurons [43,

61]. They also showed equally low or undetectable expression of *KCND1* encoding Kv4.1

potassium channels [62]. No species differences were observed in *KCNK2*, *KCNK4*, or

KCNK10 mRNA levels, and *KCNQ1*, *KCNQ2*, and *KCNQ4* mRNAs were similarly expressed

between species. However, human neurons displayed significantly higher mRNA levels of

KCNQ3 (p < 0.001) and *KCNQ5* (p < 0.001). Both human and mouse pvalb cells strongly

expressed *SCN8A* mRNA, encoding Nav1.6 sodium channels. Additionally, all cells expressed

high levels of *ACTB* and *GAPDH*. Notably, mRNA expression of *SLC6A1*, which encodes

GABA transporter 1, was significantly lower in human neurons than in mouse neurons (p =

0.002). *PVALB* mRNA expression was undetectable in 10 of 21 human neurons and 6 of 16

mouse neurons, aligning with results from the Allen Institute open access database, which

reports markedly low *PVALB* mRNA expression levels in some layer 2/3 pvalb cells (Fig 8B).

The patch-sequencing results closely aligned with the average mRNA profiles of pvalb neuron

types in the Allen Institute database (Fig 8B). The database confirmed the absence of *KCNA1*

expression in humans but not in mice (p < 0.001), as well as expression of *KCNA2* in both

species. It also confirmed that among the potassium channel genes analyzed, only Kv1 channel

genes were consistently expressed at lower levels in human pvalb neurons relative to mouse

neurons (one-way MANOVA with a *post hoc* Bonferroni test).

Kv1 channel blocker dendrotoxin lowers the AP firing threshold more prominently in mouse neurons than in human neurons

We investigated whether Kv1 channel blockade through dendrotoxin-K (DTXK)

affects AP generation differently in human and mouse pvalb neurons [53, 58]. After 5 min of

baseline whole-cell current–clamp recording (conditions similar to those shown in Fig 1), slices

were exposed to 100 nM DTXK for 10 min, and AP threshold was remeasured (Fig 9A1, 2).

DTXK significantly lowered the AP firing threshold in human neurons from −38.8 mV (IQR

= −42.3 to −32.2 mV) to −40.0 mV (IQR = −45.3 to −37.2 mV; n = 10 cells; p = 0.004;

Wilcoxon signed-rank test with Bonferroni correction). In mouse cells, the reduction was from

−31.9 mV (IQR = −35.9 to −29.5 mV) to −40.7 mV (IQR = −45.8 to −38.2 mV; n = 10 cells;

p = 0.004; Wilcoxon signed-rank test with Bonferroni correction), and more prominent

compared to human cells (p = 0.019; Mann–Whitney U test with Bonferroni correction).

Fig 9. Kv1 channel blocker dendrotoxin lowers AP threshold more prominently in mouse versus human pvalb neurons.

APs were recorded under baseline conditions (black traces) in current–clamp mode at −70 mV (as shown in Fig 1) for 5 min, followed by application of 100 nM DTXK for 10 min, after which APs were evoked again (red traces). (A) APs and corresponding AP phase plots in baseline control and after DTXK application. (A1) Left: Three superimposed APs from a human pvalb cell. Right: Corresponding phase plots under control and DTXK conditions. (A2) Left: Superimposed APs recorded in a mouse pvalb neuron. Right: Corresponding AP phase plots. (B–E) Plots show average changes in AP parameters measured at the AP threshold in human and mouse neurons (n = 10 cells each). Values represent the difference between DTXK and baseline conditions. (B) DTXK induced a larger negative shift in the AP firing threshold in mouse cells compared with human cells. (C) Reduction in rheobase current following DTXK application was greater in mouse than in human cells. (D) DTXK caused a more pronounced reduction in AP AHP in mouse compared with human cells. (E) No significant change was observed in the AP maximum fall speed between species.

In both species the AP maximum fall speed was reduced (humans: p = 0.049; mice p = 0.020), whereas in mice but not humans, the rheobase current decreased (human: p = 0.375; mice: p =

595   0.002) and the AHP amplitude declined (humans: p = 0.064; mice: p = 0.002; Wilcoxon signed-

596   rank test with Bonferroni correction). Other AP properties, Rin, and tau were unaffected. Full

597   results are provided in Table 2.

598   **Table 2. AP parameters, Rin, and tau in pvalb neurons under baseline and DTXK exposure**

599   **conditions.**

	Human n = 10			Mouse n = 10		
	Baseline median, (IQR)	DTX, median, (IQR)	P-value (baseline vs. DTX)	Baseline median, (IQR)	DTX, median, (IQR)	P-value (baseline vs. DTX)
AP Thr (mV)	−38.8 (−42.3 to −32.2)	−40.0 (−45.3 to −37.2)	**0.004**	−31.9 (−35.9 to −29.5)	−40.7 (−45.8 to −38.2)	**0.004**
Initial rise slope (ms^{-1})	39.1 (33.0 to 44.5)	41.0 (30.2 to 57.9)	0.492	45.9 (38.3 to 57.5)	49.6 (42.1 to 79.8)	0.275
AP ampl (peak value − SpkThr, mV)	56.4 (46.3 to 61.4)	52.6 (42.9 to 59.8)	0.432	57.6 (44.4 to 62.7)	62.8 (33.5 to 70.0)	0.922
AP half width (ms)	0.54 (0.43 to 0.77)	0.55 (0.46 to 0.87)	0.426	0.35 (0.30 to 0.60)	0.36 (0.32 to 0.63)	0.770
AP rheobase (pA)	83.4 (37.1 to 113.1)	77.6 (45.9 to 104.8)	0.375	280.3 (135.5 to 443.6)	121.9 (94.0 to 283.7)	**0.002**
AP latency (ms)	23.3 (20.5 to 32.1)	26.6 (17.0 to 52.7)	0.922	71.8 (37.1 to 124.0)	27.8 (12.4 to 86.6)	0.160
Spike number by step	2.6 (1.0 to 6.9)	2.1 (1.8 to 5.1)	0.742	4.2 (2.7 to 10.0)	5.0 (2.3 to 9.5)	0.770
AP AHP ampl (mV)	−16.6 (−12.7 to −18.6)	−13.6 (−10.4 to −17.2)	0.064	−18.4 (−15.2 to −21.1)	−13.4 (−8.2 to −14.8)	**0.002**
Max rise speed (mV/ms)	232.8 (151.7 to 328.9)	226.8 (118.8 to 342.7)	0.375	331.2 (233.7 to 373.2)	342.4 (132.8 to 419.1)	0.770
Max fall speed (mV/ms)	−140.6 (−168.5 to −97.9	−127.6 (−154.5 to −53.6)	**0.049**	−162.4 (−229.7 to −108.0)	−98.2 (−234.1 to −61.0)	**0.020**
Rin (MΩ)	226.5 (94.8 to 455.1)	244.7 (116.1 to 402.3)	0.432	90.3 (82.4 to 199.8)	96.5 (76.6 to 173.0)	0.492
Tau (ms)	6.8 (4.5 to 8.4)	7.0 (5.1 to 8.0)	0.770	4.7 (3.6 to 6.9)	4.7 (4.1 to 5.9)	0.695

Rin: cell input resistance; tau: membrane time constant, measured as the voltage response to a +10-mV step from −70 mV. Data are presented as medians with quartiles. P-values indicate differences within each species before and after DTXK exposure (Wilcoxon signed-rank test with Bonferroni correction).

Species comparisons of DTXK-induced changes in significantly affected AP parameters (Fig 9B–E) revealed that threshold shift ($p = 0.019$), rheobase reduction ($p < 0.001$), and AHP amplitude decrease ($p = 0.028$) were all greater in mouse cells compared with human cells, whereas the reduction in AP maximum fall speed was comparable between species ($p = 0.821$; Mann–Whitney U test with Bonferroni correction).

Fast-spiking neuron computational model with a human-type AIS reveals a lowered firing threshold and shortened time lag to AP generation

We established a single-cell computational model with axonal and dendritic cable properties and an AIS replicating the electrical characteristics of human pvalb neurons. We simulated the effects of species-specific AIS differences, namely AIS length and Kv1 channel content, on AP firing threshold and the time lag between somatic excitation and AP initiation. The model's intrinsic properties were tuned to match the Vm waveforms and AP firing threshold observed in human pvalb cells during voltage step protocols (Fig 10A1, 2). Specifically, the model reproduced a whole-cell recorded human pvalb cell (code h7, Table S1) with Hodgin–Huxley-type APs and incorporated voltage-sensitive low-threshold Na^+ current (Nav1.6 in the AIS), high-threshold Na^+ current (Nav1.2 in the soma and proximal dendrites), and K^+ currents (Kv3.1 and Kv3.2 in the AIS, soma, and dendrites). The soma and dendrites

623 also included voltage-insensitive K^+ leak current, M-type and Kir-type K^+ currents [63], and

624 hyperpolarization-activated nonspecific cation Ih current [28]. Variable Kv1 activity

625 (conductance) was applied in the AIS. Under standard conditions, the AIS length was 20 μm,

626 initiation distance from the soma was 10 μm, and the AP firing threshold was −43 mV.

Fig 10. Fast-spiking neuron computational model with human or mouse AIS.

(A) Single-cell computational model with axon and dendrite cable properties replicating the human

pvalb cell's somatic electrical behavior. (A1) Vm responses in a real human pvalb neuron (blue) and

the model cell (gray) to current steps from −200 to +200 pA. (A2) Left: First AP elicited by the lowest

suprathreshold step in the real neuron (blue) and model cell (gray). Right: Corresponding AP phase

plots. (B) Increasing AIS length from 10 to 30 μm, within the observed range in real pvalb cells, lowered

the AP firing threshold in the model (Kv1 and Nav1.6 channel conductance = 20 nS/μm², AIS-to-soma

distance = 10 μm, axon Ra = 0.9 MΩ·μm, axon diameter = 0.8 μm). (B1) Model APs and phase plots

with AIS lengths of 10 μm (purple) and 30 μm (red). (B2) AP firing threshold simulated for AIS lengths

of 10 μm (purple), 20 μm (brown), and 30 μm (red) (illustrated in schematic inset). Threshold sensitivity

was highest at 10–20 μm, a range common to both human and mouse pvalb neurons in reality. (C)

Reducing Kv1 channel activity in the AIS lowered the AP firing threshold in the model. (C1) Model

APs and phase plots at different Kv1 conductances in the AIS: gKv1 = 0 (yellow) and 15 nS/μm² (red).

(C2) AP firing threshold at Kv1 conductance levels of 0, 5, 10, and 15 nS/μm² (abscissa; Nav1.6

conductance = 20 nS/μm², AIS length = 20 μm, AIS-to-soma distance = 10 μm). (D) Combined AIS

elongation and reduced Kv1 activity synergistically lowered the AP threshold in the model. Heatmap

illustrates the combined effect of AIS length (abscissa) and Kv1 conductance (ordinate) on the AP firing

threshold (color scale, right). Total AP threshold range across conditions was 11 mV. (E–G) Human-

type AIS shortened the AP generation lag. (E) Real human and mouse neurons differed in depolarization

kinetics and AP thresholds but showed similar AP generation lags. Human pvalb neuron (blue) had a

slower membrane depolarization (E1) and lower AP threshold (E2) relative to the mouse pvalb neuron

(gray). (E3) AP generation lag, defined from the onset of a depolarizing step (black dotted lines) to AP

initiation (gray dotted lines) in four representative traces (AP onset defined at 10-mV/ms rise speed).

(E4) AP generation lag versus excitatory current steps in the soma of human (blue) and mouse (gray)

neurons. Human neuron fired with a lower current (abscissa; vertical dotted lines) but showed a similar

time lag (horizontal dotted line) relative to the mouse neuron. (F) Switching AIS properties in the model

from mouse-like to human-like reduced the AP generation lag to somatic excitation. (F1) The human-

type AIS lowered the AP threshold in response to simulated EPSCs in the model neuron soma. Vm

traces showed excitatory postsynaptic potentials (EPSP) of increasing strength that eventually triggered

an AP in model neurons with human (blue) and mouse (gray) AIS. EPSP-generating conductances with

peak amplitudes of 6–nS are presented on the left (black traces). Scale bars: 40 ms and 50 mV. (F2)

Human-type AIS shortened the AP generation lag in response to EPSPs. Vm traces showed APs

triggered by equal-strength excitatory postsynaptic currents (EPSC) (8.5-nS peak conductance) in

model neurons with human-type (blue) or mouse-type (gray) AIS. (F3) AP generation lag in the model

cell plotted against increasing EPSC conductance in model neurons with human-type (blue dots) and

mouse-type (gray dots) AIS. Response to the 8.5-nS EPSC is highlighted in red. (G) Heatmap showing

the combined effect of AIS length (abscissa) and AIS Kv1 conductance (ordinate) on the AP generation

time lag in response to a fixed-strength EPSC (8.5 nS) in the model neuron. Color scale (right) indicates

time lag in milliseconds.

First, we simulated the AP firing thresholds by varying AIS lengths (10–30 μm), within the

anatomical range observed in real pvalb neurons. Increased AIS length linearly scaled Nav1.6

and Kv1 channel activity in the AIS (conductances: gNav1.6 and gKv.1 = 20 nS/μm²). The

axon diameter was 0.8 μm, with a longitudinal resistance (Ra) of 0.9 MΩ·μm. Increasing AIS

length from 10 to 20 and 30 μm lowered the AP firing threshold (elicited by a 250-ms square-

pulse current at the soma of −70 mV) from −36.8 to −42.3 and −45.2 mV, respectively (Fig

10B1, 2), demonstrating that AIS length (with fixed gNav1.6 and gKv.1 per μm²) strongly

modulates AP threshold.

Next, we examined how varying Kv1 conductance in the AIS affects AP threshold while fixing

the AIS length (20 μm), location along the axon (10 μm from the soma, Ra = 0.9 MΩ·μm),

and gNav1.6 density (20 nS/μm^2). Depolarizing steps were applied to elicit APs. Increasing

gKv1 from 0 to 5, 10, and 15 nS/μm^2 in the AIS (Fig 10C1, 2) raised the AP threshold from

−45.3 to −44.2, −42.4, and −41.6 mV, respectively (Fig 10D). This simulation confirmed that

Kv1 conductance in the AIS increases firing threshold independently of AIS geometry. We

also simulated increased Nav1.6 conductance in the AIS, and found that it effectively reduced

AP threshold in the model cell (Fig S1A).

We then ran simulations incorporating cell-to-cell variability parameters shared across species.

With fixed AIS length (20 μm) and channel conductances (gNav1.6 and gKv1 = 20 nS/μm^2),

we varied the AIS initiation site distance (0 or 10 μm from the soma). AP threshold was lower

when AIS was 10 μm from the soma (−43.2 mV) compared with 0 μm (−41.5 mV; Fig S1B,

C). We also simulated the impact of axon diameter variation (0.6–0.9 μm), which alters both

the Ra (0.8–1.1 MΩ·μm) and extracellular membrane surface area (calculated as diameter × π,

with gNav1.6 and gKv1 = 20 nS/μm^2). Narrower axons (0.6 μm) with higher Ra (1.1 MΩ·μm)

yielded the lowest AP threshold (−44.7 mV) in the model cell, whereas wider axons (0.9 μm)

with lower Ra (0.8 MΩ·μm) had the highest threshold (−42.2 mV). Varying Ra alone (with a

fixed axon surface area) shifted the threshold by <1 mV; changing surface area alone (while

keeping Ra constant), thereby increasing total ion channel conductance, altered the threshold

by ~2 mV. Notably, when axon diameter was increased over the range observed in the

anatomical study, thereby lowering Ra while increasing total Nav1.6 and Kv1 channel

conductance, the AP firing threshold moderately increased only by ~2.5 mV. These

relationships are detailed in the heatmap shown in Fig S1A–C.

Finally, we used the model cell to explore how AIS-dependent changes in firing threshold influence the time lag between EPSC onset AP generation in pvalb neurons. The simulation aimed to address why human pvalb neurons, compared with mouse pvalb neurons, despite having a slower somatic membrane tau (Fig 10E1) and a lower AP firing threshold (Fig 10E2), can generate APs with an equal or shorter time lag in response to a fixed-amplitude excitatory current in the soma (Fig 10E3, 4) [38]. In the model neuron, we generated an EPSC in the soma at −70 mV and incrementally increased its peak amplitude to evoke a gradually increasing EPSP that eventually reached the AP firing threshold [28]. The EPSC was applied with peak conductance values from 6 to 11 nS in steps of 0.25 nS, reaching the AP firing threshold with the human- and mouse-type AIS configurations at EPSC conductances of 6.75 and 8.50 nS, respectively (Fig 10F1–3). Shifting from the "mouse-type AIS" (Kv1 density = 20 nS/μm^2, AIS length = 10 μm) to the "human-type AIS" (Kv1 density = 0 nS/μm^2, AIS length = 30 μm) resulted in a shorter time lag to AP generation when the EPSC strength was fixed (Fig 10F2, 3). The EPSP–AP time lag simulation with varying AIS lengths and Kv1 densities is shown in a heatmap in Fig 10G. The simulation revealed a range of AP generation time lags from 4.8 and 7.8 ms in response to a fixed EPSC strength (8.5 nS, corresponding to the AP threshold in the mouse-type AIS). Overall, these simulations demonstrate that the lower AP firing threshold associated with the human-type AIS compared to the mouse-type AIS in pvalb cells is accompanied by a shorter time lag to AP generation in response to synaptic current at the soma.

Discussion

Species-specific traits in organs, tissues, and cells have evolved in response to selection pressure during phylogenetic evolution [8, 64, 65]. Consequently, homologous cell types (i.e., neurons expressing the same cell-type specific marker genes) are broadly conserved across mammalian species but still exhibit interspecies differences that reflect adaptations for

functional advantages [4, 66-69]. Our findings suggest that fast-spiking pvalb neurons in the

human neocortex evolved a lower AP firing threshold to support their role as fast in–fast out

circuits. This enhanced input-to-output signal transformation speed is especially important in

human neurons, which have electrically slower somata compared with rodent neurons.

Molecular, anatomical, and pharmacological data, together with model simulations, showed

that the reduced AP threshold in human pvalb neurons results from diminished Kv1 inhibitory

potassium channel content in the AIS and an elongated AIS.

Low AP firing threshold helps electrically slower human interneurons to function as fast in–fast out circuits

A short delay in AP generation in response to excitatory synaptic input (i.e., input–

output transformation) is a key feature of fast-spiking pvalb interneurons in the cortex across

mammals [40, 43]. These neurons, the most numerous inhibitory population in the cortex [6,

70, 71], exert fast, temporally precise inhibition on various other neuron types [43]. In the

human neocortex, rapid signal transformation poses a challenge for many pvalb neurons owing

to their slower somatic membrane potential kinetics. However, a lowered AP threshold

compensates for this, reducing the time lag in input–output transformation. This helps preserve

the archetypical fast in–fast out function of pvalb neurons in humans, as demonstrated through

our computational model.

Slowed inhibitory signaling in the human neocortex would be detrimental, as theoretical

models and *in vivo* animal studies have linked such changes to reduced cortical computation

speed and cognitive impairment [40, 72-75]. Therefore, human fast-spiking interneurons have

evolved distinct adaptations that maintain rapid electrical input–output transformations. These

include simplified dendritic morphology that facilitates synaptic potential propagation from

distal dendrites to the soma [2, 31]. Moreover, human pvalb neurons express more voltage-

sensitive excitatory cation channels in the soma and proximal dendrites to accelerate membrane

potential dynamics [28]. We also observed an adaptation in AP generation at the soma. Some

of these features likely generalize to primate pvalb neurons, as macaque neocortical neurons

also exhibit slower Vm kinetics and reduced AP thresholds compared with rat neurons [36].

Why do human pvalb neurons have electrically slower somata relative to rodent neurons?

The slower soma membrane dynamics in human pvalb neurons result from increased

cell Rin, which reflects reduced transmembrane ion leakage. This is supported by findings

showing that cell Cm, a parameter that combines with Rin to determine membrane tau, is

similar between primate and rodent fast-spiking neurons [36, 37]. In mice, somatic tau

correlates with Rin, but in humans, this link is weak, indicating high variability in membrane

ion channel density between individual human neurons (hence, in electrical resistivity per

plasma membrane area), which governs ion leakage [76].

The reason high Rin evolved in human pvalb neurons remains unclear, as it is inherently

unfavorable for fast in–fast out function. One hypothesis is that evolutionary pressure favored

neurons that produce Vm changes with reduced electric current demands, minimizing

transmembrane ion flux to conserve metabolic energy [77, 78]. Fast-spiking interneurons are

energetically demanding [79-81], and their metabolic cost may be heightened in the human

brain due to the higher proportion of interneurons in the cortex compared with that in rodents

[14, 82, 83]. However, the increased electrical resistivity of the plasma membrane in human

pvalb neurons likely necessitated an adjustment of the AP threshold to preserve their rapid

electrical input–output transformation.

Mechanisms underlying the lowered firing threshold in human fast-spiking neurons

Experiments using a Kv1 channel blocker directly demonstrated the strong role of Kv1 channels in setting a higher AP threshold in mouse pvalb neurons compared with human pvalb neurons. Immunofluorescence analysis revealed a deficiency of Kv1.1 channels in human neocortical pvalb neurons, and *KCNA1* expression was absent in all human pvalb neurons, despite being present in mouse cells. These findings aligned with results from the Allen Institute transcriptomic database on averaged mRNA levels in human and mouse pvalb neuron subtypes. Although Kv1.2 channels were similarly expressed in both species, immunostaining showed weak AIS labeling in each, and the Allen Institute dataset indicated lower average Kv1.2 levels in human-type compared with mouse-type pvalb cells. However, the combined mRNA levels of Kv1.1 and Kv1.2 did not correlate with AP threshold across individual cells. Larger datasets in future studies may help clarify this relationship. Both datasets showed that the Kv1 channel gene group was the only potassium channel group expressed at lower levels in human pvalb neurons compared with mouse neurons. We argue that species differences in Kv1.1 channel density are the major determinant of the lower AP threshold in humans. Although this work does not determine the exact AP initiation site in the AIS, it shows that the low-threshold ion channels Nav1.6 and Kv1 are, on average, evenly distributed along the AIS, without a consistent intensity peak at either end.

Human pvalb neurons have a longer AIS on average relative to their mouse counterparts, and simulations showed that AIS length strongly modulates AP threshold within the physiological range observed. AIS length appears to contribute to AP threshold variability across individual pvalb cells in both species and helps partly explain the interspecies difference. The large cell-to-cell variability of AIS length and localization was also evident in the correlation between

AP threshold and AIS structure within the same cells: due to the variability in both measures,

the results remained less clear and partially diverged from the correlation findings in the larger,

independent datasets.

AIS elongation lowers the AP threshold by increasing the total number of Na channels in the

initial segment [41, 48], and this effect is especially pronounced in human AIS with low

inhibitory Kv1 channel content. Increasing AIS length reduces AP threshold more effectively

than expanding axon diameter, although both increase the number of AIS ion channels.

However, expanding axon diameter also lowers Ra, which shifts AP threshold in the opposite

direction and limits its effect [48]. In rodents, AIS elongation coincides with changes in

voltage-gated potassium channel composition to enhance excitability [45]. Human pvalb

neurons similarly exploit both geometry and reduced Kv1 activity to lower AP threshold.

Mechanisms underlying cell-to-cell variability in the AP threshold without interspecies differences

The distance from the AIS initiation site to the soma, the axon diameter, and the axon

Nav1.6 channel immunofluorescence intensity vary across individual neurons but do not

account for the species difference in AP threshold. However, these features likely contribute to

cell-to-cell variability in pvalb neuron AP thresholds in both species, as indicated by our model

cell simulations.

Increases in axon diameter [48] typically occur with reduced Ra, which raises the AP threshold

in pvalb interneurons. Given the evolutionary pressure to lower the AP threshold in human

neurons, it follows that axon diameter in human pvalb neurons is not markedly larger than that

in mouse cells, as enlargement would counteract AP threshold reduction.

The AIS is not a static cellular microdomain [84], and the observed cell-to-cell variability in

AIS properties likely results from plasticity mechanisms [85], enabling the brain to adapt, learn,

and repair. Whether these mechanisms differ between humans and animal models remains

unknown.

Conclusions

Results and conclusions regarding neocortical neural networks are likely influenced by

species-specific neuronal features. Studies that rely solely on animal models may provide an

inaccurate picture of human brain network physiology and aging. The same concerns apply to

degenerative mechanisms underlying disease processes. Therefore, identifying human-specific

neural features is crucial to developing effective therapeutic strategies for degenerative brain

conditions, in which PVALB neurons are often emphasized.

Study limitations

The lack of biotechnological tools to selectively and rapidly manipulate gene

expression, and consequently the AIS ion channel composition and microstructure, in

genetically defined neuron types, such as pvalb neurons, in human brain slices continues to

constrain experimental design. In addition, immunostaining-based localization and

quantification of proteins are limited by antibody validation, which currently relies on

knockouts only feasible in nonhuman species. Future availability of such techniques in human

tissue will enable a more detailed understanding of the human-type AIS in health and disease.

Materials and Methods

Ethics statement and licenses

All procedures were approved by the University of Szeged Ethics Committee, the

Regional Human Investigation Review Board (ref. 75/2014), and the Scientific and Research

Ethics Committee (ETT TUKEB) BM/25042-1/2024 and were conducted in accordance with

the tenets of the Declaration of Helsinki. Written informed consent was obtained from all

patients before surgery. For underage patients, consent was obtained from a parent or

guardian.

Human brain slices

Neocortical slices were prepared from frontal, temporal, or other cortical samples

resected during deep-brain surgeries. Patients ranged in age from 11 to 85 years, and samples

were obtained from both hemispheres of male and female patients (Table S1). Anesthesia was

induced with intravenous midazolam (0.03 mg/kg) and fentanyl (1–2 μg/kg) after a bolus

intravenous injection of propofol (1–2 mg/kg). Rocuronium (0.5 mg/kg) was administered to

facilitate endotracheal intubation. During surgery, patients were ventilated with a 1:2 O_2/N_2O

mixture, and anesthesia was maintained with sevoflurane. After resection, tissue blocks were

immediately immersed in an ice-cold solution containing (in mM) 130 NaCl, 3.5 KCl, 1

NaH_2PO_4, 24 $NaHCO_3$, 1 $CaCl_2$, 3 $MgSO_4$, and 10 d-(+)-glucose, which was aerated with 95%

O_2/5% CO_2 in a glass container. The container was kept on ice in an insulated box and

transported within 20 min to the electrophysiology laboratory under continuous aeration (95%

O_2/5% CO_2). Slices (350-μm thick) were cut using a vibrating blade microtome (Microm HM

650 V, Thermo Fisher Scientific, Cambridge, UK) and incubated for 1 h at 22°C–24°C in

slicing solution. This solution was gradually replaced by recording solution (180 mL) delivered

at 6 mL/min by pump. The recording solution matched the slicing solution except for 3 mM

$CaCl_2$ and 1 mM $MgSO_4$.

Mouse brain slices

Transverse slices (350 μm) were prepared from the somatosensory (n = 45) and frontal

(n = 50) cortices of 5–7-week-old heterozygous B6.129P2-Pvalbtm1(cre)Arbr/J mice (stock

017320, B6 PVcre line; Jackson Laboratory, Bar Harbor, ME, USA) crossed with Ai9 reporter

mice to express the tdTomato fluorophore in pvalb GABAergic neurons, which aided cell

selection. Cells were tentatively identified electrophysiologically through fast spike kinetics

and high-frequency, nonaccommodating firing in response to 500-ms suprathreshold

depolarizing pulses [38].

Electrophysiological recordings and data analysis

Recordings were performed in a submerged chamber perfused at 8 mL/min with

recording solution maintained at 36°C–37°C. Cells were patched under visual guidance using

infrared differential interference contrast video microscopy with a water-immersion ×20

objective, additional zoom (up to ×4), and a dichroic mirror and filter (Cy3) for TdTomato

fluorescence in mouse tissue. All recordings were conducted within 30 min of entering whole-

cell mode. Micropipettes (5–8 MΩ) were filled with intracellular solution containing (in mM)

126 K-gluconate, 8 NaCl, 4 ATP-Mg, 0.3 Na$_2$-GTP, 10 HEPES, and 10 phosphocreatine (pH

7.0–7.2; 300 mOsm) supplemented with 0.3% (w/v) biocytin for *post hoc* staining with

fluorophore-conjugated streptavidin. Recordings were performed with a Multiclamp 700B

amplifier (Axon Instruments, Scottsdale, AZ, USA) and low-pass–filtered at 6–8 kHz (Bessel

filter). Series resistance and pipette capacitance were compensated in current–clamp mode. Vm

values were not corrected for the liquid junction potential. Somatic tau was measured in

current–clamp mode with +10 mV Vm steps (250–500 ms) from −70 mV. AP threshold was

measured in dV/dt versus Vm phase plots as the Vm at which AP rise speed reached 10 mV/ms

[31, 44]. All parameters were based on at least five traces. Data were acquired with Clampex

software (Axon Instruments), digitized at 35–50 kHz, and analyzed offline using pClamp

(v10.5; Axon Instruments), Spike2 (v8.1; Cambridge Electronic Design, Milton, UK),

OriginPro (v9.5; OriginLab Corporation, Northampton, MA, USA), and SigmaPlot (v14;

Grafiti, Palo Alto, CA, USA).

Drugs

Brain slices were perfused with 100 nM dendrotoxin-K (Cat. No. D4813, Merck/Sigma-Aldrich).

Tissue fixation and cell visualization

Cells loaded with biocytin during whole-cell patch-clamp recordings were visualized using Alexa 488–conjugated streptavidin (1:1000) or Cy3-conjugated streptavidin (1:1000; both Jackson ImmunoResearch, West Grove, PA, USA). Immediately following recordings, slices were fixed in 4% paraformaldehyde and 15% picric acid in 0.1 M phosphate buffer (PB; pH 7.4) at 4°C for at least 12 h, followed by storage at 4°C in 0.1 M PB containing 0.05% sodium azide as a preservative. All slices were embedded in 20% gelatin and cut into 50-μm-thick sections using a vibratome (VT1000S, Leica Microsystems, Wetzlar, Germany) in ice-cold PB. Sections were rinsed in 0.1 M PB (3 × 10 min), cryoprotected in 10%–20% sucrose in 0.1 M PB, flash-frozen in liquid nitrogen, and thawed in 0.1 M PB. Slices were then incubated in 0.1 M Tris-buffered saline (TBS; pH 7.4) containing fluorophore-conjugated streptavidin for 2 h 30 min at 22°C–24°C, rinsed in 0.1 M PB (3 × 10 min), mounted in Vectashield mounting medium (Vector Laboratories, Burlingame, CA, USA), placed under coverslips, and examined under an epifluorescence microscope (Leica DM 5000 B, Milton Keynes, UK) at ×20–60 objectives.

For single-cell reconstructions, selected sections were incubated overnight at 4°C in a conjugated avidin–biotin horseradish peroxidase solution (1:300; Vector Labs) in TBS (pH 7.4). The enzyme reaction was developed using the glucose oxidase–DAB–nickel method with 3′3-diaminobenzidine tetrahydrochloride (0.05%) and 0.01% H_2O_2 as the chromogen and oxidant, respectively. Sections were postfixed in 1% OsO_4 in 0.1 M PB, rinsed in distilled water, and stained with 1% uranyl acetate. Dehydration was performed in a graded ethanol series, followed by overnight infiltration with epoxy resin (Durcupan, Merck kft., Budapest,

Hungary) and mounting on glass slides. Light microscopic reconstructions were performed

with the Neurolucida system (MBF Bioscience, Williston, VT, USA) using a ×100 objective

(Olympus BX51, Olympus UPlanFI, Olympus, Tokyo, Japan). Images were collapsed along

the z-axis for presentation.

Immunohistochemistry

Free-floating sections were washed in TBS containing 0.3% Triton-X (TBST) three

times over 1 h at 20°C–24°C, followed by incubation in a blocking solution containing 20%

horse serum in TBST. Antigen retrieval was performed by treating the slices with 1 mg/mL

pepsin (catalog #S3002; Dako, Glostrup, Denmark) in 1 M HCl with 0.1 M PB at 37°C for 5–

6 min, followed by rinsing in 0.1 M PB. Sections were then incubated with primary antibodies

diluted in TBST for three nights at 4°C with corresponding fluorochrome-conjugated

secondary antibodies (2% blocking serum in TBST) overnight at 4°C. Between antibody

incubations, sections were washed in TBST (3 × 15 min), followed by rinses in 0.1 M PB (2 ×

10 min). Slices were then mounted on glass slides with Vectashield mounting medium (Vector

Laboratories) for imaging.

The following primary antibodies were used for immunolabeling: goat anti-pvalb (1:1000;

PVG-213, Swant, Burgdorf, Switzerland), mouse anti-spectrin beta-4 (1:100; Invitrogen,

Thermo Fisher Scientific, Waltham, MA, USA), rabbit anti-Nav1.6 (SCN8A, 1:300; Alomone

Labs, Jerusalem, Israel), rabbit anti-Kv1.1 (KCNA1, 1:100; Alomone Labs), and rabbit anti-

Kv1.2 (KCNA2 APC-010, 1:100; Alomone Labs). Prior to confocal microscopy observations,

immunolabeling was visualized using the following secondary antibodies: DAM Alexa 488–

conjugated donkey anti-mouse (1:400; Jackson ImmunoResearch), DARb Cy3-conjugated

donkey anti-rabbit (1:400; Jackson ImmunoResearch), and DaGt A647-conjugated donkey

anti-goat (1:200; Thermo Fisher Scientific). Immunoreactivity was evaluated using an

epifluorescence microscope with a quasiconfocal option (Zeiss ApoTome 2, Zeiss, Oberkochen, Germany).

Immunofluorescence image acquisition and analysis

Immunofluorescently labeled sections were imaged using a Leica Stellaris 8 laser-scanning confocal microscope (Leica Microsystems, Biomarker Kft. Budapest, Hungary) with 499-, 554-, and 649-nm lasers to excite the fluorophores, using 504–553-, 559–654-, and 654–750-nm emission filters, respectively.

To quantify immunofluorescence around the soma, 24 radial measurement lines (50 pixels/μm) were placed on the soma such that all lines projected radially from its center into the extracellular space. Immunofluorescence intensity for pvalb (Cy5) and one of the three ion channels (Cy3) was measured along radial lines in single-plane confocal images of the cell soma. The location of the extracellular membrane was defined by the onset of the pvalb signal when moving from the extracellular to the intracellular position along each line. This position was marked as 0 and defined as follows: (average extracellular fluorescence pixel value − average intracellular fluorescence pixel value) / 2. For pooled analysis, all lines were aligned at this 0-point.

Ion channel fluorescence from the extracellular space (recorded from the 24 radial lines−3.0 to −0.5 μm relative to the 0-point, in total ca. 3000 pixels per cell) was averaged and used as a specific reference signal for each cell. This average extracellular value was subtracted from pixel intensities at the membrane (−0.5 to +0.5 μm from the 0-point) and in the intracellular cytoplasm (+0.5 to +1.5 μm from the 0-point). The extracellular site fluorescence (average intensity) did not differ significantly between species (Kv1.1, n = 31 cells, p = 0.06; Kv1.2, n = 32 and 59, p = 0.835, Mann–Whitney U test). The intracellular measurement zone was chosen near the extracellular membrane to sample the cytoplasm while avoiding the nucleus, which

was not visualized in staining. For axonal analysis, the same extracellular reference value (from 24 lines per cell) was subtracted from fluorescence intensity measurements pixels along the axonal lines.

Z-stack images were acquired from the same cells using a HC PL APO CS2 63×/1.40 oil immersion objective in unidirectional scanning mode. Axonal fluorescence measurement lines were sampled at 50 pixels/μm. AIS immunofluorescence in z-stack images was analyzed using a custom ImageJ macro (US National Institutes of Health, Bethesda, MD, USA) and a script developed by Dr. Adam Tiszlavicz [86] to determine the AIS's length and its proximal and terminal points (visualized via beta-IV-spectrin immunoreaction). This was achieved by manually marking the points and applying the Euclidean distance formula. Fluorescence intensity was then measured along a polyline connecting these points, capturing the full signal profile of the immunolabeled channels. AIS lengths were normalized by dividing into 10 bins (from proximal to distal). Ion channel immunoreactivity (Cy3) was measured in parallel with pvalb (Cy5) and beta-IV-spectrin (Alexa 488), and axonal fluorescence values were referenced to the average extracellular signal of the corresponding cell, using the same normalization approach applied for somatic radial lines.

Nucleus collection for patch-sequencing

Nuclei were collected from cells exhibiting a fast-spiking phenotype. Prior to data collection, all surfaces, equipment, and materials were thoroughly cleaned using DNA AWAY™ (Thermo Fisher Scientific) and RNaseZap™ (Sigma-Aldrich, St. Louis, MO, USA). The perfusion system was flushed with 50 mL of 70% ethanol. Following electrophysiological recording, the pipette was centered on the soma and a small, constant negative pressure was applied to extract the cytosol and draw the nucleus toward the tip. The pipette was slowly retracted under continuous negative pressure while monitoring seal integrity. Once the nucleus

appeared at the pipette tip, retraction was accelerated to fully withdraw the pipette from the slice. The pipette containing the internal solution, cytosol, and nucleus was removed from the holder and expelled into a 0.2-mL tight-lock PCR tube (TubeOne, StarLab, Hamburg, Germany) preloaded with 2 μL of lysis buffer comprising 0.8% Triton X-100; (Sigma-Aldrich) and 1 U/μL RNase inhibitor (Takara, Kyoto, Japan). The internal solution contained the following: 110 mM K-gluconate (Sigma-Aldrich), 4 mM KCl solution, RNase-free (Invitrogen), 10 mM HEPES solution (Sigma-Aldrich), 1 mM ATP-Mg (Sigma-Aldrich), 0.3 mM GTP-Na (Sigma-Aldrich), 10 mM sodium phosphocreatine (Sigma-Aldrich), 0.2 mM EGTA (Sigma-Aldrich), 20 μg/mL RNA-grade glycogen solution (Thermo Fisher Scientific), and 0.5 U/μL RNase inhibitor (Takara). The 4-μL sample was briefly spun (2 s), placed on dry ice, and stored at −80°C until in-tube reverse transcription.

cDNA amplification and library construction

Libraries were prepared using a protocol based on Smart-seq2 [87, 88]. Briefly, cDNA was generated and amplified using 24 PCR cycles for all samples, including negative controls. Indexes and adapter sequences were added to the cDNA, which was pooled equimolarly and sequenced on an Illumina MiSeq platform (San Diego, CA, USA) in 50-bp single-end read mode. Each sample was sequenced to a depth of ~1–2 million reads.

Sequencing data processing

Sequencing used 66-bp paired-end reads aligned to the GRCh38 reference genome with annotation file GRCh38_r93 from Ensembl. Alignment was performed using STAR v2.7.11a [89] with the following parameters: --soloType SmartSeq --soloUMIdedup Exact --soloStrand Unstranded --soloFeatures GeneFull. Full gene counts, including introns and exons, produced via STAR were used to compute TPM values for downstream analysis.

Identification of pvalb neurons based on single-nucleus transcriptomics

Single-nucleus transcriptomic data were analyzed using the Allen Institute's MapMyCells tool (Allen Institute for Brain Science, 2024; RRID:SCR_024672, https://knowledge.brain-map.org/mapmycells/process/). This tool uses gene expression signatures to assign neurons to cortical cell types, enabling identification of pvalb neurons even when *PVALB/pvalb* expression is undetectable (Fig S1). For human cells, classification used the Seattle Alzheimer's Disease Brain Cell Atlas reference taxonomy with the Deep Generative Mapping algorithm; for mouse cells, the 10X Whole Mouse Brain reference taxonomy was employed with the hierarchical mapping algorithm. Only cells classified as pvalb neurons at p > 0.9 (supertype_bootstrapping_probability) were retained for further analysis. Human single-nucleus gene expression profiles showed strong concordance with the Allen Institute's high-resolution reference dataset for pvalb subtypes (subclass correlation coefficient > 0.5 using the 10X Whole Human Brain dataset and correlation mapping algorithm; Fig 8).

Single-cell model

The three-compartment neuron model included cylindrical dendritic and axonal compartments and followed a previously established formalism [28]. The model comprised a soma; a 140-μm-long, 0.8-μm-diameter axon; and a 200-μm-long, 1-μm-diameter dendrite. Axonal cable parameters were as follows: axon Ra = 0.8–1.1 MΩ·μm; axon Cm = 6 fF/μm^2. Dendritic Cm was set to 12 fF/μm^2, i.e., 2-fold higher than that of the axon. The somatic compartment had a resistance of 700 MΩ and a Cm of 25 pF. The leakage reversal potential was set to −68 mV. Rheobase and voltage sag parameters were chosen to match the basic response properties and carefully adjusted using the experimental data to match the simulated

1030 and recorded voltage traces. All intrinsic voltage-dependent currents were calculated using the

1031 following equation:

1032

1033
$$I_i \;=\; g_i m_i^p h_i (E_i - V),$$

1034

1035 where i represents individual current type; g_i is the maximal conductance of the current; m_i

1036 and h_i are the activation and inactivation (either first-order or absent) variables, respectively;

1037 p is the activation exponent; and E_i is the reversal potential. Activation (m) and inactivation

1038 (h) kinetics were modeled based on the following equation:

1039

1040
$$\frac{dx}{dt} \;=\; \frac{x_\infty(V) - x}{\tau_x(V)},$$

1041

1042 where x represents m or h. The voltage-dependent steady-state activation and inactivation are

1043 described by the following sigmoid function:

1044

1045
$$x_\infty(V) \;=\; \frac{1}{2} + \frac{1}{2}\tanh\left(\frac{V - V_{x,1/2}}{V_{x,sl}}\right).$$

1046

1047 The midpoint $V_{x,1/2}$ and slope $V_{x,sl}$ parameters of the sigmoids and other kinetic parameters were

1048 presented in previous study [63]. The tau (τ) values of activation and inactivation were modeled

1049 as bell-shaped functions of the Vm (V) according to the following equation:

1050

$$\tau_x(V) = \left(\tau_{x,max} - \tau_{x,min}\right)\left[1 - \tanh\left(\frac{V - V_{\tau x,1/2}}{V_{\tau x,sl}}\right)^2\right] + \tau_{x,min}.$$

EPSP waveforms were generated by EPSCs with a time to peak of 0.5 ms, a decay time constant of 3 ms, a reversal potential of 0 mV, and peak conductances of 6–11 nS.

Statistical analysis

Results are presented as medians and IQRs. Statistical comparisons were performed using the Student's t-test, the Mann–Whitney U test, Wilcoxon signed-rank test with Bonferroni correction, ANOVA on ranks (Kruskal–Wallis H test) with Dunn's *post hoc* pairwise test, or PERMANOVA and the Mann–Whitney U test with Bonferroni correction. Proportions were compared using the chi-squared test with Benjamini–Hochberg adjustment at a 5% significance level. Correlations were assessed using Spearman's rank-order method.

Acknowledgments

We thank both Dr. Ferhan Ayaydin at the Functional Cell Biology and Immunology Advanced Core Facility, Hungarian Center of Excellence for Molecular Medicine, and Szeged University and Dr. Fanni Nedenyi at HCEMM for help with statistical analyses. The funders played no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. The authors would like to thank Enago (www.enago.com) for the English language review.

References

1. Galakhova AA, Hunt S, Wilbers R, Heyer DB, de Kock CPJ, Mansvelder HD, et al. Evolution of cortical neurons supporting human cognition. Trends Cogn Sci. 2022;26: 909-922.

2. Nehme R, Pietiläinen O, Barrett LE. Genomic, molecular, and cellular divergence of the human brain. Trends Neurosci. 2024;47: 491-505.

3. Warming H, Pegasiou C-M, Pitera AP, Kariis H, Houghton SD, Kurbatskaya K, et al. A primate-specific short GluN2A-NMDA receptor isoform is expressed in the human brain. Mol Brain. 2019;12: 64.

4. Herculano-Houzel S, Catania K, Manger PR, Kaas JH. Mammalian brains are made of these: a dataset of the numbers and densities of neuronal and nonneuronal cells in the brain of glires, primates, scandentia, eulipotyphlans, afrotherians and artiodactyls, and their relationship with body mass. Brain Behav Evol. 2015;86: 145-163.

5. Berg J, Sorensen SA, Ting JT, Miller JA, Chartrand T, Buchin A, et al. Human neocortical expansion involves glutamatergic neuron diversification. Nature. 2021;598: 151-158.

6. Hodge RD, Bakken TE, Miller JA, Smith KA, Barkan ER, Graybuck LT, et al. Conserved cell types with divergent features in human versus mouse cortex. Nature. 2019;573: 61-68.

7. Zeng H, Shen EH, Hohmann JG, Oh SW, Bernard A, Royall JJ, et al. Large-scale cellular-resolution gene profiling in human neocortex reveals species-specific molecular signatures. Cell. 2012;149: 483-496.

8. Suresh H, Crow M, Jorstad N, Hodge R, Lein E, Dobin A, et al. Comparative single-cell transcriptomic analysis of primate brains highlights human-specific regulatory evolution. Nat Ecol Evol. 2023;7: 1930-1943.

9. Rossi F. Evolutionary mechanisms and neural adaptation: selective versus constructive strategies in the development and plasticity of the nervous system. In: Fasolo A, editor. The theory of evolution and its impact. Milano: Springer; 2012. pp. 159-174.

10. Beauchamp A, Yee Y, Darwin BC, Raznahan A, Mars RB, Lerch JP. Whole-brain comparison of rodent and human brains using spatial transcriptomics. Elife. 2022;11.

11. Boldog E, Bakken TE, Hodge RD, Novotny M, Aevermann BD, Baka J, et al. Transcriptomic and morphophysiological evidence for a specialized human cortical GABAergic cell type. Nat Neurosci. 2018;21: 1185-1195.

12. Campagnola L, Seeman SC, Chartrand T, Kim L, Hoggarth A, Gamlin C, et al. Local connectivity and synaptic dynamics in mouse and human neocortex. Science. 2022;375: eabj5861.

13. Chittajallu R, Auville K, Mahadevan V, Lai M, Hunt S, Calvigioni D, et al. Activity-dependent tuning of intrinsic excitability in mouse and human neurogliaform cells. Elife. 2020;9: e57571.

14. Clowry GJ. An enhanced role and expanded developmental origins for gamma-aminobutyric acidergic interneurons in the human cerebral cortex. J Anat. 2015;227: 384-393.

15. Dehay C, Huttner WB. Development and evolution of the primate neocortex from a progenitor cell perspective. Development. 2024;151: dev199171.

16. Heyer DB, Wilbers R, Galakhova AA, Hartsema E, Braak S, Hunt S, et al. Verbal and general IQ associate with supragranular layer thickness and cell properties of the left temporal cortex. Cereb Cortex. 2022;32: 2343-2357.

17. Hu X, Rocco BR, Fee C, Sibille E. Cell type-specific gene expression of alpha 5 subunit-containing gamma-aminobutyric acid subtype A receptors in human and mouse frontal cortex. Mol Neuropsychiatry. 2019;4: 204-215.

18. Jorstad NL, Song JH, Exposito-Alonso D, Suresh H, Castro N, Krienen FM, et al.

Comparative transcriptomics reveals human-specific cortical features. Science. 2023;382:

eade9516.

19. Kalmbach BE, Buchin A, Long B, Close J, Nandi A, Miller JA, et al. h-channels

contribute to divergent intrinsic membrane properties of supragranular pyramidal neurons in

human versus mouse cerebral cortex. Neuron. 2018;100: 1194-1208.e5.

20. Shepherd GM, Rowe TB. Neocortical lamination: insights from neuron types and

evolutionary precursors. Front Neuroanat. 2017;11: 100.

21. Lim L, Mi D, Llorca A, Marín O. Development and functional diversification of cortical

interneurons. Neuron. 2018;100: 294-313.

22. Tremblay R, Lee S, Rudy B. GABAergic interneurons in the neocortex: from cellular

properties to circuits. Neuron. 2016;91: 260-292.

23. Lee BR, Dalley R, Miller JA, Chartrand T, Close J, Mann R, et al. Signature

morphoelectric properties of diverse GABAergic interneurons in the human neocortex.

Science. 2023;382: eadf6484.

24. Mihaljević B, Larrañaga P, Bielza C. Comparing the electrophysiology and morphology

of human and mouse layer 2/3 pyramidal neurons with bayesian networks. Front

Neuroinform. 2021;15: 580873.

25. Moradi Chameh H, Rich S, Wang L, Chen FD, Zhang L, Carlen PL, et al. Diversity

amongst human cortical pyramidal neurons revealed via their sag currents and frequency

preferences. Nat Commun. 2021;12: 2497.

26. Olah S, Komlosi G, Szabadics J, Varga C, Toth E, Barzo P, et al. Output of

neurogliaform cells to various neuron types in the human and rat cerebral cortex. Front

Neural Circuits. 2007;1: 4.

27. Poorthuis RB, Muhammad K, Wang M, Verhoog MB, Junek S, Wrana A, et al. Rapid

neuromodulation of layer 1 interneurons in human neocortex. Cell Rep. 2018;23: 951-958.

28. Szegedi V, Bakos E, Furdan S, Kovács BH, Varga D, Erdélyi M, et al. HCN channels at

the cell soma ensure the rapid electrical reactivity of fast-spiking interneurons in human

neocortex. PLoS Biol. 2023;21: e3002001.

29. Testa-Silva G, Rosier M, Honnuraiah S, Guzulaitis R, Megias AM, French C, et al. High

synaptic threshold for dendritic NMDA spike generation in human layer 2/3 pyramidal

neurons. Cell Rep. 2022;41: 111787.

30. Verhoog MB, Goriounova NA, Obermayer J, Stroeder J, Hjorth JJ, Testa-Silva G, et al.

Mechanisms underlying the rules for associative plasticity at adult human neocortical

synapses. J Neurosci. 2013;33: 17197-17208.

31. Wilbers R, Galakhova AA, Driessens SLW, Heistek TS, Metodieva VD, Hagemann J, et

al. Structural and functional specializations of human fast spiking neurons support fast

cortical signaling. Sci Adv. 2023;9: eadf0708.

32. Krubitzer L. In search of a unifying theory of complex brain evolution. Ann N Y Acad

Sci. 2009;1156: 44-67.

33. Rich S, Moradi Chameh H, Sekulic V, Valiante TA, Skinner FK. Modeling reveals

human-rodent differences in H-current kinetics influencing resonance in cortical layer 5

neurons. Cereb Cortex. 2021;31: 845-872.

34. Libé-Philippot B, Lejeune A, Wierda K, Louros N, Erkol E, Vlaeminck I, et al.

LRRC37B is a human modifier of voltage-gated sodium channels and axon excitability in

cortical neurons. Cell. 2023;186: 5766-5783.e25.

35. Mertens EJ, Leibner Y, Pie J, Galakhova AA, Waleboer F, Meijer J, et al. Morpho-

electric diversity of human hippocampal CA1 pyramidal neurons. Cell Rep. 2024;43: 114100.

36. Povysheva NV, Zaitsev AV, Rotaru DC, Gonzalez-Burgos G, Lewis DA, Krimer LS.

Parvalbumin-positive basket interneurons in monkey and rat prefrontal cortex. J

Neurophysiol. 2008;100: 2348-2360.

37. Wang B, Ke W, Guang J, Chen G, Yin L, Deng S, et al. Firing frequency maxima of fast-

spiking neurons in human, monkey, and mouse neocortex. Front Cell Neurosci. 2016;10: 239.

38. Szegedi V, Molnár G, Paizs M, Csakvari E, Barzó P, Tamás G, et al. High-precision fast-

spiking basket cell discharges during complex events in the human neocortex. eNeuro.

2017;4: ENEURO.0260-17.2017.

39. Szegedi V, Paizs M, Baka J, Barzo P, Molnar G, Tamas G, et al. Robust perisomatic

GABAergic self-innervation inhibits basket cells in the human and mouse supragranular

neocortex. Elife. 2020;9: e51691.

40. Hijazi S, Smit AB, van Kesteren RE. Fast-spiking parvalbumin-positive interneurons in

brain physiology and Alzheimer's disease. Mol Psychiatry. 2023;28: 4954-4967.

41. Huang CY, Rasband MN. Axon initial segments: structure, function, and disease. Ann N

Y Acad Sci. 2018;1420: 46-61.

42. Leterrier C. The axon initial segment: an updated viewpoint. J Neurosci. 2018;38: 2135-

2145.

43. Hu H, Gan J, Jonas P. Fast-spiking, parvalbumin(+) GABAergic interneurons: from

cellular design to microcircuit function. Science. 2014;345: 1255263.

44. Li T, Tian C, Scalmani P, Frassoni C, Mantegazza M, Wang Y, et al. Action potential

initiation in neocortical inhibitory interneurons. PLoS Biol. 2014;12: e1001944.

45. Yamada R, Kuba H. Structural and functional plasticity at the axon initial segment. Front

Cell Neurosci. 2016;10: 250.

46. Yang Y, Ogawa Y, Hedstrom KL, Rasband MN. βIV spectrin is recruited to axon initial

segments and nodes of Ranvier by ankyrinG. J Cell Biol. 2007;176: 509-519.

47. Jones SL, Svitkina TM. Axon initial segment cytoskeleton: architecture, development, and role in neuron polarity. Neural Plast. 2016;2016: 1-19.

48. Goethals S, Brette R. Theoretical relation between axon initial segment geometry and excitability. Elife. 2020;9: e54342.

49. Kole MH, Stuart GJ. Signal processing in the axon initial segment. Neuron. 2012;73: 235-247.

50. Kuba H, Ohmori H. Roles of axonal sodium channels in precise auditory time coding at nucleus magnocellularis of the chick. J Physiol. 2009;587: 87-100.

51. Lorincz A, Nusser Z. Cell-type-dependent molecular composition of the axon initial segment. J Neurosci. 2008;28: 14329-14340.

52. O'Brien JE, Meisler MH. Sodium channel SCN8A (Nav1.6): properties and de novo mutations in epileptic encephalopathy and intellectual disability. Front Genet. 2013;4: 213.

53. Goldberg EM, Clark BD, Zagha E, Nahmani M, Erisir A, Rudy B. K+ channels at the axon initial segment dampen near-threshold excitability of neocortical fast-spiking GABAergic interneurons. Neuron. 2008;58: 387-400.

54. Kole MH, Letzkus JJ, Stuart GJ. Axon initial segment Kv1 channels control axonal action potential waveform and synaptic efficacy. Neuron. 2007;55: 633-647.

55. Gu C, Barry J. Function and mechanism of axonal targeting of voltage-sensitive potassium channels. Prog Neurobiol. 2011;94: 115-132.

56. D'Adamo MC, Liantonio A, Rolland J-F, Pessia M, Imbrici P. Kv1.1 channelopathies: pathophysiological mechanisms and therapeutic approaches. Int J Mol Sci. 2020;21: 2935.

57. Ogawa Y, Horresh I, Trimmer JS, Bredt DS, Peles E, Rasband MN. Postsynaptic density-93 clusters Kv1 channels at axon initial segments independently of Caspr2. J Neurosci. 2008;28: 5731-5739.

58. Higgs MH, Spain WJ. Kv1 channels control spike threshold dynamics and spike timing in

cortical pyramidal neurones. J Physiol. 2011;589: 5125-5142.

59. Shao M, Zhang W, Li Y, Tang L, Hao ZZ, Liu S. Patch-seq: advances and biological

applications. Cell Mol Neurobiol. 2023;44: 8.

60. Cadwell CR, Tolias AS. Patch-seq: multimodal profiling of single-cell morphology,

electrophysiology, and gene expression. Methods Mol Biol. 2024;2752: 227-243.

61. Chow A, Erisir A, Farb C, Nadal MS, Ozaita A, Lau D, et al. K(+) channel expression

distinguishes subpopulations of parvalbumin- and somatostatin-containing neocortical

interneurons. J Neurosci. 1999;19: 9332-9345.

62. Martina M, Schultz JH, Ehmke H, Monyer H, Jonas P. Functional and molecular

differences between voltage-gated K+ channels of fast-spiking interneurons and pyramidal

neurons of rat hippocampus. J Neurosci. 1998;18: 8111-8125.

63. Szegedi V, Tiszlavicz Á, Furdan S, Douida A, Bakos E, Barzo P, et al. Aging-associated

weakening of the action potential in fast-spiking interneurons in the human neocortex. J

Biotechnol. 2024;389: 1-12.

64. Krubitzer L, Kaas J. The evolution of the neocortex in mammals: how is phenotypic

diversity generated? Curr Opin Neurobiol. 2005;15: 444-453.

65. Blekhman R, Oshlack A, Chabot AE, Smyth GK, Gilad Y. Gene regulation in primates

evolves under tissue-specific selection pressures. PLoS Genet. 2008;4: e1000271.

66. Vallender EJ, Mekel-Bobrov N, Lahn BT. Genetic basis of human brain evolution.

Trends Neurosci. 2008;31: 637-644.

67. Rosales-Reynoso MA, Juárez-Vázquez CI, Barros-Núñez P. Evolution and genomics of

the human brain. Neurología. 2018;33: 254-265.

68. Caglayan E, Ayhan F, Liu Y, Vollmer RM, Oh E, Sherwood CC, et al. Molecular features

driving cellular complexity of human brain evolution. Nature. 2023;620: 145-153.

69. Zhou Y, Song H, Ming GL. Genetics of human brain development. Nat Rev Genet.

2024;25: 26-45.

70. Kawaguchi Y, Kubota Y. GABAergic cell subtypes and their synaptic connections in rat

frontal cortex. Cereb Cortex. 1997;7: 476-486.

71.Gonchar Y, Wang Q, Burkhalter AH. Multiple distinct subtypes of GABAergic neurons in

mouse visual cortex identified by triple immunostaining. Front Neuroanat. 2007;1: 3.

72. Ruden JB, Dugan LL, Konradi C. Parvalbumin interneuron vulnerability and brain

disorders. Neuropsychopharmacol. 2021;46: 279-287.

73. Lewis DA, Curley AA, Glausier JR, Volk DW. Cortical parvalbumin interneurons and

cognitive dysfunction in schizophrenia. Trends Neurosci. 2012;35: 57-67.

74. Nahar L, Delacroix BM, Nam HW. The role of parvalbumin interneurons in

neurotransmitter balance and neurological disease. Front Psychiatry. 2021;12: 679960.

75. Marin O. Parvalbumin interneuron deficits in schizophrenia. Eur Neuropsychopharmacol.

2024;82: 44-52.

76. Brickley SG, Revilla V, Cull-Candy SG, Wisden W, Farrant M. Adaptive regulation of

neuronal excitability by a voltage- independent potassium conductance. Nature. 2001;409:

88-92.

77. Hall CN, Klein-Flugge MC, Howarth C, Attwell D. Oxidative phosphorylation, not

glycolysis, powers presynaptic and postsynaptic mechanisms underlying brain information

processing. J Neurosci. 2012;32: 8940-8951.

78. Howarth C, Gleeson P, Attwell D. Updated energy budgets for neural computation in the

neocortex and cerebellum. J Cereb Blood Flow Metab. 2012;32: 1222-1232.

79. McCasland JS, Hibbard LS. GABAergic neurons in barrel cortex show strong, whisker-

dependent metabolic activation during normal behavior. J Neurosci. 1997;17: 5509-5527.

80. Kann O, Papageorgiou IE, Draguhn A. Highly energized inhibitory interneurons are a

central element for information processing in cortical networks. J Cereb Blood Flow Metab.

2014;34: 1270-1282.

81. Ghirardini E, Sagona G, Marquez-Galera A, Calugi F, Navarron CM, Cacciante F, et al.

Cell-specific vulnerability to metabolic failure: the crucial role of parvalbumin expressing

neurons in creatine transporter deficiency. Acta Neuropathol Commun. 2023;11: 34.

82. Fonseca-Azevedo K, Herculano-Houzel S. Metabolic constraint imposes tradeoff between

body size and number of brain neurons in human evolution. Proc Natl Acad Sci USA.

2012;109: 18571-18576.

83. Pontzer H, Brown MH, Raichlen DA, Dunsworth H, Hare B, Walker K, et al. Metabolic

acceleration and the evolution of human brain size and life history. Nature. 2016;533: 390-

392.

84. Höfflin F, Jack A, Riedel C, Mack-Bucher J, Roos J, Corcelli C, et al. Heterogeneity of

the axon initial segment in interneurons and pyramidal cells of rodent visual cortex. Front

Cell Neurosci. 2017;11: 332.

85. Hamada MS, Goethals S, de Vries SI, Brette R, Kole MH. Covariation of axon initial

segment location and dendritic tree normalizes the somatic action potential. Proc Natl Acad

Sci USA. 2016;113: 14841-14846.

86. Ollion J, Cochennec J, Loll F, Escudé C, Boudier T. TANGO: a generic tool for high-

throughput 3D image analysis for studying nuclear organization. Bioinformatics. 2013;29:

1840-1841.

87. Picelli S, Faridani OR, Björklund ÅK, Winberg G, Sagasser S, Sandberg R. Full-length

RNA-seq from single cells using Smart-seq2. Nat Protoc. 2014;9: 171-181.

88. Hennig BP, Velten L, Racke I, Tu CS, Thoms M, Rybin V, et al. Large-scale low-cost NGS library preparation using a robust Tn5 purification and tagmentation protocol. G3. 2018;8: 79-89.

89. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29: 15-21.

Funding: Project no. TKP-2021-EGA-05 was implemented with support provided by the Ministry of Culture and Innovation of Hungary from the National Research, Development, and Innovation Fund, financed under the TKP2021-EGA funding scheme (K.L). Project no. 2022-2.1.1-NL-2022-00005 was implemented with support provided by the Ministry of Culture and Innovation of Hungary from the National Research, Development, and Innovation Fund, financed under the 2022-2.1.1-NL funding scheme (K.L). Additional funding was received from the EU's Horizon 2020 research and innovation program under grant agreement no. 739593 (K.L); Nemzeti Kutatási, Fejlesztési és Innovaciós Alap OTKA K 134279 and ADVANCED_150382 Magyar Tudományos Akadémia (K.L); the Hungarian Scientific Research Foundation grant number ANN-135291 (K.L); Eötvös Loránd Research Network grants ELKH-SZTE Agykérgi Neuronhálózatok Kutatócsoport and KÖ-36/2021 (G.T.); Ministry of Human Capacities Hungary (20391-3/2018/FEKUSTRAT (G.T.) and NKP 16-3-VIII-3) (G.T.); National Research, Development and Innovation Office grants GINOP 2.3.2-15-2016-00018 (G.T); Élvonal KKP 133807 (G.T.), ÚNKP-20-5 - SZTE-681 (G.T.), 2019-2.1.7-ERA-NET-2022-00038 (G.T.), TKP2021-EGA-09 (G.T.), TKP-2021-EGA-28 (G.T.). and OTKA K128863 (G.T.); National Institutes of Health awards U01MH114812 (G.T.) and UM1MH130981 (G.T.); and National Research, Development, and Innovation Office grantno. ANN_135291 (A.S.).

1313   **Author contributions:** Conceptualization: K.L. and V.S. Methodology: A.S., A.T., P.B., B.B.,

1314   G.H., G.T., and V.B. Investigation: E.B., A.T., V.S., A.D., D.W., J.L., S.F., and A.S. Data

1315   curation: V.S., A.D., and E.B. Formal analysis: A.T., E.B., V.S., and S.F. Supervision: K.L.

1316   Project administration: K.L. and V.S. Funding acquisition: K.L., G.T., and A.S. Writing

1317   original draft: K.L. Writing review and editing: K.L.

1318

1319   **Competing interests:** The authors declare that they have no competing interests.

1320

1321   **Data and materials availability:** All data required to evaluate the conclusions drawn in this

1322   paper are provided in the main text and Supplementary Materials. All datasets and custom

1323   codes used in the study are stored at the Hungarian Center of Excellence for Molecular

1324   Medicine, Szeged, Hungary, and are available upon request from the corresponding author.

1325

1326

1327

Supplementary Materials

1328

Fig. S1. Model cell simulation incorporating AIS parameter variability, showing no difference

between species.

(A) Increasing AIS Nav1.6 conductance (gNav1.6, 15–40 nS/μm^2; abscissa) progressively lowered the

AP firing threshold across multiple Kv1 activity levels (0–10 nS/μm^2). AIS length = 20 μm; axon Ra =

0.8–1.1 MΩ·μm; axon diameter = 0.8 μm. (B) Proximal–distal positioning of the AIS regulated the AP

firing threshold within the range observed anatomically. Simulations used square-pulse current steps to

mimic Vm changes (as shown in Fig 8). (B1) Vm traces from simulations in the model cell with the

AIS positioned either directly at the axon hillock (0 µm; purple) or 10 µm distally (brown), with consistent AIS length (20 µm), axon Ra (0.8–1.1 MΩ·µm), axon diameter (0.8 µm), and channel conductances (20 nS/µm^2 for Nav1.6 and Kv1). (B2) Firing threshold values plotted for AIS locations and across three Ra conditions (0.8–1.1 MΩ·µm, indicated by symbols). A more distal AIS position consistently resulted in a lower AP firing threshold, replicating the real-neuron correlation observed in Fig 3C1. (C) Heatmap shows the relationship between axon longitudinal conductance (normally reduced by axon diameter increase) and axon diameter simulating a proportional (1.3-fold) increase in both Kv1 and Nav1.6 ion conductance with increasing axon diameter, uncoupling parameters that are typically correlated in real neurons. Increasing only the Ra lowered the AP firing threshold. Conversely, increasing the ion channel conductance with an increased AIS membrane area, while keeping the channel density fixed at 20 nS/µm² for Nav1.6 and Kv1, increased the firing threshold in the model cell. This indicates the strong effect of Kv1 channels on the threshold.

model cell: altered distance of the AIS initiation site to the soma

1349

1350

1351 **Table S1. List of neocortical pvalb cells included in the study.**

1352

cell code	pvalb+ or mRNA	AP Thr	gender	age (years)	hemisphere	region	diagnosis
h1	pvalb+	-42	F	69	right	occipital	subcortical neoplasia
h2	pvalb+	-37,3	M	38	right	temporal	malformation
h3	pvalb+	-46,8	M	38	right	temporal	malformation
h4	pvalb+	-38,5	M	66	right	temporal	tumor
h5	pvalb+	-40,1	M	45	left	frontal	tumor
h6	pvalb+	-48,2	M	21	right	parietal	tumor
h7	pvalb+	-39,1	M	21	right	parietal	tumor
h8	pvalb+	-32,9	M	40	right	frontal	tumor
h9	pvalb+	-45,6	F	70	left	parietal	hydrocephalus
h10	pvalb+	-40,2	M	69	right	parietal	hydrocephalus
h11	pvalb+	-36,8	M	58	right	frontal	tumor
h12	pvalb+	-46,1	M	58	right	frontal	tumor
h13	pvalb+	-47	F	66	right	frontal	tumor
h14	pvalb+	-26,6	F	21	right	frontal	hydrocephalus
h15	pvalb+	-27,5	no info	no info	left	parietal	tumor
h16	pvalb+	-35	F	72	right	temporobasal	tumor
h17	pvalb+	-41,3	F	72	right	temporobasal	tumor
h18	pvalb+	-46,4	M	47	left	frontotemporal	tumor
h19	pvalb+	-34,7	M	38	right	temporal	malformation
h20	pvalb+	-38,8	M	38	right	temporal	malformation
h21	pvalb+	-45,4	M	34	left	no info	malformation
h22	pvalb+	-49,1	M	11	left	temporal	malformation
h23	pvalb+	-40,5	F	61	right	frontal	tumor
h24	pvalb+	-27,9	F	17	right	parietal	tumor
h25	pvalb+	-31,2	F	65	right	frontal	colloid cysta
h26	pvalb+	-30,6	M	85	left	fronto-temporal	hydrocephalus
h27	pvalb+	-37,6	F	28	right	temporal	hydrocephalus
h28	pvalb+	-37	F	30	no info	no info	hydrocephalus
h29	pvalb+	-39,9	F	40	right	frontal	anaplastic ependymoma
h30	pvalb+	-36,8	F	40	right	frontal	anaplastic ependymoma
h31	pvalb+	-35,6	F	42	left	fronto-temporal	astrocytoma
h32	pvalb+	-39	F	78	right	frontal	hydrocephalus
h33	pvalb+	-41,1	F	68	right	temporal	hydrocephalus
h34	pvalb+	-49	F	68	right	temporal	hydrocephalus
h35	pvalb+	-38,9	F	19	right	frontal	subcortical neoplasia
h36	pvalb+	-55,8	F	19	right	frontal	subcortical neoplasia
h37	pvalb+	-52	F	50	left	occipital	astrocytoma
h38	pvalb+	-20,1	F	54	right	frontal	aneurysm
h39	pvalb+	-53,1	F	69	no info	no info	hydrocephalus
h40	pvalb+	-25,5	M	36	right	frontal	hydrocephalus
h41	pvalb+	-30	M	37	right	parietal	tumor
h42	pvalb+	-48,7	F	60	right	parietal	hydrocephalus
h43	pvalb+	-31	M	22	right	frontal	hydrocephalus
h44	pvalb+	-40,7	F	39	right	frontal	tumor
h45	pvalb+	-47,7	F	63	right	temporal	hydrocephalus
h46	pvalb+	-52,6	F	68	right	temporal	hydrocephalus
h47	pvalb+	-42,2	F	76	right	frontal	Hydrocephalus

1353

h48	pvalb+	-41,1	F	73	right	frontal	biopsy (MRI: multifocal lesions)
h49	pvalb+	-40,9	F	73	right	frontal	biopsy (MRI: multifocal lesions)
h50	pvalb+	-54	M	33	right	frontal	hydrocephalus
h51	pvalb+	-39	M	20	right	frontal	hydrocephalus
h52	pvalb+	-42,7	M	49	right	frontal	colloid cyst
h53	pvalb+	-47,6	M	66	right	frontal	astrocytoma
h54	pvalb+	-36,2	M	67	right	occipital	subcortical neoplasia
h55	pvalb+	-53,3	M	68	no info	temporal	hydrocephalus
h56	pvalb+	-45,5	M	51	no info	no info	hydrocephalus
h57	pvalb+	-38,4	F	47	right	frontal	colloid cyst
h58	pvalb+	-38,4	F	53	left	no info	tumor
h59	pvalb+	-45,9	F	30	right	frontal	tumor
h60	pvalb+	-36,4	F	30	right	frontal	tumor
h61	pvalb+	-34,8	M	41	right	parietal	shunt
h62	pvalb+	-32,3	M	20	right	temporal	tumor
h63	pvalb+	-37,7	M	43	right	frontal	tumor
h64	pvalb+	-37	M	65	right	temporal	tumor
h65	pvalb+	-35	M	65	right	temporal	tumor
h66	pvalb+	-38,5	F	67	right	parietal	shunt
h67	pvalb+	-40,6	F	55	left	temporal	tumor
h68	pvalb+	-39,3	F	79	right	parietal	hydrocephalus
h69	mRNA	-31,9	F	51	right	Insula	tumor
h70	mRNA	-25,5	M	no info	right	frontal	tumor
h71	mRNA	-42,8	M	no info	right	frontal	tumor
h72	mRNA	-28,9	M	no info	right	frontal	tumor
h73	mRNA	-26,6	M	no info	right	frontal	tumor
h74	mRNA	-21,9	M	no info	right	frontal	tumor
h75	mRNA	-32,8	M	70	left	temporal	cavernoma
h76	mRNA	-37,6	F	51	right	Insula	tumor
h77	mRNA	-31,1	F	51	right	insula	tumor
h78	mRNA	-33,9	F	51	right	insula	tumor
h79	mRNA	-40,5	M	no info	left	frontal	hydrocephalus
h80	mRNA	-32,8	M	59	right	frontal	tumor
h81	pvalb+	-19,1	F	46	right	frontal	tumor
h82	pvalb+	-18,3	F	46	right	frontal	tumor
h83	pvalb+	-26,1	F	46	right	frontal	tumor
h84	pvalb+	-34,9	M	68	left	occipital	tumor
h85	pvalb+	-43,3	M	68	left	occipital	tumor
h86	pvalb+	-41,9	M	68	left	occipital	tumor
h87	pvalb+	-43,3	F	31	left	temporal	tumor
h88	pvalb+	-37,3	M	70	right	temporal	tumor
h89	pvalb+	-40,4	M	70	right	temporal	tumor
h90	pvalb+	-34,2	M	70	right	temporal	tumor
h91	pvalb+	-38,2	M	70	right	temporal	tumor

1354