Screening Envelope Genes Across Primate Genomes Reveals Evolution and

Diversity Patterns of Endogenous Retroviruses

Saili Chabukswar[1,2], Nicole Grandi[2], Elena Soddu[1,2], Liang-Tzung Lin[3,4*], Enzo

Tramontano[2*]

[1]International Ph.D. Program in Medicine, College of Medicine, Taipei Medical University,

Taipei 110, Taiwan

[2]Laboratory of Molecular Virology, Department of Life and Environmental Sciences,

University of Cagliari, 09042 Cagliari, Italy

[3]Department of Microbiology and Immunology, School of Medicine, College of Medicine,

Taipei Medical University, Taipei 110, Taiwan

[4]Graduate Institute of Medical Sciences, College of Medicine, Taipei Medical University,

Taipei 110, Taiwan

*Corresponding author

Prof. Enzo Tramontano

E-mail address: tramon@unica.it

Prof. Liang-Tzung Lin

E-mail address: ltlin@tmu.edu.tw

31
32

33
34
35
36

Abstract

Endogenous Retroviruses (ERVs) are integrated into the host DNA as result of ancient germ line infections, majorly by now extinct exogenous retroviruses. Accordingly, vertebrates' genomes contain thousands of ERV copies, providing "fossil" records for the ancestral retroviral diversity and its evolution within the host. Like exogenous retroviruses, ERV proviral sequence consists of *gag, pro-pol*, and *env* genes flanked by long terminal repeats (LTRs). In particular, the characterization of *env* gene changes over time allows both to understand ERVs evolutionary trajectory and to infer their potential role in host physiology, including their pathological involvement. Despite their potential impact on the host, a detailed characterization has been performed for few domesticated *env* genes, while a comprehensive survey of their abundance and diversity among primates' genomes is still lacking. The study primarily focused on ERVs' *env* that are known to be present in hominoid species, given their well-characterized nature and availability in public databases. Hence using these *env* sequences, we reconstructed 32 Env sequences representing the prototypes of these ancestral proteins in Class I, Class II, and Class III HERVs. These reconstructed Envs were then used for similarity search, phylogenetic analysis, and examination of recombination events occurred within primates' genomes that were applied to 55 primate species across the *Catarrhini* and *Platyrrhini* parvorders. Through such comprehensive pipeline, we reconstituted a phylogenetic distribution of ERV based specifically on the *env* genes, showing that the ERVs have been prevalent and widely distributed across the primate lineage. We observed for the first time the presence of the HML (human mouse mammary tumor virus-like) group in the *Platyrrhini* parvorder, indicating initiation of spread of HML supergroup before the split between New World Monkeys (NWM) and Old World Monkeys (OWM) i.e. even before 40 mya. Importantly, we confirmed interclass and intra-class *env* recombination events confirming the phenomenon of "*env* snatching" among primates' ERVs. As a result, we demonstrate that tracing the diversity patterns of ERVs' *env* provides relevant insights into the retroviral evolutionary history of ERVs in *Catarrhini* and *Platyrrhini* parvorders. Overall, our findings reveal that *env* recombination contributes to the diversification of ERVs, thereby broadening our comprehension of retroviral and primate evolution.

Keywords
Endogenous Retroviruses, Envelope Protein, Evolution, Recombination

Introduction

79
80

Retroviruses, discovered over a century ago, are a diverse group of animal viruses characterized by a common replication mechanism (Mager & Stoye, 2015). Their lifecycle involves reverse transcription of a single-stranded RNA (ssRNA) into a double stranded DNA (dsDNA), which integrates into the host genome forming a provirus (Mager & Stoye, 2015). While the retroviruses circulating nowadays typically infect somatic cells, an abundance of ancient retroviruses have been able to integrate their genome into germline cells, becoming endogenous retroviruses (ERVs) and allowing their vertical transmission from one host to another through generations (Sverdlov, 2000). Over the span of millions of years, thousands of ERV copies have been accumulated throughout the vertebrate lineage and can range from complete proviral structures to highly fragmented remnants of the provirus (Hayward, 2017; Mager & Stoye, 2015; Weiss, 2006).

Likewise exogenous RVs, ERVs also consists of four major viral protein-coding genes i.e. *gag, pro-pol* and *env* flanked by identical long-terminal repeats (LTRs) at 5' and 3' ends of the provirus (Hayward, 2017; Johnson, 2019). Among the four viral genes *gag* and *pro-pol* are well conserved across retroviral species while *env* is the most diverse, being responsible for the specific tropism of the infection (Henzy & Johnson, 2013). Given that the reverse transcriptase (RT) domain of the *pol* gene is a highly conserved region, it is widely used to infer phylogenetic relationships and to categorize retroviruses into distinct taxonomic lineages within the Retroviridae family (Henzy & Johnson, 2013). Based on RT sequences, ERVs are classified into three main classes i.e. Class I (Gammaretroviruses and Epsilonretroviruses), Class II (related to Deltaretroviruses, Lentiviruses, Betaretroviruses, and Alpharetroviruses) and Class III (Spumaretroviruses) (Blomberg et al., 2009; Gifford et al., 2018). Various phylogenetic relationships of ERVs are also based on the comparisons between RT and the transmembrane (TM) region of the *env* gene, however discrepancies may

arise due to the diversity of the *env* gene among different ERVs of the same class (Blomberg
et al., 2009). While the analysis of RT sequences identifies the evolutionary lineage that gives
rise to a retroviral species, studying the *env* gene provides insights into the bases of such
retroviral evolution, signaling the emergence of new viruses leading to cross species
transmission.

The ERVs' Envelope (Env) glycoprotein encoded by *env* gene consists of two main
components, i.e. surface (SU) and transmembrane (TM) units, which includes some domains
used for its characterization and phylogenetic analysis: the immunosuppressive domain
(ISD), the fusion peptide (FP), the heptad repeats (HR1 and HR2), and the CX(6)C
motif(Chabukswar et al., 2023). In addition, the most recent ERV group, the HML2, is
divided into type I and type II due to the presence of alternative splicing variants of *env* thus
producing Np9 (type I) and Rec (type II), differing in the presence and absence of a 292
nucleotide (nt) deletion in the *env* gene, respectively (Chabukswar et al., 2022; Heyne et al.,
2015; Subramanian et al., 2011). The divergence in the *env* gene can be accounted by two
main factors i.e. mutations and recombination. Mutations lead to modification in *env* nt
sequence, whereas recombination events involve the transfer of *env* portions across viral
genomes. Recombination occurs between viral genomes present in the same host cell as they
exchange one or more genetic portion(s), potentially leading to the emergence of new viral
strains and viruses, expansion of viral lineages, increase in their virulence and pathogenicity,
and resistance to antiviral drugs (Pérez-Losada et al., 2015). A high degree of mosaicism in
the *env* gene has been previously reported in the form of recombinations or secondary
integrations, and tends to complicate the sequence-based HERV classification (Vargiu et al.,
2016). Similarly, we previously observed the presence of recombinant forms of HML2
elements in Old World Monkeys (OWMs), specifically in *Macaca mulatta* and *Macaca*
fascicularis (Chabukswar et al., 2022).

The analysis of the patterns of *env* diversity is an essential step to better understand the
evolution of ERVs and reconstruct the structure of the ancestral ERV viral genes, which in
turn can help in various functional analysis of the correspondent ancestral proteins, as the
changes in the genes overtime has affected their production and functions. Hence, in the
present study we aimed to reconstruct the *env* gene from the different HERV groups by
generating a representative collection of 32 full-length consensus Envs for Class I, Class II
and Class III. These proteins were used for the screening of representative species of
Catarrhini (*Hominidae, Cercopithecinae* and *Colobinae*) and *Platyrrhini* parvorders,
identifying *env* genes and investigating their diversity by implementing a similarity search

strategy, followed by the study of their phylogeny and recombination patterns. With this
approach, we provided a comprehensive description of the ERV *env* diversity in 55 primates
species, showing the presence of several class II ERVs in the *Platyrrhini* parvorder that have
not been previously reported, and detailing the occurrence of events of interclass and
intraclass *env* recombination that significantly expand our understanding of retroviral
evolution.

Methods

Generation of Representative HERV Envelope protein sequences

The *env* consensus sequences that have been previously reported in Table 4 and Table 6 of
Vargiu et al. 2016 were collected and used as a query to extract the correspondent *env* genes
for each HERV group from human genome (GRCh38/hg38) available on UCSC Genome
Browser through the BLAT search function. To build these consensus sequences, we
followed different approaches, we first extracted a total of 33 HERV elements that have been
previously reported in Table 6 of Vargiu et al.2016, as the ones including the most intact *env*
genes, belonging to the Class I groups HERV9, HERVFC, HERVH, HERVT, HERVW,
HERVE, HERV3 and the Class II groups HML2 (18 out of 33 sequences), HML5, HML6,
and HML8 For the other HERV groups not included in this list, we referred to the Table 4 of
Vargiu et al., 2016, which provides a summary of 39 canonical HERV clades found in the
human genome (GRCh37/hg19). More 190 sequences were extracted from the additional files
of Vargiu et al., 2016 for which we generated a bed file and extracted them with the flanking
of 200 nt at both 3' and 5' ends using the table browser tool in the genome browser
(Supplementary file1 Sheet1 (S1)). These 190 sequences initially extracted were divided into
canonical and non-canonical forms (Vargiu et al., 2016) where canonical refers to sequences
that are well-preserved and match the structural and functional features of complete *env*
genes, and non-canonical refers to sequences with significant structural alterations or
truncations that deviate from this typical form (Vargiu et al., 2016). These sequences were
further aligned group-specifically, to understand if any of the sequence codes for complete
Env proteins. Further, to select some sequences to be used as a reference, we extracted 19

consensus sequences from Dfam (both nt and aa), and 17 reported HERV-Env proteins from

UniProt for which the accession numbers are provided in Supplementary file 1 Sheet2 (S2).

The sequences used to perform alignments for each HERV group are described in detail in

Supplementary file 1 Sheet3 (S3), which were then aligned to references sequences taken

from Dfam, NCBI and UniProt sequences (Supplementary file 1 (S2)).

For each HERV group alignment, we translated the nt sequences in 3 forward frames to

check the presence of stop codons using the Geneious software (http://www.geneious.com).

Furthermore, in order to reconstruct the Env proteins, we translated *env* nt sequence

alignments into all three forward frames in Geneious to identify the translatable regions. The

correct reading frame was identified according to the alignment and comparison with known

Env domain and the high sequence similarity to the reference Env proteins taken from Dfam

and Vargiu et al. 2016. The insertions or deletions (indels) were retained in the initial

alignment but were carefully managed during the reconstruction process. Regions with

ambiguous or excessive indels that compromised the alignment integrity were further masked

or excluded. The correct reading frame was further validated by identifying the conserved

domains characteristic of Env proteins, such as signal peptide, SU, and TM domains using

NCBI CD prediction tool. The frames lacking these hallmark features were excluded from the

reconstruction process. The Env protein consensus sequences were generated for each group

by realigning the extracted translated portions to the reference sequence to generate a

complete representative Env sequence for each HERV group. These generated Env sequences

were then aligned to the exogenous retroviruses of all classes. For this, we collected 52

envelope protein sequences from UniProt database that included different strains as well as

subtypes of exogenous retroviruses from all the genera (i.e. alpha, beta, gamma, delta,

epsilon, intermediate-Beta like, spuma and lentiviruses). The Uniprot accession numbers for

the Envs of the exogenous retroviruses have been provided in Supplementary file 1 (S2).

ERV Env Screening in Primate Genomes

The reconstructed HERV Env sequences were used as a query to analyze 55 primate

genomes, including the representative species from both the *Catarrhini* (*Hominidae,*

Cercopithecinae and *Colobinae* families)* and *Platyrrhini* parvorders. All the genome

assemblies   were   sourced   from   NCBI   Assembly   database

(https://www.ncbi.nlm.nih.gov/assembly/). The screening of the genomes for the *env*

sequences was done by performing the tBLASTn (https://blast.ncbi.nlm.nih.gov/Blast.cgi)

search for all the HERV groups separately. The tBLASTn algorithm uses amino acid

sequences to compare the translated nt database in all six frames, and hence the nt hits

obtained were further refined. The search parameters included a minimum sequence identity

threshold of 40% and an E-value cutoff of 1E-5. Hits shorter than 40% of the complete *env*

gene were excluded from further analysis. To ensure specificity, secondary integrations and

repetitive elements were manually curated and removed. All the primate genome assemblies

that were screened during the tBLASTn search are listed in the supplementary file 2 while the

accession numbers for hits obtained for gamma-like, beta-like and spuma-like are available in

the Supplementary phylogenetic figures 3, 4, and 5.

Sequence Alignments and Phylogenetic Analysis

All the hits obtained from the tBLASTn search were filtered and were downloaded in fasta

format. For each group, sequences whose length was less than 50% of the entire *env* gene

were discarded. We then performed group specific nt sequence alignments using MAFFT

algorithm of Geneious software. These alignments were further used to carry out

phylogenetic analysis with the IQTREE software (http://www.iqtree.org/) for gamma-like,

beta-like and spuma-like *env* nt sequences. The trees were inferred using maximum

likelihood (ML) methods in IQ-TREE with a bootstrap of 1000 replicates. The trees

generated were further visualized and annotated by the Figtree V1.4.3.

software(http://tree.bio.ed.ac.uk/software/figtree/).

Detection of Recombination Events

After the phylogenetic analysis of gamma, beta and spuma-like ERVs, the recombination

events were detected using RDP, Geneconv, Bootscan, Maxchi, Chimaera, Siscan and 3SEQ

methods implemented in RDP4 software (http://web.cbio.uct.ac.za/~darren/rdp.html) (Martin

& Rybicki, 2000). The default settings of the RDP4 software that were used are (i) only

potential recombination events detected by four or more of the above-mentioned methods; (ii)

sequences were treated as linear and (iii) window size of 30 variable nt positions used for the

RDP method. The breakpoint positions and the recombinant sequences indicated for the

potential recombination event were further confirmed through comparative genomics. The

detected recombinant sequences were further used to perform BLAT search in the primate

genomes available in UCSC Genome to confirm the position and location of recombinant

sequence in the host genomes. To highlight the initiation of the recombination events, a

phylogenetic tree of all the genomes of *Catarrhini* parvorder available on the genome

browser was generated using the TimeTree server by simply uploading the list of primate

species on the server. The tree reflects host phylogeny and was based on whole-genome

sequences, except for Gorilla species, since the *Gorilla gorilla gorilla* genome was used to

generate the primate phylogeny. We further annotated the events on the phylogenetic tree

based on the genomes in which the recombinants were detected.

Results

Reconstruction of the Retroviral Envelope from Human ERVs

To understand the ERVs evolutionary dynamics, reconstructing the ancestral ERV prototypes

is a crucial step and requires to address two important issues i.e., the biases present in the

ERV data used and the changes that had occurred after each single retrovirus was integrated

into the host genome. With the purpose of providing prototype Env proteins for the different

HERVs, we reconstructed the Env sequence of 32 HERV groups, including members of

Class I (gamma- and epsilon-like, 20), II (beta-like, 10), and III (spuma-like, 2) (Table1,

Supplementary file1(S4)). To ensure accuracy, all reconstructed sequences were manually

curated to verify the presence of functional regions and sequence integrity. A schematic

representation of the Env reconstruction process is shown in Figure 1.

Since the Env reconstruction was performed based on the data available for HERVs, to

confirm the accuracy of reconstruction by its relatedness to the other RVs, we generated Env

protein phylogeny of reconstructed Envs with 58 Env proteins of exogenous RVs (Figure 2).

As expected, we observed a class specific clustering of the reconstructed Envs with that of

exogenous RVs, except for HERVS which was clustered with Class I RVs and not with the

Class III RVs, indicating that HERVS has a Class I Env that is related to PRIMA41, PABL

and HERV1_Artodact, as also previously reported (Vargiu et al., 2016).

Distribution of ERVs in Primate Genomes

To track the spreading of the above HERV groups across primates' evolution and evaluate

their conservation, we used the above HERV Env consensus sequences to identify the *env*

genes of class I, class II and class III ERVs through a group specific tBLASTn search in a the

genome of 55 primate species. From the tBLASTn similarity search, we obtained 1374 hits

for class I i.e. gamma-like *env*, 873 hits for class II i.e. beta-like *env* and 96 for class III i.e.

spuma-like *env*. For all the three classes, we generated nt alignments and performed

maximum likelihood (ML) phylogenies using IQTREE software with a bootstrap of 1000

replicates.

For the gamma-like *env* 1374 hits the best fit model generated for the phylogeny using the

IQTREE was TVM+F+R4 (Figure 3a, Supplementary file 3). The phylogenetic analysis

revealed group-specific clustering of env genes, primarily within the HERVHF supergroup,

which includes HERVH, HERVFA, HERVFB, and HERVFC. The HERVHF supergroup is

believed to have integrated into the *Catarrhini* lineage around 30−45 million years ago (Li et

al., 2022). Our study also observed a widespread distribution of this supergroup, with most

occurrences in Hominidae genomes, followed by other species. (Table 2, Figure 3a,

Supplementary file 3). Similarly, we also observed a wide distribution of ERVT, ERVW,

ERVE and ERVFRD followed by some small clusters of ERVH48, ERV9 and ERVIP.

Additionally, we noticed distinct occurrences and clusters of HERV15, along with HERVR,

ERVV1, ERVV2 and PABL, despite their exclusion from the blast analysis. Although

HERV15, HERVR, ERVV1, ERVV2 and PABL were excluded from the primary Blast

analysis, we observed distinct occurrence and clusters of these sequences in the subsequent

phylogenetic analysis. This is likely because the Env consensus sequences, even though being

group specific, share regions of homology with the excluded sequences. As a result, the

consensus sequences effectively captured related members of the supergroup, even those not

explicitly included in the initial dataset (Figure 3a). Of note, the consistent group specific

clustering of the reconstructed Env sequences observed in the phylogenetic trees supports the

robustness of the consensus sequences in representing the broader supergroup.

Likewise, phylogenetic analysis was also performed for 873 beta-like *env* hits obtained from

tBLASTn. The best fit model for the phylogeny obtained was TVM+F+I+R3, with the major

grouping obtained for HML2, which was further divided into two subclusters corresponding

to the presence of Np9 (type I) and Rec (type II) *env* splicing variants (Table 2, Figure 3b,

Supplementary file 4). The HML2 group is present in abundance followed by HML8 and

HML10 while smaller clusters were observed for HML1, HML3, HML6 and HML9 (Table 2,

Figure 3b, Supplementary file 4). A very few sequences were obtained in the *Callithrix*

jacchus that belong to *Platyrrhini* parvorder. In addition to the above clusters for HML2 Np9

and Rec, a third cluster of HML2 Rec included sequences that are annotated as ERVK-7,

ERVK-19 and ERVK-25. The tBLASTn search for HML4 (HERVK-13) in the platyrrhini

genomes, we obtained a fewer hits in *Callithrix jacchus,* and hence to confirm the presence of

HML4 group in *Callithrix jacchus* and other Platyrrhini genomes we performed the BLAT

search in two *Platyrrhini* genomes available on UCSC Genome Browser i.e. Marmoset

(*Callithrix jacchus*; Callithrix_jacchus_cj1700_1.1/calJ) and Squirrel monkey (*Saimiri*

boliviensis; Broad/saiBol1). We also performed the BLAT search for other HML supergroup

members which have not been previously reported in the Platyrrhini parvorder.

From the BLAT search, we obtained 6 copies of HML2 annotated as HERVK in the squirrel

monkey genome while no HML2 sequences were detected in *Callithrix jacchus* (Table 3).

We also identified the presence of other members of HERVK supergroup i.e. HML1, HML4,

HML8, HML9 and HML10 in the squirrel monkey genome that has not been reported

previously (Table 3). The sequences identified are in few cases full-length ERV copies while

in most of the cases are of very short length or fragmented. These results indicates two

possible hypotheses for the presence of HML1, HML2, HML4, HML8, HML9 and HML10

in squirrel money: either the onset of these HMLs began even before 30 mya i.e. before the

split between NWM and OWM, that occurred around 43 mya or they got fixed in to the

genome as a results of cross-species transmission of RVs after the split.

While performing tBLASTn search for both HERVL and HERVS, hits were mostly obtained

for gamma-like ERVs, specifically for ERV3-1, ERVFC-1, ERVFRD and ERV-W. These

results were in line with our previous studies which states that HERVL is more similar to

ERV3-1 in hyrax, tenrec, armadillo, alligator and turtle (Bénit et al., 1997; Cordonnier et al.,

1995; Vargiu et al., 2016). The hits were further refined by excluding all the gamma-like

ERVs as well as refined by excluding short elements and removing the sequence of other

repetitive elements acquired by secondary integrations. Hence, we obtained only 96 hits

specific to HERVL and HERVS. An ML phylogenetic tree with the best fit model of

K3Pu+F+R3 was generated with the bootstrap of 1000 replicates (Figure 3c, Supplementary

file 5). The phylogenetic analysis performed for these groups showed that a major

distribution took place in *Homo sapiens* followed by *Macaca mulatta* and other primate

genomes.

Apart from the group specific *env* clustering, we also observed some small clusters hinting

towards the presence of recombinant ERVs, for example, the grouping of HML4 for

Callithrix jacchus with HML2 in the beta-like *env* phylogeny.

Evidence of Recombination in the ERVs' Env

An important factor for the diversity of the *env* gene are recombination events, since

recombination is a ubiquitous process that may give rise to new variants of existing viruses,

possibly also accounting for changes of the original tropism. Hence, we aimed to detect the

potential recombination events in our gamma, beta and spuma-like *env* alignments. To this

purpose, we used RDP4 software, which implements multiple built-in methods to detect

recombination events, which are confirmed when four or more methods yield a positive

signal. The software identified multiple recombination events in the gamma and beta

alignment files generated in this study (Figure 4a). Since the spuma-like *env* alignment

includes only two groups i.e. HERVL and HERVS, and no recombination among them was

detected, the spuma-like *env* were aligned with gamma-like *env,* to check the presence of

recombination among the two classes. The RDP4 software provides an overview of each

recombination event, including the major parent (the sequence contributing the majority of

the recombinant), the minor parent (the sequence contributing smaller portions to the

recombinant), and the breakpoint position (the nt position where the recombination occurs).

Using the detected recombination events and their specific breakpoint positions, we

performed BLAT searches in the UCSC Genome Browser for all gamma-like, beta-like, and

spuma-like *envs*. While performing the BLAT search, we considered all the primate

genomes available in Genome Browser (n=15), also with the purpose of tracing back the

initiation of these recombination events along primates' evolution (Figure 4b). From the

BLAT analysis of gamma-like, beta-like, and spuma-like *env* sequences, we observed a

distinct *env*-LTR recombination pattern. In all cases, the minor parents identified by the RDP

software were LTRs within the *env* sequences.

The recombination events detected in the gamma-like *env* alignment are ERVH-ERVE-
MER11C, ERV15-LTR12, ERV15-MER11A, and PRIMA41-LTR5/LTR18. The first event
was detected in ERVH, in which ERVE was present in the SU region while MER11C that is
an LTR of HML8 group was present in the TM region of the *env* (Figure 4a). This event was
only identified in 4 genomes (Table 4, Supplementary file 6), suggesting a very recent
occurrence (i.e. less than 9 mya) (Figure 4b). In fact, the recombinant sequences were shared
only between 4 species of *Catarrhini* parvorder and were absent in more distantly related
primates (such as gibbons, orangutans, etc.). This observation, together with the presence of
shared recombination sites, suggest that the insertion occurred after the divergence of these
species, while its absence in others indicate that it is a recent recombination event. It can be
said that this recombinant form of ERVH emerged possibly due to the co-packaging of
ERVH, ERVE and HML8 and thus the recombinant form ERVH-ERVE-MER11C was fixed
in the genome of human, chimpanzee, gorilla, and bonobo with the very few copies. Other
recombination events were observed in ERV15 and PRIMA41, in which a LTR was detected
in the TM region of the *env* gene. The ERV15-LTR12/MER11A was detected in 12 primate
genomes while the PRIMA41-LTR15/LTR18 that involves the LTR of ERV15 and ERV3
groups, respectively, was observed in only 6 genomes (Table 4, Figure 4b). Of note, these
two events were present as a single copy in each genome in just one chromosome, as
indicated in supplementary file 6. These single events might have occurred due to the
template switching mechanism during the reverse transcription process and eventually were
fixed in the chromosome.
Similar events were detected in beta-like *env* alignment, where the LTRs were detected in the
TM region of the HML *env* gene. The events detected were HML10-LTR13, HML2-LTR13
and HML2-MER11A/MER11B (Figure 4a). Single chromosomal recombination event was
detected for HML10-LTR13 and HML2-LTR13, where the presence of LTR13, which is the
LTR of the HML*4* group, was detected in the HML10 and HML2 *env* gene in 9 genomes
(Table 4, Figure 4b). Similar to ERV15, these events are also present as single copies in the
primate genomes. Unlike the above mentioned single chromosomal recombination events, the
HML2-MER11A/MER11B recombination event was found in high abundance in the
genomes of species from the *Cercopithecinae* and *Colobinae* subfamilies. However, within
the *Hominidae* family, this recombination was detected in only a single species, *Nomascus*
leucogenys. Especially, abundant number of copies of this recombinant were detected in the
two Macaque genomes i.e. *Macaca fascicularis* and *Macaca mulatta*. The HML8 group
contains three types of LTR i.e. MER11A, MER11B and MER11C (Scognamiglio et al.,

2023), of which the presence of MER11A in HML2 *env* was detected in *Chlorosebus*
sabaeus, Macaca fascicularis, Macaca mulatta, Papio hamadryas, Nasalis larvatus and
Rhinopithecus roxellana while MER11B was detected in *Nomascus leucogenys, Chlorocebus*
sabaeus, and Macaca fascicularis (Table 4, Figure 4b) suggesting that the HML2 template
switching took place with two of the three above different variants of HML8. Since no
evidence of this recombination event was found in most of the species of *Hominidae* except
for *Nomascus leucogenys,* it is likely that it occurred after the split between *Colobinae*
subfamily and *Platyrrhini*, simultaneously with the abundant spread of HML2. All the
recombinations detected in the HML group are hypothesized to be emerged due to the
template switching mechanism as in the gamma-like HERVs and hence were fixed in the
primate genomes.

Lastly for the spuma-like *env,* we detected two recombination events for ERVL i.e. ERVL-
MER11C and ERVL-LTR13 (Table 4, Figure 4a). These two single chromosome
recombination events were detected in six species genomes (Table 4). As mentioned for all
the above recombination events, ERVL seems to have also followed the same mechanism of
template switching for both the LTRs.

Finally, to infer the functional effect of the presence of LTRs in the *env* region of ERVs, we
performed the conserved domain (CD) search in NCBI. We first used the reconstructed Env
to identify the domains present in the *env* and then performed the same search with the
recombinant forms. In Table 5, all the CDs are listed, reporting their position in the *env*
obtained from NCBI for the non-recombinant forms as well as their presence or absence in
the recombinant forms. Of note, a consistent pattern of loss of CD is observed for the ERVs
which have an LTR in the *env* gene. The domains are either completely absent or truncated
due to the recombination with the LTR, leading to the loss of function in the corresponding
protein. As shown in Table 3, the only domain identified for ERVH is TLV coat, but other
domains have been identified for the recombinant form of ERVH-ERVE-MER11A
suggesting an *env* acquisition event for the ERVH group by the co-packaging with ERVE.
From the overall results, it seems that the LTRs of HML8 (MER11A, MER11B and
MER11C) are the most frequently found in the *env* gene of different ERVs and are associated
with recombination in primate genomes.

Discussion

ERVs are "fossils" of the ancient exogenous RVs that have become extinct, that can still
carry out certain functional roles (Sverdlov, 2000). Indeed, the genomic integration of ERVs
and its widespread dispersion has led to the domestication of some ERV proteins for useful
functions. The best known examples of the conservation of an open reading frame (ORF)
linked to the expression in appropriate tissue are syncytin-1 and syncytin-2 encoded by
HERVW (ERVWE1) (Blond et al., 2000; Grandi & Tramontano, 2017) and HERVFRD-1
(ERVFRD-1) (Rote et al., 2004), respectively. Another recurrent ERV domestication is
aimed to its use as a restriction factor and interfering in steps of exogenous RV infection
(Frank & Feschotte, 2017). Available ERV Env studies are mostly focused on finding their
possible role in physiological or pathological conditions, but limited studies have been
performed to understand ERVs' evolutionary dynamics. For this, a focus on how various
factors may affect the Env divergence is required.

The initial step in understanding such ERVs evolutionary dynamics – which encompasses
their origin, proliferation within the host genome, and the mechanisms of inactivation and
elimination – is to precisely reconstruct their viral genes that can also be essential to define
their host-virus interactions. In addition, understanding their residual functions can help both
in elucidating their potential role in various pathological conditions, and allow their
exploitation as antivirals, gene therapy vectors, etc. In this scenario, an Env reconstruction
clearly aids in broadening the understanding and discovering of various aspects of currently
existing ERVs (Johnson, 2019). Hence, with the intention of studying the ERV Envs
divergence, we firstly reconstructed 32 ancestral ERVs Env prototypes that can help in
various functional analysis such as gene regulation and expression, pathogenicity,
evolutionary dynamics, host-ERV interaction, etc. (Johnson, 2019). The screening of 55
primate genomes in *Catarrhini* (*Hominidae, Cercopithecinae, Colobinae*) and *Platyrrhini*
parvorders using the reconstructed Env sequences for each HERV class demonstrated that the
majority of ERV *env* nt sequences hits are obtained in the *Catarrhini* parvorder specifically in
Hominidae Superfamily. In the phylogenetic analyses performed for each Class, we observed
a group-specific clustering of all the ERVs. Over time, the genomic persistence of ERVs has
been accompanied by insertions, deletions, mutations, and even the loss of the *env* gene, all
of which contribute to biases in their phylogenetic analyses. As a result, we report some
mixed clustering among the ERV classes. Apart from the 17 gamma-like *env*, we also
observe hits for ERV15, HERVR, ERVV1, ERVV2 and PABL *env* genes which were
initially not included in the dataset, meaning that the generated consensus sequences
represent very well the members of the same supergroup. Our investigation demonstrates that

the superfamilies HERVHF and HERVW9 of gamma-like *env*, HML2 group of beta-like *env*

integrated in the primate lineage over 30-45 mya, earlier than previously reported (Li et al.,

2022). In fact, several studies focused on the characterization and identification of HERVs in

the primate genome - also reported the presence of the HML supergroup in the *Catarrhini*

parvorder (Grandi et al., 2017, 2021; Scognamiglio et al., 2023) and by far HML5 has been

known to be the most ancient member of the HERVK supergroup (Lavie et al., 2004), as its

presence has been previously reported in marmoset and squirrel monkey. These reports

indicate that their integration began after the split between *Catarrhini* and *Platyrrhini* i.e. ~30

mya, as the HML sequences were not detected in *Callithrix jacchus* genome except for

HML5. However, we identified the presence of members of HERVK supergroup i.e. HML1,

HML2, HML4, HML8, HML9 and HML10 in the squirrel monkey genome that has never

been previously reported, while no HML2 sequences were detected in *Callithrix jacchus*

(Table 3). Thus, we show that the onset of integration of HML1, HML2, HML4, HML8,

HML9 and HML10 initiated in the *Platyrrhini* even before 43 mya, possibly having a lower

rate of amplification as they are less numerous in this parvorder, while they were spread more

abundantly after the split of *Catarrhini* and *Platyrrhini*. Differently from gamma-like and

beta-like ERVs, we detected only 96 hits for spuma-like *env*. the reason for such lower

prevalence of spuma-like ERVs in the genome could be that they either less frequently

integrate into the germ line or might have a shorter persistence in the genome as compared to

gamma-like or beta-like ERVs (Hayward et al., 2015). Overall, the present phylogenomic

approach suggests the widespread presence of the gamma ERVs, indicating that they may

have been integrated in the host genomes through multiple sources, and demonstrates the

onset of class II ERV even before the split between *Catarrhini* and *Platyrrhini* parvorders. A

complex evolutionary trajectory as compared to the one of spuma-like ERVs.

Numerous studies have reported multiple recombination events in ERVs in the vertebrate

lineage leading to several modifications throughout the endogenization process (Chabukswar

et al., 2023). Recombination mechanisms contributing to the observed variation in the ERVs

primarily involve template switching during reverse transcription. Other processes, such as

homologous recombination within the host genome, can also lead to the exchange of genetic

material between ERVs. While cross-species transmission and *env* acquisition or degradation

are critical processes in shaping ERV evolution and diversity, they are not direct mechanism

of recombination but rather represent the outcome or influences on recombination

(Chabukswar et al., 2023; Pérez-Losada et al., 2015). Such events can greatly increase the

diversity among the ERVs and complicate their classification. Hence, to detect the presence

of such recombination events in primates we performed recombination analysis and identified

8 potential recombination events i.e. ERVH-ERVE-MER11C, ERV-15-LTR12, ERV-15-

MER11A, HML2-LTR13, HML2-MER11A, HML2-MER11B, HML10-LTR13, ERVL-

MER11C and ERVL-LTR13 in gamma-like and beta-like and spuma-like *env* sequences

(Table 4). We confirmed these events using the genome assemblies available in Genome

Browser, hence assuming the presence of these recombinants in the other primate genomes of

the superfamily.

The potential mechanism due to which such events occur in the genomes is the co-packaging

of two different ERVs or an ERV and an exogenous RV in a virus particle, followed by the

template switch between the two genomes. In the template switching mechanism the two RV

copies are co-packaged into the virus particle, and the template switch occurs as the RT

dissociates from one template to another resulting in intermolecular i.e. homologous or

nonhomologous recombination or intramolecular i.e. deletions or insertions (Delviks-

Frankenberry et al., 2011). As in the case of the recombinants detected in the present study, a

template switch could have occurred between the highly conserved TM region of the *env* of

one ERV and the LTR of the other ERV due to sequence similarities, resulting in the

observed recombination of an LTR inserted into the TM region of the *env* gene. According to

the mechanism of template switching, the LTR found within the *env* gene has the features of

a viral RNA genome LTR (i.e. not with the complete U5-R-U3 structure that characterizes

the proviral ones), as indeed it is observed in the sequences. Additionally, it is possible that

the observed recombinant events arose from the co-packaging from ERV and a related

exogenous RV, especially if the latters were still actively circulating at the time when the

events occurred. Such interactions could have facilitated inter-viral recombination, leading to

the integration of recombinant sequences into the host genome. A prominent example of such

event is the recombination between exogenous MuLv and its endogenous form leading to the

generation of recombinant polytrophic MuLVs that utilize a different cell surface receptor for

their entry, altering the host range and increasing their pathogenic potential (Alamgir et al.,

2005; Chabukswar et al., 2023; Evans et al., 2009, Williams et al., 2024).

As in case of ERVH-ERVE-MER11C, ERVH seems to acquire the *env* gene of the ERVE

group, leading to the generation of a recombinant ERV that retains the potentially coding

ERVE Env, while for the rest of the recombinant sequences an LTR exchange with the TM

region of the *env gene* is observed, eventually leading to loss of the *env* function, further

promoting the intragenomic spreading of *env*-less ERVs (Magiorkinis et al., 2012).

Interestingly, such swapping of the *env* region can also possibly lead to the emergence of

different variants of the existing RV, as in case of HML2-MER11A/MER11B, where it is
predicted that different HML8 variants swapped their LTR with the TM region of the HML2
env, and thus spreading in abundance in the *Cercopithecinae* subfamily members and
creating diversity in the HML2 group. The recombination between HML2-MER11A/B in the
macaque genomes was previously reported (Chabukswar et al., 2022; Williams et al., 2024),
suggesting that the HML8 region appeared in HML2 *env* region due to the recombination
between HML2 and HML8 virus due to co-packaging of HML2 and HML8 RNAs during the
reverse transcription and has been experimentally verified by Williams et al., 2024. Similar to
HML2-MER11A/B recombination, various recombinant events have been reported in this
study for gamma-, beta- and spuma-like Envs, having the LTRs in the TM region of the *env*.
Hence, as suggested, it is worth considering that these events can have implications in the
biology of these viruses (Williams et al., 2024). These kind of events have also been
previously reported for Harlequin which have a complicated structure of LTR2-HERVE-
MER57I-LTR8-MER4I-HERVI-HERVE-LTR2 (Vargiu et al., 2016). The recombination
events described in the present study provide more insights of the prevalence of recombinant
ERV forms within the host genome, underlying the role of recombination as a driving force
behind the ERVs diversification.

Conclusion

ERVs are the remnants of ancestral exogenous RVs whose origin and diversity are still not
completely understood and are a great tool to study the coevolution of host and RVs. The
present study provides a phylogenomic perspective of the retroviral *env* gene diversity. With
this study approach we provide the evidence for inter- as well as intra-group recombination
events. Present results demonstrate the prevalence of ERVs across the primate lineage as well
as the diversification of the retroviral Env through the process of recombination over the
evolutionary timescale. Overall, this analysis gives further knowledge in studying the host-
virus interactions on a larger scale.

567
References

Alamgir, A. S. M., Owens, N., Lavignon, M., Malik, F., & Evans, L. H. (2005). Precise
Identification of Endogenous Proviruses of NFS/N Mice Participating in Recombination

with Moloney Ecotropic Murine Leukemia Virus (MuLV) To Generate Polytropic
MuLVs. *Journal of Virology*, *79*(8). https://doi.org/10.1128/jvi.79.8.4664-4671.2005
Bénit, L., De Parseval, N., Casella, J. F., Callebaut, I., Cordonnier, A., & Heidmann, T.
(1997). Cloning of a new murine endogenous retrovirus, MuERV-L, with strong
similarity to the human HERV-L element and with a gag coding sequence closely
related to the Fv1 restriction gene. *Journal of Virology*, *71*(7).
https://doi.org/10.1128/jvi.71.7.5652-5657.1997
Blomberg, J., Benachenhou, F., Blikstad, V., Sperber, G., & Mayer, J. (2009). Classification
and nomenclature of endogenous retroviral sequences (ERVs). Problems and
recommendations. In *Gene* (Vol. 448, Issue 2).
https://doi.org/10.1016/j.gene.2009.06.007
Blond, J.-L., Lavillette, D., Cheynet, V., Bouton, O., Oriol, G., Chapel-Fernandes, S.,
Mandrand, B., Mallet, F., & Cosset, F.-L. (2000). An Envelope Glycoprotein of the
Human Endogenous Retrovirus HERV-W Is Expressed in the Human Placenta and
Fuses Cells Expressing the Type D Mammalian Retrovirus Receptor. *Journal of*
Virology, *74*(7). https://doi.org/10.1128/jvi.74.7.3321-3329.2000
Chabukswar, S., Grandi, N., Lin, L. T., & Tramontano, E. (2023). Envelope Recombination:
A Major Driver in Shaping Retroviral Diversification and Evolution within the Host
Genome. In *Viruses* (Vol. 15, Issue 9). https://doi.org/10.3390/v15091856
Chabukswar, S., Grandi, N., & Tramontano, E. (2022). Prolonged activity of HERV-
K(HML2) in Old World Monkeys accounts for recent integrations and novel
recombinant variants. *Frontiers in Microbiology*, *13*.
https://doi.org/10.3389/fmicb.2022.1040792
Cordonnier, A., Casella, J. F., & Heidmann, T. (1995). Isolation of novel human endogenous
retrovirus-like elements with foamy virus-related pol sequence. *Journal of Virology*,
69(9). https://doi.org/10.1128/jvi.69.9.5890-5897.1995
Delviks-Frankenberry, K., Galli, A., Nikolaitchik, O., Mens, H., Pathak, V. K., & Hu, W. S.
(2011). Mechanisms and factors that influence high frequency retroviral recombination.
Viruses, *3*(9). https://doi.org/10.3390/v3091650
Evans, L. H., Alamgir, A. S. M., Owens, N., Weber, N., Virtaneva, K., Barbian, K., Babar,
A., Malik, F., & Rosenke, K. (2009). Mobilization of Endogenous Retroviruses in Mice
after Infection with an Exogenous Retrovirus. *Journal of Virology*, *83*(6).
https://doi.org/10.1128/jvi.01926-08
Frank, J. A., & Feschotte, C. (2017). Co-option of endogenous viral sequences for host cell
function. In *Current Opinion in Virology* (Vol. 25).
https://doi.org/10.1016/j.coviro.2017.07.021
Gifford, R. J., Blomberg, J., Coffin, J. M., Fan, H., Heidmann, T., Mayer, J., Stoye, J.,
Tristem, M., & Johnson, W. E. (2018). Nomenclature for endogenous retrovirus (ERV)
loci. In *Retrovirology* (Vol. 15, Issue 1). https://doi.org/10.1186/s12977-018-0442-1
Grandi, N., Cadeddu, M., Pisano, M. P., Esposito, F., Blomberg, J., & Tramontano, E.
(2017). Identification of a novel HERV-K(HML10): Comprehensive characterization
and comparative analysis in non-human primates provide insights about HML10
proviruses structure and diffusion. *Mobile DNA*, *8*(1), 1–18.
https://doi.org/10.1186/s13100-017-0099-7
Grandi, N., Pisano, M. P., Pessiu, E., Scognamiglio, S., & Tramontano, E. (2021).
HERV□K(HML7) integrations in the human genome: Comprehensive characterization
and comparative analysis in non□human primates. *Biology*, *10*(5).
https://doi.org/10.3390/biology10050439
Grandi, N., & Tramontano, E. (2017). Type W human endogenous retrovirus (HERV-W)
integrations and their mobilization by L1 machinery: Contribution to the human

transcriptome and impact on the host physiopathology. *Viruses*, *9*(7), 1–37.
https://doi.org/10.3390/v9070162

Hayward, A. (2017). Origin of the retroviruses: when, where, and how? In *Current Opinion*
in Virology (Vol. 25). https://doi.org/10.1016/j.coviro.2017.06.006

Hayward, A., Cornwallis, C. K., & Jern, P. (2015). Pan-vertebrate comparative genomics
unmasks retrovirus macroevolution. *Proceedings of the National Academy of Sciences of*
the United States of America, *112*(2). https://doi.org/10.1073/pnas.1414980112

Henzy, J. E., & Johnson, W. E. (2013). Pushing the endogenous envelope. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *368*(1626).
https://doi.org/10.1098/rstb.2012.0506

Heyne, K., Kölsch, K., Bruand, M., Kremmer, E., Grässer, F. A., Mayer, J., & Roemer, K.
(2015). Np9, a cellular protein of retroviral ancestry restricted to human, chimpanzee
and gorilla, binds and regulates ubiquitin ligase MDM2. *Cell Cycle*, *14*(16), 2619–2633.
https://doi.org/10.1080/15384101.2015.1064565

Johnson, W. E. (2015). Endogenous Retroviruses in the Genomics Era. In *Annual Review of*
Virology (Vol. 2). https://doi.org/10.1146/annurev-virology-100114-054945

Johnson, W. E. (2019). Origins and evolutionary consequences of ancient endogenous
retroviruses. In *Nature Reviews Microbiology* (Vol. 17, Issue 6).
https://doi.org/10.1038/s41579-019-0189-2

Katoh, K., Rozewicki, J., & Yamada, K. D. (2018). MAFFT online service: Multiple
sequence alignment, interactive sequence choice and visualization. *Briefings in*
Bioinformatics, *20*(4). https://doi.org/10.1093/bib/bbx108

Lavie, L., Medstrand, P., Schempp, W., Meese, E., & Mayer, J. (2004). Human Endogenous
Retrovirus Family HERV-K(HML-5): Status, Evolution, and Reconstruction of an
Ancient Betaretrovirus in the Human Genome. *Journal of Virology*, *78*(16).
https://doi.org/10.1128/jvi.78.16.8788-8798.2004

Li, Y., Zhang, G., & Cui, J. (2022). Origin and Deep Evolution of Human Endogenous
Retroviruses in Pan-Primates. *Viruses*, *14*(7). https://doi.org/10.3390/v14071370

Mager, D. L., & Stoye, J. P. (2015). Mammalian Endogenous Retroviruses. *Microbiology*
Spectrum, *3*(1). https://doi.org/10.1128/microbiolspec.mdna3-0009-2014

Magiorkinis, G., Gifford, R. J., Katzourakis, A., De Ranter, J., & Belshaw, R. (2012). Env-
less endogenous retroviruses are genomic superspreaders. *Proceedings of the National*
Academy of Sciences of the United States of America, *109*(19), 7385–7390.
https://doi.org/10.1073/pnas.1200913109

Martin, D., & Rybicki, E. (2000). RDP: Detection of recombination amongst aligned
sequences. *Bioinformatics*, *16*(6). https://doi.org/10.1093/bioinformatics/16.6.562

Pancino, G., Ellerbrok, H., Sitbon, M., & Sonigo, P. (1994). Conserved framework of
envelope glycoproteins among lentiviruses. In *Current Topics in Microbiology and*
Immunology (Vol. 188). https://doi.org/10.1007/978-3-642-78536-8_5

Pérez-Losada, M., Arenas, M., Galán, J. C., Palero, F., & González-Candelas, F. (2015).
Recombination in viruses: Mechanisms, methods of study, and evolutionary
consequences. In *Infection, Genetics and Evolution* (Vol. 30).
https://doi.org/10.1016/j.meegid.2014.12.022

Rote, N. S., Chakrabarti, S., & Stetzer, B. P. (2004). The role of human endogenous
retroviruses in trophoblast differentiation and placental development. *Placenta*, *25*(8–9).
https://doi.org/10.1016/j.placenta.2004.02.008

Scognamiglio, S., Grandi, N., Pessiu, E., & Tramontano, E. (2023). Identification,
comprehensive characterization, and comparative genomics of the HERV-K(HML8)
integrations in the human genome. *Virus Research*, *323*.
https://doi.org/10.1016/j.virusres.2022.198976

Subramanian, R. P., Wildschutte, J. H., Russo, C., & Coffin, J. M. (2011). Identification,
characterization, and comparative genomic distribution of the HERV-K (HML-2) group
of human endogenous retroviruses. *Retrovirology*, *8*, 90. https://doi.org/10.1186/1742-
4690-8-90
Sverdlov, E. D. (2000). Retroviruses and primate evolution. In *BioEssays* (Vol. 22, Issue 2).
https://doi.org/10.1002/(SICI)1521-1878(200002)22:2<161::AID-BIES7>3.0.CO;2-X
Vargiu, L., Rodriguez-Tomé, P., Sperber, G. O., Cadeddu, M., Grandi, N., Blikstad, V.,
Tramontano, E., & Blomberg, J. (2016). Classification and characterization of human
endogenous retroviruses mosaic forms are common. *Retrovirology*, *13*(1), 1–29.
https://doi.org/10.1186/s12977-015-0232-y
Weiss, R. A. (2006). The discovery of endogenous retroviruses. In *Retrovirology* (Vol. 3).
https://doi.org/10.1186/1742-4690-3-67
Weiss, R. A. (2016). Human endogenous retroviruses: Friend or foe? In *APMIS* (Vol. 124,
Issues 1–2). https://doi.org/10.1111/apm.12476
Williams, Z. H., Imedio, A. D., Gaucherand, L., Lee, D. C., Mostafa, S. M., Phelan, J. P.,
Coffin, J. M., & Johnson, W. E. (2024). Recombinant origin and interspecies
transmission of a HERV-K(HML-2)-related primate retrovirus with a novel RNA
transport element. *ELife*, *13*. https://doi.org/10.7554/eLife.80216
690

692 **Figure legends**

Figure 1: Illustration of the reconstruction process of HERV Env proteins. The first step was to extract *env* genes for both endogenous and exogenous retroviruses from Literature, Dfam, NCBI, Uniprot, and PDB (complete *env* sequence as well as defective *env* genes). Second, we performed group specific alignments for all the ERV sequences with Mafft and further divided the sequences into subtypes based on the alignments, as in case of HERV3 and HML3. Then, we translated the nt sequences into three forward frames in the alignment and extracted only the translated portions that did not have any stop codons. Lastly, the extracted aa portions were aligned to the reference sequence of each group and hence, a reconstructed env sequence was generated for HERV groups as mentioned in Table 1.

Figure 2: Phylogeny of the reconstructed Envs with the envelope protein of exogenous retroviruses. The phylogenetic tree was generated for the reconstructed HERV Env proteins with the 69 exogenous envelope proteins for Class I, Class II and Class III retroviruses. The class I exogenous and endogenous retroviruses are highlighted in green, class II in yellow and class III in orange color. The reconstructed Envs are presented in red color in the tree. The tree also contains RVs of other class with which no ERVs were clustered and hence are not highlighted.

Figure 3: Phylogenetic analysis of the envelope hits obtained from the tBLASTn analysis for the three classes i.e. Class I (gamma-like *env*), Class II (beta-like *env*) and Class III (spuma-like *env*). The trees were generated using IQ-TREE software with the bootstrap of 1000 replicates. (a)Phylogeny for all the hits obtained for Class I ERVs with the best fit model of TVM+F+R4. (b) Phylogeny for all the hits obtained for Class II ERVs with the best fit model of TVM+F+I+R3 and (c) Phylogeny for all the hits obtained for Class III ERVs with the best fit model of K3Pu+F+R3. All the groups obtained from the tBLASTn hits and further clustered together with the phylogenetic analysis are labelled in the tree. Similar trees are provided as supplementary files with the tip labels having all the annotations and the accession numbers for all the three classes (Supplementary file 2,3 and 4).

Figure 4: Detecting the recombination events in the ERV's *env* gene. (a) Schematic representation of *env* recombination events in ERVs. The diagram indicates the location of the LTRs in the recombinant *env* gene, with the coordinates provided in supplementary file 5. (b) Phylogenetic tree illustrating the initiation of recombination events in the primate lineage. This tree was generated based on the available *Catarrhini* genomes in the Genome Browser using TimeTree server. Since the genome used by the server was *Gorilla gorilla gorilla* hence is marked with *. The presence of recombination events was tested using BLAT searches in the Genome Browser to trace the timing of their initiation and hence annotated on the phylogenetic tree based on the genomes the recombinants were detected.

Tables

746 **Table1:** Dataset of the reconstructed HERV Envelope proteins

	Supergroup	Group	Subtypes	aa Length
Class I	MLLV	HERVT		623
	HERVERI	HERVE		671
		HERV3	Subtype1	666
			Subtype2	661
		HERV1		777
		HERVI		728
	HERVW9	HERVW		544
		HERV9		554
	HERVIADP	HERVIP		780
		HERVADP		672
	HERVHF	HERVH		581
		HERVH48		244
		HERVFA		561
		HERVFB		542
		HERVFC		570
	HERVFRDlike	HERVFRD		599
	HUERSP	HUERSP3		536
Class II	HERVK	HML1		667
		HML2	Type1(Np9)	588
			Type2(Rec)	684
		HML3	Subtype1	557
			Subtype2	465
			Subtype3	390
		HML4		657
		HML5		644
		HML6		639
		HML7		664
		HML8		776
		HML9		682
		HML10		625
Class III	HSERVIII	HERVL		460
		HERVS		420

760 **Table2:** Widespread Distribution of ERV *env* genes across primate genomes

Class	Supergroup	Name	Primate distribution
Class I (Gammaretroviruses)	MLLV	HERVT	*Homo sapiens, Pongo abelii, Gorilla gorilla, Pan troglodytes, Macaca nemestrina, Macaca fascicularis, Macaca thibetana, Papio anubis, Rhinopithecus bieti, Aotus nancymaae, Cebus imitator*
	HERVERI	HERV3	*Homo sapiens, Pan paniscus, Pan troglodytes, Pongo abelii, Pongo pygmaeus, Nomascus leucogenys, Macaca fascicularis, Macaca mulatta, Macaca thibetana, Cercocebus atys,* *Theropithecus gelada, Piliocolobus tephrosceles, Rhinopithecus roxellana, Rhinopithecus bieti*
		HERVE	*Homo sapiens, Gorilla gorilla, Pongo pygmaeus, Hylobates agilis, Macaca thibetana, Macaca nemestrina, Trachypithecus francoisi*
		HERV1	*Homo sapiens, Pan troglodytes*
	HERVW9	HERVW	*Homo sapiens, Gorilla gorilla, Pongo pygmaeus, Pongo abelii, Nomascus leucogenys, Hylobates pileatus,* *Macaca fascicularis, Macaca mulatta, Macaca thibetana, Papio anubis, Papio hamadryas, Chlorocebus sabaeus, Cercopithecus hamlyni, Colobus guereza, Cercocebus torquatus, Rhinopithecus roxellana, Rhinopithecus bieti, Presbytis francoisi, Symphalangus syndactylus, Cebus imitator, Aotus nancymaae, Callithrix jacchus*
		HERV9	*Homo sapiens, Nomascus leucogenys, Macaca fascicularis, Mandrillus leucophaeus*
	HERVADP	HERVIP	*Macaca nemestrina, Papio anubis, Cercopithecus aethiops*
	HERVHF	HERVH	*Homo sapiens, Gorilla gorilla, Macaca mulatta, Rhinopithecus roxellana*
		HERVH48	*Homo sapiens, Pan paniscus, Pan troglodytes, Pongo abelii, Gorilla gorilla, Hylobates moloch, Nomascus leucogenys, Cebus imitator, Callithrix jacchus*
		HERVFA	*Homo sapiens, Pan paniscus, Pan troglodytes*
		HERVFB, HERVFC	*Homo sapiens, Pan paniscus, Pan troglodytes, Macaca fascicularis, Macaca mulatta, Macaca nemestrina, Macaca thibetana, Papio anubis, Papio cynocephalus, Chlorocebus sabaeus, Rhinopithecus bieti, Piliocolobus tephrosceles, Callithrix jacchus, Aotus*

			nancymaae, Saimiri boliviensis
	HERVFRD-like	HERVFRD	*Homo sapiens, Pan troglodytes, Pongo abelii, Gorilla gorilla, Nomascus leucogenys, Rhinopithecus roxellana, Rhinopithecus bieti, Trachypithecus francoisi, Piliocolobus tephrosceles, Colobus angolensis, Saimiri boliviensis, Cebus imitator, Callithrix jacchus, Aotus nancymaae*
	Other clusters	HERV15	*Homo sapiens, Pan troglodytes, Pongo abelii, Pongo pygmaeus, Gorilla gorilla, Hylobates agilis, Macaca fascicularis, Macaca mulatta, Macaca nemestrina, Macaca thibetana, Macaca fuscata*
		HERVR	*Homo sapiens, Pan troglodytes, Pongo pygmaeus, Hylobates agilis, Macaca fuscata*
		ERVV1, ERVV2	*Homo sapiens, Macaca fascicularis, Macaca thibetana, Macaca mulatta,* *Papio anubis,Chlorocebus sabaeus, Chlorocebus aethiops, Mandrillus leucophaeus,Colobus angolensis palliatus, Theropithecus gelada, Semnopithecus entellus, Rhinopithecus roxellana, Cebus imitator, Saimiri boliviensis, Callithrix jacchus, Saguinus oedipus*
Class II (Betaretroviruses)	HML	HML1	*Homo sapiens, Callithrix jacchus*
		HML2_Np9	*Homo sapiens, Gorilla gorilla, Gorilla gorilla gorilla, Pan troglodytes, Pan paniscus*
		HML2_Rec	*Homo_sapiens, Gorilla gorilla, Nomascus leucogenys, Hylobates moloch, Pan troglodytes, Pan paniscus, Macaca nemestrina, Macaca mulatta, Macaca fascicularis, Papio anubis, Chlorocebus sabaeus,Trachypithecus francoisi, Rhinopithecus roxellana, Rhinopithecus bieti*
		HML3	*Homo sapiens, Pan troglodytes, Macaca mulatta, Macaca thibetana, Macaca nemestrina, Trachypithecus francoisi*
		HML4	*Homo sapiens, Macaca nemestrina, Callithrix jacchus*
		HML6	*Homo sapiens, Nomascus leucogenys, Piliocolobus tephrosceles*
		HML8	*Homo sapiens, Gorilla gorilla gorilla, Pan paniscus, Pan troglodytes, Macaca nemestrina, Piliocolobus tephrosceles*
		HML9	*Homo sapiens, Pan paniscus, Pan troglodytes, Macaca mulatta*
		HML10	*Homo sapiens ,Papio hamadryas* *Macaca tonkeana, Macaca maura, Macaca brunnescens, Macaca fascicularis, Macaca ochreata, Macaca nigra, Macaca nigrescens, Macaca cyclopis, Macaca silenus, Macaca mulatta, Macaca*

			fascicularis, Macaca hecki, Macaca fuscata, Macaca radiata, Macaca thibetana, Macaca sinica, Macaca assamensis, Macaca arctoides, Macaca sylvanus, Theropithecus gelada, Mandrillus sphinx, Presbytis melalophos, Trachypithecus cristatus
	Other Clusters	ERVK-7, ERVK-19, ERVK-25	*Macaca_fascicularis, Macaca mulatta, Macaca thibetana, Papio anubis, Chlorocebus sabaeus, Cercocebus atys, Rhinopithecus roxellana, Rhinopithecus bieti, Trachypithecus francoisi, Piliocolobus tephrosceles*
Class III(Spumaretroviruses)	HSERVIII	HERVL	*Homo sapiens, Pan troglodytes, Pan paniscus, Pongo abeli, Macaca mulatta, Callithrix jacchus,Saimiri boliviensis boliviensis*
		HERVS	*Homo sapiens, Pan paniscus, Pan troglodytes, Pongo pygmaeus, Macaca mulatta,Cercocebus atys, Piliocolobus tephrosceles, Aotus nancymaae*

761
762
763
764
765 **Table3:** Coordinates of HML supergroup in *Platyrrhini* (Squirrel Monkey Genome)

SUPERGROUP	HML	COORDINATES	STRAND	LENGTH
HERVK	HML1	JH378338:275937-278501	+	2564
		JH378138:15705974-15707323	-	1349
		JH378108:32563432-32571361	+	7929
		JH378224:2469781-2470931	-	1150
		JH378134:11672169-11673824	-	1655
	HML2	JH378134:10458979-10463351	-	4372
		JH378105:56691102-56691325	-	223
		JH378105:56692918-56693866	-	948
		JH378105:56695565-56696715	-	1150
		JH378105:56697213-56697652	-	439
		JH378138:15703105-15705969	-	2864
	HML4	JH378161:520931-521612	-	681
		JH378134:16151196-16152689	+	1493
		JH378134:7690429-7691554	-	1125
		JH378146:13346796-13353682	-	6886
		JH378137:7471310-7471738	-	428
		JH378106:35578339-35578392	+	53
	HML8	JH378141:3704815-3715095	+	10280*
		JH378112:24913201-24913256	-	55
	HML9	JH378336:397006-400978	-	3972
		JH378132:8543270-8546474	+	3204

		JH378113:32170092-32175530	-	5438
		JH378152:4554178-4561789	-	7611
		JH378362:166905-169335	-	2430
		JH378344:454424-459278	-	4854
		JH378332:381766-384506	-	2790
	HML10	JH378114:8677151-8678160	-	1009
		JH378114:15213478-15214880	+	1402
		JH378130:3582370-3583730	+	1360
		JH378233:1994071-1995016	-	945
		JH378110:37554414-37555367	-	953

Note: * indicates the truncated sequence

Table4: Summary of Recombination Events in Gamma-like, Beta-like, and Spuma-like Env Genes Across Primate Genomes

Recombination Event	ERV Group	Species	Observations
ERVH-ERVE-MER11C	Gamma-like	*Homo sapiens* (8), *Pan paniscus* (4), *Pan troglodytes* (6), *Gorilla gorilla* (5)	Detected only in Catarrhini species; absent in gibbons and orangutans. Suggests recent recombination.
ERV15-LTR12/MER11A	Gamma-like	*Homo sapiens, Pan troglodytes, Gorilla gorilla, Pongo abelii, Nomascus leucogenys, Chlorocebus sabaeus,* and others	Single copy per genome
PRIMA41-LTR15/LTR18	Gamma-like	*Homo sapiens, Pan paniscus, Pan troglodytes, Gorilla gorilla, Pongo abelii*	Detected in single chromosomes; involves LTRs of ERV15 and ERV3 groups.
HML10-LTR13	Beta-like	*Homo sapiens, Pan paniscus, Pan troglodytes, Gorilla gorilla, Pongo abelii, Nomascus leucogenys, Chlorocebus sabaeus, Macaca fascicularis*	Single chromosomal event.
HML2-LTR13	Beta-like	*Homo sapiens, Pan paniscus, Pan troglodytes, Gorilla gorilla, Pongo abelii, Nomascus leucogenys, Chlorocebus sabaeus, Macaca fascicularis*	Single chromosomal event.
HML2-	Beta-like	*Nomascus*	Abundant copies

MER11A/MER11B		*leucogenys* (10), *Chlorocebus sabaeus* (17), *Macaca fascicularis* (33), *Macaca mulatta* (75), *Papio anubis* (18), *Nasalis larvatus* (6)	found; co-packaging with two HML8 variants (MER11A and MER11B).
ERVL-MER11C	Spuma-like	*Chlorocebus sabaeus, Macaca fascicularis, Macaca mulatta, Papio hamadryas, Nasalis larvatus, Rhinopithecus roxellana*	Single chromosomal event.
ERVL-LTR13	Spuma-like	*Homo sapiens, Pan paniscus, Pan troglodytes, Gorilla gorilla, Pongo abelii*	Single chromosomal event.

774 **Table5**: Detection of the conserved domains in the reference ERV sequences as well as recombinant sequences.

Recombinant ERV	Conserved Domains Features								
	HERV-K_REC Super family	HERVK_env_2	TLV Coat	Ebola-like_HR1-HR2 Superfamily					
				HR1-GP1 interface	HR2	Homotrimer interface	C1 binding site	CX(6,7)C Motif	Immuno-suppressive region
*ERV15	-	-	1064-1258	1439-1552	1631-1669	1439-1669	1517-1528	1571-1598	1521-1569
ERV15-LTR12	-	-	-	√	-	√*	-	-	√
ERV15-MER11A	-	-	-	-	-	-	-	-	-
*ERVH	-	-	825-1487	-	-	-	-	-	-
ERVH-ERVE-MER11C	-	-	-	√	-	√	√	√	√
*PRIMA41	-	-	-	-	-	-	-	-	929-955
PRIMA41-LTR5/LTR18	-	-	-	-	-	-	-	-	-
*HML2	55-315	346-849	-	1501-1659	1753-1878	1507-1878	-	-	-
HML2-LTR13	-	-	-	-	-	-	-	-	-
HML2-MER11A/MER11B	√	√		√	-	√*	-	-	-
*HML10		157-636	-	1339-1437	-	1339-1497	-	-	1421-1447
HML10-LTR13		-	-	-	-	-	-	-	-
*ERVL	-	-	455-1051	716-814	908-934	716-934	-	-	797-823
ERVL-MER11C	-	-	-	-	-	-	-	-	-
ERVL-LTR13	-	-	-	-	-	-	-	-	-

775 Note: √ indicates presence of the Domain; √* indicates the domain is truncated; - indicates absence of the domain; * indicates the non-
776 recombinant ERV

Supplementary Files

Supplementary File1: Excel sheet for all the HERVs used for the Reconstruction of HERVs' Env (S1, S2 and S3). The S4 sheet contains Fasta File of gamma-like, beta-like and spuma-like reconstructed HERV Envelope sequences.

Supplementary File2: List of the primates' genomes and the genome aseemblies that were screened during the tBLASTn search for screening the ERVs' Env

Supplementary File3: Complete phylogenetic tree with the tip labels for Gamma-like ERVs' env.

Supplementary File4: Complete phylogenetic tree with the tip labels for Beta-like ERVs' env.

Supplementary File5: Complete phylogenetic tree with the tip labels for Spuma-like ERVs' env.

Supplementary File6: Excel file for the Coordinates of the recombinant sequences in the 15 genome assemblies extracted from the BLAT search.

Data availability statement

The original contributions presented in the study are included in the article/Supplementary material, further inquiries can be directed to the corresponding author.

Author contributions

SC performed the analysis and wrote the manuscript. ES participated in performing the analysis. NG supervised the analyses, checked the data, and participated in editing the manuscript. ET and LTL conceived and coordinated the study. All authors contributed to the article and approved the submitted version.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Collecting Available Data for *env* of Human ERVs

Literature

UniProt

Dfam

Defective *env* genes

Complete *env* genes

Group specific nucleotide alignments

Sequences translated in to forward frames in the alignment

Extracting the translated portions without the stop codons

| SP | SU | FP | TM | |

A representative HERV Env Consensus sequence

(a)

(b)

(c)

(a)

(b)