Catecholamines reduce choice history biases in perceptual decision making

Jan Willem de Gee[1,2,3,*], Niels A. Kloosterman[1,4,5,6], Anke Braun[1,7] & Tobias H. Donner[1,*]

[1] Department of Neurophysiology and Pathophysiology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany.
[2] Cognitive and Systems Neuroscience, Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam, NLD;
[3] Amsterdam Brain & Cognition, University of Amsterdam, Amsterdam, The Netherlands.
[4] Department of Psychology, University of Lübeck, Germany
[5] Center of Brain, Behavior and Metabolism, University of Lübeck, Germany
[6] Max Planck Institute for Human Development, Berlin, Germany
[7] Department of Psychiatry and Neurosciences, Charité - Universitätsmedizin Berlin, Berlin, Germany.

*Address correspondence to Jan Willem de Gee & Tobias H. Donner.

Email: j.w.degee@uva.nl, t.donner@uke.edu

Author Contributions: Conceptualization: JWG, NK, TD; Funding Acquisition: TD; Investigation: JWG, NK; Data Curation: JWG, NK; Formal analysis: JWG, NK, AB; Writing—original draft: JWG, TD; Writing—review and editing: JWG, NK, AB, TD; Supervision: TD; Project Administration: TD.

Competing Interest Statement: The authors declare no competing interests.

Abstract

Theoretical accounts postulate that the catecholaminergic neuromodulator noradrenaline shapes cognition and behavior by reducing the impact of prior expectations on learning, inference, and decision-making. A ubiquitous effect of dynamic priors on perceptual decisions under uncertainty is choice history bias: the tendency to systematically repeat, or alternate, previous choices, even when stimulus categories are presented in a random sequence. Here, we directly test for a causal impact of catecholamines on these priors. We pharmacologically elevated catecholamine levels in human participants through the application of the noradrenaline reuptake inhibitor atomoxetine. We quantified the resulting changes in observers' history biases in a visual perceptual decision task. Choice history biases in this task were highly idiosyncratic, tending toward choice repetition or alternation in different individuals. Atomoxetine decreased these biases (toward either repetition or alternation) compared to placebo. Behavioral modeling indicates that this bias reduction was due to a reduced bias in the accumulation of sensory evidence, rather than of the starting point of the accumulation process. Atomoxetine had no significant effect on other behavioral measures tested, including response time and choice accuracy. Atomoxetine and variations of pupil-linked arousal at slower and faster timescales had analogous effects on choice history bias. We conclude that catecholamines reduce the impact of a specific form of prior on perceptual decisions.

Introduction

Biases are prevalent in human decisions ranging from those based on abstract reasoning to judgments of the sensory environment[1–3]. One prominent source of such biases in perceptual decisions across mammalian species is the history of previous experiences[4–14]. When, as is common in natural environments, the environmental state evaluated is relatively stable, such history biases reflect adaptive expectations that are dynamically updated over time and improve performance[5,15,16]. For example, one might choose to forage for deer in the same area where previous expeditions were successful. While recent work in rodents and primates has identified correlates of such history biases in neural population activity of parietal and frontal cortical areas[9,12,17–21], little is currently known about the neurotransmitter systems involved in this important form of bias.

Influential theoretical work postulates that the catecholaminergic neuromodulator noradrenaline shapes behavior by reducing the impact of prior expectations on learning, inference, and decision-making[22,23]. The catecholaminergic (and in particular, the noradrenaline) systems of the brainstem project to large parts of the cerebral cortex, where the neuromodulators they release change the functional properties of their target networks[24–29]. Consequently, they are in an ideal position to shape cortex-wide network activity underlying decision-making in a coordinated fashion. Indeed, pupil responses (a peripheral proxy of central arousal state[30]) point to a role of neuromodulation in regulating the impact of these history-dependent priors[8]. However, pupil responses reflect the activity of multiple neuromodulatory brainstem systems[30–32], and the above result builds on correlative approaches. Direct evidence for the role of catecholaminergic neuromodulation in choice history biases is currently sparse in animal models[13,33] and absent in humans.

Here, we aimed to provide causal evidence for the role of catecholamines in regulating (suppressing) history biases in human choice behavior. Specifically, we hypothesized that catecholamines down-regulate the impact of previous experiences on decision-making, which should translate into a reduction of choice history bias. We tested this hypothesis by pharmacologically increasing central catecholamine (in particular noradrenaline) levels through the selective noradrenaline re-uptake inhibitor atomoxetine[34,35] and quantified the resulting behavioral changes in a variant of a visual perceptual decision task adopted from recent animal physiology[36]. Atomoxetine inhibits the presynaptic norepinephrine transporter, preventing the reuptake of norepinephrine throughout the brain along with inhibiting the reuptake of dopamine in specific brain regions such as the prefrontal cortex[34,35]. The administration of atomoxetine at the dosage used in the current study (40 mg) produces robust central effects on several neurophysiological markers of the state of cortical networks: suppression of low-frequency spectral power[37] as well as changes in long-range temporal correlations of local amplitude envelopes[37], in inter-area correlations during task[38] and in signatures of evidence accumulation[39,40]. We found that atomoxetine reduced choice history biases by specifically interacting with the way people accumulate noisy perceptual evidence.

Results

Atomoxetine was administered orally (40 mg per session) in a within-subject, double-blind, placebo-controlled, and randomized design (**Figure 1A**). The behavioral task entailed the presentation of visual targets (Gabor patch of fixed orientation and varying contrast), one in the left and one in the right visual hemifield; participants were asked to report the location the larger contrast target by button press (**Figure 1B**; Methods). To increase sensory uncertainty and promote integration of evidence across time[41], we added dynamic visual noise to both Gabor patches (**Figure 1B**; Methods). There were two different signal strengths (small and large difference in contrast); auditory feedback was presented 50 ms right after participant's choices. We collected many trials per participant (range, 3520-4320) across four experimental sessions, in which we systematically varied and counterbalanced drug condition and stimulus-response mapping (**Figure 1A** and **Figure S1**; Methods).

99
Figure 1. Experimental design and effects on peripheral arousal markers. (A) Design was within-subject,
double-blinded, randomized, and placebo-controlled (Methods). Additionally, the stimulus-response mapping
*was counterbalanced within participants (**Figure S1**; Methods). **(B)** Schematic sequence of events during the*
spatial contrast discrimination task (Methods). Participants reported the location of the highest contrast Gabor
*patch (difference in contrast is high for illustration only). **(C)** Heart rate measured as the average across task*
blocks (Methods), separately for placebo and atomoxetine sessions. Every connecting line is a participant;
large data points in the middle are the group averages (error bars, s.e.m. across 19 participants); stats, paired-
*sample t-test. **(D)** As C, but for pupil size measured as the average across task blocks (Methods).*
https://doi.org/10.5281/zenodo.16779024.
109
110

Atomoxetine administration at this relatively low dose had a measurable effect on the peripheral
arousal markers heart rate (**Figure 1C**) and pupil size (**Figure 1D**). Pupil size is a measure of
central arousal[30]. Non luminance-mediated changes of pupil diameter are causally related to the
activity of neuromodulatory centers in the brainstem, including the noradrenergic locus
coeruleus[31,42–44]. There was no statistically significant correlation between the individual drug
effects on pupil size and on heart rate (**Figure S1B**), in line with the idea that pupil size reflects
central arousal more closely than heart rate, a measure of peripheral arousal.
118
Atomoxetine did not significantly affect objective task performance. While signal strength lawfully
decreased reaction time (RT; $F_{1,18}$ = 21.7, p < 0.001; two-way repeated measures ANOVA) and
increased accuracy ($F_{1,18}$ = 381.0, p < 0.001) (**Figure 2A,B**), atomoxetine changed neither of those
behavioral parameters (RT: $F_{1,18}$ = 2.1, p = 0.160; accuracy: $F_{1,18}$ = 2.8, p = 0.110; no significant
interaction of signal strength and drug; **Figure 2A,B** and **Figure S2A**). Likewise, atomoxetine did
not significantly change signal detection theoretic sensitivity (d'; **Figure S2B**).
125
By contrast, atomoxetine had a robust impact on the impact of history-based (i.e., dynamically
varying throughout the course of the experiment) prior expectations on their decisions. We
operationalized these time-varying prior expectations as the "repetition probability", the probability
of repeating the choice from the previous trial, irrespective of a potential overall bias for the left or
right choice options (Methods). The stimulus repetition probability was, by design, close to 0.5
(**Figure 2C**) (group average: 0.496; Methods), thus not explaining the sequential effects in
participants' behavior. Atomoxetine reduced the mean choice repetition probability (**Figure 2D**; $F_{1,18}$
= 9.0, p = 0.008; two-way repeated measures ANOVA). Signal strength had no such effect ($F_{1,18}$ =
0.8, p = 0.381) and signal strength and drug did not interact ($F_{1,27}$ ~ 0.0, p = 0.850). Qualitatively,
we observed the same pattern after correct or incorrect decisions (**Figure S2C,D**), and there was
no significant difference between the drug effect on repetition probability after correct and error
trials (t = 0.22, p = 0.828, BF = 0.243).
138
In the placebo sessions, most participants (15 out of 19) had a repetition probability larger than 0.5,
that is, they tended to repeat their choices (**Figure 2C**). We, therefore, wondered if (i) atomoxetine
reduced the group-level choice history biases by stereotypically pushing behavior more toward
alternation (i.e., reduction of repetition probability, in line with previous pupillometry work[8]) or (ii)
through a reduction of both repetition and alternation biases (shift of repetition probability toward
0.5). Our results are in line with the second scenario. There was a robust correlation between
individual choice history biases during placebo sessions and the shift in history biases caused by
atomoxetine: participants with the strongest repetition biases exhibited the strongest shift towards
a repetition probability of 0.5 (**Figure 2E**). Further, our data are consistent with the idea that

atomoxetine also reduced alternation biases, although the latter were only observed in five
participants and weak (**Figure S2E**, data points on x-axis < 0.5). Consequently, atomoxetine
yielded an even more consistent and robust reduction of the absolute choice history bias, defined
as the absolute value of P(repeat)−0.5 (**Figure 2F**; $F_{1,18}$ = 11.0, p = 0.008; two-way repeated
measures ANOVA). Atomoxetine did not change overall choice bias, quantified as signal detection
theoretic criterion (**Figure S2F-H**). These findings show that atomoxetine specifically reduced the
impact of history-dependent prior expectation in all participants, regardless of whether they
idiosyncratically tended to repeat (i.e., assuming a relatively stable environment) or alternate
(assuming a volatile environment).

157
158

159
Figure 2. Atomoxetine reduces choice history bias. (A) Reaction time, separately for placebo and
atomoxetine sessions, and separately or weak and strong signals. Every connecting line is a participant; large
data points in the middle are the group averages (error bars, s.e.m. across 19 participants); stats, main effect
of drug in two-way repeated measures ANOVA. See Figure S2A for reaction time distributions. (B) As A, but
for accuracy. (C) Probability of repeating the previous stimulus category (left) or choice (right) during placebo
sessions. (D) As A, but for repetition probability (Methods). (E) Individual shift in repetition probability caused
by atomoxetine, plotted against individual's repetition probability during the placebo sessions. Data points are
individual participants; error bars, 68% confidence intervals across 5K bootstraps; stats, Pearson's correlation
coefficient (corrected for reversion to the mean; Figure S2E; Methods). (F) As A, but for absolute repetition
probability (Methods). https://doi.org/10.5281/zenodo.16779024.

170
171
How did participants' expectations shape their decision processes, and how did atomoxetine act
on this mechanism? Current models of perceptual decision-making posit the temporal
accumulation of sensory evidence, resulting in an internal decision variable that grows with
time[41,45–47]. When this decision variable reaches one of two decision bounds, a choice is made, and
the corresponding motor response is initiated. In this framework (**Figure 3A**), a bias can arise in
two ways: (i) by shifting the starting point of accumulation toward one of the bounds or (ii) by
changing the rate of evidence accumulation toward one choice alternative.

179
Previous work has shown that individual biases in choice repetition probabilities are generally
explained by a history-dependent bias in the drift rather than starting point, across a range of
different perceptual choice tasks[4]. We replicated this finding for the placebo condition of the present
dataset (**Figure S3A**). To this end, we fitted an accumulation-to-bound model of decision-making
to the behavioral data. The model had five free parameters: 1) boundary separation (controlling
response caution); 2) the mean drift rate (overall efficiency of evidence accumulation); 3) non-

decision time (the speed of pre-decisional evidence encoding and post-decisional translation of
choice into motor response); 4) the starting point of the decision; 5) an evidence-independent bias
in the drift (henceforth called "drift bias"). We fitted all five parameters separately per drug condition;
additionally, we fitted drift rate separately for both signal strength conditions, and both starting point
and drift bias separately for the previous choice category ('left' vs 'right'). The fitted model (**Figure
S3C**) accounted well for the overall behavior in each task (**Figure S3D,E**). Indeed, atomoxetine
also specifically reduced the history-dependent shift in drift bias, but not the history shift in starting
point (**Figure 3B**). Furthermore, this drift bias (not starting point) effect was robustly and specifically
correlated to the individual change of choice repetition probability (**Figure 3C**). Across subjects,
the history-dependent shift in starting point did not predict a drug effect therein (r = -0.34, p = 0.673;
corrected for reversion to the mean), and the same was true for shifts in drift bias (r = -0.60, p =
0.233).

Figure 3. Atomoxetine reduces choice history bias in evidence accumulation. (A) *Schematic of drift
diffusion model accounting for choices, and their associated RTs. In the equation, v is the drift rate. Purple
and cyan curves show expected RT distributions under shifts in either the 'starting point' (z; purple) or 'drift
bias' (v_{bias}; cyan). **(B)** Left: history-dependent shift in starting point bias (Methods), separately for placebo and
atomoxetine sessions. Right, as left, but for drift bias. Every connecting line is a participant; large data points
in the middle are the group averages (error bars, s.e.m. across 19 participants); stats, paired-sample t-test.
(C) Left: individual shift in repetition probability caused by atomoxetine, plotted against individual's history-
dependent shift in starting point bias. 'History shift' is the difference between a parameter estimate for previous
'left' and previous 'right' choices, irrespective of the category of the current stimulus. Right: as left, but for
history-dependent shift in drift point bias. Data points are individual participants; stats, Pearson's correlation
coefficient (difference in correlation assessed with permutation test; Methods).*
https://doi.org/10.5281/zenodo.16779024.

Previous studies linked choice history biases to evoked pupil responses during perceptual
decisions[8], a likely correlate of the task-evoked responses of locus coeruleus (potentially other
brainstem nuclei) during decision-making[24,32,33]. We, therefore, also evaluated the relationship
between choice history bias and the pupil measurements obtained in the placebo sessions of the
current study. Because our fast-paced task design was not suited to isolate trial-to-trial variations
of pre-trial baseline pupil size from the task-evoked responses (Methods), we focused this analysis
exclusively on the pupil responses. Building on previous work finding the decision-related
component of task-evoked pupil responses to be precisely locked to participants' behavioral
choice[8,32,48,49], we analyzed the pupil responses time-locked to the choice report. This showed an
early constriction likely related to the appearance of the two salient choice targets of relatively high
baseline contrast[50,51]. This response was followed by two successive dilations, whereby the first
peaked shortly after the choice report and the second followed about 530 ms later (**Figure 4A**). We
interpret the first peak to reflect the decision-related response[8,32,48] and the second peak to be
driven by the auditory feedback[52,53], which in our current task was triggered by the registration of
the button press (Methods). Accordingly, we focused our analyses of behavioral correlates of pupil
responses on the first peak (gray window in **Figure 4A**) and completed this by a time-variant
analysis (**Figure S4A**).

We found a negative correlation between the amplitude of the decision-related pupil response
component and the probability of repeating the previous choice (**Figure 4B; Figure S4A**). Likewise,

the decision-related pupil response amplitude predicted a reduction of choice repetition bias on the
next trial (**Figure S4B**), replicating previous findings[8]. Both effects mimic the effect of atomoxetine
shown in **Figure 2**, in line with the notion that a boost of central arousal and catecholamine levels
at both slow, session-related (pharmacology) and fast, trial-related (pupil responses) timescales
have consistent effects on choice history bias. Additionally, task-evoked pupil responses were
negatively related to accuracy and quadratically related to RT (**Figure S4B**).

In a final analysis, we tested if the individual effect of atomoxetine on pupil size, measured as the
average across task blocks (**Figure 1D**), predicted the individual effect of atomoxetine on choice
history bias. Participants with a larger increase in pupil size after taking atomoxetine also exhibited
larger reductions in repetition probability (**Figure 4C**). This pattern was not statistically significant
for heart rate (**Figure S4C**). In sum, we here found analogous and correlated effects on choice
history bias of pharmacological catecholamine enhancement (via atomoxetine) and of slow and
fast variations of pupil size, a marker of central arousal state.

250
Figure 4. Pupil-linked arousal reduces choice history bias. (A) Grand-average pupil response time-locked
to report (button press) during placebo. Gray window, interval for averaging task-evoked pupil response on
single trials (Methods). (B) Relationship between task-evoked pupil response and repetition probability (8 bins)
during placebo. Stats, mixed linear modeling (Methods). See Figure S4A for time-wise regression and Figure
S4B for other behavioral measures. (C) Individual shift repetition probability caused by atomoxetine, plotted
against individual shift in pupil size caused by atomoxetine. Pupil size measured as the average across task
blocks (Methods), separately for placebo and atomoxetine sessions; data points are individual participants;
error bars, 68% confidence intervals across 5K bootstraps; *stats, Pearson's correlation coefficient. Panels A*
& B: shading or error bars, s.e.m. across 19 participants. https://doi.org/10.5281/zenodo.16779024*.*

Discussion

Using selective pharmacological interventions in humans, we here imply catecholamines in the
modulation of human choice history biases: boosting catecholamines levels through atomoxetine
specifically reduced the magnitude of choice history biases. Catecholamines, and noradrenaline in
particular, have previously been implicated in modulating learning rates[54–57] and facilitating set
shifting[58] in dynamic environments. Here, we use a selective pharmacological intervention to
provide causal evidence for their role in controlling the weight of historical information in perceptual
decisions. We interpret our findings in the context of theories postulating a noradrenergic down-
weighting of prior expectations (which, we assume, drive the choice history biases analyzed here)
in combination with new sensory evidence in the evaluation of current environmental state[22,23,59].
In the current task, participants' history biases were likely due to an implicit belief that the task
environment was relatively stable, i.e., that stimuli presented on successive trials are correlated.
Our finding that atomoxetine lowered this bias is thus in line with the above theoretical
frameworks[22,23,59]. Our findings are also consistent with the idea that participants attended to one
side of the screen for several successive trials, producing choice "streaks". Thus, the drug effects
may also reflect reduced hysteresis of attentional selection. While this scenario is not mutually
exclusive with the above theoretical accounts, future research should develop approaches
specifically tailored to isolating the dynamics of attentional selection (and associated drug effects)
in choice history biases.

Our finding of a causal effect of catecholamine levels on history bias is consistent with correlative
evidence from pupillometry, obtained in previous work[8,60–64] and replicated here. In dynamic
environments, pupil responses are associated with violations of learned, top-down expectations
and reduce their weight[60–64]. In stable environments, pupil responses predict a reduction of choice
repetition probability[8]. There are, however, also some subtle but notable differences between our
current atomoxetine effects and pupil-linked effects on behavior. First, one study found the pupil-
predicted reduction of repetition probability to be a general increase in the tendency to alternate[8],
while we found atomoxetine to reduce history biases regardless of their sign (i.e., promoting a shift
of repetition probability toward 0.5), which is more consistent with a reduction in prior weight.
Second, our pharmacological intervention did not affect participant's overall choice bias (bias
towards one choice alternative, irrespective of the immediate history), even though this is another
commonly reported behavioral correlate of pupil responses[32,48,65–67]. Third, in our current dataset,
pupil responses exhibited a monotonic (negative) relationship to accuracy, without any concomitant
accuracy effect of atomoxetine. One possible explanation for these differences is that atomoxetine
likely increased both tonic and phasic catecholamines levels[68], while task-evoked pupil responses
reflect phasic neuromodulatory activity[32]. Another possibility is that while atomoxetine selectively
elevates central catecholamine levels (noradrenaline and dopamine), pupil responses track not
only noradrenaline[31,32,42,43,69,70], but also acetylcholine[31,71], serotonin[72] and orexin[73] levels. Even so,
our current analyses of pharmacological and pupil-related effects of the same data set indicate
overall analogous effects of the variations in catecholaminergic tone at different timescales on
choice history biases.

Which neuromodulator specifically mediated the effects reported here? Atomoxetine increases the
extracellular levels of noradrenaline, but also of (prefrontal) dopamine (Methods). Indeed, one
recent report suggest that striatal dopamine may play a role in adapting choice history bias to
temporal regularities in a stimulus sequence[13]. Thus, our results are consistent with a role of one
or both catecholamines. However, the noradrenergic locus coeruleus and dopaminergic midbrain
exhibit a close interplay[74,75] and are co-activated with pupil responses during decisions[32]. Thus, it
could well be that their (complex) interaction regulates choice history bias in perceptual decisions.

In sum, pharmacologically elevating central catecholamine (specifically noradrenaline) levels in
human participants specifically reduces their propensity to accumulate noisy evidence for
perceptual decisions in a manner that is biased by previously reported perceptual experience. This
supports computational theories according to which noradrenaline reduces the impact of dynamic
priors on inference and decision-making.

Materials and Methods

Participants
Nineteen participants (14 female, age range 20-26) participated in the experiment, entailing
concurrent pupillometry and MEG recordings. All had normal or corrected to normal vision and no
history or indications of psychological or neurological disorders. All participants gave written
informed consent. The experiment was approved by the ethics committee of the University Medical
Center Hamburg-Eppendorf (approval number, PV4843) and conducted in accordance with the
Declaration of Helsinki. Participants participated in four experimental sessions, in which we crossed
drug condition (atomoxetine vs. placebo) and stimulus-response mapping (press ipsi vs. contra)
(about 2 hours per session). Participants were paid 10€ per hour. One participant completed three
recording sessions.

Pharmacological intervention
We used the selective noradrenaline reuptake inhibitor atomoxetine (dose, 40 mg) to boost the
levels of catecholamines, specifically noradrenaline and (in the prefrontal cortex) dopamine[34,35].
Atomoxetine is a relatively selective inhibitor of the noradrenaline transporter, which is responsible
for the natural reuptake of noradrenaline that has been released into the extracellular space.
Consequently, atomoxetine acts to increase the extracellular levels of noradrenaline, an effect that

has been confirmed experimentally in rat prefrontal cortex[34]. The same study showed that
atomoxetine also increases the prefrontal levels of dopamine, which has a molecular structure very
similar to the one of noradrenaline and is, in fact, a direct precursor of noradrenaline. Atomoxetine
has smaller affinity to the serotonin transporter, and there are discrepant reports about the
quantitative relevance of these effects: while one study found no increases in serotonin levels under
atomoxetine[34], a recent study reports a significant atomoxetine-related occupancy of the serotonin
transporter in nonhuman primates[76] at dosages that would correspond to human dosages of 1.0–
1.8 mg/kg. Note that these dosages are substantially higher than the administered dosage in this
study (40 mg, independent of body weight). It is therefore unclear to what extent our atomoxetine
condition affected cortical serotonin levels. A mannitol-aerosil mixture was administered as
placebo. All substances were encapsulated identically to render them visually indistinguishable.
Peak plasma concentrations are reached about 1–2 hours after administration for atomoxetine, and
the half-life is about 5 hours[77]. Thus, participants received either a placebo or atomoxetine 1.5
hours before the start of the experimental session.
352
Behavioral task
Each trial consisted of three consecutive intervals (**Figure 1A**): (i) the baseline interval (uniformly
distributed between 0.6 and 1.4 s) containing a dynamic noise pattern on both the left and right
side of the green fixation dot; (ii) the decision interval (terminated by the participant's response;
max 3 s) containing the same noise patterns with Gabor patch embedded in each; (iii) the feedback
interval (uniformly distributed between 0.25 and 1.4 s). Each block consisted of 160 trials, with mini-
breaks after every 40 trials.
360
Two dynamic noise patterns were presented throughout the experiment. The luminance across all
pixels was kept constant. This pedestal noise pattern had 20% contrast. In the decision interval two
Gabor patches (sinusoidal gratings; 2 cycles per degree; phase randomly drifted left or right) were
added to each noise pattern. The Gabor patches symmetrically differed from 80% contrast. On half
of trials, the Gabor patch on right side of the screen was of higher contrast (signal, right); on the
other half the Gabor patch on the left side was of higher contrast (signal, left). Signal location was
randomly selected on each trial, under the constraint that it would occur on 50% of the trials within
each block of 160 trials. Stimuli were presented within two circular patches to the left and right of
the central fixation cross (diameter, 15 degrees of visual angle; eccentricity, 12 degrees).
370
Participants were instructed to report the location of the signal, the highest contrast Gabor patch
by pressing one of two response buttons with their left or right index finger as soon as they felt
sufficiently confident ("free response paradigm"). We did not give specific instructions to tradeoff
accuracy for speed (or vice versa). The mapping between perceptual choice and button press (e.g.,
"right" –> press right key; "left" –> press left key) was counterbalanced across the four recording
sessions. Participants received auditory feedback (high tone, correct; low tone, error) 50 ms after
they indicated their choice.
378
379 Signal strength, the difference in contrast between the left and right signals, was individually titrated
380 before the main experiment to two difficulty levels that yielded about 70% and 85% correct choices,
381 using an adaptive staircase procedure (Quest)[78]. In the placebo sessions, we measured a mean
382 accuracy of 74.07 % correct (± 0.97 % s.e.m.) and 80.83 % correct (± 1.12 % s.e.m), for weak and
383 strong signal strengths respectively.
384
Stimuli were back-projected on a transparent screen using a Sanyo PCL-XP51 projector with a
resolution of 800x600 at 24 Hz. The screen was positioned 65 cm away from their eyes. The
luminance profile was linearized by measuring and correcting for the systems gamma curve. A
doubling of contrast values therefore also produced a doubling of luminance differences. During
the first training session stimuli were presented on a VIEWPixx monitor with the same resolution
and refresh rate (also linearized). To minimize any effect of light on pupil diameter, the overall
luminance of the screen was held constant throughout the experiment.
392

393 Participants performed between 22 and 27 blocks (distributed over four sessions) yielding a total
394 of 3520–4320 trials per participant. One participant performed a total of 17 blocks (distributed over
395 three scanning sessions), yielding a total of 2720 trials.
396
Data acquisition
Recordings took place in a dimly lit magnetically shielded room. Magnetoencephalography (MEG)
was recorded using a whole-head CTF 275 MEG system (CTF Systems Inc., Canada) at a
sampling rate of 1200 Hz. In addition, eye movements and pupil diameter were recorded with an
MEG-compatible EyeLink 1000 Long Range Mount system (SR Research, Osgoode, ON, Canada).
Finally, electrocardiogram (ECG) as well as vertical and horizontal electrooculogram (EOG) were
acquired using Ag/AgCl electrodes at a sampling rate of 1200 Hz. MEG data will be the focus of a
different report.
405
406 *Analysis of electrocardiogram data*
407 ECG data were used to analyze average heart rate. We used an adaptive threshold to detect the
408 R-peak of each QRS-complex in the ECG. Heart rate was then computed by dividing the total
409 number of R-components by time. We quantified block-wise heart rate as the average heart rate
410 across entire task blocks.
411
412 *Analysis of pupil data*
413 All analyses were performed using custom-made Python scripts, unless stated otherwise.
414
Preprocessing. Periods of blinks and saccades were detected using the manufacturer's standard
algorithms with default settings. The remaining data analyses were performed using custom-made
Python scripts. We applied to each pupil timeseries (i) linear interpolation of missing data due to
blinks or other reasons (interpolation time window, from 150 ms before until 150 ms after missing
data), (ii) low-pass filtering (third-order Butterworth, cut-off: 6 Hz), (iii) removal of pupil responses
to blinks and to saccades, by first estimating these responses by means of deconvolution and then
removing them from the pupil time series by means of multiple linear regression[79], and (iv)
conversion to units of modulation (percent signal change) around the mean of the pupil time series
from each measurement session.
424
425 *Quantification of block-wise pupil size.* We quantified block-wise pupil size as the average blink-
426 interpolated pupil size (in raw Eyelink units) across entire task blocks.
427
428 *Quantification of task-evoked pupil responses.* We quantified task-evoked pupil responses for each
429 trial as the mean of the pupil size (in units of percent signal change) in the window 0 s to 0.65 s
430 from choice (gray window in **Figure 4A**), minus the pre-trial pupil size, measured as the mean pupil
431 size during the 0.25 s before trial onset.
432
Analysis and modeling of choice behavior
All analyses were performed using custom-made Python scripts, unless stated otherwise. Trials in
which participants failed to respond within 3 seconds were excluded from the analyses.
Additionally, we excluded trials for which no previous choice was available: (i) previous trial too
slow, (ii) first trial of each block, and (iii) first trial after each mini-break. We always computed
behavioral metrics separately for each participant and drug condition.
439
440 *Overt behavior.* Reaction time (RT) was defined as the time from stimulus onset until the button
441 press. Repetition probability was defined as the fraction of choices that was the same as the choice
442 on the previous trial, irrespective of a potential overall side-bias:
443
444 Eq. 1

445
$$P(repeat) \ = \frac{P(R_n = left | R_{n-1} = left) + P(R_n = right | R_{n-1} = right)}{2}$$

446
447 where *R* is the response (left or right) and *n* is trial number.

448
449 *Signal-detection theoretic modeling.* The signal detection[80] metrics sensitivity (d') and criterion (c)
450 were computed separately for each of the pupil bins. We estimated d' as the difference between z-
451 scores of hit-rates and false-alarm rates. We estimated criterion by averaging the z-scores of hit-
452 rates and false-alarm rates and multiplying the result by -1.
453
454 *Drift diffusion modeling.* We fitted the reaction time data (separately per session) with the drift
455 diffusion model[45,47], based on continuous maximum likelihood using the Python-package PyDDM[81].
456 The decision dynamics were described by (**Figure 3A**):
457
458 $$\Delta y = (s \cdot v + v_{bias}) \cdot \Delta t + cdW \qquad\qquad \text{Eq. 2}$$
459
460 where y is the decision variable, s is the stimulus category (-1 for left signals; 1 for right signals), v
461 is the drift rate and controls the overall efficiency of evidence accumulation, v_{bias} is the drift bias
462 which is and evidence-independent constant added to the drift, and cdW is Gaussian distributed
463 white noise with mean 0 and variance $c^2 \Delta t$. The starting point of evidence accumulation z was
464 defined as a fraction of the boundary separation:
465
466 $$y(0) = z \cdot a \qquad\qquad \text{Eq. 3}$$
467
468 where a is the separation between the two decision bounds. Evidence accumulation terminated at
469 0 ("left") or a ("right). The model was fitted separately for each recording session of each participant.
470 In each fit, we let drift rate vary with signal strength[47], and starting point and drift bias with previous
471 choice[4].
472
473 *Statistical comparisons*
474 We used a paired sample t-test to test for significant differences in heart rate, pupil size, and
475 behavioral measures between placebo and drug sessions (**Figure 1** and **Figure 3**). We used 2 ×
476 2 repeated measures ANOVA to test for the main effects of drug and signal strength, as well as
477 their interaction, on behavioral measures (**Figure 2**). All tests were performed two-tailed.
478
We performed permutation analyses to obtain the distribution of correlation coefficients predicted
exclusively by regression to the mean (**Figure 2E**). This was done by computing the above-
described across-subject correlation 10K, each time using randomly assigned "placebo" vs. "drug"
labels of each participant. We then compared our observed correlation coefficient
(reflecting the combined effects of regression to the mean and the relationship between overall
bias and the atomoxetine predicted shift therein) against this permutation distribution. This analysis
suggested that the relationship between choice history bias and the atomoxetine predicted shift
therein was stronger than expected based on regression to the mean (proportion of
permutations below observed correlation < 0.05).
488
489 We also used a permutation procedure to test for differences between the correlation coefficients
490 for history shifts in starting point bias and drift bias (**Figure 3C**). Again, we permuted the condition
491 labels (drug / placebo).
492
We used a mixed linear modeling approach implemented in the Python-package 'Statsmodels'[82] to
quantify the dependence of several metrics of overt behavior on pupil response bin. Our approach
was analogous to sequential polynomial regression analysis[83], but now performed within a mixed
linear modeling framework. We fitted two mixed models to test whether pupil response bin
predominantly exhibited a monotonic effect (first-order) or a non-monotonic effect (second-order)
on the behavioral metric of interest (y). The fixed effects were specified as:
499
500 $$\text{Model 1: } y \sim \beta_0 1 + \beta_1 P$$
501 $$\text{Model 2: } y \sim \beta_0 1 + \beta_1 P + \beta_2 P^2$$
502

with β as regression coefficients and P as the task-evoked pupil response bin number (demeaned).
We included random intercepts (intercepts could vary with participant). The mixed models were
fitted through restricted maximum likelihood estimation. Each model was then sequentially tested
in a serial hierarchical analysis, which was based on BIC. This analysis was performed for the
complete sample at once, and it tested whether adding the next higher order model yielded a
significantly better description of the response than the respective lower order model (difference in
BIC > 10). We tested models from the first-order (constant, no effect of pupil response) up to the
second-order (quadratic, non-monotonic).

Data and code availability
All resources, including data and code used for the analyses on this paper (in Python), are publicly
available at https://doi.org/10.5281/zenodo.16779024.

Acknowledgments

Funding: J.W. de Gee is supported by the Dutch Research Council (NWO; grant number
VI.Veni.232.210). T.H. Donner is supported by a Project grant of Amsterdam Brain & Cognition
Priority Program and by the German Research Foundation (DFG; grant number Heisenberg
Professorship DO 1240/3-1, and SFB 936, Projects A2/A3, A7, Z3). The funders had no role in
study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Fischer, J. & Whitney, D. Serial dependence in visual perception. *Nat Neurosci* **17**, 738–743 (2014).
2. Tversky, A. & Kahneman, D. Judgment under Uncertainty: Heuristics and Biases. *Science (New York, N.Y.)* **185**, 1124–1131 (1974).
3. Kloosterman, N. A. *et al.* Humans strategically shift decision bias by flexibly adjusting sensory evidence accumulation. *eLife* **8**, e37321 (2019).
4. Urai, A. E., de Gee, J. W., Tsetsos, K. & Donner, T. H. Choice history biases subsequent evidence accumulation. *eLife* **8**, e46331 (2019).
5. Braun, A., Urai, A. E. & Donner, T. H. Adaptive History Biases Result from Confidence-Weighted Accumulation of past Choices. *J. Neurosci.* **38**, 2418–2429 (2018).
6. Hermoso-Mendizabal, A. *et al.* Response outcomes gate the impact of expectations on perceptual decisions. *Nat Commun* **11**, 1057 (2020).
7. Abrahamyan, A., Silva, L. L., Dakin, S. C., Carandini, M. & Gardner, J. L. Adaptable history biases in human perceptual decisions. *PNAS* **113**, E3548–E3557 (2016).
8. Urai, A. E., Braun, A. & Donner, T. H. Pupil-linked arousal is driven by decision uncertainty and alters serial choice bias. *Nat Commun* **8**, 14637 (2017).
9. Urai, A. E. & Donner, T. H. Persistent activity in human parietal cortex mediates perceptual choice repetition bias. *Nat Commun* **13**, 6015 (2022).
10. Gold, J. I., Law, C.-T., Connolly, P. & Bennur, S. The Relative Influences of Priors and Sensory Evidence on an Oculomotor Decision Variable During Perceptual Learning. *Journal of Neurophysiology* **100**, 2653–2668 (2008).
11. Busse, L. *et al.* The Detection of Visual Contrast in the Behaving Mouse. *J. Neurosci.* **31**, 11351–11361 (2011).
12. Braun, A. & Donner, T. H. Adaptive biasing of action-selective cortical build-up activity by stimulus history. *Elife* **12**, RP86740 (2023).
13. Fritsche, M. *et al.* Temporal regularities shape perceptual decisions and striatal dopamine signals. *Nat Commun* **15**, 7093 (2024).
14. Akaishi, R., Umeda, K., Nagase, A. & Sakai, K. Autonomous Mechanism of Internal Choice Estimate Underlies Decision Inertia. *Neuron* **81**, 195–206 (2014).
15. Yu, A. J. & Cohen, J. D. Sequential effects: Superstition or rational behavior? in *Advances in Neural Information Processing Systems 21* (eds. Koller, D., Schuurmans, D., Bengio, Y. & Bottou, L.) 1873–1880 (Curran Associates, Inc., 2009).
16. Glaze, C. M., Kable, J. W. & Gold, J. I. Normative evidence accumulation in unpredictable environments. *eLife* **4**, e08825 (2015).
17. Mochol, G., Kiani, R. & Moreno-Bote, R. Prefrontal cortex represents heuristics that shape choice bias and its integration into future behavior. *Current Biology* **31**, 1234-1244.e6 (2021).
18. Akrami, A., Kopec, C. D., Diamond, M. E. & Brody, C. D. Posterior parietal cortex represents sensory history and mediates its effects on behaviour. *Nature* **554**, 368–372 (2018).

19. Hwang, E. J., Dahlen, J. E., Mukundan, M. & Komiyama, T. History-based action selection bias in posterior parietal cortex. *Nat Commun* **8**, 1242 (2017).

20. Morcos, A. S. & Harvey, C. D. History-dependent variability in population dynamics during evidence accumulation in cortex. *Nat Neurosci* **19**, 1672–1681 (2016).

21. Scott, B. B. *et al.* Fronto-parietal Cortical Circuits Encode Accumulated Evidence with a Diversity of Timescales. *Neuron* **95**, 385-398.e5 (2017).

22. Dayan, P. & Yu, A. J. Phasic norepinephrine: a neural interrupt signal for unexpected events. *Network (Bristol, England)* **17**, 335–350 (2006).

23. Yu, A. J. & Dayan, P. Uncertainty, neuromodulation, and attention. *Neuron* **46**, 681–692 (2005).

24. Aston-Jones, G. & Cohen, J. D. An Integrative Theory of Locus Coeruleus-Norepinephrine Function: Adaptive Gain and Optimal Performance. *Annu. Rev. Neurosci.* **28**, 403–450 (2005).

25. Harris, K. D. & Thiele, A. Cortical state and attention. *Nature Reviews Neuroscience* **12**, 509–523 (2011).

26. Lee, S.-H. & Dan, Y. Neuromodulation of brain states. *Neuron* **76**, 209–222 (2012).

27. McCormick, D. A., Nestvogel, D. B. & He, B. J. Neuromodulation of Brain State and Behavior. *Annu Rev Neurosci* **43**, 391–415 (2020).

28. Pfeffer, T. *et al.* Circuit mechanisms for the chemical modulation of cortex-wide network interactions and behavioral variability. *Science Advances* **7**, eabf5620 (2021).

29. Froemke, R. C. Plasticity of Cortical Excitatory-Inhibitory Balance. *Annu. Rev. Neurosci.* **38**, 195–219 (2015).

30. Joshi, S. & Gold, J. I. Pupil Size as a Window on Neural Substrates of Cognition. *Trends in Cognitive Sciences* (2020) doi:10.1016/j.tics.2020.03.005.

31. Reimer, J. *et al.* Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in cortex. *Nat Commun* **7**, 13289 (2016).

32. de Gee, J. W. *et al.* Dynamic modulation of decision biases by brainstem arousal systems. *eLife* **6**, 309 (2017).

33. Breton-Provencher, V., Drummond, G. T., Feng, J., Li, Y. & Sur, M. Spatiotemporal dynamics of noradrenaline during learned behaviour. *Nature* **606**, 732–738 (2022).

34. Bymaster, F. P. *et al.* Atomoxetine Increases Extracellular Levels of Norepinephrine and Dopamine in Prefrontal Cortex of Rat: A Potential Mechanism for Efficacy in Attention Deficit/Hyperactivity Disorder. *Neuropsychopharmacology* **27**, 699–711 (2002).

35. Robbins, T. W. & Arnsten, A. F. T. The Neuropsychopharmacology of Fronto-Executive Function: Monoaminergic Modulation. *Annu. Rev. Neurosci.* **32**, 267–287 (2009).

36. Steinmetz, N. A., Zatka-Haas, P., Carandini, M. & Harris, K. D. Distributed coding of choice, action and engagement across the mouse brain. *Nature* **576**, 266–273 (2019).

37. Pfeffer, T. *et al.* Catecholamines alter the intrinsic variability of cortical population activity and perception. *PLOS Biology* **16**, e2003453 (2018).

38. Pfeffer, T. *et al.* Circuit mechanisms for the chemical modulation of cortex-wide network interactions and behavioral variability. *Sci Adv* **7**, eabf5620 (2021).

39. Nuiten, S. A., de Gee, J. W., Zantvoord, J. B., Fahrenfort, J. J. & van Gaal, S. Catecholaminergic neuromodulation and selective attention jointly shape perceptual decision-making. *Elife* **12**, RP87022 (2023).

40. Nuiten, S. A., Gee, J. W. de, Zantvoord, J. B., Fahrenfort, J. J. & Gaal, S. van. Pharmacological Elevation of Catecholamine Levels Improves Perceptual Decisions, But Not Metacognitive Insight. *eNeuro* **11**, (2024).

41. Gold, J. I. & Shadlen, M. N. The neural basis of decision making. *Annu. Rev. Neurosci.* **30**, 535–574 (2007).

42. Joshi, S., Li, Y., Kalwani, R. M. & Gold, J. I. Relationships between Pupil Diameter and Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. *Neuron* **89**, 221–234 (2016).

43. Breton-Provencher, V. & Sur, M. Active control of arousal by a locus coeruleus GABAergic circuit. *Nat Neurosci* **22**, 218–228 (2019).

44. Maheu, M., Donner, T. H. & Wiegert, J. S. Serotonergic and Noradrenergic Interactions in Pupil-linked Arousal. 2025.03.26.644382 Preprint at https://doi.org/10.1101/2025.03.26.644382 (2025).

45. Bogacz, R., Brown, E., Moehlis, J., Holmes, P. & Cohen, J. D. The physics of optimal decision making: a formal analysis of models of performance in two-alternative forced-choice tasks. *Psychological review* **113**, 700–765 (2006).

46. Brody, C. D. & Hanks, T. D. Neural underpinnings of the evidence accumulator. *Current opinion in neurobiology* **37**, 149–157 (2016).

47. Ratcliff, R. & McKoon, G. The diffusion decision model: theory and data for two-choice decision tasks. *Neural computation* **20**, 873–922 (2008).

48. de Gee, J. W. *et al.* Pupil-linked phasic arousal predicts a reduction of choice bias across species and decision domains. *Elife* **9**, e54014 (2020).

49. Hebisch, J. *et al.* Task-irrelevant stimuli reliably boost phasic pupil-linked arousal but do not affect decision formation. *Sci Rep* **14**, 28380 (2024).

50. Tsujimura, S., Wolffsohn, J. s. & Gilmartin, B. A linear chromatic mechanism drives the pupillary response. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **268**, 2203–2209 (2001).

51. Wang, C.-A. & Munoz, D. P. Modulation of stimulus contrast on the human pupil orienting response. *European Journal of Neuroscience* **40**, 2822–2832 (2014).

52. de Gee, J. W., Correa, C. M. C., Weaver, M., Donner, T. H. & van Gaal, S. Pupil Dilation and the Slow Wave ERP Reflect Surprise about Choice Outcome Resulting from Intrinsic Variability in Decision Confidence. *Cereb Cortex* **31**, 3565–3578 (2021).

53. Colizoli, O., de Gee, J. W., Urai, A. E. & Donner, T. H. Task-evoked pupil responses reflect internal belief states. *Scientific Reports* **8**, 13702 (2018).

54. Cremer, A., Kalbe, F., Müller, J. C., Wiedemann, K. & Schwabe, L. Disentangling the roles of dopamine and noradrenaline in the exploration-exploitation tradeoff during human decision-making. *Neuropsychopharmacol.* 1–9 (2022) doi:10.1038/s41386-022-01517-9.

55. Jepma, M. *et al.* Noradrenergic and Cholinergic Modulation of Belief Updating. *Journal of Cognitive Neuroscience* **30**, 1803–1820 (2018).

56. Lawson, R. P., Bisby, J., Nord, C. L., Burgess, N. & Rees, G. The Computational, Pharmacological, and Physiological Determinants of Sensory Learning under Uncertainty. *Current Biology* **31**, 163-172.e4 (2021).

57. Reynaud, A. J. *et al.* Atomoxetine improves attentional orienting in a predictive context. *Neuropharmacology* **150**, 59–69 (2019).

58. Totah, N. K., Logothetis, N. K. & Eschenko, O. Atomoxetine accelerates attentional set shifting without affecting learning rate in the rat. *Psychopharmacology* **232**, 3697–3707 (2015).

59. Jordan, R. The locus coeruleus as a global model failure system. *Trends in Neurosciences* **0**, (2023).

60. Filipowicz, A. L., Glaze, C. M., Kable, J. W. & Gold, J. I. Pupil diameter encodes the idiosyncratic, cognitive complexity of belief updating. *Elife* **9**, e57872 (2020).

61. Findling, C., Skvortsova, V., Dromnelle, R., Palminteri, S. & Wyart, V. Computational noise in reward-guided learning drives behavioral variability in volatile environments. *Nat Neurosci* **22**, 2066–2077 (2019).

62. Krishnamurthy, K., Nassar, M. R., Sarode, S. & Gold, J. I. Arousal-related adjustments of perceptual biases optimize perception in dynamic environments. *Nature Human Behaviour* **1**, 0107 (2017).

63. Murphy, P. R., Wilming, N., Hernandez-Bocanegra, D. C., Prat-Ortega, G. & Donner, T. H. Adaptive circuit dynamics across human cortex during evidence accumulation in changing environments. *Nat Neurosci* **24**, 987–997 (2021).

64. Nassar, M. R. *et al.* Rational regulation of learning dynamics by pupil-linked arousal systems. *Nat Neurosci* **15**, 1040–1046 (2012).

65. Lewandowska, K., Gągol, A., Sikora-Wachowicz, B., Marek, T. & Fąfrowicz, M. Saying "yes" when you want to say "no" - pupil dilation reflects evidence accumulation in a visual working memory recognition task. *International Journal of Psychophysiology* **139**, 18–32 (2019).

66. Schriver, B. J., Perkins, S. M., Sajda, P. & Wang, Q. Interplay between components of pupil-linked phasic arousal and its role in driving behavioral choice in Go/No-Go perceptual decision-making. *Psychophysiology* **00**, e13565 (2020).

67. de Gee, J. W., Knapen, T. & Donner, T. H. Decision-related pupil dilation reflects upcoming choice and individual bias. *Proceedings of the National Academy of Sciences of the United States of America* **111**, E618-25 (2014).

68. Bari, A. & Aston-Jones, G. Atomoxetine modulates spontaneous and sensory-evoked discharge of locus coeruleus noradrenergic neurons. *Neuropharmacology* **64**, 53–64 (2013).

69. Murphy, P. R., O'Connell, R. G., O'Sullivan, M., Robertson, I. H. & Balsters, J. H. Pupil diameter covaries with BOLD activity in human locus coeruleus. *Human brain mapping* **35**, 4140–4154 (2014).

70. Varazzani, C., San-Galli, A., Gilardeau, S. & Bouret, S. Noradrenaline and Dopamine Neurons in the Reward/Effort Trade-Off: A Direct Electrophysiological Comparison in Behaving Monkeys. *J. Neurosci.* **35**, 7866–7877 (2015).

71. Mridha, Z. *et al.* Graded recruitment of pupil-linked neuromodulation by parametric stimulation of the vagus nerve. *Nat Commun* **12**, 1539 (2021).

72. Cazettes, F., Reato, D., Morais, J. P., Renart, A. & Mainen, Z. F. Phasic Activation of Dorsal Raphe Serotonergic Neurons Increases Pupil Size. *Current Biology* **31**, 192-197.e4 (2021).

73. Grujic, N., Tesmer, A., Bracey, E., Peleg-Raibstein, D. & Burdakov, D. Control and coding of pupil size by hypothalamic orexin neurons. *Nat Neurosci* **26**, 1160–1164 (2023).

74. Sara, S. J. The locus coeruleus and noradrenergic modulation of cognition. *Nature Reviews Neuroscience* **10**, 211–223 (2009).

75. Ranjbar-Slamloo, Y. & Fazlali, Z. Dopamine and Noradrenaline in the Brain; Overlapping or Dissociate Functions? *Front. Mol. Neurosci.* **12**, (2020).

76. Ding, Y.-S. *et al.* Clinical doses of atomoxetine significantly occupy both norepinephrine and serotonin transports: Implications on treatment of depression and ADHD. *NeuroImage* **86**, 164–171 (2014).

77. Sauer, J.-M., Ring, B. J. & Witcher, J. W. Clinical Pharmacokinetics of Atomoxetine. *Clin Pharmacokinet* **44**, 571–
590 (2005).

78. Watson, A. B. & Pelli, D. G. QUEST: A Bayesian adaptive psychometric method. *Perception & psychophysics*
(1983).

79. Knapen, T. *et al.* Cognitive and Ocular Factors Jointly Determine Pupil Responses under Equiluminance. *PLOS*
ONE **11**, e0155574 (2016).

80. Green, D. M. & Swets, J. A. *Signal Detection Theory and Psychophysics. 1966*. (New York, 1966).

81. Shinn, M., Lam, N. H. & Murray, J. D. A flexible framework for simulating and fitting generalized drift-diffusion
models. *eLife* **9**, e56938.

82. Seabold, S. & Perktold, J. Statsmodels: Econometric and Statistical Modeling with Python. in 92–96 (Austin,
Texas, 2010). doi:10.25080/Majora-92bf1922-011.

83. Draper, N. R. & Smith, H. *Applied Regression Analysis*. (Wiley-Interscience, 1998).

Supplementary figures legends

692
693
694 **Figure S1. (A)** Schematic of stimulus-response mapping rules. **(B)** Individual shift in pupil size caused by
695 atomoxetine, plotted against individual shift in heart rate caused by atomoxetine. Heart rate and pupil size
696 measured as the average across task blocks (Methods), separately for placebo and atomoxetine sessions
697 (Methods); data points are individual participants; stats, Pearson's correlation coefficient.
698 *https://doi.org/10.5281/zenodo.16779024*.
699
700

701
702
Figure S2. Atomoxetine does not affect overall choice bias. (A) Histograms of reaction times, separately
for placebo and atomoxetine sessions. **(B)** Signal detection theoretic sensitivity (Methods), separately for
placebo and atomoxetine sessions, and separately or weak and strong signals. Every connecting line is a
participant; large data points in the middle are the group averages (error bars, s.e.m. across 19 participants);
stats, main effect of drug in two-way repeated measures ANOVA. **(C)** As B, but for repetition probability after
correct choices (positive feedback). **(D)** As C, but after incorrect choices (negative feedback). **(E)** Permutation
(randomized) distribution of correlation coefficients describing relationship between individual shift in repetition
probability caused by atomoxetine and individual's repetition probability during the placebo sessions. The
observed correlation (plotted in Figure 2E) is stronger than 99% of this permutation distribution, resulting in a
p-value of 0.01. **(F)** As B, but for signal detection theoretic criterion (Methods). **(G)** Individual shift in criterion
caused by atomoxetine, plotted against individual's criterion during the placebo sessions. Data points are
individual participants; error bars, 68% confidence intervals across 5K bootstraps; stats, Pearson's correlation
coefficient (corrected for reversion to the mean; Methods). **(H)** As F, but for absolute criterion (Methods).
https://doi.org/10.5281/zenodo.16779024.
717

718
719
Figure S3. Individual choice history biases are better explained by history-dependent changes in drift
bias than starting point bias. (A) Relationship between individual choice repetition probabilities [P(repeat)]
and history shift in starting point (left column) and drift bias (right column), in the placebo sessions. Parameter
estimates were obtained from a model in which both bias terms were allowed to vary with previous choice
(Methods). Stats, Pearson's correlation coefficient. The plots show that the starting point was shifted away
from the previous response for most participants (negative values along x-axis in left panel) and drift bias was
shifted towards the previous response (positive values along x-axis in right panel. These opposing effects
have been reported before[4]. **(B)** As A, but for atomoxetine sessions. **(C)** Group-level boxplots of all parameter
estimates. Black diamonds, outliers. **(D)** Measured and predicted RT distributions, separately for 'left' and
'right' choices, across all participants. **(E)** Conditional response bias functions. We divided all measured and
model-predicted trials into five quantiles of the RT distribution and plotted for each quantile the fraction of
'right'-choices. Error bars, s.e.m. across 19 participants. *https://doi.org/10.5281/zenodo.16779024*.
732

734
735

Figure S4. (A) Beta values from time-wise regression of repetition probability and task-evoked pupil response
(same relationship as in **Figure 4B**). **(B)** Relationship between task-evoked pupil response (same time-
window as used main **Figure 4**; Methods) and accuracy (left), RT (middle) or repetition probability w.r.t next
trial (right). Stats, mixed linear modeling (Methods). **(C)** Individual shift repetition probability caused by
atomoxetine, plotted against individual shift in heart rate caused by atomoxetine. Heart rate measured as the
average across task blocks (Methods), separately for placebo and atomoxetine sessions; data points are
individual participants; error bars, 68% confidence intervals across 5K bootstraps; stats, Pearson's correlation
coefficient. *https://doi.org/10.5281/zenodo.16779024*.