Catecholamines reduce choice history biases

Jan Willem de Gee[1,2,3,*], Niels A. Kloosterman[1,4,5,6], Anke Braun[1,7] & Tobias H. Donner[1,*]

[1] Department of Neurophysiology and Pathophysiology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany.
[2] Cognitive and Systems Neuroscience, Swammerdam Institute for Life Sciences, University of Amsterdam, Amsterdam, NLD;
[3] Amsterdam Brain & Cognition, University of Amsterdam, Amsterdam, The Netherlands.
[4] Department of Psychology, University of Lübeck, Germany
[5] Center of Brain, Behavior and Metabolism, University of Lübeck, Germany
[6] Max Planck Institute for Human Development, Berlin, Germany
[7] Department of Psychiatry and Neurosciences, Charité - Universitätsmedizin Berlin, Berlin, Germany.

*Address correspondence to Jan Willem de Gee & Tobias H. Donner.

Email: j.w.degee@uva.nl, t.donner@uke.edu

Author Contributions: Conceptualization: JWG, NK, TD; Funding Acquisition: TD; Investigation: JWG, NK; Data Curation: JWG, NK; Formal analysis: JWG, NK, AB; Writing—original draft: JWG, TD; Writing—review and editing: JWG, NK, AB, TD; Supervision: TD; Project Administration: TD.

Competing Interest Statement: The authors declare no competing interests.

Abstract

Theoretical accounts postulate that the catecholaminergic neuromodulator noradrenaline shapes cognitive behavior by reducing the impact of prior expectations on learning, inference, and decision-making. A ubiquitous effect of dynamic priors on perceptual decisions under uncertainty is choice history bias: the tendency to systematically repeat, or alternate, previous choices, even when stimulus categories are presented in a random sequence. Here, we directly test for a causal impact of catecholamines on these priors. We pharmacologically elevated catecholamine levels through the application of the noradrenaline reuptake inhibitor atomoxetine. We quantified the resulting changes in observers' history biases in a visual perceptual decision task. Choice history biases in this task were highly idiosyncratic, tending toward choice repetition or alternation in different individuals. Atomoxetine decreased these biases (toward either repetition or alternation) compared to placebo. Behavioral modeling indicates that this bias reduction was due to a reduced bias in the accumulation of sensory evidence, rather than of the starting point of the accumulation process. Atomoxetine had no significant effect on other behavioral measures tested, including response time and choice accuracy. We conclude that catecholamines reduce the impact of a specific form of prior on perceptual decisions.

Introduction

Biases are prevalent in human decisions ranging from those based on abstract reasoning to judgments of the sensory environment[1–3]. One prominent source of such biases in perceptual decisions across mammalian species is the history of previous experiences[4–13]. When, as common in natural environments, the environmental state evaluated in the judgment is relatively stable, such history biases reflect adaptive expectations that are dynamically updated over time and improve performance[5,14,15]. For example, when choosing to forage for deer in the same area as during successful previous expeditions. While recent work in rodents and primates has identified correlates of such history biases in neural population activity of parietal and frontal cortical areas[9,12,16–20], little is currently known about the neurotransmitter systems involved in this important form of bias.

Influential theoretical work postulates that the catecholaminergic neuromodulator noradrenaline shapes behavior by reducing the impact of prior expectations on learning, inference, and decision-making[21,22]. The catecholaminergic (and in particular, the noradrenaline) systems of the brainstem project to large parts of the cerebral cortex, where the neuromodulators they release change the functional properties of their target networks[23–28]. Consequently, they are in an ideal position to shape cortex-wide network activity underlying decision-making in a coordinated fashion. Indeed, pupil responses (a peripheral proxy of central arousal state[29] point to a role of neuromodulation in regulating the impact of these history-dependent priors[8]. However, pupil responses reflect the activity of multiple neuromodulatory brainstem systems[29–31], and the above result builds on correlative approaches. Direct evidence for the role of catecholaminergic neuromodulation in choice history biases is currently sparse in animal models[13,32] and absent in humans.

Here, we aimed to provide causal evidence for the role of catecholamines in regulating (suppressing) history biases in human choice behavior. Specifically, we hypothesized that catecholamines down-regulate the impact of previous experiences on decision-making, which should translate into a reduction of choice history bias. We tested this hypothesis by pharmacologically increasing central catecholamine (in particular noradrenaline) levels through the selective noradrenaline re-uptake inhibitor atomoxetine[33,34] and quantified the resulting behavioral changes in a variant of a visual perceptual decision task adopted from recent animal physiology[35]. Atomoxetine inhibits the presynaptic norepinephrine transporter, preventing the reuptake of norepinephrine throughout the brain along with inhibiting the reuptake of dopamine in specific brain

72 regions such as the prefrontal cortex[33,34]. We found that atomoxetine reduced choice history biases
73 by specifically interacting with the way people accumulated noisy perceptual evidence.
74
75 **Results**
76
Atomoxetine was administered orally (40 mg per session) in a within-subject, double-blind, placebo-
controlled, and randomized design (**Figure 1A**). The behavioral task entailed the presentation of
visual targets (Gabor patch of fixed orientation and varying contrast), one in the left and one in the
right visual hemifield; participants were asked to report the location the larger contrast target by
button press (**Figure 1B**; Methods). To increase sensory uncertainty and promote integration of
evidence across time[36], we added dynamic visual noise to both Gabor patches (**Figure 1B**;
Methods). There were two different signal strengths (small and large difference in contrast);
auditory feedback was presented 50 ms right after participant's choices. We collected many trials
per participant (range, 3520-4320) across four experimental sessions, in which we systematically
varied and counterbalanced drug condition and stimulus-response mapping (**Figure 1A** and **Figure
S1**; Methods).
88
89

90
Figure 1. Experimental design and effects on peripheral arousal markers. (A) Design was within-subject,
double-blinded, randomized, and placebo-controlled (Methods). Additionally, the stimulus-response mapping
*was counterbalanced within participants (**Figure S1**; Methods). (B) Schematic sequence of events during the*
spatial contrast discrimination task (Methods). Participants reported the location of the highest contrast Gabor
patch (difference in contrast is high for illustration only). (C) Heart rate (Methods), separately for placebo and
atomoxetine sessions. Every connecting line is a participant; large data points in the middle are the group
averages (error bars, s.e.m. across 19 participants); stats, paired-sample t-test. (D) As C, but for pupil size
(Methods).
99
100
Atomoxetine administration at this relatively low dose had a measurable effect on peripheral
arousal markers, but not on the overall efficiency of the decision process for the behavioral task at
hand. Atomoxetine increased both heart rate (**Figure 1C**) and pupil size (**Figure 1D**). While signal
strength lawfully decreased reaction time (RT; $F_{1,18}$ = 16.2, p = 0.001; two-way repeated measures
ANOVA) and increased accuracy ($F_{1,18}$ = 297.3, p < 0.001) (**Figure 2A,B**), atomoxetine changed
neither of those behavioral parameters (RT: $F_{1,18}$ = 2.3, p = 0.146; accuracy: $F_{1,18}$ = 0.8, p = 0.383;
no significant interaction of signal strength and drug; **Figure 2A,B**). Likewise, atomoxetine did not
significantly change signal detection theoretic sensitivity (d'; **Figure S2A**).
109
By contrast, atomoxetine had a robust impact on the impact of history-based (i.e., dynamically
varying throughout the course of the experiment) prior expectations on their decisions. We
operationalized these time-varying prior expectations about the next target as the probability of
repeating the choice from the previous trial (**Figure 2C**). The stimulus repetition probability was, by
design, close to 0.5 (**Figure 2C**) (group average: 0.496; Methods), thus not explaining the
sequential effects in participants' behavior. Atomoxetine reduced the mean choice repetition
probability (**Figure 2D**; $F_{1,18}$ = 8.2, p = 0.010; two-way repeated measures ANOVA). Signal strength

had no such effect ($F_{1,18} = 0.1$, $p = 0.706$) and signal strength and drug did not interact ($F_{1,27} = 0.5$,
$p = 0.507$).
119
In the placebo sessions, most participants (15 out of 19) had a repetition probability larger than 0.5,
that is, they tended to repeat their choices (**Figure 2C**). We, therefore, wondered if (i) atomoxetine
reduced the group-level choice history biases by stereotypically pushing behavior more toward
alternation (i.e., reduction of repetition probability, in line with previous pupillometry work[8]) or (ii)
through a reduction of both repetition and alternation biases (shift of repetition probability toward
0.5). Our results are in line with the second scenario. There was a robust correlation between
individual choice history biases during placebo sessions and the shift in history biases caused by
atomoxetine: participants with the strongest repetition or alternation biases, exhibited the strongest
shift towards a repetition probability of 0.5 (**Figure 2E**; corrected for reversion to the mean;
Methods). Consequently, atomoxetine yielded an even more consistent and robust reduction of the
absolute choice history bias, defined as the absolute value of P(repeat)−0.5 (**Figure 2F**; $F_{1,18} =$
10.6, $p < 0.001$; two-way repeated measures ANOVA). Atomoxetine did not change overall choice
bias, quantified as signal detection theoretic criterion (**Figure S2D-F**). These findings show that
atomoxetine specifically reduced the impact of history-dependent prior expectation in all
participants, regardless of whether they idiosyncratically tended to repeat (i.e., assuming a
relatively stable environment) or alternate (assuming a volatile environment).
136
137

138
139
Figure 2. Atomoxetine reduces choice history bias. (A) *Reaction time, separately for placebo and*
atomoxetine sessions, and separately or weak and strong signals. Every connecting line is a participant; large
data points in the middle are the group averages (error bars, s.e.m. across 19 participants); stats, main effect
*of drug in two-way repeated measures ANOVA. See **Figure S2B** for reaction time distributions. **(B)** As A, but*
*for accuracy. **(C)** Probability of repeating the previous stimulus category (left) or choice (right) during placebo*
*sessions. **(D)** As A, but for repetition probability (Methods). **(E)** Individual shift in repetition probability caused*
by atomoxetine, plotted against individual's repetition probability during the placebo sessions. Data points are
individual participants; error bars, 68% confidence intervals across 5K bootstraps; stats, Pearson's correlation
*coefficient (corrected for reversion to the mean; **Figure S2C**; Methods). **(F)** As A, but for absolute repetition*
probability (Methods).
150
151
152

How did participants' expectations shape their decision processes, and how did atomoxetine act on this mechanism? Current models of perceptual decision-making posit the temporal accumulation of sensory evidence, resulting in an internal decision variable that grows with time[36–39]. When this decision variable reaches one of two decision bounds, a choice is made, and the corresponding motor response is initiated. In this framework (**Figure 3A**), a bias can arise in two ways: (i) by shifting the starting point of accumulation toward one of the bounds or (ii) by changing the rate of evidence accumulation toward one choice alternative.

Previous work has shown that individual biases in choice repetition probabilities are generally explained by a history-dependent bias in the drift rather than starting point, across a range of different perceptual choice tasks[4]. We replicated this finding for the placebo condition of the present dataset (**Figure S3A**). To this end, we fitted an accumulation-to-bound model of decision-making to the behavioral data. The model had five free parameters: 1) boundary separation (controlling response caution); 2) the mean drift rate (overall efficiency of evidence accumulation); 3) non-decision time (the speed of pre-decisional evidence encoding and post-decisional translation of choice into motor response); 4) the starting point of the decision; 5) an evidence-independent bias in the drift (henceforth called "drift bias"). We fitted all five parameters separately per drug condition; additionally, we fitted drift rate separately for both signal strength conditions, and both starting point and drift bias separately for the previous choice category ('left' vs 'right'). The fitted model (**Figure S3C**) accounted well for the overall behavior in each task (**Figure S3D,E**). Indeed, atomoxetine also specifically reduced the history-dependent shift in drift bias, but not the history shift in starting point (**Figure 3B**). Furthermore, this drift bias (not starting point) effect was robustly and specifically correlated to the individual change of choice repetition probability (**Figure 3C**).

Figure 3. Atomoxetine reduces choice history bias in evidence accumulation. (A) *Schematic of drift diffusion model accounting for choices, and their associated RTs. In the equation, v is the drift rate. Purple and cyan curves show expected RT distributions under shifts in either the 'starting point' (z; purple) or 'drift bias' (v_{bias}; cyan).* ***(B)*** *Left: history-dependent shift in starting point bias (Methods), separately for placebo and atomoxetine sessions. Right, as left, but for drift bias. Every connecting line is a participant; large data points in the middle are the group averages (error bars, s.e.m. across 19 participants); stats, paired-sample t-test.* ***(C)*** *Left: individual shift in repetition probability caused by atomoxetine, plotted against individual's history-dependent shift in starting point bias. 'History shift' is the difference between a parameter estimate for previous 'left' and previous 'right' choices, irrespective of the category of the current stimulus. Right: as left, but for history-dependent shift in drift point bias. Data points are individual participants; stats, Pearson's correlation coefficient (difference in correlation assessed with permutation test; Methods).*

Discussion

Using selective pharmacological interventions in humans, we here imply catecholamines in the modulation of human choice history biases: boosting catecholamines levels through atomoxetine specifically reduced the magnitude of choice history biases, regardless of whether those promoted choice repetition or alternation. We interpret this finding in the context of computational theory postulating a down-weighting of prior expectations (which, we assume, drive the choice history

biases analyzed here) in combination with new sensory evidence in the evaluation of current
environmental state[21,22]. Catecholamines have previously been implicated in modulating learning
rates[40–43] and facilitating set shifting[44] in dynamic environments; here, we provide causal evidence
for their role in controlling the weight of historical information in perceptual decisions.

Our finding is roughly consistent with previous correlative evidence from pupillometry[8,45–49]. Non
luminance-mediated changes of pupil diameter are an indirect marker of neuromodulatory
(including noradrenergic and cholinergic) activity[30,31,50–54]. In dynamic environments, pupil
responses are associated with violations of learned, top-down expectations and reduce their
weight[45–49]. In stable environments, pupil responses predict a reduction of choice repetition
probability[8]. There are, however, also notable differences between our current results and this
previous pupillometry work. First, the pupil-predicted reduction of repetition probability was due to
a stereotypical increase in the tendency to alternate[8]. In contrast, we here found that
catecholamines reduce both repetition and alternation biases (shift of repetition probability toward
0.5), which is more readily in line with a reduction in prior weight. Second, our pharmacological
intervention did not reduce participant's overall choice bias (bias towards one choice alternative,
irrespective of the immediate history), even though this is another commonly reported behavioral
correlate of pupil responses[31,55–58]. One possible explanation for this difference is that atomoxetine
likely increased both tonic and phasic catecholamines levels[59], while task-evoked pupil responses
reflect phasic neuromodulatory activity[31]. Another possible explanation is that while atomoxetine
selectively elevates central catecholamine levels (noradrenaline and dopamine), pupil responses
track not only noradrenaline[30,31,50–53], but also acetylcholine[30,54], serotonin[60] and orexin[61] levels.

Which neuromodulator specifically mediated the effects reported here? Atomoxetine increases the
extracellular levels of noradrenaline, but also of (prefrontal) dopamine (Methods). Indeed, one
recent report suggest that striatal dopamine may play a role in adapting choice history bias to
temporal regularities in a stimulus sequence[13]. Thus, our results are consistent with a role of one
or both catecholamines. However, the noradrenergic locus coeruleus and dopaminergic midbrain
exhibit a close interplay[62,63] and are co-activated with pupil responses during decisions[31]. Thus, it
could well be that their (complex) interaction regulates choice history bias in perceptual decisions.

In sum, pharmacologically elevating central catecholamine (specifically noradrenaline) levels in
human participants specifically reduces their propensity to accumulate noisy evidence for
perceptual decisions in a manner that is biased by previously reported perceptual experience. This
supports computational theories according to which noradrenaline reduces the impact of dynamic
priors on inference and decision-making.

Materials and Methods

Participants
Nineteen participants (14 female, age range 20-26) participated in the experiment, entailing
concurrent pupillometry and MEG recordings. All had normal or corrected to normal vision and no
history or indications of psychological or neurological disorders. The experiment was approved by
the ethics committee of the University Medical Center Hamburg-Eppendorf. Participants
participated in four experimental sessions, in which we crossed drug condition (atomoxetine vs.
placebo) and stimulus-response mapping (press ipsi vs. contra) (about 2 hours per session).
Participants were paid 10€ per hour. One participant completed three recording sessions.

Pharmacological intervention
We used the selective noradrenaline reuptake inhibitor atomoxetine (dose, 40 mg) to boost the
levels of catecholamines, specifically noradrenaline and (in the prefrontal cortex) dopamine[33,34].
Atomoxetine is a relatively selective inhibitor of the noradrenaline transporter, which is responsible
for the natural reuptake of noradrenaline that has been released into the extracellular space.
Consequently, atomoxetine acts to increase the extracellular levels of noradrenaline, an effect that

has been confirmed experimentally in rat prefrontal cortex[33]. The same study showed that
atomoxetine also increases the prefrontal levels of dopamine, which has a molecular structure very
similar to the one of noradrenaline and is, in fact, a direct precursor of noradrenaline. Atomoxetine
has smaller affinity to the serotonin transporter, and there are discrepant reports about the
quantitative relevance of these effects: while one study found no increases in serotonin levels under
atomoxetine[33], a recent study reports a significant atomoxetine-related occupancy of the serotonin
transporter in nonhuman primates[64] at dosages that would correspond to human dosages of 1.0–
1.8 mg/kg. Note that these dosages are substantially higher than the administered dosage in this
study (40 mg, independent of body weight). It is therefore unclear to what extent our atomoxetine
condition affected cortical serotonin levels. A mannitol-aerosil mixture was administered as
placebo. All substances were encapsulated identically to render them visually indistinguishable.
Peak plasma concentrations are reached about 1–2 hours after administration for atomoxetine, and
the half-life is about 5 hours[65]. Thus, participants received either a placebo or atomoxetine 1.5
hours before the start of the experimental session.
267
Behavioral task
Each trial consisted of three consecutive intervals (**Figure 1A**): (i) the baseline interval (uniformly
distributed between 0.6 and 1.4 s) containing a dynamic noise pattern on both the left and right
side of the green fixation dot; (ii) the decision interval (terminated by the participant's response;
max 3 s) containing the same noise patterns with Gabor patch embedded in each; (iii) the feedback
interval (uniformly distributed between 0.25 and 1.4 s).
274
Two dynamic noise patterns were presented throughout the experiment. The luminance across all
pixels was kept constant. This pedestal noise pattern had 20% contrast. In the decision interval two
Gabor patches (sinusoidal gratings; 2 cycles per degree; phase randomly drifted left or right) were
added to each noise pattern. The Gabor patches symmetrically differed from 80% contrast. On half
of trials, the Gabor patch on right side of the screen was of higher contrast (signal, right); on the
other half the Gabor patch on the left side was of higher contrast (signal, left). Signal location was
randomly selected on each trial, under the constraint that it would occur on 50% of the trials within
each block of 160 trials. Stimuli were presented within two circular patches to the left and right of
the central fixation cross (diameter, 15 degrees of visual angle; eccentricity, 12 degrees).
284
Participants were instructed to report the location of the signal, the highest contrast Gabor patch
by pressing one of two response buttons with their left or right index finger as soon as they felt
sufficiently confident ("free response paradigm"). We did not give specific instructions to tradeoff
accuracy for speed (or vice versa). The mapping between perceptual choice and button press (e.g.,
"right" –> press right key; "left" –> press left key) was counterbalanced across the four recording
sessions. Participants received auditory feedback (high tone, correct; low tone, error) 50 ms after
they indicated their choice.
292
293 Signal strength, the difference in contrast between the left and right signals, was individually titrated
294 before the main experiment to two difficulty levels that yielded about 70% and 85% correct choices,
295 using an adaptive staircase procedure (Quest)[66]. In the placebo sessions, we measured a mean
296 accuracy of 74.07 % correct (± 0.97 % s.e.m.) and 80.83 % correct (± 1.12 % s.e.m), for weak and
297 strong signal strengths respectively.
298
Stimuli were back-projected on a transparent screen using a Sanyo PCL-XP51 projector with a
resolution of 800x600 at 24 Hz. The screen was positioned 65 cm away from their eyes. The
luminance profile was linearized by measuring and correcting for the systems gamma curve. A
doubling of contrast values therefore also produced a doubling of luminance differences. During
the first training session stimuli were presented on a VIEWPixx monitor with the same resolution
and refresh rate (also linearized). To minimize any effect of light on pupil diameter, the overall
luminance of the screen was held constant throughout the experiment.
306

307 Participants performed between 22 and 27 blocks (distributed over four sessions) yielding a total
308 of 3520–4320 trials per participant. One participant performed a total of 17 blocks (distributed over
309 three scanning sessions), yielding a total of 2720 trials.
310
Data acquisition
Recordings took place in a dimly lit magnetically shielded room. Magnetoencephalography (MEG)
was recorded using a whole-head CTF 275 MEG system (CTF Systems Inc., Canada) at a
sampling rate of 1200 Hz. In addition, eye movements and pupil diameter were recorded with an
MEG-compatible EyeLink 1000 Long Range Mount system (SR Research, Osgoode, ON, Canada).
Finally, electrocardiogram (ECG) as well as vertical and horizontal electrooculogram (EOG) were
acquired using Ag/AgCl electrodes at a sampling rate of 1200 Hz. MEG data will be the focus of a
different report.
319
320 *Analysis of electrocardiogram data*
321 ECG data were used to analyze average heart rate. We used an adaptive threshold to detect the
322 R-peak of each QRS-complex in the ECG. Heart rate was then computed by dividing the total
323 number of R-components by time.
324
325 *Analysis of pupil data*
326 All analyses were performed using custom-made Python scripts, unless stated otherwise.
327
Preprocessing. Periods of blinks and saccades were detected using the manufacturer's standard
algorithms with default settings. The remaining data analyses were performed using custom-made
Python scripts. We applied to each pupil timeseries (i) linear interpolation of missing data due to
blinks or other reasons (interpolation time window, from 150 ms before until 150 ms after missing
data), (ii) low-pass filtering (third-order Butterworth, cut-off: 6 Hz), (iii) removal of pupil responses
to blinks and to saccades, by first estimating these responses by means of deconvolution and then
removing them from the pupil time series by means of multiple linear regression[67], and (iv)
conversion to units of modulation (percent signal change) around the mean of the pupil time series
from each measurement session.
337
338 *Quantification of pre-trial pupil size.* We quantified pre-trial pupil size as the mean pupil size during
339 the 0.25 s before trial onset.
340
341 *Analysis and modeling of choice behavior*
342 All analyses were performed using custom-made Python scripts, unless stated otherwise. The first
343 trial of each block was excluded from the analyses. We always computed behavioral metrics
344 separately for each participant and each of the four recording sessions, and then averaged over
345 the two SR-mappings within each drug condition.
346
347 *Overt behavior.* Reaction time (RT) was defined as the time from stimulus offset until the button
348 press. Repetition probability was defined as the fraction of choices that was the same as the choice
349 on the previous trial.
350
351 *Signal-detection theoretic modeling.* The signal detection[68] metrics sensitivity (d') and criterion (c)
352 were computed separately for each of the pupil bins. We estimated d' as the difference between z-
353 scores of hit-rates and false-alarm rates. We estimated criterion by averaging the z-scores of hit-
354 rates and false-alarm rates and multiplying the result by -1.
355
356 *Drift diffusion modeling.* We fitted the reaction time data with the drift diffusion model[37,39], based on
357 continuous maximum likelihood using the Python-package PyDDM[69].
358 The decision dynamics were described by (**Figure 3A**):
359

$$\Delta y = (s \cdot v + v_{bias}) \cdot \Delta t + cdW \qquad \text{Eq. 1}$$

361
362 where y is the decision variable, s is the stimulus category (-1 for left signals; 1 for right signals), v
363 is the drift rate and controls the overall efficiency of evidence accumulation, v_{bias} is the drift bias
364 which is and evidence-independent constant added to the drift, and cdW is Gaussian distributed
365 white noise with mean 0 and variance $c^2\Delta t$. The starting point of evidence accumulation z was
366 defined as a fraction of the boundary separation:
367
368                                $$y(0) = z \cdot a$$ Eq. 2
369
370 where a is the separation between the two decision bounds. Evidence accumulation terminated at
371 0 ("left") or a ("right). The model was fitted separately for each recording session of each participant.
372 In each fit, we let drift rate vary with signal strength[39], and starting point and drift bias with previous
373 choice[4].
374
375 *Statistical comparisons*
376 We used a paired sample t-test to test for significant differences in heart rate, pupil size, and
377 behavioral measures between placebo and drug sessions (**Figure 1** and **Figure 3**). We used 2 ×
378 2 repeated measures ANOVA to test for the main effects of drug and signal strength, as well as
379 their interaction, on behavioral measures (**Figure 2**). All tests were performed two-tailed.
380
We performed permutation analyses to obtain the distribution of correlation coefficients predicted
exclusively by regression to the mean (**Figure 2E**). This was done by computing the above-
described across-subject correlation 10K, each time using randomly assigned "placebo" vs. "drug"
labels of each participant. We then compared our observed correlation coefficient
(reflecting the combined effects of regression to the mean and the relationship between overall
bias and the pupil predicted shift therein) against this permutation distribution. This analysis
suggested that the relationship between choice history bias and the Atomoxetine predicted shift
therein was stronger than expected based on regression to the mean (proportion of
permutations below observed correlation < 0.05).
390
391 We also used a permutation procedure to test for differences between the correlation coefficients
392 for history shifts in starting point bias and drift bias (**Figure 3C**). Again, we permuted the condition
393 labels (drug / placebo).
394
395 *Data availability*
396 Data will be made publicly available upon publication.
397
398 *Code availability*
399 Analysis scripts will be made publicly available upon publication.
400
401 **Acknowledgments**
402

403 Funding: Project grant of Amsterdam Brain & Cognition Priority Program; German Research
404 Foundation (DFG; grant number Heisenberg Professorship DO 1240/3-1, and SFB 936, Projects
405 A2/A3, A7, Z3).

406
407 **References**
408

1. Fischer, J. & Whitney, D. Serial dependence in visual perception. *Nat Neurosci* **17**, 738–743 (2014).
2. Tversky, A. & Kahneman, D. Judgment under Uncertainty: Heuristics and Biases. *Science (New York,*
N.Y.) **185**, 1124–1131 (1974).
3. Kloosterman, N. A. *et al.* Humans strategically shift decision bias by flexibly adjusting sensory evidence
accumulation. *eLife* **8**, e37321 (2019).

4. Urai, A. E., de Gee, J. W., Tsetsos, K. & Donner, T. H. Choice history biases subsequent evidence
accumulation. *eLife* **8**, e46331 (2019).
5. Braun, A., Urai, A. E. & Donner, T. H. Adaptive History Biases Result from Confidence-Weighted
Accumulation of past Choices. *J. Neurosci.* **38**, 2418–2429 (2018).
6. Hermoso-Mendizabal, A. *et al.* Response outcomes gate the impact of expectations on perceptual
decisions. *Nat Commun* **11**, 1057 (2020).
7. Abrahamyan, A., Silva, L. L., Dakin, S. C., Carandini, M. & Gardner, J. L. Adaptable history biases in
human perceptual decisions. *PNAS* **113**, E3548–E3557 (2016).
8. Urai, A. E., Braun, A. & Donner, T. H. Pupil-linked arousal is driven by decision uncertainty and alters
serial choice bias. *Nat Commun* **8**, 14637 (2017).
9. Urai, A. E. & Donner, T. H. Persistent activity in human parietal cortex mediates perceptual choice
repetition bias. *Nat Commun* **13**, 6015 (2022).
10. Gold, J. I., Law, C.-T., Connolly, P. & Bennur, S. The Relative Influences of Priors and Sensory
Evidence on an Oculomotor Decision Variable During Perceptual Learning. *Journal of Neurophysiology*
100, 2653–2668 (2008).
11. Busse, L. *et al.* The Detection of Visual Contrast in the Behaving Mouse. *J. Neurosci.* **31**, 11351–11361
(2011).
12. Braun, A. & Donner, T. H. Adaptive biasing of action-selective cortical build-up activity by stimulus
history. *Elife* **12**, RP86740 (2023).
13. Fritsche, M. *et al.* Temporal regularities shape perceptual decisions and striatal dopamine signals. *Nat*
Commun **15**, 7093 (2024).
14. Yu, A. J. & Cohen, J. D. Sequential effects: Superstition or rational behavior? in *Advances in Neural*
Information Processing Systems 21 (eds. Koller, D., Schuurmans, D., Bengio, Y. & Bottou, L.) 1873–
1880 (Curran Associates, Inc., 2009).
15. Glaze, C. M., Kable, J. W. & Gold, J. I. Normative evidence accumulation in unpredictable
environments. *eLife* **4**, e08825 (2015).
16. Mochol, G., Kiani, R. & Moreno-Bote, R. Prefrontal cortex represents heuristics that shape choice bias
and its integration into future behavior. *Current Biology* **31**, 1234-1244.e6 (2021).
17. Akrami, A., Kopec, C. D., Diamond, M. E. & Brody, C. D. Posterior parietal cortex represents sensory
history and mediates its effects on behaviour. *Nature* **554**, 368–372 (2018).
18. Hwang, E. J., Dahlen, J. E., Mukundan, M. & Komiyama, T. History-based action selection bias in
posterior parietal cortex. *Nat Commun* **8**, 1242 (2017).
19. Morcos, A. S. & Harvey, C. D. History-dependent variability in population dynamics during evidence
accumulation in cortex. *Nat Neurosci* **19**, 1672–1681 (2016).
20. Scott, B. B. *et al.* Fronto-parietal Cortical Circuits Encode Accumulated Evidence with a Diversity of
Timescales. *Neuron* **95**, 385-398.e5 (2017).
21. Dayan, P. & Yu, A. J. Phasic norepinephrine: a neural interrupt signal for unexpected events. *Network*
(Bristol, England) **17**, 335–350 (2006).
22. Yu, A. J. & Dayan, P. Uncertainty, neuromodulation, and attention. *Neuron* **46**, 681–692 (2005).
23. Aston-Jones, G. & Cohen, J. D. An Integrative Theory of Locus Coeruleus-Norepinephrine Function:
Adaptive Gain and Optimal Performance. *Annu. Rev. Neurosci.* **28**, 403–450 (2005).
24. Harris, K. D. & Thiele, A. Cortical state and attention. *Nature Reviews Neuroscience* **12**, 509–523
(2011).
25. Lee, S.-H. & Dan, Y. Neuromodulation of brain states. *Neuron* **76**, 209–222 (2012).
26. McCormick, D. A., Nestvogel, D. B. & He, B. J. Neuromodulation of Brain State and Behavior. *Annu*
Rev Neurosci **43**, 391–415 (2020).
27. Pfeffer, T. *et al.* Circuit mechanisms for the chemical modulation of cortex-wide network interactions
and behavioral variability. *Science Advances* **7**, eabf5620 (2021).
28. Froemke, R. C. Plasticity of Cortical Excitatory-Inhibitory Balance. *Annu. Rev. Neurosci.* **38**, 195–219
(2015).
29. Joshi, S. & Gold, J. I. Pupil Size as a Window on Neural Substrates of Cognition. *Trends in Cognitive*
Sciences (2020) doi:10.1016/j.tics.2020.03.005.
30. Reimer, J. *et al.* Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in cortex.
Nat Commun **7**, 13289 (2016).
31. de Gee, J. W. *et al.* Dynamic modulation of decision biases by brainstem arousal systems. *eLife* **6**, 309
(2017).
32. Breton-Provencher, V., Drummond, G. T., Feng, J., Li, Y. & Sur, M. Spatiotemporal dynamics of
noradrenaline during learned behaviour. *Nature* **606**, 732–738 (2022).

33. Bymaster, F. P. *et al.* Atomoxetine Increases Extracellular Levels of Norepinephrine and Dopamine in
Prefrontal Cortex of Rat: A Potential Mechanism for Efficacy in Attention Deficit/Hyperactivity Disorder.
Neuropsychopharmacology **27**, 699–711 (2002).
34. Robbins, T. W. & Arnsten, A. F. T. The Neuropsychopharmacology of Fronto-Executive Function:
Monoaminergic Modulation. *Annu. Rev. Neurosci.* **32**, 267–287 (2009).
35. Steinmetz, N. A., Zatka-Haas, P., Carandini, M. & Harris, K. D. Distributed coding of choice, action and
engagement across the mouse brain. *Nature* **576**, 266–273 (2019).
36. Gold, J. I. & Shadlen, M. N. The neural basis of decision making. *Annu. Rev. Neurosci.* **30**, 535–574
(2007).
37. Bogacz, R., Brown, E., Moehlis, J., Holmes, P. & Cohen, J. D. The physics of optimal decision making:
a formal analysis of models of performance in two-alternative forced-choice tasks. *Psychological review*
113, 700–765 (2006).
38. Brody, C. D. & Hanks, T. D. Neural underpinnings of the evidence accumulator. *Current opinion in*
neurobiology **37**, 149–157 (2016).
39. Ratcliff, R. & McKoon, G. The diffusion decision model: theory and data for two-choice decision tasks.
Neural computation **20**, 873–922 (2008).
40. Cremer, A., Kalbe, F., Müller, J. C., Wiedemann, K. & Schwabe, L. Disentangling the roles of dopamine
and noradrenaline in the exploration-exploitation tradeoff during human decision-making.
Neuropsychopharmacol. 1–9 (2022) doi:10.1038/s41386-022-01517-9.
41. Jepma, M. *et al.* Noradrenergic and Cholinergic Modulation of Belief Updating. *Journal of Cognitive*
Neuroscience **30**, 1803–1820 (2018).
42. Lawson, R. P., Bisby, J., Nord, C. L., Burgess, N. & Rees, G. The Computational, Pharmacological, and
Physiological Determinants of Sensory Learning under Uncertainty. *Current Biology* **31**, 163-172.e4
(2021).
43. Reynaud, A. J. *et al.* Atomoxetine improves attentional orienting in a predictive context.
Neuropharmacology **150**, 59–69 (2019).
44. Totah, N. K., Logothetis, N. K. & Eschenko, O. Atomoxetine accelerates attentional set shifting without
affecting learning rate in the rat. *Psychopharmacology* **232**, 3697–3707 (2015).
45. Filipowicz, A. L., Glaze, C. M., Kable, J. W. & Gold, J. I. Pupil diameter encodes the idiosyncratic,
cognitive complexity of belief updating. *Elife* **9**, e57872 (2020).
46. Findling, C., Skvortsova, V., Dromnelle, R., Palminteri, S. & Wyart, V. Computational noise in reward-
guided learning drives behavioral variability in volatile environments. *Nat Neurosci* **22**, 2066–2077
(2019).
47. Krishnamurthy, K., Nassar, M. R., Sarode, S. & Gold, J. I. Arousal-related adjustments of perceptual
biases optimize perception in dynamic environments. *Nature Human Behaviour* **1**, 0107 (2017).
48. Murphy, P. R., Wilming, N., Hernandez-Bocanegra, D. C., Prat-Ortega, G. & Donner, T. H. Adaptive
circuit dynamics across human cortex during evidence accumulation in changing environments. *Nat*
Neurosci **24**, 987–997 (2021).
49. Nassar, M. R. *et al.* Rational regulation of learning dynamics by pupil-linked arousal systems. *Nat*
Neurosci **15**, 1040–1046 (2012).
50. Breton-Provencher, V. & Sur, M. Active control of arousal by a locus coeruleus GABAergic circuit. *Nat*
Neurosci **22**, 218–228 (2019).
51. Joshi, S., Li, Y., Kalwani, R. M. & Gold, J. I. Relationships between Pupil Diameter and Neuronal
Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. *Neuron* **89**, 221–234 (2016).
52. Murphy, P. R., O'Connell, R. G., O'Sullivan, M., Robertson, I. H. & Balsters, J. H. Pupil diameter
covaries with BOLD activity in human locus coeruleus. *Human brain mapping* **35**, 4140–4154 (2014).
53. Varazzani, C., San-Galli, A., Gilardeau, S. & Bouret, S. Noradrenaline and Dopamine Neurons in the
Reward/Effort Trade-Off: A Direct Electrophysiological Comparison in Behaving Monkeys. *J. Neurosci.*
35, 7866–7877 (2015).
54. Mridha, Z. *et al.* Graded recruitment of pupil-linked neuromodulation by parametric stimulation of the
vagus nerve. *Nat Commun* **12**, 1539 (2021).
55. de Gee, J. W. *et al.* Pupil-linked phasic arousal predicts a reduction of choice bias across species and
decision domains. *Elife* **9**, e54014 (2020).
56. Lewandowska, K., Gągol, A., Sikora-Wachowicz, B., Marek, T. & Fąfrowicz, M. Saying "yes" when you
want to say "no" - pupil dilation reflects evidence accumulation in a visual working memory recognition
task. *International Journal of Psychophysiology* **139**, 18–32 (2019).

57. Schriver, B. J., Perkins, S. M., Sajda, P. & Wang, Q. Interplay between components of pupil-linked
phasic arousal and its role in driving behavioral choice in Go/No-Go perceptual decision-making.
Psychophysiology **00**, e13565 (2020).
58. de Gee, J. W., Knapen, T. & Donner, T. H. Decision-related pupil dilation reflects upcoming choice and
individual bias. *Proceedings of the National Academy of Sciences of the United States of America* **111**,
E618-25 (2014).
59. Bari, A. & Aston-Jones, G. Atomoxetine modulates spontaneous and sensory-evoked discharge of
locus coeruleus noradrenergic neurons. *Neuropharmacology* **64**, 53–64 (2013).
60. Cazettes, F., Reato, D., Morais, J. P., Renart, A. & Mainen, Z. F. Phasic Activation of Dorsal Raphe
Serotonergic Neurons Increases Pupil Size. *Current Biology* **31**, 192-197.e4 (2021).
61. Grujic, N., Tesmer, A., Bracey, E., Peleg-Raibstein, D. & Burdakov, D. Control and coding of pupil size
by hypothalamic orexin neurons. *Nat Neurosci* **26**, 1160–1164 (2023).
62. Sara, S. J. The locus coeruleus and noradrenergic modulation of cognition. *Nature Reviews*
Neuroscience **10**, 211–223 (2009).
63. Ranjbar-Slamloo, Y. & Fazlali, Z. Dopamine and Noradrenaline in the Brain; Overlapping or Dissociate
Functions? *Front. Mol. Neurosci.* **12**, (2020).
64. Ding, Y.-S. *et al.* Clinical doses of atomoxetine significantly occupy both norepinephrine and serotonin
transports: Implications on treatment of depression and ADHD. *NeuroImage* **86**, 164–171 (2014).
65. Sauer, J.-M., Ring, B. J. & Witcher, J. W. Clinical Pharmacokinetics of Atomoxetine. *Clin*
Pharmacokinet **44**, 571–590 (2005).
66. Watson, A. B. & Pelli, D. G. QUEST: A Bayesian adaptive psychometric method. *Perception &*
psychophysics (1983).
67. Knapen, T. *et al.* Cognitive and Ocular Factors Jointly Determine Pupil Responses under
Equiluminance. *PLOS ONE* **11**, e0155574 (2016).
68. Green, D. M. & Swets, J. A. *Signal Detection Theory and Psychophysics. 1966.* (New York, 1966).
69. Shinn, M., Lam, N. H. & Murray, J. D. A flexible framework for simulating and fitting generalized drift-
diffusion models. *eLife* **9**, e56938.

554
555

556 **Supplementary figures**

560 *Figure S1. Schematic of stimulus-response mapping rules.*

Figure S2. Atomoxetine does not affect overall choice bias. (A) Signal detection theoretic sensitivity
(Methods), separately for placebo and atomoxetine sessions, and separately or weak and strong signals.
Every connecting line is a participant; large data points in the middle are the group averages (error bars, s.e.m.
across 19 participants); stats, main effect of drug in two-way repeated measures ANOVA. (B) Histograms of
reaction times, separately for placebo and atomoxetine sessions. (C) Permutation (randomized) distribution
of correlation coefficients describing relationship between individual shift in repetition probability caused by
atomoxetine and individual's repetition probability during the placebo sessions. The observed correlation
(plotted in Figure 2E) is stronger than 99% of this permutation distribution, resulting in a p-value of 0.01. (D)
As A, but for signal detection theoretic criterion (Methods). (E) Individual shift in criterion caused by
atomoxetine, plotted against individual's criterion during the placebo sessions. Data points are individual
participants; error bars, 68% confidence intervals across 5K bootstraps; stats, Pearson's correlation coefficient
(corrected for reversion to the mean; Methods). (F) As A, but for absolute criterion (Methods).

Figure S3. Individual choice history biases are better explained by history-dependent changes in drift bias than starting point bias. (A) *Relationship between individual choice repetition probabilities [P(repeat)] and history shift in starting point (left column) and drift bias (right column), in the placebo sessions. Parameter estimates were obtained from a model in which both bias terms were allowed to vary with previous choice (Methods). Stats, Pearson's correlation coefficient; black diamonds, outliers.* **(B)** *As A, but for atomoxetine sessions.* **(C)** *Group-level boxplots of all parameter estimates.* **(D)** *Measured and predicted RT distributions, separately for 'left' and 'right' choices, across all participants.* **(E)** *Conditional response bias functions. We divided all measured and model-predicted trials into five quantiles of the RT distribution and plotted for each quantile the fraction of 'right'-choices. Error bars, s.e.m. across 19 participants.*