Neuroligin 3 highlights sexually dimorphic circuitry in Drosophila social spacing

Authors:

Robinson JW[1], Bechard AT[1], Evans MR[1], Mosuro K[1†], Kurbaj J[1,] Isaacson JR[1], Pillay S[1†], Lin DS[1†], Sahota A[1†], de Belle JN[1†], Robinson GI[2], Moehring AJ[1], Simon, AF[1*]

Affiliations:
1. Department of Biology, University of Western Ontario, London, ON, Canada
2. Department of Biological Sciences, University of Lethbridge, Lethbridge, AB, Canada

† Undergraduate students at time of data gathering
* Correspondence: asimon28@uwo.ca (A.F.S.)

Running head
Drosophila social spacing neurocircuitry

Keywords:
Drosophila, social space, social behaviour, neural circuitry, *neuroligin*, mushroom bodies, *fruitless*, acetylcholine, dopamine, GABA, receptors

Abstract:

In *Drosophila melanogaster*, the autism-related Neuroligin 3 (Nlg3) protein is a postsynaptic membrane protein important for synapse development and regulation, which plays a role in social spacing behaviour. Here, we report the localization of Nlg3 to the calyx of the mushroom bodies (MB), optic lobes (OL), and protocerebral bridge (PB). Using RNA interference, *nlg3* knockdown in the OL increased fly social space, especially in the males, while knocking down *nlg3* independently in the MB or the PB did not change social spacing. However, hyperactivation and silencing of these neurons in the MB, but not the PB, controls social space in males and females, while hyperactivating and silencing of all *nlg3*-expressing neurons, including within the MB, PB, and OL, regulates male and female social space. Knocking down neurotransmitter biosynthesis enzymes, which decreases the amount of neurotransmitter release, showed that reducing acetylcholine release from the MB decreased female social space, whereas knocking down any dopamine receptor in the MB increased male social space. Lastly, to investigate the sexually dimorphic effects on social spacing previously seen in *nlg3* mutants, we examined a subset of sexually dimorphic *fruitless*-expressing (*fru)P1* neurons known to regulate sexually dimorphic behaviours. Hyperactivation of those *fruP1* neurons decreased social space in both sexes, while silencing those *fruP1* neurons specifically increased male social space without affecting females. Our findings highlight a sex-specific social space neural circuitry that includes the OL, MB, and *fruP1* neurons, while uncovering the underlying basis of some of the sex differences in this behaviour.

Article Summary:

In vinegar flies (*Drosophila melanogaster*), the autism-related Neuroligin 3 protein (Nlg3) controls neuronal development and regulation but also affects fly social behaviour. Nlg3 is localized to the mushroom bodies (MB), protocerebral bridge, and the optic lobes. Genetically reducing Nlg3 in the optic lobes made male flies more socially distant, while hyperactivating or silencing neurons in the MB affected social behavior in both sexes. Reducing acetylcholine release from the MB affects female social space, while reducing dopamine receptors of the MB only affect male behaviour. Finally, the *fruitless* sexually dimorphic neurons control social behavior differently in males and females.

Introduction:

Social behaviours are fundamental to the interaction and communication within a community and are influenced by environmental cues which are processed and integrated in the brain. Abnormal social behaviours, such as difficulties with communication or interactions with others, are clinical characteristics of individuals with autism spectrum disorders (ASD). Using *Drosophila melanogaster,* we can dissect the neural and genetic underpinnings of social interactions in a simplified system (Sokolowski 2001).

Flies exhibit social behaviours that, while simpler, can reflect the basic principles of social interaction found in more complex organisms (Sokolowski 2010). By examining how environmental cues are integrated and influence social behaviours in Drosophila, we can gain insights into the mechanisms that might be disrupted in ASD, providing a basis for understanding the molecular and neural circuitries involved in these disorders (Ferreira and Moita 2019).

An important component of Drosophila social interactions within a group context is social spacing – the distance of one individual to another in a group. We use the social space assay to assess the outcome of group behaviour displayed by flies as a measure of the average number of flies within four-body lengths (4BL) in the social space chamber (Simon et al. 2012). We and other research groups have used this assay as a metric for behavioural analysis to understand and reveal the basic underlying neural circuitry involved (Burg et al. 2013; Fernandez et al. 2017; Xie et al. 2018; Yost et al. 2020).

So far, we know that mutations in genes affecting eye function, have been found to make flies less social (Simon et al. 2012; Burg et al. 2013; Jiang et al. 2020). Both males and females have decreased social space with mutants associated to fly gustation and mechanosensation (Jiang et al. 2020). In contrast, *Orco* mutants, which affect olfactory perception, do not appear to influence social spacing in males, but result in decreased social space in females. Together, these suggest a sensory modality-specific genetic influence on social behaviour that can have sex-specific effects (Simon et al. 2012; Jiang et al. 2020).

Other genes expressed in the brain have also been shown to influence Drosophila social behaviour, such as genes important for synaptic transmission and synaptogenesis (Burg et al. 2013; Wise et al. 2015; Fernandez et al. 2017; Ueoka et al. 2018; Brenman-Suttner et al. 2019; Hope et al. 2019; Jiang et al. 2020; Kanellopoulos et al. 2020; Yost et al. 2020; Shilpa et al. 2021; Cao et al. 2022). At the synapse level, the concentration of neurotransmitters, which is tightly regulated by biosynthesis enzymes or vesicular packaging proteins, have been shown to affect social spacing. Tyrosine hydroxylase (TH), the rate-limiting enzyme in dopamine synthesis, and the vesicular monoamine transporter (VMAT), which packages dopamine into synaptic vesicles, both regulate dopamine levels (Simon et al. 2009; Lawal et al. 2010). Modulating the abundance or activity of these proteins influences social spacing in Drosophila, indicating that dopamine plays a role in regulating social space (Fernandez et al. 2017).

Many human homologs of neural genes have also emerged as potential candidates in neuropsychiatric disorders, bridging the gap between basic biological processes and complex

human pathologies (Grant 2012). The gene *neurobeachin*, which is implicated in ASD in
humans, encodes a structural scaffolding postsynaptic density protein (Castermans et al. 2003).
In Drosophila, mutations in the homolog *rugose* result in altered fly social spacing, indicating a
conservation of function across species (Wise et al. 2015). Furthermore, synaptic proteins
Neuroligin and Neurexin, which facilitate neurotransmission by aiding synapse formation and
receptor recruitment, have been shown to influence social behaviour (Hahn et al. 2013). In
humans, *Neuroligin* gene homologs are correlated with ASD (Jamain et al. 2003; Südhof 2008).
Similarly, in Drosophila, mutations in these genes lead to altered social behaviours, further
revealing the genetic underpinnings of social space (Corthals et al. 2017).
89
Drosophila Neuroligin (Nlg) is a family of post-synaptic membrane proteins with the conserved
function of synapse formation and maintenance by interacting with presynaptic Neurexin to
mediate cell-cell adhesion and facilitate neurotransmission (Südhof 2008). During Drosophila
neural development, larval *nlg3* mutants have a higher number of synaptic connections but less
GluRIIA recruitment and synaptic transmission compared to control flies (Xing et al. 2014).
Furthermore, *nlg3* knockouts display a sexually dimorphic and age-associated effect on fly
aggression and social space (Yost et al. 2020). Mutant *nlg3* flies had lower climbing
performance in both males and females by over 30%, however, *nlg3* deletion decreases only
male, but not female aggression; while increasing male social space in 3-4 day old flies, and in
both sexes by 7-10 days old (Yost et al. 2020).
100
The mushroom bodies (MB), also known to play a role in Drosophila social spacing, are a well-
studied invertebrate brain structure primarily involved in fly olfactory learning and memory. The
MB are comprised of approximately 2,500 neurons and its unusually structured shape suggests a
highly specialized organization and function (Erber et al. 1987). The MB have a flexible
organization in response to social isolation and are the central processing structure of the
Drosophila brain (Technau 1984). Consisting of densely packed neurons called Kenyon cells, the
MB integrates many sources of sensory information to modulate fly behavioural responses.
Social space assays have been conducted on flies with a mutation in an ion channel selectively
targeted in the MB and lead to decreased social space, observing that the cholinergic signaling
on its own can rescue this altered phenotype in social spacing (Burg et al. 2013).
111
Sexually dimorphic behaviours in Drosophila such as male courtship, female receptivity, and
social interactions are known to be regulated by *fruitless* (*fru*) neurons (Pan and Baker 2014;
Brenman-Suttner et al. 2018; Chowdhury et al. 2020). The *fruitless* gene encodes a transcription
factor important for sex determination in Drosophila development and produces male-specific
transcripts that are essential for the development and function of male-specific neural circuits
(Sato and Yamamoto 2020). Regulated by multiple promoters each expressing unique
transcripts, the *fruP1* transcript undergoes sex-specific splicing and contributes to the sexual
differentiation of neural circuits influencing behaviours such as courtship and aggression
(Wiedemann 2007; Baker et al. 2024). Promoters *p2-p5* are necessary for adult viability and fly
morphology that is however not sex specific (Ito et al. 1996). Neurons expressing *fru* are
widespread interneurons in the adult fly brain and ventral nerve cord, where fast acting *fru*
neurons release acetylcholine, GABA, and glutamate, while other *fru* neurons also release
dopamine, serotonin, tyramine, and octopamine (Palmateer et al. 2023).
125

While progress has been made in identifying individual genes, synaptic proteins, and neuropils
like the MB that influence Drosophila social spacing, the integration of molecular players within
precise circuitry or other brain structures mediating social spacing behaviour are not fully
understood. This research looks deeper into the role *nlg3* has on Drosophila social space. Based
on Nlg3 localization and the sex-specific differences observed, we elucidated part of the neural
circuitry that directly influences social spacing behaviour.
132
133 **Materials and Methods:**
134
135 *Fly Stocks and Husbandry*
136
All flies were reared mixed sex in bottles containing our own fly food recipe (brown sugar, corn
meal, yeast, agar, benzoic acid, methyl paraben, and propionic acid). Rearing conditions were
controlled during development and aging at 50% humidity, 25ºC, and on a 12:12 hour light:dark
cycle. Parents were removed after seven days of egg laying resulting in a maximum parental age
of 14 days, as parental age affects social spacing of the offspring (Brenman-Suttner et al. 2018).
Canton-S (Cs) control line was initially sourced from Seymour Benzer in 1998. UAS-effector
lines were outcrossed five times into the Cs control line to minimize behavioural variation due to
genetic background. All genotypes used are reported in **Supp. Table 1**. A generalized crossing
scheme for both the V10 and Cs backgrounds used to create the appropriate experimental
treatment and genetic controls can be found in the supplementary material (**Supp. Fig. 1**). The
results obtained including those UAS effector lines, despite being in a different genetic
background, are presented in **Supp. Fig. 7-10** where the effect of genetic background may be
observed in some experiments.
150
151 *nlg3 RNAi Generation*
152
153 The shRNA sequences targeting nlg3 were obtained by using the model provided by Vert et al.
154 (2006). The length of the *nlg3* shRNA was 60 bases for the full hairpin; and was ~7.8kbp for the
155 entire pVALIUM20 plasmid containing the shRNA was ~7.8kbp. The hairpin sequence used
156 was:
157
158 CAGCAACTCGAACTCGAACTAtagttatattcaagcataTAGTTCGAGTTCGAGTTGCTG
159
Within the hairpin sequence, uppercase letters represent the nucleotides that bind to the *nlg3*
mRNA, while lowercase letters represent the nucleotides that connect the two halves of the pin.
The shRNA constructs were cloned into the vector pVALIUM20 (vermillion eye color maker,
(Ni et al. 2011) using the protocol in Chang et al. (2014). DNA was injected into genotype *y v,*
nanos-integrase, attP40/attP40 (Bloomington Stock #: 25709) embryos and were mated to the
parental line after reaching adulthood. Then, offspring containing the transgene were
backcrossed five times into the Cs control background (Vert et al. 2006; Chang et al. 2014).
167
168 *Immunocytochemistry*
169
Anti-Nlg3

Whole brain dissections on 3 to 4 day old, male or female brains were fixed within 20 minutes of
dissection in Bouin's solution. Following fixation and washing, brains were blocked in 5% NGS
in 1X PBS and stained with guinea pig anti-Nlg3 primary antibody (1:1000, Invitrogen)
overnight at 4 °C, which is a polyclonal primary antibody kindly provided by Dr. Brian Mozer
(Yost et al. 2020). After washing, brains were stained with goat anti-guinea pig Alexa 488
(1:500, Invitrogen) for 2 hours at room temperature. Next, all samples were left in glycerol for 1
hour then were mounted on slides using Fluoroshield™ media, stored at 4 °C and imaged the
following day. Fluorescent images were visualized with a AxioImager Z1 Compound
Fluorescent microscope (Zeiss). Note: we used the entirety of our polyclonal antibody stock and
have been unsuccessful in generating another functioning antibody for Nlg3. We switched to
genetic constructs that tag Nlg3 with a GFP reporter (*MiMIC-nlg3* and *nlg3-Gal4 > mCD8-*
GFP) that both verify our findings.
183
Anti-GFP
Whole brain dissections on 3 to 4 day old males or females were fixed within 20 minutes of
dissection in 4% paraformaldehyde in 1X PBS-T (0.3% Triton-X). Following fixation and
washing, brains were blocked in 5% normal goat serum (NGS) in PBS-T and stained with rabbit
anti-GFP (1:1000, Invitrogen) and mouse anti-Brp (nc82; 1:30, Developmental Studies
Hybridoma Bank) for 48 hours at 4 °C. After washing, brains were stained with Alexa 660 anti-
mouse (1:400, Invitrogen) and Alexa 488 anti-rabbit (1:800, Invitrogen) for 48 hours at 4°C.
Samples were mounted on slides in Fluoroshield™ media, stored at 4 °C and imaged the
following day. Slides were imaged on an SP8 Lightening Confocal microscopy system (Leica).
All images were visualized with LAS X 2.0 software (Leica).
194
195　*Social Space Assay*
196
The assay was performed as previously described (McNeil et al. 2015; Yost et al. 2020). In
short, flies were collected after eclosion and group housed until aged at 3 to 4 days old. Yost et
al. (2024) have shown that the flies are no longer virgin after the age treatment we use for these
experiments. Twenty-four hours prior to the assay, flies were sexed under cold anesthesia and
separated into vials with food containing 12-17 flies for each treatment tested. Two hours before
the assay, each treatment was transferred into new vials and acclimated to standard conditions
(25°C and 50% humidity). Ten minutes prior to test start, if the standard acclimation temperature
is different than testing condition temperature, treatments were set to the testing temperature
required for neuronal hyperactivation (*TrpA1* at 27°C and 50% humidity) or silencing (*Shi^{ts}* at
29°C and 50% humidity). For those experiments we compared the effect of the treatment at
higher temperature (activation of genetic constructs and alteration of neural transmission) to the
typical room testing temperatures (25°C, control: inactivated genetic constructs and basal
transmission of neurons). So, the same genotypes were tested at 2 different temperatures. All
genetic background lines used to create the experimental lines displayed no effect of the
temperature changes (**Fig. 2**). At the time of testing, flies were loaded into the triangular
chamber and knocked to the bottom of the chamber to induce a startle response that forces the
flies into a group at the top of the chamber within the first minute. Flies are then given 20-40
minutes to settle at their preferred social space. Each treatment was loaded into the social space
apparatus and photos were taken as soon as flies have settled as a group (McNeil et al. 2015).
Social space assay photos were processed with ImageJ (National Institute of Health, Bethesda,

MA, USA) to calculate the number of flies within a four-body length (4BL or ~1cm) from
macros (as described in Yost et al. 2020). Image analysis data were imported to GraphPad Prism
10 software for statistical analysis and graphical representation.
220
221   *Climbing Assay*
222
The assay was performed as previously described (McNeil et al. 2015; Yost et al. 2020). In this
assay we are testing flies' stamina, ability to respond to startle, and locomotion. Any lines with
flies able to reach the top vial (performance index, PI, more than zero) in 15 seconds do not have
impairments preventing testing in the social space assays (McNeil et al. 2015; Yost et al. 2020).
Flies were aged to 2 to 4 days old and 24 hours prior to conducting the assay were sexed under
cold anesthesia and separated in vials with food containing 40 flies for each treatment. Two
hours before the assay, each treatment was transferred into new vials and acclimated to testing
conditions (25ºC and 50% humidity). Ten minutes before the assay, if the standard acclimation
temperature is different than testing condition temperature, treatments were set to testing
temperature required for neuronal hyperactivation (*TrpA1* at 27°C and 50% humidity) or
silencing (*Shi^{ts}* at 29°C and 50% humidity). Each treatment was transferred into a testing tube
and inserted into the climbing assay. Flies were forced to the bottom of the tube and given 15
seconds to climb to the tube on top of the apparatus. This timeframe was chosen to allow
approximately 80% of control flies to reach the top vial. After 15 seconds, the apparatus was
closed and the number of flies in the top and bottom tubes were counted to calculate a
performance index of percent flies that reached the top vial.
239
240   *Statistical Analysis*
241
Statistical analysis and graphical representation were performed in GraphPad Prism
(RRID:SCR_002798, version 7.0a for Mac, GraphPad Software, La Jolla, CA, USA,
www.graphpad.com). Normality test was used to verify normality and homoscedasticity test was
used to verify the homoscedasticity of the distribution before performing an ANOVA test. Two-
way ANOVAs were performed on data presented on the same graph, which were collected
simultaneously and had 2 variables in 2 groups. Following the ANOVAs, Šídák's post-hoc tests
were performed within each sex to compare between different genotypes or transmission states,
while correcting for multiple comparisons. The results of all the comparison tests for social
space analysis can be found in **Supp. Table 2**. Supplemental figures have all ANOVA results
and comparison tests in **Supp. Table 3**. P-values are reported on graphs if $p < 0.05$.
252
253   *Drawings:*
254   Drawings presented in figures were created on the app Procreate 5.3.5 (Savage Interactive Pty
255   Ltd, Hobart, Australia).
256

Results:

Neuroligin 3 protein is expressed at low levels in the calyx of the mushroom bodies, the protocerebral bridge, and the optic lobes

We investigated the expression patterns of the Nlg3 protein in the adult brain using an anti-Nlg3 polyclonal antibody. We found Nlg3 protein was localized within the calyx (Ca) of the mushroom bodies (MB), the protocerebral bridge (PB), and the optic lobes (OL) all in low levels compared to the background autofluorescence of brain lipids (**Fig. 1A,B**). For confirmation of the findings done with our antibody, we used two genetic constructs (*MiMIC* p-element insertion within *nlg3*, as well as an *nlg3-Gal4* insertion within the gene – see **Supp. Table 1** for specific localization). The *MiMIC-nlg3* insertion which incorporates a GFP reporter into the coding sequence of the host gene *nlg3* (Venken et al. 2011). From this construct we saw GFP expressed at low levels to the MB and OL and at very low fluorescence we can see the PB when compared to background autofluorescence of the Drosophila brain (**Fig. 1C,D**). Confirmation of expression in the calyx and PB was observed with the *nlg3-Gal4* insertion driving *mCD8-GFP* at higher levels than endogenous *nlg3* (**Fig. 1E,F**). Male and female localization of Nlg3 did not appear to be different either qualitatively or quantitatively in 3-4 day old adult flies (representative images of only the males are shown). Next, using the *nlg3-Gal4* driver to express GFP in **Fig. 1E** and **1F**, we activated an RNAi specific to *nlg3* (outcrossed in the Cs background) to all cells that express *nlg3*. In both males and females, both genetic control fly lines (*UAS/+* and *RNAi/+*) did not have different social space compared to the Cs genetic background control strain (**Fig. 1G,H**). However, in both males and females again, we did see a decrease in the number of flies within 4BL for the RNAi knocking in *nlg3*-expressing cells (One-Way ANOVA $p < 0.0001$, post-hoc comparison of ngl3>UAS-RNAi nlg3 to other controls $p < 0.0001$, see details in **Fig. 1G,H**).

Neuroligin 3 expression in the calyx of the mushroom bodies, the protocerebral bridge, and the optic lobes is required for normal social spacing

To determine the behavioural function of Nlg3 located in these structures, we targeted RNAi knock down of *nlg3* independently in each of the MB, PB, and OL and tested fly social space. We observed a behavioural effect of knocking down *nlg3* in the MB (**Fig. 2A**; Two-way ANOVA – Effect of genotype: $F_{1,39} = 14.71$, $p = 0.0004$), in the PB (**Fig. 2B**; Two-way ANOVA – Effect of genotype: $F_{1,40} = 19.02$, $p < 0.0001$; Šídák's multiple comparisons test p = 0.001), and the OL (**Fig. 2C**; Two-way ANOVA – Effect of genotype: $F_{1,31} = 7.191$, $p = 0.0116$). Then, we used a climbing assay to assess fly locomotion and their ability to respond to a startle response and climb; all to ensure the treatment flies could perform in the social space assay. While there was no reduction in climbing when knocking down *nlg3* in the MB (**Supp. Fig. 2A**), there was a reduction in climbing when knocking down *nlg3* in each of the PB and OL (**Supp. Fig. 2B,C**). The flies were able to climb and therefore were able to perform the social space assay. We also observed the variations in climbing are not correlated to the variations in social spacing.

Hyperactivation and silencing nlg3-expressing neurons, independently or simultaneously, changes fly social space in a sex-specific manner

Next, we looked at genetically manipulating neural transmission within the brain structures
where Nlg3 is localized (**Fig. 1**) to examine their effect on social space. When hyperactivating
all neurons within the MB, we saw a lower number of flies within 4BL (increased social space)
in both males and females (**Fig. 3A**; Two-way ANOVA – Effect of neural transmission: $F_{1,47} =$
7.395, $p = 0.0091$). Silencing all MB neurons showed the same reduction in the number of flies
within 4BL again in both males and females (**Fig. 3A**; Two-way ANOVA – Effect of neural
transmission: $F_{1,30} = 14.40$, $p = 0.0007$) highlighting the importance of the MB in determining
fly social space on their own. In contrast, hyperactivation of the PB neurons did not change fly
social space, nor did silencing PB neurons in males but looks to have an effect in females
specifically (**Fig. 3B**; Two-way ANOVA – Effect of neural transmission: $F_{1,31} = 3.443$, $p =$
0.0730). Finally, we used the *nlg3-Gal4* insertion to manipulate the transmission of all *nlg3*-
expressing neurons. Hyperactivation of all *nlg3*-expressing neurons decreases the number of
flies within 4BL in both males and females (**Fig. 3C**; Two-way ANOVA – Effect of neural
transmission: $F_{1,28} = 12.70$, $p = 0.0013$). Silencing these neurons only increased social space in
females but not males (**Fig. 3C**; Two-way ANOVA – Effect of interaction: $F_{1,22} = 6.169$, $p =$
0.0211). There was no reduction in climbing with a hyperactivation or silencing of the MB or PB
neurons (**Supp. Fig. 3A,B**), but we did observe a reduction in climbing when hyperactivating
nlg3-expressing neurons in males only (**Supp. Fig. 3C**).
321
322   *Reducing acetylcholine release from the mushroom bodies decreases social space in both sexes,*
323   *reducing GABA production only increases social space in females, but knocking down dopamine*
324   *receptors in the mushroom bodies increases male social space*
325
As the MB are important for fly social space, we investigated which neurotransmitters in the MB
were important for the observed effects. Using UAS-RNAi, we knocked down the rate-limiting
biosynthesis enzymes within the MB (Shih et al. 2019), including *choline acetyltransferase*
(ChAT for acetylcholine production) and *glutamate decarboxylase 1* (*GAD1* for GABA
production; Shih et al. 2019). Acetylcholine reduction in the MB decreased fly social space in
males and females (**Fig. 4A**; Two-way ANOVA – Effect of genotype: $F_{1,32} = 11.19$, $p = 0.0021$).
GABA reduction also increased the number of flies within 4BL in males, but not in females
(**Fig. 4B**; Two-way ANOVA – Effect of genotype: $F_{1,29} = 18.10$, $p = 0.0002$).
334
Previous research shows global dopaminergic changes in the adult fly brain will affect social
space (Fernandez et al. 2017; Xie et al. 2018; Yost et al. 2020), however, the anatomical
locations within the adult fly brain where alterations in dopaminergic transmission affect social
behaviour is unknown. Thus, we tested the dopaminergic inputs into the MB, and found that
knocking down every dopamine receptor within the MB – including Dop1R1 (**Fig. 5A**; Two-
way ANOVA – Effect of genotype: $F_{1,32} = 19.61$, $p < 0.0001$), Dop1R2 (**Fig. 5B**; Two-way
ANOVA – Effect of genotype: $F_{1,32} = 7.401$, $p = 0.0105$), Dop2R (**Fig. 5C**; Two-way ANOVA
– Effect of genotype: $F_{1,32} = 41.10$, $p < 0.0001$), and DopEcR (**Fig. 5D**; Two-way ANOVA –
Effect of genotype: $F_{1,32} = 8.905$, $p = 0.0054$) – decreased the number of flies within 4BL with a
larger effect size in males than females.
345
346
347
348

349 *Neurons expressing the fruitless gene under the P1 promotor are involved in sexually dimorphic*
350 *responses in fly social spacing*
351
Since male and female flies appear to have differently neural circuitry underlying their social
spacing preferences, we looked at a subset of sexually dimorphic *fruitless*-expressing neurons.
We used the *fruP1-Gal4* driver to manipulate neural transmission. While hyperactivation of
fruP1 neurons decreased the number of flies within 4BL for both males and females (**Fig. 6**;
Two-way ANOVA – Effect of neural transmission: $F_{1,33} = 22.65$, $p < 0.0001$). silencing of *fruP1*
resulted in a sex specific effect. Silencing *FruP1* neurons resulted in a higher number of flies
within 4BL for males (**Fig. 6**; Two-way ANOVA and Šídák's *post-hoc* test: $F_{1,30} = 3.451$, $p =$
0.0453) but did not change female social space. Importantly, we saw a reduction in climbing in
both males and females for hyperactivating (Two-way ANOVA – Effect of transmission: $F_{1,32} =$
153.6, $p < 0.0001$) and silencing (Two-way ANOVA – Effect of transmission: $F_{1,32} = 33.54$, $p =$
0.0102) *fruP1* neurons (**Supp. Fig. 4**).
363
To assess the effect of Nlg3 in *fruitless*-expressing neurons, we drove an RNAi against *nlg3* with
the *fruP1* driver. We saw reducing levels of *nlg3* decreased social space in flies. Knocking down
nlg3 in *fruitless*-expressing neurons resulted in an increase in the number of flies within 4BL for
males and females (**Fig. 7A**; Two-way ANOVA – Effect of genotype: $F_{1,57} = 16.27$, $p = 0.0002$).
Also, we observed a reduction of male climbing when knocking down *nlg3* in *fruP1* neurons
(**Supp. Fig. 5A**; Two-way ANOVA – Effect of genotype: $F_{1,40} = 9.735$, $p = 0.0033$). Next, we
investigated the neurotransmitters involved in fly social space in *fruP1* neurons. Knocking down
TH, the rate limiting-enzyme in the pathway of dopamine synthesis, in *fruP1* neurons did not
change male or female social space (**Fig. 7B**; Two-way ANOVA – Effect of genotype: $F_{1,29} =$
0.413, $p = 0.5255$). However, knocking down *TH* in *fruP1* neurons decreased climbing in both
males and females indicating the importance of dopamine released from *fruP1* neurons on fly
climbing (**Supp. Fig. 5B**; Two-way ANOVA – Effect of genotype: $F_{1,31} = 18.07$, $p = 0.0002$). In
contrast, a reduction of acetylcholine levels by knocking down *ChAT* in *fruP1* neurons increased
the number of flies within 4BL in both males and females (**Fig. 7C**; Two-way ANOVA – Effect
of genotype: $F_{1,47} = 99.11$, $p < 0.0001$). Acetylcholine release reduction from *fruP1* neurons also
decreases fly climbing (**Supp. Fig. 5C**). While a reduction of GABA by knocking down *GAD1*
in *fruP1* neurons decreased male and female social space (**Fig. 7D**; Two-way ANOVA – Effect
of genotype: $F_{1,42} = 8.478$, $p = 0.0057$), decreasing *fruP1* GABA resulted in no change in
climbing.
383
384

Discussion:

Social interactions within a group requires an integration of both attractive and repulsive social cues that lead to a behavioural output. Here, we expand the map of the neural basis for social behaviour by focusing on the sexually dimorphic role of *neuroligin 3* and its associated adult brain structures in male and female *Drosophila melanogaster*.

We show for the first time that *nlg3* is expressed at low levels in the mushroom bodies, optic lobes, and protocerebral bridge in the adult fly brain. Each of those structures independently requires *nlg3* for normal social spacing. The importance of Nlg3 in the OL on social behaviour was expected as vision is known to be important for proper social spacing (Simon et al. 2012; Burg et al. 2013; Jiang et al. 2013).

Additionally, manipulating total neural transmission in these brain regions highlighted the importance of the Kenyon cells for maintaining social space. Hyperactivating and silencing MB neurons in both males and females increased fly social space. However, hyperactivating and silencing the PB only affected social space in females. Reducing acetylcholine release from the MB decreased social space in both sexes, while reducing GABA only decreased social space in males and knocking down all dopamine-type receptors increased mostly male social space. These findings suggest that mapping the circuitry through the MB happens in a sexually dimorphic manner, where total acetylcholine release from the MB will modify female social space and dopaminergic inputs to the MB will change male social space. We still do not know the specific output from the MB leading to changes in males or the inputs to the MB leading to changes in females.

We also examined the involvement of *fru* neurons, which are known to regulate other sexually dimorphic behaviors such as male courtship or female receptivity (Chowdhury et al. 2020). Hyperactivating *fruP1* neurons decreased social space in both sexes, while silencing them showed to be sexually dimorphic with increased social space only occurring in males. Again, this demonstrates the influence *fruP1* neurons have on social space along with other fly social behaviours.

Lastly, manipulation of neurotransmitters within *fruP1* neurons revealed that the release of dopamine form those neurons was not important for social space. Instead, acetylcholine release from *fruP1* neurons play a role in fly social space, a phenocopy observed when knocking down acetylcholine in the MB, but GABA release decreased social space in both, with a smaller effect size in females. In addition, RNAi knockdown of *nlg3* in the *fruP1* neurons also showed sex-specific effects on social space, with a larger effect size in females.

Our antibody targeted the intracellular domain of Nlg3, which allowed us to identify protein localization specifically in the calyx (dendritic ends of the Kenyon cells) of the mushroom bodies, the medulla and lamina layers of the optic lobes, and throughout the entire protocerebral bridge (**Fig. 1**). However, there was a low signal to noise ratio when imaging with just a polyclonal anti-Nlg3 antibody (**Fig. 1A,B**) indicating a low level of expression of *nlg3* in the adult 3-4 day old brain. Attempts to create another polyclonal, or even a monoclonal antibody specific to Nlg3 were unsuccessful, so we transitioned to using two different genetic constructs

(*MiMIC-nlg3* and *nlg3-Gal4 > mCD8-GFP*) to confirm the results we observed with our polyclonal antibody. The GFP localization in this line in the mushroom bodies and optic lobes was similar to the Nlg3 protein localization (**Fig. 1C,D**). However, the GFP expression in the *MiMIC-nlg3* was also low, probably due to the endogenously low expression of *nlg3* (reported in Yost et al. 2020), and we could not see any expression in the protocerebral bridge. We then switched to a *nlg3-Gal4* driver to express Gal4 transcription factor in the same cells as *nlg3*, allowing to activated the *UAS-GFP* at higher levels than just endogenous *nlg3* (**Fig. 1F**) allowing for better visualization with immunofluorescence of the calyx and protocerebral bridge (**Fig. 1E**).

After determining the localization of *nlg3* expression in the adult brain, we examined the effects of targeted RNAi knockdown of *nlg3* on fly social space. Deletions of the whole *nlg3* gene leads to an increase in fly social space (Yost et al. 2020), which we replicated with an RNAi knockdown. Knocking down *nlg3* in the optic lobes might affect fly vision but we did not yet directly test this. The RNAi knockdown of *nlg3* leads to a reduction in climbing (**Supp. Fig. 2C**) as the climbing assay is performed under uniform light. The reduction in climbing may be influenced by impairments in vision and not solely or entirely due to impairments in locomotion. Vision is known to play a role in social spacing as well (Simon et al. 2012; Burg et al. 2013; Jiang et al. 2020), so the role of *nlg3* on vision will need to be further investigated in the future.

Notably, we observed that the knockdown of *nlg3* in all three structures individually resulted in sex differences in social space with females generally settling in a group closer together than males (**Fig. 2A-C**). This is in line with previously published results with sex differences seen in social space (Simon et al. 2012).

We then modified the transmission of the neurons within these structures to assess their individual roles in this social behaviour. Indeed, Burg et al. (2012) reported the importance of the MB in fly social space. We also observed an increase to male and female social space whether the entire MB neurons were hyperactivated or silenced (**Fig. 3A**). As the MB is not only important for learning and memory but also sensory integration, it would follow that manipulation of this circuitry would affect behavioural output.

There were no changes to social space when we hyperactivated the PB neurons but silencing the PB neurons increased female social space specifically (**Fig. 3B**) suggesting the important of the PB circuitry in regulating female social space. Next, using the *nlg3-Gal4* driver, we manipulated the transmission of all *nlg3*-expressing neurons revealing their sexually dimorphic effect on social space. Females will get further apart with hyperactivation or silencing of *nlg3*-containing neurons, but males will only get further apart with hyperactivation and not silencing of the same neurons (**Fig. 3C**). This might explain the sexually dimorphic effect of *nlg3* deletion on flies of the same age (Yost et al. 2020) are not influenced by just one of the identified brain structures *nlg3* is expressed in (MB, PB, OL), but instead may be due to the collective circuitry of all *nlg3*-expression neurons (**Fig. 1**).

The mushroom bodies influence on fly social space was clear with changes to neural transmission, so we then aimed to identify the neurotransmitter released from the mushroom bodies that affected social spacing. As expected, knocking down *ChAT* and reducing

477 acetylcholine levels in the MB decreased social space in both sexes (**Fig. 4A**). But knocking
478 down *GAD* and reducing GABA levels in the MB decreased social space only in males (**Fig.**
479 **4A**).
480
We also found that knocking down all dopamine receptor types in the mushroom bodies will
specifically increase male social space but did not affect females (**Fig. 5A-D**). Of note, Dop1R1
and Dop2R knockdown did increase female social space (**Fig. 5A,C**; Two-way ANOVA – Effect
of interaction: $F_{1,32} = 27.24$, $p < 0.0001$; Two-way ANOVA – Effect of interaction: $F_{1,32} =$
7.448, $p < 0.0001$, respectively) but this was only a small difference in the number of flies within
4BL and was at the upper end of how close flies can group together showing a ceiling effect that
produced a minimal change in their behaviour. Overall, this would indicate that the transmission
of the Kenyon cells themselves, specifically acetylcholine release, will influence female social
space but the dopaminergic inputs into the MB modify the male decision of their social space. It
was interesting to observe that all the 4 types of Drosophila dopamine receptors (Dop1R1,
Dop1R2, Dop2R, and DopEcR) all had the same effect on social space while each receptor plays
unique roles to fly behaviour such as sleep, courtship, feeding, and memory (Kim et al. 2003;
Lebestky et al. 2009; Kuo et al. 2015).
494
In Drosophila social space, many genetic and neural manipulations lead to sex differences, even
though both sexes respond to changes in their social environment in similar ways. For example,
age, mating, and isolation all lead to decreased social space in both sexes (Simon et al. 2012;
Brenman-Suttner et al. 2018; Yost et al. 2024). Due to this, we examined the influence that the
sexually dimorphic *fruP1* neuron subset had on social space. Again, we saw male-female sex
differences in response to modulation of *fruP1* neuronal transmission. For males, when *fruP1*
neurons are hyperactivated, flies get further apart, but when these neurons are silenced, males get
closer together (**Fig. 6**). Females also get further apart when *fruP1* neurons are hyperactivated,
but do not change their social space when these neurons are silenced (**Fig. 6**). This indicates that
fruP1 neurons in males can increase and decrease neuronal activity resulting in a change in their
social space, but for females only an increase in *fruP1* activity modifies their social space. The
basal transmission of these neurons that affect female social space might not be active and
therefore cannot be silenced, only hyperactivation changes this resting state and leads to females
settling further in a group setting.
509
Knowing the importance for *fruP1* neurons in Drosophila social space, we then examined the
molecular players within these neurons that are contributing to the behavioural changes.
Knocking down *nlg3* in *fruP1* neurons showed a male-female sexual dimorphism with males
getting further apart and females closer together (**Fig. 7A**). This confirms there are *fruitless*- and
nlg3-expressing neurons that are contributing to fly social space behaviour. Furthermore, our
results indicate that those neurons are not dopaminergic. However, the *fruP1* neurons affecting
social space are cholinergic, in both males and females, a phenocopy of the effect seen on
females both when reducing acetylcholine levels in the mushroom bodies and when knocking
down *nlg3* in *fruP1* neurons. Because a subset of the MB neurons is also *fruP1* neurons
(Palmateer et al. 2023), we propose that the Kenyon cells *nlg3* neurons are also *fruP1* neurons, at
least those that affect female social space (**Fig. 8**).
521

Finally, as previously reported (Jiang et al. 2020; Yost et al. 2024), we did not observe any correlation in our experiments between variations in performance index leading to any changes in social space.

Fly social behaviour relies heavily on an individual's ability to perceive and integrate social and environmental cues. Using the Drosophila social space assay, we show the roles of the OL and MB in regulating fly social spacing. The OL are essential for sensory perception, as they transduce visual cues into electrochemical signals that are integrated within MB, and the central complex, thereby enabling the fly to perceive its environment and alter its behaviour accordingly.

The significance of Nlg3 in the OL is reiterated here, as it is probably involved in the proper development and regulation of synaptic connections within these synaptically rich brain regions. Proper development of the OL layers is required to decode parallel circuitry pathways involved in processing color, shapes, and motion, which are crucial for interpreting environmental stimuli. Outside of the development and maintenance of synaptic connections that can affect fly behaviour, neural transmission through these brain regions reveals the MB to be important for social behaviour. The MB, which receives signals and integrates from multiple other brain regions (Aso et al. 2014), relies on proper development of synaptic connections where disruptions or improper regulations in these connections can lead to misinterpretation of signals, ultimately altering fly behavior through its descending output neurons.

By dissecting the molecular players within these neuronal subsets, we identify some of the inputs and outputs of the MB that influence fly social spacing. This behavioral data allows us to begin constructing a functional circuitry model of the neural pathways that regulate social space and where sexual dimorphic integration of cues might be taking place (hypothesis generating model **Fig. 8**). We have identified that dopaminergic signalling into the MB influences social space but still need to identify which dopaminergic clusters are playing a role. We can similarly test the known dopaminergic subsets to innervate the MB (PPL1, PAM, and PPL2ab). We also identified that acetylcholine signalling from the MB will affect female social space, but we have yet to identify which mushroom body output neurons (MBONs) are associated with these behavioural changes (**Fig. 8**). We can also test the known 21 MBONs to find downstream circuits that influence this social behaviour specifically.

Acknowledgments:
We want to thank Dr. Michael Strong in whose laboratory the immunocytochemistry imaging
work was performed, and for the help of his research team for the troubleshooting involved in
obtaining good quality images.
This study would not have been possible without the continued funding received by Western's
Biotron Facility and the support offered there by the staff: we are very thankful to Karen Nygard
for her technical and molecular help in immunofluorescence protocol establishment and
microscopy within the integrated microscopy unit. We also want to thank the Insect Suite within
the Biotron Facility of Western University, for housing our flies in climate-controlled walk-in
rooms (25°C). We also acknowledge Dr. Ryley T. Yost for his western blot analysis to show the
nlg3-RNAi results in a reduction of Nlg3 protein levels (data not shown).
We also want to thank Flybase (Öztürk-Çolak et al., 2024 - NHGRI Awards U41HG000739 and
U24HG010859), the Bloomington Drosophila Stock Center (NIH P40OD018537) and for the
nc82 (anti-Brp) primary antibody which was obtained from the Developmental Studies
Hybridoma Bank, created by the NICHD of the NIH and maintained at The University of Iowa,
Department of Biology, Iowa City, IA 52242, for the invaluable resources they provide to the
scientific community.
575
576 **Funding:**
577 This project was funded by internal grants to AS, internal and provincial grants to JWR, ATB
578 and JRI; internal and NSERC-USRA grant to KM; NSERC Fellowships to JWR, ATB, MRE,
579 GIR and JRI; NSERC Discovery grants 2020-06464 to AJM, and 05054-2022 to AFS.

580
581 **Ethics Statement**
582 Western's Animal Care Committee or Ontario Provincial and Federal regulatory bodies specifies
583 that ethical considerations do not apply to the study of invertebrate animals including insects
584 such as *Drosophila melanogaster*
585
586 **Data availability statement:**
587 The datasets analysed for this study will be deposited in the Dryad repository (TBD).
588
589 **Conflict of Interest:**
590 The authors declare that the research was conducted in the absence of any commercial or
591 financial relationships that could be construed as a potential conflict of interest.
592
593 **Authors contribution:**
594 Based on Oxford Academic (Genetics journal)
595
JWR: Conceptualization, Supervision, Project Administration, Investigation, Formal Analysis,
Methodology, Resources, Funding Acquisition, Writing - Original Draft Preparation, and Writing -
Review & Editing.
ATB: Conceptualization, Investigation, Formal Analysis, Funding Acquisition, and Writing -
Review & Editing.
MRE: Conceptualization, Investigation, Formal Analysis, Funding Acquisition, and Writing -
Review & Editing.
JK: Conceptualization, Investigation, Formal Analysis and Writing.
JRI: Methodology, Review & Editing.

SP: Investigation and Formal Analysis.
DSL: Investigation and Formal Analysis.
AS: Methodology.
JND: Investigation, Formal Analysis and Validation.
GIR: Writing - Review & Editing.
AJM: Methodology, Supervision, and Funding Acquisition.
AFS: Conceptualization, Supervision, Project Administration, Formal Analysis, Resources, Funding Acquisition, and Writing - Review & Editing.

References:

Aso Y, Hattori D, Yu Y, Johnston RM, Iyer NA, Ngo TTB, Dionne H, Abbott LF, Axel R, Tanimoto H, et al. 2014. The neuronal architecture of the mushroom body provides a logic for associative learning. Elife. 3:e04577. doi:10.7554/eLife.04577.

Baker CA, Guan X-J, Choi M, Murthy M. 2024. The role of *fruitless* in specifying courtship behaviors across divergent Drosophila species. Sci Adv. 10(11). doi:10.1126/sciadv.adk1273.

Brenman-Suttner D, Yost R, Frame A, Robinson J, Moehring A, Simon A. 2019. Social behaviour and aging: a fly model. Genes Brain Behav. 12598:1–18. doi:10.1111/gbb.12598.

Brenman-Suttner DB, Long SQ, Kamesan V, Belle JN De, Ryley T, Kanippayoor RL, Simon AF. 2018. Progeny of old parents have increased social space in *Drosophila melanogaster*. Sci Rep. 8(3673):1–13. doi:10.1038/s41598-018-21731-0.

Burg ED, Langan ST, Nash HA. 2013. Drosophila social clustering is disrupted by anesthetics and in narrow abdomen ion channel mutants. Genes Brain Behav. 12(3):338–347. doi:10.1111/gbb.12025.

Cao H, Tang J, Liu Q, Huang J, Xu R. 2022. Autism-like behaviors regulated by the serotonin receptor 5-HT2B in the dorsal fan-shaped body neurons of *Drosophila melanogaster*. Eur J Med Res. 27(1):203. doi:10.1186/s40001-022-00838-1.

Castermans D, Wilquet V, Parthoens E, Huysmans C, Steyaert J, Swinnen L, Fryns J-P, Van de Ven W, Devriendt K. 2003. The *neurobeachin* gene is disrupted by a translocation in a patient with idiopathic autism. J Med Genet. 40(5):352–6. doi:10.1136/jmg.40.5.352.

Chang K, Marran K, Valentine A, Hannon GJ. 2014. Generation of transgenic drosophila expressing shRNAs in the miR-1 backbone. Cold Spring Harb Protoc. 2014(5):pdb.prot080762. doi:10.1101/pdb.prot080762.

Chowdhury T, Calhoun RM, Bruch K, Moehring AJ. 2020. The *fruitless* gene affects female receptivity and species isolation. Proceedings of the Royal Society B: Biological Sciences. 287(1923):20192765. doi:10.1098/rspb.2019.2765.

651 Corthals K, Heukamp AS, Kossen R, Großhennig I, Hahn N, Gras H, Göpfert MC, Heinrich R,
652 Geurten BRH. 2017. Neuroligins Nlg2 and Nlg4 affect social behavior in *Drosophila*
653 *melanogaster*. Front Psychiatry. 8:1–13. doi:10.3389/fpsyt.2017.00113.
654
655 Fernandez RW, Akinleye AA, Nurilov M, Feliciano O, Lollar M, Aijuri RR, O'Donnell JM,
656 Simon AF. 2017. Modulation of social space by dopamine in *Drosophila melanogaster*, but no
657 effect on the avoidance of the Drosophila stress odorant. Biol Lett. 13(8):20170369.
658 doi:10.1098/rsbl.2017.0369.
659
660 Ferreira CH, Moita MA. 2019. What can a non-eusocial insect tell us about the neural basis of
661 group behaviour? Curr Opin Insect Sci. 36:1–7. doi:10.1016/j.cois.2019.09.001.
662 https://doi.org/10.1016/j.cois.2019.09.001.
663
664 Grant SG. 2012. Synaptopathies: diseases of the synaptome. Curr Opin Neurobiol. 22(3):522–
665 529. doi:10.1016/j.conb.2012.02.002.
666
667 Hahn N, Geurten B, Gurvich A, Piepenbrock D, Kästner A, Zanini D, Xing G, Xie W, Göpfert
668 MC, Ehrenreich H, et al. 2013. Monogenic heritable autism gene neuroligin impacts Drosophila
669 social behaviour. Behavioural Brain Research. 252:450–457. doi:10.1016/j.bbr.2013.06.020.
670
671 Hope KA, Flatten D, Cavitch P, May B, Sutcliffe JS, O'Donnell J, Reiter LT. 2019. The
672 Drosophila Gene Sulfateless Modulates Autism-Like Behaviors. Front Genet. 10.
673 doi:10.3389/fgene.2019.00574.
674
675 Ito H, Fujitani K, Usui K, Shimizu-Nishikawa K, Tanaka S, Yamamoto D. 1996. Sexual
676 orientation in Drosophila is altered by the satori mutation in the sex-determination gene *fruitless*
677 that encodes a zinc finger protein with a BTB domain. Proceedings of the National Academy of
678 Sciences. 93(18):9687–9692. doi:10.1073/pnas.93.18.9687.
679
680 Jamain S, Quach H, Betancur C, Råstam M, Colineaux C, Gillberg IC, Soderstrom H, Giros B,
681 Leboyer M, Gillberg C, et al. 2003. Mutations of the X-linked genes encoding *neuroligins*
682 NLGN3 and NLGN4 are associated with autism. Nat Genet. 34(1):27–29. doi:10.1038/ng1136.
683
684 Jiang L, Cheng Y, Gao S, Zhong Y, Ma C, Wang T, Zhu Y. 2020. Emergence of social cluster
685 by collective pairwise encounters in Drosophila. Elife. 9. doi:10.7554/eLife.51921.
686
687 Jiang Y, Yuen RKC, Jin X, Wang M, Chen N, Wu X, Ju J, Mei J, Shi Y, He M, et al. 2013.
688 Detection of clinically relevant genetic variants in autism spectrum disorder by whole-genome
689 sequencing. The American Journal of Human Genetics. 93(2):249–263.
690 doi:10.1016/j.ajhg.2013.06.012.
691
692 Kanellopoulos AK, Mariano V, Spinazzi M, Woo YJ, McLean C, Pech U, Li KW, Armstrong
693 JD, Giangrande A, Callaerts P, et al. 2020. Aralar sequesters GABA into hyperactive
694 mitochondria, Causing Social Behavior Deficits. Cell. 180(6):1178-1197.e20.
695 doi:10.1016/j.cell.2020.02.044.
696

Kim Y-C, Lee H-G, Seong C-S, Han K-A. 2003. Expression of a D1 dopamine receptor
dDA1/DmDOP1 in the central nervous system of *Drosophila melanogaster*. Gene Expression
Patterns. 3(2):237–245. doi:10.1016/S1567-133X(02)00098-4.

Kuo SY, Wu CL, Hsieh MY, Lin CT, Wen RK, Chen LC, Chen YH, Yu YW, Wang HD, Su YJ,
et al. 2015. PPL2ab neurons restore sexual responses in aged Drosophila males through
dopamine. Nat Commun. 6:1–10. doi:10.1038/ncomms8490.

Lawal HO, Chang HY, Terrell AN, Brooks ES, Pulido D, Simon AF, Krantz DE. 2010. The
Drosophila vesicular monoamine transporter reduces pesticide-induced loss of dopaminergic
neurons. Neurobiol Dis. 40(1):102–112. doi:10.1016/j.nbd.2010.05.008.
http://dx.doi.org/10.1016/j.nbd.2010.05.008.

Lebestky T, Chang J-SC, Dankert H, Zelnik L, Kim Y-C, Han K-A, Wolf FW, Perona P,
Anderson DJ. 2009. Two different forms of arousal in Drosophila are oppositely regulated by the
dopamine d1 receptor ortholog DopR via distinct neural circuits. Neuron. 64(4):522–536.
doi:10.1016/j.neuron.2009.09.031.

McNeil AR, Jolley SN, Akinleye AA, Nurilov M, Rouzyi Z, Milunovich AJ, Chambers MC,
Simon AF. 2015. Conditions affecting social space in *Drosophila melanogaster*. Journal of
Visualized Experiments.(105). doi:10.3791/53242.

Ni J-Q, Zhou R, Czech B, Liu L-P, Holderbaum L, Yang-Zhou D, Shim H-S, Tao R, Handler D,
Karpowicz P, et al. 2011. A genome-scale shRNA resource for transgenic RNAi in Drosophila.
Nat Methods. 8(5):405–407. doi:10.1038/nmeth.1592.

Palmateer CM, Artikis C, Brovero SG, Friedman B, Gresham A, Arbeitman MN. 2023. Single-
cell transcriptome profiles of Drosophila *fruitless*-expressing neurons from both sexes. Elife. 12.
doi:10.7554/eLife.78511.

Pan Y, Baker BS. 2014. Genetic identification and separation of innate and experience-
dependent courtship behaviors in Drosophila. Cell. 156(1–2):236–248.
doi:10.1016/j.cell.2013.11.041.

Sato K, Yamamoto D. 2020. The mode of action of *fruitless*: Is it an easy matter to switch the
sex? Genes Brain Behav. 19(2). doi:10.1111/gbb.12606.

Shih M-FM, Davis FP, Henry GL, Dubnau J. 2019. Nuclear transcriptomes of the seven
neuronal cell types that constitute the Drosophila mushroom bodies. G3
Genes|Genomes|Genetics. 9(1):81–94. doi:10.1534/g3.118.200726.

Shilpa O, Anupama KP, Antony A, Gurushankara HP. 2021. Lead (Pb)-induced oxidative stress
mediates sex-specific autistic-like behaviour in *Drosophila melanogaster*. Mol Neurobiol.
58(12):6378–6393. doi:10.1007/s12035-021-02546-z.

Simon AF, Chou M -T., Salazar ED, Nicholson T, Saini N, Metchev S, Krantz DE. 2012. A simple assay to study social behavior in Drosophila: measurement of social space within a group. Genes Brain Behav. 11(2):243–252. doi:10.1111/j.1601-183X.2011.00740.x.

Simon AF, Daniels R, Romero-Calderón R, Grygoruk A, Chang HY, Najibi R, Shamouelian D, Salazar E, Solomon M, Ackerson LC, et al. 2009. Drosophila vesicular monoamine transporter mutants can adapt to reduced or eliminated vesicular stores of dopamine and serotonin. Genetics. 181(2):525–541. doi:10.1534/genetics.108.094110.

Sokolowski MB. 2001. Drosophila: genetics meets behaviour. Nat Rev Genet. 2(11):879–890. doi:10.1038/35098592.

Sokolowski MB. 2010. Social interactions in "simple" model systems. Neuron. 65(6):780–94. doi:10.1016/j.neuron.2010.03.007.

Südhof TC. 2008. Neuroligins and neurexins link synaptic function to cognitive disease. Nature. 455(7215):903–911. doi:10.1038/nature07456.

Ueoka I, Kawashima H, Konishi A, Aoki M, Tanaka R, Yoshida H, Maeda T, Ozaki M, Yamaguchi M. 2018. Novel Drosophila model for psychiatric disorders including autism spectrum disorder by targeting of ATP-binding cassette protein A. Exp Neurol. 300:51–59. doi:10.1016/j.expneurol.2017.10.027.

Venken KJT, Schulze KL, Haelterman NA, Pan H, He Y, Evans-Holm M, Carlson JW, Levis RW, Spradling AC, Hoskins RA, et al. 2011. MiMIC: a highly versatile transposon insertion resource for engineering *Drosophila melanogaster* genes. Nat Methods. 8(9):737–743. doi:10.1038/nmeth.1662.

Vert J-P, Foveau N, Lajaunie C, Vandenbrouck Y. 2006. An accurate and interpretable model for siRNA efficacy prediction. BMC Bioinformatics. 7(1):520. doi:10.1186/1471-2105-7-520.

Wiedemann C. 2007. Fruitless aggression. Nat Rev Neurosci. 8(1):5–5. doi:10.1038/nrn2064.

Wise A, Tenezaca L, Fernandez RW, Schatoff E, Flores J, Ueda A, Zhong X, Wu C-F, Simon AF, Venkatesh T. 2015. Drosophila mutants of the autism candidate gene *neurobeachin* (*rugose*) exhibit neuro-developmental disorders, aberrant synaptic properties, altered locomotion, and impaired adult social behavior and activity patterns. J Neurogenet. 29(2–3):135–143. doi:10.3109/01677063.2015.1064916.

Xie T, Ho MCW, Liu Q, Horiuchi W, Lin C-C, Task D, Luan H, White BH, Potter CJ, Wu MN. 2018. A genetic toolkit for dissecting dopamine circuit function in Drosophila. Cell Rep. 23(2):652–665. doi:10.1016/j.celrep.2018.03.068.

Xing G, Gan G, Chen D, Sun M, Yi J, Lv H, Han J, Xie W. 2014. Drosophila *neuroligin3* regulates neuromuscular junction development and synaptic differentiation. Journal of Biological Chemistry. 289(46):31867–31877. doi:10.1074/jbc.M114.574897.

788
789 Yost RT, Robinson JW, Baxter CM, Scott AM, Brown LP, Aletta MS, Hakimjavadi R, Lone A,
790 Cumming RC, Dukas R, et al. 2020. Abnormal social interactions in a Drosophila mutant of an
791 autism candidate gene: *neuroligin 3*. Int J Mol Sci. 21(13):4601. doi:10.3390/ijms21134601.
792
793 Yost RT, Scott AM, Kurbaj JM, Walshe-Roussel B, Dukas R, Simon AF. 2024. Recovery from
794 social isolation requires dopamine in males, but not the autism-related gene *nlg3* in either sex. R
795 Soc Open Sci. 11(7). doi:10.1098/rsos.240604.
796

    Figure captions

Figure 1. Nlg3 is expressed in the calyx of the mushroom bodies, throughout the
protocerebral bridge, and in the optic lobes, and that expression is important for social
space. A-B: Immunofluorescent imaging of polyclonal antibody specific to the c-terminal
intracellular domain with anti-Nlg3 (green) on Canton-S male fly whole brains dissected at 3-4
days old. Magnification: 20X. **A.** Fluorescent imaging on posterior side of adult brain shows
three main structures expressing *nlg3* in low levels: calyx of the MB (Ca), protocerebral bridge
(PB), and the optic lobes (OL). **B.** Confocal imaging on posterior side of adult brain shows
definition of Nlg3 expression in the Ca, PB, and OL (medulla and lamina). **C-D:** Expression
patterns of MiMIC construct (*Mi{y[+mDint2]=MIC}nlg3[MI08924]*) incorporating *mCD8-GFP*
into *nlg3*. Immunofluorescence with anti-GFP (green) and nc82 (magenta). **C.** Confocal imaging
on anterior side of adult brain at 20X magnification indicates low expression of Nlg3 in the MB
and OL. **D.** Confocal imaging at 40X magnification highlighting expression in the MB. **E-F:**
Expression patters of *MiMIC nlg3-Gal4* genetic construct (*Mi{Trojan-Gal4}nlg3>UAS-*
mCD8::GFP) driving exogenous expression of *mCD8-GF*P with *nlg3*. Immunofluorescence
with anti-GFP (green) and nc82 (magenta). **E.** Confocal imaging of posterior side of adult brain
at 40X magnification indicates low expression patterns in PB and Ca. **F.** Confocal imaging of
anterior side of adult brain at 40X magnification shows expression throughout the MB. Antennal
Lobes show for reference (AL). Scale bar, 50μm. **G-H:** Social space of flies with RNAi
knockdown of *nlg3* targeted specifically to all *nlg3*-expressing cells. There is a decrease in the
average number of flies within 4 body lengths (4BL) in male (G; one-way ANOVA*: $F_{3,32}$ =*
1.771, *p* = 0.0003, post-hoc Šidák's test p<0.0001 when comparing *nlg3*>UAS-RNAi *nlg3* to all
controls) and female (H; one-way ANOVA*: $F_{3,28}$* = 3.967, *p* < 0.0001, post-hoc Šidák's test
p<0.0001 when comparing *nlg3*>UAS-RNAi *nlg3* to all controls).
822
Figure 2. Knocking down Nlg3 specifically in each of the mushroom bodies, the
protocerebral bridge or the optic lobes specifically increases fly social space in males and
females. A-C: Average number of flies within 4 body lengths (4BL) in male and female flies
with RNAi knockdown of *nlg3* targeted specifically to the mushroom bodies (A), protocerebral
bridge (B), and the optic lobes (C). **A.** The number of flies within 4BL decreased in both males
and females in flies with *nlg3* knockdown targeted to the MB (two-way ANOVA – Effect of
genotype*: $F_{1,39}$* = 14.71, *p* = 0.0004). **B.** The average number of flies within 4BL decreased in
both males and females in flies with *nlg3* knockdown targeted to the PB (two-way ANOVA –
Effect of genotype: $F_{1,40}$ = 19.02, *p* < 0.0001). **C.** Knocking down *nlg3* in the OL decreases the
number of flies within 4BL in both males and females compared to genetic control flies (two-
way ANOVA – Effect of genotype: $F_{1,31}$ = 7.191, *p* = 0.0116). Grey colour: genetic control
driver/+; pink colour: *driver/RNAi*. The results of all two-way ANOVA and *post-hoc* tests
performed can be found in **Supp. Table 2**. N = 7 – 14 for all treatments. Error bars are +/- s.e.m.
836
Figure 3. Manipulation of neural transmission within the brain structures enriched with
Nlg3 increases fly social space. A-C: Average number of flies within 4 body lengths (4BL) in
male and female flies when genetically hyperactivating or silencing neural transmission within
the MB (A), PB (B), and *nlg3*-containing neurons (C). **A.** Hyperactivation of MB neurons
decreases the number of flies within 4BL in both males and females compared to basal
transmission of MB neurons (two-way ANOVA – Effect of neural transmission: $F_{1,47}$ = 7.395, *p*

= 0.0091). Silencing MB neurons also decreases the number of flies within 4BL in both males
and females compared to basal transmission (two-way ANOVA – Effect of neural transmission:
$F_{1,30} = 14.40$, $p = 0.0007$). **B.** The number of flies within 4BL was not different in males and
females when the PB neurons are hyperactivated (two-way ANOVA - Effect of neural
transmission: $F_{1,74} = 17.72$, $p = 0.2018$) or silenced (two-way ANOVA – Effect of neural
transmission: $F_{1,31} = 3.443$, $p = 0.0730$). **C.** Hyperactivation of *nlg3*-containing neurons
decreases the number of flies within 4BL in both males and females compared to basal
transmission (two-way ANOVA – Effect of neural transmission: $F_{1,28} = 12.70$, $p = 0.0013$).
There is an interaction between sex and silencing *nlg3*-containing neurons where females have a
decrease in the number of flies within 4BL (two-way ANOVA – Effect of interaction: $F_{1,22} =$
6.169, $p = 0.0211$). Grey colour bars: inactivated genetic construct and basal transmission of
neurons; green colour bars: activation of genetic construct and hyperactivation of neurons with
Gal4 driver; pink colour bars: activation of genetic construct and silencing of neurons with *Gal4*
driver. The results of all two-way ANOVA and *post-hoc* tests performed can be found in **Supp.**
Table 2. N = 6 – 22 for all treatments. Error bars are +/- s.e.m.

Figure 4. Knocking down the acetylcholine rate-limiting biosynthesis enzyme C*hAT* in the
MB decreases fly social space. A-B: Average number of flies within 4 body lengths (4BL) in
male and female flies with RNAi knockdown of *ChAT* for acetylcholine synthesis (A) or *GAD1*
for GABA synthesis (B) specifically in the mushroom bodies. **A.** Both male and female flies
have an increased number of flies within 4BL with *ChAT* knockdown when compared to the
genetic control flies (two-way ANOVA – Effect of genotype: $F_{1,32} = 11.19$, $p = 0.0021$). **B.** The
interaction between sex and *GAD1* knockdown shows that male flies have an increased number
of flies within 4BL while females have a decreased number compared to their genetic control
flies (two-way ANOVA – Effect of interaction: $F_{1,29} = 4.276$, $p = 0.0068$). Grey colour bars:
genetic control *driver/+*; pink colour bars: *driver/RNAi*. The results of all two-way ANOVA and
post-hoc tests performed can be found in **Supp. Table 2**. N = 8 – 9 for all treatments. Error bars
are +/- s.e.m.

Figure 5. RNAi knockdown of all dopamine receptors in the MB increases social space
more in males than in females. A-D: Average number of flies within 4 body lengths (4BL) in
male and female flies knocking down dopamine receptors *Dop1R1* (A), *Dop1R2* (B), *Dop2R*
(C), and *DopEcR* (D) specifically in the mushroom bodies. **A.** *Dop1R1* knockdown in the MB of
males has a decreased number of flies within 4BL compared to genetic controls (Two-way
ANOVA – Effect of genotype: $F_{1,32} = 19.61$, $p < 0.0001$). **B.** *Dop1R2* knockdown in the MB of
males has a decreased number of flies within 4BL compared to genetic controls (Two-way
ANOVA – Effect of genotype: $F_{1,32} = 41.10$, $p < 0.0001$). **C.** *Dop2R* knockdown in the MB of
males has a decreased number of flies within 4BL compared to genetic controls (Two-way
ANOVA – Effect of genotype: $F_{1,32} = 41.10$, $p < 0.0001$). **D.** *DopEcR* knockdown in the MB of
males has a decreased number of flies within 4BL compared to genetic controls (Two-way
ANOVA – Effect of genotype: $F_{1,32} = 8.905$, $p = 0.0054$). Grey colour bars: genetic control
driver/+; pink colour bars: *driver/RNAi*. The results of all two-way ANOVA and *post-hoc* tests
performed can be found in **Supp. Table 2**. N = 9 for all treatments. Error bars are +/- s.e.m.

Figure 6. Hyperactivation of *FruP1* neurons increases social space in males and females
while silencing *FruP1* neurons only decreases social space in males. Average number of flies

within 4 body lengths (4BL) in males and females when genetically hyperactivating neural
transmission in *FruP1* neurons. Hyperactivation of *FruP1* neurons decreases the number of flies
within 4BL in both males and females compared their basal transmission (two-way ANOVA –
Effect of neural transmission: $F_{1,33}$ = 22.65, p < 0.0001). Silencing *FruP1* neurons did not alter
change female social space but increased the number of flies within 4BL in males (two-way
ANOVA with Šídák's *post-hoc* test: p = 0.0453). Grey colour bars: inactivated genetic construct
and basal transmission of neurons; green colour bars: activation of genetic construct and
hyperactivation of neurons with Gal4 driver; pink colour bars: activation of genetic construct and
silencing of neurons with Gal4 driver. The results of all two-way ANOVA and *post-hoc* tests
performed can be found in **Supp. Table 2**. N = 7 – 11 for all treatments. Error bars are +/- s.e.m.
899
Figure 7. Knocking down *nlg3* in *FruP1* neurons decreases social space, while a reduction
of acetylcholine and GABA in *FruP1* neurons both also decrease social space in male and
female flies. A-D: Average number of flies within 4 body lengths (4BL) in male and female flies
with RNAi knockdown of *nlg3* (A), *TH* for dopamine synthesis (B), *ChAT* for acetylcholine
synthesis (C) or *GAD1* for GABA synthesis (D) specifically in *FruP1* neurons. **A.** Knocking
down *nlg3* in *FruP1* has increases the number of flies within 4BL (two-way ANOVA – Effect of
genotype: $F_{1,57}$ = 16.27, p = 0.0002) **B.** *TH* knockdown in *FruP1* neurons does not change social
space in males and females (two-way ANOVA – Effect of genotype: $F_{1,29}$ = 0.413, p = 0.5255).
C. *ChAT* knockdown in *FruP1* neurons increases the number of flies within 4BL in both males
and females compared to the genetic control flies (two-way ANOVA – Effect of genotype: $F_{1,47}$
= 99.11, p < 0.0001). **D.** *GAD1* knockdown in *FruP1* neurons increases the number of flies
within 4BL in males and females (two-way ANOVA – Effect of genotype: $F_{1,42}$ = 8.478, p =
0.0057). Grey colour bars: genetic control *driver/+*; pink colour bars: *driver/RNAi*. The results of
all two-way ANOVA and *post-hoc* tests performed can be found in **Supp. Table 2**. N = 9 - 13
for all treatments. Error bars are +/- s.e.m.
915
Figure 8. Proposed theoretical model of functional neural influences on Drosophila social
space: Dopaminergic inputs into the mushroom body neurons (dark blue line) influence male
social space while cholinergic outputs from the mushroom bodies influence female social space.
Cholinergic outputs from *fruP1* neurons (light green line) also influence male and female social
space while some mushroom body neurons also express *fru*. Neuroligin protein (red dots)
enrichment is also found to be located in the optic lobes, calyx of the mushroom bodies, and
protocerebral bridge.
923
924

925   **Figure 1**
926
927

935
936

Effect of sex (p = 0.0296)
Effect of neural transmission (p < 0.0001)

Effect of interaction (p = 0.0730)

FruP1 > TrpA1

Basal Transmission   Hyperactivation   Silencing

 Figure 7

941   **Figure 8**

943
944

Supplemental Figure Captions

Supplemental Table 1. Genotypes of fly lines used. Fly line names, full genotypes, and sources are listed.

Supplemental Table 2. ANOVA table for all Two-Way ANOVAs performed for social space experiments presented in Figures 2-6. All p values with a result less than 0.05 are bolded.

Supplemental Table 3. ANOVA table for all Two-Way ANOVAs performed for climbing experiments presented in Supplemental Figures 2-5. All p values with a result less than 0.05 are bolded.

Supplemental Figure 1: Schematic of crossing scheme used to generate genotype controls for all RNAi knockdown experiments. Bolded genotypes are presenting in main figures shown. Results for both genetic controls are included in **Supplemental Figures 7-10.** (A) Genetic crossing scheme used for Cs background and exclusion of inappropriate genetic background as a control. (B) Genetic crossing scheme used for V10 background and exclusion of inappropriate genetic background as a control.

Supplemental Figure 2: Knocking down Nlg3 specifically in the Protocerebral bridge and optic lobes decreases fly climbing in males and females but not when knocking down in the mushroom bodies specifically. A-C: Startle-induced climbing represented by the mean percentage of flies reaching the top vial (performance index). Genetic constructs include RNAi knockdown of *nlg3* targeted specifically to the mushroom bodies (A), protocerebral bridge (B), and the optic lobes (C). Grey colour: genetic control *driver/+*; pink colour: *driver/RNAi*. The results of all two-way ANOVA performed are shown on the figure. N = 5 – 10 for all treatments. Errors bar are +/- s.e.m.

Supplemental Figure 3: Hyperactivation of all the brain structures enriched with Nlg3 decreases fly climbing. A-C: Startle-induced climbing represented by the mean percentage of flies reaching the top vial (performance index). Hyperactivation or silencing neurons specifically to the mushroom bodies (A), protocerebral bridge (B), and all *nlg3-expressing* neurons (C). Grey colour bars: inactivated genetic construct and basal transmission of neurons; green colour bars: activation of genetic construct and hyperactivation of neurons with *Gal4* driver; pink colour bars: activation of genetic construct and silencing of neurons with *Gal4* driver. The results of all two-way ANOVA performed are shown on the figure. N = 4 – 7 for all treatments. Error bars are +/- s.e.m.

Supplemental Figure 4: Hyperactivation and silencing of *FruP1* neurons decreases climbing in males and females. Startle-induced climbing represented by the mean percentage of flies reaching the top vial (performance index). Hyperactivation or silencing neurons specifically in *FruP1* neuron subset. Grey colour bars: inactivated genetic construct and basal transmission of neurons; green colour bars: activation of genetic construct and hyperactivation of neurons with *Gal4* driver; pink colour bars: activation of genetic construct and silencing of neurons with

990   *Gal4* driver. The results of all two-way ANOVA performed are shown on the figure. N = 9 for
991   all treatments. Error bars are +/- s.e.m.
992
Supplemental Figure 5: Knocking down *nlg3* in *FruP1* neurons decreases male climbing,
while a reduction of dopamine, acetylcholine, and GABA in *FruP1* neurons all decrease
climbing in male and female flies. A-D: Startle-induced climbing represented by the mean
percentage of flies reaching the top vial (performance index). Genetic constructs include RNAi
knockdown targeted specifically to the *FruP1* subset of neurons of *nlg3* (A), knocking down
dopamine (B), acetylcholine (C), and GABA (D). Grey colour: genetic control *driver/+*; pink
colour: *driver/RNAi*. The results of all two-way ANOVA performed are shown on the figure. N
= 6 – 15 for all treatments. Errors bar are +/- s.e.m.
1001
1002 **Supplemental Figure 6: Elevated temperature on Cs, *Trp1A/+*, and *Shi^{ts}/+* does not affect**
1003 **social space or climbing.** The genetic controls of the temperature sensitive constructs were
1004 tested in social space with the average number of flies within 4 body lengths **A-C:** and climbing
1005 with a performance of percentage of flies reaching the top vial **D-F:** in both male and female
1006 which reveals the construct alone does not affect social space or climbing.
1007
Supplemental Figure 7: Knocking down Nlg3 specifically in the optic lobes increases fly
social space in males and females but no changes are seen when knocking down in the
protocerebral bridge or mushroom bodies specifically when looking at both genetic
controls with different genetic backgrounds. Average number of flies within 4 body lengths
(4BL) in male and female flies with RNAi knockdown of *nlg3* targeted specifically to the
mushroom bodies (A), protocerebral bridge (B), and the optic lobes (C). **A.** The number of flies
within 4BL does not change in both males and females in flies with *nlg3* knockdown targeted to
the MB. Comparing the knockdown line to both control lines shows an intermediate phenotype.
B. The average number of flies within 4BL does not change in both males and females in flies
with *nlg3* knockdown targeted to the PB. Comparing the knockdown line to both control lines
shows an intermediate phenotype. **C.** Knocking down *nlg3* in the OL decreases the number of
flies within 4BL in both males and females compared to genetic control flies (two-way ANOVA
– Effect of genotype: $p = 0.0318$). Grey colour: genetic control *driver/+* and *Gal4/+*; pink
colour: *driver/RNAi*. N = 7 – 14 for all treatments. Error bars are +/- s.e.m.
1022
Supplemental Figure 8. Knocking down the acetylcholine rate-limiting biosynthesis enzyme
C*hAT* in the MB decreases female fly social space when looking at both genetic controls
with different genetic backgrounds. Average number of flies within 4 body lengths (4BL) in
male and female flies with RNAi knockdown of *ChAT* for acetylcholine synthesis (A) or *GAD1*
for GABA synthesis (B) specifically in the mushroom bodies. **A.** The interaction between sex
and *ChAT* knockdown shows that female flies have an increased number of flies within 4BL
when compared to both the genetic control flies (two-way ANOVA – Effect of interaction: $p = $
0.0079). **B.** *GAD1* knockdown shows no difference in male and female fly social space when
compared to both genetic control flies. Comparing the female *ChAT* knockdown line to both
control lines shows an intermediate phenotype. Grey colour bars: genetic control *driver/+* and
Gal4/+; pink colour bars: *driver/RNAi*. N = 8 – 9 for all treatments. Error bars are +/- s.e.m.
1034

Supplemental Figure 9: RNAi knockdown of all dopamine receptors in the MB increases male social space when looking at both genetic controls with different genetic backgrounds. The main conclusions remain unchanged when looking at both genetic controls from the V10 background even with different genetic backgrounds. Average number of flies within 4 body lengths (4BL) in male and female flies knocking down dopamine receptors *Dop1R1* (A), *Dop1R2* (B), *Dop2R* (C), and *DopEcR* (D) specifically in the mushroom bodies. **A.** *Dop1R1* knockdown in the MB of males has a decreased number of flies within 4BL compared to genetic controls (Two-way ANOVA – Effect of genotype: $F_{1,32} = 19.61$, $p < 0.0001$). **B.** *Dop1R2* knockdown in the MB of males has a decreased number of flies within 4BL compared to genetic controls (Two-way ANOVA – Effect of genotype: $F_{1,32} = 41.10$, $p < 0.0001$). **C.** *Dop2R* knockdown in the MB of males has a decreased number of flies within 4BL compared to genetic controls (Two-way ANOVA – Effect of genotype: $F_{1,32} = 41.10$, $p < 0.0001$). **D.** *DopEcR* knockdown in the MB of males has a decreased number of flies within 4BL compared to genetic controls (Two-way ANOVA – Effect of genotype: $F_{1,32} = 8.905$, $p = 0.0054$). Grey colour bars: genetic control *driver/+* and *Gal4/+*; pink colour bars: *driver/RNAi*. N = 9 for all treatments. Error bars are +/- s.e.m.

Supplemental Figure 10: Knocking down *nlg3* in *FruP1* neurons decreases female social space, while a reduction of acetylcholine in *FruP1* neurons decreases social space in male and female flies when looking at both genetic controls with different genetic backgrounds. Average number of flies within 4 body lengths (4BL) in male and female flies with RNAi knockdown of *nlg3* (A), *TH* for dopamine synthesis (B), *ChAT* for acetylcholine synthesis (C) or *GAD1* for GABA synthesis (D) specifically in *FruP1* neurons. **A.** The interaction between sex and *nlg3* knockdown shows that female flies have a decreased number of flies within 4BL when compared to both the genetic control flies (two-way ANOVA – Effect of interaction: $p = 0.0204$). Comparing the male *nlg3* knockdown to both control lines shows an intermediate phenotype. **B.** *TH* knockdown in *FruP1* neurons does not change social space in males and females. **C.** *ChAT* knockdown in *FruP1* neurons increases the number of flies within 4BL in both males and females compared to the genetic control flies (two-way ANOVA – Effect of genotype: $p = 0.0030$). **D.** The interaction between sex and *GAD1* knockdown shows that female flies have a increased number of flies within 4BL when compared to both the genetic control flies (two-way ANOVA – Effect of interaction: $p = 0.0041$). Male *GAD1* knockdown in *FruP1* neurons shows an intermediate phenotype when compared to both genetic controls. Grey colour bars: genetic control *driver/+* and *Gal4/+*; pink colour bars: *driver/RNAi*. N = 9 - 13 for all treatments. Error bars are +/- s.e.m.

1071 **Supp. Table 1**

1072

1073

1074

Fly Line Name	Stock #	Source	Genotype	*Description*
Control Strain				
Cs	64346	Seymour Benzer	+/+	Genetic control
nlg3 Mutant				
MiMIC-nlg3	58509	BDSC	*y[1]w[*];Mi{y[+mDint2]=MIC}Nlg3[MI08924]*	P-element insertion in 1st intron
Gal4 Drivers (Outcrossed 5X in Cs)				
MB-Gal4	49265	BDSC	*w[1118]; P{y[+t7.7] w[+mC]=GMR15E01-GAL4}attP2*	Mushroom body expression under the enhancer of *rut*
PB-Gal4	50422	BDSC	*w[1118]; P{y[+t7.7] w[+mC]=GMR55G08-GAL4}attP2*	Protocerebral bridge expression under the enhancer of *cmpy*
OL-Gal4	49234	BDSC	*w[1118]; P{y[+t7.7] w[+mC]=GMR10B01-GAL4}attP2*	Optic lobe expression under the enhancer of *crp*
nlg3-Gal4	76134	BDSC	*y[1] w[*]; Mi{Trojan-GAL4.2}Nlg3[MI00445-TG4.2]*	Gal4 in *nlg3*-expressing cells
fruP1-Gal4	66696	BDSC	*TI{GAL4}fru[GAL4.P1.D]/TM3, Sb[1]*	Gal4 in *fruP1*-expressing cells
UAS Effectors				
TrpA1	26263	BDSC	*w[*]; P{y[+t7.7] w[+mC]=UAS-TrpA1(B).K}attP16*	Temperature control of neuron activation
shi[ts]	44222	BDSC	*w[*]; P{w[+mC]=UAS-shi[ts1].K}3*	Temperature control of neuron silencing
ChAT25-RNAi	25856	BDSC	*y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.JF01877}attP2*	RNAi knockdown of acetylcholine receptors
Gad51-RNAi	51794	BDSC	*y[1] v[1]; P{y[+t7.7] v[+t1.8]=TRiP.HMC03350}attP40*	RNAi knockdown of glutamate receptors
TH-RNAi	n/a	Mark Wu (Johns Hopkins)	*w;;UAS-TH RNAi 3#G / TM6, Sb*	RNAi knockdown of dopamine receptors

Supp. Table 1. Genotype table of fly lines used.

Supp. Table 2

Experiment	Figure	Effect	Df	F	p	Šídák's post hoc		
Social space						Comparison Group	Compared with	p
MB>nlg3-RNAi	2A	**Sex**	(1,39)	13.97	0.0006	*Male*		
		Genotype	(1,39)	14.71	0.0004		MB>RNAi	RNAi/+	0.3492
		Sex * Genotype	(1,39)	4.009	0.0523	*Female*		
						MB>RNAi	**RNAi/+**	**0.0005**
PB>nlg3-RNAi	2B	**Sex**	(1,40)	14.90	0.0004	*Male*		
		Genotype	(1,40)	19.02	<0.0001		PB>RNAi	RNAi/+	0.2497
		Sex * Genotype	(1,40)	5.607	0.0228	*Female*		
						PB>RNAi	**RNAi/+**	**0.0001**
OL>nlg3-RNAi	2C	**Sex**	(1,31)	31.29	<0.0001	*Male*		
		Genotype	(1,31)	7.191	0.0116		OL>RNAi	RNAi/+	0.4048
		Sex * Genotype	(1,31)	0.9424	0.3392	*Female*		
						OL>RNAi	**RNAi/+**	**0.0320**
MB>TrpA1	3A	**Sex**	(1,47)	5.382	0.0247	*Male MB>Trp*		
		Neural Transmission	(1,47)	7.395	0.0091	Activated	Inactivated	0.2437
		Sex * Neural Transmission	(1,47)	0.172	0.6796	*Female MB>Trp*		
						Activated	**Inactivated**	**0.0442**
MB>Shi^{ts}	3B	Sex	(1,30)	0.680	0.4161	*Male MB>Shi*		
		Neural Transmission	(1,30)	14.40	0.0007	**Activated**	**Inactivated**	**0.0105**
		Sex * Neural Transmission	(1,30)	0.344	0.5617	*Female MB>Shi*		
						Activated	Inactivated	0.0518
PB>TrpA1	3C	Sex	(1,74)	39.94	0.0868	*Male PB>Trp*		
		Neural Transmission	(1,74)	17.72	0.2018	Activated	Inactivated	0.9999
		Sex * Neural Transmission	(1,74)	13.62	0.1953	*Female PB>Trp*		
						Activated	Inactivated	0.1382
PB>Shi^{ts}	3D	Sex	(1,31)	0.6335	0.4321	*Male PB>Shi*		
		Neural Transmission	(1,31)	3.443	0.0730	Activated	Inactivated	0.8897
		Sex * Neural Transmission	(1,31)	1.483	0.2325	*Female PB>Shi*		
						Activated	Inactivated	0.0596
Trojan>TrpA1	3E	Sex	(1,28)	2.380	0.1341	*Male Trojan>Trp*		
		Neural Transmission	(1,28)	12.70	0.0013	Activated	Inactivated	0.2214
		Sex * Neural Transmission	(1,28)	1.366	0.2523	*Female Trojan>Trp*		
						Activated	**Inactivated**	**0.0029**
Trojan>Shi^{ts}	3F	**Sex**	(1,22)	4.666	0.0419	*Male MB>Trp*		
		Neural Transmission	(1,22)	5.998	0.0228	Activated	Inactivated	0.9996
		Sex * Neural Transmission	(1,22)	6.169	0.0211	*Female MB>Trp*		
						Activated	**Inactivated**	**0.0042**
MB>CHAT25-RNAi	4A	**Sex**	(1,32)	41.09	<0.0001	*Male*		
		Genotype	(1,32)	11.19	0.0021		MB>RNAi	RNAi/+	0.4588
		Sex * Genotype	(1,32)	3.020	0.0919	*Female*		
						MB>RNAi	**RNAi/+**	**0.0022**
MB>GAD51-RNAi	4B	**Sex**	(1,29)	15.22	0.0005	*Male*		
		Genotype	(1,29)	18.10	0.0002		MB>RNAi	RNAi/+	0.2389
		Sex * Genotype	(1,29)	4.276	0.0477	*Female*		
						MB>RNAi	**RNAi/+**	**0.0003**
MB>Dop1R1-RNAi	5A	**Sex**	(1,32)	65.51	<0.0001	*Male*		
		Genotype	(1,32)	19.61	<0.0001	**MB>RNAi**	**MB/+**	**<0.0001**
		Sex * Genotype	(1,32)	27.24	<0.0001	*Female*		
						MB>RNAi	MB/+	0.0505
MB>Dop1R2-RNAi	5B	**Sex**	(1,32)	139.4	<0.0001	*Male*		
		Genotype	(1,32)	7.401	0.0105	**MB>RNAi**	**MB/+**	**0.0225**
		Sex * Genotype	(1,32)	1.168	0.2880	*Female*		
						MB>RNAi	MB/+	0.4447
MB>Dop2R-RNAi	5C	**Sex**	(1,32)	120.7	<0.0001	*Male*		
		Genotype	(1,32)	41.10	<0.0001	**MB>RNAi**	**MB/+**	**<0.0001**
		Sex * Genotype	(1,32)	7.448	0.0102	*Female*		
						MB>RNAi	**MB/+**	**0.0276**
MB>DopEcR-RNAi	5D	**Sex**	(1,32)	139.3	<0.0001	*Male*		
		Genotype	(1,32)	8.905	0.0054	**MB>RNAi**	**MB/+**	**0.0103**
		Sex * Genotype	(1,32)	1.586	0.2167	*Female*		
						MB>RNAi	MB/+	0.4098
FruP1>TrpA1	6A	**Sex**	(1,33)	5.168	0.0296	*Male FruP1>Trp*		
		Neural Transmission	(1,33)	22.65	<0.0001	**Activated**	**Inactivated**	**0.0241**
		Sex * Neural Transmission	(1,33)	0.6987	0.4092	*Female FruP1>Trp*		
						Activated	**Inactivated**	**0.0004**
FruP1>Shi^{ts}	6B	Sex	(1,30)	0.5671	0.4573	*Male FruP1>Trp*		
		Neural Transmission	(1,30)	2.350	0.1358	**Activated**	**Inactivated**	**0.0453**
		Sex * Neural Transmission	(1,30)	3.451	0.0730	*Female FruP1>Trp*		
						Activated	Inactivated	0.9676
FruP1>nlg3-RNAi	7A	**Sex**	(1,57)	15.57	0.0002	*Male*		
		Genotype	(1,57)	16.27	0.0002		FruP1>RNAi	RNAi/+	0.3363
		Sex * Genotype	(1,57)	4.090	0.0478	*Female*		
						FruP1>RNAi	**RNAi/+**	**<0.0001**
FruP1>TH-RNAi	7B	**Sex**	(1,29)	32.36	<0.0001	*Male*		
		Genotype	(1,29)	0.413	0.5255		FruP1>RNAi	RNAi/+	0.9938
		Sex * Genotype	(1,29)	0.603	0.4436	*Female*		
						FruP1>RNAi	RNAi/+	0.5709
FruP1>CHAT25-RNAi	7C	Sex	(1,47)	16.12	0.0002	*Male*		
		Genotype	(1,47)	99.11	<0.0001	**FruP1>RNAi**	**RNAi/+**	**<0.0001**
		Sex * Genotype	(1,47)	0.7134	0.4026	*Female*		
						FruP1>RNAi	**RNAi/+**	**<0.0001**
FruP1>GAD51-RNAi	7D	**Sex**	(1,42)	11.81	0.0013	*Male*		
		Genotype	(1,42)	8.478	0.0057	**FruP1>RNAi**	**RNAi/+**	**0.0363**
		Sex * Genotype	(1,42)	0.2264	0.6367	*Female*		
						FruP1>RNAi	RNAi/+	0.1897

Supp. Table 2: **ANOVA table for all Two-way ANOVAs performed.** All p-values with a result less than 0.05 are bolded.

Supp. Table 3

Experiment	Sup Figure	Effect	Df	F	p	Šídák's post hoc		
						Comparison Group	Compared with	p
Social space								
MB>nlg3-RNAi	2A	Sex	(1,26)	0.1363	0.7150	*Male*		
		Genotype	(1,26)	2.146	0.1549	MB>RNAi	RNAi/+	0.1714
		Sex * Genotype	(1,26)	0.8841	0.3557	*Female*		
						MB>RNAi	RNAi/+	0.9231
PB>nlg3-RNAi	2B	Sex	(1,23)	1.349	0.2573	*Male*		
		Genotype	**(1,23)**	**52.07**	**<0.0001**	**PB>RNAi**	**RNAi/+**	**<0.0001**
		Sex * Genotype	**(1,23)**	**4.505**	**0.0448**	*Female*		
						PB>RNAi	RNAi/+	0.0034
OL>nlg3-RNAi	2C	Sex	(1,16)	3.924	0.0651	*Male*		
		Genotype	**(1,16)**	**88.04**	**<0.0001**	**OL>RNAi**	**RNAi/+**	**<0.0001**
		Sex * Genotype	**(1,16)**	**10.45**	**0.0052**	*Female*		
						OL>RNAi	RNAi/+	0.0010
MB>TrpA1	3A	Sex	(1,18)	2.591	0.1249	*Male MB>Trp*		
		Neural Transmission	(1,18)	4.359	0.0513	Activated	Inactivated	0.4658
		Sex * Neural Transmission	(1,18)	0.1717	0.6835	*Female MB>Trp*		
						Activated	Inactivated	0.1574
MB>Shi^{ts}	3A	**Sex**	**(1,19)**	**18.95**	**0.0003**	*Male MB>Shi*		
		Neural Transmission	(1,19)	0.0243	0.8776	Activated	Inactivated	0.9857
		Sex * Neural Transmission	(1,19)	0.0029	0.9575	*Female MB>Shi*		
						Activated	Inactivated	0.9969
PB>TrpA1	3B	Sex	(1,12)	0.5633	0.4674	*Male PB>Trp*		
		Neural Transmission	(1,12)	0.3293	0.5767	Activated	Inactivated	0.7708
		Sex * Neural Transmission	(1,12)	0.1301	0.7246	*Female PB>Trp*		
						Activated	Inactivated	0.9862
PB>Shi^{ts}	3B	Sex	(1,15)	3.534	0.0797	*Male PB>Shi*		
		Neural Transmission	(1,15)	0.3622	0.5563	Activated	Inactivated	0.8414
		Sex * Neural Transmission	(1,15)	0.0168	0.8985	*Female PB>Shi*		
						Activated	Inactivated	0.9375
Trojan>TrpA1	3C	Sex	(1,20)	0.5228	0.4780	*Male Trojan>Trp*		
		Neural Transmission	**(1,20)**	**4.775**	**0.0410**	Activated	Inactivated	0.0750
		Sex * Neural Transmission	(1,20)	1.173	0.2917	*Female Trojan>Trp*		
						Activated	Inactivated	0.6693
Trojan>Shi^{ts}	3C	Sex	(1,19)	0.1124	0.7411	*Male MB>Trp*		
		Neural Transmission	(1,19)	0.2331	0.6348	Activated	Inactivated	0.9245
		Sex * Neural Transmission	(1,19)	0.0011	0.9736	*Female MB>Trp*		
						Activated	Inactivated	0.9367
FruP1>TrpA1	4	**Sex**	**(1,32)**	**25.10**	**<0.0001**	*Male FruP1>Trp*		
		Neural Transmission	**(1,32)**	**153.6**	**<0.0001**	**Activated**	**Inactivated**	**<0.0001**
		Sex * Neural Transmission	(1,32)	3.408	0.0742	*Female FruP1>Trp*		
						Activated	**Inactivated**	**<0.0001**
FruP1>Shi^{ts}	4	Sex	(1,32)	0.0873	0.7695	*Male FruP1>Trp*		
		Neural Transmission	**(1,32)**	**33.54**	**<0.0001**	**Activated**	**Inactivated**	**<0.0001**
		Sex * Neural Transmission	**(1,32)**	**4.278**	**0.0468**	*Female FruP1>Trp*		
						Activated	**Inactivated**	**0.0257**
FruP1>nlg3-RNAi	5A	**Sex**	**(1,40)**	**3.688**	**0.0140**	*Male*		
		Genotype	**(1,40)**	**9.735**	**0.0033**	**FruP1>RNAi**	**RNAi/+**	**0.0031**
		Sex * Genotype	(1,40)	6.601	0.0620	*Female*		
						FruP1>RNAi	RNAi/+	0.6103
FruP1>TH-RNAi	5B	Sex	(1,31)	0.3099	0.5817	*Male*		
		Genotype	**(1,31)**	**18.07**	**0.0002**	FruP1>RNAi	RNAi/+	0.0712
		Sex * Genotype	(1,31)	1.229	0.2762	*Female*		
						FruP1>RNAi	**RNAi/+**	**0.0011**
FruP1>CHAT25-RNAi	5C	Sex	(1,25)	18.82	0.1822	*Male*		
		Genotype	**(1,25)**	**26.26**	**<0.0001**	**FruP1>RNAi**	**RNAi/+**	**0.0078**
		Sex * Genotype	(1,25)	0.2810	0.6007	*Female*		
						FruP1>RNAi	**RNAi/+**	**0.0008**
FruP1>GAD51-RNAi	5D	Sex	(1,38)	3.336	0.0757	*Male*		
		Genotype	(1,38)	2.111	0.1544	FruP1>RNAi	RNAi/+	0.5132
		Sex * Genotype	(1,38)	0.0010	0.9742	*Female*		
						FruP1>RNAi	RNAi/+	0.5367

Supp. Table 3: Two-way ANOVAs performed for climbing assays. All p-values with a result less than 0.05 are bolded.

Supp. Figure 1

Climbing

Climbing

Climbing

Males   Females

Effect of sex (p = 0.0431)
Effect of genotype (p = 0.0008)
Effect of interaction (p = 0.0370)

A

FruP1 > *nlg3*-RNAi

Effect of genotype (p < 0.0001)

FruP1 > *TH*-RNAi

Effect of genotype (p < 0.0001)

FruP1 > *CHAT*-RNAi

D

Effect of genotype (p = 0.0026)

FruP1 > *GAD*-RNAi

Gal4 Control   RNAi Knockdown

 Supp Figure 7

Males    Females

Effect of sex (p = 0.0009)
Effect of genotype (p < 0.0001)
Effect of interaction (p = 0.0002)

A

MB > *nlg3*-RNAi

Effect of sex (p < 0.0001)
Effect of genotype (p < 0.0001)
Effect of interaction (p = 0.0379)

B

PB > *nlg3*-RNAi

Effect of sex (p < 0.0001)
Effect of genotype (p = 0.0318)
Effect of interaction (p = 0.2616)

C

OL > *nlg3*-RNAi

Gal4 Control  RNAi Knockdown  UAS Control

 Supp Figure 8

Effect of sex (p < 0.0001)
Effect of genotype (p = 0.0016)
Effect of interaction (p = 0.0079)

MB > *CHAT*-RNAi

Effect of sex (p < 0.0001)
Effect of genotype (p = 0.0002)
Effect of interaction (p = 0.0041)

MB > *GAD*-RNAi

Number of Flies Within 4BL

Gal4 Control   RNAi Knockdown   UAS Control

Supp Figure 10

Males   Females

Effect of sex (p = 0.0024)
Effect of genotype (p = 0.0001)
Effect of interaction (p = 0.0204)

A

FruP1 > *nlg3*-RNAi

Effect of sex (p < 0.0001)
Effect of genotype (p = 0.2472)
Effect of interaction (p = 0.1246)

B

FruP1 > *TH*-RNAi

Effect of sex (p = 0.1510)
Effect of genotype (p = 0.0030)
Effect of interaction (p = 0.0034)

C

FruP1 > *CHAT*-RNAi

Effect of sex (p < 0.0001)
Effect of genotype (p = 0.0002)
Effect of interaction (p = 0.0041)

D

FruP1 > *GAD*-RNAi

Gal4 Control   RNAi Knockdown   UAS Control