Neuroligin 3 highlights sexually dimorphic circuitry in Drosophila social spacing

Authors:

Robinson JW[1], Bechard AT[1], Evans MR[1], Mosuro K[1†], Isaacson JR[1], Pillay S[1†], Lin DS[1†], Sahota A[1†], de Belle JN[1†], Robinson GI[2], Strong MJ[3,4], Simon, AF[1*]

Affiliations:
1. Department of Biology, University of Western Ontario, London, ON, Canada
2. Department of Biological Sciences, University of Lethbridge, Lethbridge, AB, Canada
3. Molecular Medicine Group, Robarts Research Institute, Schulich School of Medicine & Dentistry, University of Western Ontario, London, ON, Canada
4. Department of Clinical Neurological Sciences, Schulich School of Medicine & Dentistry, University of Western Ontario, London, ON, Canada

† Undergraduate students at time of data gathering
* Correspondence: asimon28@uwo.ca (A.F.S.)

Running head
Drosophila social spacing neurocircuitry

Keywords:
Drosophila, social space, social behaviour, neural circuitry, *neuroligin*, mushroom bodies, *fruitless*, acetylcholine, dopamine, GABA, receptors

Abstract:

In *Drosophila melanogaster*, the autism-related Neuroligin 3 (Nlg3) protein is a postsynaptic membrane protein important for synapse development and regulation, which plays a role in social spacing behaviour. Here, we report the localization of Nlg3 to the calyx of the mushroom bodies (MB), optic lobes (OL), and protocerebral bridge (PB). Using RNA interference, *nlg3* knockdown in the OL increased fly social space, especially in the males, while knocking down *nlg3* independently in the MB or the PB did not change social spacing. However, hyperactivation and silencing of these neurons in the MB, but not the PB, controls social space in males and females, while hyperactivating and silencing of all *nlg3*-expressing neurons, including within the MB, PB, and OL, regulates male and female social space. Knocking down neurotransmitter biosynthesis enzymes, which decreases the amount of neurotransmitter release, showed that reducing acetylcholine release from the MB decreased female social space, whereas knocking down any dopamine receptor in the MB increased male social space. Lastly, to investigate the sexually dimorphic effects on social spacing previously seen in *nlg3* mutants, we examined a subset of sexually dimorphic *fruitless*-expressing *(fru)P1* neurons known to regulate sexually dimorphic behaviours. Hyperactivation of those *fruP1* neurons decreased social space in both sexes, while silencing those *fruP1* neurons specifically increased male social space without affecting females. Our findings highlight a sex-specific social space neural circuitry that includes the OL, MB, and *fruP1* neurons, while uncovering the underlying basis of some of the sex differences in this behaviour.

Article Summary:

In vinegar flies (*Drosophila melanogaster*), the autism-related Neuroligin 3 protein (Nlg3) controls neuronal development and regulation, but also affects fly social behaviour. Nlg3 is localized to the mushroom bodies (MB), protocerebral bridge, and the optic lobes. Genetically reducing Nlg3 in the optic lobes made male flies more socially distant, while hyperactivating or silencing neurons in the MB affected social behavior in both sexes. Reducing acetylcholine release from the MB affects female social space, while reducing dopamine receptors of the MB only affect male behaviour. Finally, the *fruitless* sexually dimorphic neurons control social behavior differently in males and females.

Introduction:

Social behaviours are fundamental to the interaction and communication within a community and are influenced by environmental cues which are processed and integrated in the brain. Abnormal social behaviours, such as difficulties with communication or interactions with others, are clinical characteristics of individuals with autism spectrum disorders (ASD). Using *Drosophila melanogaster* we can dissect the neural and genetic underpinnings of social interactions in a simplified system (Sokolowski 2001).

Flies exhibit social behaviours that, while simpler, can reflect the basic principles of social interaction found in more complex organisms (Sokolowski 2010). By examining how environmental cues are integrated and influence social behaviours in Drosophila, we can gain insights into the mechanisms that might be disrupted in ASD, providing a basis for understanding the molecular and neural circuitries involved in these disorders (Ferreira and Moita 2019).

An important component of Drosophila social interactions within a group context is social spacing – the distance of one individual to another in a group. We use the social space assay to assess the outcome of group behaviour displayed by flies as a measure of the average number of flies within four-body lengths (4BL) in the social space chamber (Simon et al. 2012). We and other research groups have used this assay as a metric for behavioural analysis to understand and reveal the basic underlying neural circuitry involved (Burg et al. 2013; Fernandez et al. 2017; Xie et al. 2018; Yost et al. 2020).

So far, we know that mutations in genes affecting eye function, have been found to make flies less social (Simon et al. 2012; Burg et al. 2013; Jiang et al. 2020). Both males and females have decreased social space with mutants associated to fly gustation and mechanosensation (Jiang et al. 2020). In contrast, *Orco* mutants, which affect olfactory perception, do not appear to influence social spacing in males, but result in decreased social space in females. Together, these suggest a sensory modality-specific genetic influence on social behaviour that can have sex-specific effects (Simon et al. 2012; Jiang et al. 2020).

Other genes expressed in the brain have also been shown to influence Drosophila social behaviour, such as genes important for synaptic transmission and synaptogenesis (Burg et al. 2013; Wise et al. 2015; Fernandez et al. 2017; Ueoka et al. 2018; Brenman-Suttner et al. 2019; Hope et al. 2019; Jiang et al. 2020; Kanellopoulos et al. 2020; Yost et al. 2020; Shilpa et al. 2021; Cao et al. 2022). At the synapse level, the concentration of neurotransmitters, which is tightly regulated by biosynthesis enzymes or vesicular packaging proteins, have been shown to affect social spacing. Tyrosine hydroxylase (TH), the rate-limiting enzyme in dopamine synthesis, and the vesicular monoamine transporter (VMAT), which packages dopamine into synaptic vesicles, both regulate dopamine levels (Simon et al. 2009; Lawal et al. 2010). Modulating the abundance or activity of these proteins influences social spacing in Drosophila, indicating that dopamine plays a role in regulating social space (Fernandez et al. 2017).

Many human homologs of neural genes have also emerged as potential candidates in neuropsychiatric disorders, bridging the gap between basic biological processes and complex

human pathologies (Grant 2012). The gene *neurobeachin*, which is implicated in ASD in
humans, encodes a structural scaffolding postsynaptic density protein (Castermans et al. 2003).
In Drosophila, mutations in the homolog *rugose* result in altered fly social spacing, indicating a
conservation of function across species (Wise et al. 2015). Furthermore, synaptic proteins
Neuroligin and Neurexin, which facilitate neurotransmission by aiding synapse formation and
receptor recruitment, have been shown to influence social behaviour (Hahn et al. 2013). In
humans, *Neuroligin* gene homologs are correlated with ASD (Jamain et al. 2003; Südhof 2008).
Similarly, in Drosophila, mutations in these genes lead to altered social behaviours, further
revealing the genetic underpinnings of social space (Corthals et al. 2017).
89
Drosophila Neuroligin (Nlg) is a family of post-synaptic membrane proteins with the conserved
function of synapse formation and maintenance by interacting with presynaptic Neurexin to
mediate cell-cell adhesion and facilitate neurotransmission (Südhof 2008). During Drosophila
neural development, larval *nlg3* mutants have a higher number of synaptic connections but less
GluRIIA recruitment and synaptic transmission compared to control flies (Xing et al. 2014).
Furthermore, *nlg3* knockouts display a sexually dimorphic and age-associated effect on fly
aggression and social space (Yost et al. 2020). Mutant *nlg3* flies had significantly less climbing
in both males and females by over 30%, however, *nlg3* deletion decreases only male, but not
female aggression; while increasing male social space in 3-4 day old flies, and in both sexes by
7-10 days old (Yost et al. 2020).
100
The mushroom bodies (MB), also known to play a role in Drosophila social spacing, are a well-
studied invertebrate brain structure primarily involved in fly olfactory learning and memory. The
MB are comprised of approximately 2,500 neurons and its unusually structured shape suggests a
highly specialized organization and function (Erber et al. 1987). The MB have a flexible
organization in response to social isolation and are the central processing structure of the
Drosophila brain (Technau 1984). Consisting of densely packed neurons called Kenyon cells, the
MB integrates many sources of sensory information to modulate fly behavioural responses.
Social space assays have been conducted on flies with a mutation in an ion channel selectively
targeted in the MB and lead to decreased social space, observing that the cholinergic signaling
on its own can rescue this altered phenotype in social spacing (Burg et al. 2013).
111
Sexually dimorphic behaviours in Drosophila such as male courtship, female receptivity, and
social interactions are known to be regulated by *fruitless* (*fru*) neurons (Pan and Baker 2014;
Brenman-Suttner et al. 2018; Chowdhury et al. 2020). The *fruitless* gene encodes a transcription
factor important for sex determination in Drosophila development and produces male-specific
transcripts that are essential for the development and function of male-specific neural circuits
(Sato and Yamamoto 2020). Regulated by multiple promoters each expressing unique
transcripts, the *fruP1* transcript undergoes sex-specific splicing and contributes to the sexual
differentiation of neural circuits influencing behaviours such as courtship and aggression
(Wiedemann 2007; Baker et al. 2024). Promoters *p2-p5* are necessary for adult viability and fly
morphology that is however not sex specific (Ito et al. 1996). Neurons expressing *fru* are
widespread interneurons in the adult fly brain and ventral nerve cord, where fast acting *fru*
neurons release acetylcholine, GABA, and glutamate, while other *fru* neurons also release
dopamine, serotonin, tyramine, and octopamine (Palmateer et al. 2023).
125

While progress has been made in identifying individual genes, synaptic proteins, and neuropils
like the MB that influence Drosophila social spacing, the integration of molecular players within
precise circuitry or other brain structures mediating social spacing behaviour are not fully
understood. This research looks deeper into the role *nlg3* has on Drosophila social space. Based
on Nlg3 localization and the sex-specific differences observed, we elucidated part of the neural
circuitry that directly influences social spacing behaviour.

Materials and Methods:

Fly Stocks and Husbandry

All flies were reared mixed sex in bottles containing our own fly food recipe (brown sugar, corn
meal, yeast, agar, benzoic acid, methyl paraben, and propionic acid). Rearing conditions were
controlled during development and aging at 50% humidity, 25ºC, and on a 12:12 hour light:dark
cycle. Parents were removed after seven days of egg laying resulting in a maximum parental age
of 14 days, as parental age affects social spacing of the offspring (Brenman-Suttner et al. 2018).
Canton-S (Cs) control line was initially sourced from Seymour Benzer in 1998. UAS-effector
lines were outcrossed five times into the Cs control line to minimize behavioural variation due to
genetic background. A generalized crossing scheme used to create the appropriate experimental
treatments and genetic controls can be found in the supplementary material, figure S1.

nlg3 RNAi Generation

The shRNA sequences targeting nlg3 were obtained by using the model provided by Vert et al.
(2006). The length of the *nlg3* shRNA was 60 bases for the full hairpin; and was ~7.8kbp for the
entire pVALIUM20 plasmid containing the shRNA was ~7.8kbp. The hairpin sequence used
was:

CAGCAACTCGAACTCGAACTAtagttatattcaagcataTAGTTCGAGTTCGAGTTGCTG

Within the hairpin sequence, uppercase letters represent the nucleotides that bind to the *nlg3*
mRNA, while lowercase letters represent the nucleotides that connect the two halves of the pin.
The shRNA constructs were cloned into the vector pVALIUM20 (vermillion eye color maker,
(Ni et al. 2011) using the protocol in Chang et al. (2014). DNA was injected into genotype *y v,*
nanos-integrase, attP40/attP40 embryos and were mated to the parental line after reaching
adulthood. Then, offspring containing the transgene were backcrossed five times into the Cs
control background (Vert et al. 2006; Chang et al. 2014).

Immunocytochemistry

Anti-Nlg3
Whole brain dissections on 3 to 4 day old, male or female brains were fixed within 20 minutes of
dissection in Bouin's solution. Following fixation and washing, brains were blocked in 5% NGS
in 1X PBS and stained with guinea pig anti-Nlg3 primary antibody (1:1000, Invitrogen)
overnight at 4 °C, which is a polyclonal primary antibody kindly provided by Dr. Brian Mozer
(Yost et al. 2020). After washing, brains were stained with goat anti-guinea pig Alexa 488

(1:500, Invitrogen) for 2 hours at room temperature. Next, all samples were left in glycerol for 1
hour then were mounted on slides using Fluoroshield™ media, stored at 4 °C and imaged the
following day. Fluorescent images were visualized with a AxioImager Z1 Compound
Fluorescent microscope (Zeiss). Note: we used the entirety of our polyclonal antibody stock and
have been unsuccessful in generating another functioning antibody for Nlg3. We switched to
genetic constructs that tag Nlg3 with a GFP reporter (*MiMIC-nlg3* and *nlg3-Gal4 > mCD8-*
GFP) that both verify our findings.

Anti-GFP
Whole brain dissections on 3 to 4 day old males or females were fixed within 20 minutes of
dissection in 4% paraformaldehyde in 1X PBS-T (0.3% Triton-X). Following fixation and
washing, brains were blocked in 5% normal goat serum (NGS) in PBS-T and stained with rabbit
anti-GFP (1:1000, Invitrogen) and mouse anti-Brp (nc82; 1:30, Developmental Studies
Hybridoma Bank) for 48 hours at 4 °C. After washing, brains were stained with Alexa 660 anti-
mouse (1:400, Invitrogen) and Alexa 488 anti-rabbit (1:800, Invitrogen) for 48 hours at 4°C.
Samples were mounted on slides in Fluoroshield™ media, stored at 4 °C and imaged the
following day. Slides were imaged on an SP8 Lightening Confocal microscopy system (Leica).
All images were visualized with LAS X 2.0 software (Leica).

Social Space Assay

The assay was performed as previously described (McNeil et al. 2015; Yost et al. 2020). In
short, flies were collected under CO_2 after adult eclosion and aged at 3 to 4 days old. Twenty-
four hours prior to the assay, flies were sexed under cold anesthesia and separated into vials with
food containing 12-17 flies for each treatment tested. Two hours before the assay, each treatment
was transferred into new vials and acclimated to standard conditions (25°C and 50% humidity).
Ten minutes prior to test start, if the standard acclimation temperature is different than testing
condition temperature, treatments were set to the testing temperature required for neuronal
hyperactivation (*TrpA1* at 27°C and 50% humidity) or silencing (*Shi^{ts}* at 29°C and 50%
humidity). At the time of testing, flies were loaded into the triangular chamber and knocked to
the bottom of the chamber to induce a startle response that forces the flies into a group at the top
of the chamber. Flies are then given time to settle at their preferred social space. Each treatment
was loaded into the social space apparatus and photos were taken between 20-40 minutes or as
soon as flies have settled as a group (McNeil et al. 2015). Social space assay photos were
processed with ImageJ (National Institute of Health, Bethesda, MA, USA) to calculate the
number of flies within a four-body length (4BL or ~1cm) from macros (as described in Yost et
al. 2020). Image analysis data were imported to GraphPad Prism 10 software for statistical
analysis and graphical representation.

Climbing Assay

The assay was performed as previously described (McNeil et al. 2015; Yost et al. 2020). In
short, flies were aged to 2 to 4 days old and 24 hours prior to conducting the assay were sexed
under cold anesthesia and separated in vials with food containing 40 flies for each treatment.
Two hours before the assay, each treatment was transferred into new vials and acclimated to
testing conditions (25ºC and 50% humidity). Ten minutes before the assay, if the standard

acclimation temperature is different than testing condition temperature, treatments were set to
testing temperature required for neuronal hyperactivation (*TrpA1* at 27°C and 50% humidity) or
silencing (*Shi^{ts}* at 29°C and 50% humidity). Each treatment was transferred into a testing tube
and inserted into the climbing assay. Flies were forced to the bottom of the tube and given 15
seconds to climb to the tube on top of the apparatus. This timeframe was chosen to allow
approximately 80% of control flies to reach the top vial. After 15 seconds, the apparatus was
closed and the number of flies in the top and bottom tubes were counted to calculate a
performance index of percent flies that reached the top vial.
226
227   *Statistical Analysis*
228
Statistical analysis and graphical representation were performed in GraphPad Prism
(RRID:SCR_002798, version 7.0a for Mac, GraphPad Software, La Jolla, CA, USA,
www.graphpad.com). Normality test was used to verify normality and homoscedasticity test was
used to verify the homoscedasticity of the distribution before performing an ANOVA test. Two-
way ANOVAs were performed on data presented on the same graph, which were collected
simultaneously and had 2 variables in 2 groups. Following the ANOVAs, Šídák's post-hoc tests
were performed within each sex to compare between different genotypes or transmission states,
while correcting for multiple comparisons. For all graphs, an alpha level of 0.05 was used and
only significant values ($p < 0.05$) were shown for the two-way ANOVAs and post hoc tests. The
results of all the comparison tests for social space analysis, significant or not, can be found in
Table 2. Supplemental figures have ANOVA results on graphs.
240
241   *Drawings:*
242   Drawings presented in figures were created on the app Procreate 5.3.5 (Savage Interactive Pty
243   Ltd, Hobart, Australia).
244

Results:

Neuroligin 3 protein is expressed at low levels in the calyx of the mushroom bodies, the protocerebral bridge, and the optic lobes

We investigated the expression patterns of the Nlg3 protein in the adult brain using an anti-Nlg3 polyclonal antibody. We found Nlg3 protein was localized within the calyx (Ca) of the mushroom bodies (MB), the protocerebral bridge (PB), and the optic lobes (OL) all in low levels compared to the background autofluorescence of brain lipids (**Fig. 1A,B**). For confirmation of the findings done with our antibody, we used two genetic constructs (*MiMIC* p-element insertion within *nlg3*, as well as an *nlg3-Gal4* insertion within the gene – see Table 1 for specific localization). The *MiMIC-nlg3* insertion which incorporates a GFP reporter into the coding sequence of the host gene *nlg3* (Venken et al. 2011). From this construct we saw GFP expressed at low levels to the MB and OL and at very low fluorescence we can see the PB when compared to background autofluorescence of the Drosophila brain (**Fig. 1C,D**). Confirmation of expression in the calyx and PB was observed with the *nlg3-Gal4* insertion driving *mCD8-GFP* at higher levels than endogenous *nlg3* (**Fig. 1E,F**). Male and female localization of Nlg3 did not appear to be different either qualitatively or quantitatively in 3-4 day old adult flies (representative images of only the males are shown).

Neuroligin 3 expression in the optic lobes alters fly social spacing, mostly in males

To determine the behavioural function of Nlg3 located in these structures, we targeted RNAi knock down of *nlg3* independently in each of the MB, PB, and OL and tested fly social space. We observed no effect of knocking down *nlg3* in the MB (**Fig. 2A**; Two-way ANOVA – Effect of genotype: $F_{1,39}$ = 1.005, p = 0.3223) and PB (**Fig. 2B**; Two-way ANOVA – Effect of genotype: $F_{1,37}$ = 0.068, p = 0.7951) when compared to control flies. However, knocking down *nlg3* in the optic lobes did reduce the number of flies within 4BL and increased fly social space (**Fig. 2C**; Two-way ANOVA – Effect of genotype: $F_{1,37}$ = 4.837, p = 0.0354), an effect mostly driven by the males. Then, we used a climbing assay to assess fly locomotion and their ability to respond to a startle response and climb; all to ensure the treatment flies could perform in the social space assay. While there was no reduction in climbing when knocking down *nlg3* in the MB (**Supp. Fig. 2A**), there was a reduction in climbing when knocking down *nlg3* in each of the PB and OL (**Supp. Fig. 2B,C**). We observed that the flies were still able to climb and therefore would not confound the fly decision making for social distance, but rather show a decrease in their performance index on climbing. We also observed the presence or absence of defects in climbing is not correlated to any changes in social space, however, we continued to check climbing to ensure the flies were healthy enough to be able to climb.

Hyperactivation and silencing nlg3-expressing neurons, independently or simultaneously, changes fly social space in a sex-specific manner

Next, we looked at genetically manipulating neural transmission within the brain structures where Nlg3 is localized (**Fig. 1**) to examine their effect on social space. When hyperactivating all neurons within the MB, we saw a lower number of flies within 4BL (increased social space) in both males and females (**Fig. 3A**; Two-way ANOVA – Effect of neural transmission: $F_{1,47}$ =

7.395, $p = 0.0091$). Silencing all MB neurons showed the same reduction in the number of flies
within 4BL again in both males and females (**Fig. 3A**; Two-way ANOVA – Effect of neural
transmission: $F_{1,30} = 14.40$, $p = 0.0007$) highlighting the importance of the MB in determining
fly social space on their own. In contrast, hyperactivation of the PB neurons did not change fly
social space, nor did silencing PB neurons in males but looks to have an effect in females
specifically (**Fig. 3B**; Two-way ANOVA – Effect of neural transmission: $F_{1,31} = 3.443$, $p =$
0.0730). Finally, we used the *nlg3-Gal4* insertion to manipulate the transmission of all *nlg3*-
expressing neurons. Hyperactivation of all *nlg3*-expressing neurons decreases the number of
flies within 4BL in both males and females (**Fig. 3C**; Two-way ANOVA – Effect of neural
transmission: $F_{1,28} = 12.70$, $p = 0.0013$). Silencing these neurons only increased social space in
females but not males (**Fig. 3C**; Two-way ANOVA – Effect of interaction: $F_{1,22} = 6.169$, $p =$
0.0211). There was no reduction in climbing with a hyperactivation or silencing of the MB or PB
neurons (**Supp. Fig. 3A,B**), but we did observe a reduction in climbing when hyperactivating
nlg3-expressing neurons in males only (**Supp. Fig. 3C**).
305
306   *Reducing acetylcholine release from the mushroom bodies decreases female social space, but*
307   *knocking down dopamine receptors in the mushroom bodies increases male social space*
308
As the MB are important for fly social space, we investigated which neurotransmitters in the MB
were important for the observed effects. Using UAS-RNAi, we knocked down the rate-limiting
biosynthesis enzymes within the MB (Shih et al. 2019), including *choline acetyltransferase*
(ChAT for acetylcholine production) and *glutamate decarboxylase 1* (*GAD1* for GABA
production; Shih et al. 2019). Acetylcholine reduction in the MB decreased fly social space but
only in females (**Fig. 4A**; Two-way ANOVA – Effect of interaction: $F_{1,30} = 8.465$, $p = 0.0068$).
GABA reduction did not change the number of flies within 4BL in either males and females
(**Fig. 4B**; Two-way ANOVA – Effect of genotype: $F_{1,31} = 0.021$, $p = 0.8851$).
317
Previous research shows global dopaminergic changes in the adult fly brain will affect social
space (Fernandez et al. 2017; Xie et al. 2018; Yost et al. 2020), however, the anatomical
locations within the adult fly brain where alterations in dopaminergic transmission affect social
behaviour is unknown. Thus, we tested the dopaminergic inputs into the MB, and found that
knocking down every dopamine receptor within the MB – including Dop1R1 (**Fig. 5A**; Two-
way ANOVA and Šídák's *post-hoc* test: $p < 0.0001$), Dop1R2 (**Fig. 5B**; Two-way ANOVA and
Šídák's *post-hoc* test: $p = 0.0225$), Dop2R (**Fig. 5C**; Two-way ANOVA and Šídák's *post-hoc*
test: $p < 0.0001$), and DopEcR (**Fig. 5D**; Two-way ANOVA and Šídák's *post-hoc* test: $p =$
0.0103) – decreased the number of male flies within 4BL but did not change female social space.
327
328   *Neurons expressing the fruitless gene under the P1 promotor are involved in sexually dimorphic*
329   *responses in fly social spacing*
330
Since male and female flies appear to have differently neural circuitry underlying their social
spacing preferences, we looked at a subset of sexually dimorphic *fruitless*-expressing neurons.
We used the *fruP1-Gal4* driver to manipulate neural transmission. While hyperactivation of
fruP1 neurons decreased the number of flies within 4BL for both males and females (**Fig. 6**;
Two-way ANOVA – Effect of neural transmission: $F_{1,33} = 22.65$, $p < 0.0001$). silencing of *fruP1*
resulted in a sex specific effect. Silencing *FruP1* neurons resulted in a higher number of flies

within 4BL for males (**Fig. 6**; Two-way ANOVA and Šídák's *post-hoc* test: $F_{1,30}$ = 3.451, p =
0.0453) but did not change female social space. Importantly, we saw a reduction in climbing in
both males and females for hyperactivating and silencing *fruP1* neurons (**Supp. Fig. 5**).
340
To assess the effect of Nlg3 in *fruitless*-expressing neurons, we drove an RNAi against *nlg3* with
the *fruP1* driver. We saw reducing levels of Nlg3 had a sexually dimorphic effect on flies.
Knocking down *nlg3* in *fruitless*-expressing neurons resulted in a decreased number of flies
within 4BL for males, while females had an increased number of flies within 4BL (**Fig. 7A**;
Two-way ANOVA – Effect of interaction: $F_{1,58}$ = 7.056, p = 0.0102). Also, we observed a
reduction of male climbing when knocking down *nlg3* in *fruP1* neurons (**Supp. Fig. 6A**). Next,
we investigated the neurotransmitters involved in fly social space in *fruP1* neurons. Knocking
down *TH*, the rate limiting-enzyme in the pathway of dopamine synthesis, in *fruP1* neurons did
not change male or female social space (**Fig. 7B**; Two-way ANOVA – Effect of genotype: $F_{1,31}$
= 1.134, p = 0.2951). However, knocking down *TH* in *fruP1* neurons decreased climbing in both
males and females indicating the importance of dopamine released from *fruP1* neurons on fly
climbing (**Supp. Fig. 6B**). In contrast, reduction of acetylcholine levels by knocking down *ChAT*
in *fruP1* neurons increased the number of flies within 4BL in both males and females (**Fig. 7C**;
Two-way ANOVA – Effect of genotype: $F_{1,48}$ = 38.01, p < 0.0001). Acetylcholine release
reduction from *fruP1* neurons also decreases fly climbing (**Supp. Fig. 6C**). While a reduction of
GABA by knocking down *GAD1* in *fruP1* neurons decreased female social space but did not
change male social space (**Fig. 7D**; Two-way ANOVA – Effect of interaction: $F_{1,39}$ = 9.327, p =
0.0041), reduction of GABA resulted in a small reduction in climbing of male and female flies
(**Supp. Fig. 6D**).
360
361

Discussion:

Social interactions within a group requires an integration of both attractive and repulsive social cues that lead to a behavioural output. Here, we expand the map of the neural basis for social behaviour by focusing on the sexually dimorphic role of *neuroligin 3* and its associated adult brain structures in male and female *Drosophila melanogaster*.

We show for the first time that *nlg3* is expressed at low levels in the mushroom bodies, optic lobes, and protocerebral bridge in the adult fly brain. Individually, *nlg3* in the OL is important to maintain fly social space, specifically in males, while knockdown of *nlg3* in the MB and PB did not have any effect. Next, we need to assess the effect of *nlg3* knockdown in all *nlg3* neurons simultaneously, to determine whether we see a stronger effect than what we observed only with the OL-driver. The effects of Nlg3 in the OL on social behaviour was expected as vision is known to be important for proper social spacing (Simon et al. 2012; Burg et al. 2013; Jiang et al. 2013).

Additionally, manipulating total neural transmission in these brain regions highlighted the importance of the Kenyon cells for maintaining social space. Hyperactivating and silencing MB neurons in both males and females increased fly social space. Reducing acetylcholine release from the MB decreased female social space, whereas knocking down all dopamine-type receptors increased male social space. These findings suggest that mapping the circuitry through the MB happens in a sexually dimorphic manner, where total acetylcholine release from the MB will modify female social space and dopaminergic inputs to the MB will change male social space. We still do not know the specific output from the MB leading to changes in males or the inputs to the MB leading to changes in females.

We also examined the involvement of *fru* neurons, which are known to regulate other sexually dimorphic behaviors such as male courtship or female receptivity (Chowdhury et al. 2020). Hyperactivating *fruP1* neurons decreased social space in both sexes, while silencing them showed to be sexually dimorphic with increased social space only occurring in males. Again, this demonstrates the influence *fruP1* neurons have on social space along with other fly social behaviours.

Lastly, manipulation of neurotransmitters within *fruP1* neurons revealed that the release of dopamine was not important for social space. Instead, both acetylcholine and GABA release from *fruP1* neurons play a role in fly social space, a phenocopy observed when knocking down acetylcholine in the MB. In addition, RNAi knockdown of *nlg3* in the *fruP1* neurons also showed sex-specific effects on social space.

Our antibody targeted the intracellular domain of Nlg3, which allowed us to identify protein localization specifically in the calyx (dendritic ends of the Kenyon cells) of the mushroom bodies, the medulla and lamina layers of the optic lobes, and throughout the entire protocerebral bridge (**Fig. 1**). However, there was a low signal to noise ratio when imaging with just a polyclonal anti-Nlg3 antibody (**Fig. 1A,B**) indicating a low level of expression of *nlg3* in the adult 3-4 day old brain. Attempts to create another polyclonal, or even a monoclonal antibody specific to Nlg3 were unsuccessful, so we transitioned to using two different genetic constructs

(*MiMIC-nlg3* and *nlg3-Gal4 > mCD8-GFP*) to confirm the results we observed with our
polyclonal antibody. The GFP localization in this line in the mushroom bodies and optic lobes
was similar to the Nlg3 protein localization (**Fig. 1C,D**). However, the GFP expression in the
MiMIC-nlg3 was also low, probably due to the endogenously low expression of *nlg3* (reported in
Robinson, Yost et al. 2020), and we could not see any expression in the protocerebral bridge. We
then switched to an *nlg3-Gal4* driver to express Gal4 transcription factor in the same cells as
nlg3, allowing to activated the *UAS-GFP* at higher levels than just endogenous *nlg3* (**Fig. 1F**)
allowing for better visualization with immunofluorescence of the calyx and protocerebral bridge
(**Fig. 1E**).

After determining the localization of *nlg3* expression in the adult brain, we examined the effects
of targeted RNAi knockdown of *nlg3* on fly social space. Deletions of the whole *nlg3* gene leads
to an increase in fly social space (Yost et al. 2020), thus, we expected to see a similar phenotype
with RNAi knockdown. However, the knockdown of *nlg3* only increased fly social space in the
optic lobes with the males contributing to most of the difference between experimental and
control treatments (**Fig. 2C**). Knocking down *nlg3* in the optic lobes might affect fly vision but
we did not yet directly test this. The RNAi knockdown of *nlg3* leads to a reduction in climbing
(**Supp. Fig. 2C**) as the climbing assay is performed under uniform light. The reduction in
climbing may be influenced by impairments in vision and not solely or entirely due to
impairments in locomotion. Vision is known to play a role in social spacing as well (Simon et al.
2012; Burg et al. 2013; Jiang et al. 2020), so the role of *nlg3* on vision will need to be further
investigated in the future.

Notably, we observed that the knockdown of *nlg3* in all three structures individually resulted in
sex differences in social space with females generally settling in a group closer together than
males (**Fig. 2A-C**). This is in line with previously published results with sex differences seen in
social space (Simon et al. 2012).

As we've seen that knocking down *nlg3* individually in the protocerebral bridge or mushroom
bodies isn't necessary for social space, we modified the transmission of the neurons within these
structures to assess their potential roles in this social behaviour. Indeed, Burg et al. (2012)
reported the importance of the MB in fly social space. We also observed an increase to male and
female social space whether the entire MB neurons were hyperactivated or silenced (**Fig. 3A**).
This suggests that it's not *nlg3* expression that influences social space but the MB transmission
itself. As the MB is not only important for learning and memory but also sensory integration, it
would follow that manipulation of this circuitry would affect behavioural output.

There were no changes to social space when we hyperactivated the PB neurons, but silencing the
PB neurons increased female social space specifically (**Fig. 3B**) suggesting the important of the
PB circuitry in regulating female social space. Next, using the *nlg3-Gal4* driver, we manipulated
the transmission of all *nlg3*-expressing neurons revealing their sexually dimorphic effect on
social space. Females will get further apart with hyperactivation or silencing of *nlg3*-containing
neurons, but males will only get further apart with hyperactivation and not silencing of the same
neurons. This might explain the sexually dimorphic effect of *nlg3* deletion on flies of the same
age (Yost et al. 2020) are not influenced by just one of the identified brain structures *nlg3* is

expressed in (MB, PB, OL), but instead may be due to the collective circuitry of all *nlg3*-
expression neurons (**Fig. 1**).
455
The mushroom bodies influence on fly social space was clear with changes to neural
transmission, so we then aimed to identify the neurotransmitter released from the mushroom
bodies that affected social spacing. As expected, knocking down *ChAT* and reducing
acetylcholine levels in the MB decreased social space, but only in females and did not affect
males (**Fig. 4A**). This indicates that acetylcholine release throughout the entire MB will lead to
behavioural changes in group formation for females but not males. We also found that knocking
down all dopamine receptor types in the mushroom bodies will specifically increase male social
space but did not significantly affect females (**Fig. 5A-D**). Of note, Dop1R1 and Dop2R
knockdown did increase female social space (**Fig. 5A,C**; Two-way ANOVA and Šídák's *post-*
hoc test: $p < 0.0505$; Two-way ANOVA and Šídák's *post-hoc* test: $p < 0.0276$, respectively) but
this was only a small difference in the number of flies within 4BL and was at the upper end of
how close flies can group together showing a ceiling effect that produced a minimal change in
their behaviour. Overall, this would indicate that the transmission of the Kenyon cells
themselves, specifically acetylcholine release, will influence female social space but the
dopaminergic inputs into the MB modify the male decision of their social space. It was
interesting to observe that all the 4 types of Drosophila dopamine receptors (Dop1R1, Dop1R2,
Dop2R, and DopEcR) all had the same effect on social space while each receptor plays unique
roles to fly behaviour such as sleep, courtship, feeding, and memory (Kim et al. 2003; Lebestky
et al. 2009; Kuo et al. 2015).
475
In Drosophila social space, many genetic and neural manipulations lead to sex differences,
despite the fact that both sexes respond to changes in their social environment in similar ways.
For example, age, mating, and isolation all lead to decreased social space in both sexes (Simon et
al. 2012; Brenman-Suttner et al. 2018; Yost et al. 2024). Due to this, we examined the influence
that the sexually dimorphic *fruP1* neuron subset had on social space. Again, we saw male-female
sex differences in response to modulation of *fruP1* neuronal transmission. For males, when
fruP1 neurons are hyperactivated, flies get further apart, but when these neurons are silenced,
males get closer together (**Fig. 6**). Females also get further apart when *fruP1* neurons are
hyperactivated, but do not change their social space when these neurons are silenced (**Fig. 6**).
This indicates that *fruP1* neurons in males can increase and decrease neuronal activity resulting
in a change in their social space, but for females only an increase in *fruP1* activity modifies their
social space. The basal transmission of these neurons that affect female social space might not be
active and therefore cannot be silenced, only hyperactivation changes this resting state and leads
to females settling further in a group setting.
490
Knowing the importance for *fruP1* neurons in Drosophila social space, we then examined the
molecular players within these neurons that are contributing to the behavioural changes.
Knocking down *nlg3* in *fruP1* neurons showed a male-female sexual dimorphism with males
getting further apart and females closer together (**Fig. 7A**). This confirms there are *fruitless*- and
nlg3-expressing neurons that are contributing to fly social space behaviour. Furthermore, our
results indicate that those neurons are not dopaminergic. However, the *fruP1* neurons affecting
social space are cholinergic, in both males and females, a phenocopy of the effect seen on
females both when reducing acetylcholine levels in the mushroom bodies and when knocking

down *nlg3* in *fruP1* neurons. Because a subset of the MB neurons are also *fruP1* neurons
(Palmateer et al. 2023), we propose that the Kenyon cells *nlg3* neurons are also *fruP1* neurons, at
least those that affect female social space (**Fig. 8**).
502
503 Finally, as previously reported (Jiang et al. 2020; Yost et al. 2024), we did not observe any
504 correlation in our experiments between abnormalities in performance index leading to any
505 changes in social space. Even when climbing was reduced, flies still were able to climb up to the
506 upper vial of counter current apparatus and are not necessarily consistent with changes to social
507 space.
508
Fly social behaviour relies heavily on an individual's ability to perceive and integrate social and
environmental cues. Using the Drosophila social space assay, we show the roles of the OL and
MB in regulating fly social spacing. The OL are essential for sensory perception, as they
transduce visual cues into electrochemical signals that are integrated within MB, and the central
complex, thereby enabling the fly to perceive its environment and alter its behaviour
accordingly.
515
The significance of Nlg3 in the OL is reiterated here, as it is probably involved in the proper
development and regulation of synaptic connections within these synaptically rich brain regions.
Proper development of the OL layers is required to decode parallel circuitry pathways involved
in processing color, shapes, and motion, which are crucial for interpreting environmental stimuli.
Outside of the development and maintenance of synaptic connections that can affect fly
behaviour, neural transmission through these brain regions reveals the MB to be important for
social behaviour. The MB, which receives signals and integrates from multiple other brain
regions (Aso et al. 2014), relies on proper development of synaptic connections where
disruptions or improper regulations in these connections can lead to misinterpretation of signals,
ultimately altering fly behavior through its descending output neurons.
526
By dissecting the molecular players within these neuronal subsets, we identify some of the inputs
and outputs of the MB that influence fly social spacing. This behavioral data allows us to begin
constructing a functional circuitry model of the neural pathways that regulate social space and
where sexual dimorphic integration of cues might be taking place. We have identified that
dopaminergic signalling into the MB influences social space but still need to identify which
dopaminergic clusters are playing a role. We can similarly test the known dopaminergic subsets
to innervate the MB (PPL1, PAM, and PPL2ab). We also identified that acetylcholine signalling
from the MB will affect female social space, but we have yet to identify which mushroom body
output neurons (MBONs) are associated with these behavioural changes (**Fig. 8**). We can also
test the known 21 MBONs to find downstream circuits that influence this social behaviour
specifically.
538
539
540

Acknowledgments:
We want to thank Dr. Amanda J. Moehring, in whose laboratory the plasmid carrying the *nlg3-RNAi* was generated and injected.
This study would not have been possible without the continued funding received by Western's Biotron Facility and the support offered there by the staff: we are very thankful to Karen Nygard for her technical and molecular help in immunofluorescence protocol establishment and microscopy within the integrated microscopy unit. We also want to thank the Insect Suite within the Biotron Facility of Western University, for housing our flies in climate-controlled walk-in rooms (25°C). We also acknowledge Dr. Ryley T. Yost for his western blot analysis to show the *nlg3-RNAi* results in a reduction of Nlg3 protein levels (data not shown).
We also want to thank Flybase (Öztürk-Çolak et al., 2024 - NHGRI Awards U41HG000739 and U24HG010859), the Bloomington Drosophila Stock Center (NIH P40OD018537) and for the nc82 (anti-Brp) primary antibody which was obtained from the Developmental Studies Hybridoma Bank, created by the NICHD of the NIH and maintained at The University of Iowa, Department of Biology, Iowa City, IA 52242, for the invaluable resources they provide to the scientific community.

Ethics Statement
Western's Animal Care Committee or Ontario Provincial and Federal regulatory bodies specifies that ethical considerations do not apply to the study of invertebrate animals including insects such as *Drosophila melanogaster*

Data availability statement:
The datasets analysed for this study can be found in the Dryad repository (TBD).

Conflict of Interest:

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Authors contribution:
Based on Oxford Academic (Genetics journal)

JWR: Conceptualization, Supervision, Project Administration, Investigation, Formal Analysis, Methodology, Resources, Funding Acquisition, Writing - Original Draft Preparation, and Writing - Review & Editing.
ATB: Conceptualization, Investigation, Formal Analysis, Funding Acquisition, and Writing - Review & Editing.
MRE: Conceptualization, Investigation, Formal Analysis, Funding Acquisition, and Writing - Review & Editing.
JRI: Methodology
SP: Investigation and Formal Analysis.
DSL: Investigation and Formal Analysis.
AS: Methodology.
JND: Investigation, Formal Analysis and Validation.
GIR: Writing - Review & Editing.
MBS: Supervision, Resources and Funding Acquisition - Review

AFS: Conceptualization, Supervision, Project Administration, Formal Analysis, Resources, Funding Acquisition, and Writing - Review & Editing.

References:

Aso Y, Hattori D, Yu Y, Johnston RM, Iyer NA, Ngo TTB, Dionne H, Abbott LF, Axel R, Tanimoto H, et al. 2014. The neuronal architecture of the mushroom body provides a logic for associative learning. Elife. 3:e04577. doi:10.7554/eLife.04577.

Baker CA, Guan X-J, Choi M, Murthy M. 2024. The role of fruitless in specifying courtship behaviors across divergent Drosophila species. Sci Adv. 10(11). doi:10.1126/sciadv.adk1273.

Brenman-Suttner D, Yost R, Frame A, Robinson J, Moehring A, Simon A. 2019. Social behaviour and aging: a fly model. Genes Brain Behav. 12598:1–18. doi:10.1111/gbb.12598. https://onlinelibrary.wiley.com/doi/abs/10.1111/gbb.12598.

Brenman-Suttner DB, Long SQ, Kamesan V, Belle JN De, Ryley T, Kanippayoor RL, Simon AF. 2018. Progeny of old parents have increased social space in Drosophila melanogaster. Sci Rep. 8(3673):1–13. doi:10.1038/s41598-018-21731-0. http://dx.doi.org/10.1038/s41598-018-21731-0.

Burg ED, Langan ST, Nash HA. 2013. Drosophila social clustering is disrupted by anesthetics and in narrow abdomen ion channel mutants. Genes Brain Behav. 12(3):338–347. doi:10.1111/gbb.12025.

Cao H, Tang J, Liu Q, Huang J, Xu R. 2022. Autism-like behaviors regulated by the serotonin receptor 5-HT2B in the dorsal fan-shaped body neurons of Drosophila melanogaster. Eur J Med Res. 27(1):203. doi:10.1186/s40001-022-00838-1.

Castermans D, Wilquet V, Parthoens E, Huysmans C, Steyaert J, Swinnen L, Fryns J-P, Van de Ven W, Devriendt K. 2003. The neurobeachin gene is disrupted by a translocation in a patient with idiopathic autism. J Med Genet. 40(5):352–6. doi:10.1136/jmg.40.5.352.

Chang K, Marran K, Valentine A, Hannon GJ. 2014. Generation of Transgenic Drosophila Expressing shRNAs in the miR-1 Backbone. Cold Spring Harb Protoc. 2014(5):pdb.prot080762. doi:10.1101/pdb.prot080762.

Chowdhury T, Calhoun RM, Bruch K, Moehring AJ. 2020. The fruitless gene affects female receptivity and species isolation. Proceedings of the Royal Society B: Biological Sciences. 287(1923):20192765. doi:10.1098/rspb.2019.2765.

Corthals K, Heukamp AS, Kossen R, Großhennig I, Hahn N, Gras H, Göpfert MC, Heinrich R, Geurten BRH. 2017. Neuroligins Nlg2 and Nlg4 affect social behavior in Drosophila melanogaster. Front Psychiatry. 8:1–13. doi:10.3389/fpsyt.2017.00113.

634 Fernandez RW, Akinleye AA, Nurilov M, Feliciano O, Lollar M, Aijuri RR, O'Donnell JM,
635 Simon AF. 2017. Modulation of social space by dopamine in Drosophila melanogaster, but no
636 effect on the avoidance of the Drosophila stress odorant. Biol Lett. 13(8):20170369.
637 doi:10.1098/rsbl.2017.0369.
638
639 Ferreira CH, Moita MA. 2019. What can a non-eusocial insect tell us about the neural basis of
640 group behaviour? Curr Opin Insect Sci. 36:1–7. doi:10.1016/j.cois.2019.09.001.
641 https://doi.org/10.1016/j.cois.2019.09.001.
642
643 Grant SG. 2012. Synaptopathies: diseases of the synaptome. Curr Opin Neurobiol. 22(3):522–
644 529. doi:10.1016/j.conb.2012.02.002.
645
646 Hahn N, Geurten B, Gurvich A, Piepenbrock D, Kästner A, Zanini D, Xing G, Xie W, Göpfert
647 MC, Ehrenreich H, et al. 2013. Monogenic heritable autism gene neuroligin impacts Drosophila
648 social behaviour. Behavioural Brain Research. 252:450–457. doi:10.1016/j.bbr.2013.06.020.
649
650 Hope KA, Flatten D, Cavitch P, May B, Sutcliffe JS, O'Donnell J, Reiter LT. 2019. The
651 Drosophila Gene Sulfateless Modulates Autism-Like Behaviors. Front Genet. 10.
652 doi:10.3389/fgene.2019.00574.
653
654 Ito H, Fujitani K, Usui K, Shimizu-Nishikawa K, Tanaka S, Yamamoto D. 1996. Sexual
655 orientation in Drosophila is altered by the satori mutation in the sex-determination gene fruitless
656 that encodes a zinc finger protein with a BTB domain. Proceedings of the National Academy of
657 Sciences. 93(18):9687–9692. doi:10.1073/pnas.93.18.9687.
658
659 Jamain S, Quach H, Betancur C, Råstam M, Colineaux C, Gillberg IC, Soderstrom H, Giros B,
660 Leboyer M, Gillberg C, et al. 2003. Mutations of the X-linked genes encoding neuroligins
661 NLGN3 and NLGN4 are associated with autism. Nat Genet. 34(1):27–29. doi:10.1038/ng1136.
662
663 Jiang L, Cheng Y, Gao S, Zhong Y, Ma C, Wang T, Zhu Y. 2020. Emergence of social cluster
664 by collective pairwise encounters in Drosophila. Elife. 9. doi:10.7554/eLife.51921.
665
666 Jiang Y, Yuen RKC, Jin X, Wang M, Chen N, Wu X, Ju J, Mei J, Shi Y, He M, et al. 2013.
667 Detection of Clinically Relevant Genetic Variants in Autism Spectrum Disorder by Whole-
668 Genome Sequencing. The American Journal of Human Genetics. 93(2):249–263.
669 doi:10.1016/j.ajhg.2013.06.012.
670
671 Kanellopoulos AK, Mariano V, Spinazzi M, Woo YJ, McLean C, Pech U, Li KW, Armstrong
672 JD, Giangrande A, Callaerts P, et al. 2020. Aralar Sequesters GABA into Hyperactive
673 Mitochondria, Causing Social Behavior Deficits. Cell. 180(6):1178-1197.e20.
674 doi:10.1016/j.cell.2020.02.044.
675
676 Kim Y-C, Lee H-G, Seong C-S, Han K-A. 2003. Expression of a D1 dopamine receptor
677 dDA1/DmDOP1 in the central nervous system of Drosophila melanogaster. Gene Expression
678 Patterns. 3(2):237–245. doi:10.1016/S1567-133X(02)00098-4.
679

Kuo SY, Wu CL, Hsieh MY, Lin CT, Wen RK, Chen LC, Chen YH, Yu YW, Wang HD, Su YJ,
et al. 2015. PPL2ab neurons restore sexual responses in aged Drosophila males through
dopamine. Nat Commun. 6:1–10. doi:10.1038/ncomms8490.

Lawal HO, Chang HY, Terrell AN, Brooks ES, Pulido D, Simon AF, Krantz DE. 2010. The
Drosophila vesicular monoamine transporter reduces pesticide-induced loss of dopaminergic
neurons. Neurobiol Dis. 40(1):102–112. doi:10.1016/j.nbd.2010.05.008.
http://dx.doi.org/10.1016/j.nbd.2010.05.008.

Lebestky T, Chang J-SC, Dankert H, Zelnik L, Kim Y-C, Han K-A, Wolf FW, Perona P,
Anderson DJ. 2009. Two Different Forms of Arousal in Drosophila Are Oppositely Regulated
by the Dopamine D1 Receptor Ortholog DopR via Distinct Neural Circuits. Neuron. 64(4):522–
536. doi:10.1016/j.neuron.2009.09.031.

McNeil AR, Jolley SN, Akinleye AA, Nurilov M, Rouzyi Z, Milunovich AJ, Chambers MC,
Simon AF. 2015. Conditions Affecting Social Space in Drosophila melanogaster. Journal of
Visualized Experiments.(105). doi:10.3791/53242.

Ni J-Q, Zhou R, Czech B, Liu L-P, Holderbaum L, Yang-Zhou D, Shim H-S, Tao R, Handler D,
Karpowicz P, et al. 2011. A genome-scale shRNA resource for transgenic RNAi in Drosophila.
Nat Methods. 8(5):405–407. doi:10.1038/nmeth.1592.

Palmateer CM, Artikis C, Brovero SG, Friedman B, Gresham A, Arbeitman MN. 2023. Single-
cell transcriptome profiles of Drosophila fruitless-expressing neurons from both sexes. Elife. 12.
doi:10.7554/eLife.78511.

Pan Y, Baker BS. 2014. Genetic Identification and Separation of Innate and Experience-
Dependent Courtship Behaviors in Drosophila. Cell. 156(1–2):236–248.
doi:10.1016/j.cell.2013.11.041.

Sato K, Yamamoto D. 2020. The mode of action of Fruitless: Is it an easy matter to switch the
sex? Genes Brain Behav. 19(2). doi:10.1111/gbb.12606.

Shih M-FM, Davis FP, Henry GL, Dubnau J. 2019. Nuclear Transcriptomes of the Seven
Neuronal Cell Types That Constitute the Drosophila Mushroom Bodies. G3
Genes|Genomes|Genetics. 9(1):81–94. doi:10.1534/g3.118.200726.

Shilpa O, Anupama KP, Antony A, Gurushankara HP. 2021. Lead (Pb)-induced oxidative stress
mediates sex-specific autistic-like behaviour in Drosophila melanogaster. Mol Neurobiol.
58(12):6378–6393. doi:10.1007/s12035-021-02546-z.

Simon AF, Chou M -T., Salazar ED, Nicholson T, Saini N, Metchev S, Krantz DE. 2012. A
simple assay to study social behavior in Drosophila: measurement of social space within a group.
Genes Brain Behav. 11(2):243–252. doi:10.1111/j.1601-183X.2011.00740.x.
724

Simon AF, Daniels R, Romero-Calderón R, Grygoruk A, Chang HY, Najibi R, Shamouelian D,
Salazar E, Solomon M, Ackerson LC, et al. 2009. Drosophila vesicular monoamine transporter
mutants can adapt to reduced or eliminated vesicular stores of dopamine and serotonin. Genetics.
181(2):525–541. doi:10.1534/genetics.108.094110.

Sokolowski MB. 2001. Drosophila: genetics meets behaviour. Nat Rev Genet. 2(11):879–890.
doi:10.1038/35098592.

Sokolowski MB. 2010. Social interactions in "simple" model systems. Neuron. 65(6):780–94.
doi:10.1016/j.neuron.2010.03.007. http://www.ncbi.nlm.nih.gov/pubmed/20346755.

Südhof TC. 2008. Neuroligins and neurexins link synaptic function to cognitive disease. Nature.
455(7215):903–911. doi:10.1038/nature07456.

Ueoka I, Kawashima H, Konishi A, Aoki M, Tanaka R, Yoshida H, Maeda T, Ozaki M,
Yamaguchi M. 2018. Novel Drosophila model for psychiatric disorders including autism
spectrum disorder by targeting of ATP-binding cassette protein A. Exp Neurol. 300:51–59.
doi:10.1016/j.expneurol.2017.10.027.

Venken KJT, Schulze KL, Haelterman NA, Pan H, He Y, Evans-Holm M, Carlson JW, Levis
RW, Spradling AC, Hoskins RA, et al. 2011. MiMIC: a highly versatile transposon insertion
resource for engineering Drosophila melanogaster genes. Nat Methods. 8(9):737–743.
doi:10.1038/nmeth.1662.

Vert J-P, Foveau N, Lajaunie C, Vandenbrouck Y. 2006. An accurate and interpretable model
for siRNA efficacy prediction. BMC Bioinformatics. 7(1):520. doi:10.1186/1471-2105-7-520.

Wiedemann C. 2007. Fruitless aggression. Nat Rev Neurosci. 8(1):5–5. doi:10.1038/nrn2064.

Wise A, Tenezaca L, Fernandez RW, Schatoff E, Flores J, Ueda A, Zhong X, Wu C-F, Simon
AF, Venkatesh T. 2015. Drosophila mutants of the autism candidate gene neurobeachin (rugose)
exhibit neuro-developmental disorders, aberrant synaptic properties, altered locomotion, and
impaired adult social behavior and activity patterns. J Neurogenet. 29(2–3):135–143.
doi:10.3109/01677063.2015.1064916.

Xie T, Ho MCW, Liu Q, Horiuchi W, Lin C-C, Task D, Luan H, White BH, Potter CJ, Wu MN.
2018. A Genetic Toolkit for Dissecting Dopamine Circuit Function in Drosophila. Cell Rep.
23(2):652–665. doi:10.1016/j.celrep.2018.03.068.

Xing G, Gan G, Chen D, Sun M, Yi J, Lv H, Han J, Xie W. 2014. Drosophila Neuroligin3
Regulates Neuromuscular Junction Development and Synaptic Differentiation. Journal of
Biological Chemistry. 289(46):31867–31877. doi:10.1074/jbc.M114.574897.

Yost RT, Robinson JW, Baxter CM, Scott AM, Brown LP, Aletta MS, Hakimjavadi R, Lone A,
Cumming RC, Dukas R, et al. 2020. Abnormal Social Interactions in a Drosophila Mutant of an
Autism Candidate Gene: Neuroligin 3. Int J Mol Sci. 21(13):4601. doi:10.3390/ijms21134601.

Yost RT, Scott AM, Kurbaj JM, Walshe-Roussel B, Dukas R, Simon AF. 2024. Recovery from

social isolation requires dopamine in males, but not the autism-related gene nlg3 in either sex. R

Soc Open Sci. 11(7). doi:10.1098/rsos.240604.

 Figure captions

 Figure 1. Nlg3 is expressed in the calyx of the mushroom bodies, throughout the
 protocerebral bridge, and in the optic lobes. A-B: Immunofluorescent imaging of polyclonal
 antibody specific to the c-terminal intracellular domain with anti-Nlg3 (green) on Canton-S male
 fly whole brains dissected at 3-4 days old. Magnification: 20X. **A.** Fluorescent imaging on
 posterior side of adult brain shows three main structures expressing *nlg3* in low levels: calyx of
 the MB (Ca), protocerebral bridge (PB), and the optic lobes (OL). **B.** Confocal imaging on
 posterior side of adult brain shows definition of Nlg3 expression in the Ca, PB, and OL (medulla
 and lamina). **C-D:** Expression patterns of MiMIC construct
 (*Mi{y[+mDint2]=MIC}nlg3[MI08924]*) incorporating *mCD8-GFP* into *nlg3*.
 Immunofluorescence with anti-GFP (green) and nc82 (magenta). **C.** Confocal imaging on
 anterior side of adult brain at 20X magnification indicates low expression of Nlg3 in the MB and
 OL. **D.** Confocal imaging at 40X magnification highlighting expression in the MB. **E-F:**
 Expression patters of *MiMIC nlg3-Gal4* genetic construct (*Mi{Trojan-Gal4}nlg3>UAS-*
 mCD8::GFP) driving exogenous expression of *mCD8-GF*P with *nlg3*. Immunofluorescence
 with anti-GFP (green) and nc82 (magenta). **E.** Confocal imaging of posterior side of adult brain
 at 40X magnification indicates low expression patterns in PB and Ca. **F.** Confocal imaging of
 anterior side of adult brain at 40X magnification shows expression throughout the MB. Antennal
 Lobes show for reference (AL). Scale bar, 50μm.

 Figure 2. Knocking down Nlg3 specifically in the optic lobes increases fly social space in
 males and females but changes are seen when knocking down in the protocerebral bridge
 or mushroom bodies specifically. A-C: Average number of flies within 4 body lengths (4BL)
 in male and female flies with RNAi knockdown of *nlg3* targeted specifically to the mushroom
 bodies (A), protocerebral bridge (B), and the optic lobes (C). **A.** The number of flies within 4BL
 was not different in both males and females in flies with *nlg3* knockdown targeted to the MB
 (two-way ANOVA – Effect of genotype*: $F_{1,39}$* = 1.005, *p* = 0.3223). **B.** The average number of
 flies within 4BL was not different in both males and females in flies with *nlg3* knockdown
 targeted to the PB (two-way ANOVA – Effect of genotype: $F_{1,37}$ = 0.068, *p* = 0.7951), but there
 is an increase in the number of flies within 4BL in female *nlg3* knockdown flies when compared
 to males (two-way ANOVA – Effect of sex: $F_{1,37}$ = 10.12, *p* = 0.0030). **C.** Knocking down *nlg3*
 in the OL decreases the number of flies within 4 body lengths in both males and females
 compared to genetic control flies (two-way ANOVA – Effect of genotype: $F_{1,37}$ = 4.837, *p* =
 0.0354). Grey colour: genetic control *driver/+*; pink colour: *driver/RNAi*. The results of all two-
 way ANOVA and *post-hoc* tests performed can be found in table 1. N = 7 – 14 for all treatments.
 Error bars are +/- s.e.m.

 Figure 3. Manipulation of neural transmission within the brain structures enriched with
 Nlg3 increases fly social space. A-C: Average number of flies within 4 body lengths (4BL) in
 male and female flies when genetically hyperactivating or silencing neural transmission within
 the MB (A), PB (B), and *nlg3*-containing neurons (C). **A.** Hyperactivation of MB neurons
 decreases the number of flies within 4BL in both males and females compared to basal
 transmission of MB neurons (two-way ANOVA – Effect of neural transmission: $F_{1,47}$ = 7.395, *p*
 = 0.0091). Silencing MB neurons also decreases the number of flies within 4BL in both males
 and females compared to basal transmission (two-way ANOVA – Effect of neural transmission:

$F_{1,30}$ = 14.40, p = 0.0007). **B.** The number of flies within 4BL was not different in males and
females when the PB neurons are hyperactivated (two-way ANOVA - Effect of neural
transmission: $F_{1,74}$ = 17.72, p = 0.2018) or silenced (two-way ANOVA – Effect of neural
transmission: $F_{1,31}$ = 3.443, p = 0.0730). **C.** Hyperactivation of *nlg3*-containing neurons
decreases the number of flies within 4BL in both males and females compared to basal
transmission (two-way ANOVA – Effect of neural transmission: $F_{1,28}$ = 12.70, p = 0.0013).
There is an interaction between sex and silencing *nlg3*-containing neurons where females have a
decrease in the number of flies within 4BL (two-way ANOVA – Effect of interaction: $F_{1,22}$ =
6.169, p = 0.0211). Grey colour bars: inactivated genetic construct and basal transmission of
neurons; green colour bars: activation of genetic construct and hyperactivation of neurons with
Gal4 driver; pink colour bars: activation of genetic construct and silencing of neurons with *Gal4*
driver. The results of all two-way ANOVA and *post-hoc* tests performed can be found in table 1.
N = 6 – 22 for all treatments. Error bars are +/- s.e.m.
835
Figure 4. Knocking down the acetylcholine rate-limiting biosynthesis enzyme C*hAT* in the
MB decreases female fly social space. A-B: Average number of flies within 4 body lengths
(4BL) in male and female flies with RNAi knockdown of *ChAT* for acetylcholine synthesis (A)
or *GAD1* for GABA synthesis (B) specifically in the mushroom bodies. **A.** The interaction
between sex and *ChAT* knockdown shows that female flies have an increased number of flies
within 4BL compared to female genetic control flies (two-way ANOVA – Effect of interaction:
$F_{1,30}$ = 8.465, p = 0.0068). **B.** Knockdown of *GAD1* in the MB does not change the number of
flies in 4BL in both males and females compared to genetic control flies (two-way ANOVA –
Effect of genotype: $F_{1,31}$ = 0.021, p = 0.8851). Grey colour bars: genetic control *driver/+*; pink
colour bars: *driver/RNAi*. The results of all two-way ANOVA and *post-hoc* tests performed can
be found in table 1. N = 8 – 9 for all treatments. Error bars are +/- s.e.m.
847
Figure 5. RNAi knockdown of all dopamine receptors in the MB increases male social
space. A-D: Average number of flies within 4 body lengths (4BL) in male and female flies
knocking down dopamine receptors *Dop1R1* (A), *Dop1R2* (B), *Dop2R* (C), and *DopEcR* (D)
specifically in the mushroom bodies. **A.** *Dop1R1* knockdown in the MB of males has a decreased
number of flies within 4BL compared to genetic controls (two-way ANOVA with Šídák's *post-*
hoc test: p < 0.0001). **B.** *Dop1R2* knockdown in the MB of males has a decreased number of
flies within 4BL compared to genetic controls (two-way ANOVA and Šídák's *post-hoc* test: p =
0.0225). **C.** *Dop2R* knockdown in the MB of males has a decreased number of flies within 4BL
compared to genetic controls (two-way ANOVA and Šídák's *post-hoc* test: p < 0.0001). **D.**
DopEcR knockdown in the MB of males has a decreased number of flies within 4BL compared
to genetic controls (two-way ANOVA and Šídák's *post-hoc* test: p = 0.0103). Grey colour bars:
genetic control *driver/+*; pink colour bars: *driver/RNAi*. The results of all two-way ANOVA and
post-hoc tests performed can be found in table 1. N = 9 for all treatments. Error bars are +/-
s.e.m.
862
863   **Figure 6. Hyperactivation of *FruP1* neurons increases social space in males and females**
864   **while silencing *FruP1* neurons only decreases social space in males.** Average number of flies
865   within 4 body lengths (4BL) in males and females when genetically hyperactivating neural
866   transmission in *FruP1* neurons. Hyperactivation of *FruP1* neurons decreases the number of flies
867   within 4BL in both males and females compared their basal transmission (two-way ANOVA –

Effect of neural transmission: $F_{1,33} = 22.65$, $p < 0.0001$). Silencing *FruP1* neurons did not alter change female social space but increased the number of flies within 4BL in males (two-way ANOVA with Šídák's *post-hoc* test: $p = 0.0453$). Grey colour bars: inactivated genetic construct and basal transmission of neurons; green colour bars: activation of genetic construct and hyperactivation of neurons with Gal4 driver; pink colour bars: activation of genetic construct and silencing of neurons with Gal4 driver. The results of all two-way ANOVA and *post-hoc* tests performed can be found in table 1. N = 7 – 11 for all treatments. Error bars are +/- s.e.m.

Figure 7. Knocking down *nlg3* in *FruP1* neurons decreases female social space, while a reduction of acetylcholine in *FruP1* neurons decreases social space in male and female flies. **A-D:** Average number of flies within 4 body lengths (4BL) in male and female flies with RNAi knockdown of *nlg3* (A), *TH* for dopamine synthesis (B), *ChAT* for acetylcholine synthesis (C) or *GAD1* for GABA synthesis (D) specifically in *FruP1* neurons. **A.** Knocking down *nlg3* in *FruP1* has an interaction between sex for number of flies within 4BL (two-way ANOVA – Effect of interaction: $F_{1,58} = 7.056$, $p = 0.0102$) **B.** *TH* knockdown in *FruP1* neurons does not change social space in males or females (two-way ANOVA – Effect of genotype: $F_{1,31} = 1.134$, $p = 0.2951$). **C.** *ChAT* knockdown in *FruP1* neurons increases the number of flies within 4BL in both males and females compared to the genetic control flies (two-way ANOVA – Effect of genotype: $F_{1,48} = 38.01$, $p < 0.0001$). **D.** *GAD1* knockdown in *FruP1* neurons increases the number of flies within 4BL in only females (two-way ANOVA – Effect of interaction: $F_{1,39} = 9.327$, $p = 0.0041$). Grey colour bars: genetic control *driver/+*; pink colour bars: *driver/RNAi*. The results of all two-way ANOVA and *post-hoc* tests performed can be found in table 1. N = 9 - 13 for all treatments. Error bars are +/- s.e.m.

Figure 8. Working model of functional neural influences on Drosophila social space: Dopaminergic inputs into the mushroom body neurons (dark blue line) influence male social space while cholinergic outputs from the mushroom bodies influence female social space. Cholinergic outputs from *fruP1* neurons (light green line) also influence male and female social space while some mushroom body neurons also express *fru*. Neuroligin protein (red dots) enrichment is also found to be located in the optic lobes, calyx of the mushroom bodies, and protocerebral bridge.

901 **Figure 1**
902

903
904
905
906

Figure 2

Males   **Females**

Effect of sex (p = 0.0980)

A

MB > *nlg3*-RNAi

Effect of sex (p = 0.0030)

B

PB > *nlg3*-RNAi

Effect of sex (p = 0.0004)
Effect of genotype (p = 0.0354)

C

OL > *nlg3*-RNAi

Gal4 Control   RNAi Knockdown

Figure 3

 Figure 5

Males   **Females**

Effect of sex (p < 0.0001)
Effect of genotype (p < 0.0001)
Effect of interaction (p < 0.0001)

Effect of sex (p < 0.0001)
Effect of genotype (p = 0.005)

MB > *Dop2R*-RNAi

MB > *DopEcR*-RNAi

Gal4 Control   RNAi Knockdown

919
920

Number of Flies Within 4BL

Figu

Hyper

14   9

14   9   15   13

8   8   9   9

Effect of sex (p = 0.0296)
Effect of neural transmission (p < 0.0001)

Effect of interaction (p = 0.0730)

0.0453

19   20   17   22

8   8   10   9

10   7   11   9

FruP1 > TrpA1

9   8

6   7   7   6

■ **Basal Transmission**   ■ **Hyperactivation**   **Silencing**

      Figure 8

Supplemental Figure Captions

Supplemental Table 1. ANOVA table for all Two-Way ANOVAs performed. Statistically significant results are bolded.

Supplemental Figure 1: Schematic of crossing scheme used to generate genotype controls for all RNAi knockdown experiments. Bolded genotypes are presenting in main figures. Results for all genotypes are in the supplemental data below. Main conclusions are unchanged when adding or removing the *UAS/+* genetic control.

Supplemental Figure 2: Knocking down Nlg3 specifically in the Protocerebral bridge and optic lobes decreases fly climbing in males and females but not when knocking down in the mushroom bodies specifically. A-C: Startle-induced climbing represented by the mean percentage of flies reaching the top vial (performance index). Genetic constructs include RNAi knockdown of *nlg3* targeted specifically to the mushroom bodies (A), protocerebral bridge (B), and the optic lobes (C). Grey colour: genetic control *driver/+*; pink colour: *driver/RNAi*. The results of all two-way ANOVA and *post-hoc* tests performed can be found in table 1. N = 5 – 10 for all treatments. Errors bar are +/- s.e.m.

Supplemental Figure 3: Hyperactivation of all the brain structures enriched with Nlg3 decreases fly climbing. A-C: Startle-induced climbing represented by the mean percentage of flies reaching the top vial (performance index). Hyperactivation or silencing neurons specifically to the mushroom bodies (A), protocerebral bridge (B), and all *nlg3-expressing* neurons (C). Grey colour bars: inactivated genetic construct and basal transmission of neurons; green colour bars: activation of genetic construct and hyperactivation of neurons with *Gal4* driver; pink colour bars: activation of genetic construct and silencing of neurons with *Gal4* driver. The results of all two-way ANOVA and *post-hoc* tests performed can be found in table 1. N = 4 – 7 for all treatments. Error bars are +/- s.e.m.

Supplemental Figure 4: Hyperactivation and silencing of *FruP1* neurons decreases climbing in males and females. Startle-induced climbing represented by the mean percentage of flies reaching the top vial (performance index). Hyperactivation or silencing neurons specifically in *FruP1* neuron subset. Grey colour bars: inactivated genetic construct and basal transmission of neurons; green colour bars: activation of genetic construct and hyperactivation of neurons with *Gal4* driver; pink colour bars: activation of genetic construct and silencing of neurons with *Gal4* driver. The results of all two-way ANOVA and *post-hoc* tests performed can be found in table 1. N = 9 for all treatments. Error bars are +/- s.e.m.

Supplemental Figure 5: Knocking down *nlg3* in *FruP1* neurons decreases male climbing, while a reduction of dopamine, acetylcholine, and GABA in *FruP1* neurons all decrease climbing in male and female flies. A-D: Startle-induced climbing represented by the mean percentage of flies reaching the top vial (performance index). Genetic constructs include RNAi knockdown targeted specifically to the *FruP1* subset of neurons of *nlg3* (A), knocking down dopamine (B), acetylcholine (C), and GABA (D). Grey colour: genetic control *driver/+*; pink colour: *driver/RNAi*. The results of all two-way ANOVA and *post-hoc* tests performed can be found in table 1. N = 6 – 15 for all treatments. Errors bar are +/- s.e.m.

976   **Supplemental Figure 6: Elevated temperature on Cs, *Trp1A/+*, and *Shi^{ts}/+* does not affect**
977   **social space or climbing.**
978
979   **Supplemental Figure 7: Knocking down Nlg3 specifically in the optic lobes increases fly**
980   **social space in males and females but changes are seen when knocking down in the**
981   **protocerebral bridge or mushroom bodies specifically (both genetic controls shown).**
982
983   **Supplemental Figure 8. Knocking down the acetylcholine rate-limiting biosynthesis enzyme**
984   **C*hAT* in the MB decreases female fly social space (both genetic controls shown).**
985
986   **Supplemental Figure 9: RNAi knockdown of all dopamine receptors in the MB increases**
987   **male social space (both genetic controls shown).**
988
989   **Supplemental Figure 10: Knocking down *nlg3* in *FruP1* neurons decreases female social**
990   **space, while a reduction of acetylcholine in *FruP1* neurons decreases social space in male**
991   **and female flies (both genetic controls shown).**
992

Supp. Table 1

Experiment	Figure	Effect	Df	F	p	Šídák's post hoc		
Social space						Comparison Group	Compared with	p
MB>nlg3-RNAi	2A	Sex	(1,39)	2.873	0.0980	*Male*		
		Genotype	(1,39)	1.005	0.3223	MB>RNAi	MB/+	0.9602
		Sex * Genotype	(1,39)	1.849	0.1817	*Female*		
						MB>RNAi	MB/+	0.1992
PB>nlg3-RNAi	2B	**Sex**	(1,37)	10.12	**0.0030**	*Male*		
		Genotype	(1,37)	0.068	0.7951	PB>RNAi	PB/+	0.4937
		Sex * Genotype	(1,37)	1.328	0.2566	*Female*		
						PB>RNAi	PB/+	0.8051
OL>nlg3-RNAi	2C	**Sex**	(1,31)	15.93	**0.0004**	*Male*		
		Genotype	(1,31)	4.837	**0.0354**	OL>RNAi	OL/+	0.0845
		Sex * Genotype	(1,31)	0.542	0.4670	*Female*		
						OL>RNAi	OL/+	0.5322
MB>TrpA1	3A	**Sex**	(1,47)	5.382	**0.0247**	*Male MB>Trp*		
		Neural Transmission	(1,47)	7.395	**0.0091**	Activated	Inactivated	0.2437
		Sex * Neural Transmission	(1,47)	0.172	0.6796	*Female MB>Trp*		
						Activated	**Inactivated**	**0.0442**
MB>Shi^{ts}	3B	Sex	(1,30)	0.680	0.4161	*Male MB>Shi*		
		Neural Transmission	(1,30)	14.40	**0.0007**	**Activated**	**Inactivated**	**0.0105**
		Sex * Neural Transmission	(1,30)	0.344	0.5617	*Female MB>Shi*		
						Activated	Inactivated	0.0518
PB>TrpA1	3C	Sex	(1,74)	39.94	0.0868	*Male PB>Trp*		
		Neural Transmission	(1,74)	17.72	0.2018	Activated	Inactivated	0.9999
		Sex * Neural Transmission	(1,74)	13.62	0.1953	*Female PB>Trp*		
						Activated	Inactivated	0.1382
PB>Shi^{ts}	3D	Sex	(1,31)	0.6335	0.4321	*Male PB>Shi*		
		Neural Transmission	(1,31)	3.443	0.0730	Activated	Inactivated	0.8897
		Sex * Neural Transmission	(1,31)	1.483	0.2325	*Female PB>Shi*		
						Activated	Inactivated	0.0596
Trojan>TrpA1	3E	Sex	(1,28)	2.380	0.1341	*Male Trojan>Trp*		
		Neural Transmission	(1,28)	12.70	**0.0013**	Activated	Inactivated	0.2214
		Sex * Neural Transmission	(1,28)	1.366	0.2523	*Female Trojan>Trp*		
						Activated	**Inactivated**	**0.0029**
Trojan>Shi^{ts}	3F	**Sex**	(1,22)	4.666	**0.0419**	*Male MB>Trp*		
		Neural Transmission	(1,22)	5.998	**0.0228**	Activated	Inactivated	0.9996
		Sex * Neural Transmission	(1,22)	6.169	**0.0211**	*Female MB>Trp*		
						Activated	**Inactivated**	**0.0042**
MB>CHAT25-RNAi	4A	**Sex**	(1,30)	22.26	**<0.0001**	*Male*		
		Genotype	(1,30)	6.805	**0.0140**	MB>RNAi	MB/+	0.9721
		Sex * Genotype	(1,30)	8.465	**0.0068**	*Female*		
						MB>RNAi	**MB/+**	**0.0010**
MB>GAD51-RNAi	4B	**Sex**	(1,31)	42.89	**<0.0001**	*Male*		
		Genotype	(1,31)	0.021	0.8851	MB>RNAi	MB/+	0.4656
		Sex * Genotype	(1,31)	2.012	0.1651	*Female*		
						MB>RNAi	MB/+	0.6167
MB>Dop1R1-RNAi	5A	**Sex**	(1,32)	65.51	**<0.0001**	*Male*		
		Genotype	(1,32)	19.61	**<0.0001**	**MB>RNAi**	**MB/+**	**<0.0001**
		Sex * Genotype	(1,32)	27.24	**<0.0001**	*Female*		
						MB>RNAi	MB/+	0.0505
MB>Dop1R2-RNAi	5B	**Sex**	(1,32)	139.4	**<0.0001**	*Male*		
		Genotype	(1,32)	7.401	**0.0105**	**MB>RNAi**	**MB/+**	**0.0225**
		Sex * Genotype	(1,32)	1.168	0.2880	*Female*		
						MB>RNAi	MB/+	0.4447
MB>Dop2R-RNAi	5C	**Sex**	(1,32)	120.7	**<0.0001**	*Male*		
		Genotype	(1,32)	41.10	**<0.0001**	**MB>RNAi**	**MB/+**	**<0.0001**
		Sex * Genotype	(1,32)	7.448	**0.0102**	*Female*		
						MB>RNAi	**MB/+**	**0.0276**
MB>DopEcR-RNAi	5D	**Sex**	(1,32)	139.3	**<0.0001**	*Male*		
		Genotype	(1,32)	8.905	**0.0054**	**MB>RNAi**	**MB/+**	**0.0103**
		Sex * Genotype	(1,32)	1.586	0.2167	*Female*		
						MB>RNAi	MB/+	0.4098
FruP1>TrpA1	6A	**Sex**	(1,33)	5.168	**0.0296**	*Male FruP1>Trp*		
		Neural Transmission	(1,33)	22.65	**<0.0001**	**Activated**	**Inactivated**	**0.0241**
		Sex * Neural Transmission	(1,33)	0.6987	0.4092	*Female FruP1>Trp*		
						Activated	**Inactivated**	**0.0004**
FruP1>Shi^{ts}	6B	Sex	(1,30)	0.5671	0.4573	*Male FruP1>Trp*		
		Neural Transmission	(1,30)	2.350	0.1358	**Activated**	**Inactivated**	**0.0453**
		Sex * Neural Transmission	(1,30)	3.451	0.0730	*Female FruP1>Trp*		
						Activated	Inactivated	0.9676
FruP1>nlg3-RNAi	7A	**Sex**	(1,58)	6.969	**0.0107**	*Male*		
		Genotype	(1,58)	0.087	0.7690	FruP1>RNAi	FruP1/+	0.2303
		Sex * Genotype	(1,58)	7.056	**0.0102**	*Female*		
						FruP1>RNAi	FruP1/+	0.0563
FruP1>TH-RNAi	7B	**Sex**	(1,31)	69.88	**<0.0001**	*Male*		
		Genotype	(1,31)	1.134	0.2951	FruP1>RNAi	FruP1/+	0.9788
		Sex * Genotype	(1,31)	1.750	0.1956	*Female*		
						FruP1>RNAi	FruP1/+	0.2013
FruP1>CHAT25-RNAi	7C	Sex	(1,48)	0.777	0.3824	*Male*		
		Genotype	(1,48)	38.01	**<0.0001**	**FruP1>RNAi**	**FruP1/+**	**0.0150**
		Sex * Genotype	(1,48)	4.919	**0.0313**	*Female*		
						FruP1>RNAi	**FruP1/+**	**<0.0001**
FruP1>GAD51-RNAi	7D	Sex	(1,39)	0.008	0.9279	*Male*		
		Genotype	(1,39)	11.47	**0.0016**	FruP1>RNAi	FruP1/+	0.9649
		Sex * Genotype	(1,39)	9.327	**0.0041**	*Female*		
						FruP1>RNAi	**FruP1/+**	**0.0001**

Table 1. ANOVA table for all Two-way ANOVAs performed. Statistically significant results are bolded

995   **Supp. Figure 1**
996

997
998

Climbing

Climbing

Hyperactivation

A

Number of Flies Within 4BL

MB > TrpA1

Effect of sex (p = 0.0003)

MB > Shi^ts

B

Number of Flies Within 4BL / Climbing (Performance Index)

PB > TrpA1

PB > Shi^ts

C

Climbing (Performance Index)

Effect of transmission (p = 0.0410)

Trojan > TrpA1

Trojan > Shi^ts

■ **Basal Transmission**   ■ **Hyperactivation**   ■ **Silencing**

Supp. Figure 5

Climbing

Males   Females

Effect of sex (p = 0.0431)
Effect of genotype (p = 0.0008)
Effect of interaction (p = 0.0370)

FruP1 > *nlg3*-RNAi
Effect of genotype (p < 0.0001)

FruP1 > *TH*-RNAi
Effect of genotype (p < 0.0001)

FruP1 > *CHAT*-RNAi

Effect of genotype (p = 0.0026)

FruP1 > *GAD*-RNAi

Gal4 Control   RNAi Knockdown

1018
1019

Males   Females

Effect of sex (p = 0.0024)
Effect of genotype (p = 0.0001)
Effect of interaction (p = 0.0204)

A

FruP1 > *nlg3*-RNAi

Effect of sex (p < 0.0001)
Effect of genotype (p = 0.2472)
Effect of interaction (p = 0.1246)

B

FruP1 > *TH*-RNAi

Effect of sex (p = 0.1510)
Effect of genotype (p = 0.0030)
Effect of interaction (p = 0.0034)

C

FruP1 > *CHAT*-RNAi

Effect of sex (p < 0.0001)
Effect of genotype (p = 0.0002)
Effect of interaction (p = 0.0041)

D

FruP1 > *GAD*-RNAi

■ Gal4 Control   ■ RNAi Knockdown   ■ UAS Control