Reduction of *Grin2a* in adolescent rat dopamine neurons confers a phenotype
relevant to psychosis

Michelle L. Kielhold[1], David S. Jacobs[1], Alejandro Torrado-Pacheco[1], Merridee J. Lefner [1],
Joseph J. Lebowitz[2], Angela J. Langdon[3], John T. Williams[2], Larry S. Zweifel[4], and Bita
Moghaddam[1]

[1]Department of Behavioral Neuroscience, Oregon Health & Science University, Portland,
Oregon

[2]Vollum Institute, Oregon Health & Science University, Portland, Oregon

[3]Intramural Research Program, National Institute of Mental Health, National Institutes of
Health, Bethesda, MD

[4]Department of Pharmacology, University of Washington, Seattle, Washington

Corresponding author: Bita Moghaddam bita@ohsu.edu

SUMMARY

Psychosis is a hallmark of schizophrenia. It involves a collection of symptoms that are typically associated with disrupted dopamine signaling and emerges during adolescence or early adulthood. Most schizophrenia-associated genes, however, involve glutamatergic or other ubiquitous targets that do not explain the latent expression of psychosis or dopaminergic abnormalities. Here, we describe an etiologically relevant model for the adolescent onset of dopamine-related dysfunction in schizophrenia. We focused on *GRIN2A,* the gene encoding the GluN2A subunit of the NMDA receptor, as both the common loss-of-function variants and the rare missense variants in this gene are risk factors for schizophrenia. We find that GluN2A levels distinctly decline in dopamine neuron-containing regions throughout adolescence while remaining stable in other regions. This suggested that adolescent dopamine neurons may be particularly vulnerable to further reductions in GluN2A caused by a damaging variant of *GRIN2A*. Consistent with this idea, we find that selective knockout of *Grin2a* in adolescent rat dopamine neurons results in a psychosis-relevant behavioral phenotype. This manipulation also reduced dopamine release in response to unexpected outcomes in young adults, reflective of prediction error signaling abnormalities observed in the clinical population. These data provide mechanistic insight into how *GRIN2A* mutations may contribute to the delayed onset of dopamine-related symptoms and provide a model for identifying course altering treatments for schizophrenia.

INTRODUCTION

NMDA receptor-mediated glutamatergic neurotransmission has long been implicated in the pathophysiology of schizophrenia (SCZ)[1,2]. This association initially stemmed from postmortem findings[3-5] and the discovery that the drug phencyclidine, which transiently mimics SCZ symptoms, is an antagonist of the NMDA receptor[6,7]. Recent genetic findings have strengthened evidence implicating the NMDA receptor in the etiology of SCZ and, more specifically, have directed the focus toward the GluN2A subunit of this receptor. Both common and rare variants that reduce the function of the gene encoding this subunit (*GRIN2A*) have been identified as genetic risk factors for SCZ[8-13]. In line with this discovery, *Grin2a* mutant mice display a wide range of aberrant cortical activity and locomotor abnormalities[14-16].

Dopamine neurotransmission is equally implicated in SCZ pathophysiology and psychosis[17-19]. Notably, all known antipsychotic drugs act on dopamine receptors[20-23]. While genetic vulnerability to develop SCZ generally does not suggest a primary role for dopamine receptors or related proteins, extensive imaging and pharmacological studies have implicated the dopamine system in the positive symptoms of the disorder. PET studies, for

example, have reported disrupted dopamine neurotransmission in individuals with SCZ or
young, at-risk individuals who go on to develop the disorder[24-26].

The involvement of *GRIN2A* in the etiology of SCZ may explain its distinct temporal pattern
of symptom expression. SCZ is characterized by three types of symptoms: cognitive,
affective (negative), and psychotic (positive) symptoms. Cognitive and affective symptoms
often emerge in late childhood, whereas psychotic symptoms typically manifest in late
adolescence or early adulthood and are associated with disruptions in dopamine
neurotransmission[17]. In humans and rodents *GRIN2A/Grin2a* cortical expression is absent
or undetected early in development but steeply increases with age, beginning perinatally
(humans) or after birth (rodents), reaching adult levels in early childhood[27-31]. Thus,
phenotypes related to a reduction in functional *GRIN2A* would emerge during this early
developmental stage, which coincides with the timeline of cognitive and affective symptom
expression in SCZ. But why would psychosis remain latent until late adolescence or early
adulthood? Moreover, why are these late-emerging positive symptoms—and not the
negative or cognitive symptoms—alleviated by dopamine receptor antagonists[21-25,32-38].
These questions led us to investigate how reduced function of *GRIN2A* in adolescent
dopamine neurons may lead to presentation of behavioral phenotypes that are relevant to
psychosis.

We found that, unlike cortical regions where GluN2A remains stable during adolescence,
levels in dopamine neuron-containing regions decline at this age. This suggested that a
loss of function in *Grin2a* may exaggerate this natural decline and be most detrimental to
dopamine neurotransmission during adolescence. Consistent with this mechanism,
targeted knockout of *Grin2a* in adolescent dopamine neurons was sufficient to produce a
selective phenotype relevant to aspects of psychosis, without affecting spontaneous or
simple reward seeking behaviors.

RESULTS

GluN2A expression is dynamic across postnatal development in a region-specific manner

Brain regions including the prefrontal cortex and hippocampus do not express GluN2A in
rodents at birth, but levels rapidly increase postnatally, reaching adult levels around
weaning[30,31,39]. It is, however, not known if GluN2A levels remain stable between weaning
and adulthood. We quantified GluN2A protein levels in male and female rats immediately
after weaning on postnatal day 21 (P21), during adolescence (P35, P45), and adulthood
(P80) (**Figure 1a**), focusing on the medial prefrontal cortex (mPFC), hippocampus, and
regions containing dopamine neurons (i.e. ventral tegmental area (VTA) and substantia
nigra (SN)). We found that GluN2A levels in the mPFC and hippocampus remain stable

between ages P21 and P80 (**Figure 1b**). In contrast, GluN2A levels were labile in dopamine
neuron-rich regions during adolescence. In both the VTA and SN, GluN2A levels declined
after P21, reaching adult levels during mid-to-late adolescence (**Figure 1c**).

This observation led us to hypothesize that the damaging variants of *GRIN2A* implicated in
SCZ may exacerbate this natural reduction of GluN2A in dopamine neurons during
adolescence, providing a mechanism for the emergence of psychosis at this
developmental stage (**Figure 1d**). We tested this idea by establishing a model of
adolescent dopamine neuron-selective knockout of *Grin2a* (DA-*Grin2a* KO) through the
infusion of a Cre-inducible CRISPR/Cas9 virus targeting *Grin2a* into the VTA of TH::Cre rats.
This virus was co-infused with a Cre-dependent virus that would express green fluorescent
protein (GFP) for visualization (**Figure 1e**). Viral infusions were performed on the day of
weaning (P21) and expression was assessed following a 2-week incubation period.
Immunohistochemistry confirmed cell-specific virus expression, as staining for the
hemagglutinin (HA)-tag on the DA-*Grin2a* KO virus significantly overlapped with GFP
(**Figure 1f**) and this expression was specific to dopamine neurons as there was significant
co-expression of GFP and tyrosine hydroxylase (TH) (**Figure 1g**). Functional reduction of
GluN2A in GFP+ neurons was established with recordings in slice preparation (**Figure 1h**).
Bath application of a GluN2A-specific antagonist (TCN201) resulted in a reduction of the
glutamate-induced current in controls whereas DA-*Grin2a* KO cells showed a significantly
blunted response (**Figure 1i**). This reduction was specific to GluN2A-containing NMDA
receptors as a GluN2B-targeting antagonist (ifenprodil) produced similar effects in cells of
both control and DA-*Grin2a* KO animals (**Figure 1j**). The extent of this regional loss of
GluN2A was determined through Western blot analysis, where we observed an
approximate 50% reduction of GluN2A in the VTA of DA-*Grin2a* KO animals compared to
controls (**Figure 1k**).

Figure 1: GluN2A loss in the adolescent midbrain was augmented with virally mediated *Grin2a* selective knockout in dopamine neurons.

a, Timeline of tissue collection ages for postnatal day 21 (P21), adolescent (P35, P45), and adult (P80) samples. **b,** Medial prefrontal cortex (mPFC) and hippocampus showed no effect of age on normalized GluN2A levels. **c,** Ventral tegmental area (VTA) showed significant effect of age, with reductions in normalized GluN2A at P21 vs P45 (*p=0.0157) and vs P80 (**p=0.0058). The substantia nigra (SN) also showed an effect of age with reductions from P21 vs P35 (**p=0.0036), vs P45 (*p=0.0318), and vs P80 (**p=0.0033); (n=5-6 brains). Significance testing: one-way ANOVA; Dunnett post hoc. **d,** Representation of correlation between adolescent emergence of psychosis and midbrain GluN2A loss. **e,** Schematic representation of virus infusion protocol and timeline of phenotyping. **f,** Representative

image of GFP overlap with hemagglutinin (HA)-tagged CRISPR/Cas9 virus, quantification shows significantly more cells expressed both GFP and HA than only GFP (***p=0.0002) or only HA (***p=0.0001) **g,** Representative image showing GFP overlap with dopaminergic cells, quantification shows significantly more cells expressed both GFP and tyrosine hydroxylase (TH) than only GFP (*p=0.0152); (n=3 brains). Significance testing: one-way ANOVA; Tukey post hoc. **h,** Schematic representation of experimental design and example traces of slice electrophysiology experiments in adolescent brain. **i,** Quantification of peak amplitudes showed dopamine cells in DA-*Grin2a* KO brains were significantly less affected by TCN201 than controls (*p=0.049); (control n=9, DA-*Grin2a* KO n=11 cells). **j,** Quantification of peak amplitudes showed dopamine cells in DA-*Grin2a* KO brains were similarly affected by ifenprodil as controls; (control n=9, DA-*Grin2a* KO n=10 cells). **k,** Western blot analysis of adult brain tissue showed a significant reduction of normalized GluN2A in the VTA of DA-*Grin2a* KO animals compared to controls (**p =0.0082); (control n=4, DA-*Grin2a* KO n=3 brains). Significance testing: unpaired t-test. Error bars denote ±SEM.

DA-Grin2a KO adolescents retain typical spontaneous behaviors but have hypersensitivity to amphetamine

To assess the effect of DA-*Grin2a* KO on spontaneous unrewarded behaviors, a battery of tests was run to quantify spatial working memory, innate anxiety, and locomotion in adolescent animals (P45-55). Spontaneous alternation in a short-walled plus maze was used to assess spatial working memory. Since rats tend to explore novel arms over previously visited ones, impairments in working memory would emerge as reduced alternation in this task[40]. There were no significant between-group differences in spontaneous alternation in the plus maze, suggesting no effect of the DA-*Grin2a* KO on this form of spatial working memory in adolescents (**Figure 2a**). This was not due to differences in overall exploration of the maze as total arm entries were similar between groups (**Figure 2b**). In the elevated plus maze (EPM), avoidance of the open arms of the maze and preference for the walled-in, closed arms can be used to ascertain anxiety-like behavior in rodents[41]. There were no differences in anxiety-like behavior in the EPM as DA-*Grin2a* KO adolescents spent a similar percentage of time in the open and closed arms compared to controls (**Figure 2c,d**). Similar to EPM, avoidance of the exposed center portion of the open field maze is related to anxiety in rodents[42]. With this outcome measure, control and DA-*Grin2a* KO rats showed no significant difference in anxiety-like behavior in the open field task (**Figure 2e**). This was not a result of an impaired locomotor ability, as both treatment groups traveled similar distances in the open field (**Figure 2f**). We then performed a pharmacological challenge with amphetamine to determine the effect of the psychostimulant on locomotion. After the baseline period in the open field, animals received an injection of 1.0 mg/kg amphetamine and were returned to the field to

determine their responsivity to the locomotor effects of the drug (**Figure 2g**). Amphetamine elicits hyperlocomotion by increasing dopamine levels, and hyperreactivity to the locomotor effects of this drug is considered a behavioral correlate of psychosis[43,44]. When we analyzed distance traveled in the open field under amphetamine in 10-min bins, we found DA-*Grin2a* KO rats displayed hypersensitivity to amphetamine (**Figure 2h**). Thus while "baseline" spontaneous behavior was not affected in DA-*Grin2a* KO animals, a difference emerged when the system was challenged by amphetamine.

Figure 2: DA-*Grin2a* KO adolescents maintain typical spontaneous maze behaviors but show hypersensitivity to amphetamine

a, b, DA-*Grin2a* KO animals had no significant differences from controls in percent alternation or total arm entries in the plus maze; (control *n*=5, DA-*Grin2a* KO *n*=8 rats). **c, d,** Control and DA-*Grin2a* KO animals showed similar percent of the time spent in either the open or closed arms of the Elevated Plus Maze; (control *n*=9, DA-*Grin2a* KO *n*=8 rats). **e, f,** DA-*Grin2a* KO animals showed no significant differences from control animals in time spent in the center of the open field or in general locomotor activity in this maze; (control *n*=9, DA-*Grin2a* KO *n*=8 rats). Significance testing: unpaired t-test. **g,** Schematic depicting amphetamine (AMPH) administration in the open field. **h,** Distance traveled in 10 min time

bins showed overall significant effect of treatment (*p=0.0397) with DA-*Grin2a* KO animals
having higher distance traveled than controls; (control *n*=11, DA-*Grin2a* KO *n*=17 rats).
Baseline (BL) bin was determined from the final 10 min of open field testing in (f). Significant
testing: two-way ANOVA. Error bars denote ± SEM.

DA-Grin2a KO disrupts adolescent reward-guided behavior during operant and Pavlovian
conditioning in a construct-selective manner

The dopamine system is implicated in effort-, reward-, and punishment-related learning
and associated behaviors. We used two distinct conditioning paradigms to assess the
effect of *DA-Grin2a* KO on these learned behaviors.

<u>Effort related operant behavior</u>

Abnormalities in effort expenditure during reward seeking are a clinically observed
phenomena in people with SCZ. When asked to perform high or low effort tasks for reward,
individuals with SCZ show no deficits in ability or want to expend effort but inefficiently
scale high effort choices with high reward magnitude or probability. Specifically, compared
to controls, they have reduced capacity to rely on expected value to optimize effort
allocation[45-47]. To investigate this in our model, we ran a series of reward-based operant
tasks. Adolescent animals (P35±1 day) first learned to execute a nose poke in response to a
light cue to receive a sucrose pellet reward (FR1), then progressed to 2 sessions of
executing 5 nose pokes to receive the reward (FR5) (**Figure 3a**). In these low-effort operant
sessions, control and DA-*Grin2a* KO animals performed similarly. Cue-action latencies,
action-reward latencies, and total completed trials were not significantly different between
groups (**Figure 3b-d**), suggesting basic operant reward association learning was conserved
in the DA-*Grin2a* KO animals. Animals were then tested on a progressive ratio task for 4
sessions. The task began with an FR1 trial, but in each subsequent trial the response
requirement to receive a reward increased exponentially (**Figure 3e**). Progressive ratio
allows for measures of motivation through breakpoint (or highest completed ratio) and
effort optimization through response rate and post reinforcement pause (PRP) duration[48].
DA-*Grin2a* KO adolescents did not differ from controls in their breakpoint (**Figure 3f**).
However, while animals in both treatment groups completed a similar number of trials,
their overall response rates significantly differed in these sessions (**Figure 3g**). Overall
response rate was calculated as total responses divided by total trial time, so increased
overall response rates could reflect either enhanced motor capacity (i.e., faster
responding) or increased motivation to engage in effortful behavior (i.e., shorter initiation
delays)[48]. To dissociate these contributions, we calculated running response rates (total
responses per second following the first active response) and PRPs (latency from cue onset
to trial initiation). We observed no difference in running response rates between groups

(**Figure 3h**), but a significant difference in PRPs (**Figure 3i**). Typically, PRPs increase as a
function of an increasing response requirement[48] as observed in control animals. The DA-
Grin2a KO adolescents, however, did not flexibly change this response and maintained
stable, and shorter, response latencies despite the increase in action requirement. Thus,
while general motivational responding was normal in the DA-*Grin2a* KO animals, their
abnormal post reinforcement behavior in high effort trials suggests disrupted effort
optimization and impaired ability to use feedback to modify motivated actions.

Figure 3: Adolescent DA-*Grin2a* KO results in disrupted effort optimization in
progressive ratio
a, Representative schematic of rat behaving in operant box during the progressive ratio (PR)
task. **b, c, d,** Data collected during FR1 and FR5 training sessions show no group differences
in basic operant behavior. Cue-action latency (b), action-reward latency (c), and total trials
completed (d) were similar between groups. **e,** Graphic representation of exponential
increase in responses required with each trial during the progressive ratio (PR) schedule. **f,**
DA-*Grin2*a KO animals showed no differences in breakpoint from controls across the four PR

testing days. **g,** DA-*Grin2a* KOs had higher overall response rates compared to control
animals, main effect of treatment (*p=0.0335). **h,** No difference between groups in running
response rate. **i,** Significant interaction between treatment and ratio requirement
(**p=0.0065) in average post-reinforcement pause (PRP), post hoc analysis revealed
significant difference in PRP when ratio requirement was 77 (**p=0.0050) and 95
(****p<0.0001); (control n=10, DA-*Grin2a* KO n=10 rats). Significance testing: two-way
ANOVA: Sidak post hoc. Error bars denote ± SEM.
245

<u>Pavlovian learning and reversal learning</u>

Individuals with SCZ display impairments in reversal learning[49,50]. To test whether this
deficit occurs in our adolescent rat model, valence-specific behavioral flexibility was
assessed with a flexible contingency learning (FCL) task[51-53]. Adolescent animals (P35±1
day) were exposed to two novel conditioned stimuli (CS), which appeared in random order
during the behavior session, each of which predicted the delivery of an unconditioned
stimulus (US): either a reward or shock. After 5 sessions, during which associations with
each CS were learned, the contingencies were switched so the CS that was initially
associated with reward (referred to as CS_A) now predicted shock and the CS that previously
predicted shock (referred to as CS_B) was now followed by reward delivery (**Figure 4a**). The
key behavioral readout for this task was discrimination index (DI), which was the number of
reward port entries an animal made during each CS normalized to entries made during the
inter-trial-interval (ITI). The reward-predictive CS_A and its subsequent reversal to shock-
predictive was learned similarly by both DA-*Grin2a* KO and control animals (**Figure 4b**). The
shock-predictive CS_B was also learned similarly between groups as demonstrated through
suppressed DI values. After reversing the CS_B association to reward, however, DA-*Grin2a*
KO animals developed higher levels of conditioned discrimination compared to controls
(**Figure 4c**). Additional analysis of these data showed this difference in DI was not due to
faster latencies to respond to the cue and, while not significant, may be more influenced by
approach probability (**Figure S2**). To determine if the observed behavioral difference was
driven by distinct learning processes, we applied computational models rooted in
Pavlovian learning theory to the FCL data. We employed two distinct models: the Rescorla-
Wagner model, which infers learning from immediate prediction errors with a fixed learning
rate, and the Pearce-Hall model, which predicts learning from stimulus associability based
on prior prediction errors and allows for more flexible learning rates. These models both
work to explain how associations form between stimuli and outcomes, but the more
flexible Pearce-Hall model accounts for the influence of earlier stimuli presentations on
information updating and related phenomena such as latent inhibition that are not
captured with the Rescorla-Wagner model[54,55]. Models were fit to data collected during the
initial 5 reward association learning sessions and fit both control and DA-*Grin2a* KO data

with no clear preference for either learning mechanism (**Figure 4d, Figure S3**). Data
simulated using individual parameter estimates successfully recapitulated our observed
data during acquisition (**Figure 4e**). These similarities were expected because with no prior
learning about the reward-predicting stimulus, both models relied on similar principles of
associative learning. When the models were applied to data from reward association
learning *after* the contingency reversal, we observed that the control group's behavior was
better explained by Pearce-Hall whereas the DA-*Grin2a* KO animals' behavior was well
described by Rescorla-Wagner (**Figure 4f, Figure S3f,h**). Importantly, data simulated with
parameter estimates from the preferred model were able to recover the group differences
we initially observed (**Figure 4g**), while the unpreferred models tended to misestimate
behavioral trajectories within groups (**Figure S3e,g**). These results indicate that DA-*Grin2a*
KO animals expressed a reduced capacity to gate associative processes based on previous
learning, and this led to altered behavioral adaptation when outcome contingencies
switched.

Figure 4: Computational modeling of valence specific behavioral flexibility reveals
altered associative learning in DA-*Grin2a* KO adolescents

a, Schematic depicting flexible conditioning learning (FCL) task. **b,** Discrimination index (DI)
over sessions and averaged across sessions were used as a behavioral readout for the FCL

task. Both treatment groups responded to CS_A similarly, increasing food port entries relative
to baseline during CS_A presentation, then decreasing once the CS_A contingency was
switched on Session 6. **c,** DI over sessions and averaged across sessions in response to CS_B.
There were no significant differences between treatment groups in response to CS_B over all
sessions, but when DIs were pooled across initial learning (Session 1-5) and reversal
(Session 6-10) DA-*Grin2a* KO animals had significantly higher DIs than controls after the
contingency switch (*p=0.0204). **d,** Models were fit to data from acquisition (Sessions 1-5)
of the reward association (CS_A). Average and individual values (symbols) for difference in
model fit between Rescorla Wagner (RW) and Pearce-Hall (PH). Values <0 indicate a better
fit from PH mechanisms, >0 indicate a better fit from RW. No model was preferred for the
initial learning data. **e,** Observed (solid lines) and simulated (dashed lines) DIs for the
simplest (RW) model for each group recovered behavioral trajectories over acquisition
across both groups. **f,** Models were fit to data around the contingency reversal (Session 4-
10) for CS_B (shock to reward association). Average and individual values (symbols) for
difference in model fit between RW and PH. The RW model was preferred for all DA-*Grin2a*
KO animals, while the PH model was preferred for control animals (**p=0.0016). **g,** Observed
(solid lines) and simulated (dashed lines) DIs for the best fit model for each group recovered
differences in behavioral trajectories over reversal between groups; (control n = 6, DA-*Grin2a*
KO n = 5 rats). Significance testing: unpaired t-test. Error bars and shading denote ± SEM.

DA-Grin2a KO disrupts dopamine release in young adult animals

Disruptions in striatal dopamine neurotransmission are consistently reported in people
with SCZ and in individuals at high risk to develop this disorder[24,36,37,56]. Notably, these
disruptions are subtle and do not involve overt changes in dopamine related measures in
postmortem or imaging studies. Instead, PET and fMRI studies have reported small but
consistent disruptions in presynaptic activity[57] and in prediction error signaling in people
with SCZ[58,59]. To determine the long-term consequences of our DA-*Grin2a* KO on prediction
error signaling, we measured dopamine in the ventral striatum (VS) of young adult animals
(P60+) preforming the FCL task using fiber photometry with a fluorescent dopamine
reporter ($GRAB_{DA}$) (**Figure 5a,b**)[60]. This event-rich task allowed us to assess phasic
dopamine responses to novel stimuli, their transition to predictive stimuli, as well as to
expected and unexpected outcomes. Of note, these measures have two caveats. First, the
time constraint of rodent adolescence and other technical considerations did not allow us
to run these studies in adolescent animals, when the above behavioral measures were
performed. Second, the animals were tethered during the task, potentially influencing their
movement in the operant box, which may have contributed to no observed difference in DI
in the young adult animals (**Figure S4a**). Given these limitations, we primarily focused on
analyses of fiber photometry data that would provide fundamental information about CS
processing as opposed to correlation of dopamine signals with behavioral events.
Dopamine responses to novelty were assessed by quantifying phasic responses to CS_A and
CS_B onset in early trials of Session 1—before any association learning had occurred (**Figure
5c**). To quantify this response, we binned data from the first 25 presentations of CS_A into 5

trial bins (**Figure 5d**), then calculated the average z-score across the 2 s following CS$_A$
onset. Both groups' response to CS$_A$ decreased over time, and we found no significant
differences between control and DA-*Grin2a* KO animals (**Figure 5e**). A similar pattern was
observed in response to CS$_B$ (**Figure 5f,g**), suggesting novelty signaling was not disrupted by
DA-*Grin2a* KO manipulation.

Figure 5: Dopaminergic response to novelty in the ventral striatum is conserved in
young adult animals with the DA-*Grin2a* KO virus

a, Schematic of virus infusion and fiber implant locations. **b,** Schematic of fiber photometry
paradigm in operant box during FCL. **c,** Representative heatmaps from single animal
showing first 25 presentations of CS$_A$ and CS$_B$ during Session 1. **d,** Average fluorescent
signal, binned into 5 trial sections for the first 25 presentations of CS$_A$ in Session 1. **e,** Average
z-score across 2 s following CS$_A$ onset showed reduction in response from first 5
presentations to final 5 presentations (***p=0.0003). **f,** Average fluorescent signal for each
animal, binned into 5 trial sections for the first 25 presentations of CS$_B$ in Session 1. **g,**
Average z-score across 2 s following CS$_B$ onset showed reduction in response from first 5
presentations to final 5 presentations(**p=0.0073); (control n = 7, DA-*Grin2a* KO n = 8 rats).
Significance testing: two-way ANOVA. Error bars and shading denote ±SEM.

To determine if there were differences in dopaminergic responses once initial associations
were learned, we quantified signals during CS onset and US delivery in Session 5 (**Figure**

6a, S5b). Both groups displayed increases in ventral striatal dopamine in response to the
reward-predicting cue (CS_A) and reduced dopamine in response to the punishment-
predictive cue (CS_B) (**Figure 6b**). We observed a similar pattern in response to the two USs,
with reward delivery resulting in increased dopamine and a reduced dopamine response
following shock delivery (**Figure 6c**). Thus, we observed expected dopamine signaling[61] in
DA-*Grin2a* KOs during initial association learning. We then quantified the dopaminergic
response when the cue-outcome contingencies were reversed in Session 6. We split
Session 6 into Early trials (first 25 presentations) and Late trials (final 25 presentations) to
determine differences as the new associations were formed. During CS_A trials where
animals were expecting a reward but instead received a shock (**Figure 6d, S5c**), we
observed the expected decrease in dopamine signal to CS_A onset across Early to Late trials
as the cue began to lose its positive association (**Figure 6e**). In response to the unexpected
shock associated with CS_A, control animals showed the expected heightened
dopaminergic response in Early trials, which decreased by the Late trials, suggesting this
was a response to the mismatch between expected and actual outcome (**Figure 6f**).
Interestingly, DA-*Grin2a* KO rats showed a significantly muted response to the unexpected
shock during Early Session 6 trials (**Figure 6f**). During CS_B trials where the cue that was
initially associated with shock switched to reward (**Figure 6g, S5c**), we similarly observed
the expected increase in response to CS_B onset across Early to Late trials as the cue began
to lose its negative association (**Figure 6h**). As with CS_A trials, we observed a deficit in
dopaminergic response to the unexpected reward in DA-*Grin2a* KOs compared to controls
(**Figure 6i**). Together these data suggest that young adult animals with DA-*Grin2a* KO have
disrupted dopamine signaling in response to unexpected outcomes, regardless of valence.
As previously discussed, the behavioral consequences of this signaling deficit are unclear.
Dopamine levels differed between control and DA-*Grin2a* KOs in Early trials of Session 6;
however, there were no significant differences between the groups in terms of probability or
latency to approach the food port during these trials (**Figure S6**). In later sessions of FCL,
control animals showed increased probability to approach during CS_B presentation (**Figure
S4c**). However, fiber recording data from Session 10 revealed no significant signaling
differences between groups at that time (**Figure S7**). These findings suggest that while
manipulating *Grin2a* expression led to disruptions in dopaminergic signaling, the effect of
these differences on behavior remains unclear under the tested conditions. Future work
will explore how this altered dopamine modulation affects behavior in different contexts.

Figure 6: DA-*Grin2a* KO young adults show abnormal dopaminergic response to unexpected outcomes during FCL

a, Fluorescent signal of CS$_A$ and CS$_B$ trials on Session 5 of FCL. **b,** Average z-score across 2 s period at CS delivery showed positive responses to the reward-predicting cue (CS$_A$) and negative responses to the shock-predicting cue (CS$_B$) in both groups. **c,** Average z-score across 4 s period at US onset showed a similar pattern. **d,** Average fluorescence response during Early and Late CS$_A$ trials of Session 6. **e,** Average z-score across 2 s period at CS$_A$ onset, no significant group differences. **f,** Average z-score across 4 s period at shock delivery, control animals had significantly higher responses than DA-*Grin2a* KOs in early trials that fell by late trials (**p=0.0081). **g,** Average fluorescence response during Early and Late CS$_B$ trials of Session 6. **h,** Average z-score across 2 s period at CS$_B$ onset, no significant group differences. **i,** Average z-score across 4 s period at reward delivery, control animals had significantly higher responses than DA-*Grin2a* KOs in early trials that fell by late trials (**p=0.0336); (control *n* = 7, DA-*Grin2a* KO *n* = 8 rats). Significance testing: two-way ANOVA: Sidak post hoc. Error bars and shading denote ±SEM.

DISCUSSION

Informed by both GWAS and exome sequencing studies[9-11,62], we developed a rodent model with construct and face validity that provides insight about how variants of *GRIN2A* can lead to delayed onset of a collection of behavioral and dopamine abnormalities relevant to positive symptoms of SCZ. Our approach was guided by the well-established clinical literature showing a relationship between dopamine neurotransmission and psychosis[24,25,36-38,56,63-65], which often emerges later in development than cognitive and affective symptoms. *GRIN2A* encodes the GluN2A subunit of the NMDA receptor and has a cortical developmental expression profile that corresponds with the emergence of cognitive symptoms in childhood and early adolescence. We found that—unlike cortical regions where GluN2A levels remain stable throughout adolescence—there was significant loss of GluN2A in the dopamine neuron-containing regions of the midbrain during this developmental period. This led us to reason that the *GRIN2A* variants observed in the clinical population may exacerbate this developmental decline and contribute to the emergence of psychosis. We find that a genetically driven deficit of *Grin2a* in midbrain dopamine neurons of adolescent rats is sufficient to recapitulate some behaviors that are consistent with psychosis. These findings provide an etiologically relevant model for latent expression of psychosis which may aid with the development of novel course alternating treatments for SCZ.

GRIN2A and schizophrenia

Genetic variants in *GRIN2A* have been associated with SCZ diagnosis[9-11,62], providing further credence to the long-standing theory that abnormalities in NMDA receptor function are a core component of SCZ pathophysiology[1,66]. Global heterozygous loss of *Grin2a* in mice produces sustained phenotypes such as motor abnormalities, disrupted cortical activity, and changes in expression of multiple genes relevant to SCZ[15]. These global KO studies prompted our investigation into the consequences of an age- and cell type-specific *Grin2a* manipulation. We were particularly interested in how a *GRIN2A* loss of function may lead to latent expression of psychosis and dopamine processing deficits[24,25,35-38]. This led us to ask if selective loss of function of *Grin2a* in the dopamine neurons of adolescents is sufficient to produce phenotypes that are relevant to psychosis. We used a Cre-driven CRISPR/Cas9 virus in adolescent Th::Cre male and female rats to induce a deficit in *Grin2a* in midbrain dopamine neurons. This approach was successfully implemented despite the short duration of rodent adolescence. Moreover, the animals lacked any gross motor or learning deficits as measured by operant, Pavlovian, and maze tasks. The behavioral and dopamine release changes we observed were highly selective to constructs relevant to psychosis

including hypersensitivity to psychostimulants, inefficient effort optimization, and reduced
capacity to utilize feedback to guide behavior.

DA-*Grin2a* KO results in adolescent phenotype relevant to positive SCZ symptoms

Clinically, SCZ is defined as a heterogeneous psychiatric disorder characterized by the
presence of positive, negative, and cognitive symptoms. To determine the extent of the
effect of our manipulation, we included a variety of SCZ-related behavioral tasks in our
battery. We found a specific behavioral phenotype in which adolescent animals with the
DA-*Grin2a* KO showed abnormalities in line with early emerging positive symptoms.
Consistent with clinical studies showing the exacerbation of positive symptoms in people
with SCZ by psychostimulants, we determined DA-*Grin2a* KO adolescents had increased
sensitivity to the locomotor effects of amphetamine. When effort-based reward learning
was assessed with an exponential progressive ratio, we observed control rats suppressed
their stimulus-driven responding in high effort trials, while DA-*Grin2a* KO animals did not. In
the classical conditioning task, behavioral differences between control and DA-*Grin2a* KO
animals were observed selectively during contingency reversal sessions. These behavioral
data, reinforced with modeling, revealed DA-*Grin2a* KO animals showed reduced capacity
to use feedback to guide behavior by incorporating previous learning with new experiences.
Together these data reflect clinically observed phenomena relevant to positive SCZ
symptoms.
464
DA-*Grin2a* KO results in disrupted dopamine signaling

Abnormalities in dopamine neurotransmission in people with SCZ are consistently
reported. Evidence for faulty prediction error signaling in schizophrenia comes from fMRI
studies showing abnormalities in regional activity during reinforcement learning.
Specifically, people with SCZ have reduced activity in the midbrain and ventral striatum
when presented with an unexpected outcome, while maintaining a typical level of activity
in response to expected outcomes[67,68].

To drive learning, dopamine neurons increase activity when an unexpected event
occurs[69,70]. After the contingency reversal in the FCL, the robust increases in synaptic
dopamine in Early trials observed in control animals was attenuated in the DA-*Grin2a* KOs.
This suggests that the DA-*Grin2a* KO selectively impairs the ability of dopamine neurons to
increase signaling when there is a difference between expected vs actual outcomes. These
data demonstrate a specific role for GluN2A-containing NMDA receptors on midbrain
dopamine neurons in prediction error signaling in response to unexpected stimuli. This is
consistent with clinical data[67,68], providing a potential mechanism through which a genetic

variant of *GRIN2A* could contribute to the deficits in prediction error encoding observed in
people with SCZ.

Conclusion and Limitations

Our findings provide mechanistic insight into how variants of *GRIN2A* may lead to the latent
presentation of abnormalities in dopamine dynamics in SCZ. Our aim was not to produce a
full model of *GRIN2A* loss of function, but to tease apart how specific cell-types and neural
systems may be affected by this mutation. While the mutation is not confined to a single
region in humans, our approach is mechanistically informative and provides a novel
potential model for designing and testing course-altering treatments for transition to SCZ in
vulnerable individuals. Decades of translational studies focused on the NMDA receptor
have led to the discovery of potential treatments for SCZ[1,71,72], some of which showed
clinical efficacy for symptom amelioration only in younger individuals[73]. Clinical trials with
these drugs may have failed in the past because they were tested in chronically ill patients
with decades of antipsychotic drug treatment, which may have remodeled and changed
the nature of dopamine abnormalities. In younger people with—or at high risk to develop—
SCZ, these targets or newer GluN2A-selective positive allosteric modulators may provide a
precise approach for preventing or altering the course of the illness.

METHODS

Subjects

499 All animal procedures were approved by the Institutional Animal Care and Use Committee
500 at Oregon Health & Science University. Long Evan's male and female rats were bred in
501 house and maintained on a reverse 12hr light/dark cycle with ad libitum access to food and
502 water unless otherwise stated.

Tissue collection and protein extraction

Brain levels of GluN2A were examined in animals aged P21, P35, P45, and P80. Tissue
collection methods were adapted from Wager-Miller[74]. Briefly, at each age of interest,
brains were harvested, and flash frozen in isopentane that was pre-chilled with dry ice.
Whole brains were stored in conical tubes at -80°C until dissection.

While maintaining a frozen condition, brains were sliced into 1mm sections using a pre-
chilled brain matrix and razor blades. Bilateral micropunches of tissue were collected from
regions of interest using a rat brain atlas as a reference[75]. To extract protein, tissue
samples were homogenized in a lysis buffer containing protease and phosphatase
inhibitors. Protein concentration was quantified with a Pierce BCA Protein Assay Kit
(ThermoFisher). Homogenates were aliquoted and stored at -80°C until use.

Protein quantification

Automated immuno-quantification was run on a Wes instrument (Bio-Techne) according to
the manufacturer's protocol[76]. Briefly, each tissue sample was diluted to a concentration
of 0.08 µg/µL protein and mixed with dithiothreitol and a fluorescent molecular weight
marker at a ratio of 5:1. These samples were heated to 95°C for 5 min for protein
denaturation and then loaded into the allocated wells of the Wes plate. Other wells were
loaded with blocking reagent, primary antibody (Rabbit-anti-NMDAR2A (Sigma Aldrich,
#M264)), horseradish peroxidase (HRP)-conjugated secondary antibody, and a mix of
Luminol-S and Peroxide (Bio-Techne Anti-Rabbit Detection Module, #DM-001). Size-based
separation electrophoresis, immobilization, and immunodetection were automatically run
using the capillary array system of the Wes (**Figure S1**). All samples were normalized to
total protein as a loading control (Bio-Techne Total Protein Detection Module for
Chemiluminescence, #DM-TP01). Densitometric analysis was performed using the
Compass software (v5.0.1) from Bio-Techne.

Tissue dissection, micropunching, protein extraction, and Western blots were run as
described above on punches collected from the VTA of virus infused animals.

530
Viral production

The Cre-inducible recombinant adeno-associated virus (AAV) vector constructs utilized in
this study were produced and gifted by Larry Zweifel[77]. Viruses in these experiments were:
AAV1-FLEX-Cas9-U6-sgGrin2a for dopamine neuron-specific *Grin2a* knockout (DA-*Grin2a*
KO) (Addgene, #124850) which targets the first conserved exon (exon 6) across splice
variants in both rat and mouse (sgRNA: 5' GAGCAGGCAACCGGCTTGCCC 3'). AAV1-FLEX-
Cas9-U6-sgRosa26 was used as a control (Addgene, #159914) (sgRNA 5'
CTCGATGGAAAATACTCCGAG 3'). These viruses were co-infused with AAV1-hSyn-DIO-
EGFP (Addgene, #50457) at a 3:1 ratio to aid with visualization in subsequent experiments.

Stereotaxic surgeries

Juvenile rats (P21±1), were anesthetized with isoflurane and placed in a stereotaxic
apparatus and body temperature was maintained using a water heating circulation pump
(E-Z Systems). The scalp was incised and two bilateral craniotomies over the VTA were
performed. Four 250 nL infusions at the bilateral VTA sites (AP -4.9 mm, ML ±0.7 mm, DV -
6.5 and -6.0 mm from dura) were delivered using a syringe pump (World Instruments) and
micro infusion syringe (Hamilton) at a rate of 100 nL/min. To prevent backflow, the syringe
was not removed until 5 min after infusions. The incision was then closed with surgical
sutures, treated with triple antibiotic, and animals were administered 5 mg/kg carprofen
subcutaneously to relieve pain and inflammation. Animals remained on heat and were
administered oxygen until consciousness was regained, then returned to their home cage.
There was a minimum of a two-week incubation period before experiments were run.

For fiber photometry experiments, juvenile (P21±1) animals were infused with either the
control or DA-*Grin2a* KO virus mixed with AAV9-hsyn-GRAB_DA2m (GRAB$_{DA}$) (addgene,
#140553) at a 1:1 ratio. These were infused bilaterally into the VTA as described above.

Once animals reached adulthood (P60-70), they were implanted with a 400 µm optical fiber
(Doric Lenses Inc) into the ventral striatum (AP +1.6 mm, ML 1.5 mm, DV -7.0 mm). Animals
were given at least one week to recover from the implant surgery before behavioral training.

Immunohistochemistry

Animals were euthanized with an intraperitoneal (IP) injection of 400 mg/kg chloral hydrate
and perfused via the vascular system with 0.1 M phosphate buffer solution (PBS) followed
by 4% paraformaldehyde (PFA). Brains were removed and stored in 4% PFA at 4°C
overnight, then transferred to a 30% sucrose solution and stored at 4°C until sectioning.

Coronal sections (35 µm) of each brain were collected using a cryostat (Lecia) and stored
in 0.1 M PBS with 0.05% sodium azide. Free-floating sections were blocked and
permeabilized at room temperature for 2 hrs in 0.1 M PBS, 0.25% Triton X, 10% normal
donkey serum. Sections were then incubated overnight at 4˚C with the following primary
antibodies: chicken anti-TH (1:500, #76442, abcam), rabbit anti-GFP (1: 500, #290,
abcam), or mouse anti-HA (1:500, MMS-101P, Biolegend). Sections were washed, then
incubated in secondary antibodies (AlexaFluor-594; 488, 1:1000) for 2 hrs at room
temperature. Sections were washed again and mounted onto Super Frost+ slides.
Coverslips were applied using Vectashield HardSet antifade mounting media.
Fluorescence was imaged using a Ziess Axiovert 200 microscope and cell counting was
done in Zen 2 software (Zeiss).

For fiber photometry experiments, sections containing the fiber tract were mounted onto
Super Frost+ slides. Coverslips were applied using Vectashield HardSet antifade mounting
media containing DAPI and imaged using a Ziess Axiovert 200 microscope and cell
counting was done in Zen 2 software (Zeiss). To confirm fiber placement in the ventral
striatum, images were superimposed with a brain atlas[75] (See hit map, **Figure S4a**).

Slice Electrophysiology

Two to three weeks following virus infusion (P35-42), rats were anesthetized with
isoflurane and decapitated. The brain was collected and deposited in artificial
cerebrospinal fluid (ACSF) at 32–35°C containing (in mM): 126 NaCl, 2.5 KCl, 1.2 $MgCl_2$, 2.4
$CaCl_2$, 1.4 NaH_2PO_4, 25 $NaHCO_3$, and 11 Dextrose. The ACSF used for brain extraction,
slicing and recovery also contained 2 mM kynurenic acid (Sigma) to prevent NMDA-
mediated excitotoxic damage. Horizontal slices (220 µm) containing the midbrain were cut
using a vibratome (Leica) in warm ACSF bubbled with 95% O_2/5% CO_2. Slices were allowed
to recover in the same buffer at 30°C for at least 30 min prior to recording. Hemisected
slices were transferred to the recording chamber, which was continuously perfused at 2 to
3 mL/min with bubbled ACSF at 34°C. In most experiments the recording buffer was
identical to the collection buffer except it did not contain any $MgCl_2$, to allow for NMDA
receptor activation by glutamate iontophoresis. In a subset of recordings, experiments
were started in Mg-containing ACSF and Mg-free ACSF was perfused later. All experiments
were completed within 7 h of slicing.

VTA dopamine neurons were identified by their morphology and the presence of GFP.
Recordings were made using glass pipettes (1.0 – 2.5 MΩ resistance) filled with an internal
solution containing (in mM): 100 K-gluconate, 20 NaCl, 1.5 $MgCl_2$, 10 HEPES (K), 2 ATP, 0.2
GTP, 10 phosphocreatine, and 10 BAPTA (4K). Recordings were made in whole-cell
configuration. After break-in, cells were voltage-clamped at −55 mV and resistance and
capacitance were monitored; cells with a series resistance ≥12 MΩ were discarded. The
dopaminergic identity of the patched cells was further confirmed by the presence of a
hyperpolarization-induced depolarizing current (I_h). Cells with an I_h < 200 pA were
discarded. Another glass pipette was filled with a 1 M solution of monosodium glutamate
(Sigma Aldrich). This pipette was lowered into the slice and placed in proximity to the
patched cell, while a continuous backing current of +1.0 nA was applied. Glutamate was
iontophoretically delivered by the application of 20 ms negative current pulses (-40.0 to -
80.0 nA). Location and current intensity were adjusted to produce a stable depolarizing
current in the patched cell.

Data were acquired with AxoGraph software (Berkeley, CA) and recordings were monitored
with LabChart (AD instruments, Colorado Springs, CO). Current responses in Mg-free
ACSF were recorded until at least 10 stable responses were obtained. For experiments
where MG-free ACSF was perfused after patching this took up to 25 min. The slice was then
perfused with Mg-free ACSF containing the GluN2A-specific antagonist TCN201 (30 μM).
Responses to glutamate puffs were then recorded until a stable state was obtained or for a
maximum of 20 minutes. The slice was then perfused with Mg-free ACSF containing 10 μM
ifenprodil, a GluN2B-specific antagonist. Responses were measured until a stable state
was reached, or a maximum of 20 min. Lastly, we bath applied 50 μM AP5 (Sigma Aldrich),
a non-selective NMDA blocker, to confirm the identity of the measured currents. Because
both TCN201 and ifenprodil do not readily wash out, each slice was only used for one
attempted recording. Recordings were analyzed post-hoc using AxoGraph by averaging the
last 5 stable pulses for each condition and measuring the peak amplitude of the resulting
trace. Data are expressed as a fraction of the Mg-free baseline.

Spontaneous behaviors

All behavioral testing was administered at P45-P55. Animals were habituated to handling
and the testing environment for two days preceding behavior sessions. All behaviors were
run in a dim, white light. On the day of testing, animals were habituated to the room for 2 hr
before trials were run. The assays were run as follows:

Elevated plus maze (EPM)

The EPM consisted of two opposing open arms and two opposing arms bordered
with 48 cm walls. Animals were placed in the center of the maze and allowed to
explore for 10 min and their movement tracked with PanLab SMART v3.0 software
(Harvard Apparatus). Percent of time spent in open and closed arms was evaluated.

Spontaneous alternation test

A four-arm, plus maze was used to assess spontaneous alternation as a measure of
spatial working memory. In this task, distinct visual cues were positioned on all four
walls of the behavioral testing room. Animals were placed in the center of the maze
and allowed to explore for 10 min. The order of arm entries was recorded manually
during the task. Percent alternation and total number of arm entries were
evaluated. An alteration consisted of four distinct arm choices out of five
consecutive arm entries. A 4/5 alternation score was determined by dividing the
total number of alternations into overlapping quintuplets by the number of possible
alternations and multiplying by 100.

Open field (OF) with Pharmacological Challenge

Animals were placed in an opaque square arena and allowed to explore for 20 min.
Movement was tracked with PanLab SMART v3.0 software. During this time,
baseline locomotion and percent time spent in center (defined as a 5 cm distance
from the walls of the arena) was calculated.

After OF habituation, animals received IP injections of 1.0 mg/kg amphetamine
(AMPH) and placed back into the arena for 1 hr while video recordings were
collected. Horizontal locomotion in cm was tracked with PanLab SMART v3.0
software as animals explored the maze.

Operant and Pavlovian conditioning

Adolescent behavioral training experiments were performed on animals aged P35-P45.
Rats were mildly food restricted and habituated to operant boxes (Coulbourn Instruments)
two days prior to experimentation. All behavioral tasks were run in red light and behavior
recorded with Graphic State software (Coulbourn Instruments). Each animal received one
of the following tasks:

Progressive ratio of reinforcement

Operant chambers contained one wall with a nose poke hole that could be illuminated and
an opposing wall with a food port where sucrose pellets were dispensed. Animals were
initially trained on a fixed ratio one (FR1) schedule where one nose poke in response to a
light cue resulted in the delivery of one sucrose pellet. After five days of training, animals
advanced to an FR5 schedule for two days. For the final four days, animals were tested
with a progressive ratio schedule, where the response ratio increased according to the
formula $5e^{(0.2n)-5}$, where n = trial number[78]. This resulted in an exponential increase in
response requirement (1, 2, 4, 6, 9, 12 etc.) when rounded to the nearest whole number.
Sessions were terminated if there was a 5 min cessation in responding, if a trial took longer
than 45 min to complete, or after 3 hr total. Latencies to respond to the cue, to retrieve a
reward, and total rewards received were calculated during FR1 and FR5 sessions. We
determined the breakpoint for each animal across the four progressive ratio sessions. For
each trial, the overall response rate was calculated as total responses divided by *total* trial
time, while running response rate was calculated as total responses per second following
trial initiation. Post-reinforcement pause (PRP) duration was defined as the interval from

cue onset to the initiation of the next trial after the preceding trial's completion. Overall
response rate, running response rate, PRP were calculated as an average of the testing
days. For trial-by-trial analyses in progressive ratio, only trials where 80% of animals
completed a given ratio were analyzed.

Flexible contingency learning (FCL)

Operant chambers contained one wall with a light cue and speaker and an opposing wall
with a food port where sucrose pellets were dispensed. We utilized a modified version of a
previously published task[51-53] abbreviated to better fit within the adolescent time window.
Briefly, animals were presented with two conditioned stimuli (CS): a 10 sec light or tone. In
a counterbalanced fashion, animals were conditioned to pair one CS with the delivery of a
sucrose pellet and the other with a mild foot shock (180 ms, 0.2 mA). These unconditioned
stimuli (US) immediately followed the termination of either CS. After five days of
conditioning, the initial associations were reversed for Sessions 6-10. In each session,
animals underwent 100 stimulus pairings (50 trials of each CS) delivered randomly with a
20 sec inter-trial interval (ITI). CS_A represents the cue that was initially appetitive and then
became aversive, CS_B represents the cue that was initially aversive and then switched to
appetitive.

Behavior was analyzed by normalizing the amount of nose pokes into the food port during
CS to baseline food port entries during the ITI. The following equation was used to
calculate this discrimination index (DI):

$$\left(\frac{\text{CS entries} - \text{last 10s ITI entries}}{\text{CS entries} + \text{last 10s ITI entries}}\right)$$

With this formula, a positive DI suggests conditioned approach, a null DI suggests no
conditioning has occurred, and a negative DI suggests conditioned suppression. Average
DIs across Sessions 1-5 were considered initial learning, while average DIs across
Sessions 6-10 were considered contingency reversal learning.

Additional analyses included probability of approaching the food port during the 10 s CS
presentation and 10 s following US delivery as well as average latency to approach the
food port following CS or US onset.

Computational modeling

We utilized computational models of learning to further dissect the pattern of responding
in individual subjects in each group as measured by the DI during the FCL task (see above).

Models

We compared two models of associative learning that explain learning as a function of
immediate prediction error (Rescorla-Wagner, RW)[79] or stimulus associability (Pearce-
Hall, PH)[80]. In each model, the conditioned strength (V) of a stimulus evolves trial-by-trial

according to experience; we predicted the DI as a proxy for the conditioned strength (V) of
the stimulus presented on each trial.

RW – Learning in the RW model depends on prediction error (δ) and a fixed learning rate. V
develops for each trial according to the difference between actual (λ_n) and expected
outcome (V_n) for a stimulus on a given trial (n). This prediction error is multiplied by a fixed
learning rate (α, free parameter constrained between 0-1) which controls the rate of
learning. This can be formulated as:

717
$$\delta = (\lambda_n - V_n)$$
718
$$V_n = V_n + \alpha\delta$$
719

PH - The PH model accounts for learning as a function of the associability (α) of a stimulus.
We used an extended formulation of the PH model with gradual learning[81] where α is
updated on each trial from its previous level by a prediction error (δ) that tracks the
absolute value of the difference between actual (λ_{n-1}) and expected outcomes (V_{n-1}) for that
stimulus on the prior trial (n-1). This update to the associability of the stimulus is
controlled by a free weighting parameter (γ, free parameter constrained between 0-1). The
update to V is formulated as:

727
$$\delta = |\lambda_{n-1} - V_{n-1}|$$
728
$$\alpha_n = \alpha_{n-1} + \gamma(\delta - \alpha_{n-1})$$
729
$$V_n = V_n + \alpha\lambda_n$$
Model fitting

To address the high trial-by-trial response variability, we smoothed responses using a 5-
trial moving average and fit each of the RW and PH models to the DI values for the
acquisition (Sessions 1-5) and reversal (Sessions 6-10) epochs for each subject. Initial
conditioning strength (V_0) and value of an outcome (λ) was determined for each subject to
account for subject differences in initial bias and outcome sensitivity. Thus, the initial
conditioning strength was determined by taking the average DI for the first half of Session 1
(for acquisition) or from pre-reversal (i.e. Sessions 4-5). Value of an outcome was
determined by the mean DI observed in the last session of acquisition or reversal. This
effectively normalized the initial and maximum 'strength' of the cue, leaving the learning
trajectory to be fit. Best-fitting model parameters (α and γ for RW and PH, respectively)
were estimated through grid search across a range of 1,000 equally spaced values and
selected via a least-squares approach.

Model comparison

Models were assessed by the total sum of squared residual error (SS) and models with the
lowest total SS were selected for each group. We also assessed our model selection with a
paired permutation approach. Briefly a null distribution was generated by shuffling
condition labels within subject 10,000 times (i.e. the within group paired nature of the data
was kept intact) and calculating the mean difference between PH and RW residual error.
We then assessed the proportion of instances where a value in the null distribution was

equal to or greater than the observed PH/RW difference from the unshuffled data to obtain
a p-value (one-tailed). To compare between groups, we quantified the difference in
residual sum of squares between RW and PH fits for each subject and compared groups
using a two-tailed t-test.

Data simulations

We validated parameter estimates for α or γ by simulating learning trajectories for each
subject from their best fit parameters. We then plotted the group-averaged response of the
preferred model to validate the model(s) could recover key trends in the data.

Statistical analysis

Statistical analyses were performed in GraphPad Prism (Version 10.0.1). Unpaired t-tests
were run in instances where two groups were compared. Otherwise, the data were
analyzed with either a one- or two-way ANOVA with Tukey or Sidak post hoc corrections,
respectively.

Fiber photometry recording

Two days preceding fiber photometry experiments, implanted animals were mildly food
restricted and habituated to handling, being attached to the recording patch cord, and to
the testing environment. Animals were then run on the FCL task as described above. Fiber
photometry experiments were executed using an RZ10X fiber photometry system (Tucker-
Davis Technologies). The Quick-Release Interconnect (Thorlabs) was used to mate the
implanted fiber optic cannula and the patch cord. Excitation light was passed through a
400 µm patch cord from 465 and 405 nm LEDs (Doric Lenses Inc), sinusoidally modulated
at 220 and 310-Hz, respectively. Signals were demodulated at 1kHz with a 6Hz low pass
filter in real-time using Synapse software (Tucker-Davis Technologies). Time-locked
behavioral events from the operant box were recorded through the RZ10X digital input
ports. Recording was performed on each day of the behavioral task.

Data analysis

Fiber photometry data were analyzed using Python 3 and the TDT Python package. Raw
fluorescence signals from 465 nm ($GRAB_{DA}$) and 405 nm (isosbestic) channels were
segmented into trials, each 30 s (10 s before CS onset to 10 s after US onset). For each trial,
the $GRAB_{DA}$ signal was fit to the isosbestic signal using a least-squares linear fit (numpy
polyfit) to correct for motion artifacts and photobleaching. The fitted signal was used to
compute the change in fluorescence ($\Delta F/F$) for each trial.

To assess event-related changes in striatal dopamine, peri-event z-scores were calculated
by normalizing the $\Delta F/F$ signal to the 10 s baseline period prior to CS onset (within the ITI).
Average responses were then computed across trials for each animal. Dopamine
responses to specific events were calculated as follows:

Novelty: First 25 CS_A presentations and first 25 CS_B presentations were grouped into 5 bins
of 5 trials each.

Initial association: Responses were averaged across all 50 CS_A trials and 50 CS_B trials in
Session 5.

Association reversal: following reversal in Session 6, responses to unexpected outcomes
were assessed by averaging the first 25 trials (Early) and the last 25 trials (Late) of each CS-
US pairing.

To quantify differences in synaptic dopamine between the treatment groups, we
determined the average z-score across two periods. The CS time bin consisted of the first 2
s following cue onset and US time bin consisted of the first 4 s following the onset of the
reward/shock.

Statistical analyses were performed in GraphPad Prism (Version 10.0.1). Average z-scores
were analyzed with a two-way ANOVA (with factors Session and Treatment) with Sidak post
hoc corrections if there was a significant interaction ($p<0.05$).

REFERENCES

1. Moghaddam, B., and Javitt, D. (2012). From revolution to evolution: the glutamate hypothesis of schizophrenia and its implication for treatment. Neuropsychopharmacology *37*, 4-15. 10.1038/npp.2011.181.

2. Coyle, J.T., Ruzicka, W.B., and Balu, D.T. (2020). Fifty Years of Research on Schizophrenia: The Ascendance of the Glutamatergic Synapse. Am J Psychiatry *177*, 1119-1128. 10.1176/appi.ajp.2020.20101481.

3. Law, A.J., and Deakin, J.F. (2001). Asymmetrical reductions of hippocampal NMDAR1 glutamate receptor mRNA in the psychoses. Neuroreport *12*, 2971-2974. 10.1097/00001756-200109170-00043.

4. Weickert, C.S., Fung, S.J., Catts, V.S., Schofield, P.R., Allen, K.M., Moore, L.T., Newell, K.A., Pellen, D., Huang, X.F., Catts, S.V., and Weickert, T.W. (2013). Molecular evidence of N-methyl-D-aspartate receptor hypofunction in schizophrenia. Mol Psychiatry *18*, 1185-1192. 10.1038/mp.2012.137.

5. Meador-Woodruff, J.H., Clinton, S.M., Beneyto, M., and McCullumsmith, R.E. (2003). Molecular abnormalities of the glutamate synapse in the thalamus in schizophrenia. Ann N Y Acad Sci *1003*, 75-93. 10.1196/annals.1300.005.

6. Javitt, D.C., and Zukin, S.R. (1991). Recent advances in the phencyclidine model of schizophrenia. Am J Psychiatry *148*, 1301-1308. 10.1176/ajp.148.10.1301.

7. Moghaddam, B., and Krystal, J.H. (2012). Capturing the angel in "angel dust": twenty years of translational neuroscience studies of NMDA receptor antagonists in animals and humans. Schizophr Bull *38*, 942-949. 10.1093/schbul/sbs075.

8. Singh, T., Poterba, T., Curtis, D., Akil, H., Al Eissa, M., Barchas, J.D., Bass, N., Bigdeli, T.B., Breen, G., Bromet, E.J., et al. (2022). Rare coding variants in ten genes confer substantial risk for schizophrenia. Nature *604*, 509-516. 10.1038/s41586-022-04556-w.

9. Schizophrenia Working Group of the Psychiatric Genomics, C. (2014). Biological insights from 108 schizophrenia-associated genetic loci. Nature *511*, 421-427. 10.1038/nature13595.

10. Ripke, S., O'Dushlaine, C., Chambert, K., Moran, J.L., Kahler, A.K., Akterin, S., Bergen, S.E., Collins, A.L., Crowley, J.J., Fromer, M., et al. (2013). Genome-wide association analysis identifies 13 new risk loci for schizophrenia. Nat Genet *45*, 1150-1159. 10.1038/ng.2742.

11. Stefansson, H., Ophoff, R.A., Steinberg, S., Andreassen, O.A., Cichon, S., Rujescu, D., Werge, T., Pietilainen, O.P., Mors, O., Mortensen, P.B., et al. (2009). Common variants conferring risk of schizophrenia. Nature *460*, 744-747. 10.1038/nature08186.

12. Trubetskoy, V., Pardinas, A.F., Qi, T., Panagiotaropoulou, G., Awasthi, S., Bigdeli, T.B., Bryois, J., Chen, C.Y., Dennison, C.A., Hall, L.S., et al. (2022). Mapping genomic loci implicates genes and synaptic biology in schizophrenia. Nature *604*, 502-508. 10.1038/s41586-022-04434-5.

847   13.   Hall, J., and Bray, N.J. (2022). Schizophrenia Genomics: Convergence on Synaptic
848   Development, Adult Synaptic Plasticity, or Both? Biol Psychiatry *91*, 709-717.
849   10.1016/j.biopsych.2021.10.018.
850   14.   Farsi, Z., Nicolella, A., Simmons, S.K., Aryal, S., Shepard, N., Brenner, K., Lin, S.,
851   Herzog, L., Moran, S.P., Stalnaker, K.J., et al. (2023). Brain-region-specific changes in
852   neurons and glia and dysregulation of dopamine signaling in Grin2a mutant mice.
853   Neuron *111*, 3378-3396 e3379. 10.1016/j.neuron.2023.08.004.
854   15.   Herzog, L.E., Wang, L., Yu, E., Choi, S., Farsi, Z., Song, B.J., Pan, J.Q., and Sheng, M.
855   (2023). Mouse mutants in schizophrenia risk genes GRIN2A and AKAP11 show EEG
856   abnormalities in common with schizophrenia patients. Transl Psychiatry *13*, 92.
857   10.1038/s41398-023-02393-7.
858   16.   Camp, C.R., Vlachos, A., Klockner, C., Krey, I., Banke, T.G., Shariatzadeh, N.,
859   Ruggiero, S.M., Galer, P., Park, K.L., Caccavano, A., et al. (2023). Loss of Grin2a
860   causes a transient delay in the electrophysiological maturation of hippocampal
861   parvalbumin interneurons. Commun Biol *6*, 952. 10.1038/s42003-023-05298-9.
862   17.   van Os, J., and Kapur, S. (2009). Schizophrenia. Lancet *374*, 635-645.
863   10.1016/S0140-6736(09)60995-8.
864   18.   Abi-Dargham, A. (2014). Schizophrenia: overview and dopamine dysfunction. J Clin
865   Psychiatry *75*, e31. 10.4088/JCP.13078tx2c.
866   19.   Grace, A.A. (2016). Dysregulation of the dopamine system in the pathophysiology of
867   schizophrenia and depression. Nat Rev Neurosci *17*, 524-532. 10.1038/nrn.2016.57.
868   20.   Kapur, S., and Seeman, P. (2001). Does fast dissociation from the dopamine d(2)
869   receptor explain the action of atypical antipsychotics?: A new hypothesis. Am J
870   Psychiatry *158*, 360-369. 10.1176/appi.ajp.158.3.360.
871   21.   Matthysse, S. (1973). Antipsychotic drug actions: a clue to the neuropathology of
872   schizophrenia? Fed Proc *32*, 200-205.
873   22.   Seeman, M.V. (2021). History of the dopamine hypothesis of antipsychotic action.
874   World J Psychiatry *11*, 355-364. 10.5498/wjp.v11.i7.355.
875   23.   Seeman, P., Lee, T., Chau-Wong, M., and Wong, K. (1976). Antipsychotic drug doses
876   and neuroleptic/dopamine receptors. Nature *261*, 717-719. 10.1038/261717a0.
877   24.   Abi-Dargham, A., Gil, R., Krystal, J., Baldwin, R.M., Seibyl, J.P., Bowers, M., van Dyck,
878   C.H., Charney, D.S., Innis, R.B., and Laruelle, M. (1998). Increased striatal dopamine
879   transmission in schizophrenia: confirmation in a second cohort. Am J Psychiatry
880   *155*, 761-767. 10.1176/ajp.155.6.761.
881   25.   Abi-Dargham, A., Rodenhiser, J., Printz, D., Zea-Ponce, Y., Gil, R., Kegeles, L.S.,
882   Weiss, R., Cooper, T.B., Mann, J.J., Van Heertum, R.L., et al. (2000). Increased
883   baseline occupancy of D2 receptors by dopamine in schizophrenia. Proc Natl Acad
884   Sci U S A *97*, 8104-8109. 10.1073/pnas.97.14.8104.
885   26.   Howes, O.D., Williams, M., Ibrahim, K., Leung, G., Egerton, A., McGuire, P.K., and
886   Turkheimer, F. (2013). Midbrain dopamine function in schizophrenia and depression:
887   a post-mortem and positron emission tomographic imaging study. Brain *136*, 3242-
888   3251. 10.1093/brain/awt264.

27. Bagasrawala, I., Memi, F., N, V.R., and Zecevic, N. (2017). N-Methyl d-Aspartate
Receptor Expression Patterns in the Human Fetal Cerebral Cortex. Cereb Cortex *27*,
5041-5053. 10.1093/cercor/bhw289.
28. Bar-Shira, O., Maor, R., and Chechik, G. (2015). Gene Expression Switching of
Receptor Subunits in Human Brain Development. PLoS Comput Biol *11*, e1004559.
10.1371/journal.pcbi.1004559.
29. Maussion, G., Diallo, A.B., Gigek, C.O., Chen, E.S., Crapper, L., Theroux, J.F., Chen,
G.G., Vasuta, C., and Ernst, C. (2015). Investigation of genes important in
neurodevelopment disorders in adult human brain. Hum Genet *134*, 1037-1053.
10.1007/s00439-015-1584-z.
30. Monyer, H., Burnashev, N., Laurie, D.J., Sakmann, B., and Seeburg, P.H. (1994).
Developmental and regional expression in the rat brain and functional properties of
four NMDA receptors. Neuron *12*, 529-540. 10.1016/0896-6273(94)90210-0.
31. Wenzel, A., Scheurer, L., Kunzi, R., Fritschy, J.M., Mohler, H., and Benke, D. (1995).
Distribution of NMDA receptor subunit proteins NR2A, 2B, 2C and 2D in rat brain.
Neuroreport *7*, 45-48.
32. Creese, I., Burt, D.R., and Snyder, S.H. (1976). Dopamine receptor binding predicts
clinical and pharmacological potencies of antischizophrenic drugs. Science *192*,
481-483. 10.1126/science.3854.
33. Snyder, S.H. (1976). The dopamine hypothesis of schizophrenia: focus on the
dopamine receptor. Am J Psychiatry *133*, 197-202. 10.1176/ajp.133.2.197.
34. Yun, S., Yang, B., Anair, J.D., Martin, M.M., Fleps, S.W., Pamukcu, A., Yeh, N.H.,
Contractor, A., Kennedy, A., and Parker, J.G. (2023). Antipsychotic drug efficacy
correlates with the modulation of D1 rather than D2 receptor-expressing striatal
projection neurons. Nat Neurosci *26*, 1417-1428. 10.1038/s41593-023-01390-9.
35. Angrist, B., Rotrosen, J., and Gershon, S. (1980). Responses to apomorphine,
amphetamine, and neuroleptics in schizophrenic subjects. Psychopharmacology
(Berl) *67*, 31-38. 10.1007/BF00427592.
36. Abi-Dargham, A., van de Giessen, E., Slifstein, M., Kegeles, L.S., and Laruelle, M.
(2009). Baseline and amphetamine-stimulated dopamine activity are related in
drug-naive schizophrenic subjects. Biol Psychiatry *65*, 1091-1093.
10.1016/j.biopsych.2008.12.007.
37. Laruelle, M., Abi-Dargham, A., van Dyck, C.H., Gil, R., D'Souza, C.D., Erdos, J.,
McCance, E., Rosenblatt, W., Fingado, C., Zoghbi, S.S., et al. (1996). Single photon
emission computerized tomography imaging of amphetamine-induced dopamine
release in drug-free schizophrenic subjects. Proc Natl Acad Sci U S A *93*, 9235-9240.
10.1073/pnas.93.17.9235.
38. Egerton, A., Chaddock, C.A., Winton-Brown, T.T., Bloomfield, M.A., Bhattacharyya,
S., Allen, P., McGuire, P.K., and Howes, O.D. (2013). Presynaptic striatal dopamine
dysfunction in people at ultra-high risk for psychosis: findings in a second cohort.
Biol Psychiatry *74*, 106-112. 10.1016/j.biopsych.2012.11.017.
39. Watanabe, M., Inoue, Y., Sakimura, K., and Mishina, M. (1992). Developmental
changes in distribution of NMDA receptor channel subunit mRNAs. Neuroreport *3*,
1138-1140. 10.1097/00001756-199212000-00027.

40. Kraeuter, A.K., Guest, P.C., and Sarnyai, Z. (2019). The Y-Maze for Assessment of
Spatial Working and Reference Memory in Mice. Methods Mol Biol *1916*, 105-111.
10.1007/978-1-4939-8994-2_10.
41. Montgomery, K.C. (1955). The relation between fear induced by novel stimulation
and exploratory behavior. J Comp Physiol Psychol *48*, 254-260. 10.1037/h0043788.
42. Broadhurst, P.L. (1969). Psychogenetics of emotionality in the rat. Ann N Y Acad Sci
159, 806-824. 10.1111/j.1749-6632.1969.tb12980.x.
43. Powell, C.M., and Miyakawa, T. (2006). Schizophrenia-relevant behavioral testing in
rodent models: a uniquely human disorder? Biol Psychiatry *59*, 1198-1207.
10.1016/j.biopsych.2006.05.008.
44. Bondi, C., Matthews, M., and Moghaddam, B. (2012). Glutamatergic animal models
of schizophrenia. Curr Pharm Des *18*, 1593-1604. 10.2174/138161212799958576.
45. McCarthy, J.M., Treadway, M.T., Bennett, M.E., and Blanchard, J.J. (2016). Inefficient
effort allocation and negative symptoms in individuals with schizophrenia.
Schizophr Res *170*, 278-284. 10.1016/j.schres.2015.12.017.
46. Treadway, M.T., Peterman, J.S., Zald, D.H., and Park, S. (2015). Impaired effort
allocation in patients with schizophrenia. Schizophr Res *161*, 382-385.
10.1016/j.schres.2014.11.024.
47. Kirschner, M., Hager, O.M., Bischof, M., Hartmann-Riemer, M.N., Kluge, A., Seifritz,
E., Tobler, P.N., and Kaiser, S. (2016). Deficits in context-dependent adaptive coding
of reward in schizophrenia. NPJ Schizophr *2*, 16020. 10.1038/npjschz.2016.20.
48. Bradshaw, C.M., and Killeen, P.R. (2012). A theory of behaviour on progressive ratio
schedules, with applications in behavioural pharmacology. Psychopharmacology
(Berl) *222*, 549-564. 10.1007/s00213-012-2771-4.
49. Reddy, L.F., Waltz, J.A., Green, M.F., Wynn, J.K., and Horan, W.P. (2016). Probabilistic
Reversal Learning in Schizophrenia: Stability of Deficits and Potential Causal
Mechanisms. Schizophr Bull *42*, 942-951. 10.1093/schbul/sbv226.
50. Culbreth, A.J., Gold, J.M., Cools, R., and Barch, D.M. (2016). Impaired Activation in
Cognitive Control Regions Predicts Reversal Learning in Schizophrenia. Schizophr
Bull *42*, 484-493. 10.1093/schbul/sbv075.
51. Kim, Y., Wood, J., and Moghaddam, B. (2012). Coordinated activity of ventral
tegmental neurons adapts to appetitive and aversive learning. PLoS One *7*, e29766.
10.1371/journal.pone.0029766.
52. Kim, Y.B., Matthews, M., and Moghaddam, B. (2010). Putative gamma-aminobutyric
acid neurons in the ventral tegmental area have a similar pattern of plasticity as
dopamine neurons during appetitive and aversive learning. Eur J Neurosci *32*, 1564-
1572. 10.1111/j.1460-9568.2010.07371.x.
53. Lefner, M.J., and Moghaddam, B. (2025). Flexible updating of reward and
punishment contingencies by VTA GABA neurons. Curr Biol.
10.1016/j.cub.2025.07.021.
54. Lubow, R.E. (2005). Construct validity of the animal latent inhibition model of
selective attention deficits in schizophrenia. Schizophr Bull *31*, 139-153.
10.1093/schbul/sbi005.

55. Lubow, R.E., and Gewirtz, J.C. (1995). Latent inhibition in humans: data, theory, and
implications for schizophrenia. Psychol Bull *117*, 87-103. 10.1037/0033-
2909.117.1.87.
56. Laruelle, M., and Abi-Dargham, A. (1999). Dopamine as the wind of the psychotic
fire: new evidence from brain imaging studies. J Psychopharmacol *13*, 358-371.
10.1177/026988119901300405.
57. Howes, O.D., Kambeitz, J., Kim, E., Stahl, D., Slifstein, M., Abi-Dargham, A., and
Kapur, S. (2012). The nature of dopamine dysfunction in schizophrenia and what this
means for treatment. Arch Gen Psychiatry *69*, 776-786.
10.1001/archgenpsychiatry.2012.169.
58. Millard, S.J., Bearden, C.E., Karlsgodt, K.H., and Sharpe, M.J. (2022). The prediction-
error hypothesis of schizophrenia: new data point to circuit-specific changes in
dopamine activity. Neuropsychopharmacology *47*, 628-640. 10.1038/s41386-021-
01188-y.
59. Katthagen, T., Kaminski, J., Heinz, A., Buchert, R., and Schlagenhauf, F. (2020).
Striatal Dopamine and Reward Prediction Error Signaling in Unmedicated
Schizophrenia Patients. Schizophr Bull *46*, 1535-1546. 10.1093/schbul/sbaa055.
60. Sun, F., Zeng, J., Jing, M., Zhou, J., Feng, J., Owen, S.F., Luo, Y., Li, F., Wang, H.,
Yamaguchi, T., et al. (2018). A Genetically Encoded Fluorescent Sensor Enables
Rapid and Specific Detection of Dopamine in Flies, Fish, and Mice. Cell *174*, 481-
496 e419. 10.1016/j.cell.2018.06.042.
61. Bromberg-Martin, E.S., Matsumoto, M., and Hikosaka, O. (2010). Dopamine in
motivational control: rewarding, aversive, and alerting. Neuron *68*, 815-834.
10.1016/j.neuron.2010.11.022.
62. Ohi, K., Shimada, T., Nitta, Y., Kihara, H., Okubo, H., Uehara, T., and Kawasaki, Y.
(2016). Specific gene expression patterns of 108 schizophrenia-associated loci in
cortex. Schizophr Res *174*, 35-38. 10.1016/j.schres.2016.03.032.
63. Howes, O.D., Montgomery, A.J., Asselin, M.C., Murray, R.M., Valli, I., Tabraham, P.,
Bramon-Bosch, E., Valmaggia, L., Johns, L., Broome, M., et al. (2009). Elevated
striatal dopamine function linked to prodromal signs of schizophrenia. Arch Gen
Psychiatry *66*, 13-20. 10.1001/archgenpsychiatry.2008.514.
64. Howes, O.D., and Nour, M.M. (2016). Dopamine and the aberrant salience
hypothesis of schizophrenia. World Psychiatry *15*, 3-4. 10.1002/wps.20276.
65. Fletcher, P.C., and Frith, C.D. (2009). Perceiving is believing: a Bayesian approach to
explaining the positive symptoms of schizophrenia. Nat Rev Neurosci *10*, 48-58.
10.1038/nrn2536.
66. Olney, J.W., and Farber, N.B. (1995). Glutamate receptor dysfunction and
schizophrenia. Arch Gen Psychiatry *52*, 998-1007.
10.1001/archpsyc.1995.03950240016004.
67. Morris, R.W., Vercammen, A., Lenroot, R., Moore, L., Langton, J.M., Short, B.,
Kulkarni, J., Curtis, J., O'Donnell, M., Weickert, C.S., and Weickert, T.W. (2012).
Disambiguating ventral striatum fMRI-related BOLD signal during reward prediction
in schizophrenia. Mol Psychiatry *17*, 235, 280-239. 10.1038/mp.2011.75.

68. Corlett, P.R., Murray, G.K., Honey, G.D., Aitken, M.R., Shanks, D.R., Robbins, T.W.,
Bullmore, E.T., Dickinson, A., and Fletcher, P.C. (2007). Disrupted prediction-error
signal in psychosis: evidence for an associative account of delusions. Brain *130*,
2387-2400. 10.1093/brain/awm173.
69. Schultz, W., Dayan, P., and Montague, P.R. (1997). A neural substrate of prediction
and reward. Science *275*, 1593-1599. 10.1126/science.275.5306.1593.
70. Schultz, W. (1998). Predictive reward signal of dopamine neurons. J Neurophysiol
80, 1-27. 10.1152/jn.1998.80.1.1.
71. Balu, D.T., and Coyle, J.T. (2015). The NMDA receptor 'glycine modulatory site' in
schizophrenia: D-serine, glycine, and beyond. Curr Opin Pharmacol *20*, 109-115.
10.1016/j.coph.2014.12.004.
72. Moghaddam, B., and Adams, B.W. (1998). Reversal of phencyclidine effects by a
group II metabotropic glutamate receptor agonist in rats. Science *281*, 1349-1352.
10.1126/science.281.5381.1349.
73. Kinon, B.J., Millen, B.A., Zhang, L., and McKinzie, D.L. (2015). Exploratory analysis for
a targeted patient population responsive to the metabotropic glutamate 2/3
receptor agonist pomaglumetad methionil in schizophrenia. Biol Psychiatry *78*, 754-
762. 10.1016/j.biopsych.2015.03.016.
74. Wager-Miller, J., Murphy Green, M., Shafique, H., and Mackie, K. (2020). Collection
of Frozen Rodent Brain Regions for Downstream Analyses. J Vis Exp. 10.3791/60474.
75. Paxinos, G., & Watson, C. (1998). A stereotaxic atlas of the rat brain. (New York:
Academic).
76. Luck, C., Haitjema, C., and Heger, C. (2021). Simple Western: Bringing the Western
Blot into the Twenty-First Century. Methods Mol Biol *2261*, 481-488. 10.1007/978-1-
0716-1186-9_30.
77. Hunker, A.C., Soden, M.E., Krayushkina, D., Heymann, G., Awatramani, R., and
Zweifel, L.S. (2020). Conditional Single Vector CRISPR/SaCas9 Viruses for Efficient
Mutagenesis in the Adult Mouse Nervous System. Cell Rep *30*, 4303-4316 e4306.
10.1016/j.celrep.2020.02.092.
78. Richardson, N.R., and Roberts, D.C. (1996). Progressive ratio schedules in drug self-
administration studies in rats: a method to evaluate reinforcing efficacy. J Neurosci
Methods *66*, 1-11. 10.1016/0165-0270(95)00153-0.
79. Rescorla, R.A.W., A.R. (1972). A theory of Pavlovian conditioning: Variations in the
effectiveness of reinforcement and nonreinforcement. In Classical conditioning II:
Current research and theory, A.H.P. Black, W. F., ed. (Appleton-Century-Crofts), pp.
64–99.
80. Pearce, J.M., and Hall, G. (1980). A model for Pavlovian learning: variations in the
effectiveness of conditioned but not of unconditioned stimuli. Psychol Rev *87*, 532-
552.
81. Roesch, M.R., Esber, G.R., Li, J., Daw, N.D., and Schoenbaum, G. (2012). Surprise!
Neural correlates of Pearce-Hall and Rescorla-Wagner coexist within the brain. Eur J
Neurosci *35*, 1190-1200. 10.1111/j.1460-9568.2011.07986.x.

1061