Adaptive Gaze and Hand Coordination while Manipulating and Monitoring the Environment in Parallel

Jolande Fooken[1,2]*, Roland S. Johansson[3]†, and J. Randall Flanagan[2]†

[1]Department of Psychology and Centre for Cognitive Science, Technical University of Darmstadt, Darmstadt, Germany

[2]Department of Psychology and Centre for Neuroscience Studies, Queen's University, Kingston, Ontario, Canada

[3]Department of Medical and Translational Biology, Umeå University, Umeå, Sweden

*Corresponding author: jolande.fooken@tu-darmstadt.de

† Equal contribution

Credit authorship contribution statement

Conceptualization (JF, RSJ, JRF), Methodology (JF, RSJ, JRF), Validation (JF, RSJ, JRF), Formal analysis (JF, RSJ, JRF), Resources (RSJ, JRF), Data Curation (JF, RSJ), Writing – Original Draft (JF, RSJ, JRF), Writing – review & editing (JF, RSJ, JRF), Visualization (JF, RSJ, JRF), Funding acquisition (JF, RSJ, JRF)

Declaration of Competing Interest

The authors have no competing financial interests to disclose.

Acknowledgments

This work was supported by a Deutsche Forschungsgemeinschaft (DFG) Research Fellowship to JF (FO 1347/1-1), the Swedish Research Council Project (grant 22209 to RSJ), the Canadian Institutes for Health Research (grant RGPIN/05944-2019 to JRF), and the Natural Science and Engineering Research Council of Canada (grant 156173 to JRF).

Abstract

In real-world tasks, visually guided manual actions are often performed in parallel with visual
monitoring of the environment, creating competition for gaze. This study examined how
participants coordinated gaze and hand movements while performing a manipulation task—
grasping a small ball with either fingertips or tweezers and depositing it into a slot—while
concurrently tasked with monitoring a display for probabilistically occurring letter changes.
Gaze fixations away from the text display were exclusively directed to the ball and slot, and
were temporally linked to key contact events (ball contact and slot entry). In fingertip trials,
action-task fixations, when present, were brief and could support directing the hand using
peripheral vision. In tweezer trials, action-task fixations were more frequent and helped both
directing the hand with peripheral vision and guiding it with central vision. Participants
identified and leveraged the temporal regularities in the monitoring task to reduce competition
for gaze resources between tasks. Specifically, they adjusted both their gaze behaviour and hand
movement timing to lower the likelihood that action task fixations would coincide with letter
changes. These findings demonstrate how individuals integrate sensorimotor demands with
environmental statistics to manage competition for gaze when concurrently acting on and
monitoring the environment.

Introduction

Gaze fixations, occurring between eye movements, play a vital role in both perceiving the world

and in planning and controlling actions (Hayhoe, 2017; Kowler, 2011; Land, 2006; Yarbus,

1967). Although research has extensively examined gaze control in action and perception tasks

independently, real-life scenarios often demand concurrent performance of visually guided

actions and visual perception tasks (Fooken et al., 2023; Land and Furneaux, 1997), leading to

a competition for gaze resources. For instance, at a dinner party, diners use their gaze to control

their manual actions, such as handling objects, while also using their gaze to survey their

surroundings and engage in conversations. Under such circumstances, one would expect that

gaze would only be directed to the action task when the support of gaze is most important.

To our knowledge, no prior research has delved into the control policies and strategies

governing gaze allocation when there are competing demands for gaze from manual actions and

environmental monitoring. To address this gap, we designed an experiment where participants

simultaneously performed an object manipulation task and a visual monitoring task that

required central vision. The manipulation task involved repeatedly grasping a small ball situated

on a platform and inserting it into a slot in a vertical tube, from which the ball returned to the

platform (Fig. 1A). To manipulate the visuomotor demands, participants were required to

perform the task using both their fingertips and tweezers. At the same time, participants

monitored a text display for randomly occurring letter changes. Successful ball drops were

rewarded, while failures to detect letter changes incurred penalties.

This study aimed to examine three novel hypotheses regarding how gaze allocation might be

optimized during the concurrent execution of an action task and a perception task that compete

for gaze. First, we hypothesized that the prevalence and function of gaze fixations directed to

the action task would depend on the specific task phase and the visuomotor demands, influenced

by the end-effector employed (fingertips and tweezers). Second, we hypothesized that, by

observing the temporal statistics of relevant visual events in the environment, individuals would

anticipate when events that require central vision will occur and use this information to allocate

gaze more efficiently between tasks. Third, we hypothesized that individuals would reduce

competition between tasks for gaze resources by adjusting the timing of the action task and,

consequently, the timing of required action task fixations. The rationale and motivation for these

hypotheses are elucidated below.

Object manipulation tasks involve a sequence of action phases delimited by kinematic or

mechanical events (Flanagan et al., 2006; Johansson et al., 2001). The action phases in our ball-

drop task are illustrated in Fig. 1B. When action tasks are performed in isolation, gaze is almost

exclusively directed to targets of action—the ball and slot in our ball-drop task—with the

function of gaze changing across action phases. When moving the hand, or object in hand,

towards the vicinity of a target—as in the reach and transport phases of our task—individuals

typically fixate the target. Such fixations can enable fast, automatic feedback control

mechanisms that use peripheral vision and gaze-related signals to *direct* the hand toward the

target (Goodale et al., 1986; Saunders and Knill, 2004, 2003). Once the hand gets close to the

object, and more slowly approaches it—as in the ball and slot approach phases in our task—

gaze may remain on the target, in which case central vision can be used to *guide* the hand, or

grasped object, to the target through more deliberate closed-loop feedback control (Ballard et

al., 1992; Johansson et al., 2001; Land, 2006). Once the hand or object in hand contacts the

target—ball contact and slot entry marking the start of the grasp and slots phases in our task—

tactile feedback becomes available. This tactile feedback provides confirmation that contact

was achieved. If gaze remains on the target until contact, central vision can also be used to *check*

that contact was achieved (Säfström et al., 2014). Within this framework of gaze functions in

isolated tasks, the current study addresses whether and how corresponding gaze functions are

expressed when there is competition for central vision.

We expected that when the ball-drop is performed concurrently with the visual monitoring task,

all fixations directed to the action task would be located at the ball and slot. Regarding our first

hypothesis, we predicted that when using the fingertips, action task fixations, if observed, would

primarily serve the purpose of *directing* the hand via peripheral vision. We predicted that

fixations involved in *guiding* the hand via central vision could be omitted when using fingertips

because, once the hand (or ball in hand) arrives at the target, haptic feedback can be used to

correct positioning errors. In contrast, we predicted that when using tweezers, gaze fixations, in

addition to being involved in *directing* the hand, would also be involved in *guiding* the hand, as

tweezers require greater spatial precision, particularly in grasping the ball, and offer limited

tactile feedback about the contact state. Note that impaired tactile sensibility is known to

increase reliance on visual feedback for object manipulation (Brink and Mackel, 1987;

Chemnitz et al., 2013; Jenmalm and Johansson, 1997; Jerosch-Herold, 1993).

Previous work has shown that when concurrently monitoring two locations to detect

probabilistic events, individuals can optimize their gaze allocation by learning the temporal

regularities of the events at each location and adjusting their gaze accordingly (Hoppe and

Rothkopf, 2016). This raises the question of whether people can similarly learn and exploit

temporal regularities of environmental visual events when concurrently performing a visual

monitoring task and a visually guided action task. In the monitoring task, letter changes

occurred at intervals randomly sampled from a uniform distribution between 1.5 to 6.5 seconds

(s). Thus, each letter change was followed by a 1.5 s 'silent period', during which no change

could occur. After this period, the probability of the next letter change (i.e., the hazard rate)

linearly increased. With respect to our second hypothesis, we predicted that participants would

exploit this temporal structure to primarily allocate their gaze to the manipulation task during

silent periods and shortly afterwards, when the probability of a letter change was low. In

contrast to visual monitoring of environmental events, where timing demands on central vision

are externally determined, individuals can, in principle, adjust the timing of their own actions

and, consequently, the timing of the required action task fixations. For our third hypothesis, we

predicted that individuals would mitigate competition for gaze resources between tasks by

strategically adjusting the timing of the action task. This hypothesis relies on the assumption

that participants not only learn the statistical properties of letter changes in the monitoring task,

but also possess knowledge of when and where fixations are required for the action task as it

unfolds.

Figure 1. Apparatus and action phases in the ball-drop task. (A) Experimental setup from the participant's perspective showing the path of the end-effector tip during a single tweezer trial. Movements were executed in a plane parallel to the participant's coronal plane, positioned 40 cm from the participant's eyes. (B) Corresponding velocity profile of the end-effector. (A, B) The task was divided into 7 consecutive action phases separated by distinct kinematic or contact events (see Methods). (1) *Reach phase*: Begins when the hand departs from its starting position and is characterized by a bell-shaped velocity profile. (2) *Ball approach phase*: Begins at a minimum (or inflexion point) in the velocity profile. (3) *Grasp phase*: Begins upon first contact with the ball. (4) *Transport phase*: Begins when the ball is lifted from the platform and characterized by a bell-shaped velocity profile. (5) *Slot approach phase*: Begins at a minimum (or inflexion point) in the speed profile and is characterized by slow movement. (6) *Slot phase*: Begins when the tips of the end-effector holding the ball enter the slot. (7) *Return phase*: Begins when the ball is released and exhibits a bell-shaped velocity profile. The trial concludes when the hand returns to its starting position.

Results

Eleven participants performed, at their own pace, 30 consecutive trials of the ball-drop task in each of four experimental conditions. Participants performed the task using either their fingertips or tweezers, either as a standalone activity (referred to as 'single task' conditions) or concurrently with the visual monitoring task (referred to as 'dual task' condition). In each ball-drop trial, participants reached for and grasped a ball positioned on a platform adjacent to the base of a vertical tube. They then transported the ball to one of three slots within the tube, inserted it, and released it before returning their hand to its starting position. After the ball was released, it descended through the tube and returned to its starting position on the platform. One second after the ball returned to the start position (or was already located in the start position in the first trial), an auditory signal instructed the participant about which slot to use.

Figure 1A shows the end-effector movement path during a tweezer trial, from the participant's perspective. The motion occurred in a plane parallel to the participant's coronal plane, situated in front of the body. Each ball-drop trial was divided into seven consecutive action phases, delimited by distinct kinematic or mechanical contact events (see Figure 1B). The visual monitoring task involved detecting changes in a letter displayed on a text screen positioned in the upper right quadrant of the participant's scene (Fig. 1A; for details see Methods). Participants received rewards for successful ball drops and were penalized for missing letter changes, which were signaled by an auditory tone and visual feedback on the display.

We first examine gaze and end-effector coordination in the single task conditions performed using either fingertips or tweezers. The conditions provide baselines for comparison with the corresponding dual task conditions.

Single task conditions

Figures 2 A-B show gaze and end-effector paths during representative trials with the fingertips and tweezers, respectively. In both fingertip and tweezer trials, participants typically fixated the ball while reaching for it, shifted their gaze to the slot around the time the ball was grasped, and maintained the fixation of the slot until around the time the ball was dropped. Gaze then returned to the ball's start position at the tube's base. Figures 2 C-D show for each end-effector, the average speed of the tip of the end-effector (top panels) and the instantaneous probabilities of gaze fixating the ball at its start position or the target slot, as a function of time (bottom panels). Note that these figures are based on aggregate data from all trials and participants. To temporally align trials while preserving phase-specific information, we normalized the duration of each phase in each trial to the median phase duration across all trials within each condition.

In both fingertip and tweezer trials, participants predominantly fixated the ball throughout the
reach phase, though gaze was occasionally located at the slot during the reach phase of fingertip
trials. Similarly, during the transport phase, participants primarily fixated the slot. The timing
of the gaze shift from the ball to the slot differed significantly between fingertip and tweezer
trials ($t_{10} = 13.24$; $p < 0.001$; $d = 3.99$). In fingertip trials, the shift occurred just before contact
(-0.06 ± 0.05 s; mean ± sem), typically during either the late reach phase or the ball-approach
phase. However, in tweezer trials, the shift occurred well after ball contact (0.29 ± 0.11 s),
mainly during the grasp phase. This finding supports the prediction that tweezers require greater
reliance on central vision to establish stable grasp on the ball.

Figure 2. Gaze-hand coordination in the single-task condition. (A, B) Gaze and end-effector paths in single fingertip (A) and tweezer (B) trials. Fixations are color-coded according to landmark (ball, slot), and end-effector paths (tip of the fingers or tweezers) are color-coded based on the current gaze state: fixating the ball, fixating the slot, or executing a saccade. Note that the ball was fixated during the reach and return phases while the slot was fixated during the transport phase. (C, D) Average end-effector speed (black) and the probabilities of fixating the ball (orange) and slot (green) as a function of time relative to lift-off, for fingertip (C) and tweezer (D) trials. Separate traces are shown for each slot (top, middle, and bottom coded dark to light). Alternating white, brown, and grey-blue regions indicate movement phases labelled in D. Plots aggregate data from all trials conducted by all participants, with phase durations in each trial normalized to the median duration of the phase. Fixations were almost always directed to the landmarks (ball and slot); however, the combined probabilities of these fixations may be less than 1 due to saccades occurring between the landmarks.

Dual task conditions

In the dual task conditions, participants divided their gaze fixations between the text display

and the action-related landmarks, consistent with the fact that detecting letter changes in the

visual monitoring task required central vision (Fig. 3A-B). As a result, the likelihood of fixating

the ball or slot during the ball-drop task was reduced compared to the single task conditions,

regardless of end-effector used (Fig. 3C–D). Additionally, the durations of occurring action

landmark fixations were consistently shorter.

Propensity and patterns of action fixations depend on the end-effector

In fingertip trials, participants primarily fixated the display during the reach, ball approach, and

grasp phases (88% of trials), suggesting that grasping the ball generally did not require central

vision. In trials where participants did fixate the ball, these fixations occurred during the reach

phase, with gaze typically shifting away before the ball approach phase and almost never

remaining on the ball after contact. Transporting the ball and inserting it into the slot were also

frequently performed with gaze remaining on the display. However, participants briefly fixated

the slot in about half of the trials (51%), as in the example shown in Fig. 3A. The probability of

fixating the slot peaked midway through the transport phase, then steadily declined,

approaching zero by the end of the slot phase (Fig. 3C). These findings show that in fingertip

trials, inserting the ball into the slot often did not require central vision. However, the likelihood

of fixating the slot during the slot approach and slot phase was higher than fixating the ball

during the ball approach and grasp phase.

Consistent with the fixation probabilities described above, we observed two predominant gaze

patterns in fingertip trials: 'display-only', where gaze remained on the display throughout the

trial, and 'slot', where gaze shifted from the display to the slot and then back to the display (see

Figs. 3E-F). To examine the relationship between gaze pattern and manual performance, we

compared the action phase durations in display-only and slot trials. Only participants (N=9)

who demonstrated both fixation patterns were included. The duration of the slot phase was

shorter when participants fixated the slot (0.30 ± .054 s) compared to when they did not (0.378

± .056 s; t_8 = 3.76, Holm-Bonferroni adjusted p = 0.03). No other phase durations were affected

by gaze pattern (adjusted p > 0.51 in all cases).

In tweezer trials, participants were much more likely to fixate both the ball and the slot

compared to fingertip trials (Fig. 3D). Participants almost always fixated the ball before contact

(88% of trials), and the slot before slot entry (89% of trials). The probability of fixating the ball

peaked towards the end of the reach phase and remained relatively high during the ball approach

and most of the grasp phase. Similarly, the likelihood of fixating the slot peaked towards the
end of the transport phase and remained elevated during the slot approach and slot phases. These
findings suggest that, in general in tweezer trials, central vision was crucial for both grasping
the ball and inserting it into the slot.

Figure 3. Gaze-hand coordination during dual-task conditions. (A, B) Gaze and end-effector paths in single fingertip (A) and tweezer (B) trials. Fixations are color-coded by landmark (ball, slot, or display), and end-effector paths (tip of fingers or tweezers) are color-coded based on the current gaze state (fixating the ball, slot or display, or executing a saccade). (C, D) Average end-effector speed (black) and the probabilities of fixating the ball (orange), slot (green), and display (blue) as a function of time relative to lift-off, for fingertip (C) and tweezer (D) trials. Separate traces are shown for each slot (top to bottom coded dark to light). Alternating white, brown, and grey-blue regions indicate movement phases labelled in D. Plots combine all trials from all participants with phase durations in each trial normalized to the median duration of the phase. (E) The five distinct gaze patterns observed during dual-task trials are illustrated with data from single trials. (F) Average percentage of each gaze pattern (color-coded as in E) observed across participants in fingertip and tweezer trials. Individual participants are represented by dots.

The predominant gaze pattern in tweezer trials was 'ball-display-slot', where participants shifted
their gaze from the display to the ball, back to the display, then to the slot and finally returned
to the display (see example in Fig. 3B). The second most common pattern was 'ball-slot', where
participants shifted their gaze from the display to the ball and then directly to the slot before
returning to the display (see Figs. 3E-F). To examine the relationship between gaze pattern and
manual performance, we compared the action phase durations between the ball-display-slot and
ball-slot trials. Only participants who exhibited both gaze patterns ($N = 8$) were included in this
analysis. We found that the transport phase was shorter when gaze shifted directly from the ball
to the slot ($0.32 \pm .051$ s) compared to when gaze fixated the display between the ball and slot
fixations (0.55 ± 0.17 s; $t_7 = 4.71$, Holms-Bonferroni adjusted $p = 0.01$). No other phase
durations were associated with gaze pattern (Holms-Bonferroni adjusted $p > 0.22$ in all cases).

As in the single-task conditions, in the dual task conditions the ball-drop task was performed
more slowly with the tweezers (2.77 ± 0.4 s) than with the fingertips (2.11 ± 0.23 s; $t_{10} = 5.61$,
$p < 0.001$; $d = 1.69$). Similar to the single task conditions, the greater time taken to perform the
task with tweezers was due to increased durations of the reach, grasp, and transport phases
(Holms-Bonferroni adjusted $p < .01$ in all three cases).

Action fixations are temporally anchored to contact events

To examine the trial-by-trial coordination between the timing of action fixations and specific
kinematic or contact events, we conducted a multiple linear regression analysis to assess how
well the onset and offset times of ball and slot fixations could be predicted by the timing of the
following six events delimiting action phases: (1) start of the reach phase, (2) start of the ball
approach phase, (3) time of first ball contact (start of the ball grasp phase), (4) time of ball liftoff
(start of the ball transport phase), (5) start of the slot approach phase and (6) time for slot entry
(start of the slot phase). To mitigate structural multicollinearity, predictors were centered
individually for each participant by subtracting their mean. Furthermore, participants were
included as a categorical nuisance factor to account for variance due to differences in task
performance speed. Separate regression analyses were conducted for fixation onsets and offsets,
as well as for each action landmark (ball and slot) and end-effector (fingertips and tweezers).

In both fingertip and tweezer trials, the initiation and termination of ball fixations were best
predicted by the time of first ball contact, while the initiation and termination of slot fixations
were best predicted by the time of slot entry. For fixation onset times, in fingertip trials, ball
fixation onset was predicted solely by the time of first ball contact ($t_{1,32} = 2.84$; $p = 0.008$), and

slot fixation onset solely by the time of slot entry ($t_{1,158} = 9.28$; $p < 0.001$). Similarly, in tweezer trials, the ball fixation onset was primarily predicted by the time of first ball contact ($t_{1,243} = 6.26$; $p < 0.001$), while slot fixation onset was solely predicted by the time of slot entry ($t_{1,244} = 11.2$; $p < 0.001$). Comparable patterns were observed for fixation offset times. In fingertip trials, ball fixation offset was solely predicted by the time of first ball contact ($t_{1,32} = 3.08$; $p = 0.004$), while slot fixation offset was best predicted by the time of slot entry ($t_{1,158} = 8.30$; $p < 0.001$). Similarly, in tweezer trials, ball fixation offset was best predicted by the time of first ball contact ($t_{1,243} = 5.18$; $p < 0.001$), and slot fixation offset was solely predicted by the time of slot entry ($t_{1,244} = 9.04$; $p < 0.001$). These findings demonstrate a robust temporal coupling between the initiation and termination of action landmark fixations and their associated contact events, consistent across both end-effectors.

Figure 4 illustrates this temporal relationship, showing that the timing of ball and slot fixation periods relative to ball contact and slot entry, respectively, remained quite consistent across trials. The left panels of Figs. 4A and B show the timing of ball fixations and associated action phases relative to the time of first ball contact (time = 0). Each row represents an individual trial and data from all participants are shown. Orange lines indicate periods of ball fixation, while dots mark the onset of the reach, ball approach and transport phases in each trial. The trials are arranged in order of ball approach phase duration. Similarly, the left panels of Figs. 4C and D, show the timing of slot fixation and associated action phases relative to the time of slot entry. Green lines represent periods of slot fixation, and dots mark the onset of the transport, slot approach, and return phases for each trial. These trials are sorted by slot approach phase duration. In some trials, the ball or slot were re-fixated (depicted by purple lines in Figs. 4A-D). Re-fixartions typically occurred when multiple attempts were required to successfully grasp and lift the ball or insert it into the slot. Notably, the durations of both the ball and slot approach phases could vary considerably across trials.

The scatter plots on the right side of each panel in Fig. 4 illustrates the position of each trial in its test block (No. 1 – 30), ranked by the duration of the ball or slot approach phase, as well as the position of each participant's trials in the same ranking (Pt. Nos. 1 -11). Colored dots (orange for ball and green for slot) represent trials with an action landmark fixation, while black dots indicate trials without such fixations. The absence of apparent structure in these scatter plots suggests that neither the decision to fixate the action landmark nor the variation in ball and slot approach phase durations were influenced by trial position. Furthermore, these scatter plots indicate that, overall, participants exhibited similar behaviour.

Figure 4. Timing of action fixations. (A-D) These panels show timing data for all trials stacked vertically. (A, B) The left column shows periods of initial (orange lines) and secondary (purple lines) ball fixations, aligned to the time of initial ball contact (grey vertical line), in fingertip (A) and tweezer (B) trials. For each trial, the onsets of the reach and transport phases are marked by small black dots, and the onset of the ball approach phase is marked by larger black dots. Trials are sorted by ball approach phase duration. The middle and right columns show trial number and participant number, with orange and black dots representing trials with and without a ball fixation, respectively. (C, D) Corresponding plots for initial (green) and secondary (purple) slot fixations, aligned to the time of slot contact. Small black dots mark the onsets of the transport and return phases, and the larger black dots indicate the onset of the slot approach phase. Data are sorted by the slot approach phase duration. In the middle and right columns, the green and black dots depict trials with and without a slot fixation. (E) Cumulative distributions of ball and slot fixation onsets and offsets, aligned to the initial ball contact and slot entry, respectively, in fingertip trials. (F) Corresponding distributions in tweezer trials.

The timing of action fixation onsets, relative to contact events, showed remarkable consistency

across action landmarks and end-effectors. Both ball and slot fixations typically began

approximately 0.4 s before ball contact and slot entry, respectively (solid line curves in Figs. 4E

and F). The timing of fixation offsets was also consistent across action landmarks but influenced

by the end effector used. In fingertip trials, gaze generally shifted away from both the ball and

the slot before the contact event, with an average lead time of about 0.15 s (dashed line curves

in Fig. 4E). In contrast, in tweezer trials, gaze typically shifted shortly after the contact event,

with an average lag of about 0.05 s (dashed line curves in Fig. 4F).

Variability in the function of action fixations across trials

The variability in the timing of ball and slot fixations, relative to their associated action phases,

suggests that the functions of these fixations—such as *directing* and *guiding*—may vary across

trials. To investigate this possibility, we analyzed the function of each individual ball and slot

fixation, acknowledging that an individual fixation could serve multiple functions. A fixation

was classified as *directing* if the ball or slot was fixated for at least 100 ms during the reach or

transport phase, respectively. Similarly, a fixation was classified as *guiding* if the ball or slot

was fixated for at least 100 ms between the start of the ball or slot approach phase and the end

of the grasp or slot phase, respectively (i.e., the combination of the approach and manipulation

phases). Additionally, fixations were considered as checking when the ball or slot was fixated

for any period of time after the end of the grasp or slot phase, respectively, allowing visual

confirmation of the action's completion (Säfström et al., 2014). Figure 5A illustrates examples

of slot fixations in tweezer trials demonstrating these distinct functions.

In fingertip trials, ball fixations primarily served a *directing* function, whereas slot fixations

also frequently played a *guiding* role (Fig. 5B). In contrast, in tweezer trials, both ball and slot

fixations were more evenly distributed between *directing* and *guiding* (Fig. 5C). In both

conditions, a small proportion of fixations were involved in *checking* (Fig. 5B and C).

Interestingly, in fingertip trials, most of the ball and slot fixations served only one function (80%

overall), whereas in tweezer trials, this proportion was lower (48% overall). This distinction is

highlighted by the thin solid bars within each wide bar in Figs. 5B and C.

Figure 5. Classification of fixation functions. (A) Examples of slot fixations from tweezer trials that serve different functions. (B, C) Wide bars show the mean percentage of ball (orange) and slot (green) fixations engaged in *directing*, *guiding*, and *checking* across participants in fingertip (B) and tweezer (C) trials. Note that a given fixation could be engaged in more than one function. Circles represent individual participants, except for the circles at zero which represent all participants with zero. The thin bars represent percentages of single-function fixations.

Impact of letter change statistics on the allocation of gaze resources

In the following section, we investigate whether participants can learn and exploit the statistical
properties of letter changes (LCs) to mitigate competition for gaze between the visual
monitoring and manipulation tasks. Our findings reveal that participants adapt their gaze
behaviour both directly, by selecting different gaze patterns, and indirectly, by modifying their
manual actions in response to LC statistics (see next section).

In the dual-task condition, the interval between LCs was drawn from a uniform distribution
spanning 1.5 to 6 s. As a consequence, there was a 1.5 s "silent period" following a LC when
the next LC could not occur. Following the silent period, the likelihood of the next LC occurring
(i.e., the hazard rate) gradually increased over the subsequent 5 s rate. On average, there were
1.08 and 1.41 LCs per trial in fingertip and tweezer trials, respectively. Importantly, participants
were very good at detecting these LCs and therefore could, in principle, exploit the LC statistics.
Participants detected these LCs with an accuracy of $88.8 \pm 11.8\%$ and $87.1 \pm 9.1\%$, respectively
(mean ± standard deviation across participants).

To analyze whether LC detection influenced the timing action task fixations, we compared the
observed frequency distributions of ball and slot fixation onsets—relative to the most recently
detected LC preceding each fixation—with the distributions expected if fixation onsets occurred
randomly with respect to LCs. For each end-effector, we conducted separate analyses for trials
exhibiting each of the primary gaze patterns: ball-display-slot, ball-slot, and slot trials. In both

fingertip and tweezer trials in which participants fixated both the ball and slot (i.e., ball-slot and

ball-display-slot trials), the distribution of ball fixations onsets deviated from the expected

random distribution (Kolmogorov-Smirnov test, $p \leq 0.01$ in all four cases). Specifically, ball

fixation onsets during the silent period occurred more frequently than expected by chance (top

panels of Fig. 6A).

In tweezer trials, the choice of gaze pattern was strongly influenced by the timing of ball fixation

onset relative to the preceding LC. When the ball was fixated during the silent period,

participants were equally likely to adopt either the ball-slot or ball-display-slot pattern.

However, if the ball was fixated after the silent period, the ball-display-slot pattern was

predominantly chosen. That is, participants rarely shifted their gaze directly from the ball to the

slot (ball-slot pattern) unless the ball was fixated within the silent period. Conversely, when ball

fixation took place after the silent period, participants almost always fixated the display before

fixating the slot (ball-display-slot pattern).

As expected, in both fingertip and tweezer trials where gaze shifted directly from the ball to the

slot (ball-slot pattern), a peak in the distribution of slot fixation onsets followed the peak in the

distribution of ball fixation onsets in the silent period (bottom panels of Fig. 6A). This occurred

because, in the vast majority of these trials (96.4%), the last LC detected before the ball fixation

was also the last LC before the slot fixation. Both of these distributions differed from the

expected random distributions (KS test, $p < 0.02$ in both cases). Similarly, in fingertip and

tweezer trials with the ball-display-slot pattern, there is a peak the distribution of slot fixation

onsets that is shifted to the right of the peak in the distribution of ball fixations, also representing

trials in which last LC detected before the ball fixation was also the last LC before the slot

fixation. In fingertip trials with the ball-display-slot pattern, the distribution of slot fixations

differed from the expected random distribution (KS test, $p < 0.01$). Note that in tweezer trials

with the ball-display-slot pattern, there is also an earlier peak in the distribution of slot fixations

occurring during the silent period, representing trials where a LC was detected while gaze

fixated the display prior to shifting to the slot. As a consequence of this earlier peak, the

distribution of slot fixations did not differ from the expected random distribution (KS test, $p = $

0.4). In both fingertip and tweezer trials, where only the slot was fixated (slot gaze pattern), the

distribution of slot fixation onsets did not differ significantly from the expected random

distribution (KS test, $p > 0.06$ in both cases; bottom panels of Fig. 6A).

Together, these results demonstrate that participants could adapt their gaze behaviour to exploit

the statistical properties of LCs to mitigate competition for gaze resources between the visual

monitoring and object manipulation tasks.

Figure 6. Impact of timing of letter changes (LCs) on gaze fixations and hand movements. (A) Frequency distributions, combining trials from all participants, of ball (top) and slot (bottom) fixation onsets—relative to the time of the last detected LC before the fixation onset—in fingertip (left) and tweezer (right) trials. Yellow regions indicate the silent period. Separate distributions are shown for trials with the main gaze patterns. Dashed lines show expected distributions assuming fixation onsets occurred randomly and independently LC timing. The expected frequency is constant within the silent period, during which the hazard rate (the probability of a LC occurring if one has not yet occurred) remains at 0, and then decreases, at a constant rate, over the next 5 s as the hazard rate increases from 0 to 1. (B) Corresponding distributions of reach start times relative to the last detected antecedent LC in fingertip (left) and tweezer (right) trials. Separate plots are shown for the four main gaze patterns. (C) The top row shows, for fingertip and tweezer trials, the relationship between ball fixation onset and ball contact times, relative to the last detected LC before fixation onset. Dots represent trials from all participants and are colour-coded by gaze pattern. The bottom row shows the corresponding relationships between slot fixation onset and slot entry times. (D) Relationship between reach start time and time of nearest detected LC, both relative to cue onset, in fingertip and tweezer trials with ball-slot and ball-display-slot patterns. (E) Frequency distributions of the reach start times in trials with ball-slot and ball-display-slot gaze patterns, aligned with the data shown in (D). These distributions combine fingertip and tweezer trials. Separate distributions are shown for trials in which the LC was within or outside the 'ready period' (1 s period prior to the cue).

Impact of letter change statistics on reach timing

We found that the LC statistics influenced manual performance in the ball-drop task, particularly the timing of reach initiation (Fig. 6B). In both fingertip and tweezer trials where participants fixated the ball (i.e., the ball-slot and ball-display-slot gaze patterns), the distribution of reach onset times relative to the antecedent detected LC differed from the random distribution (K-S test, $p < 0.05$ in all four cases). Specifically, reach onsets were biased towards the silent period (top two rows of Fig. 6B), mirroring the bias observed in ball fixations. In contrast, in fingertip trials where the ball was not fixated (i.e., the slot and display-only gaze patterns), the distribution of reach onset times did not differ from the random distribution (K-S test, $p \geq 0.4$; bottom two rows of Fig. 6B). Because of the limited number tweezer trials with the slot and display-only gaze patterns, we did not test whether the distributions of reach onsets in these trials differed from the random distribution. These findings suggest that initiating the reach during the silent period was a strategy to reduce the risk of missing LCs.

Importantly, we found that there was a robust temporal coupling between action landmark fixations and contact events, irrespective of the gaze pattern that participant selected. Thus, for all gaze patterns involving a ball fixation, and across a wide range of ball fixation onset times relative to the last detected LC, we observed strong temporal coupling between ball fixation onset and ball contact (top row of Fig. 6C). Similarly, across all gaze patterns involving a slot fixation, we observed a strong temporal coupling between slot fixation onset and slot entry (bottom row of Fig. 6C).

The intercept and slope of the relationship between ball fixation onset and ball contact time were 0.43 s and 1.00 in fingertip trials and 0.41 s and 0.97 in tweezer trials. The intercept and slope of the relationship between slot fixation onset and slot entry time were 0.368 s and 1.00 in fingertip trials and 0.38 s and 0.99 in tweezer trials. These intercepts values are consistent with our earlier observation that both ball and slot fixations typically began approximately 0.4 s before the corresponding contact event, irrespective of the end-effector used (Figs. 4E and F).

In the ball drop task, participants typically initiated their reach movement towards the ball in response to the auditory cue signaling the active slot for that trial. These 'reactive reaches' generally began about 0.5 s after the cue. However, in a notable proportion of trials, participants initiated their reach in anticipation of the cue, starting either before or shortly after the cue (within 0.5 s). If participants generated these 'anticipatory reaches' in response to a LC occurring shortly before the cue, and fixated the ball when doing so, it could contribute to the greater-than-expected frequency of both reach onsets and ball fixation onsets during the silent period in trials with the ball-slot and ball-display-slot gaze patterns.

To examine this further, for fingertip and tweezer trials involving ball fixation, we analyzed the

relationship between the timing of reach onset relative to the cue and the timing of the detected

LC relative to the cue. For each trial, we identified the detected LC closest in time to the

midpoint of the 'ready period', defined as the 1-second interval between the ball returning to its

starting position and the auditory cue. This LC could occur either before or after the midpoint

of the ready period. We observed that most anticipatory reaches—characterized by reach onset

times near or preceding the cue—occurred when the detected LC happened within the ready

period. This relationship is evident in the left two panels of Fig. 6D, which show reach onset

times relative to the cue plotted against the timing of the detected LC. These findings suggest

that the decision to initiate an anticipatory reach is closely linked to the detection of a LC during

the ready period.

The frequency distributions of reach onset times relative to the cue, shown in the right two

panels of Fig. 6D for trials with the ball-slot and ball-display-slot patterns, revealed earlier reach

onset times when the LC occurred within the ready period compared to when it occurred outside

of it. Due to the relatively small number of fingertip trials, fingertip and tweezer trials were

combined for these analyses. Importantly, for both gaze patterns, the distributions within and

outside the ready period differed (KS test, $p < 0.02$ in both cases). This suggests that the timing

of detected LCs had a distinct influence on reach initiation during the ball drop task.

Overall, these results demonstrate two strategies that participants used to exploit the statistical

properties of LCs to mitigate competition for gaze resources between the tasks. First, they

adjusted the timing of their reaching movements so that the ball could be preferentially fixated

during the silent period. Second, they adapted their gaze patterns on a trial-by-trial basis to

increase the likelihood that gaze could be allocated to the action task with minimal disruption

of the LC monitoring task.

Discussion

This study investigated three hypotheses about how individuals allocate gaze during object

manipulation while also monitoring the environment. These hypotheses concerned: (1) how

often, when, and where fixations related to the action task occur when performing the task with

end-effectors with different visuomotor demands; (2) whether participants learn and exploit the

temporal statistics of the monitoring task to guide gaze allocation; and (3) whether they adjust

manual action timing based on these statistics to reduce competition for gaze resources between

the tasks. Our findings supported all three hypotheses, as detailed below.

Gaze allocation to the ball-drop task: Effects of end-effector

As expected, when the ball-drop task was performed in isolation using either the fingertips or

tweezers, gaze was directed exclusively to the ball and slot, arriving ahead of the end-effector

and departing around the time of its arrival or shortly afterward. This finding aligns with prior

research on eye-hand coordination in visually guided tasks (Flanagan and Johansson, 2003;

Fooken et al., 2021; Hayhoe, 2017; Illamperuma and Fooken, 2024; Johansson et al., 2001;

Land et al., 1999). However, when participants performed the ball-drop alongside the

monitoring task, we expected gaze to be allocated to the action task when visuomotor control

was most critical. Consistent with this expectation, we found that while the ball and slot were

almost always fixated in tweezer trials, ball fixations—and, to a lesser extent, slot fixations—

were frequently absent in fingertip trials. The increased reliance on central vision with tweezers

is likely due to several factors. First, the smaller contact surfaces of the tweezer tips require

greater spatial precision, particularly during ball grasping. Second, unlike the more malleable

fingertips, the rigid tweezer tips do not conform to the ball's surface, reducing grasp stability.

Note that in robotic manipulation, soft contact surfaces are used to enhance stability and reduce

spatial precision demands (Bicchi, 2000; Bicchi and Kumar, 2002; Billard and Kragic, 2019).

Third, the tweezers provide limited tactile feedback about the contact state, and impaired tactile

sensibility is known to increase reliance on vision during object manipulation (Brink and

Mackel, 1987; Chemnitz et al., 2013; Jenmalm and Johansson, 1997; Jerosch-Herold, 1993).

These findings suggest that analyzing gaze behaviour during object manipulation tasks can offer

insights into tactile impairments and the effectiveness of tools in transmitting tactile information

to the user, an important consideration for applications such as teleoperation and robot assisted

surgery.

Timing and role of action task fixations across end-effectors

Independent of the end-effector employed, the timing of ball and slot fixation onsets and offsets

were most closely correlated with ball contact and slot entry, respectively, compared to other

events in the ball-drop task. For both action task landmarks (ball and slot), and both end-

effectors, gaze typically arrived around 0.4 s prior to contact, a pattern consistent with

visuomotor coordination in natural action tasks (Ballard et al., 1995; Keshava et al., 2024; Land

and Hayhoe, 2001). However, as hypothesized, the timing of gaze shifts differed based on the

end-effector. In fingertip trials, gaze shifted away from the ball and slot well ahead of contact

(0.15 s on average), whereas in tweezer trials, gaze shifted just after contact (0.05 s on average).

This difference supports our hypothesis that the role of action task fixations varies depending

on the end effector. Specifically, when using the fingertips, ball and slot fixations primarily

direct the hand (or object in hand) toward the target landmark using peripheral vision and gaze position-related signals (de Brouwer et al., 2018; Prablanc et al., 1986; Wilmut et al., 2006). In contrast, when using tweezers, the delayed gaze shift away from the target suggests that these fixations also assist in *guiding* the end effector with central vision when close to the target. Once the fingertips or tweezer tips contact the ball or slot, tactile feedback becomes available, facilitating a transition from visuomotor control to haptic sensorimotor control. This transition aligns with the shift from motion control to force control, which is thought to involve distinct control processes (Casadio et al., 2015; Chib et al., 2009; Kolesnikov et al., 2011; Piovesan et al., 2019). In the ball drop task, tactile information can not only guide force adjustments, as when grasping the ball, but can also drive kinematic corrections, as when repositioning the ball when inserting it into the slot. In manipulation tasks, tactile information enables rapid adjustments (90-120 ms) to both force (Johansson and Flanagan, 2009) and kinematics (Pruszynski et al., 2018, 2016) via automatic feedback control mechanisms.

Gaze patterns and task performance across end-effectors

We found that across both fingertip and tweezer trials, participants used different gaze patterns when performing the ball drop task alongside the monitoring task. In fingertip trials, two main gaze patterns were observed: "display only" and "slot", distinguished by whether the participant fixated the slot or kept their gaze on the display throughout the trial. In tweezer trials, two main patterns were also observed: "ball-slot" and "ball-display-slot", distinguished by whether the participant decided to fixate the display between the ball and slot fixations. Importantly, the choice of gaze patterns was linked to task performance. In fingertip trials, the duration of the slot phase was shorter when participants fixated the slot, and in tweezer trials, the duration of transport phase was shorter when participants shifted their gaze directly from the ball to the slot, bypassing the display. These results suggest a trade-off between the action and visual monitoring tasks; allocating gaze to the action task improves performance but increases the risk of missing a letter change (LC). However, as we will discuss next, participants can mitigate this trade-off by leveraging the LC statistics to guide gaze pattern selection and execution of the manipulation task.

Exploiting temporal statistics of the monitoring task for gaze allocation

Our findings support the hypothesis that participants exploited the temporal statistics of LCs in the visual monitoring task to guide gaze allocation decisions. In fingertip trials, the decision to fixate the ball was strongly influenced by the LC statistics, with almost all observed ball fixations occurring during the silent period when no LC could occur. Similarly, in tweezer trials, the choice to fixate the display between fixating the ball and slot was also influenced by LC

statistics. Specifically, participants often skipped the display when the ball fixation occurred within the silent period, but almost always fixated the display when the ball fixation occurred outside this period. These findings align with previous research demonstrating that human gaze behaviour is sensitive to probabilistic regularities in the environment (Jovancevic-Misic and Hayhoe, 2009). For example, individuals adjust the timing of gaze shifts based on learned temporal statistics of relevant visual events to optimize detection when events occur at different intervals across two spatial locations (Hoppe and Rothkopf, 2016). In addition, during visual search tasks, gaze is allocated strategically based on spatial statistics to optimize exploration (Eckstein, 2017; Hoppe and Rothkopf, 2019; Najemnik and Geisler, 2005; Renninger et al., 2007). Our study adds a novel perspective on how environmental statistics influence gaze behaviour by demonstrating that humans can learn and exploit the temporal patterns of externally determined events in the visual environment while concurrently performing an action task that relies on visual guidance. This result suggests that the processes involved in using visual information in sensorimotor control—including peripheral and central vision and gaze-related signals—are largely independent of those used to extract and use statistical regularities from the visual environment.

Timing of manual actions to mitigate gaze competition

Unlike visual monitoring tasks where the timing of relevant events is often externally determined, in manual action tasks, the timing of events that make demands on central vision can generally be controlled or adjusted by the actor. We hypothesized that participants would adjust the timing of their manual actions to reduce competition for gaze resources between the ball drop and visual monitoring tasks. Supporting this hypothesis, we found that participants adjusted their reach onset times so that ball fixations—supporting the reaching movement and ball grasping—occurred during the silent period more often than expected if reach timing was independent of LC occurrences. Importantly, this indicates that participants not only learned the statistical properties of LCs but exploited predictions of when and where action task fixations are required during the unfolding task.

We found that in tweezer trials, participants often initiated their reach movement in anticipation of the cue when they observed a LC during the ready period (i.e., the period from when the ball returned to its start position to the auditory cue). This allowed them to both preferentially initiate their reach movement, and fixate the ball, during the silent period. In contrast, in fingertip trials in which the ball was rarely fixated, participants generally did not generate anticipatory reaches, indicating that anticipatory reaches are linked to fixating the ball, and are not simply a consequence of seeing a LC during the ready period. Indeed, in the few fingertip trials in which

participants did fixate the ball, they tended to generate anticipatory reaches, allowing them to
fix the ball "for free" during the silent period.

In fingertip trials, participants almost always maintained fixation at the display while reaching
towards and grasping the ball. Importantly, even though they were not fixating the target of the
action (i.e., the ball), we know from previous work that they could still use peripheral vision to
direct the hand towards the action target; i.e., the ball (de Brouwer et al., 2018; Neggers and
Bekkering, 2001, 2000). The question arises as to whether the use of peripheral vision for
controlling action is affected by the concurrent use of central vision to monitor external
environmental events. Our results suggest that these two visual functions may be independent.
If the processing demands associated with visual monitoring are reduced during the silent
period, and if visual monitoring interferes with the use of peripheral vision for action, we might
have expected participants to preferentially initiate their reaches during the silent period, which
they did not.

Overall, our results suggest that participants learned LC statistics to strategically adjust the
timing of reaching movements to mitigate potential conflicts between the monitoring and object
manipulation tasks, thereby supporting both tasks.

Conclusions

This study offers novel insights into the control and coordination of eye and hand movements
in real-world action tasks. First, our findings support the hypothesis that when gaze competition
arises, participants prioritize key control points—specifically, contact events between the hand,
or tool in hand, and target object in the environment—when allocating gaze during action tasks.
Second, our results demonstrate that participants exploit temporal regularities in the external
environment to enhance task performance by adapting both hand and eye movements.

Methods

Participants

501 Eleven right-handed participants (8 male; aged 22 to 33 yr) took part in the study. All
502 participants reported normal or corrected-to-normal vision and were naive to the purpose of the
503 study. The study was approved by the ethics committee of the University of Umeå and
504 participants gave written informed consent before participating in the study.

Apparatus and general procedure

Participants were seated at a table with the ball-drop apparatus positioned in front of them (Fig.

1A). It consisted of a 15 cm high vertically oriented Perspex tube (inner diameter = 14 mm;

wall thickness = 3 mm) mounted on a wooden platform. The tube was positioned about 2.5 cm

to the left of the participant's mid-sagittal plane, with its top aligned at eye level. The tube

contained three slots, the centers of which were positioned at heights of 5, 8, and 11 cm above

the platform surface.

Participants repeatedly performed the ball-drop task which required them to reach for and grasp

a small ball (12 mm diameter polished brass sphere) from a designated start position on the

platform, transport it towards the target slot, insert it into the tube and release it, and then return

the hand to a support plate located 20 cm to the right of the platform. A pre-recorded verbal

instruction ("bottom", "middle", or "top"), indicating which slot to use, was played 1 s after the

ball had rolled back to its start position. The ball's start position was 3 cm to the right of the

tube's vertical midline, and all movements occurred within a frontal plane at 40 cm distance

from the participants' eyes. The platform surface was slightly lower at the ball start position,

ensuring that the ball rolled back to its start position when exiting the tube. In different blocks

of trials, the ball was grasped either with the fingertips or with a pair of tweezers held by the

right hand. The tweezers, made of plastic, were 14 cm long with cylindrical tips (4 mm

diameter) coated for 12 mm with thin plastic tubing to increase the friction against the ball.

In the single task conditions, participants performed the ball-drop task only. In the dual task

conditions, participants completed a visual monitoring task that involved detecting letter

changes (LCs) on an LED text display, which required central vision. The text display was

positioned in the upper right quadrant of the scene (Fig. 1 C and D), 7 cm behind the frontal

plane in which the tube was located. The visual angle between the center of the letter area and

the center of the top slot and ball were 24° and 28°, respectively. The letter height and width

corresponded to 0.5° × 0.7° visual angle. At randomly distributed intervals (1.5—6.5 s, uniform

distribution), the letter M was changed to W for 300 ms before reverting to M. The participants

were instructed to press a button held in their left hand as soon as they detected a LC. If they

failed to respond within 1 second after a change, the LC was considered missed. Missed

detections were signaled by a brief beep and flashing hash marks on the display for 600 ms. To

ensure that the participants relied on central vision to detect LCs rather than peripheral visual

cues, the letter M randomly shifted its horizontal position by 0.6° visual angle at intervals

between 1 and 3 s (uniform distribution). Pilot tests confirmed that participants had to foveate

the display to detect LCs.

To encourage engagement with both tasks, participants earned 1 Swedish krona (SEK) for each ball-drop and lost 3 SEK for each missed LC. Participants were informed of these reward and penalty rules before testing.

Participants completed four conditions in the following order: single-task with fingertips, single-task with tweezers, dual-task with fingertips, and dual-task with tweezers. Each condition included 30 trials (10 per target slot). The target slot order varied unpredictably across trials.

Note that each participant completed three versions of the experiment that differed in terms of how the target slot was cued. In the version reported in this paper, the slot was cued with an auditory cue. In the visual cue version, the slots were indicated by illuminating LEDs located on the tubes, and in the 'memory' version, the participants were instructed to follow a specific target slot sequence that they had to remember. The order in which these three versions were completed was counterbalanced across participants. Importantly, we observed no clear differences in manual or gaze behaviour across these experiments and therefore decided to focus on one of them.

Data collection and analysis

Gaze position was recorded at 120 samples/s using an infrared video-based eye-tracking system (RK-726PCI pupil/corneal tracking system, ISCAN Inc., Burlington, MA). Head stability was maintained with an adjustable chin support and a forehead support to which the head was secured with Velcro straps. The standard deviations of gaze position measurement errors were 0.50° (horizontal) and 0.52° (vertical) of visual angle, corresponding to 0.35 cm and 0.36 cm in the vertical plane—parallel to the participant's coronal plane—in which the ball was moved during the task.

The position of the right index finger and the tweezers tips was tracked at 60 samples/s using miniature electromagnetic position-angle sensors with six degrees of freedom (RX1-D miniature receiver; FASTRAK, Polhemus, Colchester, VT). The fingertip sensor, attached to the nail, recorded the preferred contact point with the ball. The mapping between the position and orientation of the sensor and the preferred contact points was established during calibration trials in which the participants grasped the stationary ball at a known position. The position of the tweezer sensor, attached to the proximal end of the tweezers, was offset to the midpoint between the tweezer tips. When the ball was at its start position, it rested on a rectangular 'force plate' that formed part of the platform surface. A six-axis force-torque transducer (Nano F/T transducer, ATI Industrial Automation, Apex, NC; sampling rate 400 Hz), mounted under the

plate, detected when the impacted on the platform after being dropped through the tube, when the ball was first contacted by the fingertips or tweezers, and when the ball lifted off the platform. An optical reflex detector (SG-2BC, Kodenshi, Japan), mounted in a small hole beneath the ball's start position in the platform, provided a digital signal that indicated when the ball was at its start position. Additional reflex detectors, positioned at the lower edge of each slot, registered when the ball had fallen ~5 mm below the slot's lower edge. To estimate the position of the grasped ball during the transport phase, we used the sensors that recorded the fingertip position and the tips of the tweezers.

All data were sampled using the SC/ZOOM software (Department of Medical and Translational Biology, Umeå University). Signals from the various sensors were time synchronized and stored at 200 samples/s using linear interpolation between consecutive samples.

Gaze fixations were identified based on previously established criteria (Johansson et al., 2001). Fixations were classified into three categories according to their location, each defined as a centroid with a 2.5 cm radius: the ball at its start position, the target slot, and the text display. Note that fixation data were pooled across all slot positions (bottom, middle, and top). To qualify as a fixation, gaze had to remain within a designated zone for at least 100 ms. Notably, practically all fixations occurred within these zones (~95 %). Consistent with previous findings on object manipulation, participants almost never fixated their hand, or tweezers in hand, during hand movement.

To characterize the movement sequence in the ball-drop task we defined seven action phases based on the speed of the fingertip and tweezers (Fig. 1B)). Speed was computed as the vector sum of the first-time derivative of filtered horizontal and vertical position signals (2nd order Butterworth low-pass filter, 10 Hz cut-off frequency). (2nd order Butterworth low-pass filter, 10 Hz cut-off frequency). The start of the reach phase began when the end-effector speed exceeded 2 cm/s. The start of the subsequent ball approach phase was defined as the first minimum (or inflexion point) in the speed profile of the end-effector following the initial reach movement. The ball grasp phase spanned from the first contact with the ball to when its was lifted off the platform. The following transport phase began at lift off and ended at the notch in the speed profile marking the transition to the slot approach phase. The slot entry phase started when the ball was 1 cm to the right of the position where it was released inside the tube and ended when the ball was dropped, as detected by the optical sensor in the tube. Finally, the slot entry phase was followed by the return phase and the offset of the return phase were defined at the times at which the end-effector speed exceeded 2 cm/s.

We used paired t-tests with Holm-Bonferroni correction, Kolmogorov-Smirnov (KS) tests, and
multiple linear regression analyses in our statistical analyses as detailed in the Results. We
defined a statistically significant outcome if $p < 0.05$. All statistical analyses were conducted in
R (R Core Team, 2022; www.r-project.org). Trials in which participants dropped the ball after
initiating the transport phase were excluded. A total of 64 trials (4.7 %) were removed, with no
participant losing more than 7 trials.

Citation diversity statement

Recent work in several fields of science has identified a bias in citation practices such that
papers from women and other minority scholars are under-cited relative to the number of such
papers in the field (Bertolero et al., 2020; Caplar et al., 2017; Chatterjee & Werner, 2021; Dion
et al., 2018; Dworkin et al., 2020; Fulvio et al., 2021; Maliniak et al., 2013; Mitchell et al.,
2013; Wang et al., 2021). Here we sought to proactively consider choosing references that
reflect the diversity of the field in thought, form of contribution, gender, race, ethnicity, and
other factors. First, we obtained the predicted gender of the first and last author of each reference
by using databases that store the probability of a first name being carried by a woman (Dworkin
et al., 2020; Zhou et al., 2020). By this measure (and excluding self-citations to the first and last
authors of our current paper), our references contain 22.73% woman(first)/woman(last),
11.36% man/woman, 16.71% woman/man, and 49.19% man/man. This method is limited in
that a) names, pronouns, and social media profiles used to construct the databases may not, in
every case, be indicative of gender identity and b) it cannot account for intersex, non-binary, or
transgender people. Second, we obtained the predicted racial/ethnic category of the first and
last author of each reference by databases that store the probability of a first and last name being
carried by an author of colour (Ambekar et al., 2009; Chintalapati et al., 2023). By this measure
(and excluding self-citations), our references contain 5.3% author of colour (first)/author of
colour(last), 12.04% white author/author of colour, 16.71% author of colour/white author, and
65.95% white author/white author. This method is limited in that a) names and Florida Voter
Data to make the predictions may not be indicative of racial/ethnic identity, and b) it cannot
account for Indigenous and mixed-race authors, or those who may face differential biases due
to the ambiguous racialization or ethnicization of their names. We look forward to future work
that could help us to better understand how to support equitable practices in science.

References

Ambekar A, Ward C, Mohammed J, Male S, Skiena S. 2009. Name-ethnicity classification from open sourcesProceedings of the 15th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining. Presented at the KDD09: The 15th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining. Paris France: ACM. pp. 49–58. doi:10.1145/1557019.1557032

Ballard DH, Hayhoe MM, Li F, Whitehead SD. 1992. Hand-eye coordination during sequential tasks. *Philos Trans R Soc Lond B Biol Sci* **337**:331–339. doi:10.1098/rstb.1992.0111

Ballard DH, Hayhoe MM, Pelz JB. 1995. Memory Representations in Natural Tasks. *J Cogn Neurosci* **7**:66–80. doi:10.1162/jocn.1995.7.1.66

Bertolero MA, Dworkin JD, David SU, Lloreda CL, Srivastava P, Stiso J, Zhou D, Dzirasa K, Fair DA, Kaczkurkin AN, Marlin BJ, Shohamy D, Uddin LQ, Zurn P, Bassett DS. 2020. Racial and ethnic imbalance in neuroscience reference lists and intersections with gender. doi:10.1101/2020.10.12.336230

Bicchi A. 2000. Hands for dexterous manipulation and robust grasping: a difficult road toward simplicity. *IEEE Trans Robot Autom* **16**:652–662. doi:10.1109/70.897777

Bicchi A, Kumar V. 2000. Robotic grasping and contact: a reviewProceedings 2000 ICRA. Millennium Conference. IEEE International Conference on Robotics and Automation. Symposia Proceedings (Cat. No.00CH37065). Presented at the 2000 ICRA. IEEE International Conference on Robotics and Automation. San Francisco, CA, USA: IEEE. pp. 348–353. doi:10.1109/ROBOT.2000.844081

Billard A, Kragic D. 2019. Trends and challenges in robot manipulation. *Science* **364**:eaat8414. doi:10.1126/science.aat8414

Brink EE, Mackel R. 1987. Sensorimotor performance of the hand during peripheral nerve regeneration. *J Neurol Sci* **77**:249–266. doi:10.1016/0022-510X(87)90127-4

Caplar N, Tacchella S, Birrer S. 2017. Quantitative evaluation of gender bias in astronomical publications from citation counts. *Nat Astron* **1**:0141. doi:10.1038/s41550-017-0141

Casadio M, Pressman A, Mussa-Ivaldi FA. 2015. Learning to push and learning to move: the adaptive control of contact forces. *Front Comput Neurosci* **9**. doi:10.3389/fncom.2015.00118

Chatterjee P, Werner RM. 2021. Gender Disparity in Citations in High-Impact Journal Articles. *JAMA Netw Open* **4**:e2114509. doi:10.1001/jamanetworkopen.2021.14509

Chemnitz A, Dahlin LB, Carlsson IK. 2013. Consequences and adaptation in daily life – patients' experiences three decades after a nerve injury sustained in adolescence. *BMC Musculoskelet Disord* **14**:252. doi:10.1186/1471-2474-14-252

Chib VS, Krutky MA, Lynch KM, Mussa-Ivaldi FA. 2009. The Separate Neural Control of Hand Movements and Contact Forces. *J Neurosci* **29**:3939–3947. doi:10.1523/JNEUROSCI.5856-08.2009

Chintalapati R, Laohaprapanon S, Sood G. 2023. Predicting Race and Ethnicity From the
Sequence of Characters in a Name.

de Brouwer AJ, Gallivan JP, Flanagan JR. 2018. Visuomotor feedback gains are modulated by
gaze position. *J Neurophysiol* **120**:2522–2531. doi:10.1152/jn.00182.2018

Dion ML, Sumner JL, Mitchell SM. 2018. Gendered Citation Patterns across Political Science
and Social Science Methodology Fields. *Polit Anal* **26**:312–327.
doi:10.1017/pan.2018.12

Dworkin JD, Linn KA, Teich EG, Zurn P, Shinohara RT, Bassett DS. 2020. The extent and
drivers of gender imbalance in neuroscience reference lists. *Nat Neurosci* **23**:918–926.
doi:10.1038/s41593-020-0658-y

Eckstein MP. 2017. Probabilistic Computations for Attention, Eye Movements, and Search.
Annu Rev Vis Sci **3**:319–342. doi:10.1146/annurev-vision-102016-061220

Flanagan JR, Bowman MC, Johansson RS. 2006. Control strategies in object manipulation
tasks. *Curr Opin Neurobiol* **16**:650–659. doi:10.1016/j.conb.2006.10.005

Flanagan JR, Johansson RS. 2003. Action plans used in action observation. *Nature* **424**:769–
771. doi:10.1038/nature01861

Fooken J, Baltaretu BR, Barany DA, Diaz G, Semrau JA, Singh T, Crawford JD. 2023.
Perceptual-Cognitive Integration for Goal-Directed Action in Naturalistic Environments.
J Neurosci **43**:7511–7522. doi:10.1523/JNEUROSCI.1373-23.2023

Fooken J, Kreyenmeier P, Spering M. 2021. The role of eye movements in manual
interception: A mini-review. *Vision Res* **183**:81–90. doi:10.1016/j.visres.2021.02.007

Fulvio JM, Akinnola I, Postle BR. 2021. Gender (Im)balance in Citation Practices in
Cognitive Neuroscience. *J Cogn Neurosci* **33**:3–7. doi:10.1162/jocn_a_01643

Goodale MA, Pelisson D, Prablanc C. 1986. Large adjustments in visually guided reaching do
not depend on vision of the hand or perception of target displacement. *Nature* **320**:748–
750. doi:10.1038/320748a0

Hayhoe MM. 2017. Vision and Action. *Annu Rev Vis Sci* **3**:389–413. doi:10.1146/annurev-
vision-102016-061437

Hoppe D, Rothkopf CA. 2019. Multi-step planning of eye movements in visual search. *Sci*
Rep **9**:144. doi:10.1038/s41598-018-37536-0

Hoppe D, Rothkopf CA. 2016. Learning rational temporal eye movement strategies. *Proc Natl*
Acad Sci **113**:8332–8337. doi:10.1073/pnas.1601305113

Illamperuma NH, Fooken J. 2024. Towards a functional understanding of gaze in goal-
directed action. *J Neurophysiol* **132**:767–769. doi:10.1152/jn.00342.2024

Jenmalm P, Johansson RS. 1997. Visual and Somatosensory Information about Object Shape
Control Manipulative Fingertip Forces. *J Neurosci* **17**:4486–4499.
doi:10.1523/JNEUROSCI.17-11-04486.1997

Jerosch-Herold C. 1993. Measuring Outcome in Median Nerve Injuries. *J Hand Surg* **18**:624–
628. doi:10.1016/0266-7681(93)90019-C

Johansson RS, Flanagan JR. 2009. Coding and use of tactile signals from the fingertips in
object manipulation tasks. *Nat Rev Neurosci* **10**:345–359. doi:10.1038/nrn2621

Johansson RS, Westling G, Bäckström A, Flanagan JR. 2001. Eye–Hand Coordination in
Object Manipulation. *J Neurosci* **21**:6917–6932. doi:10.1523/JNEUROSCI.21-17-
06917.2001

Jovancevic-Misic J, Hayhoe MM. 2009. Adaptive Gaze Control in Natural Environments. *J*
Neurosci **29**:6234–6238. doi:10.1523/JNEUROSCI.5570-08.2009

Keshava A, Nezami FN, Neumann H, Izdebski K, Schüler T, König P. 2024. Just-in-time:
Gaze guidance in natural behavior. *PLOS Comput Biol* **20**:e1012529.
doi:10.1371/journal.pcbi.1012529

Kolesnikov M, Piovesan D, Lynch KM, Mussa-Ivaldi FA. 2011. On force regulation strategies
in predictable environments2011 Annual International Conference of the IEEE
Engineering in Medicine and Biology Society. Presented at the 2011 33rd Annual
International Conference of the IEEE Engineering in Medicine and Biology Society.
Boston, MA: IEEE. pp. 4076–4081. doi:10.1109/IEMBS.2011.6091013

Kowler E. 2011. Eye movements: The past 25years. *Vision Res* **51**:1457–1483.
doi:10.1016/j.visres.2010.12.014

Land M, Mennie N, Rusted J. 1999. The Roles of Vision and Eye Movements in the Control
of Activities of Daily Living. *Perception* **28**:1311–1328. doi:10.1068/p2935

Land MF. 2006. Eye movements and the control of actions in everyday life. *Prog Retin Eye*
Res **25**:296–324. doi:10.1016/j.preteyeres.2006.01.002

Land MF, Furneaux S. 1997. The knowledge base of the oculomotor system. *Philos Trans R*
Soc Lond B Biol Sci **352**:1231–1239. doi:10.1098/rstb.1997.0105

Land MF, Hayhoe M. 2001. In what ways do eye movements contribute to everyday
activities? *Vision Res* **41**:3559–3565. doi:10.1016/S0042-6989(01)00102-X

Maliniak D, Powers R, Walter BF. 2013. The Gender Citation Gap in International Relations.
Int Organ **67**:889–922. doi:10.1017/S0020818313000209

Mitchell SM, Lange S, Brus H. 2013. Gendered Citation Patterns in International Relations
Journals. Int Stud Perspect **14**:485–492. doi:10.1111/insp.12026

Najemnik J, Geisler WS. 2005. Optimal eye movement strategies in visual search. *Nature*
434:387–391. doi:https://doi.org/10.1038/nature03390

Neggers SFW, Bekkering H. 2001. Gaze Anchoring to a Pointing Target Is Present During the
Entire Pointing Movement and Is Driven by a Non-Visual Signal. *J Neurophysiol*
86:961–970. doi:10.1152/jn.2001.86.2.961

Neggers SFW, Bekkering H. 2000. Ocular Gaze is Anchored to the Target of an Ongoing
Pointing Movement. *J Neurophysiol* **83**:639–651. doi:10.1152/jn.2000.83.2.639

Piovesan D, Kolesnikov M, Lynch K, Mussa-Ivaldi FA. 2019. The Concurrent Control of
Motion and Contact Force in the Presence of Predictable Disturbances. *J Mech Robot*
11:060903. doi:10.1115/1.4044599

Prablanc C, Pélisson D, Goodale MA. 1986. Visual control of reaching movements without
vision of the limb: I. Role of retinal feedback of target position in guiding the hand. *Exp*
Brain Res **62**. doi:10.1007/BF00238848

Pruszynski JA, Flanagan JR, Johansson RS. 2018. Fast and accurate edge orientation
processing during object manipulation. *eLife* **7**:e31200. doi:10.7554/eLife.31200

Pruszynski JA, Johansson RS, Flanagan JR. 2016. A Rapid Tactile-Motor Reflex
Automatically Guides Reaching toward Handheld Objects. *Curr Biol* **26**:788–792.
doi:10.1016/j.cub.2016.01.027

Renninger LW, Verghese P, Coughlan J. 2007. Where to look next? Eye movements reduce
local uncertainty. *J Vis* **7**:6. doi:10.1167/7.3.6

Säfström D, Johansson RS, Flanagan JR. 2014. Gaze behavior when learning to link
sequential action phases in a manual task. *J Vis* **14**:3–3. doi:10.1167/14.4.3

Saunders JA, Knill DC. 2004. Visual Feedback Control of Hand Movements. *J Neurosci*
24:3223–3234. doi:10.1523/JNEUROSCI.4319-03.2004

Saunders JA, Knill DC. 2003. Humans use continuous visual feedback from the hand to
control fast reaching movements. *Exp Brain Res* **152**:341–352. doi:10.1007/s00221-003-
1525-2

Wang X, Dworkin JD, Zhou D, Stiso J, Falk EB, Bassett DS, Zurn P, Lydon-Staley DM. 2021.
Gendered citation practices in the field of communication. *Ann Int Commun Assoc*
45:134–153. doi:10.1080/23808985.2021.1960180

Wilmut K, Wann JP, Brown JH. 2006. How active gaze informs the hand in sequential
pointing movements. *Exp Brain Res* **175**:654–666. doi:10.1007/s00221-006-0580-x

Yarbus AL. 1967. Eye Movements and Vision. Boston, MA: Springer US. doi:10.1007/978-1-
4899-5379-7

Zhou D, Bertolero MA, Stiso J, Cornblath EJ, Teich EG, Blevins AS, Virtualmario Camp,
Dworkin J, Bassett DS. 2020. Gender diversity statement and code notebook.