1 **Temperate phages increase antibiotic effectiveness in a *Caenorhabditis elegans* infection**
2 **model**
3

Authors
Rabia Fatima[1], Lesley T. MacNeil[1,2,3], Alexander P. Hynes[1,2,3,4]

[1] Department of Biochemistry and Biomedical Sciences, McMaster University.

[2] Farncombe Family Digestive Health Research Institute, McMaster University

[3] Michael G. DeGroote Institute for Infectious Disease Research, McMaster University

[4] Department of Medicine, McMaster University.

Correspondence
Alexander P Hynes, hynes@mcmaster.ca

Abstract

The bactericidal nature of obligately lytic bacterial viruses (phages) is of increasing interest for the treatment of drug-resistant bacterial infections, either administered alone or in combination with antibiotics. In contrast, temperate phages are largely ignored in a therapeutic context due to their ability to lie dormant within the bacterial host. However, these phages often undergo a lytic cycle. Furthermore, even in their dormant state their carriage can be a considerable burden to the bacterium – most famously by their ability to switch to lytic replication in response to environmental triggers, such as antibiotics, that stress the bacterial host. Recent reports of antibiotics synergizing uniquely with temperate phages *in vitro*, termed "temperate phage antibiotic synergy" (tPAS), present a potentially scalable opportunity to make use of these abundant entities for the treatment of bacterial infections. Here we employ *Caenorhabditis elegans* as a robust *in vivo* animal model for testing the efficacy of temperate phages as adjuvants to antibiotics. *In vivo*, the antibiotic ciprofloxacin can abolish dormancy of temperate phage Hali - infecting a ciprofloxacin resistant *Pseudomonas aeruginosa* clinical strain - while in the absence of the antibiotic dormancy events explain 60% of the bacterial survivors. The Phage Hali-ciprofloxacin pairing increased the lifespan of *P. aeruginosa* infected worms to that of the uninfected control, at doses where neither the phage nor the antibiotic had any effect alone. Complete rescue was also observed in worms infected with a phage-carrying strain treated with the otherwise ineffective antibiotic, supporting that the phage - even in its dormant form - can greatly enhance antibiotic effectiveness. This illustrates potential "accidental" phage therapy when antibiotics are normally prescribed. Our work establishes *C. elegans* as a suitable model for studying the *in vivo* efficacy of tPAS and is the first *in vivo* demonstration of this synergy, greatly expanding the therapeutic potential of temperate phages.

Introduction

In light of the growing antimicrobial resistance crisis, bacteriophages (phages) are increasingly being explored as adjuvants or alternatives to antibiotics. Most of this work has focused on a particular type of phage called virulent or strictly lytic, given their immediate bactericidal nature (1–12). However, it is estimated that 50% - 75% of all bacteria contain at least one dormant phage, termed temperate or lysogenic (13, 14). Much of the resistance to the use of temperate phages in therapy comes from concerns of further spread of fitness-benefitting genes within the bacterial population, especially the spread of toxins or antibiotic resistance cassettes. While on average 3.06 ± 1.84 dormant phages were predicted per clinical isolate of *Pseudomonas aeruginosa*, they were not associated with antibiotic resistance of the bacteria (15) and estimate of phage-encoded antibiotic resistance genes appears to be overestimated due to inherent biases in bioinformatic based virome studies (16). In addition, temperate phages are not inherently better transducers than virulent phages, many of which are capable of generalized transduction (17–19).

A key concern specific to temperate phages comes at the level of the integrated phage (prophage) conferring to its host immunity to subsequent phage infections through a variety of mechanisms (20, 21). This leads to a rapid regrowth of phage-generated resistant 'mutants' and could readily contribute to treatment failure. However, carrying a large genome such as a prophage, especially if it does not confer direct fitness benefit, can pose a heavy metabolic burden on the bacteria. Lysogens of several *Pseudomonas aeruginosa* phages demonstrate defects in bacterial motility, swarming and twitching (22). As a result, the prophage can have lasting influence on their host physiology, virulence, and pathogenesis, which could be exploited for the treatment of bacterial infections.

One of the most common physiological changes in the prophage-carrying lysogen is the one from which it derives its name; the prophage can be highly responsive to environmental cues that stress their bacterial host in a process referred to as induction (23). In the well-characterized phage model Lambda infecting *Escherichia coli,* the phage can sense host DNA damage to switch from lysogenic to lytic replication, resulting in cell lysis (24). In the gastrointestinal tract of monoxenic mice, 1 to 2% of Lambda lysogenic bacteria were lysed *per* generation due to spontaneous prophage induction (25). Hence, prophage excision itself likely presents a higher burden to the bacteria, especially in environments of high stress. Lysogens can be exhibit increased sensitivity to antibiotics, monitored as decreases in minimum inhibitory concentration (MIC), compared to the parent strain (26–28). Temperate phages of *Burkholderia cepacia* (29)*, E. coli* (28, 30)*,* and *P. aeruginosa* (27) can also function synergistically with antibiotics, resulting in increases in phage production (29) or a decrease in the functionally effective dose of the antibiotic (28). When the mechanism of action specifically interacts with the phage lysis-lysogeny decision, this is referred to as temperate phage-antibiotic synergy (tPAS). Given that temperate phages are already dominate within the human body (31), there is a strong case in favour of their therapeutic use - they are likely already an influential player in health and treatment outcomes.

There are currently several popular animal models for studying efficacy of phage therapy. Specifically for temperate phages, *Burkholderia* phage AP3 was able to significantly increase survival of wax moth larvae (15% survival at 72 h compared to 0% for untreated at 48 h) (32). In

a hamster model, a four-phase cocktail targeting *Clostridioides difficile* reduced the colonization
load and delayed the onset of symptoms (33). More recently, temperate phage of *Acinetobacter*
baumannii, vB_AbaM_ABMM1, was able to reduce mortality in a zebrafish model (34). While
useful, these models present several scalability concerns, each animal requiring individual
attention or manipulation.

In this study, we develop the nematode *Caenorhabditis elegans* as an infection model for
studying the *in vivo* efficacy of a temperate phage-antibiotic combination against the multidrug-
resistant bacteria *P. aeruginosa*. Due to its short and measurable lifespan and the ability to
generate large, germ-free synchronous populations in the lab, *C. elegans* is a well-established
model for the study of bacterial pathogenesis (35). Many human pathogens, including *P.*
aeruginosa (36) are also pathogenic to *C. elegans*. Accordingly, *C. elegans* are also commonly
employed as an *in vivo* model for studying the efficacy of new antibacterial drugs (37–39).
Pathogenesis in *C. elegans* of *P. aeruginosa* has been studied in three different contexts. Agar
based methods are differentiated by fast and slow killing, dependent on media composition (40).
Fast killing occurs within hours, does not require live bacteria, and is mediated by bacterial
secreted toxic phenazines (40, 41). In contrast, slow killing occurs over the course of several
days with active intestinal colonization being a key determinant (40). A third *P. aeruginosa*-
mediated *C. elegans* killing method is observed in liquid, with the same formulation as that of
slow killing medium (SKM) (42). While not as well understood, this is independent of phenazine
production or intestinal colonization, however, is reported to be mediated by *P. aeruginosa*
secreted siderophore pyoverdine which induces a hypoxic response and subsequent worm death.
100
Only a few studies have used *C. elegans* to investigate phage therapy, focusing on virulent
phages of *B. pseudomallei* (43), *Salmonella* Enteritidis (44), *Staphylococcus aureus(45)*, *E. coli*
(46), *Klebsiella pneumoniae* (46), *Enterobacter cloacae* (46), with only one specifically for *P.*
aeruginosa (47). The primary outcome measured in these studies was the ability of phages to
reduce bacterial load and/or increase survival post infection. However, none of these studies
tested antibiotics alone or in combination with their phage of interest in this model. In addition,
in vitro results are not always sufficient indicators of *in vivo* efficacy (48, 49), especially in cases
of antibiotic potency which can be highly sensitive to media composition, pathogen, and host
factors (50, 51). As a result, here we employ a simple animal model to be able to quickly iterate
and screening a large set of temperate phage-antibiotic pairings that synergized *in vitro* against
P. aeruginosa (27).

Results/Discussion

Validation of C. elegans as an infection model

To investigate if co-administration of temperate phage and antibiotic can result in bacterial
killing in an *in vivo C. elegans* model, we first validated *P. aeruginosa* infection with a clinical
strain of interest, C0400. C0400 is a ciprofloxacin resistant strain where co-administration of
temperate phage Hali and ciprofloxacin was able to re-sensitize the strain to the antibiotic *in*
vitro (27). The strain PA14 is commonly used in this model for studying *P. aeruginosa*
pathogenesis (40, 41) and served as our control. For our experiments, we employed the *glp*-4
(*bn2*) worm strain that carries a temperature sensitive defect in germ line proliferation such that
animals maintained at a restrictive temperature of 25$^{\text{o}}$C are sterile (52). In slow killing assays,

dependent on host colonization, both *P. aeruginosa* strains tested colonized the intestine at levels
higher than *E. coli* OP50 control, with no differences between PA14 and C0400 (Fig. S1a).
Similarly, while OP50 does not result in worm

Fig 1. Representation of the assay used for testing efficacy of temperate PAS in a *C. elegans* model
of *P. aeruginosa* infection. Worms are bleached to collects eggs which are hatched and synchronized to
L1 stage. Animals are developed on *E. coli* OP50 and infected with *P. aeruginosa,* followed by phage
and/or antibiotic treatment. Outcome of treatment is monitored by bacterial colonization and lifespan on
regular bacterial diet. Each assay was done in independent biological replicates, with technical triplicates
plated for assessing bacterial load.

killing, *P. aeruginosa* fed worms maintained on SKM begin dying after three days with complete
population death observed by day 10 (Fig. S1b). The lifespan data closely matches previously
reported survival data of the *glp-4* strain exposed to OP50 and PA14 (52), validating that the
clinical *P. aeruginosa* strain C0400 is suitable for investigating tPAS in this *in vivo* model.
135

Ciprofloxacin selects against lysogens in vivo

To test whether tPAS can reduce *P. aeruginosa* loads *in vivo*, we used two different treatment
durations, 4 h and 18 h, after infection on SKM (Fig. 2). We treated the worms with phage Hali
and ciprofloxacin across a range of antibiotic concentrations (0.25 µg/mL – 100 µg/mL), alone
and in combination, only showing the data for highest concentration (100 µg/mL) in Fig 2a as
there was no effect observed at lower doses. There was no reduction in colonization when treated
with phage alone and only 1-2 $\log_{10}$ reduction observed with the highest ciprofloxacin
concentration tested. Similarly, there was no synergistic reduction in bacterial load for tPAS for
either treatment duration.

Despite no reduction in bacterial loads observed, it was still possible that tPAS biased the
survivor profile, selecting against lysogens. The phage alone results in an increase in lysogens
from 20% to 60% as treatment duration increased (Fig. 2b). This regrowth of lysogens is
expected as they are immune to further phage infection due to superinfection immunity and
aligns very closely with our previous report that phage Hali results in 75% lysogens after an
overnight challenge *in vitro* (27). In comparison, tPAS resulted in primarily lysogen population
within 4 h treatment (80% compared to 20% for phage alone) and a reduction in frequency of
lysogeny to zero with longer treatment duration. Compared to administering phage by itself,
tPAS selectively biases the phage away from lysogeny *in vivo,* completely preventing the
regrowth of lysogens that would otherwise form 60% of the population. This formation of

lysogens (4 h) that are then depleted is consistent with the previously reported mechanism of this interaction to be operating through induction (27).

Fig 2. Phage Hali and ciprofloxacin selects against lysogen survivors *in vivo*. (a) *C. elegans* intestinal colonization measured as log10 cfu/mL/worm after 48 h C0400 infection followed by 4 h or 18 h treatment with phage Hali (1 x 10^9 pfu/mL) $\pm$ ciprofloxacin. Data shown as mean $\pm$ SD, each data point is a biological replicate plated in technical triplicates, except ones denoted by triangle which are single technical replicate, with of ~50-100 worms each condition. The two treatment durations were tested in independent trials. Values were compared using two-way ANOVA with a multiple comparison Tukey test with conditions only compared within their own treatment duration. All comparisons are not significant. **(b)** Frequency of lysogeny within survivors of phage Hali alone and phage Hali + ciprofloxacin (tPAS). Twenty survivors were tested for each condition.

In short, while all treatments had – at best – modest effects on the bacterial load, combining phage and antibiotic drastically altered the nature of those bacteria, selecting against lysogens. We sought to determine whether this bias was enough to rescue the worms after treatment.

Combined temperate phage and ciprofloxacin rescues P. aeruginosa infected worms

To test if selecting for a different survivor population would be enough to rescue the worms, we carried out the same assay but moved the animals to their standard *E. coli* OP50 diet after 18 h treatment to measure lifespan. We performed the lifespan assay with a shorter 4 h infection as we did not observe any rescue from even the antibiotic at high doses with a longer infection duration, suggesting that worms were moribund (data not shown). With a shorter infection, when left untreated, the worms died within maximum thirteen days compared to eighteen plus days for the uninfected OP50 control as they reach the end of their lifespan (Fig. 3). Treatment with phage Hali alone was not only ineffective, but it also showed a worse survival outcome compared to untreated in two of the three trials (Fig. 3). Ciprofloxacin alone at low antibiotic dose of 2 µg/mL did not reliably rescue alone or in combination with phage (Fig. S2a). At a much higher antibiotic dose, 92 µg/mL, the antibiotic alone was sufficient to rescue (Fig. S2b). However, we could not reliably discern the effect of tPAS as both antibiotics alone and tPAS survival curve matched the uninfected OP50 control in two trials.

We were primarily interested in antibiotic concentrations that were lower than the effective dose and was therefore insufficient to increase survival. Ciprofloxacin alone at 32 µg/mL could not rescue survival (Fig. 3). In comparison, tPAS increased lifespan to levels comparable to the

186　uninfected OP50 control in two of the three trials, and a marked improvement in survival over
187　either treatment alone in the third. In short, the *in vitro* work that had shown synergy between

**Fig 3. Phage Hali and low dose ciprofloxacin increase survival of *P. aeruginosa* infected worms on
standard *E. coli* OP50 diet after treatment.** Representative life span of worms on OP50 after 4 h *P.
aeruginosa* infection followed by 18 h phage Hali　ciprofloxacin treatment**.** Approximately 50 worms
were added to each condition at the time of treatment, tested in three biological replicates. Survival curves
were compared using the Log-rank (Mantel Cox) test with one pair compared at a time, ns = not
significant, * P ≤ 0.05, *** P ≤ 0.001, and **** P ≤ 0.0001. Phage alone and antibiotic alone were
compared to the untreated, and tPAS to the phage alone and antibiotic alone, as denoted by colour of the
significance value.

phage Hali and ciprofloxacin translated to a robust, reproducible increase in worm survival at concentrations where neither agent alone had any detectable effect.

We wished to correlate survival with bacterial colonization assays as carried out earlier (Fig 2) but found that treatment outcomes for these shorter 4 h infections did not correlate to bacterial loads (Fig S3). For instance, 32 µg/mL ciprofloxacin with or without phage were both successful in eradicating *Pseudomonas*, even though the antibiotic-alone treatment had no effect on lifespan (Fig. S3). The most likely explanation is that the phage-antibiotic combination is benefiting the worms through either faster eradication, qualitatively different endpoints for the bacterium (phage-mediated lysis rather than antibiotic-mediated death), or that the presence of transient lysogeny is in some way mitigating the virulence of the strain.

The lysogen is more sensitive to ciprofloxacin in vivo

To determine whether the formed lysogens were differentially virulent, we infected worms with wild type C0400 and a C0400 phage Hali lysogen, followed by ciprofloxacin treatment and subsequently monitoring survival on the standard *E. coli* OP50 diet. Without treatment, the pathogenicity of the lysogen and wild type strains did not differ (Fig. 4a). Intriguingly, lysogen-infected worms could be completely rescued with 32 µg/mL ciprofloxacin , which was insufficient to rescue worms infected with the wild type C0400. This strongly supports the notion that the tPAS observed by co-administration of the phage and antibiotic is primarily driven by the interaction between the prophage and the antibiotic, as any phages induced (spontaneously or by the antibiotic) would have little effect on the lysogens, which are immune to superinfection.

This finding is particularly important mechanistically, because one possible explanation for tPAS would be that the phage administration results in a modest reduction of the bacterial load prior to the regrowth of lysogens, and that at this lower bacterial density the bacteria are more sensitive to the antibiotic – a concept known as density-dependent antibiotic sensitivity (53, 54). However, here the phage and antibiotic are truly synergistically interacting. This similar rescue observed to tPAS (Fig. 3) also means that irrespective of whether the phage is co-administered or if it exists as a prophage, it can increase the effectiveness of the antibiotic if it is antibiotic inducible.

Considering that lysogens are often more sensitive to environmental stressors because of carrying prophages that are responsive to these triggers (26–28), we assessed if carrying a prophage substantially burdens the bacteria *in vivo*, especially in the presence of antibiotic, by measuring intestinal bacterial colonization of worms infected with wild type or lysogen and treated with ciprofloxacin. Ciprofloxacin results in a higher reduction in bacterial load in lysogen infected animals compared to wild type (Fig. 4b). This was associated with a higher phage count inside the worms after antibiotic treatment, consistent with induction (Fig. 4c). These data are consistent with *in vitro* report that phage Hali lysogens shows 4-fold lower ciprofloxacin MIC compared to wild type, and that this is due to prophage induction (27).

The increase survival of phage Hali lysogen infected worms treated with ciprofloxacin taken together with a greater reduction in bacterial load observed for the lysogen suggests that carrying an inducible phage likely imposes a major fitness cost under mild antibiotic stress. Given the abundance of prophages within bacterial hosts (14, 31) and that pathogens are well-known reservoirs of prophages (55, 56), this effect is likely already taking place during antibiotic treatments, in a form of "accidental phage therapy". It is important to note that this is likely phage dependent, not just with regards to its integration site and receptor, but especially in its

ability to respond to antibiotic triggers. The *P. aeruginosa* LES strain common in cystic fibrosis patients is known to contain several prophages that can be induced with ciprofloxacin *in vitro* (57). In

Fig 4. Phage Hali lysogen is more sensitive to ciprofloxacin *in vivo*. **(a)** Representative life span of worms on OP50 after 4 h *P. aeruginosa* C0400 wild type and phage Hali lysogen infection followed by 18 h ciprofloxacin treatment**.** Approximately 50 worms were added to each condition at the time of treatment, tested in three biological replicates. Survival curves were compared using the Log-rank (Mantel Cox) test with one pair compared at a time, ns = not significant, * P ≤ 0.05, *** P ≤ 0.001, and **** P ≤ 0.0001. Phage alone and antibiotic alone were compared to the untreated, and tPAS to the phage alone and antibiotic alone, as denoted by colour of the significance value. **(b)** *C. elegans* intestinal C0400 wild type and phage Hali lysogen colonization measured as log10 cfu/mL/worm after 48 h infection followed by 18 h treatment with ciprofloxacin. **(c)** Corresponding phage quantification shows as log10 pfu/mL/worm for the lysogen infected, and ciprofloxacin treated worms. All bar graph data shown as mean SD where each data point is a biological replicate. Colonization assay carried out with roughly 100 worms per condition.

addition, from patient data, the induction of these phages was shown to play a crucial role in regulation of the pathogen in CF lungs (58). However, it is challenging to correlate the effects to long term antibiotic prescription since the data are incomplete. However, James et al. (2014) do propose that treatments that result in phage induction could be promising approaching to tackling

259 *P. aeruginosa* load in these patients, not without caution in case of phage encoded virulence
260 factors (58).

Conclusion

In phage therapy, temperate phages are discarded due to their ability to lysogenize. However,
these have an enormous untapped potential due to their ability to respond to host stressing
environmental cues, such as antibiotics, to switch into lytic replication, killing its host in the
process. Here we show that *C. elegans* is a suitable model for studying *in vivo* efficacy of tPAS,
but this can easily be extended to traditional PAS with virulent phages. While we did not observe
synergistic reduction in bacterial load, ciprofloxacin strongly selects against phage Hali
lysogens. In addition, even if lysogens do arise, they can be more sensitive to the antibiotic *in*
vivo, compared to the parental strain. While phage Hali is completely insufficient to rescue the
worms, phage and ciprofloxacin combined increased the lifespan of *P. aeruginosa* infected
worms to levels comparable to the uninfected control, at a similarly ineffective antibiotic dose.
Complete rescue can be seen for the lysogen-infected worms treated with the antibiotic. The
phage, even in its prophage form, greatly enhanced antibiotic effectiveness. This suggests to us
that this kind of 'accidental' phage therapy already often occurs when antibiotics are prescribed.

Acknowledgement

The authors would like to Mercedes DiBernardo for helping set up the worm strains and sharing
her expertise on the colonization assay, as well as Sommer Chou for providing training on the
worm sorter. We would also like to thank Dr. Gerry Wright for access to the *P. aeruginosa*
clinical isolates from the Wright Clinical Isolate Collection at the Institute for Infectious Disease
Research (McMaster University, Ontario) and Dr. Lori Burrows for PA14.

R.F was supported by an NSERC Canadian Graduate Scholarship, Master's and Doctoral
scholarship. A.P.H. acknowledges funding through the Natural Sciences and Engineering
Council of Canada (NSERC) Discovery Grant 2018-05996 and the Farncombe Family Chair in
Phage Biology.

R.F. performed all the assays, and, jointly with A.P.H., conceived the study. L.M. provided the
worm strains, *E. coli* OP50, and all the experiments were performed at L.M. laboratory. R.F. and
A.P.H contributed to the writing of the manuscript.

Material and Methods

Bacterial strains and growth conditions

P. aeruginosa strain PA14 was obtained from Dr. Lori Burrows, McMaster University and *P.*
aeruginosa C0400 was from the McMaster IIDR Wright clinical isolate collection. Bacterial
strains were grown in 10 mL of lysogeny broth (LB) at 37°C with 250 rpm shaking (Ecotron,
Infors HT, Quebec, Canada). For growth on solid media, 1% (w/v) of LB (lysogeny broth) agar
and 0.75% (w/v) of LB soft agar was used. All plates were incubated at 37°C from overnight
growth.

To prepare *C. elegans* plates, bacterial strains were streaked out on 1% LB plates and incubated overnight. A single colony was inoculated into 10 mL LB broth, followed by overnight incubation with shaking. To seed plates, 200 µL of overnight culture was used for 100 mm plates and 100 µL was used for 60 mm plates. The plates were incubated overnight at 37°C prior to transferring worms on to them.

Phage propagation and titration

Phage Hali infecting *P. aeruginosa* PA14 and C0400 was used in this study. Phage lysates were prepared either using a primary or secondary amplification. Briefly, primary amplification was performed by inoculating 10 mL LB broth with bacteria and phage from frozen glycerol stocks for amplification to take place overnight. Secondary amplification was performed as needed to increase phage titer by inoculating 10 mL of same day culture prepared in LB with 50 µL of primary amplification. Phages in filtrate were quantified by standard spot test or full plate plaque assay.

Multiplicity of infection (M.O.I) was calculated based on titer and volume of bacteria and phage being combined using the following formula: $\frac{\text{phage titer}\left(\frac{\text{pfu}}{\text{mL}}\right) \times \text{phage volume (mL)}}{\text{bacterial titer}\left(\frac{\text{cfu}}{\text{mL}}\right) \times \text{volume(mL)}}$. The bacterial titer was determined as previously described (27).

C. elegans **strains and growth conditions**

All experiments were performed using *C. elegans* SS104, a temperature sensitive sterile strain, *glp-4* (*bn2*) available from the *Caenorhabditis* Genetic Center (CGC, https://cgc.umn.edu/). Worms were maintained at 15°C on 100 mm nematode growth media (NGM) plates (59) seeded with overnight cultures of *E. coli* OP50, prepared as indicated above. Eggs were collected by hypochlorite treatment (60). Eggs were rocked approximately 18 h at 20°C to synchronize to L1 larval stage. L1 worms were plated on 100 mm NGM plates seeded with *E. coli* OP50 and developed to young adults at 25°C, restrictive temperature at which they are sterile, for approximately 40 h.

Antibiotic stock preparation

Ciprofloxacin (hydrochloride) was obtained from Cayman Chemicals (Catalog 14286-5, Ann Arbor, Michigan, USA). Working stocks were prepared in nuclease free water. The minimum inhibitory concentration of each newly prepared batch was re-evaluated prior to use as reported previously (27).

C. elegans P. aeruginosa **infection**

For *P. aeruginosa* infection, young adults were washed off NGM OP50 plates with 10 mL M9 and washed three times to remove excess bacteria. Worms were transferred to 100 mm Slow Killing Media plates (SKM; 0.35% peptone, 50 mM NaCl, 2% agar, 1 mM CaCl$_2$, 5 µg/ml cholesterol, 1 mM MgSO$_4$, 20 mM KH$_2$PO$_4$, 5 mM K$_2$HPO$_4$) seeded with different *P. aeruginosa* strains for appropriate infection duration as indicated in figure legends. Worms were also plates on *E. coli* OP50 for the life span assay as a control.

Preparing *P. aeruginosa* infected worms for treatment

After infection with *P. aeruginosa,* worms were washed off SKM plates with 10 mL M9, followed by another three M9 washes. Each wash step consisted of pelleting the worm 2 min at 12 min and removing the supernatant, except for the third wash in which they were allowed to settle by gravity. To further dilute the bacteria, 1 mL of M9 with 10 ug/mL of tobramycin was added and worms were pelleted using centrifugation, followed by transfer to an unseeded 100 mm 10 ug/mL SKM tobramycin plate for 45 min in the biosafety cabinet to allow remove surface attached bacteria. After crawling, they were washed off with SKM tobramycin plates with 10 mL M9 tobramycin and subsequently washed another three times with 10 mL M9 tobramycin and final three times with just M9 to dilute out the tobramycin.

Colonization assay

To measure the efficacy of tPAS on bacterial colonization, approximately 100 infected worms were added to 2 mL tube manually, along with ciprofloxacin (final concentration tested 0.5 µg/mL – 32 µg/mL at 2-fold increments, then 32 µg/mL - 100 µg/mL in increments of 10 µg/mL) and/or phage Hali at 1×10^9 pfu. The final volume was topped up to 1 mL using SKM broth. SKM alone was used a negative control. Tubes were incubated for 18 h with 150 rpm horizontal shaking and colonization assay was carried out subsequently.

Following treatment, worms were settled by centrifugation for 2 min at 1200 rpm or by gravity when working with more than 10 samples. After removing the supernatant, worms were washed three times with 1 mL M9, followed by once with 1 mL 10 µg/mL M9 tobramycin. To remove surface attached bacteria, they were transferred to 60 mm 10 µg/mL SKM tobramycin plates for 45 min. At this point, the total number of animals for each condition was counted.

Worms were washed off plates with 2 mL 120 µM levamisole in M9 (LM buffer). The number of worms remaining on the plate was counted to determine the number of animals per condition going into the lyses step. Animals were washed with 1 mL 10 µg/mL tobramycin LM buffer; once for 4 h treatment and three times for 18 h treatment. To dilute out the tobramycin, worms were additionally washed three times with 1mL M9. In the last wash, 500 µL of supernatant was transferred into a new tube.

Approximately ten sterile silica carbide beads were added to each tube and worms were lysed for 1 min at speed 5 using the Bead Mill 4 (Fisherbrand, catalog 15340164). Lysed fractions and the last wash supernatants were serial diluted 10-folds in LB up to 10^{-8} and 3 µL was spotted onto a 1% LB plate in technical triplicates, along with one replicate for the last wash. LB only and M9 with beads were spotted as control. Plates were incubated at 32°C to prevent overgrowth and colony forming unit were counted after an overnight incubation. Bacterial load was determined by subtracting the counts from the last wash and the cfu/mL was normalized to the number of animals per condition.

Frequency of lysogeny

Frequency of lysogeny in the survivors was determined as previously described (27). Twenty colonies for each condition were streak purified three times on 1% LB plate and inoculated in

200 – 250 µL LB broth overnight in a narrow 96 well plate. Wild type culture, C0400 phage Hali
lysogen, and LB broth were added as control. The plates were incubated for 18-24 h at 37^{o}C with
no agitation. The inoculated plates were stamped on an agar overlay of wild type C0400 using a
disposable pin replicator. Plates were also stamped onto 1% LB with no agar overlay to serve as
growth controls. Frequency of lysogeny was calculated as percent of the total number of
survivors that resulted in clearing of the wild type host.

Survival on OP50 life span assay

C. elegans were infected with *P. aeruginosa* strains, or OP50 as control, for 4 h and washed off
as indicated outline above. The volume of infected worms per strain was adjusted to 1 worm/µL
and approximately 100 worms were sorted into a 96 well using the COPAS FP (Union
Biometrica, Holliston, MA). These were then transferred to a 2 mL tube, along with 50 µL of
20x ciprofloxacin stock (final concentration tested 0.25 µg/mL – 32 µg/mL at 2-fold increments,
then 32 µg/mL - 100 µg/mL in increments of 10 µg/mL), phage Hali at 1×10^{9} pfu. Nuclease
free water was used as substitute for antibiotic and LB in place for phage. The final volume was
topped up with SKM broth to 1 mL. The worms were incubated in treatment for 18 h shaking
horizontally at 150 rpm.

Following treatment, animals were washed with 1 mL M9 three times, allowed to settle by
gravity in the third wash, and once with 1 mL 10 ug/mL M9 tobramycin. They were transferred
to 60 mm SKM tobramycin plates for 45 min to crawl, after which they were washed off with 2
mL M9 tobramycin. They were washed once more with 1 mL M9 tobramycin, ending with final
three washes with M9 alone.

After the last wash, animals were moved to 60 mm SKM plates seeded with OP50. Plates were
incubated at 25^{o}C and scored daily starting day three for survival, with dead worms removed
each day. Survival was plotted using the Kaplan–Meier method.

Statistical tests

Details of the statistical tests can be found in the figure legend. N denotes a biological replicate
performed from an independently grown culture with appropriate technical replicates indicated
in the figure caption. Quantitative values are represented by mean $\pm$ SD. All statistical analysis
was done using GraphPad Prism 10.3.0 (GraphPad Software, Inc., CA, US), with P value ≤ 0.05
is   considered   significant.

References

1. Akturk E, Oliveira H, Santos SB, Costa S, Kuyumcu S, Melo LDR, Azeredo J. 2019. Synergistic action of phage and antibiotics: parameters to enhance the killing efficacy against mono and dual-species biofilms. Antibiotics 8(3):103.

2. Chaudhry WN, Concepcion-Acevedo J, Park T, Andleeb S, Bull JJ, Levin BR. 2017. Synergy and order effects of antibiotics and phages in killing *Pseudomonas aeruginosa* biofilms. PLoS One 12(1):e0168615.

3. Henriksen K, Rørbo N, Rybtke ML, Martinet MG, Tolker-Nielsen T, Høiby N, Middelboe M, Ciofu O. 2019. *P. aeruginosa* flow-cell biofilms are enhanced by repeated phage treatments but can be eradicated by phage-ciprofloxacin combination - monitoring the phage - *P. aeruginosa* biofilms interactions. Pathog Dis 77(2):ftz011.

4. Kamal F, Dennis JJ. 2015. *Burkholderia cepacia* complex phage-antibiotic synergy (PAS): antibiotics stimulate lytic phage activity. Appl Environ Microbiol 81(3):1132-8.

5. Kebriaei R, Lev K, Morrisette T, Stamper KC, Abdul-Mutakabbir JC, Lehman SM, Morales S, Rybaka MJ. 2020. Bacteriophage-antibiotic combination strategy: an alternative against methicillin-resistant phenotypes of *Staphylococcus aureus*. Antimicrob Agents Chemother 64(7):e00461-20.

6. Kim M, Jo Y, Hwang YJ, Hong HW, Hong SS, Park K, Myung H. 2018. Phage antibiotic synergy via delayed lysis. Appl Environ Microbiol 84(22):e02085-18.

7. Liu CG, Green S, Min L, Clark J, Salazar K, Terwilliger A, Kaplan H, Trautner B, Ramig R, Maresso A. 2020. Phage-antibiotic synergy is driven by a unique combination of antibacterial mechanism of action and stoichiometry. mBio 11(4):e01462-20.

8. Rodriguez-Gonzalez RA, Leung CY, Chan BK, Turner PE, Weitz JS. 2020. Quantitative models of phage-antibiotic combination therapy. mSystems 5(1):e00756-19.

9. Aslam S, Lampley E, Wooten D, Karris M, Benson C, Strathdee S, Schooley RT. 2020. Lessons learned from the first 10 consecutive cases of intravenous bacteriophage therapy to treat multidrug-resistant bacterial infections at a single center in the United States. Open Forum Infect Dis 7(9):ofaa389.

10. Chan BK, Turner PE, Kim S, Mojibian HR, Elefteriades JA, Narayan D. 2018. Phage treatment of an aortic graft infected with *Pseudomonas aeruginosa*. Evol Med Public Health 2018(1):60-66.

11. Gainey AB, Burch AK, Brownstein MJ, Brown DE, Fackler J, Horne B, Biswas B, Bivens BN, Malagon F, Daniels R. 2020. Combining bacteriophages with cefiderocol and meropenem/vaborbactam to treat a pan-drug resistant *Achromobacter* species infection in a pediatric cystic fibrosis patient. Pediatr Pulmonol 55(11):2990-2994.

12. Comeau AM, Tétart F, Trojet SN, Prère MF, Krisch HM. 2007. Phage-antibiotic synergy (PAS): β-lactam and quinolone antibiotics stimulate virulent phage growth. PLoS One 2(8):e799.

13. Touchon M, Bernheim A, Rocha EPC. 2016. Genetic and life-history traits associated with the distribution of prophages in bacteria. ISME J 10(11):2744-2754.

14. López-Leal G, Camelo-Valera LC, Hurtado-Ramírez JM, Verleyen J, Castillo-Ramírez S, Reyes-Muñoz A. 2022. Mining of thousands of prokaryotic genomes reveals high abundance of prophages with a strictly narrow host range. mSystems 7(4):e0032622.

15. Chang TH, Pourtois JD, Haddock NL, Furkuawa D, Kelly KE, Amanatullah DF, Burgener E, Milla C, Banaei N, Bollyky PL. 2024. Prophages are infrequently associated with antibiotic resistance in *Pseudomonas aeruginosa* clinical isolates. bioRxiv 2024.06.02.595912.

16. Enault F, Briet A, Bouteille L, Roux S, Sullivan MB, Petit MA. 2016. Phages rarely encode antibiotic resistance genes: a cautionary tale for virome analyses. ISME J 2017 11(1):237-247.

17. Petty NK, Toribio AL, Goulding D, Foulds I, Thomson N, Dougan G, Salmond GPC. 2007. A generalized transducing phage for the murine pathogen *Citrobacter rodentium*. Microbiology (Reading) 153(Pt 9):2984-2988.

18. Canosi U, Lüder G, Trautner TA. 1982. SPP1-mediated plasmid transduction. J Virol 44(2):431-6.

19. Wilson GG, Young KKY, Edlin GJ, Konigsberg W. 1979. High-frequency generalised transduction by bacteriophage T4. Nature 280(5717):80-2.

20. Bondy-Denomy J, Qian J, Westra ER, Buckling A, Guttman DS, Davidson AR, Maxwell KL. 2016. Prophages mediate defense against phage infection through diverse mechanisms. ISME J 10(12):2854-2866.

21. Mavrich TN, Hatfull GF. 2019. Evolution of superinfection immunity in cluster A mycobacteriophages. mBio 10(3):e00971-19.

22. Chung IY, Sim N, Cho YH. 2012. Antibacterial efficacy of temperate phage-mediated inhibition of bacterial group motilities. Antimicrob Agents Chemother 56(11):5612-7.

23. Howard-Varona C, Hargreaves KR, Abedon ST, Sullivan MB. 2017. Lysogeny in nature: mechanisms, impact and ecology of temperate phages. ISME J 11(7):1511-1520.

24. Sauer RT, Rosslll MJ, Ptashnel M. 1982. Cleavage of the λ and P22 repressors by RecA protein. J Biol Chem 257(8):4458-62.

25. De Paepe M, Tournier L, Moncaut E, Son O, Langella P, Petit MA. 2016. Carriage of λ latent virus is costly for its bacterial host due to frequent reactivation in monoxenic mouse intestine. PLoS Genet 12(2):e1005861.

26. Goerke C, Koller J, Wolz C. 2006. Ciprofloxacin and trimethoprim cause phage induction and virulence modulation in *Staphylococcus aureus*. Antimicrob Agents Chemother 50(1):171-7.

27. Fatima R, Hynes AP. 2024. Temperate phage-antibiotic synergy is widespread, but varies by phage, host, and antibiotic pairing. mBio (in press).

28. Al-Anany AM, Fatima R, Hynes AP. 2021. Temperate phage-antibiotic synergy eradicates bacteria through depletion of lysogens. Cell Rep 35(8):109172.

29. Kamal F, Dennis JJ. 2015. *Burkholderia cepacia* complex phage-antibiotic synergy (PAS): antibiotics stimulate lytic phage activity. Appl Environ Microbiol 81(3):1132.

30. Al-Anany AM, Fatima R, Nair G, Mayol JT, Hynes AP. 2024. Temperate phage-antibiotic synergy across antibiotic classes reveals new mechanism for preventing lysogeny. mBio 15(6):e0050424.

31. Shkoporov AN, Clooney AG, Sutton TDS, Ryan FJ, Daly KM, Nolan JA, McDonnell SA, Khokhlova E V., Draper LA, Forde A, Guerin E, Velayudhan V, Ross RP, Hill C. 2019. The human gut virome is highly diverse, stable, and individual specific. Cell Host Microbe 26(4):527-541.e5.

32. Roszniowski B, Latka A, Maciejewska B, Vandenheuvel D, Olszak T, Briers Y, Holt GS, Valvano MA, Lavigne R, Smith DL, Drulis-Kawa Z. 2017. The temperate *Burkholderia* phage AP3 of the Peduovirinae shows efficient antimicrobial activity against *B. cenocepacia* of the IIIA lineage. Appl Microbiol Biotechnol 101(3):1203-1216.

33. Nale JY, Spencer J, Hargreaves KR, Trzepiński P, Douce GR, Clokie MRJ. 2016. Bacteriophage combinations significantly reduce *Clostridium difficile* growth *in vitro* and proliferation *in vivo*. Antimicrob Agents Chemother 60(2):968-81.

34. Mardiana M, Teh SH, Tsai YC, Yang HH, Lin LC, Lin NT. 2023. Characterization of a novel and active temperate phage vB_AbaM_ABMM1 with antibacterial activity against *Acinetobacter baumannii* infection. Sci Rep 13(1):11347.

35. Corsi AK. 2006. A biochemist's guide to *Caenorhabditis elegans*. Anal Biochem 359(1):1-17.

36. Darby C. 2005. Interactions with microbial pathogens. *In* WormBook: the online review of *C. elegans* biology. WormBook, Pasadena (CA).

37. Yao H, Xu A, Liu J, Wang F, Yao H, Chen J. 2022. Evaluation of *in vivo* antibacterial drug efficacy using *Caenorhabditis elegans* infected with carbapenem-resistant *Klebsiella pneumoniae* as a model host. Front Pharmacol 13:973551

38. Palacios-Gorba C, Pina R, Tortajada-Girbés M, Jiménez-Belenguer A, Siguemoto É, Ferrús MA, Rodrigo D, Pina-Pérez MC. 2020. *Caenorhabditis elegans* as an *in vivo* model to assess fucoidan bioactivity preventing *Helicobacter pylori* infection. Food Funct 11(5):4525-4534.

39. Son SJ, Park MR, Ryu SD, Maburutse BE, Oh NS, Park J, Oh S, Kim Y. 2016. Short communication: *in vivo* screening platform for bacteriocins using *Caenorhabditis elegans* to control mastitis-causing pathogens. J Dairy Sci 99(11):8614-8621.

40. Tan MW, Mahajan-Miklos S, Ausubel FM. 1999. Killing of *Caenorhabditis elegans* by *Pseudomonas aeruginosa* used to model mammalian bacterial pathogenesis. Proc Natl Acad Sci U S A 96(2):715-20.

41. Cezairliyan B, Vinayavekhin N, Grenfell-Lee D, Yuen GJ, Saghatelian A, Ausubel FM. 2013. Identification of *Pseudomonas aeruginosa* phenazines that kill *Caenorhabditis elegans*. PLoS Pathog 9(1):e1003101.

42. Kirienko N V., Kirienko DR, Larkins-Ford J, Wählby C, Ruvkun G, Ausubel FM. 2013. *Pseudomonas aeruginosa* disrupts *Pseudomonas aeruginosa* iron homeostasis, causing a hypoxic response and death. Cell Host Microbe 13(4):406-16.

43. Wang Y, Li X, Dance DAB, Xia H, Chen C, Luo N, Li A, Li Y, Zhu Q, Sun Q, Wu X, Zeng Y, Chen L, Tian S, Xia Q. 2022. A novel lytic phage potentially effective for phage therapy against *Burkholderia pseudomallei* in the tropics. Infect Dis Poverty 11(1):87.

44. Augustine J, Gopalakrishnan M V., Bhat SG. 2014. Application of ΦSP-1 and ΦSP-3 as a therapeutic strategy against *Salmonella Enteritidis* infection using *Caenorhabditis elegans* as model organism. FEMS Microbiol Lett 356(1):113-7.

45. Glowacka-Rutkowska A, Gozdek A, Empel J, Gawor J, Zuchniewicz K, Kozinska A, Debski J, Gromadka R, Lobocka M. 2019. The ability of lytic staphylococcal podovirus vB_SauP_phiAGO1.3 to coexist in equilibrium with its host facilitates the selection of host mutants of attenuated virulence but does not preclude the phage antistaphylococcal activity in a nematode infection model. Front Microbiol 9:3227.

46. Manohar P, Loh B, Elangovan N, Loganathan A, Nachimuthu R, Leptihn S. 2022. A multiwell-plate *Caenorhabditis elegans* assay for assessing the therapeutic potential of bacteriophages against clinical pathogens. Microbiol Spectr 10(1):e0139321.

47. Manohar P, Loh B, Turner D, Tamizhselvi R, Mathankumar M, Elangovan N, Nachimuthu R, Leptihn S. 2022. *In vitro* and *in vivo* evaluation of the biofilm-degrading *Pseudomonas* phage Motto, as a candidate for phage therapy. Front Microbiol 15:15:134496.

48. Hulsart-Billström G, Dawson JI, Hofmann S, Müller R, Stoddart MJ, Alini M, Redl H, El Haj A, Brown R, Salih V, Hilborn J, Larsson S, Oreffo ROC. 2016. A surprisingly poor correlation between *in vitro* and *in vivo* testing of biomaterials for bone regeneration: results of a multicentre analysis. Eur Cell Mater 31:312-22.

49. Kotani N, Ito K. 2023. Translatability of *in vitro* potency to clinical efficacious exposure: A retrospective analysis of FDA-approved targeted small molecule oncology drugs. Clin Transl Sci 16(8):1359-1368.

50. Heithoff DM, Barnes V L, Mahan SP, Fried JC, Fitzgibbons LN, House JK, Mahan MJ. 2023. Re-evaluation of FDA-approved antibiotics with increased diagnostic accuracy for assessment of antimicrobial resistance. Cell Rep Med 4(5):101023.

51. Radlinski L, Conlon BP. 2018. Antibiotic efficacy in the complex infection environment. Curr Opin Microbiol 42:19–24.

52. Tekippe M, Aballay A. 2010. *C. elegans* germline-deficient mutants respond to pathogen infection using shared and distinct mechanisms. PLoS One 5(7):e11777.

53. Udekwu KI, Parrish N, Ankomah P, Baquero F, Levin BR. 2009. Functional relationship between bacterial cell density and the efficacy of antibiotics. J Antimicrob Chemother 63(4):745-57.

54. Gutierrez A, Jain S, Bhargava P, Hamblin M, Lobritz MA, Collins JJ. 2017. Understanding and sensitizing density-dependent persistence to quinolone antibiotics. Mol Cell 68(6):1147-1154.e3.

55. González de Aledo M, Blasco L, Lopez M, Ortiz-Cartagena C, Bleriot I, Pacios O, Hernández-García M, Cantón R, Tomas M. 2023. Prophage identification and molecular analysis in the genomes of *Pseudomonas aeruginosa* strains isolated from critical care patients. mSphere 8(4):e0012823.

56. Ambroa A, Blasco L, López-Causapé C, Trastoy R, Fernandez-García L, Bleriot I, Ponce-Alonso M, Pacios O, López M, Cantón R, Kidd TJ, Bou G, Oliver A, Tomás M. 2020. Temperate bacteriophages (prophages) in *Pseudomonas aeruginosa* isolates belonging to the international cystic fibrosis clone (CC274). Front Microbiol 11:556706.

57. Fothergill JL, Mowat E, Walshaw MJ, Ledson MJ, James CE, Winstanley C. 2011. Effect of antibiotic treatment on bacteriophage production by a cystic fibrosis epidemic strain of *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 55(1):426-8.

58. James CE, Davies E V., Fothergill JL, Walshaw MJ, Beale CM, Brockhurst MA, Winstanley C. 2014. Lytic activity by temperate phages of *Pseudomonas aeruginosa* in long-term cystic fibrosis chronic lung infections. ISME J 9(6):1391-8.

59. Porta-de-la-Riva M, Fontrodona L, Villanueva A, Cerón J. 2012. Basic *Caenorhabditis elegans* methods: synchronization and observation. J Vis Exp (64):e4019.

60. Stiernagle T. 2006. Maintenance of *C. elegans*, p. 1–11. *In* WormBook: the online review of *C. elegans* biology. WormBook, Pasadena (CA).

Fig S1. *P. aeruginosa C0400* **can colonize and kill** *C. elegans*. **(a)** *C. elegans* intestinal colonization measured as cfu/mL after 24 h growth on *E. coli* OP50 or *P. aeruginosa* strains (PA14 or C0400). Data shown as mean ± SD of three biological replicates, each in single technical replicate. Values were compared using a one-way ANOVA with a Tukey multiple comparison test with *P ≤ 0.05, **P ≤ 0.01, and ns representing not significant. **(b)** Representative survival curve of worms exposed to OP50, PA14, or C0400 in slow killing assay. Curves were compared using the Log-rank (Mantel Cox) test with one pair compared at a time, *P ≤ 0.05 and **** P ≤ 0.0001. Each condition is 150-200 animals. The colour of the significance value represents the comparison performed, with black represent comparison to control and green to PA14.

Fig S2. Ciprofloxacin alone at 92 ug/mL could rescue life span of *P. aeruginosa* infected worms on standard *E. coli* OP50 diet after treatment. Survival curve of worms on OP50 after 4 h *P. aeruginosa* infection followed by 18 h phage Hali   ciprofloxacin treatment, **(a)** low dose 2 µg/mL and (b) high dose 92 µg/mL**.** Approximately 50 worms were added to each condition at the time of treatment. Independent biological replicates are shown for each concentration. Survival curves were compared using the Log-rank (Mantel Cox) test with one pair compared at a time, ns = not significant, * P ≤ 0.05, *** P ≤ 0.001, and **** P ≤ 0.0001. Phage alone and antibiotic alone were compared to the untreated, and tPAS to the phage alone and antibiotic alone, as denoted by colour of the significance value.

670 **Fig S3. Shorter infection prior to treatment does not provide enough resolution to measuring**
671 **bacterial colonization.** *C. elegans* intestinal colonization measured as log10 cfu/mL/worm after 4h
672 C0400 infection followed by 18 h treatment with phage Hali (1 x 10^9 pfu/mL) $\pm$ ciprofloxacin. UN
673 denotes untreated. Data shown as mean $\pm$ SD where each data point is a biological replicate performed in
674 technical triplicates with ~100 worms each.