TITLE: APOBEC3A, not APOBEC3B, drives deaminase mutagenesis in human gastric

epithelium

Running title: APOBEC3A, not APOBEC3B, is mutagenic in human cell

AUTHORS

Yohan An[1,10], Ji-Hyun Lee[2,10], Joonoh Lim[1,3,10], Jeonghwan Youk[1], Seongyeol Park[1], Ji-

Hyung Park[1,4], Kijong Yi[1,3], Taewoo Kim[1], Chang Hyun Nam[1], Won Hee Lee[1], Soo A Oh[3],

Yoo Jin Bae[5], Junehwak Lee[6], Jung Woo Park[6], Jie-Hyun Kim[5], Hyunki Kim[7], Hugo Snippert[8],

Bon-Kyoung Koo[2,9]*, and Young Seok Ju[1,3]*

1 Graduate School of Medical Science and Engineering, Korea Advanced Institute of

Science and Technology (KAIST), Daejeon 34141, Republic of Korea

2 Center for Genome Engineering, Institute for Basic Science, 55, Expo-ro, Yuseong-gu,

Daejeon, 34126, Republic of Korea

3 Inocras Inc., 6330 Nancy Ridge Drive, San Diego, CA 92121, United States

4 Department of Neurology, Seoul National University Bundang Hospital, Seongnam, Korea

5 Department of Internal Medicine, Division of Gastroenterology, Gangnam Severance

Hospital, Yonsei University College of Medicine, 20, Eonju-ro 63-gil, Gangnam-gu, Seoul

06229, Korea

6 Korea Institute of Science and Technology Information, Daejeon 34141, Republic of Korea

7 Department of Pathology, Yonsei University College of Medicine, Seoul, Republic of Korea

8 Oncode Institute, Center for Molecular Medicine, University Medical Center Utrecht,

Utrecht, the Netherlands

9 Department of Life Sciences, Pohang University of Science and Technology (POSTECH),

Pohang, 37673, Republic of Korea

10 These authors contributed equally

* Correspondence: ysju@kaist.ac.kr(Y.S.J.); koobk@ibs.re.kr(B.-K.K)

Corresponding authors

Bon-Kyoung Koo, PhD.

koobk@ibs.re.kr

Director

Center for Genome Engineering, Institute for Basic Science, 55, Expo-ro, Yuseong-gu,

Daejeon, 34126, Republic of Korea

Young Seok Ju, MD, PhD.

ysju@kaist.ac.kr

Professor

Graduate School of Medical Science and Engineering, Korea Advanced Institute of Science

and Technology (KAIST), Daejeon 34141, Republic of Korea

ABSTRACT

Cancer genomes frequently carry APOBEC (apolipoprotein B mRNA editing catalytic polypeptide-like)-associated DNA mutations, suggesting APOBEC enzymes as innate mutagens during cancer initiation and evolution. However, the pure mutagenic impacts of the specific enzymes among this family that are responsible for APOBEC-associated mutagenesis remain unclear in human normal cell lineage, particularly the comparative mutagenic activities of *APOBEC3A* and *APOBEC3B*. Here, we investigated the mutagenic contributions of these enzymes through whole-genome sequencing of human normal gastric organoid lines carrying doxycycline-inducible *APOBEC3A* or *APOBEC3B* cassettes. Our findings demonstrated that transcriptional *APOBEC3A* upregulation led to the acquisition of a massive number of genomic mutations in a few cell cycles. By contrast, *APOBEC3B* upregulation did not generate a substantial number of mutations in gastric epithelium. *APOBEC3B*-associated mutagenesis remained insignificant even after a combined inactivation of *TP53*. Based on the spectrum of acquired mutations after *APOBEC3A* upregulation, we further analyzed *APOBEC3A*-associated mutational signatures, encompassing indels mainly composed of 1bp deletions, characteristics of clustered mutations, and selective pressures operative on cells carrying the mutations. Our observations provide a clear foundation for understanding the mutational impact of APOBEC enzymes in human cells.

INTRODUCTION

Large-scale cancer genome studies have revealed various mutational processes in human somatic cells (Alexandrov et al. 2020). According to the COSMIC database (Sondka et al. 2024), APOBEC-associated mutations are detected in ~75% of human cancer types, including bladder transitional cell carcinoma (98%, 381 of 389 samples), breast adenocarcinoma (83%; 759 of 915 samples), and stomach adenocarcinoma (21%; 101 of 486 samples). In addition, APOBEC-associated mutational activity is also observed in non-neoplastic normal cells, particularly within the epithelium of the bladder, bronchus, and small intestine (Lawson et al. 2020; Wang et al. 2023; Yoshida et al. 2020).

APOBEC enzymes are well known for their role in innate immunity, where they destabilize pathogens by inducing DNA and RNA deamination (Vieira and Soares 2013). Recently, they have gained wide recognition as mutagens that also affect host genomes in breast cancer cells (Nik-Zainal et al. 2012b). APOBEC-associated DNA mutations are characterized by predominantly C>T and C>G base substitutions enriched in TpCpN context (with the mutated cytosine is underlined; Nik-Zainal et al. 2012a). The mutational spectrum has been deconvoluted by two mutational signatures (SBS2 and SBS13), which are consistent across different cancer types (Alexandrov et al. 2020). From the characteristic sequence-context dependency of the mutations, their mutagenic origin was associated with APOBEC enzyme activity (Nik-Zainal et al. 2012b; Lawrence et al. 2013; Roberts et al. 2013). Of the 11 APOBEC family genes in the human genome, *APOBEC3A* (**A3A**) and *APOBEC3B* (**A3B**) have been suggested as major potential contributors to the mutations attributable to SBS2 and SBS13 in most human cell types (Chan et al. 2015; Roberts et al. 2013). In the small intestine, *APOBEC1* was also proposed as a main mutagenic enzyme (Wang et al. 2023).

The mutagenic potential of A3A and A3B has been investigated through a range of human cancer cell lines (Burns et al. 2013; Carpenter et al. 2023; Petljak et al. 2022). APOBEC-associated mutagenic activity was suggested to be episodic in cancer cell line models

(Petljak et al. 2019). In addition, non-human model systems, such as *Mus musculus,*

Saccharomyces cerevisiae, and *Gallus gallus domesticus*-derived cell lines were also

utilized for validating the mutagenic impact of APOBEC enzymes (Durfee et al. 2023;

DeWeerd et al. 2022; Naumann et al. 2023; Law et al. 2020; Chan et al. 2015). However,

these systems are suboptimal to understand the mutagenicity of APOBEC enzymes in

human normal cells, as these models harbor confounding factors, such as additional

oncogenic alterations, or non-human genomic backgrounds.

In this regard, we established human gastric organoid lines carrying doxycycline-inducible

APOBEC cassettes to evaluate the pure mutagenic activity of A3A and A3B in human

normal cells. Combining single-cell cloning and whole-genome sequencing (**WGS**; Jager et

al. 2019; Pleguezuelos-Manzano et al. 2020; Youk et al. 2024) as well as duplex DNA

sequencing (Hoang et al. 2016), we investigated mutations attributable to A3A and A3B

activation in human normal cells.

RESULTS

Doxycycline-inducible *APOBEC3A* and *APOBEC3B* in normal gastric organoids

To control the transcription of A3A and A3B in non-neoplastic human cells, we established

human gastric organoid lines with doxycycline-inducible expression of either A3A or A3B

(referred to as hGO$_{iA3A}$ and hGO$_{iA3B}$, respectively). In these lines, doxycycline treatment

simultaneously triggers the upregulation of (1) mCherry fluorescence, and (2) either the A3A

or A3B enzyme (**Fig. 1A**). To this end, we utilized *PiggyBac* transposon system (Woodard

and Wilson 2015; Wilson et al. 2007) to integrate two independent constructs into the

genome of gastric organoids: (1) a cassette expressing rtTA and hygromycin B resistance

protein (CMV-rtTA-HygR), and (2) a cassette expressing APOBEC enzymes and

fluorescence protein (TRE-APOBEC (A3A or A3B)-IRES-mCherry). Successfully engineered

organoids were selected through hygromycin treatment and subsequently constructed into

clonal lines. WGS confirmed their clonality (**Supplemental Fig. S1**), copy number and

genomic positions of integrated cassettes (**Supplemental Tables S1, S2**), and correctness

of the sequence and reading frame of APOBEC gene in the inserted constructs in the

hGO$_{iA3A}$ and hGO$_{iA3B}$ lines (**Supplemental Fig. S2**).

Figure 1. Overview of conditional APOBEC (APOBEC3A or APOBEC3B)

overexpression models with human gastric organoids.

(A) Experimental design of the study. Schematic illustration of genetic engineering, cloning,

and sequencing (whole-genome sequencing, duplex sequencing, and bulk RNA-seq) is

shown. Dox, doxycycline.

(B) Changes of morphology and mCherry fluorescence during 48 hours with 0.1 µg/ml

doxycycline treatment. (top) hGO$_{iA3A}$ lines and (bottom) hGO$_{iA3B}$ lines. Scale bars represent

1mm.

(C) Expression levels of APOBEC3A (A3A) or APOBEC3B (A3B) in each line following 0.1

µg/ml and 3 µg/ml doxycycline treatment for 48 hours. (left) hGO$_{iA3A}$ lines (n=3 in each

condition) and (right) hGO$_{iA3B}$ lines (n=3 in each condition). Data are presented as mean ±

95% confidence intervals. Statistical significance was determined using a t-test: *$p < 0.05$,

**$p < 0.005$, ***$p < 0.0005$, ****$p < 0.00005$.

(D) Changes of viability of each line following 0.1 µg/ml and 3 µg/ml doxycycline treatment

for 48 hours. (left) hGO$_{iA3A}$ lines (n=3 in each condition) and (right) hGO$_{iA3B}$ lines (n=3 in each

condition). Data are presented as mean ± 95% confidence intervals.

(E) Differentially expressed genes upon APOBEC (A3A or A3B) overexpression following 0.1

µg/ml doxycycline treatment for 48 hours.

(F) Subcellular localization of overexpressed A3A and A3B following 3 µg/ml doxycycline

treatment for 48 hours in each line. (top) hGO$_{iA3A}$ line and (bottom) hGO$_{iA3B}$ line. Scale bars

represent 50µm.

(G) γ-H2AX foci following 3 µg/ml doxycycline treatment for 48 hours in each line. (left)

hGO$_{iA3A}$ line and (right) hGO$_{iA3B}$ line. Scale bars represent 50µm.

Upon doxycycline treatment, upregulation of mCherry fluorescence and APOBEC transcripts

were clearly observed within 48 hours (**Fig. 1B**). Here, we treated 0.1 µg/ml and 3 µg/ml of

doxycycline for low- and high-level induction, respectively. Bulk RNA-seq analysis revealed

that gene expression levels of A3A or A3B ranged from 400 to 6,000 transcripts per millions

(**TPMs**), representing a several hundred-fold increase compared to background levels (**Fig.**

1C; Supplemental Fig.S3). Of note, transcripts from the endogenous *A3A* and *A3B* genes

accounted for only ~0.2% of total A3A or A3B expression levels following induction,

indicating minimal contribution of the innate APOBEC gene copies (**Supplemental Table**

S3). The range of expression levels was comparable to the transcriptional levels in single

cancer cells with APOBEC expression bursts in many cancer types, including lung

adenocarcinoma, head-and-neck squamous cell carcinoma, triple negative breast cancer,

esophageal adeno- and squamous cell carcinomas (**Supplemental Fig. S4;** Karaayvaz et al.

2018; Maynard et al. 2020; Puram et al. 2017; Wu et al. 2018). Unfortunately, the levels in

gastric adenocarcinoma cells are uncertain, due to the lack of public single-cell

transcriptomes through Smart-seq in the cancer type.

Following APOBEC induction, the cell viability of the lines was substantially compromised,

suggesting a negative impact of APOBEC enzymes on cell survival (**Fig. 1D**). Differentially

expressed genes after APOBEC induction included genes related to cell cycle (e.g.,

CDKN1A and *CDC20*) and immune response (e.g., *IL8* and *CCL20*) in both the hGO$_{iA3A}$ and

hGO$_{iA3B}$ models (**Fig. 1E;** Cazzalini et al. 2010; Amador et al. 2007; Matsushima et al. 2022;

Wu et al. 2007).

We also observed the nuclear localization of A3A and A3B enzymes following protein

translation (**Fig. 1F; Supplemental Fig. S5**). The increased number of γ-H2AX foci after

APOBEC upregulation confirmed that the APOBEC enzymes led to DNA damage, such as

replication stall or double-strand breaks (**DSBs**) as previously observed (**Fig. 1G;**

Supplemental Fig. S5; Burns et al. 2013; Green et al. 2016).

APOBEC3A, not APOBEC3B, induces genomic mutations

To assess the mutagenic impacts of A3A and A3B, we investigated acquired mutations after

48 hours from doxycycline treatments in the hGO$_{iA3A}$ and hGO$_{iA3B}$ lines. Interestingly, a

substantial number of base substitutions, whose spectrum is concordant with a linear

combination of SBS2 and SBS13, were detected in the hGO$_{iA3A}$ clones (**Fig. 2A, F;**

Supplemental Tables S4, S5). On average, the hGO$_{iA3A}$ lines with 0.1 µg/ml and 3 µg/ml

doxycycline exhibited 267 and 2,448 base substitutions genome-wide, respectively (0.1

µg/ml: 95% CI, 107-505; 3 µg/ml: 95% CI, 1,052-3,708; **Fig. 2A, B; Supplemental Table**

S5). By contrast, the hGO_{iA3B} clones showed a negligible number of APOBEC-associated

mutations (0.1 µg/ml: 95% CI, 0-1; **Fig. 2A, D; Supplemental Table S5**).

Figure 2. Mutational impact of APOBEC-associated single base substitutions in DNA

and RNA.

(A) Mutational burdens of single nucleotide variants (SNVs) after overexpression of

APOBEC (A3A or A3B) measured by whole-genome sequencing of the clones and duplex

DNA sequencing classified. (left) hGO_{iA3A} linesand (right) hGO_{iA3B} lines.

(B) Mutational burdens of APOBEC-associated SNVs in the hGO_{iA3A} and TP53KO-hGO_{iA3A}

clone sequencing**.** The number of A3A-associated SNVs (SBS2+SBS13) in hGO_{iA3A} and

TP53KO-hGO_{iA3A} clones under each condition. Statistical significance was determined using

a t-test: *$p < 0.05$, **$p < 0.005$, ***$p < 0.0005$,****$p < 0.00005$.

(C) The corrected number of A3A-associated SNVs (SBS2+SBS13) in BotSeqS results for

hGO_{iA3A} lines under each doxycycline treatment condition. Black lines indicate confidence

intervals calculated with *Poisson* distribution.

(D) Mutational burdens of APOBEC-associated SNVs in the hGO_{iA3B} and TP53KO-hGO_{iA3B}

clone sequencing**.** The number of A3A-associated SNVs (SBS2+SBS13) in hGO_{iA3B} and

TP53KO-hGO_{iA3B} clones under each condition. Statistical significance was determined using

a t-test: *$p < 0.05$, **$p < 0.005$, ***$p < 0.0005$,****$p < 0.00005$.

(E) The corrected number of A3B-associated SNVs (SBS2+SBS13) in BotSeqS results for

hGO_{iA3B} lines under each doxycycline treatment condition. Black lines indicate confidence

intervals calculated with *Poisson* distribution.

(F) Mutational burdens and spectra of APOBEC-associated SNVs in each experimental

condition. (left) hGO_{iA3A} lines and (right) hGO_{iA3B} lines.

(G) The number of C>U RNA editing counts in bulk RNA-seq in hGO_{iA3A} lines. n=3 in each

condition.

(H) The spectra of RNA editing in trinucleotide contexts in hGO_{iA3A} lines

(I) The number of C>U RNA editing counts in bulk RNA-seq in hGO_{iA3A} lines. n=3 in each

condition.

(J) The spectra of RNA editing in trinucleotide contexts in hGO$_{iA3B}$ lines.

Then, we assessed the potential for detection bias in our single-cell cloning system, which

could underestimate the mutational impact of APOBEC enzymes. Here, if heavily damaged

cells exhibit reduced proliferative potential, they may fail to expand into clones and thus be

excluded from subsequent genome sequencing. To bypass the cloning step and directly

detect mutations, we employed duplex DNA sequencing technique, enabling the capture of

subclonal mutations across the entire cell population, including those present in single cells

that could not proliferate further (Hoang et al. 2016).

After 48 hours of doxycycline treatment, the duplex DNA sequencing technique revealed a

higher mutational burden of APOBEC-associated base substitutions in hGO$_{iA3A}$ (0.1 µg/ml:

95% CI, 4,590-6,038; 3 µg/ml: 95% CI, 9,316-10,077; **Fig. 2A**; **Supplemental Table S6**).

We observed a time-dependent increase in A3A-associated mutational burden following

doxycycline treatment (**Fig. 2C**). Our observation confirms that (1) A3A is a potent mutator

enzyme, and (2) its quantitative mutational impact may be underestimated in clonal analyses.

In contrast, APOBEC-associated base substitutions remained minimal in cell populations

exposed to A3B (0.1 µg/ml: 95% CI, 136-352; 3 µg/ml: 95% CI, 79-133; **Fig. 2A**;

Supplemental Table S6). Of note, we did not detect A3B-associated mutations even at

earlier time points, likely prior to cellular clearance (**Fig 2E**), suggesting that the absence of

mutations in hGO$_{iA3B}$ is unlikely to be due to negative selection against hypermutated cells.

Despite the minimal mutational impact observed in the organoids, the A3B enzyme was fully

functional, exhibiting strong cytosine deamination activity *in vitro*. First, cell lysates from

hGO$_{iA3B}$ induced a substantial number of C>T artifacts on extracellular double-strand DNA

fragments, particularly in the 5' head region (~50bp), with a clear positive correlation to the

level of doxycycline concentration (**Supplemental Fig. S6**). As previously reported (Abascal

et al. 2021), such artifacts likely arise at single-stranded overhangs of DNA fragments during

cell lysis or early stages of library preparation, driven by APOBEC activity in the lysates.

Second, recombinant A3B enzymes synthesized from the A3B sequence introduced C>T

mutations in cell-free denatured genomic DNA fragments, with nearly all (~99%)

unmethylated (non-CpG) cytosines converted to thymines in approximately 60% of the

fragments likely exposed to A3B (**Fig. 3A, B**). In contrast, the remaining ~40% of the

fragments did not show any C>T conversion, suggesting these were not engaged by A3B

enzymes. Finally, the increase of γ-H2AX foci in the hGO$_{iA3B}$ following doxycycline induction

(**Fig. 1G; Supplemental Figure S5**) also indicates DNA damage consistent with A3B activity.

Collectively, our findings suggest that while A3B is enzymatically active, gastric organoids

efficiently repair A3B-induced lesions or possess intrinsic mechanisms that suppress A3B-

associated mutation in the nucleus, thus limiting its mutagenic impact in the cellular DNA.

Figure 3. Deamination activity of APOBEC3B *in vitro.*

(A) Sequences in randomly selected 100 read pairs in APOBEC3B exposed cell-free

denatured genomic single stranded DNA for 60 seconds.

(B) Deamination rate of unmethylated cytosines in A3B-exposed cell-free DNA fragments.

CTRL, negative control, 30s, DNA exposed for 30 seconds, 60s, DNA exposed for 60

seconds.

To further investigate whether APOBEC-associated mutagenesis depends on the

inactivation of tumor suppressor genes, we generated TP53-knockout derivatives of the

hGO$_{iA3A}$ and hGO$_{iA3B}$ lines (hereafter referred to as TP53KO-hGO$_{iA3A}$ and TP53KO-hGO$_{iA3B}$,

respectively; **Fig. 1A**). From the TP53KO-hGO$_{iA3A}$ lines, we observed a similar burden of

APOBEC-associated mutations to what was observed in the hGO$_{iA3A}$ lines (0.1 μg/ml: 95%

CI, 785-1,917; 3 μg/ml: 95% CI, 152-3,091; **Fig. 2A, B; Supplemental Table S5**). Similarly,

the TP53KO-hGO$_{iA3B}$ lines also showed results consistent with the hGO$_{iA3B}$ lines, with a lack

of APOBEC-associated mutations (0.1 μg/ml: 95% CI, 11-19; 3 μg/ml: 95% CI, 2-12; **Fig. 2A,**

D; **Supplemental Table S5**). Taken together, these findings imply that the inactivation of

TP53 does not promote APOBEC-associated mutagenesis in gastric organoids.

Despite the increased γ-H2AX foci following A3A or A3B overexpression (**Fig. 1G;**

Supplemental Figure S5), which indicates DNA damage and may reflect double-strand

breaks in DNA, we did not observe a corresponding increase in structural variations (**SVs**;

Supplemental Table S5). The absence of APOBEC-induced SVs was also observed in the

TP53KO-hGO$_{iA3A}$ and TP53KO-hGO$_{iA3B}$ lines. We speculate that APOBEC-induced DSBs or

replication stalls were mostly repaired correctly within the normal cells via the error-free

homologous recombination pathway in non-neoplastic gastric cells.

Both APOBEC3A and APOBEC3B induce C>U RNA editing

Despite their differing mutagenic impact on DNA, both A3A and A3B exhibited deaminase

activity on RNA, leading to C>U RNA editing events in the clones. In both the hGO$_{iA3A}$ and

hGO$_{iA3B}$ lines, we observed an approximately 8-fold and 30-fold increase in C>U RNA editing

following treatment with 0.1 μg/ml and 3 μg/ml doxycycline for 48 hours, respectively,

compared to baseline levels (**Fig. 2G, I**). These findings indicate that both constructs

possess potent enzymatic activity.

The spectra of C>U RNA-editing deviated from SBS2 for DNA mutations and varied between

A3A and A3B (**Fig. 2H, J**). Specifically: (1) both A3A and A3B exhibited reduced specificity

for the UpCpU context (equivalent to TpCpT in DNA); (2) A3A displayed enhanced specificity

for the UpCpG context (equivalent to TpCpG in DNA); and (3) A3B showed decreased

specificity for the UpCpN context (equivalent to TpCpN in DNA). Further *de novo* extraction

of RNA editing signatures revealed distinct context-specific characteristics of the A3A and

A3B-associated RNA-editing (**Supplemental Fig. S7**), in line with previous reports

(Martínez-Ruiz et al. 2023; Fixman et al. 2024; Alonso de la Vega et al. 2023). Expansion of

context analysis up to pentanucleotide sequences revealed that A3A-mediated RNA editing

were preferentially enriched in ApUpCpApN (equivalent to ApTpCpApN in DNA) and

ApUpCpGpN (equivalent to ApTpCpGpN in DNA) contexts (**Supplemental Fig. S8**),

whereas A3B-mediated editing did not show such an enrichment.

To further investigate the RNA editing activity of A3A and A3B, we systematically compared

the recurrent sites for C>U RNA editing of each enzyme. Among the identified sites, 28,500

were commonly edited by both A3A and A3B, while 558 sites were unique to A3A and

13,078 were specific to A3B (**Supplemental Fig. S9A**). Sequence context analysis revealed

that ~92% of A3A-specific editing events occurred within the UpCpN (equivalent to TpCpN in

DNA) context, compared to ~69% for A3B-specific sites, suggesting a broader sequence

tolerance by A3B (**Supplemental Fig. S9B**). We next examined the local RNA secondary

structures at editing sites. Consistent with previous literature (Jalili et al. 2020), A3A

preferentially targeted cytosines positioned within defined loop structures, with a clear

dependency on both loop size and cytosine position. A3B exhibited similar structural

preference; however, the magnitude of these biases was markedly attenuated, suggesting

weaker structural constraints on A3B-mediated editing (**Supplemental Fig. S9C**).

Characteristics of APOBEC-associated mutations

Using the pure A3A-mediated mutational profiles from hGO_{iA3A} lines, we further examined

the characteristics of A3A-associated mutational signatures. A3A-associated mutations in

the TpCpA context were 2.7 times more abundant following pyrimidine bases (YpTpCpA)

compared to purine bases (RpTp$\underline{C}$pA; **Fig. 4A**) as previously reported (Sanchez et al. 2024;

Chan et al. 2015). Of note, the RpTpCpA sequence context has been reported as a

preferred motif for A3B-associated mutations in the yeast and *in vitro* model (Chan et al.

2015; Sanchez et al. 2024). The YpTpCpA preference in the gastric organoids closely

mirrors the enrichment level observed in human cancer tissues carrying APOBEC-induced

hypermutations (YpTpCpA/RpTpCpA = 2.5), indicating that A3A could be a key enzyme of

the APOBEC-associated hypermutations in most human cancers.

Figure 4. Characteristics of A3A-associated mutational signatures.

(A) Context preference of A3A between YpTp$\underline{C}$pA and RpTp$\underline{C}$pA context in hGO$_{iA3A}$ lines

and APOBEC-associated mutations in hypermutant cancer samples. Only PCAWG cancer

samples with a combined APOBEC-associated mutational burden (SBS2 + SBS13) greater

than 5,000 were selected (n=63) among eight cancer types with a high prevalence of

APOBEC mutational activity. lung adenocarcinoma (n=15), breast adenocarcinoma (n=12),

bladder urothelial carcinoma (n=11), head and neck squamous cell carcinoma (n=13), lung

adenocarcinoma (n=6), uterine corpus endometrial carcinoma (n=3), esophageal

adenocarcinoma (n=2), and stomach adenocarcinoma (n=1). Dashed black line, expected;

orange line, hGO$_{iA3A}$; dark brown line, cancer.

(B) Correlation between A3A-associated base substitutions and ID9 contributing indels.

(C) Associations between A3A-associated (SBS2 and SBS13) and age-associated (SBS5

and SBS40) base substitutions among hGO$_{iA3A}$ lines and TP53KO-hGO$_{iA3A}$ lines.

(D) Gene expression changes of *POLH* (translesion DNA polymerase) after A3A induction in

hGO$_{iA3A}$ and TP53KO-hGO$_{iA3A}$ lines.

Further, in line with previous reports from cancer (DeWeerd et al. 2022), indels attributable

to the COSMIC ID9 signature (Sondka et al. 2024) showed a suggestive positive correlation

with the number of A3A-associated base substitutions in hGO$_{iA3A}$ clones (**Fig. 4B**; p-value =

0.051). The relative rate of ID9-associated indels was one for every 333 SBS2/13 base

substitutions in the gastric organoids.

Interestingly, base substitutions attributable clock-like mutational signatures (SBS5 and

SBS40) also demonstrated a positive correlation with the burden of A3A-associated base

substitutions (SBS2 and SBS13; **Fig. 4C**). This suggests that A3A-induced genomic damage

indirectly promotes error-prone DNA repair processes across the genome. However, this

association was absent in hGO$_{iA3A}$ clones carrying *TP53* truncating mutations (**Fig. 4C**). We

speculate that the difference stems from differential DNA repair pathways according to the

activities of TP53 (Kim et al. 2016). Notably, among the polymerases encoded in the human

genome, increased transcription of DNA polymerase eta (*POLH*), which is involved in

translesion synthesis (TLS; Choi and Pfeifer 2005; Delbos et al. 2005) was exclusively

observed in clones without TP53 knockout after induction (**Fig. 4D; Supplemental Fig. S10**).

Of note, *REV1*, a component of the translesion synthesis (TLS) pathway, was implicated in

the generation of SBS5 and SBS40 (Petljak et al. 2022). These findings collectively suggest

that multiple TLS polymerases may contribute to the mutational processes underlying SBS5

and SBS40.

Clusters of APOBEC-associated mutations

In cancer genomes, APOBEC-induced localized hypermutation events are frequently

observed (Nik-Zainal et al. 2012a; The ICGC/TCGA Pan-Cancer Analysis of Whole

Genomes Consortium 2020). In our clones, approximately 5% of the 29,650 acquired single-

nucleotide variants were clustered within 1 kbp, a frequency 100 times higher than random

chance (**Fig. 5A**). Such clustered mutation events can be classified into *omikli* and *kataegis*

(usually composed of 2-3 and >3 mutations, respectively) based on statistical models, not by

absolute distance (Bergstrom et al. 2022b). A3A-associated clustered mutation events were

distributed across 615 *omikli* and 109 *kataegis* events. The observed *kataegis* events ranged

from 4 to 22 base substitutions (median = 5). Contrary to the ~36% of *kataegis* events in

cancers reportedly being observed with nearby SVs (The ICGC/TCGA Pan-Cancer Analysis

of Whole Genomes Consortium 2020; Bergstrom et al. 2022b; Nik-Zainal et al. 2012a),

rearrangement-associated hypermutations were rare in our data (2.8%; 3 out of 109 events).

Similarly, the *kataegis* in our clones were independent of other known *kataegis*-shaping

genomic events, such as anaplastic DNA bridge (Maciejowski et al. 2020) and

extrachromosomal DNA (ecDNA; Bergstrom et al. 2022b). Therefore, our results indicate

that the clustered mutations found in our clones were driven by pure activity of A3A.

The relative frequencies of *omikli* and *kataegis* remained consistent at 2.4% (95% CI: 1.7-

3.1%) and 0.4% (95% CI: 0.22-0.58%) of all A3A-associated mutation events, respectively,

with each isolated and clustered mutations counted as a single event (**Fig. 5B**;

 Supplemental Table S7). These ratios were more or less constant in clones, regardless of

the A3A expression levels and *TP53* mutational status. Interestingly, the frequencies of

omikli and *kataegis* were comparable to the SV-unrelated *omikli* and *kataegis* frequencies in

cancers (**Fig. 5B**; **Supplemental Table S7**).

Figure 5. Landscape of APOBEC3A-associated clustered mutation.

(A) Observed-to-expected ratio of the intermutational distances between substitutions in

each clone. Data are presented as mean ± standard error.

(B) Frequency of clustered mutation events (*omikli* and *kataegis*) among A3A-associated

base substitutions in clones and cancer genomes.

(C) Proportion of classic (in TpCpN context) and non-classical mutations (in non-TpCpN and

NpTpN) in SBS2, SBS13, and *kataegis* regions in the clones.

(D) Comparison of expected and observed SNV counts in the non-TpCpN context within

kataegis regions.

(E) Schematic representation of strand-switching *kataegis*.

(F) Frequency of strand-switching *omikli* and *kataegis.*

(G) A complex strand-switching *kataegis* with five switches found in clone A3A_1st_C3_3µg-

5.

(H) Enrichments of minor strand mutations in the TpCpN context in strand-switching *kataegis*

events.

Within *kataegis* regions, cytosine alterations did not always occur in the TpCpN context. We

observed that ~7% of cytosine substitutions occurred in non-TpCpN context (39 out of 556

mutations; **Fig. 5C**), approximately 2-fold higher than expectation based on SBS2 and

SBS13 signatures (7.00% vs. 3.33%; chi-square test, p<0.005). Our data indicated these

non-TpCpN mutations arose simultaneously with the *kataegis* for two reasons: (1) Non-

TpCpN mutations were 267-fold more abundant than observed in the background regions

(**Fig. 5D**), making contamination by *kataegis*-unrelated mutations unlikely; and (2) Non-

TpCpN mutations within *kataegis* regions were completely phased with other classical

TpCpN mutations on the same chromosome (27 out of 27), where 50% was expected for

random contaminating mutations. This suggests that DNA repair mechanisms for cytosine

deamination in non-TpCpN contexts are less accurate during the clustered mutagenesis

compared to isolated deamination events.

Of the 615 *omikli* and 106 *kataegis* events (excluding 3 *kataegis* events associated with

rearrangements*)*, approximately 5% (35 *omikli* and 5 *kataegis* events) exhibited strand-

switching of the mutated cytosines between parental and daughter strands during replication

(**Fig. 5E, F**). In one *kataegis* event composed of 9 base substitutions (in clone

A3A_1st_C3_3µg-5), we observed five strand-switching events (**Fig. 5G**). Phasing analysis

suggested that clustered mutations across both DNA strands originated from single mutation

events (26 out of all 26 informative events). Additionally, the mutational spectrum for the

mutations in the minorly contributing strand was predominantly composed of mainly C>T or

C>G mutations in the TpCpN context (62%; 13 out of 21 mutations; **Fig. 5C**), implying

APOBEC-associated mutations. Besides, the rate of TpCpN mutations in the minor strand

within strand-switching *kataegis* regions was 124-fold higher compared to outside of the

strand-switching *kataegis* region (**Fig. 5H**). Although the underlying mechanism is unclear,

strand-switching of the A3A enzyme should be possible when generating clustered

mutations during replication.

Epigenetic contexts associated with A3A-associated mutations

Mutational processes are often influenced by the epigenetic contexts of the genome (Otlu et

al. 2023). Of the 14 genomic features examined, three features (replication timing, local

transcription level, and H3K27me3) showed potential associations with local A3A-associated

mutational burdens (**Fig. 6A**). Of the three features, the replication timing and local gene

transcription level demonstrated consistent trends correlating with A3A-associated mutation

rates (**Supplemental Table S8**).

Figure 6

Figure 6. Genomic and epigenomic distribution of APOBEC3A-associated mutations.

(A) Correlation between epigenetic markers and A3A-associated substitutions.

(B) Fold change of mutation rates of A3A-associated SNVs according to the replication

timing of the genomic region. Data are presented as mean ± 95% confidence intervals.

(C) Mutation rates depending on the DNA strands during replication (leading and lagging

strands). Statistical significance was determined using chi-square test: ****p<0.00005.

(D) Fold change of mutation rates of A3A-associated SNVs according to the local

transcription levels in the genic region. Data are presented as mean ± 95% confidence

intervals.

(E) Mutation rates depending on the DNA strands during transcription (transcribed and

untranscribed strands). Statistical significance was determined using a chi-square test:

***p<0.0005.

(F) Mutation rates within sub-genic regions (5`UTR, intron, protein coding sequences (CDS),

and 3`UTR) in (left) hGO$_{iA3A}$ clones and (right) TP53KO-hGO$_{iA3A}$ clones. Red dashed line,

average genome-wide mutation rate.

For the replication timing, the latest-replicating regions showed a 1.26-fold higher rate of

A3A-associated mutation compared to the earliest-replicating regions (**Fig. 6B**) as previously

reported (Kazanov et al. 2015). This may be attributed to the DNA repair mechanisms,

including base excision repair, which are particularly active in early-replicating regions

consisting of open chromatin (Amouroux et al. 2010; Rhind and Gilbert 2013). Further, the

A3A-associated mutation rate in the lagging strand of DNA replication was 1.26-fold higher

compared to the leading strand (**Fig. 6C**), which presumably originated from more frequent

exposure of single-stranded DNA (ssDNA) induced by Okazaki fragments in the lagging

strand (Wu et al. 2020), similar with previous elucidation (Hoopes et al. 2016).

A similar pattern was observed in gene transcription. Mutation rates in genic regions were

correlated with expression levels (**Fig. 6D**), showing a 1.37-fold higher mutation rate in

actively transcribed genomic regions compared to silent genic regions consistent with the

previous study (Kazanov et al. 2015; Nordentoft et al. 2014). Of note, the lagging strand and

highly transcribed genes tend to be more frequently single-stranded than the leading strand

and silent genes (Okazaki et al. 1968; Gnatt et al. 2001), which may make them more

susceptible to A3A-induced DNA damage, respectively. The results demonstrated that DNA

regions with frequent ssDNA exposure have a higher chance of being damaged by A3A. In

line with the observation, the non-transcribed genic strand was mutated 1.13-fold more

frequently than the transcribed strand (**Fig. 6E**), consistent with the previous report (Saini et

al. 2017).

Compared with non-coding sequences, protein-coding sequences showed much lower

mutation rates, at 0.79-fold the genome average, suggesting a selective pressure against

mutations that could alter amino-acid-changing mutations (**Fig. 6F**). Interestingly, in TP53-

inactivated clones (TP53KO-hGO$_{iA3A}$ clones), mutation rates in protein-coding regions

increased to 0.834-fold of the genome average, potentially due to reduced negative selection

pressures in the absence of functional *TP53*.

DISCUSSION

Our study clearly demonstrated the qualitative and quantitative mutational impact of A3A and

A3B in human non-neoplastic cells using a gastric organoid culture system. The gastric

organoid model was particularly suitable for this study based on three key criteria: (1) the

biological relevance, supported by the presence of significant APOBEC-associated

mutations in corresponding gastric cancer tissues, (2) the robust proliferative capacity under

culture conditions, enabling experimental feasibility, and (3) the availability of standardized

protocols for genetic manipulation. Although APOBEC-associated mutations are also

frequently observed in breast and lung cancers, we did not include normal breast or lung

epithelial cell models for this study due to their limitations in the cell proliferation capacity in

organoid culture condition and genetic engineering compatibility.

When A3A was induced by 3 µg/ml of doxycycline in hGO$_{iA3A}$ lines, transcription was

activated for ~2-3 days, reaching peak gene expression levels of ~800 TPM. Under these

conditions, we detected ~2,500 A3A-induced base substitutions in the proliferative clones,

suggesting that ~1,000 base substitutions could be acquired in a day. As demonstrated by

duplex sequencing, the mutational burden is likely even higher in less proliferative cells.

These findings support the notion that a transcriptional burst of A3A can also lead to a

massive number of mutations in normal gastric epithelium, consistent with previous report in

cancer (Petljak et al. 2019).

Our models indicated increased mutational burdens of SBS5 and SBS40 in proportion to the

overall burden of A3A-associated mutations. These signatures were significantly enriched in

late-replicating regions and correlated with multiple epigenomic features, including

replication timing and transcriptional activity, consistent with previous reports (**Supplemental**

Fig. S11; Sondka et al. 2024). Together, these findings support a model in which A3A-

induced DNA damage perturbs global DNA repair processes, thereby promoting the

accumulation of SBS5 and SBS40 mutational signatures.

Previous studies have highlighted the mutagenic potential of A3B across various model

systems (Dananberg et al. 2024; Carpenter et al. 2023; Durfee et al. 2023; Chan et al. 2015).

In contrast, a recent study reported only a modest reduction in APOBEC-associated

mutations after A3B inactivation in human cancer cell lines, suggesting a more limited role

for A3B in mutagenesis (Petljak et al. 2022). To our knowledge, our study is the first to

directly assess the mutagenic activity of A3B in human normal cells. Our findings indicate

that A3B alone does not act as a major mutator, at least in the context of human normal

gastric epithelium.

Understanding the potential cell-type specificity of the mutagenic activity of A3B, as well as

the underlying mechanisms, represents an important direction for future research. Applying

similar experimental approaches to organoids derived from a wider range of tissue types, in

both normal and neoplastic contexts, will be essential to clarify the broader relevance of A3B

activity. Furthermore, emerging technologies that enable simultaneous whole-genome and

transcriptome sequencing at the single-cell level will be invaluable for resolving the timing,

cellular context, and consequences of APOBEC-associated mutagenesis in a tissue- and

model-independent manner.

Although this study successfully evaluated the mutagenic activity of A3A and A3B in human

gastric epithelium, it has a few technical limitations that warrant consideration. First, the

endogenous copies of A3A and A3B were not inactivated in the hGO_{iA3A} and hGO_{iA3B} models,

raising the possibility that a small fraction of the observed APOBEC-associated mutations

may have originated from these native enzymes. In addition, the inclusion of further control

models, including catalytically inactive mutants of A3A and A3B (i.e., A3A-E72A and A3B-

E255A; Carpenter et al. 2023), would help more definitively rule out the contribution of

indirect mechanisms triggered by APOBEC overexpression. In parallel, treatment with

APOBEC-enzyme inhibitors to our model could also help to validate A3A as a directly

mutagenic enzyme. However, we consider these experiments to be of limited necessity, as

(1) the exogenously introduced APOBEC enzymes is quantitatively predominant in our

system (**Supplemental Table S3**), and (2) it is widely accepted that APOBEC enzymes are

direct mutators with intrinsic mutagenic activity. Lastly, though our gastric organoid models

reproduced comparable transcriptional bursts of APOBEC enzymes *in vivo*, they cannot fully

replicate the physiological complexity of *in vivo* systems.

METHODS

Materials Availability

Organoids established in this study will be available under a material transfer agreement. To do so, please contact the lead author (ysju@kaist.ac.kr).

Human normal gastric samples

Normal gastric tissues were obtained via endoscopic biopsy from a female undergoing routine screening. The protocol for this study was approved by the institutional review board of Yonsei University Gangnam Severance Hospital (3-2018-0207) and KAIST (KH2022-211).

Confirmation of non-neoplasticity of normal gastric organoids with whole-genome sequencing

Whole-genome sequencing analysis confirmed the absence of potent cancer driver mutations and copy number variations of DNA (**Supplemental Fig. S12**) in the normal gastric organoids.

Human stomach organoid culture

Organoid culture methods and media compositions were adopted from previous research with slight modification (Bartfeld et al. 2015). Wnt3A and R-spondin-1 conditioned media was produced with HEK293 cell line producing Afamin-Wnt3a (Mihara et al. 2016) and Cultrex HA-R-Spondin 1-Fc 293T cell line (Trevigen). Complete medium composed of Wnt3A conditioned medium, R-spondin-1 conditioned medium, Advanced DMEM/F-12 (Gibco, Cat No.12634028), HEPES (1M) (Gibco, Cat No.15630-080), Penicillin/streptomycin (10,000 U/mL) (Gibco, Cat No.15140122), GlutaMax (Gibco, Cat No.35050061), Human EGF Recombinant Protein (Invitrogen, Cat No.PHG0311), hNoggin (Peprotech, Cat No.120-10C), hFGF10 (Peprotech, Cat No.3100-26), B27 supplement (50x) serum free (Gibco, Cat No.17504044), N-acetylcysteine (Sigma-Aldrich, Cat No.A9165), Gastrin (Sigma-Aldrich, Cat

No.G9145), Y27632 (Sigma-Aldrich, Cat No.Y0503), TGF-b R kinase inhibitor IV (Biogems,

Cat No.3014193; **Supplemental Table S9**).

Tissues were dissociated into 10~15 clustered cells by pipetting with 1000µl tips after 30

minutes of incubation in 4°C in TrypLE (Gibco, Cat No.12604013). Tissues were washed

twice with PBS, and the pellet was isolated by removing the supernatant after centrifugation

at 300g for 5 minutes at 4°C. The resuspended pellets with cold Matrigel (Corning, Cat

No.BDL356231) were seeded on the 12- or 24-well plates (Merck) in dorm shape. After

incubation of plates in the humidified incubator (5% CO_2, 37□) for 10 minutes to polymerize

Matrigel, prewarmed 0.5-1 ml of culture medium was added to the well. Organoids were

cultured in the humidified incubator. The medium was changed every two to three days.

Organoid passaging was conducted for about two weeks. Medium in the well was replaced

by cell recovery solution (Corning, Cat No.354253), and the plates were incubated for 40~60

minutes at 4°C to dissolve Matrigel. Organoid pellet was isolated by removing supernatants

after centrifugation at 300g for 5 min at 4°C. Organoids followed by incubation for 5 minutes

at 37°C with resuspension with Accutase (Stemcell Technology, Cat No.07922) were

dissociated into 10-15 cell clusters by pipetting 10-20 times. After diluting Accutase by

adding 1 ml of ADF media (Gibco), the pellet was isolated by centrifugation at 300g for 5

minutes at 4°C. Organoids were seeded in a 12- or 24-well plate at a ratio of 1:4 to 1:6 within

Matrigel.

Preparation of vectors for transfection

Two vector systems: (1) CMV-rtTA-HygR vector (Addgene, Cat No.102423) / and (2)

CRISPR-*Cas9* vectors containing gRNA sequence for TP53 (Addgene, Cat No.121917)

were purchased from Addgene. Basically, mini-prep and maxi-prep were conducted with

commercial competent cells (Biosearch Technologies, Cat No.6010) and mini-prep kit

(QIAGEN, Cat No.27104) and maxi-prep kit (QIAGEN, Cat No.12123) according to the

manufacturers' protocols. To create pPB-CMVmin-A3A/A3B-IRES-mCherry vectors,

APOBEC constructs were designed following these steps. NCBI reference sequences for

APOBEC3A (NM_145699.4) and *APOBEC3B* (NM_004900.4) were utilized for the design.

Kozak consensus sequences (GCCACC) were added at the 5' end of the A3A/A3B cDNA

sequences, while HA tag sequences (5'-TAC-CCA-TAC-GAT-GTT-CCA-GAT-TAC-GCT-3')

were appended to the 3' ends of the A3A/B cDNA sequences. Subsequently, *XhoI* and *NotI*

restriction enzyme cut sites (CTCGAG and GCGGCCGC) were added to the 5' and 3' ends

of the construct, respectively. *De novo* gene synthesis from GenScript (Piscataway, NJ) was

used for the synthesis of the two constructs. Following the acquisition of the A3A and A3B

constructs, the constructs were cloned into the backbone vector (pPB-CMVmin-TRE-IRES-

mCherry (Lee et al. 2022) to make pPB-CMVmin-A3A/B-IRES-mCherry cassette.

Transfection of organoids

Organoids were prepared with the same dissociation protocol of passaging organoids.

Transfection methods were adopted from previously reported protocols (Gaebler et al. 2020;

Fujii et al. 2015). A mixture of three types of vectors were utilized for transfection: having (1)

TRE-APOBEC (A3A or A3B)-IRES-mCherry cassette and (2) CMV-rtTA-HygR cassette (3)

pPiggyBac transposase cassette. For hGO$_{iA3A}$ lines, organoids were suspended in 90µl of

Opti-MEM (Gibco, Cat No.31985062), and mixed with about 30 µg of each vector.

Electroporation was conducted using a previously described program from the literature

(Fujii et al. 2015; **Supplemental Table S10**). Transfected organoids were cultured for seven

days with the medium composition described in the previously reported protocol. Organoids

were incubated with the medium containing 1 µg/ml hygromycin (InvivoGen, Cat No.ant-hg-1)

for seven days after splitting. To isolate the transfected organoids having insertions of two

vector cassettes for conditional overexpression, organoids were incubated with doxycycline

(Sigma-Aldrich., Cat No.D9891-1G) in a 3 µg/ml containing medium for 12-16 hours.

mCherry-positive organoids were manually isolated by pipetting under a fluorescent

microscope, and single-cell cloning was conducted according to a previously reported

method by using FACSAria□ (BD Biosciences) and manually picking single cell originated

organoids by pipetting (Youk et al. 2021). Isolated organoids were dissociated in the same

way to passage. Then, organoids were then filtered through a 40 µm strainer (Falcon, Cat

No.352340). Using FACSDiva software, pure single cells were isolated. After seeding the

organoids, the single-cell-originated organoids were isolated. For hGO$_{iA3B}$ lines, organoids

were suspended in 100 µl of BTXpress buffer (BTX), and 10 µg of vector mixtures were

mixed. The following steps were carried out in the reported protocol (Fujii et al. 2015).

Selection and single-cell cloning steps were conducted using the same method as for

hGO$_{iA3A}$ lines.

Doxycycline treatment

A doxycycline stock solution was prepared by dissolving doxycycline in the DMSO. The

stock solution was added to the medium to achieve final concentrations of 0.1 µg/ml and 3

µg/ml. Before treatment, dissociated 10k viable cells were seeded in 24-well the plates. After

7 days, doxycycline solution was added to the medium, and the organoids were incubated

with doxycycline.

Viability assay of organoids

According to the manufacturer's protocol, the viability of organoids was measured using a

Celltiter-Glo 3D Assay kit (Promega, Cat No.G9681). Each well was washed twice with 1 ml

of PBS by repeatedly removing the medium and adding PBS. After washing, PBS was

replaced with fresh medium. An equivalent volume of reaction solution was added to each

well. Matrigel was dissolved in the reaction solution by incubating at 4□ for 30 minutes with

shaking. The amounts of ATP in each well were measured with a luminometer (BERTHOLD

Technologies GmbH, Cat No.TriStar LB 942). The viability percentage was calculated by

dividing luminescence after doxycycline treatment by the average luminescence of untreated

control groups.

Capturing fluorescent images

Fluorescents of mCherry after doxycycline treatment were monitored under the fluorescent

microscope (LEICA, DMi8; excitation = 541-551 nm; emission = 565-605 nm). Fluorescent

images were captured using Las X programs provided by the manufacturer, and

brightness/contrast were adjusted using the same Las X program.

Immunostaining and imaging of organoids

Whole-mount staining of human gastric organoids was performed as previously described

(van Ineveld et al. 2020). Briefly, organoids were fixed with 4 % paraformaldehyde (PFA)

after depolymerizing the 3D matrix using an ice-cold cell recovery solution (Corning). After

washing the cells with 0.1% PBS-Tween-20 and blocking with organoid washing buffer (0.1 %

Triton X-100, 0.02 % SDS, 0.2 % bovine serum albumin (BSA) in 1X PBS), immunolabeling

was performed with mouse anti-HA tag antibody (Santa Cruz, Cat No.sc-7392; 1:50) and

rabbit anti-phospho-histone H2A.X antibody (CST, Cat No.2577S; 1:400) to detect inserted

HA-tagged APOBEC proteins and double-strand breaks, respectively. Goat anti-mouse

Alexa Fluor 488 (Thermo Fisher Scientific, Cat No.A-11001; 1:500) and donkey anti-rabbit

Alexa Fluor 647 (Thermo Fisher Scientific, Cat No.A32795; 1:500) were used as secondary

antibodies, and nuclei were counter-stained with DAPI (Sigma-aldrich., Cat No.D9542;

1:1000). After washing the samples, FUnGI clearing solution was added to the organoids.

The samples with the clearing solution were directly mounted between two coverslips with a

0.25 mm-deep iSpacer (SunJin Lab, Cat No.IS213), and the imaging was performed at least

1 hour after slide preparation.

Imaging was performed with a Leica Stellaris 8 Confocal Microscope at the Research

Solution Center (RSC) at the Institute for Basic Science (IBS). Alexa Fluor 488 (excitation,

479 nm; emission, 482–554 nm), Alexa Fluor 647 (excitation, 580nm; emission, 663-716 nm),

and DAPI (excitation, 405 nm; emission, 425-489 nm) signals were obtained. Either 40X or

63X objectives with a digital zoom factor of 1 to 7-fold were used. The X/Y resolution was set

to 1024 x 1024 pixels. Images were processed and analyzed using Adobe Photoshop.

Standard whole genome sequencing and alignment

Organoids were isolated by removing matrigel with a cell recovery solution (Corning), similar

to the method used for passaging organoids. According to the manufacturer's protocol, DNA

from clonal organoids was extracted using DNeasy Blood & Tissue Kit (QIAGEN, Cat

No.69506). Libraries were constructed with Truseq DNA PCR-Free Library Prep Kits

(Illumina, Cat No.20015963) according to the manufacturer's protocols. Whole genome

sequencing was conducted by the NovaSeq☐6000. Organoid clones were whole-genome

sequenced with a mean 30-fold depth of coverage. Adapter sequence in the FASTA files

was removed by Cutadapt software (https://cutadapt.readthedocs.io/en/stable/; Martin 2011).

Sequencing reads were aligned to the human reference genome, GRCh37, with bwa-mem

v0.7.17 (https://github.com/lh3/bwa; Li and Durbin 2010). Further processing, including

sorting, marking duplication, and indel realignment, was conducted with samtools v1.9

(https://www.htslib.org/; Li et al. 2009), PICARD v2.1.0 (https://broadinstitute.github.io/picard/;

McKenna et al. 2010), and GATK tools v3.8.0 (https://gatk.broadinstitute.org/hc/en-

us/categories/360002369672-Tool-Index; McKenna et al. 2010).

BotSeqS library construction

DNA for the BotSeqS library was extracted with All-Prep DNA/RNA Mini kit (Qiagen, Cat

No.80204) according to the manufacturer's protocol. Constructed libraries with Truseq DNA

PCR-Free Library Prep Kits (Illumina) were utilized for subsequent steps. The construction of

BotSeqS libraries was based on the previous literature with slight modifications (Hoang et al.

2016). Briefly, the quantification of DNA libraries was conducted with KAPA Library

Quantification Kit ILLUMINA® Platforms (Roche, Cat No.KK48247) according to the

manufacturer's protocol. An effective library concentration was calculated considering the

size of added DNA length during library construction. PCR amplification of libraries

equivalent to 4 pg of DNA was performed with the KAPA HiFi Uracil+ kit (Roche, Cat

No.KK2802) and primers having a Y-adaptor sequence with a phosphorothioate bond(*) at

the 3' end from IDT (Coralville, IA):

Forward: 5'-AATGATACGGCGACCACCGAG*A-3'

Reverse: 5'-CAAGCAGAAGACGGCATACGA*G-3'

PCR programs were adopted from the protocol of KAPA Library Amplification Kits (Roche),

and PCR was conducted with 20 cycles. Library clean-up was performed with SPRIselect

beads (BECKMAN COULTER life Sciences) according to the manufacturer's protocol. The

libraries were sequenced as paired-end sequencing (2x151 bps) using NovaSeq 6000.

In vitro **deamination of DNA with synthesized APOBEC3B**

The NEBNext Enzymatic Methyl-seq (EM-seq) Kit (NEB, Cat No.E7120S) and the

manufacturer's protocol were utilized for the construction of the DNA library with slight

modification. Firstly, the DNA library was constructed with DNA extracted from a doxycycline

treatment-free hGO$_{iA3B}$ line. DNA libraries equivalent to 10 pg of input DNA were prepared

following the process used for BotSeqS library construction. Oxidation of DNA by TET2 and

denaturation steps were performed according to the manufacturer's instructions. For

deamination steps, synthesized recombinant APOBEC3B from EUROPROTEIN INC. (North

Brunswick Township, NJ) was used instead of the APOBEC protein included in the EM-seq

kit. Additionally, the reaction time was modified from 3 hours to 30 and 60 seconds. Library

amplification steps followed the BotSeqS library construction.

Whole transcriptome sequencing library construction

RNA was isolated during the extraction of DNA for BotSeqS libraries with AllPrep DNA/RNA

Mini kit according to the manufacturer's instructions. RIN value of raw RNA was measured

with Agilent 4200 TapeStation (Agilent, Cat No.G2991BA). Libraries were constructed with

NEBNext Ultra II Directional DNA Library Prep Kit for Illumina (NEB, Cat No.E7760) and

QiAseq FastSelect -rRNA HMR kit (Qiagen, Cat No.334388) according to manufacturers'

instructions. Clean-up process was conducted with SPRIselect beads according to the

manufacturer's protocol. Libraries were sequenced with Novaseq 6000 with paired-end

sequencing.

Confirming non-neoplasticity in primary human gastric organoid

The germline mutation calling mode in GATK v4.0.12.0 (https://gatk.broadinstitute.org/hc/en-

us/categories/360002369672-Tool-Index; McKenna et al. 2010) was utilized for the

identification of mutation in the organoid lines. The primary call set was first filtered by in-

house scripts using the pysam module in Python

(https://pysam.readthedocs.io/en/latest/index.html; Li et al. 2009) with the following

conditions: (1) the number of reference reads ≥ 5; (2) the number of variant reads ≥ 3; (3)

median miss-matched bases in variant reads < 5; (4) median mismatched bases in variant

reads < 5 (5) median mapping quality of reference reads > 20 (6) median mapping quality of

variant reads > 20 (7) median distance from the ends of reads > 4. The impact of variants in

the protein-coding region was annotated with ANNOVAR

(https://annovar.openbioinformatics.org/en/latest/; Wang et al. 2010). Non-silent mutations

located in the protein-coding and splicing regions were isolated. The functional impact in

cancer driver genes (the list can be downloaded at https://cancer.sanger.ac.uk/cosmic) was

evaluated with PolyPhen-2 (Adzhubei et al. 2013) and SIFT (Ng and Henikoff 2003) scores

annotated by ANNOVAR, AlphaMissence (https://alphamissense.hegelab.org/; Cheng et al.

2023), and MutPred-Indel (http://mutpred2.mutdb.org/mutpredindel/about.html; Pagel et al.

2019). Considering the role of cancer driver genes, the impact of variants was finally

assessed.

Calling copy number variations

719   Copy number variations were accessed by Sequenza (https://sequenza-

720   utils.readthedocs.io/en/latest/#; Favero et al. 2015). Copy number variations in segments

721   smaller than 1 Mbp were considered false positives. After removing the copy number

722   variations in short segments, final copy number variations were investigated by a second run

723   of the Sequenza.

Detection of somatic mutations

To detect single nucleotide variants and indels in clonal organoids, GATK Mutect2 v4.1.9

(https://gatk.broadinstitute.org/hc/en-us/categories/360002369672-Tool-Index; McKenna et

al. 2010) and Strelka2 v2.9.2 (https://github.com/Illumina/strelka; Kim et al. 2018) were

utilized. Bulk whole-genome sequencing of mother clones, hGO_{iA3A}, $TP53KO\text{-}hGO_{iA3A}$,

hGO_{iA3B}, and $TP53KO\text{-}hGO_{iA3B}$ organoids, were utilized as matched normal for calling

somatic mutations in doxycycline treated organoids. In each call set, false-positive variants

were filtered out with in-house python scripts annotating information within BAM files with the

pysam, a python module. The filtered call sets from two callers, Mutect2 and Strelka2, were

merged, and the union call set was utilized for downstream analysis. To exclude mutations

generated during the culture of mother clones, recurrent somatic mutations observed across

multiple doxycycline-treated clones, including control samples, were removed from each call

set.

In the case of BotSeqS, VarScan2 v2.3.9 (https://varscan.sourceforge.net/; Koboldt et al.

2012) was used to increase the sensitivity of calls. Similarly, in-house Python scripts were

utilized to filter out false-positive calls. Specifically, F1R2 and F2R1 reads from same DNA

was grouped and filtered with the following criteria: (1) total depth of each type of read ≥ 3;

(2) the number of variant reads ≥ 3 or 90% of reads; (3) distances of mutations from each

end of each read > 5 bp; (4) distances of mutations from the end of reads > 100 bp

considering the DNA strand (e.g., excluding C>T near the end of F1R2 reads, but not G>A);

(5) median mapping quality of variant reads ≥ 20; (6) median base quality of variant reads ≥

30; (7) the number of variants in WGS of HEK293T <3; (8) the number of variants in WGS of

control < 1. Unlike with standard whole genome sequencing, C>T mutations were frequently

observed at the end of the reads, and these were considered false positives and removed.

Additionally, rare variants from HEK293T cells, which were used to generate conditioned

medium, were observed. Thus, variants observed in HEK293T cell line BAM files at least

three times were removed from the call set.

Calculation of mutation rate in BotSeqS

Unlike standard WGS, the length of the effective covered region was calculated for each

BAM file. First, each read was evaluated using in-house scripts that considered DNA strand

orientation and applied the following criteria to isolate effective DNA fragments: (1) median

mapping quality > 20 (2) total depth of each type of reads ≥ 3. Only regions where both

F1R2 and F2R1 reads were aligned were included in the covered region calculation. To

account for the exclusion of mutations located within 100 bp of extreme of read ends during

variant filtering (considering the reference genome strand), the total length of the covered

region was adjusted by multiplying it by (151x2-110)/(151x2)). Mutation rates were

calculated by dividing the number of observed mutations by this adjusted covered length. To

normalize mutational burdens against the standard genome length, mutation rates were then

multiplied by the total genome length excluding repeat regions (3,041,373,115 bp) and

further doubled.

Detection of deamination during *in vitro* denomination with APOBEC3B

The calling pipeline for BotSeqS results was utilized with slight modifications. Among the

criterias, the filtering criteria for distances from the ends of reads were neglected. Initially, all

of the filtered mutations were collected. Among the mutations in grouped DNA, only those in

DNA fragments where only one strand of DNA was mapped were counted.

To calculate the genome-wide mutation rates, the BotSeqS pipeline was also used, with the

exception of counting cytosine or guanine considering the strand of DNA. Since original

methylated CpG sites were preserved during the library construction, total counts and

mutation counts in all of CpG contexts were excluded. To calculate mutation rates in DNA

fragments with at least one C>T variant, ranges of DNA fragments overlapping with the

variants were isolated while considering the DNA strands using BEDtools

(https://bedtools.readthedocs.io/en/latest/; Quinlan and Hall 2010). After this step, cytosine

or guanine outside of CpG contexts were counted.

Analysis of mutational signatures

Mutational signature analysis of single base substitutions (SBS) and small indels were

carried out using non-negative least squares method. The mutational signature was

represented by 96 patterns of SBS and 83 patterns of small indels (Alexandrov et al. 2020).

Pre-learned catalogs of mutational signatures in the COSMIC (Sondka et al. 2024) were

used to fit individual samples with a known set of signatures for each tissue type. SBS2 and

SBS13 (known APOBEC signatures) were included for all cases.

Analysis of mutation rates depending on epigenetic marker

Risk ratio of mutation rates in non-signal regions and signal-detected regions depending on

each epigenetic marker was analyzed based on previous researches (Nam et al. 2023;

Supek and Lehner 2017). Genome-wide signals of each marker, including replication timing,

across eight cell types (E017,E114, E117, E118, E119, E122, E125, and E127) were

downloaded from Roadmap Epigenomics Consortium. Fold-enrichment signal was averaged

across the cell types, and the regions showing fold-enrichment signal lower than 1 identified

as bin 0, non-signal regions. The other regions were classified as the signal-detected

regions. Based on the defined regions, the number of SNVs were counted, and

subsequently, risk ratio was calculated considering 3bp genomic context. For APOBEC-

associated mutations, only cytosines within the TCN context were considered as background,

and C>T and C>G substitutions at these sites were counted. For SBS5 and SBS40, thymine

bases outside the TCN context were used as the reference, and T>A,T>G,T>C, C>T and

C>G substitutions were counted. C>A mutations were excluded to avoid confounding effects

from overlapping sequence contexts associated with SBS1 and SBS18.

To analyze fold-enrichment of mutation rates, in case of replication timing and H3K27me3

signals, the signal-detected regions were further divided into four groups, considering the

equal distribution of length of each group. In case of expression level, expression level with

TPM unit in each base calculated from merged whole-transcriptome sequencing data of

doxycycline treated organoids for 48 hours under 3 µg/ml doxycycline was utilized. First of all,

non-genic regions and genic regions were classified based on gene information from the

"hg19_refGene.txt" file. Only genic regions were included in the analysis. Genic regions

having 0 of TPM were defined as bin0 regions. TPM>0 regions were classified into 4 groups

(cutoff; bin1=0.05, bin2=1.73, and bin3&bin4=9.68) based on quartiles of TPM. Considering

interference among H3K27me3, replication timing, and transcription, glm.nb() function in

MASS, R package (https://cran.r-project.org/web/packages/MASS/index.html; Venables and

Ripley 2003), was utilized in enrichment analysis as described in previous research (Supek

and Lehner 2017).

Analysis of mutation rates depending on genomic location

SNVs located in previously described mappable regions were utilized throughout the

analysis. Gene information included in "hg19_refGene.txt" file from ANNOVAR tools were

utilized to match the additional information of position of SNVs. Classification of sub-genic

(5`/3`UTR, protein coding sequence (CDS), and intron) and transcription strand were also

based on the gene information. All of the merged genic regions were used as standard for

discrimination of genic and intergenic regions. Instead of that, genic regions that do not

overlap with other genic regions and CDS regions located between intron regions were

utilized for comparing mutation rates in the sub-genic region as previous research reported

(Frigola et al. 2017).

Calling structural variations

Structural variations were called using Delly v0.7.6 (https://github.com/dellytools/delly;

Rausch et al. 2012). Raw calls were filtered using in-house scripts. The final call set was

manually reviewed by Integrative Genomics Viewer (https://igv.org/; Robinson et al. 2011).

Calculating RNA expression level

Bulk RNA-seq reads were aligned to GRCh37 using STAR2 v2.6.1d

(https://github.com/hbctraining/Intro-to-rnaseq-hpc-O2; Dobin et al. 2013). TPM and reads

count were calculated with RSEM v1.3.1 (https://github.com/deweylab/RSEM; Li and Dewey

2011). Differential expression gene analysis was conducted using the DEseq2 package in R

(https://bioconductor.org/packages/release/bioc/html/DESeq2.html; Love et al. 2014).

Calculation of ratio between endogenous APOBEC mRNA and overexpressed

APOBEC mRNA

To distinguish transcripts originating from endogenous APOBEC genes versus the

exogenous construct, we utilize sequence differences between the two sources. For A3A,

two heterozygous variants at chr22:39,357586 (C/T) and 39,357,599 (C/T) were used. In the

endogenous A3A allele, reads carried the same base at both positions-either C/C or T/T-

while the inserted construct contained C/T bases at the two positions, respectively. For A3B,

a homozygous variant at chr22:39,381,999 (C) was present in the endogenous gene, while

the construct carried only T at the same locus. The proportion of endogenous mRNA was

estimated by calculating the ratio of supporting reads at the positions.

Calling RNA editings

VarScan2 was utilized to call RNA editing. WGS of mother clones was utilized as paired

normal. False positives that did not meet the following criteria were filtered out using an in-

house Python script with the pysam module: (1) excluding HLA regions; (2) DNA depth ≥ 5;

(3) number of variant reads in DNA = 0; (4) RNA depth ≥ 4; (5) number of variant reads ≥ 3;

(6) median number of NM tags in the variant reads < 4; (7) percentage of variant reads with

NM tags greater than 3 < 0.5; (8) distance from the end of reads ≥ 5; (9) variants in only one

strand; (10) median base quality of variants > 20; (11) percentage of variant reads with clips

< 0.5; (12) median mapping quality of variant reads ≥ 30; (13) single-base variant. RNA

editings that did not meet all criteria but were commonly observed with all conditions fulfilled

in more than two samples were rescued.

To compare the RNA editings, the number of RNA editings was normalized with total

sequencing output size. The target output size was determined by the sample having the

lowest sequencing output. RNA editings were filtered again by adjusting the ratio of

sequencing output to the total depth and variant reads. Variants that met the following two

criteria were used in the analysis: (1) corrected depth of RNA ≥ 4 (2) corrected the number

of variants.

Analysis of RNA editing signatures

RNA editing signatures were obtained by a modified version of the mutational signature

extraction method used in (Youk et al. 2024; Alexandrov et al. 2013). Briefly, this method

uses non-negative matrix factorization (NMF) to disentangle an individual RNA editing

spectrum based on a notion of mixed spectra (Cichocki et al. 2006).

RNA editing spectrum was represented as 192 patterns of single-base change with

immediately adjacent bases on the 5' and 3'-end in the canonical mRNA sequence. The

number of features is doubled of DNA mutational signature (Alexandrov et al. 2013),

because RNA does not necessarily have symmetric base change due to general non-

strandedness.

Total 18 experimental data were used as input, split into two subsets: A3A and A3B sets.

Each of the A3A and A3B sets included three batches from experiments with 0μg/ml, 0.1

μg/ml, and 3 μg/ml APOBEC3 exposure, totaling 9 batches for each set. Initially, the

signature extraction method was used with no additional procedures from the previous

method (Youk et al. 2024), and then the optimal number of independent RNA editing

signatures was determined. We observed that there were 4 potential candidates for RNA

editing signature that are attributable to APOBEC3A, APOBEC3B, ADAR, and the other

endogenous editing processes.

APOBEC3A and APOBEC3B aligned well with editing patterns of C>T found in experiments

with varying exposure amounts. However, the ADAR signature co-appeared in these

signatures at significant proportion. This "leakage" phenomenon occurs due to the constant

activity of the ADAR editing process, causing its signal to always appear in RNA editing

spectra regardless of the experimental settings. This can be problematic when trying to

accurately quantify the attributes of each RNA editing process in a given sample.

To mitigate the potential effect of the leakage phenomenon, an L1 constraint was imposed

on the signature matrix (W) (Youk et al. 2024). For that, the NMF update algorithm was

adopted from a work on sparse NMF (Le Roux et al. 2015). It presents a method that puts L1

constraint on the exposure matrix (H). Therefore, we tweaked its objective function by

transposing the data matrix (V) as well as W and H and back-transposing them to get W.

Afterwards, the matrix W was normalized so that each column vector sums to one. The

sparsity coefficient (μ) was set to 0.6, after trying out several values in a range [0.001, 10].

The A3A set resulted in ADAR and APOBEC3A that were now isolated from each other. The

A3B set led to ADAR and APOBEC3B signatures that were also well-separated. ADAR

signatures learned from the A3A and A3B set were consistent with cosine similarity 0.999

and were merged into a single ADAR signature by taking average. Endogenously occurring

RNA editing other than ADAR was negligible in amount and did not add much value to

overall analysis. Thus, we did not seek further signatures.

Lastly, the three RNA editing signatures learned from the data were refitted to individual

samples to evaluate the attributes of the RNA editing processes in each sample.

Analysis of secondary structure of RNA editing sites.

Secondary structure of RNA editing sites were predicted as described previously (Jalili et al.

2020). Briefly, a 41 bp sequence centered on each RNA editing site in the canonical mRNA

was used to assess the potential for forming secondary structures. Stem strength was

calculated as 3*G/C pair + 1*A/T pair in stem. Among candidate structures, the most

probable one was selected based on the following hierarchical criteria: (1) highest stem

strength, (2) greatest number of G/C pairs in the stem, and (3) smallest loop size.

Calling clustered mutation

SigProfilerClusters (https://github.com/AlexandrovLab/SigProfilerClusters; Bergstrom et al.

2022a), a python module, was utilized to identify clustered mutations. FlexMix, R package

(https://cran.r-project.org/web/packages/flexmix/index.html; Leisch 2004), was utilized to

classify the identified clustered mutations into *omikli* and *kataegis*. Since the tools determine

the standard intermutation distance for clustering through simulations that randomly

distribute SNVs, the total number of SNVs influence the detection rate of clustered mutation

events. To correct for the number of clustered mutation events, simulations were conducted.

Identified clustered mutations from our analysis and 703,858 SNVs from 125 samples in

PCAWG database, where the contribution of age-related mutational signatures (SBS1,

SBS5, and SBS40) is over 0.8, were utilized for simulation. The target number of SNVs was

set to 33 (increased by 2,000 from 2,000 to 10,000; increased by 5,000 from 15,000 to

100,000; increased by 10,000 from 110,000 to 200,000). From the observed 615 *omikli*

(1,369 SNVs) and 109 *kataegis* (559 SNVs) events, the clustered mutation events were

randomly sampled based on the following conditions for 50 iterations: 50, 100, 250, 500

omikli, and 25, 50, 75, 100 *kataegis*. Background mutations were selected to match the

number remaining after excluding the number of clustered mutations from the target number

of SNVs. With the same SigProfilerClusters module, clustered mutations, *omikli and kataegis,*

were identified, and the detection rate was calculated. By combining data from the two types

of clustered mutations, the relationship between the number of mutations and the detection

rate of clustered mutations was estimated (**Supplemental Fig. S13**) using the "drm" function

with the fct=AR.3() option in the drc, R package (https://cran.r-

project.org/web/packages/drc/index.html; Ritz et al. 2019).

To compare clustered mutation events, 146 samples from the PCAWG database were

selected based on the following criteria: (1) nine cancer types showing high prevalence of

APOBEC mutational signatures; (2) samples with fewer than 500 SBS2+SBS13 were

excluded in both our samples and PCAWG cancer samples to avoid bias.

Analysis of 4bp context mutations

SNVs found in 10 hGO$_{iA3A}$ samples (excluding control samples) were utilized in the analysis.

Among 567 PCAWG samples across eight cancer types with a high prevalence of APOBEC

mutational activity - breast adenocarcinoma (BRCA; n=195), esophageal adenocarcinoma

(ESAD; n=97), stomach adenocarcinoma (STAD; n=68), head and neck squamous cell

carcinoma (HNSC; n=56), lung squamous cell carcinoma (LUSC; n=47), uterine corpus

endometrial carcinoma (UCEC; n=44), lung adenocarcinoma (LUAD; n=37), bladder

urothelial Carcinoma (BLCA; n=23) - only those with a combined APOBEC-associated

mutational burden (SBS2 + SBS13) greater than 5,000 were selected for this analysis (n=63).

This subject included: LUSC (n=15), BRCA (n=12), BLCA (n=11), HNSC (n=13), LUAD

(n=6), UCEC (n=3), ESAD (n=2), and STAD (n=1).

Analysis of A3A and A3B expression levels in single cancer cells with public scRNA

data

Publicly available single-cell RNA-seq datasets, generated with Smart-seq library, from lung

adenocarcinoma, triple negative breast cancer, esophageal adenocarcinoma and

esophageal squamous cell carcinoma were analyzed using the same pipeline applied to our

bulk RNA-seq data. Epithelial cell populations were first identified following the tutorial

workflow provided in the Seurat R package (https://satijalab.org/seurat/; Satija et al. 2015),

with the exception of the cell filtering step. Within the epithelial population, cancer cells were

distinguished based on the presence of large-scale copy number variations (CNVs) inferred

using the inferCNVpy Python package

(https://infercnvpy.readthedocs.io/en/latest/index.html). CNV profiles were calculated using

fibroblast and endothelial cell populations as reference non-malignant cells. For the head

and neck squamous cell carcinoma dataset, where raw data were not available, we utilized a

publicly provided summary table containing TPM values and annotated cell types for the

analysis.

Publicly available datasets

Publicly available whole-genome sequencing data were utilized to demonstrate copy number

variation in the normal human gastric organoid. WGS of blood from HC05 sample was

utilized as the unpaired normal sample (https://ega-archive.org/studies/EGAS00001006213;

Nam et al. 2023). To compare the 4bp-context preference and frequency of clustered

mutation, SNV calls from the ICGC/TCGA Pan-Cancer Analysis of Whole-Genome (PCAWG)

Consortium were utilized for the analysis. The call set data is available for download at

https://dcc.icgc.org/releases/PCAWG.

To compare in vivo transcriptional burst of A3A and A3B, we analyzed publicly available

Smart-seq-based single-cell RNA-seq datasets from five cancer types: (1) lung

adenocarcinoma (https://www.ncbi.nlm.nih.gov/bioproject/591860; Maynard et al. 2020) (2)

head and neck squamous cell carcinoma

(https://www.ncbi.nlm.nih.gov/bioproject/?term=GSE103322; Puram et al. 2017) (3) triple

negative breast cancer (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE118390;

Karaayvaz et al. 2018) (4) esophageal adenocarcinoma and esophageal squamous cell

carcinoma (https://www.ncbi.nlm.nih.gov/bioproject/PRJNA401501; Wu et al. 2018).

Quantification and statistical analysis

All statistical analyses were performed with R version 4.1.3 (R Core Team, Vienna, Austria).

A two-tailed one-sample t-test was used to evaluate p-values for comparing APOBEC-

associated SNVs and expression levels between the groups. Linear regressions were

conducted using the basic 'lm' function in R to analyze the association among APOBEC-

associated SNVs, ID9, SBS5, and SBS40. A chi-square test was utilized to evaluate p-

values for comparing replication strand bias and transcription strand bias. A 95% confidence

interval was used to determine the statistical range of continuous data.

DATA ACCESS

Whole-genome sequencing, EM-seq, BotSeqS, and bulk RNA-seq raw data produced in this

study were deposited in the Korean Nucleotide Archive (KoNA;

https://www.kobic.re.kr/kona/). The project accession ID is KAP240815

(https://kbds.re.kr/KRA/browse/Experiment?keyword=KAP240815;

https://kbds.re.kr/bioproject_run_all?from_uid=2585828). All data are provided for review

purposes upon request by reviewers. Essential in-house scripts used in this study are

available on Zenodo (DOI: 10.5281/zenodo.12771074;

https://zenodo.org/records/12771074?preview=1&token=eyJhbGciOiJIUzUxMiJ9.eyJpZCI6Ij

diNDQ2YzU4LTIxNDMtNDViOS1hYTQ5LTE1MTY4OTEzMmQ2NyIsImRhdGEiOnt9LCJyY

W5kb20iOiIwZTg1YmFiYjAzNDRmMzgwOTQ1ZGI2MGI0NDYxMmZkZiJ9.pxeRUXdTXwcoJ

rUMkx2ZGlMZNnOyNai2-

mcrqyiOlbJkB10lgSOhcvc5Z6mrNKc4TSicAm7NrSgqNops57_SWg).

COMPETING INTEREST STATEMENT

Y.S.J. is a genomic co-founder of Inocras Inc..

ACKNOWLEDGEMENTS

The authors thank Youngwon Cho (Epithelial Biology Center, Vanderbilt University Medical

Center) for helping with the cell viability test, Myungsuk Choi (KAIST) for technical help for

organoids, and Mia Petljak (New York University) for her valuable advice and feedback on

the manuscript. This work was supported by the Young Investigator Grants from the Human

Frontier Science Program (RGY0071/2018 to Y.S.J., B.-K.K., and H.S.); the National

Research Foundation of Korea (Leading Researcher Program NRF-2020R1A3B2078973 to

Y.S.J.); the Korea Bio Data Station (N24NM016-24 to J.L and J.W.P); and the Institute for

Basic Science (IBS-R021-D1 to B.-K.K).

Author Contributions

Y.S.J., B.-K.K., and H.S. designed the experiments. J.-H.K. provided human gastric

epithelium sample from biopsy. Y.J.B. established organoids from primary human gastric

tissues. J.L. and J.W.P. helped basic bioinformatic works (alignment and calling somatic

variants). J-H.L. and J.Y. conducted vector cloning. Y.A. and T.K. conducted the

transformation of organoids, with J-H.L. and J.Y. providing advice. Y.A. performed organoid

culture, clonal expansion, and DNA/RNA extraction. B.-K.K. and H.K. provided training in

organoids culture technologies. Y.A. performed cell viability analysis. J-H.L. performed

immunohistochemistry. S.A.O. conducted *in vitro* deamination experiments and most of the

library construction including WGS and RNA-seq. Y.A., J.-H.P., and K.Y. conducted library

construction for duplex sequencing analysis. Y.A. conducted most bioinformatic analyses,

including alignment, mutation calling, duplex sequencing analysis, and analysis of

expression levels with bulk RNA-seq data, with S.P., J.Y., and Y.S.J. providing advice. Y.A.

REFERENCE

Abascal F, Harvey LMR, Mitchell E, Lawson ARJ, Lensing SV, Ellis P, Russell AJC,
Alcantara RE, Baez-Ortega A, Wang Y, et al. 2021. Somatic mutation landscapes at
single-molecule resolution. *Nature* **593**: 405–410.

Adzhubei I, Jordan DM, Sunyaev SR. 2013. Predicting functional effect of human missense
mutations using PolyPhen-2. *Curr Protoc Hum Genet* **Chapter 7**: Unit7.20.

Alexandrov LB, Kim J, Haradhvala NJ, Huang MN, Tian Ng AW, Wu Y, Boot A, Covington
KR, Gordenin DA, Bergstrom EN, et al. 2020. The repertoire of mutational signatures in
human cancer. *Nature* **578**: 94–101.

Alexandrov LB, Nik-Zainal S, Wedge DC, Campbell PJ, Stratton MR. 2013. Deciphering
signatures of mutational processes operative in human cancer. *Cell Rep* **3**: 246–259.

Alonso de la Vega A, Temiz NA, Tasakis R, Somogyi K, Salgueiro L, Zimmer E, Ramos M,
Diaz-Jimenez A, Chocarro S, Fernández-Vaquero M, et al. 2023. Acute expression of
human APOBEC3B in mice results in RNA editing and lethality. *Genome Biol* **24**: 267.

Amador V, Ge S, Santamaría PG, Guardavaccaro D, Pagano M. 2007. APC/C(Cdc20)
controls the ubiquitin-mediated degradation of p21 in prometaphase. *Mol Cell* **27**: 462–
473.

Amouroux R, Campalans A, Epe B, Radicella JP. 2010. Oxidative stress triggers the
preferential assembly of base excision repair complexes on open chromatin regions.
Nucleic Acids Res **38**: 2878–2890.

Bartfeld S, Bayram T, van de Wetering M, Huch M, Begthel H, Kujala P, Vries R, Peters PJ,
Clevers H. 2015. In vitro expansion of human gastric epithelial stem cells and their
responses to bacterial infection. *Gastroenterology* **148**: 126–136.e6.

Bergstrom EN, Kundu M, Tbeileh N, Alexandrov LB. 2022a. Examining clustered somatic
mutations with SigProfilerClusters. *bioRxiv* 2022.02.11.480117.
https://www.biorxiv.org/content/10.1101/2022.02.11.480117v2.abstract (Accessed July
7, 2024).

Bergstrom EN, Luebeck J, Petljak M, Khandekar A, Barnes M, Zhang T, Steele CD, Pillay N,
Landi MT, Bafna V, et al. 2022b. Mapping clustered mutations in cancer reveals
APOBEC3 mutagenesis of ecDNA. *Nature* **602**: 510–517.

Burns MB, Lackey L, Carpenter MA, Rathore A, Land AM, Leonard B, Refsland EW,
Kotandeniya D, Tretyakova N, Nikas JB, et al. 2013. APOBEC3B is an enzymatic
source of mutation in breast cancer. *Nature* **494**: 366–370.

Carpenter MA, Temiz NA, Ibrahim MA, Jarvis MC, Brown MR, Argyris PP, Brown WL,
Starrett GJ, Yee D, Harris RS. 2023. Mutational impact of APOBEC3A and APOBEC3B
in a human cell line and comparisons to breast cancer. *PLoS Genet* **19**: e1011043.

Cazzalini O, Scovassi AI, Savio M, Stivala LA, Prosperi E. 2010. Multiple roles of the cell
cycle inhibitor p21(CDKN1A) in the DNA damage response. *Mutat Res* **704**: 12–20.

Chan K, Roberts SA, Klimczak LJ, Sterling JF, Saini N, Malc EP, Kim J, Kwiatkowski DJ,
Fargo DC, Mieczkowski PA, et al. 2015. An APOBEC3A hypermutation signature is
distinguishable from the signature of background mutagenesis by APOBEC3B in human

cancers. *Nat Genet* **47**: 1067–1072.

Choi J-H, Pfeifer GP. 2005. The role of DNA polymerase eta in UV mutational spectra. *DNA*
Repair **4**: 211–220.

Cichocki A, Zdunek R, Amari S. 2006. New algorithms for non-negative matrix factorization
in applications to blind source separation. In *2006 IEEE International Conference on*
Acoustics Speed and Signal Processing Proceedings, IEEE
http://ieeexplore.ieee.org/document/1661352/.

Dananberg A, Striepen J, Rozowsky JS, Petljak M. 2024. APOBEC Mutagenesis in Cancer
Development and Susceptibility. *Cancers* **16**.
http://dx.doi.org/10.3390/cancers16020374.

Delbos F, De Smet A, Faili A, Aoufouchi S, Weill J-C, Reynaud C-A. 2005. Contribution of
DNA polymerase eta to immunoglobulin gene hypermutation in the mouse. *J Exp Med*
201: 1191–1196.

DeWeerd RA, Németh E, Póti Á, Petryk N, Chen C-L, Hyrien O, Szüts D, Green AM. 2022.
Prospectively defined patterns of APOBEC3A mutagenesis are prevalent in human
cancers. *Cell Rep* **38**: 110555.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**: 15–
21.

Durfee C, Temiz NA, Levin-Klein R, Argyris PP, Alsøe L, Carracedo S, Alonso de la Vega A,
Proehl J, Holzhauer AM, Seeman ZJ, et al. 2023. Human APOBEC3B promotes tumor
development in vivo including signature mutations and metastases. *Cell Rep Med* **4**:
101211.

Favero F, Joshi T, Marquard AM, Birkbak NJ, Krzystanek M, Li Q, Szallasi Z, Eklund AC.
2015. Sequenza: allele-specific copy number and mutation profiles from tumor
sequencing data. *Ann Oncol* **26**: 64–70.

Fixman B, Díaz-Gay M, Qiu C, Margaryan T, Lee B, Chen XS. 2024. Validation of the
APOBEC3A-Mediated RNA Single Base Substitution Signature and Proposal of Novel
APOBEC1, APOBEC3B, and APOBEC3G RNA Signatures. *J Mol Biol* **436**: 168854.

Frigola J, Sabarinathan R, Mularoni L, Muiños F, Gonzalez-Perez A, López-Bigas N. 2017.
Reduced mutation rate in exons due to differential mismatch repair. *Nat Genet* **49**:
1684–1692.

Fujii M, Matano M, Nanki K, Sato T. 2015. Efficient genetic engineering of human intestinal
organoids using electroporation. *Nat Protoc* **10**: 1474–1485.

Gaebler A-M, Hennig A, Buczolich K, Weitz J, Welsch T, Stange DE, Pape K. 2020.
Universal and Efficient Electroporation Protocol for Genetic Engineering of
Gastrointestinal Organoids. *J Vis Exp*. http://dx.doi.org/10.3791/60704.

Gnatt AL, Cramer P, Fu J, Bushnell DA, Kornberg RD. 2001. Structural basis of
transcription: an RNA polymerase II elongation complex at 3.3 A resolution. *Science*
292: 1876–1882.

Green AM, Landry S, Budagyan K, Avgousti DC, Shalhout S, Bhagwat AS, Weitzman MD.
2016. APOBEC3A damages the cellular genome during DNA replication. *Cell Cycle* **15**:

998–1008.

Hoang ML, Kinde I, Tomasetti C, McMahon KW, Rosenquist TA, Grollman AP, Kinzler KW, Vogelstein B, Papadopoulos N. 2016. Genome-wide quantification of rare somatic mutations in normal human tissues using massively parallel sequencing. *Proc Natl Acad Sci U S A* **113**: 9846–9851.

Hoopes JI, Cortez LM, Mertz TM, Malc EP, Mieczkowski PA, Roberts SA. 2016. APOBEC3A and APOBEC3B Preferentially Deaminate the Lagging Strand Template during DNA Replication. *Cell Rep* **14**: 1273–1282.

Jager M, Blokzijl F, Kuijk E, Bertl J, Vougioukalaki M, Janssen R, Besselink N, Boymans S, de Ligt J, Pedersen JS, et al. 2019. Deficiency of nucleotide excision repair is associated with mutational signature observed in cancer. *Genome Res* **29**: 1067–1077.

Jalili P, Bowen D, Langenbucher A, Park S, Aguirre K, Corcoran RB, Fleischman AG, Lawrence MS, Zou L, Buisson R. 2020. Quantification of ongoing APOBEC3A activity in tumor cells by monitoring RNA editing at hotspots. *Nat Commun* **11**: 2971.

Karaayvaz M, Cristea S, Gillespie SM, Patel AP, Mylvaganam R, Luo CC, Specht MC, Bernstein BE, Michor F, Ellisen LW. 2018. Unravelling subclonal heterogeneity and aggressive disease states in TNBC through single-cell RNA-seq. *Nat Commun* **9**: 3588.

Kazanov MD, Roberts SA, Polak P, Stamatoyannopoulos J, Klimczak LJ, Gordenin DA, Sunyaev SR. 2015. APOBEC-Induced Cancer Mutations Are Uniquely Enriched in Early-Replicating, Gene-Dense, and Active Chromatin Regions. *Cell Rep* **13**: 1103–1109.

Kim J, Mouw KW, Polak P, Braunstein LZ, Kamburov A, Kwiatkowski DJ, Rosenberg JE, Van Allen EM, D'Andrea A, Getz G. 2016. Somatic ERCC2 mutations are associated with a distinct genomic signature in urothelial tumors. *Nat Genet* **48**: 600–606.

Kim S, Scheffler K, Halpern AL, Bekritsky MA, Noh E, Källberg M, Chen X, Kim Y, Beyter D, Krusche P, et al. 2018. Strelka2: fast and accurate calling of germline and somatic variants. *Nat Methods* **15**: 591–594.

Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER, Ding L, Wilson RK. 2012. VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing. *Genome Res* **22**: 568–576.

Law EK, Levin-Klein R, Jarvis MC, Kim H, Argyris PP, Carpenter MA, Starrett GJ, Temiz NA, Larson LK, Durfee C, et al. 2020. APOBEC3A catalyzes mutation and drives carcinogenesis in vivo. *J Exp Med* **217**. http://dx.doi.org/10.1084/jem.20200261.

Lawrence MS, Stojanov P, Polak P, Kryukov GV, Cibulskis K, Sivachenko A, Carter SL, Stewart C, Mermel CH, Roberts SA, et al. 2013. Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature* **499**: 214–218.

Lawson ARJ, Abascal F, Coorens THH, Hooks Y, O'Neill L, Latimer C, Raine K, Sanders MA, Warren AY, Mahbubani KTA, et al. 2020. Extensive heterogeneity in somatic mutation and selection in the human bladder. *Science* **370**: 75–82.

Lee J-H, Kim S, Han S, Min J, Caldwell B, Bamford A-D, Rocha ASB, Park J, Lee S, Wu S-HS, et al. 2022. p57 imposes the reserve stem cell state of gastric chief cells. *Cell Stem Cell* **29**: 826–839.e9.

Leisch F. 2004. FlexMix: A general framework for finite mixture models and latent class
regression inR. *J Stat Softw* **11**. http://www.jstatsoft.org/v11/i08/.

Le Roux, J., Weninger, F.J. and Hershey, J.R., 2015. Sparse NMF–half-baked or well
done?. Mitsubishi Electric Research Labs (MERL), Cambridge, MA, USA, Tech. Rep.,
no. TR2015-023, 11:13-15. https://www.merl.com/publications/docs/TR2015-023.

Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* **12**: 323.

Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler
transform. *Bioinformatics* **26**: 589–595.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin
R, 1000 Genome Project Data Processing Subgroup. 2009. The Sequence
Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078–2079.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**: 550.

Maciejowski J, Chatzipli A, Dananberg A, Chu K, Toufektchan E, Klimczak LJ, Gordenin DA,
Campbell PJ, de Lange T. 2020. APOBEC3-dependent kataegis and TREX1-driven
chromothripsis during telomere crisis. *Nat Genet* **52**: 884–890.

Martínez-Ruiz C, Black JRM, Puttick C, Hill MS, Demeulemeester J, Larose Cadieux E, Thol
K, Jones TP, Veeriah S, Naceur-Lombardelli C, et al. 2023. Genomic-transcriptomic
evolution in lung cancer and metastasis. *Nature* **616**: 543–552.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet.journal* **17**: 10–12.

Matsushima K, Yang D, Oppenheim JJ. 2022. Interleukin-8: An evolving chemokine.
Cytokine **153**: 155828.

Maynard A, McCoach CE, Rotow JK, Harris L, Haderk F, Kerr DL, Yu EA, Schenk EL, Tan
W, Zee A, et al. 2020. Therapy-Induced Evolution of Human Lung Cancer Revealed by
Single-Cell RNA Sequencing. *Cell* **182**: 1232–1251.e22.

McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K,
Altshuler D, Gabriel S, Daly M, et al. 2010. The Genome Analysis Toolkit: a MapReduce
framework for analyzing next-generation DNA sequencing data. *Genome Res* **20**: 1297–
1303.

Mihara E, Hirai H, Yamamoto H, Tamura-Kawakami K, Matano M, Kikuchi A, Sato T, Takagi
J. 2016. Active and water-soluble form of lipidated Wnt protein is maintained by a serum
glycoprotein afamin/α-albumin. https://elifesciences.org/articles/11621 (Accessed
August 30, 2024).

Nam CH, Youk J, Kim JY, Lim J, Park JW, Oh SA, Lee HJ, Park JW, Won H, Lee Y, et al.
2023. Widespread somatic L1 retrotransposition in normal colorectal epithelium. *Nature*
617: 540–547.

Naumann JA, Argyris PP, Carpenter MA, Gupta HB, Chen Y, Temiz NA, Zhou Y, Durfee C,
Proehl J, Koniar BL, et al. 2023. DNA Deamination Is Required for Human APOBEC3A-
Driven Hepatocellular Carcinoma In Vivo. *Int J Mol Sci* **24**.
http://dx.doi.org/10.3390/ijms24119305.

Ng PC, Henikoff S. 2003. SIFT: Predicting amino acid changes that affect protein function.
Nucleic Acids Res **31**: 3812–3814.

Nik-Zainal S, Alexandrov LB, Wedge DC, Van Loo P, Greenman CD, Raine K, Jones D,
Hinton J, Marshall J, Stebbings LA, et al. 2012a. Mutational processes molding the
genomes of 21 breast cancers. *Cell* **149**: 979–993.

Nik-Zainal S, Van Loo P, Wedge DC, Alexandrov LB, Greenman CD, Lau KW, Raine K,
Jones D, Marshall J, Ramakrishna M, et al. 2012b. The life history of 21 breast cancers.
Cell **149**: 994–1007.

Nordentoft I, Lamy P, Birkenkamp-Demtröder K, Shumansky K, Vang S, Hornshøj H, Juul M,
Villesen P, Hedegaard J, Roth A, et al. 2014. Mutational context and diverse clonal
development in early and late bladder cancer. *Cell Rep* **7**: 1649–1663.

Okazaki R, Okazaki T, Sakabe K, Sugimoto K, Sugino A. 1968. Mechanism of DNA chain
growth. I. Possible discontinuity and unusual secondary structure of newly synthesized
chains. *Proc Natl Acad Sci U S A* **59**: 598–605.

Otlu B, Díaz-Gay M, Vermes I, Bergstrom EN, Zhivagui M, Barnes M, Alexandrov LB. 2023.
Topography of mutational signatures in human cancer. *Cell Rep* **42**: 112930.

Pagel KA, Antaki D, Lian A, Mort M, Cooper DN, Sebat J, Iakoucheva LM, Mooney SD,
Radivojac P. 2019. Pathogenicity and functional impact of non-frameshifting
insertion/deletion variation in the human genome. *PLoS Comput Biol* **15**: e1007112.

Petljak M, Alexandrov LB, Brammeld JS, Price S, Wedge DC, Grossmann S, Dawson KJ, Ju
YS, Iorio F, Tubio JMC, et al. 2019. Characterizing Mutational Signatures in Human
Cancer Cell Lines Reveals Episodic APOBEC Mutagenesis. *Cell* **176**: 1282–1294.e20.

Petljak M, Dananberg A, Chu K, Bergstrom EN, Striepen J, von Morgen P, Chen Y, Shah H,
Sale JE, Alexandrov LB, et al. 2022. Mechanisms of APOBEC3 mutagenesis in human
cancer cells. *Nature* **607**: 799–807.

Pleguezuelos-Manzano C, Puschhof J, Rosendahl Huber A, van Hoeck A, Wood HM,
Nomburg J, Gurjao C, Manders F, Dalmasso G, Stege PB, et al. 2020. Mutational
signature in colorectal cancer caused by genotoxic pks E. coli. *Nature* **580**: 269–273.

Puram SV, Tirosh I, Parikh AS, Patel AP, Yizhak K, Gillespie S, Rodman C, Luo CL, Mroz
EA, Emerick KS, et al. 2017. Single-Cell Transcriptomic Analysis of Primary and
Metastatic Tumor Ecosystems in Head and Neck Cancer. *Cell* **171**: 1611–1624.e24.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**: 841–842.

Rausch T, Zichner T, Schlattl A, Stütz AM, Benes V, Korbel JO. 2012. DELLY: structural
variant discovery by integrated paired-end and split-read analysis. *Bioinformatics* **28**:
i333–i339.

Rhind N, Gilbert DM. 2013. DNA replication timing. *Cold Spring Harb Perspect Biol* **5**:
a010132.

Ritz C, Jensen SM, Gerhard D, Streibig JC. 2019. *Dose-Response Analysis Using R.* CRC
Press.

Roberts SA, Lawrence MS, Klimczak LJ, Grimm SA, Fargo D, Stojanov P, Kiezun A,
Kryukov GV, Carter SL, Saksena G, et al. 2013. An APOBEC cytidine deaminase

mutagenesis pattern is widespread in human cancers. *Nat Genet* **45**: 970–976.

Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, Mesirov JP.
2011. Integrative genomics viewer. *Nat Biotechnol* **29**: 24–26.

Saini N, Roberts SA, Sterling JF, Malc EP, Mieczkowski PA, Gordenin DA. 2017.
APOBEC3B cytidine deaminase targets the non-transcribed strand of tRNA genes in
yeast. *DNA Repair (Amst)* **53**: 4–14.

Sanchez A, Ortega P, Sakhtemani R, Manjunath L, Oh S, Bournique E, Becker A, Kim K,
Durfee C, Temiz NA, et al. 2024. Mesoscale DNA features impact APOBEC3A and
APOBEC3B deaminase activity and shape tumor mutational landscapes. *Nat Commun*
15: 2370.

Satija R, Farrell JA, Gennert D, Schier AF, Regev A. 2015. Spatial reconstruction of single-
cell gene expression data. *Nat Biotechnol* **33**: 495–502.

Sondka Z, Dhir NB, Carvalho-Silva D, Jupe S, Madhumita, McLaren K, Starkey M, Ward S,
Wilding J, Ahmed M, et al. 2024. COSMIC: a curated database of somatic variants and
clinical data for cancer. *Nucleic Acids Res* **52**: D1210–D1217.

Supek F, Lehner B. 2017. Clustered Mutation Signatures Reveal that Error-Prone DNA
Repair Targets Mutations to Active Genes. *Cell* **170**: 534–547.e23.

The ICGC/TCGA Pan-Cancer Analysis of Whole Genomes Consortium. 2020. Pan-cancer
analysis of whole genomes. *Nature* **578**: 82–93. https://doi.org/10.1038/s41586-020-
1969-6.

van Ineveld RL, Ariese HCR, Wehrens EJ, Dekkers JF, Rios AC. 2020. Single-Cell
Resolution Three-Dimensional Imaging of Intact Organoids. *J Vis Exp*.
http://dx.doi.org/10.3791/60709.

Venables WN, Ripley BD. 2003. *Modern Applied Statistics with S*. Springer Science &
Business Media.

Vieira VC, Soares MA. 2013. The role of cytidine deaminases on innate immune responses
against human viral infections. *Biomed Res Int* **2013**: 683095.

Wang K, Li M, Hakonarson H. 2010. ANNOVAR: functional annotation of genetic variants
from high-throughput sequencing data. *Nucleic Acids Res* **38**: e164.

Wang Y, Robinson PS, Coorens THH, Moore L, Lee-Six H, Noorani A, Sanders MA, Jung H,
Katainen R, Heuschkel R, et al. 2023. APOBEC mutagenesis is a common process in
normal human small intestine. *Nat Genet* **55**: 246–254.

Wilson MH, Coates CJ, George AL Jr. 2007. PiggyBac transposon-mediated gene transfer in
human cells. *Mol Ther* **15**: 139–145.

Woodard LE, Wilson MH. 2015. piggyBac-ing models and new therapeutic strategies.
Trends Biotechnol **33**: 525–533.

Wu H, Yu J, Li Y, Hou Q, Zhou R, Zhang N, Jing Z, Jiang M, Li Z, Hua Y, et al. 2018. Single-
cell RNA sequencing reveals diverse intratumoral heterogeneities and gene signatures
of two types of esophageal cancers. *Cancer Lett* **438**: 133–143.

Wu T, Lyu R, You Q, He C. 2020. Kethoxal-assisted single-stranded DNA sequencing
captures global transcription dynamics and enhancer activity in situ. *Nat Methods* **17**:

515–523.

Wu Y-Y, Tsai H-F, Lin W-C, Hsu P-I, Shun C-T, Wu M-S, Hsu P-N. 2007. Upregulation of
CCL20 and recruitment of CCR6+ gastric infiltrating lymphocytes in Helicobacter pylori
gastritis. *Infect Immun* **75**: 4357–4363.

Yoshida K, Gowers KHC, Lee-Six H, Chandrasekharan DP, Coorens T, Maughan EF, Beal
K, Menzies A, Millar FR, Anderson E, et al. 2020. Tobacco smoking and somatic
mutations in human bronchial epithelium. *Nature* **578**: 266–272.

Youk J, Kwon HW, Kim R, Ju YS. 2021. Dissecting single-cell genomes through the clonal
organoid technique. *Exp Mol Med* **53**: 1503–1511.

Youk J, Kwon HW, Lim J, Kim E, Kim T, Kim R, Park S, Yi K, Nam CH, Jeon S, et al. 2024.
Quantitative and qualitative mutational impact of ionizing radiation on normal cells. *Cell*
Genom **4**: 100499.

Figure 1

Figure 2

Figure 3

A

B

Likeley A3B exposed DNA fragments

Figure 4

Figure 5

Figure 6