Microbial lysates as low-cost serum replacements in cellular

agriculture media formulation

James Dolgin [1,4]*, Damayanti Chakravarty[2,4]*, Sean F. Sullivan[2,4], Yiming Cai[3,4],

Taehwan Lim[1,4], Pomaikaimaikalani Yamaguchi[2,4], Joseph E. Balkan[2], Licheng Xu[1,4],

Aaron D. Olawoyin[1,4], Kyongbum Lee[2,4], David L. Kaplan[1,4], Nikhil U. Nair[2,4]†

* equal contribution

† correspondence: nikhil.nair@tufts.edu

[1] Department of Biomedical Engineering, Tufts University, Medford, MA

[2] Department of Chemical & Biological Engineering, Tufts University, Medford, MA

[3] Department of Biology, Tufts University, Medford, MA

[4] Tufts University Center for Cellular Agriculture (TUCCA), Tufts University, Medford, MA

ABSTRACT

Cultivated meat, the process of generating meat in vitro without sacrificing animals, is a

promising alternative to the traditional practice of livestock agriculture. However, the

success of this field depends on finding sustainable and economical replacements for

animal-derived and expensive fetal bovine serum (FBS) that is typically used in cell

culture processes. Here, we outline an effective screening process to vet the suitability

of microbial lysates to support the growth of immortalized bovine satellite cells (iBSCs)

and mackerel (Mack1) cells. We show that easily producible, low-cost whole-cell lysates

from *Vibrio natriegens* can be used to create serum-free media for the long-term growth

of iBSCs. The optimized medium, named "VN40" (basal B8 media containing Vibrio

natriegens lysate proteins at 40 µg/mL), outperforms previously established serum-free

media while maintaining cell phenotype and myogenicity. Overall, this study shows a

novel approach to producing serum-free media for cultivated meat production using

microbially-derived lysates.

KEYWORDS

30 cultured meat, animal free meat, cultivated meat, serum free, media replacement,

31 proliferation, differentiation

1. INTRODUCTION

The environmental impact of meat consumption is expected to continue to rise with increasing demand and the increasing global population (OECD, Food, & Nations, 2021). The production of beef using conventional livestock agriculture is unsustainable due to greenhouse gas emissions, eutrophication, deforestation, overuse of water, and the spread of zoonotic disease (Hayek, 2022; Poore & Nemecek, 2018). Similarly, modern fishing practices have resulted in 90% of the world's fish stocks to be overexploited or collapse (Boyd et al., 2020; Grimaldo et al., 2023).

Cultivated meat – meat products generated from *in vitro* cultures of animal cells – represents a potentially sustainable alternative to conventional meat and fish production, which can reduce negative externalities and benefit animal welfare (Sinke et al. 2023; Treich, 2021). While conventional meat is responsible for 15% of global greenhouse gas emissions, cultivated meat can reduce this almost 13-fold (Sinke et al., 2023), especially through the use of renewable energy sources. Much progress has been made recently in developing stable immortalized lines of both bovine (Stout et al., 2023a) and mackerel (Saad et al., 2023) muscle satellite cells, leading to the creation of "iBSC" (Immortalized Bovine Satellite Cell) and "Mack1" cell lines, respectively. However, mammalian and marine cell cultures still largely depend on the use of fetal bovine serum (FBS), an expensive and animal-based ingredient with significant lot-to-lot variability (Zheng et al., 2006) and risks of contamination factors (Urz et al., 2022), making its use in cultivated meat infeasible. With 90% of the cost of cultivated meat production stemming from growth media (Specht, 2020), inexpensive FBS alternatives are needed to bring cultivated meat towards commercial production. Many attempts have been made to create serum-free media (SFM) for cultivated beef and fish with varying levels of success. Of note is recent work adapting serum-free "B8" media designed for human iPSC culture (Kuo et al., 2020) to the culture of bovine satellite cells (Stout et al., 2022) via the addition of recombinant albumin (rAlbumin), leading to the creation of "Beefy 9" medium. Still, efforts to replace FBS with recombinant albumin (Kolkmann, Van Essen, Post, & Moutsatsou, 2022; Stout et al., 2022) or other mitogenic proteins like fetuin (Skrivergaard et al., 2023) for cultivated meat production require expensive purified recombinant protein on the g/L scale. Current estimates show cultivated meat may require as much as 42 liters of media per kilogram to produce (Robert Vergeer, 2021), making the dependence on such quantities of recombinant proteins economically infeasible. Other methodologies using crude extracts or hydrolysates from plant (Stout et al., 2023) or microalgal (Dong et al., 2024; Okamoto et al., 2022) sources to supplement SFM have the potential to reduce raw material costs considerably. However, these methods can be labor-intensive, (Stout et al., 2023b; Ho et al., 2021), require corrosive chemicals (Okamoto et al., 2022), or are otherwise still

dependent on animal serum (Dong et al., 2024) or animal by-products (Andreassen et
al., 2020).

Microbes such as yeast, bacteria, and algae may provide a promising solution to the
longstanding problem of creating serum-free media formulations, as they are animal-
free, inexpensive to produce, highly renewable, and rich in proteins and other nutrients.
Microbes can also be engineered to express growth factors, which currently represent
99% of the cost of cultured meat media at scale (Specht, 2020). Yeast hydrolysates
have longstanding use in creating serum-free media (Chevalot et al., 2010; Sung et al.,
2004), and recent reports detail using extracts from *Saccharomyces cerevisiae* (Lei et
al., 2023), *Enterococcus hirae* (Celebi-Birand et al., 2023), cyanobacteria (Ghosh et al.,
2023; Jeong et al., 2021; Tuomisto & de Mattos, 2011), and *Chlorella* algae (Dong et al.,
2023; Okamoto et al., 2022) in cultured meat production. However, current research has
not demonstrated success in identifying a non-pathogenic, rapidly replicating microbial
extract capable of stimulating long-term cellular growth in serum-free conditions.
Microalgae are slow-growing, taking several days to culture (Okamoto et al., 2022),
while cyanobacteria can have toxic effects on cells (Chia et al., 2018). Yeast and
bacterial extracts have not been shown to completely substitute for FBS or support
long-term growth of animal cells (Celebi-Birand et al., 2023; Chia et al., 2018). A more
comprehensive list of previous attempts to replace serum in cultivated meat media is in
Table S1.

Media to produce cultivated meat must strive to minimize time, labor, and material costs
in order to make cultivated meat economically viable. Such media must also be animal-
free and safe to consume and produce. Finally, cultivated meat media must stimulate
long-term cell growth with minimal impacts on cell characteristics. While microbial
extracts are a promising ingredient for replacing FBS in cell culture media, there have
not been any results that satisfy the criteria necessary for commercial use in cultivated
meat production. Further, it is known that media requirements vary significantly
depending on cell type (O'Neill et al., 2022).

In this work, we tested multiple lysates from Gram-positive and Gram-negative bacteria,
as well as a yeast, to determine their suitability as replacements for FBS in bovine and
fish cell culture media. Based on an initial screen, we found that lysates derived from
Vibrio natriegens could serve as an alternative to FBS for iBSCs. A similar screen also
identified that lysates from *Saccharomyces cerevisiae* could support growth of Mack1
cells in reduced serum conditions. We show that iBSCs can be adapted to grow rapidly
in *Vibrio natriegens* lysate-containing media by long-term and short-term serial
passaging. Cells cultured in this microbial lysate media (called VN40) demonstrate
robust growth, preserve their satellite cell phenotype, and the ability to differentiate into

multinucleated myotubes. These results highlight the promise of the whole-cell lysate
derived from the fast-growing non-pathogenic marine bacterium, *V. natriegens*, as an
easily producible, low-cost supplement for long-term serum-free growth of iBSCs for
cultivated meat. Further, our approach may be generalizable to help identify novel
microbial lysates as FBS substituents for different cultivated meat cell lines.
116
117

2. MATERIAL & METHODS

2.1 Culture and extraction of microbial lysate

Escherichia coli BL21(DE3) and Nissle 1917 ("BL21" and "ECN" respectively), *Vibrio natriegens* ATCC14048 ("VN"), *Lactobacillus plantarum* WCFS1 ("LP"), *Bacillus subtilis* 168 ("BS")*, Lactococcus lactis* MG1363 ("LL") and *Saccharomyces cerevisiae* CEN.PK ("CEN.PK") were streaked out on plates using their respective solid media (**Table 1**). Single colonies from the plates were used to inoculate 25 mL of overnight cultures. The next day, cells at stationary phase were harvested by centrifuging at 4000 rpm for 5 min (ThermoScientific Sorvall Legend X1R Centrifuge). Harvested cells were prepared for sonication by washing and resuspending in 1 mL sterile phosphate buffer saline (PBS), without any additives. To generate microbial lysates, cells were sonicated (Branson 150 sonicator and 10 s Pulse, 30 s gap between each pulse and a total 7 min at 45 % amplitude) on ice. After sonication, the soluble lysate was obtained by centrifuging the sonicated solution at 21000 × g for 20 min. The supernatant was then filtered through a 0.2 µm syringe filter (Corning, #431219). The total protein content of the lysate was determined using Bradford Assay (Coomasie Plus^{TM} Protein Assay Reagent, #1856210) (**Figure 1A**).

Microorganisms	Medium	Growth Conditions
E. coli BL21(DE3) (BL21), *E. coli Nissle 1917* (ECN)*, Bacillus subtilis* 168 (BS)	Lysogeny Broth (LB)	Overnight at 37 °C
Vibrio natriegens ATCC14048 (VN)	LB with 5 % NaCl	Overnight at 37 °C
Lactiplantibacillus plantarum WCFS1 (LP), *Lactococcus Lactis* MG1363 (LL)	De Man–Rogosa–Sharpe (MRS)	Overnight at 37 °C
S. cerevisiae CEN.PK (CEN.PK)	Yeast peptone adenine dextrose (YPAD)	Overnight at 30 °C

Table 1. Media and growth conditions for microbes cultivated for lysate generation.

2.2 SDS-PAGE

Denaturing SDS-PAGE was performed following the Invitrogen NuPAGE® protocol. Lysate samples containing 30 µg of protein were adjusted to a final volume of 50 µL with PBS and mixed with 10 µL of 6× SDS loading buffer. The samples were incubated at 95 °C for 10 min. Subsequently, 10 µL of the sample was loaded onto a pre-cast NuPAGE Novex 4-12% Bis-Tris 1.0 mm mini-gel (Invitrogen). For each gel run, 5 µL of pre-stained SDS-PAGE standards (Bio-Rad) were included. Electrophoresis was

146 conducted at a constant voltage (200 V) in 1× MES (2-morpholinoethanesulphonic acid)
147 buffer solution at room temperature for approximately 45 min.
148
2.3 Lysate fractionation
Vibrio natriegens lysate was fractionated using a 3 kDa molecular weight cutoff filter
(MilliporeSigma™ Amicon™ Ultra-15 Centrifugal Filter) at 4000 × g. This resulted in two
fractions, i.e., protein rich fraction greater than 3 kDa (HMW fraction) and low molecular
weight fraction lower than 3 kDa (LMW fraction). The fractions were then tested
individually on iBSC cells. To test for additive effect of fractions, we used **Equation 1**
and **Equation 2** to determine the sum of means (*µ*) and standard deviations (*σ*),
respectively, of fractions' impact on cell growth. These effects were compared to the
impact of whole cell lysates, which are HMW and LMW fractions combined
158

$$\sum \mu(LMW, HMW) = \mu(LMW) + \mu(HMW) \qquad \textbf{(Equation 1)}$$

$$\sum \sigma(LMW, HMW) = \sqrt{\sigma^2(LMW) + \sigma^2(HMW)} \qquad \textbf{(Equation 2)}$$

161
2.4 iBSC and Mack1 culture and maintenance
iBSCs were previously isolated from the semitendinosus of a 5-week-old Simmental calf
at the Tufts Cummings School of Veterinary Medicine (Stout et al., 2022).These cells
were engineered to constitutively express bovine telomerase reverse transcriptase
(TERT) and cyclin-dependent kinase 4 (CDK4) in prior studies (Stout et al., 2023a).
iBSCs at passage 40 were thawed from liquid nitrogen stock and passaged in iBSC
growth media (iBSCGM) containing DMEM+Glutamax (ThermoFisher #10566024,
Waltham, MA, USA) with 20 % FBS (ThermoFisher #26140079), 1 ng/mL human
fibroblast growth factor 2 (FGF-2) (PeproTech #100-18B, Rocky Hill, NJ, USA),
supplemented with 1 % antibiotic/antimycotic (ThermoFisher #1540062) and 2.5 µg/mL
puromycin (ThermoFisher #A1113803) as a selection compound. iBSCs were cultured
at 37 °C and 5 % humidity on tissue culture polystyrene (TCPS) and routinely passaged
at 80–90 % confluency using 0.25 % trypsin-EDTA (ThermoFisher #25200056) to
disassociate cells from TCPS. During passaging, cells were dissociated, centrifuged at
300 × g, and seeded at 1,500–2,000 cells/cm^2. Mack1 cells (Kerafast, #ETU008-FP)
isolated from Atlantic mackerel were cultured in Mack growth media (MackGM)
containing Leiovitz's L-15 medium (ThermoFisher #11415064) with 20 % FBS, 20 mM
HEPES (Sigma Aldrich #H4034, St. Louis, MO, USA) buffer solution (pH 7.4), 1 %
antibiotic/antimycotic, 1 ng/mL human FGF-2, and 10 µg/mL gentamicin (Sigma Aldrich
#G1397)(Boyd et al., 2020; Saad et al., 2023). Mack1 cells were incubated at 27 °C
without CO_2 on TCPS, routinely detached at 70–80 % confluency using 0.05% trypsin-
EDTA (ThermoFisher #25300054), centrifuged at 300 × g, and seeded at 3,500–4,000
cells/cm^2. Both cell lines (iBSCs and Mack1) were fed every 2-3 days with iBSCGM and
MackGM, respectively.

2.5 Short-term iBSC and Mack1 cell growth screening with microbial lysates

For iBSC short-term screening on microbial lysate-based media, iBSCs were harvested and resuspended into serum-free B8 media containing Hi-Def B8 aliquots (Defined Bioscience #LSS-204, San Diego, CA, USA) in DMEM/F12 (ThermoFisher #11320032) supplemented with 1 % antibiotic/antimycotic and 2.5 µg/mL puromycin. Cells were seeded onto 96-well plates at 2,500 cells/cm^2 with 1.5 µg/cm^2 of truncated recombinant human vitronectin (rhVTN-N; ThermoFisher #A31804). Cells were allowed to adhere for >3 h, then were washed once with DPBS (ThermoFisher #14190250) and media was changed to B8 + microbial lysates at given protein concentrations. Negative control conditions were fed B8 alone. Positive control conditions were fed Beefy 9, containing B8 supplemented with 0.8 mg/mL of recombinant albumin (rAlbumin, Sigma Aldrich #A9731)(Stout et al., 2022). Mack1 cells were harvested and resuspended in Mack1GM and plated onto 96-well plates at 2,500 cells/cm^2. After more than 3 h incubation, cells were washed once with DPBS and media was changed to 2.5 % FBS-containing L-15 media (other media components, including HEPES, antibiotic/antimycotic, FGF-2, and gentamicin, were added at the same concentrations as Mack1GM) + microbial lysates at give protein concentrations. Negative controls were Mack1GM without FBS and 2.5% FBS-containing L-15 media. Media formulations without cells were used as blanks to account for media autofluorescence. All samples were allowed to grow undisturbed for 4 (Mack1) and 5 days (iBSCs), at which point wells were washed once with DPBS and frozen overnight at –80°C. Samples were then analyzed for short-term growth using direct DNA quantification with CyQUANT Cell Proliferation Assay (ThermoFisher #C7026) using manufacturer's protocol. Emission/excitation was read at 480/520 nm on a ThermoFisher Varioskan LUX multimode microplate reader. Blank values from media-only samples were subtracted from experimental values, and these values were normalized to growth using either B8 alone (iBSCs) or 2.5 % FBS-containing L-15 media (Mack1) to compare the growth between conditions. Experiments were repeated using a narrower range of lysate concentration and using serum-free adapted iBSCs passaged 4 times consecutively in Beefy 9.

2.6 Adaptation and long-term growth in *Vibrio natriegens*-based media

After choosing *V. natriegens* 10 mg/mL serum-free media as a starting point for adaptation based on short-term screens, iBSCs, now at passage 60, were resuspended and seeded at 2,000 cells/cm^2 in B8 media with 1.5 µg/cm^2 vitronectin in a 6-well plate. After adhering for >3 h, cells were fed 2 mL B8 medium supplemented with 10 mg/mL *V. natriegens* lysate. Cells were fed *V. natriegens* lysate-based media every 2-3 days until they reached 80-90 % confluence. At this point, cells were passaged using TrypLE Express enzyme (ThermoFisher #12604013) and counted using a NucleoCounter NC-

225 200 (Chemometec, Allerød, Denmark). Doubling time (DT) was calculated using
226 **Equation 3**.
227
$$DT = \frac{\Delta t}{\log_2\left(\frac{C_f}{C_i}\right)}$$ (**Equation 3**)

with Δt being time between passages, C_f being cell count at passage, and C_i being total
cells seeded. Cells were then centrifuged at 300 × g and seeded in a new well. This
process was repeated continuously. When doubling time reached a minimum or
plateaued, *V. natriegens* concentration was doubled, until doubling time was
comparable to that of iBSCGM, which was when *V. natriegens* was at 40 mg/mL (VN40).
At this point, VN40-adapted cells were seeded in triplicate in a 6-well plate and fed
VN40. Cell doublings were measured at each passage for each replicate. This same
process was repeated simultaneously for Beefy 9-adapted cells fed with Beefy 9 (as
control). Cells were periodically imaged using phase contrast microscopy (KEYENCE,
BZ-X700, Osaka, Japan; Olympus, CKX53, Tokyo, Japan)
239
2.7 Characterization and differentiation of VN40 adapted cells
After 20 passages in VN40 media, iBSCs were allowed to adhere in triplicate in 48-well
plates at 6000 cells/cm^2 and their satellite cell identity was characterized via staining for
Paired-box 7 (Pax7). Cells were fixed with 4 % paraformaldehyde in PBS
(ThermoFisher #AAJ61899AK) for 20 min, washed with DPBS containing 0.1 % Tween-
20 (ThermoFisher, BP337), permeabilized for 10 min with DPBS containing 0.1 % Triton
X-100 (ThermoFisher, BP151), washed again with DPBS+Tween-20, and blocked for 30
min using a blocking solution containing DPBS with 5 % goat serum (ThermoFisher
#16210064) and 0.05 % sodium azide (Sigma #S2002). A primary antibody solution
containing 1:500 Pax7 antibody (ThermoFisher #PA5-68506) and 1:1000 actin stain
Phalloidin 488 (ThermoFisher A12379) in blocking solution was added to cells and
incubated overnight at 4°C. Cells were then washed with DPBS+Tween-20 and blocked
for 30 min once again. A solution containing 1:500 Pax7 secondary antibody
(ThermoFisher #A11072) and 1:1000 nuclear stain DAPI (ThermoFisher #D1306) in
blocking solution was added to cells and incubated for 1 h at room temperature. Cells
were then washed with DPBS+Tween-20 and kept in DPBS for imaging. Imaging was
carried out using fluorescence microscopy (KEYENCE, BZ-X700, Osaka, Japan).
Stemness was quantified using a Celigo automated image cytometer (Revvity, Waltham,
MA) to determine cells positive for Pax7. This was repeated for cells maintained in
iBSCGM without VN40 adaptation to determine if adaptation altered iBSC satellite cell
phenotype. Cells were differentiated by seeding at 5,000 cells/cm^2 in 48-well plates,
feeding with VN40 media, and initiating differentiation when cells appeared confluent ~4
days after initial seeding. Differentiation media contained DMEM+Glutamax
supplemented with 2 % horse serum (ThermoFisher #16050130), 0.5 mg/mL

recombinant human albumin, 1× ITS-X (ThermoFisher #51400056), 0.5 mM
LDN193189 (Sigma #SML0559), 1 % antibiotic/antimycotic, and 2.5 µg/mL puromycin.
Differentiation continued for 18 days with cells fed fresh differentiation media every 2–3
days. After mature myotubes were observed, cells were fixed, permeabilized, and
blocked as described. Cells were then stained using myosin heavy chain (MHC;
Developmental studies hybridoma bank #MF-20, Iowa City, IA, USA) and 1:200 desmin
(Abcam #ab15200, Cambridge, UK) primary antibodies diluted in blocking solution.
Secondary antibodies for MHC (ThermoFisher #A32723; 1:200) and desmin
(ThermoFisher #A11072; 1:500) along with 1:1,000 DAPI, diluted in blocking buffer,
were then applied as described. Imaging was carried out using fluorescence microscopy
(KEYENCE, BZ-X700, Osaka, Japan).
275
2.8 Sample preparation for metabolomics analysis
A 100 % methanol extraction was performed on the lysate. Methanol and VN lysate
were mixed in a 3:1 proportion and vortexed. The mixture was centrifuged at 15,000 × g
at 4 °C for 15 min. Then, 250 µL of the supernatant was transferred to each Eppendorf
tube. The samples were dried by speed vacuuming (Eppendorf Vacufuge^{TM}) and stored
at –20 °C. Prior to the LC-MS run, samples were reconstituted by adding methanol and
water at 1:1 proportion to dried pellet, mixed by vortexing, and then centrifuged at
15,000 x g at 4 °C for 15 min (ThermoScientific Sorvall Legend X1R Centrifuge). Finally,
100 µL of the supernatant was transferred to sample tubes for LC-MS.
285
2.9 Liquid chromatography-tandem mass spectrometry (LC-MS/MS)
Untargeted analysis of metabolites present in the extracted lysate samples was
preformed using information-dependent acquisition (IDA) experiments on a quadrupole
time-of flight (TOF) mass spectrometer (TripleTOF 5600+, AB Sciex, Framingham, MA,
USA) with an electrospray ionization (ESI) source. IDA experiments include a single
TOF MS scan corresponding to four dependent product ion (MS/MS) scans based on
the highest intensity unique mass. Fragmentation was triggered when precursor ion
counts rose quickly over several scans to ensure ions were selected near the top of
their LC peaks. A total of four methods was applied to each sample using either a
hydrophilic interaction chromatography (HILIC) column (Phenomenex Luna NH2,
Torrance, CA, USA) or a reverse-phase column (Phenomenex Synergi Hydro-RP,
Torrance, CA, USA) in both positive and negative ionization modes. The
chromatographic separation was performed with a binary pump HPLC system (1260
Infinity, Agilent, Santa Clara, CA, USA). Chromatographic gradient methods and mobile
phases described in supplementary information.
301
302 **2.10 Feature annotation**

LC-MS features were annotated using the Biologically Consistent Annotation of Metabolomics Data computational tool (BioCAn)(Alden et al., 2017). Briefly, BioCAn assess potential annotations of *m/z* values and MS/MS spectra using the following annotation tools: CFM-ID(Allen, Pon, Wilson, Greiner, & Wishart, 2014), NIST17 (NIST/EPA/NIH, 2017), MONA (https://massbank.us/), and HMDB(Wishart et al., 2013) Features could be assigned multiple annotations either by a single annotation tool or by disagreement between assignments designated by different tools. To determine the most likely annotation for a feature, the feature was mapped to a metabolite in the *V. natriegens* metabolic network and assigned an annotation score by BioCAn. Annotation scores were based on agreement of the annotation assignment by the integrated databases as well as confidence in annotation of neighboring metabolites within the metabolic network.

2.11 Metabolomics data analysis

The peak area corresponding to the integrated area under the curve (AUC) of each ion chromatograph was blank subtracted to reduce noise. Each peak area was then normalized to the total ion chromatogram (TIC) for each replicate (n = 3). Unannotated features were subsequently removed. In addition, features showing annotation scores < 1.0 were removed due to the low confidence in the annotation. Finally, annotated features showing negative (blank subtracted) peak areas were removed as their concentration is likely lower than the limit of detection for the applied method. The remaining metabolites were then sorted by peak area.

2.12 Cost analysis

We compared the cost of VN lysate containing media with BSCGM and B9 media. The details of the cost comparison are in supplementary information.

3. RESULTS

3.1 Mechanical lysis of gram-negative bacteria yields extracts with high protein content.

The overall process of lysate production and screening on iBSCs and the Mack1 cell line is illustrated in the schematic diagram shown in **Figure 1A**. The Gram-negative bacteria *E. coli* Nissle ("ECN"), *E. coli* BL21 ("BL21"), and *V. natriegens* ("VN"), along with the yeast strain *S. cerevisiae* CEN.PK ("CEN.PK"), demonstrate relatively higher protein yields per liter of culture compared to Gram-positive bacteria *L. lactis* ("LL") and *L. plantarum* ("LP") (**Figure 1B**). This discrepancy can be attributed to their thick peptidoglycan cell wall, which makes them more resistant to lysis (de Bruin et al., 2019).

We analyzed the protein profiles of lysates obtained from the microbial strains using SDS-PAGE to assess their composition. **Figure 1C** illustrates the results for the lysates

from *V. natriegens*, both *E. coli* strains, *B. subtilis*, *L. plantarum*, *L. lactis*, and *S. cerevisiae*. Each lane shows distinct protein banding patterns, highlighting the differences in protein composition across species.

Figure 1. Workflow – screening of lysates from various microorganisms for muscle satellite cell performance. A) Overview of microbial workflow: microbial biomass production, lysate generation by mechanical lysis, and screening for proliferative activity on animal cells. After filter sterilization, the lysate is added to basal media, which is followed by estimating cell proliferation and differentiation. **B)** Higher protein yield resulted from Gram negative bacteria and yeast compared to gram positive bacteria. **C)** SDS-PAGE profile of seven lysates used in the initial screening. (L-R) Ladder, *B. subtilis* 168 (BS), *E. coli* Nissle 1917 (ECN), *V. natriegens* ATCC14048 (VN), *E. coli* BL21(DE3) (BL21), *L. lactis* MG1363 (LL), *S. cerevisiae* CEN.PK (CEN.PK), and *L. plantarum* WCFS1 (LP).

3.2 Screening helps identify compatibility between lysates and cell lines.
The iBSCs used in this study were extensively characterized in our previous studies
(Stout et al., 2023a; Stout et al., 2024). As an initial screen for iBSC growth in serum-
free lysate-based media, cells were tested directly after passaging in serum-containing
growth medium (iBSCGM). Performance (cell growth) was compared to previously
established Beefy 9 serum-free medium (Stout et al., 2022), which contains similar
ingredients as our tested formulations, except the recombinant human albumin present
in Beefy 9 was replaced with microbial lysates. The results (**Figure 2A**) show that many
of the lysates were inhibitory to cell growth after 5 days of exposure. All lysates elicited
decreased cell growth at higher concentration (up to 500 μg/mL). Lysates from VN (10
μg/mL) and LP (10 and 100 μg/mL) exhibited short-term cell growth similar to that of the
Beefy 9 control.

We also screened the short-term growth-promoting effects of lysate-based reduced-
serum (2.5% FBS) media on Mack1 cells (**Figure 2B**). Given that all lysates at 500
μg/mL showed detrimental effects on iBSC growth, we tested a lower range of lysate
protein concentrations for Mack1 cells. Unlike with iBSCs, VN at concentrations above
10 μg/mL and LP at all concentrations inhibited short-term Mack1 cell growth. However,
CEN.PK, BL21, and ECN lysates significantly improved the short-term growth of the
cells. Specifically, Mack1 cells performed best in the presence of CEN.PK (up to 100
μg/mL), BL21 (10 μg/mL), and ECN (10 μg/mL and 50 μg/mL) lysates. At 100 μg/mL,
CEN.PK lysate nearly doubled short-term growth of Mack1 cells, having a similar effect
as 7.5% FBS-containing media as shown by Lim et al. (2024), suggesting this lysate
can replace 5% FBS in media without detrimental effect.

Despite our success with Mack1 cells, we found greater success in adaptation of iBSCs
to serum-free media using lysates and proceeded with subsequent studies using only
iBSCs. Since serum-free adaptation significantly alters cellular phenotype and gene
expression (J.Y. Lee et al., 2022), transferring cells directly from serum-containing
media to lysates may not be an appropriate strategy to assess the long-term potential of
lysates. So, next we re-screened the lysates using iBSCs grown in Beefy 9 (serum-free)
medium (4 passages) with select lysates (**Figure 2C).** The bovine cells were grown in
lysate concentrations lower than those of the first screen, since concentrations above
10 μg/mL were shown to be cytotoxic. The results showed that cells cultured in serum-
free conditions responded differently to lysates than those cultured only in serum-
containing media. Cells cultured in VN media at 10 μg/mL outperformed those grown at
this concentration of BS and LP. Cell growth improved with increased VN lysate
concentration between 2.5 and 10 μg/mL, while the opposite effect was seen in cells
grown in LP and BS. While initial screens found that non-adapted cells proliferated in LP
media, these results did not translate to cells adapted to serum-free media. Given that

386   VN performed well with serum adapted and un-adapted cells, we chose it for further

387   investigations.

Figure 2. Short-term proliferation assay of iBSCs and Mack1 cells using various microbial lysates. A) iBSC (not serum-free adapted) proliferation screen (5-day incubation) on microbial lysate-based serum-free media shows VN and LP lysates performing similarly to Beefy 9 medium (positive control). B8 medium (Beefy 9 without rAlbumin) was used as negative control. Lysate-based media created through addition of respective lysates to B8 media. Data is from CyQuant DNA staining kit. n=4 (experimental groups) or 10 (controls). **B)** Mack1 (not serum-free adapted) proliferation screen using an L-15 medium supplemented with 2.5 % FBS (negative control) and additional microbial lysates. n=3. CEN.PK was well tolerated at different concentrations. **C)** iBSC (4× serum-free Beefy 9 passages) proliferation screen using selected lysates at lower concentrations. n=4 (experimental groups) or 10 (controls). Lysate formulations were compared to one another to determine candidates for long-term growth studies, showing VN as top contender. All presented data is mean values ± SD. Statistical analysis by unpaired t-test with Welch's correction in which p > 0.05 (NS), p < 0.05 (*), 0.01 (**), and 0.001(***).

3.3 iBSCs can adapt to long-term growth on *V. natriegens* lysate containing media.
iBSCs were cultured directly from BSCGM into serum-free B8 medium supplemented with 10 μg/mL VN lysate. Growth was monitored throughout several passages and lysate concentration in feeding media was increased when cell growth slowed. This strategy led to increases in cell growth rate and was used to slowly adapt cells to a medium with VN lysate protein concentration of 40 μg/mL (**Figure 3A**), dubbed "VN40". At this concentration, cell growth rate plateaued, and doubling time reached 24.9 h, similar to those observed using serum-containing BSCGM (21.9 ± 3.1 h) during the course of the study. Overall growth kinetics of iBSCs grown in VN40 over 25 days were similar to those grown in 20% FBS BSCGM medium – ~19 total cell doublings in VN40 compared to ~22 in BSCGM (Stout et al., 2022). This is a significant increase from the ~14 doublings shown in Beefy-9 and implies VN lysate can completely replace FBS in iBSC growth media. Cells showed an increase in viability between 10 μg/mL VN feeding (avg. 71.5 % viability) versus 40 μg/mL feeding (avg. 93.9 % viability), though this may have been a factor of initial population enrichment during adaptation to lysate-based medium. In all, the process of adapting cells to VN40 took approximately 2.5 months (passage 1 to passage 14), in line with previous durations of adapting mammalian cells to serum-free media (Hartman et al., 2007; Hua et al., 2022; Miki & Takagi, 2015). Adaptation of iBSCs to VN40 was expedited by passaging cells directly into 20 μg/mL VN medium (**Figure 3B**), reducing total adaptation time to 4 weeks. After adaptation of cells in VN40 medium, cells were grown in triplicate to compare to growth in serum-free Beefy 9 media. Both cell populations were passaged continuously over 24 days with collective cell doublings recorded at each passage (**Figure 3C**). Cells grown in VN40 significantly outgrew cells grown in Beefy 9, with higher collective cell doublings and lower average doubling times (31.2 ± 4.8 h for VN40; 43.1 ± 8.7 h for Beefy 9). An increase of VN lysate content in the media beyond 40 μg/mL hindered cell growth compared to VN40 (**Figure S1**). Cells had higher confluence after 3 days of growth in VN40 compared to Beefy 9 (**Figure 3D-E**). VN40 also resulted in less observable lipid accumulation compared to Beefy 9 (**Figure 3F-G**). Beefy 9 has been shown to cause aberrant lipid accumulation (Stout et al., 2022), potentially due to induced insulin resistance from the high concentration of insulin in B8, or via adipogenic signaling. Components of VN40 may be preventing such insulin resistance and/or adipogenic signaling from taking place, which could be beneficial in long-term maintenance of myogenicty and proliferative capacity. Cells in VN40 have maintained proliferation over 108 passages and can survive long-term storage and thawing from liquid nitrogen.

3.4 High and low molecular weight components of *V. natriegens* lysate are additive in promoting iBSC proliferation.
To determine if a specific molecular weight fraction contributed to the mitogenic effect of VN lysate-based media, we fractionated the lysate by ultrafiltration using a 3 kDa

Figure 3. Adaptation and long-term growth of iBSCs in VN40. A) iBSCs grown over multiple passages in B8 medium containing increasingly concentrated *V. natriegens* lysate (corresponding to 10, 20, 40 µg/mL protein concentrations). The cells were grown in T-25 flasks and passaged at 80–90 % confluence. Dotted line representing doubling times in Beefy 9 & BSCGM (General Media for iBSCs containing 20 % FBS) collected throughout this study. **B)** iBSC adaptation to *V. natriegens* lysate-containing media beginning with 20 µg/mL protein concentration. **C)** iBSCs grown continuously in VN40 and Beefy 9 media with cells counted at each passage to determine cumulative cell doublings. Cells grown in triplicates in 6 well plates for each condition. Significance indicated for final cumulative cell doublings. All presented data is mean values ± SD. Statistical analysis by unpaired t-test with Welch's correction in which p < 0.001 (***). **D-E)** Cells after the final experimental passage (passage 82 overall, passage 22 in serum-free media) grown for 3 days and imaged by phase contrast microscopy, showing improved growth of VN40 (**D**) compared to Beefy 9 (**E**). Scale bar = 200 µm. **F-G)** Magnification increased to show additional detail, with orange arrows indicating increased cytoplasmic lipid accumulation in cells grown in Beefy 9 (**G**) compared to VN40 (**F**). Cells were quantified manually or using an automated cell counter. Scale bar = 100 µm.

428
molecular weight cutoff membrane. iBSC growth was determined via the same protocols

as earlier, with short term growth compared to that in B8 (negative control) and Beefy 9
(positive control). Formulations using the protein-rich high molecular weight (HMW) >3
kDa fraction, low molecular weight (LMW) <3 kDa fraction, and whole-cell VN lysate,
were prepared at various concentrations in B8 media. For total lysate
and HMW fractions, protein concentrations were determined using Bradford assay, as
described previously. Since the protein component of the LMW fraction was below the
detection limit for Bradford assay, we used the same composition as would have been
required for the whole cell lysate. The results (**Figure 4A**) showed that for
concentrations of VN protein equal to or above 40 µg/mL, whole cell lysates out-
performed HMW or LMW fractions alone. It can be concluded from this data that no
individual fraction of VN lysate was cytotoxic compared to whole-cell lysates, and that
larger proteins act together with small molecule metabolites to have an overall pro-
mitogenic effect on iBSCs. This has been shown in prior work in which metabolomics
and proteomics showed the indispensability of both amino acid and protein fractions in
Auxenochlorella pyrenoidosa extracts for goldfish muscle cell growth (Dong et al., 2023).
Whole-cell lysates of VN40 outperformed individual fractions, suggesting that the
proteins and metabolite fractions have an additive impact (**Figure 4B**). Ultimately, our
data shows the importance of using unfractionated VN40 in stimulating iBSC growth.
3.5 VN40 cultured iBSC maintain stemness and can be differentiated.

Figure 4. Assessing the role of HMW and LMW *Vibrio natriegens* lysate fractions on iBSC growth. A) The growth of iBSCs using a short-term screen (5 days) demonstrates that whole-cell lysates support growth more effectively than either the >3 kDa (HMW) or <3 kDa (LMW) fractions alone. HMW and LMW samples represent retentate and filtrate, respectively, that were added to B8 media at protein concentrations indicated (<3kDa concentration assessed by adding low molecular weight fraction representative of corresponding whole cell lysate). Beefy-9 media used as positive control, B8 media used as negative control. Data is from CyQuant DNA staining kit. **B)** iBSC proliferation in whole-cell VN40 medium compared to the sum of HMW and LMW growth indicates additivity of their relative contributions to iBSC growth. All presented data is mean values ± SD. Statistical analysis by unpaired t-test with Welch's correction in which $p > 0.05$ (NS), $p < 0.01$ (**).

We validated continued satellite cell phenotype via Pax7 expression (Relaix et al., 2005)
of iBSCs after 22 passages in VN40, showing that expression was maintained
throughout serum-free adaptation (**Figure 5A-B**). According to quantitative imaging
cytometry, cells were 99.7% positive for Pax7 after adaptation (**Figure 5C**), indicating
maintenance of a pure satellite cell population. After validation of phenotype, we
successfully differentiated VN-40 adapted cells into mature multi-nucleated myotubes
expressing both Desmin and Myosin Heavy Chain proteins (**Figures 5D-K**), indicative of
early-stage and late-stage differentiation respectively (Capetanaki et al., 2022). This
implies that VN40-adapted cells maintain their myogenicity after extensive serum-free
passaging, essential for creating functional cultivated meat products (M. Lee et al.,
2024) . Cells differentiated after slow, gradual adaptation (10 μg/mL → 20 μg/mL → 40
μg/mL VN lysate) (**Figure 5D-G**) differentiated more slowly and less robustly than cells
differentiated after a quicker adaptation (20 μg/mL → 40 μg/mL) (**Figure 5H-K**). This is
likely due to the impact of higher passage number on myogenicity (Park et al., 2023).
Differentiation media was optimized to contain ITS-X (Messmer et al., 2022) and LDN-
193189 (Horbelt et al., 2015), which were necessary for differentiating these overall high
passage (>p70) cells. Ultimately, we have arrived at a bacterial lysate-based serum-free
formulation which allows for successful proliferation and differentiation of iBSCs.
467
3.6 LC-MS/MS identifies major metabolites in *V. natriegens* lysate.
VN lysate was analyzed for metabolite composition using a series of untargeted LC-
MS/MS assays. The untargeted analysis detected a total of 11,306 features using a
combination of HILIC and reverse phase chromatography in both positive and negative
ionization modes (four total methods) The number of detected features in our study
aligns with typical results from untargeted metabolomic analyses. This is due to the
generation of degenerate features, such as those resulting from adduct formation,
dimerization, isotopic variants, and fragmentation variation, which can inflate the total
number of features derived from single unique metabolites by up to 90% (Mahieu et al.,
2017). A previously developed automated tool (BioCAn) was used to annotate the
features, resulting in a total of 403 metabolite annotations that mapped to known
metabolites in VN. Similar network-based feature annotation tools demonstrate
comparable identification capacities to BioCAn. For instance, MetDNA utilizes a reaction
network-based recursive algorithm to sequentially identify features utilizing a surrogate
MS/MS spectra from an initial set of seed metabolites. In a study on *Drosophila* samples,
MetDNA annotated 1,314 metabolites out of 18,320 detected features with an accuracy
of 77.5% (Shen et al., 2019). A significant limitation of network-based annotation
algorithms is the dependence on the understanding of the organism's metabolic network.
As a result, model organisms with more comprehensive and curated genomic
information and metabolic function tend to achieve greater metabolite coverage, while
lesser studied organisms, such as *V. natriegens,* may face more limited metabolite

annotations. Although BioCAn uses biological network data to reduce false positives,
errors may still occur. Thus, only annotations with a score greater than 1.0 were further
considered. An annotation score >1.0 indicates strong agreement across integrated
databases, as well as similar agreement in the annotation of metabolites within two
reactions of the annotation in the metabolic network. A total of 155 metabolites were
identified across the 4 methods with annotation scores ≥1.0 (**Table 2**), which included
80 unique metabolites. An analysis of commonality between top metabolites detected by
each method can be seen in **Figure S2.** Interestingly, purine and pyrimidine derivates
and aromatic amino acids show the greatest area under the curve, suggesting
potentially elevated concentrations of these metabolites in VN lysate extracts.
499
3.7 VN40 component costs at lab scale are competitive relative with Beefy 9.
Overall production cost of cultivated meat includes at least three main factors –
bioreactors, labor, and cell culture medium. For this study, we have focused on the
economics of FBS, a significant factor in cell culture medium content and costs.
Understanding the cost of VN lysate compared to FBS is necessary to predict the
sustainability of cultivated meat made using lysates as key media components. VN
lysate at 40 µg/mL used as a replacement of FBS is sustainable and cost effective at $2
per liter compared to $264 of FBS per liter, a 99.2 % price reduction, at lab scale (**Table
S2 & S3**). Thus, based on the calculations, VN40 reduces media cost from $323.30 to
$158.60, a 51% reduction (**Figure 6**). This accounts for only reagent costs involved in
the production of *V. natriegens* lysate in a lab. On an industrial scale, while the relative
advantage of VN lysate over FBS may remain consistent, the absolute cost of mass-
producing VN lysate should be lower than that calculated here.

Figure 5. Stemness and differentiation of VN40-adapted iBSCs. **A-B)** Immunofluorescent staining of iBSCs before (**A**) and after (**B**) adaptation and serial passaging (22 passages) in VN40 medium show no change in satellite cell phenotype after adaptation. Cells were stained with phalloidin (green) to indicate actin expression and anti-Pax7 (orange). Scale bar = 100 μm. **C)** Quantification of Pax 7+ cells before and after adaptation and serial passaging (22 passages) in VN40 medium. % Cells Pax7+ determined by (number of Pax7+ cells)/(number of actin+ cells). Statistical analysis by unpaired t-test with Welch's correction in which p>0.05 (NS). **D-G)** Immunofluorescent staining of gradually adapted iBSCs (p98) after 18 days differentiation in differentiation media (2 % horse serum, 50 μg/mL rAlbumin, 0.5 μM LDN-193189, ITS-X). **H-K)** Immunofluorescent staining of rapidly adapted iBSCs (p70) after 13 days differentiation in differentiation media. Results show successful differentiation of cells expressing early-stage (Desmin, green) and late-stage (myosin heavy chain, red) differentiation markers, with more robust differentiation of rapidly adapted cells. Cells were stained with anti-desmin, MF20 (for myosin heavy chain), and DAPI (blue) to stain nuclei. **D,F,H,J:** low magnification, Scale bar = 200 μm; **E,G,I,K:** high magnification, Scale bar = 100 μm.

Method	Metabolite	*M/Z*	RT	Annotation Score	Normalized Average AUC
HILIC (-)	Uracil	111.020	251.478	1.2	1.00
HILIC (-)	Hypoxanthine	135.031	531.488	6.4	0.26
HILIC (-)	Xanthine	151.026	791.072	6.3	0.17
HILIC (-)	L-Phenylalanine	164.072	535.862	1.6	0.14
HILIC (-)	Guanine	150.042	532.663	4.8	0.05
HILIC (-)	L-Arogenate	226.075	510.862	1.3	0.05
HILIC (-)	L-Glutamate	146.046	980.112	5	0.04
HILIC (-)	L-Citrulline	174.088	645.793	2.2	0.03
HILIC (-)	D-Xylulose 5-phosphate	229.012	1584.138	2.4	0.02
HILIC (-)	L-Methionine	148.044	547.405	1.7	0.02
HILIC (+)	5-Aminopentanoate	118.082	504.533	1	1.00
HILIC (+)	Uracil	113.030	249.740	1.2	0.04
HILIC (+)	Cytosine	112.048	297.273	1.2	0.003
HILIC (+)	L-Proline	116.066	991.431	2.1	0.002
HILIC (+)	Inosine	269.090	3112.661	3.3	0.002
HILIC (+)	L-Serine	106.046	659.801	3.6	0.001
HILIC (+)	Benzonitrile	104.050	546.203	1.1	0.001
HILIC (+)	Portulacaxanthin II	375.117	257.946	1.3	0.0003
RP (-)	Xanthine	151.026	709.452	6.3	1.00
RP (-)	L-Phenylalanine	164.072	1220.405	1.6	0.87
RP (-)	Inosine	267.074	1253.153	3.3	0.55
RP (-)	Uracil	111.020	299.736	1.2	0.44
RP (-)	Hypoxanthine	135.031	498.233	6.4	0.31
RP (-)	L-Tyrosine	180.066	736.295	1.5	0.26
RP (-)	L-Tryptophan	203.083	1607.048	3.2	0.26
RP (-)	Guanosine	282.084	1257.624	2.8	0.25
RP (-)	Xanthosine	283.069	1298.537	3.7	0.19
RP (-)	L-Citrulline	174.088	187.141	2.2	0.14
RP (+)	L-Phenylalanine	166.086	1218.830	1.6	1.00
RP (+)	L-Tyrosine	182.080	721.470	1.5	0.56
RP (+)	Guanine	152.056	310.407	4.8	0.55
RP (+)	Hypoxanthine	137.045	560.229	6.4	0.37
RP (+)	Uracil	113.034	321.318	1.2	0.30
RP (+)	Guanosine	284.097	1265.559	2.8	0.29
RP (+)	L-Methionine	150.057	286.392	1.7	0.28
RP (+)	Xanthine	153.040	776.350	6.3	0.17
RP (+)	Guanine	152.056	1288.269	4.8	0.15
RP (+)	Inosine	269.087	1248.965	3.3	0.11

515 **Table 2**. **Annotated metabolites showing greatest AUC detected in each LC-MS/MS method.** Each
516 metabolite's area is normalized to that with the highest AUC for each LC-MS method (n=3). Top 10
517 greatest AUC metabolites shown for each method, except HILIC (+) which found only 8 metabolites with
518 annotation score >1.0.

Figure 6. VN lysate is less expensive at lab scale than FBS, making VN40 media more economically sustainable than BSCGM. Media costs for BSC-GM, Beefy-9 and VN40 were assessed for reagents purchased at USD/L. VN media is the most economical followed by B9 and BSC-GM. FBS in BSCGM and rAlbumin in B9 are major cost-drivers for respective media compared to negligible cost of VN lysate in VN media.

4. DISCUSSION

Issues of increasing global food insecurity and climate crisis (Boyd et al., 2020; Hayek, 2022; Poore & Nemecek, 2018), coupled with the growing demand for meat (OECD et al., 2021), create an urgent need for more sustainable approaches to meat production. Cultivated meat, the process of growing meat *in vitro* without the need to raise livestock, is a possible alternative approach (Sinke et al., 2023; Treich, 2021). However, production of cultivated meat needs to be inexpensive, reliable, and relatively simple to compete with the low costs of traditional meat products from industrial farming. Moreover, the reliance on FBS for cell culture is a significant roadblock to scaling and commercializing cultivated meat due to FBS's animal origin, high cost, lot-to-lot variability, and potential as a source of contamination (Urzì et al., 2022; Zheng et al., 2006). Recombinant proteins (Kolkmann et al., 2022; Skrivergaard et al., 2023; Stout et al., 2023a; Venkatesan et al., 2022) or hydrolysates from organisms (Andreassen et al., 2020; Okamoto et al., 2022; Stout et al., 2023b; Teng, Lee, & Chen, 2023) are costly, labor intensive, and at times, still reliant on animal products or serum. Microbial extracts can be a protein-rich, inexpensive, and scalable alternatives to generating FBS-replacing media supplements. Microbial extracts have been utilized for cultivated meat production (Celebi-Birand et al., 2023; Ghosh et al., 2023; Lei et al., 2023), but still depended on serum (Lei et al., 2023), toxic bacteria (Ghosh et al., 2023), or otherwise animal-derived microbes (Celebi-Birand et al., 2023) to stimulate growth. Further, manufacturing these microbe-based serum-free media (SFM) formulations is limited by relatively slow growth rates of the chosen microbes.

In screening microbial lysates for cell line growth, we anticipated positive outcomes in either Gram positive bacteria (lacking lipopolysaccharides), or food-associated/probiotic microbes such as *S. cerevisiae* and *Lactococcus lactis.* Growth of Mack1 cells in 2.5% FBS medium supplemented with *S. cerevisiae* lysate 100 µg/mL was equivalent to that shown in 7.5% FBS-containing media by Lim et al. (2024). This means the yeast lysate can replace 5% FBS in medium without detrimental effects. The beneficial impact of *V. natriegens* lysate on iBSCs was a surprise, highlighting the importance of individualized screens for separate cell lines, potentially due to unique nutritional and signaling requirements (O'Neill et al., 2022). After screening cell growth using lysates from seven microbial strains, *V. natriegens* (a fast-growing marine bacterium) was chosen for use in long-term adaptation of iBSCs to SFM. By sequentially increasing *V. natriegens* lysate concentration in SFM, we were able to achieve maximal doubling rates of iBSCs using the VN40 formulation. This led to cell growth rates exceeding previously established Beefy 9 medium (Stout et al., 2022), while preserving iBSC phenotype and myogenicity. VN40 was able to stimulate long-term growth in iBSCs at up to 24.9 h doubling times, rivaling that of serum-containing media. The lysates were simple to produce and highly potent, stimulating cell growth at a mere 40 µg/mL. Further results indicated the

importance of both low- and high-molecular weight components in the microbial extract, suggesting there is no need for process-intensive ultrafiltration of lysate fractions to achieve optimal performance as in other studies (Ho et al., 2021; Stout et al., 2023b; Teng et al., 2023). LC-MS analysis of VN lysate revealed the presence of aromatic amino acids, purine derivatives and nucleoside intermediates which are also present in FBS (D. Y. Lee et al., 2022). These compounds aid in the proliferation of iBSCs through known mechanisms. Aromatic amino acids are synthesized in cow's rumen by fermentation (Khan et al., 2002), and can be found in FBS and basal media used for mammalian cell culture (Liu et al., 2023). The aromatic amino acids found in VN lysate could serve as nutrients for iBSCs and aid protein synthesis, cellular signaling, and/or oxidative stress response (Liu et al., 2023). The three aromatic amino acids found in VN lysate (tyrosine, phenylalanine, and tryptophan) are specifically known to be ubiquitously involved in protein synthesis, as well as synthesis of a host of metabolites vital for cell growth (Parthasarathy et al., 2018). They are also important for stimulating cell proliferation via the MAPK pathway (El Refaey et al., 2015), a pathway similarly stimulated by growth factors such as FGF, EGF (epidermal growth factor) and VEGF (vascular endothelial growth factor) (Rodrigues et al., 2010) found in FBS. Purines, pyrimidines, and their derivatives could contribute to the growth of iBSC in absence of any serum. These nucleotides found in the VN lysate, such as uracil, guanine, and cytosine, supplement precursors for DNA and RNA via the nucleotide salvage pathway in serum-free conditions lacking these metabolites (Morrison et al., 2019). Supplementation of the same nucleosides found in VN lysate (eg. hypoxanthine, inosine, and guanosine) have shown to improve growth rates in CHO cell cultures (Morrison et al., 2019). Inosine specifically can serve as a carbon source in media lacking the nutrients of FBS (Wang et al., 2020). While our analysis did uncover several compounds with pro-mitogenic properties in the absence of serum, any imbalance of these compounds can cause cytotoxicity. This explains why a high concentration of lysates may have inhibited cell growth. An imbalance of nucleotides can cause cell growth inhibition (Diehl et al., 2022). Overabundance of the aromatic amino acids found in VN lysate can lead to cell cycle arrest (Xavier et al., 2024). Numerous other compounds in VN lysate may have become cytotoxic to cells at higher concentrations. Further optimization of individual compounds in lysate-based media is worth investigation.

V. natriegens is a promising ingredient in cultivated meat media, as it is the fastest growing non-pathogenic bacteria known, with a generation time of < 10 min (H. H. Lee et al., 2019). Further, its nutritional needs are both minimal and flexible (Ellis et al., 2019). Our work shows its potential to supply the cultivated meat industry with an inexpensive on-demand serum-free supplement in a potent form. Previous environmental impact assessments using microbial lysates assumed a need for microbial protein on the scale of milligrams per milliliter of media (Tuomisto & de Mattos,

2011). Our media, only requiring 40 µg/mL protein, would reduce such needs by a factor
of one hundred. This would result in significant decreases in water usage, energy usage,
and greenhouse gas emissions associated with cultivated meat production, which
largely stem from production and processing of media ingredients (Sinke et al., 2023).
This high potency would also significantly reduce cost of cultivated meat production,
which are largely driven by media and additional growth factor costs (Specht, 2020).
606

AUTHOR CONTRIBUTIONS

James Dolgin, Damayanti Chakravarty – Conceptualization; formal analysis;
investigation; methodology; data visualization; writing & editing. Yiming Cai, Taehwan
Lim, Pomaikaimaikalani Yamaguchi – Formal analysis; investigation; methodology; data
visualization; writing. Sean F. Sullivan, Joseph E. Balkan, Licheng Xu, Aaron D.
Olawoyin – investigation; Kyongbum Lee – funding acquisition; supervision; writing—
review. David L. Kaplan and Nikhil U. Nair – Conceptualization; funding acquisition;
supervision; writing – reviewing & editing.

ACKNOWLEDGMENTS

617 This work was supported by funds from Tufts University Center for Cellular Agriculture
618 Industry Consortium (to N.U.N., D.L.K., K.L.) and the USDA (2021-05678) (to D.L.K.).

CONFLICT OF INTEREST

621 A patent application is pending based on this work.

REFERENCES

Alden, N., Krishnan, S., Porokhin, V., Raju, R., McElearney, K., Gilbert, A., & Lee, K. (2017). Biologically Consistent Annotation of Metabolomics Data. *Analytical Chemistry, 89*(24), 13097-13104. doi:10.1021/acs.analchem.7b02162

Allen, F., Pon, A., Wilson, M., Greiner, R., & Wishart, D. (2014). CFM-ID: a web server for annotation, spectrum prediction and metabolite identification from tandem mass spectra. *Nucleic Acids Research, 42*(W1), W94-W99. doi:10.1093/nar/gku436

Andreassen, R. C., Pedersen, M. E., Kristoffersen, K. A., & Ronning, S. B. (2020). Screening of by-products from the food industry as growth promoting agents in serum-free media for skeletal muscle cell culture. *Food & Function, 11*(3), 2477-2488. doi:10.1039/c9fo02690h

Batish, I., Zarei, M., Nitin, N., & Ovissipour, R. (2022). Evaluating the Potential of Marine Invertebrate and Insect Protein Hydrolysates to Reduce Fetal Bovine Serum in Cell Culture Media for Cultivated Fish Production. *Biomolecules, 12*(11). doi:10.3390/biom12111697

Boyd, R., Thorpe, R., Hyder, K., Roy, S., Walker, N., & Sibly, R. (2020). Potential Consequences of Climate and Management Scenarios for the Northeast Atlantic Mackerel Fishery. *Frontiers in Marine Science, 7*. doi:10.3389/fmars.2020.00639

Capetanaki, Y., Milner, D. J., & Weitzer, G. (1997). Desmin in muscle formation and maintenance: Knockouts and consequences. *Cell Structure and Function, 22*(1), 103-116. doi:10.1247/csf.22.103

Celebi-Birand, D., Genc, K., Agun, I., Erikci, E., Akcali, K. C., & Kiran, F. (2023). Microbiota-Derived Postbiotics Enhance the Proliferative Effects of Growth Factors on Satellite Cells in Cultivated Meat Applications. *Sustainability, 15*(23). doi:ARTN 16164 10.3390/su152316164

Chevalot, I., Dziukala, C., Olmos, E., Marc, I., Druaux-Borreill, D., Oriol, E., & Marc, A. (2010). Use of Yeast Derived Nutrients for Cell Culture in Serum-Free Media. *Cells and Culture, 4*, 843-846. doi:10.1007/978-90-481-3419-9_148

Chia, M. A., Jankowiak, J. G., Kramer, B. J., Goleski, J. A., Huang, I. S., Zimba, P. V., Bittencourt-Oliveira, M., Gobler, C. J. (2018). Succession and toxicity of and blooms are controlled by nutrient-dependent allelopathic interactions. *Harmful Algae, 74*, 67-77. doi:10.1016/j.hal.2018.03.002

Dai, W., Chen, Y., Xiong, W., Li, S., Tan, W. S., & Zhou, Y. (2024). Development of a serum-free medium for myoblasts long-term expansion and 3D culture for cell-based meat. *J Food Sci, 89*(2), 851-865. doi:10.1111/1750-3841.16884

de Bruin, O. M., Chiefari, A., Wroblewski, D., Egan, C., & Kelly-Cirino, C. D. (2019). A novel chemical lysis method for maximum release of DNA from difficult-to-lyse bacteria. *Microb Pathog, 126*, 292-297. doi:10.1016/j.micpath.2018.11.008

Diehl, F. F., Miettinen, T. P., Elbashir, R., Nabel, C. S., Darnell, A. M., Do, B. T., Manalis, S. R., Lewis, C. A., & Vander Heiden, M. G. (2022). Nucleotide imbalance decouples cell growth from cell proliferation. *Nature Cell Biology, 24*(8), 1252–1264. https://doi.org/10.1038/s41556-022-00965-1

Dong, N. N., Jiang, B. X., Chang, Y. G., Wang, Y. C., & Xue, C. H. (2024). Integrated Omics Approach: Revealing the Mechanism of Protein Extract Replacing Fetal

Bovine Serum for Fish Muscle Cell Culture. *Journal of Agricultural and Food*
Chemistry, 72(11), 6064-6076. doi:10.1021/acs.jafc.4c00624

Dong, N. N., Xue, C. H., Yang, Y. X., Chang, Y. G., Wang, Y. M., Guo, H. R., Liu, Y.,
Wang, Y. C. (2023). extract supplementation replacing fetal bovine serum for
muscle cell culture under low-serum conditions. *Food Research International,*
164. doi:ARTN 112438 10.1016/j.foodres.2022.112438

El Refaey, M., Watkins, C. P., Kennedy, E. J., Chang, A., Zhong, Q., Ding, K.-H., Shi, X.,
Xu, J., Bollag, W. B., Hill, W. D., Johnson, M., Hunter, M., Hamrick, M. W., &
Isales, C. M. (2015). Oxidation of the aromatic amino acids tryptophan and
tyrosine disrupts their anabolic effects on bone marrow mesenchymal stem cells.
Molecular and Cellular Endocrinology, 410, 87–96.
https://doi.org/10.1016/j.mce.2015.01.034

Ellis, G. A., Tschirhart, T., Spangler, J., Walper, S. A., Medintz, I. L., & Vora, G. J. (2019).
Exploiting the Feedstock Flexibility of the Emergent Synthetic Biology Chassis
for Engineered Natural Product Production. *Marine Drugs, 17*(12). doi:ARTN 679
10.3390/md17120679

Ghosh, J., Haraguchi, Y., Asahi, T., Nakao, Y., & Shimizu, T. (2023). Muscle cell
proliferation using water-soluble extract from nitrogen-fixing cyanobacteria
Anabaena sp. PCC 7120 for sustainable cultured meat production. *Biochemical*
and Biophysical Research Communications, 682, 316-324.
doi:10.1016/j.bbrc.2023.10.018

Grimaldo, E., Karl, C. W., Alvestad, A., Persson, A. M., Kubowicz, S., Olafsen, K.,
Hatlebrekke, H., Lilleng, G., Brinkhof, I. (2023). Reducing plastic pollution caused
by demersal fisheries. *Marine Pollution Bulletin, 196*. doi:ARTN 115634
10.1016/j.marpolbul.2023.115634

Hartman, T. E., Sar, N., Genereux, K., Barritt, D. S., He, Y. M., Burky, J. E., Wesson, M.
C., Tso, J.Y., Tsurushita, N., Zhou, W., Sauer, P. W. (2007). Derivation and
characterization of cholesterol-independent non-GS NS0 cell lines for production
of recombinant antibodies. *Biotechnology and Bioengineering, 96*(2), 294-306.
doi:10.1002/bit.21099

Hayek, M. N. (2022). The infectious disease trap of animal agriculture. *Sci Adv, 8*(44),
eadd6681. doi:10.1126/sciadv.add6681

Ho, Y. Y., Lu, H. K., Lim, Z. F. S., Lim, H. W., Ho, Y. S., & Ng, S. K. (2021). Applications
and analysis of hydrolysates in animal cell culture. *Bioresources and*
Bioprocessing, 8(1). doi:ARTN 93 10.1186/s40643-021-00443-w

Horbelt, D., Boergermann, J. H., Chaikuad, A., Alfano, I., Williams, E., Lukonin, I.,
Timmel, T., Bullock, A. N., Knaus, P. (2015). Small Molecules Dorsomorphin and
LDN-193189 Inhibit Myostatin/GDF8 Signaling and Promote Functional Myoblast
Differentiation. *Journal of Biological Chemistry, 290*(6), 3390-3404.
doi:10.1074/jbc.M114.604397

Hua, J. L., Wei, Y. X., Zhang, Y., Xu, H. L., Ge, J. L., Liu, M. Z., Wang, Y., Shi, Y., Hou, L.,
Jiang, H. (2022). Adaptation process of engineered cell line FCHO/IL-24 stably
secreted rhIL-24 in serum-free suspension culture. *Protein Expression and*
Purification, 199. doi:ARTN 106154 10.1016/j.pep.2022.106154

Iberite, F., Gruppioni, E., & Ricotti, L. (2022). Skeletal muscle differentiation of human iPSCs meets bioengineering strategies: perspectives and challenges. *Npj Regenerative Medicine, 7*(1). doi:ARTN 23 10.1038/s41536-022-00216-9

Jeong, Y., Choi, W. Y., Park, A., Lee, Y. J., Lee, Y., Park, G. H., Lee, S. J., Lee, W. K., Ryu, Y. K., Kang, D. H. (2021). Marine cyanobacterium as an alternate to the animal cell culture medium supplement. *Scientific Reports, 11*(1). doi:ARTN 4906 10.1038/s41598-021-84558-2

Khan, R. I., Onodera, R., Amin, M. R., & Mohammed, N. (2002). Aromatic amino acid biosynthesis and production of related compounds from hydroxyphenylpyruvic acid by rumen bacteria, protozoa and their mixture. *Amino Acids, 22*(2), 167-177. doi:DOI 10.1007/s007260200005

Kolkmann, A. M., Van Essen, A., Post, M. J., & Moutsatsou, P. (2022). Development of a Chemically Defined Medium for Expansion of Primary Bovine Satellite Cells. *Frontiers in Bioengineering and Biotechnology, 10.* doi:ARTN 895289 10.3389/fbioe.2022.895289

Kuo, H. H., Gao, X. Z., DeKeyser, J. M., Fetterman, K. A., Pinheiro, E. A., Weddle, C. J., Fonoudi, H., Orman, M. V., Romero-Tejeda, M., Jouni, M., Blancard, M., Magdy, T., Epting, C. L., George Jr., A. L., Burridge, P. W. (2020). Negligible-Cost and Weekend-Free Chemically Defined Human iPSC Culture. *Stem Cell Reports, 14*(2), 256-270. doi:10.1016/j.stemcr.2019.12.007

Lee, D. Y., Lee, S. Y., Yun, S. H., Jeong, J. W., Kim, J. H., Kim, H. W., Choi, J. S., Kim, G., Joo, S. T., Choi, I., Hur, S. J. (2022). Review of the Current Research on Fetal Bovine Serum and the Development of Cultured Meat. *Food Science of Animal Resources, 42*(5), 775-799. doi:10.5851/kosfa.2022.e46

Lee, H. H., Ostrov, N., Wong, B. G., Gold, M. A., Khalil, A. S., & Church, G. M. (2019). Functional genomics of the rapidly replicating bacterium by CRISPRi. *Nature Microbiology, 4*(7), 1105-1113. doi:10.1038/s41564-019-0423-8

Lee, J. Y., Kang, M. H., Jang, J. E., Lee, J. E., Yang, Y., Choi, J. Y., Kang, H. S., Lee, U., Choung, J. W., Jung, H., Yoon, Y., Jung, K. H., Hong, S., Yi, E. C., Park, S. G. (2022). Comparative analysis of mesenchymal stem cells cultivated in serum free media. *Scientific Reports, 12*(1). doi:ARTN 8620 10.1038/s41598-022-12467-z

Lee, M., Park, S., Choi, B., Choi, W., Lee, H., Lee, J. M., Lee, S. T., Yoo, K. H., Han, D., Bang, G., Hwang, H., Koh, W., Lee, S., Hong, J. (2024). Cultured meat with enriched organoleptic properties by regulating cell differentiation. *Nature Communications, 15*(1). doi:ARTN 77 10.1038/s41467-023-44359-9

Lei, Q. Z., Ma, J., Du, G. C., Zhou, J. W., & Guan, X. (2023). Efficient expression of a cytokine combination in Saccharomyces cerevisiae for cultured meat production. *Food Research International, 170.* doi:10.1016/j.foodres.2023.113017

Lim, T., Chang, H., Saad, M. K., Joyce, C. M., Park, B., O'Beirne, S. X., Cohen, M. A., & Kaplan, D. L. (2024). Development of Serum-Reduced Medium for Mackerel Muscle Cell Line Cultivation. *ACS Sustainable Chemistry & Engineering, 12*(31), 11683–11691. https://doi.org/10.1021/acssuschemeng.4c03345

Liu, S., Yang, W., Li, Y., & Sun, C. (2023). Fetal bovine serum, an important factor affecting the reproducibility of cell experiments. *Sci Rep, 13*(1), 1942. doi:10.1038/s41598-023-29060-7

Mahieu, N. G., & Patti, G. J. (2017). Systems-Level Annotation of a Metabolomics Data Set Reduces 25 000 Features to Fewer than 1000 Unique Metabolites. *Analytical Chemistry*, *89*(19), 10397–10406. https://doi.org/10.1021/acs.analchem.7b02380

Messmer, T., Klevernic, I., Furquim, C., Ovchinnikova, E., Dogan, A., Cruz, H., Post, M. J., Flack, J. E. (2022). A serum-free media formulation for cultured meat production supports bovine satellite cell differentiation in the absence of serum starvation. *Nature Food, 3*(1), 74-+. doi:10.1038/s43016-021-00419-1

Miki, H., & Takagi, M. (2015). Design of serum-free medium for suspension culture of CHO cells on the basis of general commercial media. *Cytotechnology, 67*(4), 689-697. doi:10.1007/s10616-014-9778-0

Morrison, C., Bandara, K., Wang, W., Zhang, L., & Figueroa, B., Jr. (2019). Improvement of growth rates through nucleoside media supplementation of CHO clones. *Cytotechnology, 71*(3), 733-742. doi:10.1007/s10616-019-00319-0

O'Neill, E. N., Ansel, J. C., Kwong, G. A., Plastino, M. E., Nelson, J., Baar, K., & Block, D. E. (2022). Spent media analysis suggests cultivated meat media will require species and cell type optimization. *Npj Science of Food, 6*(1). doi:ARTN 46 10.1038/s41538-022-00157-z

OECD, Food, & Nations, A. O. o. t. U. (2021). *OECD-FAO Agricultural Outlook 2021-2030*.

Okamoto, Y., Haraguchi, Y., Yoshida, A., Takahashi, H., Yamanaka, K., Sawamura, N., .Asahi, T., Shimizu, T. (2022). Proliferation and differentiation of primary bovine myoblasts using extract for sustainable production of cultured meat. *Biotechnology Progress, 38*(3). doi:ARTN e3239 10.1002/btpr.3239

Park, S., Park, G., Oh, S., Park, Y., Kim, Y., Kim, J., & Choi, J. (2023). Investigating proliferation and differentiation capacities of Hanwoo steer myosatellite cells at different passages for developing cell-cultured meat. *Scientific Reports, 13*(1). doi:ARTN 15614 10.1038/s41598-023-40800-7

Parthasarathy, A., Cross, P. J., Dobson, R. C. J., Adams, L. E., Savka, M. A., & Hudson, A. O. (2018). A Three-Ring Circus: Metabolism of the Three Proteogenic Aromatic Amino Acids and Their Role in the Health of Plants and Animals. *Frontiers in Molecular Biosciences, 5*. https://doi.org/10.3389/fmolb.2018.00029

Poore, J., & Nemecek, T. (2018). Reducing food's environmental impacts through producers and consumers. *Science, 360*(6392), 987-992. doi:10.1126/science.aaq0216

Radosevic, K., Dukic, B., Andlar, M., Slivac, I., & Gaurina Srcek, V. (2016). Adaptation and cultivation of permanent fish cell line CCO in serum-free medium and influence of protein hydrolysates on growth performance. *Cytotechnology, 68*(1), 115-121. doi:10.1007/s10616-014-9760-x

Relaix, F., Rocancourt, D., Mansouri, A., & Buckingham, M. (2005). A Pax3/Pax7-dependent population of skeletal muscle progenitor cells. *Nature, 435*(7044), 948-953. doi:10.1038/nature03594

Robert Vergeer, P. S., Ingrid Odegard. (2021). TEA of cultivated meat: Future projections of different scenarios -corrigendum. Retrieved from https://cedelft.eu/wp-content/uploads/sites/2/2021/02/CE_Delft_190254_TEA_of_Cultivated_Meat_FINAL_corrigendum.pdf

Rodrigues, M., Griffith, L. G., & Wells, A. (2010). Growth factor regulation of proliferation and survival of multipotential stromal cells. *Stem Cell Research & Therapy, 1*(4), 32. https://doi.org/10.1186/scrt32

Saad, M. K., Yuen, J. S. K. Y., Joyce, C. M., Li, X. X., Lim, T., Wolfson, T. L., Wu, J., Laird, J., Vissapragada, S., Calkins, O. P., Ali, A., Kaplan, D. L. (2023). Continuous fish muscle cell line with capacity for myogenic and adipogenic-like phenotypes. *Scientific Reports, 13*(1). doi:ARTN 5098 10.1038/s41598-023-31822-2

Shen, X., Wang, R., Xiong, X., Yin, Y., Cai, Y., Ma, Z., Liu, N., & Zhu, Z.-J. (2019). Metabolic reaction network-based recursive metabolite annotation for untargeted metabolomics. *Nature Communications, 10*(1), 1516. https://doi.org/10.1038/s41467-019-09550-x

Sinke, P., Swartz, E., Sanctorum, H., van der Giesen, C., & Odegard, I. (2023). Ex-ante life cycle assessment of commercial-scale cultivated meat production in 2030 (vol 28, pg 234, 2023). *International Journal of Life Cycle Assessment, 28*(9), 1225-1228. doi:10.1007/s11367-023-02183-9

Skrivergaard, S., Young, J. F., Sahebekhtiari, N., Semper, C., Venkatesan, M., Savchenko, A., Stogios, P. J., Thirkildsen, M., Rasmussen, M. K. (2023). A simple and robust serum-free media for the proliferation of muscle cells. *Food Research International, 172*. doi:ARTN 113194 10.1016/j.foodres.2023.113194

Specht, L. (2020). An analysis of culture medium cost and production volumes for cultivated meat. Retrieved from https://gfi.org/wp-content/uploads/2021/01/clean-meat-production-volume-and-medium-cost.pdf

Stout, A. J., Arnett, M. J., Chai, K. S., Guo, T. A., Liao, L. S., Mirliani, A. B., White, E. C., Yuen Jr., J. S. K., Zhang, X., Kaplan, D. L. (2023). Immortalized Bovine Satellite Cells for Cultured Meat Applications. *Acs Synthetic Biology, 12*(5), 1567-1573. doi:10.1021/acssynbio.3c00216

Stout, A. J., Mirliani, A. B., Rittenberg, M. L., Shub, M., White, E. C., Yuen, J. S. K., & Kaplan, D. L. (2022). Simple and effective serum-free medium for sustained expansion of bovine satellite cells for cell cultured meat. *Communications Biology, 5*(1). doi:ARTN 466 10.1038/s42003-022-03423-8

Stout, A. J., Rittenberg, M. L., Shub, M., Saad, M. K., Mirliani, A. B., Dolgin, J., & Kaplan, D. L. (2023). A Beefy-R culture medium: Replacing albumin with rapeseed protein isolates. *Biomaterials, 296*. doi:ARTN 122092 10.1016/j.biomaterials.2023.122092

Stout, A. J., Zhang, X., Letcher, S. M., Rittenberg, M. L., Shub, M., Chai, K. M., Kaul, M., Kaplan, D. L. (2024). Engineered autocrine signaling eliminates muscle cell FGF2 requirements for cultured meat production. *Cell Reports Sustainability, 1(1)*. https://doi.org/10.1016/j.crsus.2023.100009

Sung, Y. H., Lim, S. W., Chung, J. Y., & Lee, G. M. (2004). Yeast hydrolysate as a low-cost additive to serum-free medium for the production of human thrombopoietin in suspension cultures of Chinese hamster ovary cells. *Applied Microbiology and Biotechnology, 63*(5), 527-536. doi:10.1007/s00253-003-1389-1

Teng, T. S., Lee, J. J. L., & Chen, W. N. (2023). Ultrafiltrated Extracts of Fermented Okara as a Possible Serum Alternative for Cell Culturing: Potential in Cultivated

Meat Production. *Acs Food Science & Technology, 3*(4), 699-709.
doi:10.1021/acsfoodscitech.2c00401
Treich, N. (2021). Cultured Meat: Promises and Challenges. *Environmental & Resource*
Economics, 79(1), 33-61. doi:10.1007/s10640-021-00551-3
Tuomisto, H. L., & de Mattos, M. J. T. (2011). Environmental Impacts of Cultured Meat
Production. *Environmental Science & Technology, 45*(14), 6117-6123.
doi:10.1021/es200130u
Urzì, O., Bagge, R. O., & Crescitelli, R. (2022). The dark side of foetal bovine serum in
extracellular vesicle studies. *Journal of Extracellular Vesicles, 11*(10). doi:ARTN
12271 10.1002/jev2.12271
Venkatesan, M., Semper, C., Skrivergaard, S., Di Leo, R., Mesa, N., Rasmussen, M. K.,
Young, J. F., Therkildsen, M., Stogios, P. J., Savchenko, A. (2022). Recombinant
production of growth factors for application in cell culture. *Iscience, 25*(10).
doi:ARTN 105054 10.1016/j.isci.2022.105054
Wang, T., Gnanaprakasam, J. N. R., Chen, X., Kang, S., Xu, X., Sun, H., Liu, L.,
Rodgers, H., Miller, E., Cassel, T. A., Sun, Q., Vicente-Muñoz, S., Warmoes, M.
O., Lin, P., Piedra-Quintero, Z. L., Guerau-de-Arellano, M., Cassady, K. A., Zheng,
S. G., Yang, J., … Wang, R. (2020). Inosine is an alternative carbon source for
CD8+-T-cell function under glucose restriction. *Nature Metabolism*, *2*(7), 635–647.
https://doi.org/10.1038/s42255-020-0219-4
Wishart, D. S., Jewison, T., Guo, A. C., Wilson, M., Knox, C., Liu, Y. F., Djoumbou, Y.,
Mandal, R., Aziat, F., Dong, E., Bouatra, S., Sinelnikov, I., Arndt, D., Xia, J., Liu,
P., Yallou, F., Bjorndahl, T., Perez-Pineiro, R., Eisner, R., Allen, F., Neveu, V.,
Greiner, R., Scalbert, A. (2013). HMDB 3.0-The Human Metabolome Database in
2013. *Nucleic Acids Research, 41*(D1), D801-D807. doi:10.1093/nar/gks1065
Xavier, A., & Dikic, I. (2024). Feeding cancer to death—A triad of aromatic acids
reduces tumor growth. *Cell Death & Differentiation*, *31*(10), 1239–1241.
https://doi.org/10.1038/s41418-024-01372-9
Yamanaka, K., Haraguchi, Y., Takahashi, H., Kawashima, I., & Shimizu, T. (2023).
Development of serum-free and grain-derived-nutrient-free medium using
microalga-derived nutrients and mammalian cell-secreted growth factors for
sustainable cultured meat production. *Sci Rep, 13*(1), 498. doi:10.1038/s41598-
023-27629-w
Zheng, X. Y., Baker, H., Hancock, W. S., Fawaz, F., McCaman, M., & Pungor, E. (2006).
Proteomic analysis for the assessment of different lots of fetal bovine serum as a
raw material for cell culture. Part IV. Application of proteomics to the manufacture
of biological drugs. *Biotechnology Progress, 22*(5), 1294-1300.
doi:10.1021/bp060121o
885

Supplemental Information

Microbial lysates as low-cost serum replacements in cellular
agriculture media formulation

James Dolgin [1,4]*, Damayanti Chakravarty[2,4]*, Sean F. Sullivan[2,4], Yiming Cai[3,4],
Taehwan Lim[1,4], Pomaikaimaikalani Yamaguchi[2,4], Joseph E. Balkan[2], Licheng Xu[1,4],
Aaron D. Olawoyin[1,4], Kyongbum Lee[2,4], David L. Kaplan[1,4], Nikhil U. Nair[2,4]†

* equal contribution
† correspondence: nikhil.nair@tufts.edu
[1] Department of Biomedical Engineering, Tufts University, Medford, MA
[2] Department of Chemical & Biological Engineering, Tufts University, Medford, MA
[3] Department of Biology, Tufts University, Medford, MA
[4] Tufts University Center for Cellular Agriculture (TUCCA), Tufts University, Medford, MA

900
901

Ref.	Active ingredient for serum replacement	Low-cost	Minimally labor-intensive production	Rapidly replicating source organism	Non-toxic	Free of FBS or other animal ingredients	Long-term cell growth & myogenic capacity
(Stout et al., 2022)	Recombinant albumin		X	X	X	X	X
(Kolkmann et al., 2022)	Recombinant human serum albumin		X	X	X	X	X
(Skrivergaard et al., 2023)	Fetuin + other growth factors		X	X	X	X	X
(Seo et al., 2024)	Recombinant heme				X	X	
(Dai et al., 2024)	Recombinant growth factors, bovine albumin, ethanolamine		X				X
(Stout, Rittenberg, et al., 2023)	Rapeseed protein isolates	X			X	X	X
(Dong et al., 2024)	*A. pyrenoidosa* Extract	X			X		
(Okamoto et al., 2022)	*C. vulgaris* extract	X			X		
(Yamanaka, Haraguchi, Takahashi, Kawashima, & Shimizu, 2023)	*C. vulgaris* extract	X			X	X	
(Teng et al., 2023)	Ultrafiltered Okara extract	X			X	X	
(Radosevic, Dukic, Andlar, Slivac, & Gaurina Srcek, 2016)	Soy and gluten hydrolysates	X			X	X	
(Batish, Zarei, Nitin, & Ovissipour, 2022)	Invertebrate hydrolysates	X			X		
(Lei et al., 2023)	Growth factor-expressing *S. cerevisiae*		X	X	X		
(Celebi-Birand et al., 2023)	*E. hirae* – secreted postbiotics	X	X	X	X		
(Ghosh et al., 2023)	*Anabaena* extract	X				X	

(Jeong et al., 2021)	S. maxima extract	X	X		X		
This work	V. natriegens lysate	X	X	X	X	X	X

903 **Table S1.** Previous research exploring cultivated meat media, with advantages of
904 current work indicated.
905

906
907 **Figure S1.** iBSCs fed serum-free media with *V. natriegens* lysate at 40 (VN40), 60, and 80 μg/mL, and
908 resultant doubling times.
909

Figure S2. Venn diagram indicating the top 10 highest AUC (8 for HILIC+) metabolites detected by each method (HILIC (-), HILIC (+), RP (+) and RP (-) and the common metabolites among the method groups.

BSCGM								
Compone nt	**Unit/ L**	**Amount/ L (rxn)**	**Supplier**	**Cat#**	**Units/orde r**	**Cost (USD)/orde r**	**Cost/ Unit**	**Cost/ L rxn**
DMEM+ HEPES+ Sodium bicarbonate	ml	800	ThermoFish er	10569044	5000	365	0.073	58.4
FBS	ml	200	ThermoFish er	26140079	500	660	1.32	264
FGF-2	ug	1	Peprotech	100-18B	1000	902	0.902	0.902
Total								**323.302**

B9								
Component	**Unit/ L**	**Amount/ L (rxn)**	**Supplier**	**Cat#**	**Units/orde r**	**Cost (USD)**	**Cost/ Unit**	**Cost/ L rxn**
DMEM+ HEPES+ Sodium bicarbonate	ml	1000	ThermoFish er	11330057	5000	401	0.0802	80.2
L-ascorbic acid 2-phosphate	mg	200	Sigma	49752-10G	10000	85.8	0.00858	1.716
Insulin	mg	20	Sigma	91077C-250MG	250	349	1.396	27.92
Transferrin	mg	20	inVitria	777TRF029	1000	332.75	0.33275	6.655
Sodium selenite	ug	20	Sigma	S5261-10G	10000000	40.3	0.00000403	0.0000806
FGF2	ug	40	Peprotech	100-18B	1000	902	0.902	36.08
NRG1	ng	100	Peprotech	100-03	10000	89.5	0.00895	0.895
TGFb3	ng	100	R&D Systems	s 8420-B3-005/CF	5000	249	0.0498	4.98
rAlbumin	mg	800	Sigma	A9731-1G	1000	133	0.133	106.4
Total								**264.846081**

VN40 media								
Component	**Unit/ L**	**Amount/ L (rxn)**	**Supplier**	**Cat#**	**Units/orde r**	**Cost (USD)**	**Cost/ Unit**	**Cost/L (rxn)**
DMEM+ HEPES+ Sodium bicarbonate	ml	1000	ThermoFish er	11330057	5000	401	0.0802	80.2

Component	Unit	Amount/L (rxn)	Supplier	Cat #	Unit per order	Cost (USD)	Cost/Unit	Cost/L (rxn)
L-ascorbic acid 2-phosphate	mg	200	Sigma	49752-10G	10000	85.8	0.00858	1.716
Insulin	mg	20	Sigma	91077C-250MG	250	349	1.396	27.92
Transferrin	mg	20	inVitria	777TRF029	1000	332.75	0.33275	6.655
Sodium selenite	ug	20	Sigma	S5261-10G	10000000	40.3	0.00000403	0.0000806
FGF2	ug	40	Peprotech	100-18B	1000	902	0.902	36.08
NRG1	ng	100	Peprotech	100-03	10000	89.5	0.00895	0.895
TGFb3	ng	100	R&D Systems	s 8420-B3-005/CF	5000	249	0.0498	4.98
VN Lysate	mg	40						2
Total								158.446081

Table S2. Cost comparison of itemized media components and total cost of iBSCGM and B9 with VN40 medium. The cost calculation for VN lysate was based on LB media (BD Difco, 244620). One gram of LB costs 0.346 USD. We prepare 25 mL of VN culture in LB, yielding approximately 4 mg/mL of lysate (total protein content). For 25 mL of culture, we use 0.625 grams of LB. Therefore, 0.625 grams of LB yields 4 mg/mL of lysate, and 40 µg/mL lysate is produced from 0.006 grams of LB. The cost of 0.006 grams of LB is $0.002, making the cost for 40 µg/mL VN lysate $0.002. This translates to $0.00005 per 1 µg/mL of VN lysate, and thus, 40,000 µg/mL would cost ~$2.

Component	Unit	Amount/L (rxn)	Supplier	Cat #	Unit per order	Cost (USD)	Cost/Unit	Cost/L (rxn)
FBS	mL	200 mL	ThermoFisher Scientific	26140079	500	660	1.32	264
V. natriegens lysate	µg	40 mg/L	NA	NA	NA	NA	0.0005	2

Table S3: VN lysate is 99.2% cheaper than FBS.

931 **Additional details on LC-MS/MS methods.**
932
Reverse phase (RP) chromatography method
(Phenomenex Synergi Hydro-RP Column)
• Solvents:
A: 0.1% formic acid in water
B: 0.1% formic acid in methanol
• Column temperature: 15°C
• Flow rate: 0.2 mL/min
• Ion source: Turbo spray (ESI)
• Ion source Gas 1: 35
• Ion source Gas 2: 45
• Curtain Gas: 25
• Temperature: 450C
• IonSpray Voltage Floating: ±4500 V
946
947 Gradient:

Time (min)	%B
0-8	3
8-38	3 → 95
38-45	95
45-47	95 → 3
47-55	3

948
Hydrophilic interaction chromatography (HILIC) method
(Phenomenex Luna NH2 Column)
• Solvents
A: 95:5 water:acetonitrile
+ 20mM ammonium acetate, pH to 9.45 using ammonium hydroxide
B: 100% acetonitrile
• Column temperature: 25°C
• Ion source: Turbo spray (ESI)
• Ion source Gas 1: 35
• Ion source Gas 2: 45
• Curtain Gas: 25
• Temperature: 450C
• IonSpray Voltage Floating: ±5500 V
962
963 Gradient:

Time (min)	%B
0-15	85 → 0
15-28	0
28-30	0 → 85
30-60	85

964
965

References

Batish, I., Zarei, M., Nitin, N., & Ovissipour, R. (2022). Evaluating the Potential of Marine Invertebrate and Insect Protein Hydrolysates to Reduce Fetal Bovine Serum in Cell Culture Media for Cultivated Fish Production. *Biomolecules, 12*(11). doi:10.3390/biom12111697

Celebi-Birand, D., Genc, K., Agun, I., Erikci, E., Akcali, K. C., & Kiran, F. (2023). Microbiota-Derived Postbiotics Enhance the Proliferative Effects of Growth Factors on Satellite Cells in Cultivated Meat Applications. *Sustainability, 15*(23). doi:ARTN 16164 10.3390/su152316164

Dai, W., Chen, Y., Xiong, W., Li, S., Tan, W. S., & Zhou, Y. (2024). Development of a serum-free medium for myoblasts long-term expansion and 3D culture for cell-based meat. *J Food Sci, 89*(2), 851-865. doi:10.1111/1750-3841.16884

Dong, N. N., Xue, C. H., Yang, Y. X., Chang, Y. G., Wang, Y. M., Guo, H. R., Liu, Y., Wang, Y. C. (2023). extract supplementation replacing fetal bovine serum for muscle cell culture under low-serum conditions. *Food Research International, 164*. doi:ARTN 112438 10.1016/j.foodres.2022.112438

Ghosh, J., Haraguchi, Y., Asahi, T., Nakao, Y., & Shimizu, T. (2023). Muscle cell proliferation using water-soluble extract from nitrogen-fixing cyanobacteria Anabaena sp. PCC 7120 for sustainable cultured meat production. *Biochemical and Biophysical Research Communications, 682*, 316-324. doi:10.1016/j.bbrc.2023.10.018

Jeong, Y., Choi, W. Y., Park, A., Lee, Y. J., Lee, Y., Park, G. H., Lee, S. J., Lee, W. K., Ryu, Y. K., Kang, D. H. (2021). Marine cyanobacterium as an alternate to the animal cell culture medium supplement. *Scientific Reports, 11*(1). doi:ARTN 4906 10.1038/s41598-021-84558-2

Kolkmann, A. M., Van Essen, A., Post, M. J., & Moutsatsou, P. (2022). Development of a Chemically Defined Medium for Expansion of Primary Bovine Satellite Cells. *Frontiers in Bioengineering and Biotechnology, 10*. doi:ARTN 895289 10.3389/fbioe.2022.895289

Lei, Q. Z., Ma, J., Du, G. C., Zhou, J. W., & Guan, X. (2023). Efficient expression of a cytokine combination in Saccharomyces cerevisiae for cultured meat production. *Food Research International, 170*. doi: 10.1016/j.foodres.2023.113017

Okamoto, Y., Haraguchi, Y., Yoshida, A., Takahashi, H., Yamanaka, K., Sawamura, N., .Asahi, T., Shimizu, T. (2022). Proliferation and differentiation of primary bovine myoblasts using extract for sustainable production of cultured meat. *Biotechnology Progress, 38*(3). doi:ARTN e3239 10.1002/btpr.3239

Radosevic, K., Dukic, B., Andlar, M., Slivac, I., & Gaurina Srcek, V. (2016). Adaptation and cultivation of permanent fish cell line CCO in serum-free medium and influence of protein hydrolysates on growth performance. *Cytotechnology, 68*(1), 115-121. doi:10.1007/s10616-014-9760-x

Seo, Y. A., Cha, M. J., Park, S., Lee, S., Lim, Y. J., Son, D. W., Lee, E. J., Kim, P., & Chang, S. (2024). Development of a Normal Porcine Cell Line Growing in a Heme-Supplemented, Serum-Free Condition for Cultured Meat. *International Journal of Molecular Sciences*, *25*(11), 5824. https://doi.org/10.3390/ijms25115824

Skrivergaard, S., Young, J. F., Sahebekhtiari, N., Semper, C., Venkatesan, M.,
Savchenko, A., Stogios, P. J., Thirkildsen, M., Rasmussen, M. K. (2023). A simple
and robust serum-free media for the proliferation of muscle cells. *Food Research*
International, 172. doi:ARTN 113194 10.1016/j.foodres.2023.113194
Stout, A. J., Rittenberg, M. L., Shub, M., Saad, M. K., Mirliani, A. B., Dolgin, J., & Kaplan,
D. L. (2023). A Beefy-R culture medium: Replacing albumin with rapeseed
protein isolates. *Biomaterials, 296*. doi:ARTN 122092
10.1016/j.biomaterials.2023.122092
Stout, A. J., Mirliani, A. B., Rittenberg, M. L., Shub, M., White, E. C., Yuen, J. S. K., &
Kaplan, D. L. (2022). Simple and effective serum-free medium for sustained
expansion of bovine satellite cells for cell cultured meat. *Communications Biology,*
5(1). doi:ARTN 466 10.1038/s42003-022-03423-8
Teng, T. S., Lee, J. J. L., & Chen, W. N. (2023). Ultrafiltrated Extracts of Fermented
Okara as a Possible Serum Alternative for Cell Culturing: Potential in Cultivated
Meat Production. *Acs Food Science & Technology, 3*(4), 699-709.
doi:10.1021/acsfoodscitech.2c00401
Yamanaka, K., Haraguchi, Y., Takahashi, H., Kawashima, I., & Shimizu, T. (2023).
Development of serum-free and grain-derived-nutrient-free medium using
microalga-derived nutrients and mammalian cell-secreted growth factors for
sustainable cultured meat production. *Sci Rep, 13*(1), 498. doi:10.1038/s41598-
023-27629-w
1032
1033

A

1: Culturing of microorganisms

2: Lysate extraction by sonication.

3: Filter sterilization of lysate.

4: Addition of lysates to B8 media for iBSCs and MACK1 cells.

5: Analyzing cell proliferation and differentiation

B

Protein yield (mg) per L of culture

BS   ECN   VN   BL21   LL   CEN.PK   LP

Strain

C

KDa

Ladder   BS   ECN   VN   BL21   LL   CEN.PK   LP