Comparative spatiotemporal single cell transcriptomes reveal rewiring of pre-existing regulations during emergence of Kranz anatomy in C₄ grasses

Caiyao Zhao,[1*] Jianzhao Yang,[2,4*] Anting Ni,[1,3,11*] Hanyang Chen,[1,3*] Hong Su,[2,4*] Yating Cheng,[2,4*] Xiaoya Li,[2,8] Liyuan Zhong,[1] Dongming Fang,[1] Hao-Ran Sun,[4,6] Juan Yi,[2,4] Ying Wang,[2,4] Yanjie Wang,[2] Ming-ju Amy Lyu,[2] Xiaoxiang Ni,[2] Lichuan Chen,[5] Huanjin Li,[1,3] Guichao Yu,[2,4] Yumeng Wang,[2,4] Rulei Chen,[2,4,10] Hanyue Yu,[2,4] Wenwen Shao,[1,3] Haixi Sun,[4,5] Yinqi Bai,[5] Chuanshun Li,[7] Yang Dong,[4,6] Changsong Zou,[8] Xiaoyu Tu,[7] Shisong Ma,[9] Xing Guo,[1,3] Gengyun Zhang,[3] Peng Wang,[2] Huan Liu,[3] Tong Wei,[1#] Ying Gu,[1#] Xun Xu, [5#] Xin-Guang Zhu[2#]

BGI Research, Wuhan, 430047, China

National Key Laboratory of Carbon Capture, Center for Excellence in Molecular Plant Sciences, Institute of Plant Physiology and Ecology, Chinese Academy of Sciences, Shanghai, 200032, China

State Key Laboratory of Agricultural Genomics, BGI Research, Shenzhen, 518083, China

University of Chinese Academy of Sciences, Beijing, 100049, China

Key Laboratory of Genomics, Ministry of Agriculture, BGI Research, Shenzhen, 518083, China

State Key Laboratory of Plant Diversity and Specialty Crops, Institute of Botany, Chinese Academy of Sciences, 20 Nanxincun, Xiangshan, Beijing 100093, China

Joint Center for Single Cell Biology, School of Agriculture and Biology, Shanghai Jiao Tong University, Shanghai 200240, China

School of Life Sciences, Henan University, Kaifeng 475001, China

School of Life Sciences, Division of Life Sciences and Medicine, University of Science and Technology of China, Innovation Academy for Seed Design, Chinese Academy of Sciences, Hefei 230027, China

National Center for Gene Research, Key Laboratory of Plant Design, Centre for Excellence in Molecular Plant Sciences, Chinese Academy of Sciences, Shanghai 200032, China

College of Biological & Pharmaceutical Sciences, China Three Gorges University, Yichang, 443002, China

* These authors contributed equally to this work.

To whom correspondence should be addressed. Email: weitong@genomics.cn, guying@genomics.cn, xuxun@genomics.cn, zhuxg@cemps.ac.cn

Summary

Many of the world's most productive food and bioenergy crops use C_4 photosynthesis, which
have high photosynthetic efficiency due to a Kranz anatomy-based CO_2 concentrating
mechanism. Here, we took a comparative transcriptomics approach using single cell spatial
transcriptomes of leaf primordia for maize (*Zea mays*) and single cell RNA-seq (scRNA-seq)
atlases of corresponding leaf tissues for three C_4 species (*Zea mays*, *Sorghum bicolor*, *Setaria*
viridis) and one C_3 rice (*Oryza sativa*) to study regulatory networks involved in the
development and evolution of Kranz anatomy. We show that the formation of Kranz anatomy
involves extensive recruitment and modification of pre-existing regulatory modules,
especially the SHR-SCR module and the auxin signaling pathway. Members of the
INDETERMINATE DOMAIN (*IDD*) family transcription factors, including *IDD7* and *IDDP1*,
contribute to the modification of the involved modules. These extensive recruitment and
modification of existing genetic regulatory programs underlie the repeated emergence of C_4
photosynthesis during evolution.
54

55 **Keywords**
56 C_4 plants, photosynthesis, Kranz anatomy, Spatiotemporal transcriptome, scRNA-seq, gene
57 regulatory network
58

Introduction

Given the high light, nitrogen and water use efficiency, C_4 plants hold great potential in protecting global food security.[1] C_4 plants have evolved independently over 60 times historically.[2] Understanding the mechanisms of C_4 plant development and evolution could potentially enable the engineering of C_4 photosynthesis in C_3 crops.[3] The superior photosynthetic properties of C_4 plants are primarily due to a CO_2 concentrating mechanism based on Kranz anatomy, which facilitates the compartmentalized metabolism in bundle sheath cell (BSC) and mesophyll cell (MC). So far, metabolic structure and morphological features associated with C_4 photosynthesis are well established.[3,4] In contrast, regulatory mechanisms underlying these different properties are largely unknown, although a number of regulators controlling Kranz anatomy have been identified, such as *GOLDEN2* (*G2*), *SHOOT ROOT1* (*SHR1*), *SCARECROW1* (*SCR1*), *TOO MANY LATERALS1* (*TML1*)) through traditional forward genetics approaches. [5-12]

High-throughput RNA-seq methods emerged as a promising approach to reveal the mechanisms controlling Kranz anatomy development.[13-29] For example, Wang et al (2013, 2017) performed comparative analysis on different leaf primordia of maize and identified *GLK* as a major regulator of BSC development, which might have facilitated the transition from C_3 to C_3-C_4 transition.[18,30] Camo-Escobar et al. (2024) used scRNA-seq to show a significant reduction in the proportion of BSC and a downregulation of the C_4 shuttle system in the *shr2* mutant.[24] By using multiplexed in situ hybridization, Perico et al. (2024) showed the spatiotemporal expression dynamics of auxin- and other vein-related genes during the development of different-rank veins in the maize leaf, suggesting auxin plays a crucial role during Kranz development.[22] Given that the development of Kranz anatomy is highly coordinated between different cells types in space, spatial-temporal single-cell transcriptomes of leaf primordia, in contrast to the earlier whole-tissue based transcriptomics approach, hold a great promise to reveal molecular mechanisms controlling morphogenesis of the Kranz anatomy in C_4 leaves.

In this study, we generated spatial enhanced resolution omics-sequencing (Stereo-seq) and scRNA-seq atlases for leaf primordia in maize to study the development of Kranz anatomy. With these spatial scRNA atlases, we constructed a gene regulatory network for different cell types and identified regulatory cascades that control the vein initiation and Kranz anatomy development. By also collecting and analyzing scRNA-seq atlases for leaf primordia in sorghum, foxtail-grass, and rice, we identified the new regulatory mechanisms controlling features of Kranz anatomy in C_4 grasses. Furthermore, we show that the well-known regulatory modules including the SHR-SCR module, were recruited and modified during the emergence of C_4 photosynthesis in grasses, with the *INDETERMINATE DOMAIN* (*IDD*) family gene members involved in the altered regulation. This study shows that there is extensive

96   rewiring the regulatory programs during the emergence of C_4 photosynthesis in the grass
97   family.

Results

Spatial transcriptome atlas for early developmental leaves of maize differentiates the subtle differences among cell types

To profile the transcriptome of cells of early stage Kranz anatomy, we conducted the
spatial transcriptome of maize leaf primordia, consisting of P3-P6 on the section of base stem
at the stage of 12-day-old maize seedling (Figure 1A). We extracted *in situ* transcriptome of
individual cells based on their cell wall outlines stained with fluorochrome (see details in
materials and methods, Figure 1B) and obtained the transcriptomes of 14,037 cells (Figures
1A and 1B). The expression patterns of known marker genes are in strong alignment with
previously established knowledge (Figure 1C), confirming the reliability of the spatial
transcriptome data. Within the 17 unsupervised clusters, we identified most of the cell types of
maize primordium based on expression patterns of established marker genes (Figure 1D,
Figure S1), including two clusters of pavement cells (classified as epidermis), two clusters of
guard cells and/or subsidiary cells (GC/SC); five clusters of veins including phloem (cluster 7,
C7, companion cells), xylem cells (C13), procambium (C12) located in the leaf primordium,
older veins (C14) and sclerenchyma cells (C11) located in the leaf sheath (Figures S1B and
1C). There are also three clusters of dividing cells undefined, including two clusters of S-stage
cells and one cluster of G2M-stage cells (Figures 1D, Figure S1C). Among these cell types,
the two distinct clusters of epidermal cells are notably positioned within the inner and outer
regions of the leaf primordia, respectively. This spatial segregation indicates that the nuanced
differences between these two developmental stages of the epidermis can indeed be
accurately differentiated.

The mesophyll cells (MC) and bundle sheath cells (BSC) are mixed in two clusters (C5
and C0) located in inner and outer primordium (Figure S1C). Given the morphologically distinct
Kranz anatomy and the clear spatial arrangement of BSC and MC in an inner and outer
circular pattern around the vein, respectively, we directly labelled and extracted their
transcriptomes from Stereo-seq spatial data (Figure 1A and S2). Beside the cells of the leaf
sheath, we recognized 14 (sub-)cell types in one section of the leaf primordium, including three
BSC sub-cell types around three sub-cell types of veins (named main vein, rank1 vein and
rank2 vein, respectively) and three sub-cell types of MC (side MC is next to epidermis,
double-cell MC and single-cell MC are two and one cells of MC between veins) (Figure 1E,
Figure S2). Then we identified differential expressed genes (DEGs) of all (sub-)cell types
(Figure S3, Table S1). Based the fine classification of sub-cell type in the spatial transcriptome,
we observed that single-cell MC preferentially expressed a higher number of genes associated
with cell cycle processes, while double-cell MC and side MC exhibited higher expression
levels of genes involved in photosynthesis (Figure S4, Tables S1-2). This result indicates that

the three sub-cell types of MC are developmentally distinct in the primordium, which highlights

the value of our single cell resolution spatial transcriptome.

Identifying gradient expressed genes within the Kranz anatomy by a

pseudo-Kranz strategy

In the leaf primordium section with spatial transcriptome data, three distinct stages of

Kranz anatomy were identified: 544 Kranz anatomies with four BSC surrounding veins as

4BS-early stage, 813 Kranz anatomies with five unenlarged BSC as 5BS-middle stage, and

278 Kranz anatomies with enlarged BSC as 5BS-late stage (Figure 2A). [31] Since Kranz

anatomy at the same developmental stages are distributed throughout leaves, it is challenging

to derive the spatial relationships between genes, particularly low-expressed transcription

factors (TFs). To better visualize and identify gene expression patterns within the Kranz

anatomy, we superimposed the distributed Kranz anatomy of individual stages to create a

unified "pseudo-Kranz", with the center of each Kranz anatomy aligned at the origin of the

coordinate system (Figure 2B, see Materials and Methods for further details). Unlike Kranz

anatomy expression patterns in the spatial atlas, the expression of marker genes in the

pseudo-Kranz is clearer and more vivid (Figure 2C; Figure S5).

The pseudo-Kranz analysis enables the identification of genes that exhibits a gradient of

expression, progressing from the center to the periphery within the Kranz anatomical structure.

To do this, we segmented the pseudo-Kranz into eight concentric rings with uniform width and

calculated the average expression level of genes in each region. We then utilized weighted

gene co-expression network analysis (WGCNA) to classify these genes into separate modules

based on their expression patterns in the eight regions. We identified four modules of genes

preferentially expressed in the center of pseudo-Kranz, encompassing 435 genes highly

expressed at the center in all three stages (Figures 2D and 2E, Table S3). In comparison to

DEGs of three vein cell types in the spatial atlas, a greater number of marker genes for veins

were identified, for example, *ZmPIN1a*, *ZmSHR1*, *ZmGLK49*, *ZmHB21, ZmDOF11*, with the

first two known for their role in regulating vascular bundle differentiation (Figures 2F). [9,32,33]

Integrating spatial transcriptome with scRNA-seq atlas reveal the

co-expression patterns of *SHR* and *SCR* genes in early BSC

To comprehensively understand the continuous development trajectory of Kranz anatomy

across different stages, we complemented our Stereo-seq analysis with single-cell RNA

sequencing (scRNA-seq) on identical sections (see details in materials and methods). After

background RNA filtration with SoupX, 33,309 nuclei were obtained, and 30,548 genes were

detected. These nuclei are clustered distinctly into 29 clusters, most of which are annotated

with known marker genes (Figure 3A, Figure S6). In total, we found 8 clusters of epidermal

pavement cells at different developmental stages (C2, C3, C10, C14, C8, C15, C23, C28) and

two clusters of stomata cells (C22, stomatal precursor and C26, mature GC&SC). By

integrating scRNA-seq data with Stereo-seq, we distinguished different vascular cell types:

C16 and C27 represent xylem on the adaxial side of the vascular vessel; C7, C9, C11, C13,
C19, C20 and C21 represent phloem on the abaxial side (Figure S7). C12 with the high
expression levels of markers, such as *ZmLAX2*, *ZmMPa*, represents procambium cells; C18
with *ZmRS1*, *ZmLG3*, *ZmRss3* represents meristem; C1 with *ZmMAD2*, *ZmUBC19*,
ZmCYC23 represent G2M stage cells. C5 is assigned as MC expressing gene encoding *CA*,
PEPC and *NADP-MDH* and C25 as BSC expressing genes encoding NADP-ME, SBPase and
FBPase. C5 and C25 are more similar to MC and BSC in leaf sheath, respectively, rather than
that of leaf primordium in spatial transcriptomics atlas (Figure S6).

Besides the MC and BSC clusters, we found that C0 also expressed relatively low levels
of photosynthetic genes and relatively high levels of cell division related genes (Figure S6). We
also found that C0 is more similar to the cells in primordium than that of leaf sheath (Figure S7).
These results indicate that C0 possibly represents the mixed cells including BSC and MC at
early developmental stages in primordium (Figure S7). To resolve the early MC and BSC, we
subset four clusters at early developmental stages, including C0, C1, C4, C12, as well as MC
of C5 and BSC of C25 by supervised clustering (see details in methods). We combined the
2000 variable genes identified by Seurat in these subset cells and DEGs identified in spatial
transcriptome to conduct the dimensionality reduction of PCA analysis and then clustered
these cells using top30 PCAs (Figure 3B). We assigned all 10 sub-clusters and distinguished
MC and BSC at early developmental stages (Figures 3B-E). The sub-cluster 2 (sub-C2), a
subset of cells comprising of cells within C4 and C0, was characterized by the high expression
of marker genes preferentially expressed in the BSC identified in the spatial atlas (Figure 3E).
We also found that sub-C2 is closely connected with sub-cluster of mature BSC in the
pseudo-time trajectory analysis (Figure 3D) and have relatively high similarity with the cells
around veins in the primordium (Figure S8). Furthermore, DEGs of sub-C2 were preferentially
highly expressed in the region of BSC in pseudo-Kranz (Fig. 3F). The early MC were
subdivided into Sub-C0 (inner early MC) and Sub-C9 (outer early MC), which are more similar
to MC of very inner leaf primordium and outer leaf primordium, respectively (Figure S8).

The identification of these BSC at different developmental stages allowed us to study the
developmental trajectory of early stage BSC (Figure 3G). There are total 5193 genes
(including 255 TFs) expressed variably along the developmental pseudo-time trajectory
(Figure 3G, Tables S4). As expected, BSC at early stage of pseudo-time expressed cell
cycle-related genes, such as *ZmHIS2B5*, *ZmSMR1* and *ZmROA1* and enriched GO terms,
such as "chromosome organization" and "DNA packaging" (Figure 3H, Tables S4-5). There are
also a few of TFs, such as *ZmC78*, *ZmRS2*, *ZmGATA29* (*ZmGNC*), preferentially highly
expressed at the early stage of pseudo-time. At the late stages of pseudo-time, there was a
progressive upregulation of photosystem genes and genes related to cell wall modification,
such as *ZmPLT2*, *ZmXTH9*, *ZmEXPA4*, *ZmXT2* (Figure 3H, Table S4). Interestingly, we found
that *ZmSHR1*, *ZmSHR2*, and *ZmSCR1* genes were both expressed at the early stage of
pseudo-time in BSC (Figure 3H). Although *SHR* genes and *SCR* genes were predominantly

expressed in early veins and MC, respectively, we observed their co-expression in early BSC
(sub-C2) of scRNA-seq data (Figure 3I) and notably within confined single cells in spatial atlas
(Figure S9). This suggests that, beyond their established roles in veins and MC[8-10,12,34], *SHR*
and *SCR* genes maybe also play important roles in establishing the identity of BSC in Kranz
anatomy.

Spatiotemporal co-expression network reveals the regulatory cascades underlying vein initiation and development

After we identified single cell transcriptome of most cell types of early-stage leaf,
especially BSC at the early stage of Kranz anatomy, we conducted the co-expression network
analysis of genes with Hotspot[35] in these cells to mine key regulatory genes over leaf
morphogenesis. Informative genes were filtered by gene autocorrelation and were clustered
into 25 co-expression modules (Figure 4A, see details in methods). We identified specific
modules associated with each cell type through mapping the expression patterns of each
module in two ways: by mapping module scores of individual cells onto the UMAP of the
scRNA-seq transcriptome and by displaying the total gene counts for each module on the
pseudo-Kranz (Figure S10, Table S6). For instance, module 13 (M13, including *ZmSMXL3* and
ZmSHR1, *ZmSHR2*) is associated with veins, which have high module scores in sub-C3 and
sub-C4 and only highly expressed in the center of pseudo-Kranz. M7 (including marker genes,
ZmSSU2 and *G2*) is associated with BSC, which is featured with high module scores in the
sub-C2 and preferentially highly expressed in the region of BSC in the pseudo-Kranz. M9
(including markers, such as *ZmGLP1* and *ZmCP29*) is associated with MC of Kranz anatomy
(Figure 4B, Table S6).

Interestingly, there were at least 6 modules (M3, M4, M5, M11, M12, M13) associated with
veins, all of which have preferentially high expression levels in center of the pseudo-Kranz
(Figure S10). To elucidate the spatial heterogeneity of gene expression for each module, we
computed and graphed the mean normalized gene expression for each of the six
vein-associated modules across the spatial atlas (Figure S11). By plotting the expression
profiles of vein-associated modules across the spatial atlas, we found that M11 and M13 are
predominantly higher expressed in small veins throughout the inner primordium, whereas M5
and M3 are mainly expressed in relatively mature lateral veins and midveins (Figure 4C).
Additionally, M4 shows higher expression in more mature veins across the leaf sheath (Figure
4C). Consistent with the observed spatial heterogeneity, our analysis reveals that these
modules are preferentially expressed in different sub-clusters of veins. For example, M3 and
M4 have higher module scores in sub-C6 of vein, and M12 in sub-C6 and sub-C3, M13 in
sub-C3, sub-C4 and sub-C5 (Figure S10).

We constructed the gene regulatory network (GRN) of veins within the leaf primordium by
integrating co-expression relationships with potential TF-target interactions (see details in
methods). The GRN clearly illustrates the regulatory cascades originating from TFs within

modules M11 and M13, which subsequently influence modules M12 and M5, leading to

downstream effects on modules M3 and M4. (Figure S12). It is noteworthy that key markers of

vein initiation, including auxin-responsive factors (*ZmMPa* and *ZmMPb*) and auxin transporters

(*ZmLAX2*, *ZmLAX3*, *ZmPIN1a* and *ZmPIN2*), are grouped in modules M12 and M5. This

suggests that upstream regulators from M11 (such as *DOF* family genes) and M13 (such as

EREB family genes, *ZmSHR1* and *ZmSHR2)* might influence vein initiation, thereby potentially

controlling vein density. Additionally, there is a gene related to IAA transport, *BNLG1484*,

which is the ortholog of *WAG1* and is potentially involved in establishing the auxin maximum

during the initiation of veins and Kranz formation (Figure S12). This temporal regulatory

network, spanning from vein initiation to development, provides a rich resource to support

studying vein differentiation in maize (Figure S12).

To further dissect the regulatory cascades underlying vein development, we integrated

our co-expression analysis with the construction of a developmental trajectory. This trajectory

delineates the pseudo-time of developmental progression, tracing the path from median

ground meristem (mGM, sub-C4) through early vein stages (sub-C3) to more mature veins

(sub-C6) (Figure 4D). By mapping the module scores onto this trajectory for each cell, we

observed distinct temporal expression patterns: M11 and M13 exhibits peak expression at

early developmental stages, with expression levels subsequently declining as veins developed.

Conversely, M12, M5, M3, and M4 display an inverse pattern, with expression increasing over

the course of vein development (Figure 4E). The 4626 significantly variable genes along with

pseudo-time can be divided into five clades by their expression pattern through pseudo-time

(Table S8). By cross-referencing the genes from various vein-associated modules with those

that exhibit pseudo-time-related expression patterns, we refined the pool of candidate genes

involved in vein initiation and development (Figure 4F). Specifically, M11 and M13 overlap

exclusively with genes expressed during the early pseudo-time stages, confined to clusters C1

and C2. In contrast, M12 and M5 show the overlap with genes expressed across stages C2,

C3 and C4, while M3 and M4 aligned only with genes expressed during the later pseudo-time

stages, specifically within clusters C4 and C5. These findings reinforce the existence of

regulatory cascades spanning the six vein-associated modules and highlight potential master

regulators governing the critical early phases of vein initiation (Figure 4F).

Gene regulatory network identifies regulators of Kranz anatomy development in

leaf primordium

For a systemic overview to identify new regulators involved in Kranz anatomy, we

constructed a GRN of leaf primordium that includes all TFs within the co-expression modules

(Figure 4G). As an application of this new transcriptomics atlas and cell specific GRN, in

addition to a few of established markers, we identified many new regulators related to

morphogenesis of Kranz anatomy. The *ZmSCR1* gene exhibits predominant expression in the

MC of C_4 plants (Figure S13), and its role in influencing the number of MC during Kranz

anatomy development has been substantiated.[12] Therefore, it is plausible to hypothesize that
TFs co-expressed with *ZmSCR1* might act within the same regulatory pathway that regulates
Kranz anatomy development. Among the most highly correlated TFs with *ZmSCR1* and/or
ZmSCR2 are *ZmNKD1*, *ZmIDD7*, *ZmIDDP1*, *ZmMYBR35*, *ZmHSFTF19*, *ZmBHLH23*, and
ZmNKD2 (also known as *IDD22*, Figure 4G). These candidate regulators were mainly
expressed in early MC of leaf primordium but not in mature MC, suggesting that these TFs are
very likely involved in morphogenesis and development of early-stage MC. It has been verified
that *ZmSHR1* and *ZmSHR2* play crucial roles in the regulation of vein and MC development in
Kranz anatomy.[7,9,17,34] Within the GRN, a lot of TFs are predicted to directly regulate *ZmSHR1*
and *ZmSHR2*. Notably, the auxin response factor *ZmARFTF23*, *ZmGLK49* and *YABBY*
transcription factor *ZmYAB10* predominantly expressed within the mGM cluster (sub-C4), may
affect the early expression of *ZmSHR1* and *ZmSHR2* during the initial stages of vein
development in the mGM region (Figure 4G, Figure S12). Interestingly, we found that many
TFs predicted to regulate *ZmSHR1* and *ZmSHR2*, such as *ZmGLK6*, *ZmYAB15* and *ZmDOF22*,
exhibit dual expression in both vein and BSC, while some of the TFs in M7, the module
particularly associated with BSC, were co-expressed in both BSC and MC (Figure 4G, Figure
S13). The expression patterns indicate that there are possible regulatory cascades affecting
the Kranz anatomy development from vein through BSC to MC.

Comparing the scRNA-seq atlas between C$_4$ and C$_3$ leaves identifies the
innovations for the developmental programs of Kranz anatomy
To identify new inventions during the evolution of C$_4$ Kranz anatomy, we collected more
scRNA-seq atlases for early developing leaves of C$_4$ plants in grass, including *Sorghum*
bicolor (hereafter called sorghum) and *Setaria viridis* (hereafter called Setaria) and one C$_3$
plant of *Oryza sativa* (hereafter called rice) as control group. Similar to the annotations of
maize scRNA-seq data, we identified most of the clusters of these three atlases through the
expression patterns of known marker gene orthologs in each of these species (Figures S14-16,
Table S9). In total, we obtained 24 (sub-)cell types in the four atlases, which include most of
cell types in the sampled tissues (Figures 5A-D, Table S9). Even though each cell type
occupied different proportion of cells in each species, we found highly conserved shared
genes in each cell type, such as *PIN1a* and *LAX2* genes in procambium, *SUT1* in phloem
parenchyma, *XCP1* in xylem (Figure D). Besides these shared marker genes in each cell type
among these four species, we also identified many new marker genes for each cell type, which
are conserved within four species (Table S10).
The high level of overlap between the homologous genes in different cell types among the
four species promoted us to further identify the new regulators involved in Kranz anatomy
evolution. To do this, we first identified the conserved DEGs specifically present in each cell
type of Kranz anatomy in C$_4$ plants, then we compared these to that of corresponding cell type
in rice to distinguish conserved genes among C$_3$ and C$_4$ plants or C$_4$-specific markers. There

were 106 DEGs shared by MC&BS among C_4 and C_3 plants (since here we could not
differentiate MC and BSC in sorghum, Setaria, we used MC&BSC to represent the group
including these two cell types), such as *NKD1*, *TCP23*, *G2*, *IDDP1*, *IDD7* and *ARFTF25*, that
may perform the basic function during the development of MC and/or BSC (Figure S17, Table
S10). In addition, we also found DEGs specifically identified in BSC&MC of C_4 plant, such as
SCR1, *SCR2*, *bHLH93*, and *IAA5* (Figure S17). The procambium is a key cell type whose
initiation influences the vein density and development of Kranz anatomy. We found that
procambium of three C_4 species exhibited 174 common DEGs, with a large portion of 116
genes also present in rice procambium, while only a small subset of 58 genes was unique to
the C_4 plants (Figure 5E, Table S10). This result indicates a high degree of conservation in the
developmental programs of procambium between C_4 and C_3 plants. We found not only many
well-known marker genes associated with vein development, such as *LAX2*, *MPa*, *PHB*,
WOX13A, but also many new markers, such as *HB124*, *DOF49*, *DOF17*, *bHLH48*, *NACTF46*,
LBD26, *LBD12*, *RLD1* and *CCHH25*, which were shared among C3 and C4 plant. We also
found some C_4 specific marker genes of procambium, such as *SHR1*, *SHR2*, *HB78* and
ARFTF35 (Figure 5E).
Surprisingly, expression level of *SHR1* and *SHR2* genes in the procambium and G2M
phase cells of C_4 plants is significantly higher than that in C_3 rice, just as the expression levels
of *SCR* genes in MC&BS of C_4 species are higher than that in C_3 rice (Figure 5F and S17).
Indeed, *SHR* genes are expressed in most cell types of vein in the three C_4 plants, including
phloem sieve, phloem parenchyma, xylem, and xylem parenchyma, procambium and even
mGM, while rice *SHR* genes are only highly expressed in clusters of phloem sieve and xylem
parenchyma in leaf and rarely expressed in procambium and other early-stage cells (Figure
S18), indicating that *SHR* genes are initiated earlier and/or higher, resulting in a wider range of
expression in C_4 leaves than C_3 leaves. Using public data of transcriptomics
(https://www.plantrnadb.com/), we further found that the ratio of expression levels of maize
SHR genes in leaf to that in root was higher than that of rice *SHR* genes (Figure S19).
The level of auxin in leaves is crucial for controlling vein patterning and leaf
morphogenesis. We investigated key genes related to auxin, including biosynthesis, transport,
and signaling (Tables S11-12). Contrary to expectations, we did not find significantly higher
expression of biosynthesis genes or distinct expression patterns of signaling genes in C_4
compared to C_3 plants, which might be applied to explain the more veins in C_4 than C_3 plants
(Figures S20-21). However, most IAA transport genes, including *PIN* genes responsible for
auxin efflux and *LAX* genes for influx, showed preferential high expression in specific cell types
(Figure S22). For instance, *PIN1a* gene were expressed in the procambium of both C_4 and C_3
plants, with relatively higher expression in C_4 plants. In contrast, *PIN1b/c* genes were more
expressed in C_3 rice procambium. Notably, *PIN8* gene expression in C_4 plant procambium was
significantly lower compared to C_3 rice, potentially influencing auxin levels and vein initiation in
C_4 plants (Figure S22).

The developmental programs of Kranz anatomy are mostly recruited from C$_3$ leaves with modifications

The GRN of early-stage leaves in the other three species were constructed using the same method as in maize. We then established a GRN for the developing leaves by comparing the conserved network in the three C$_4$ plants with that of corresponding cells in the C$_3$ rice (see details in methods). In this network, all known TFs involved in the development of Kranz anatomy are included. For instance, *DOT5*, also known as *TML1*, which was predicted to directly regulate *SHR1* and *SHR2* in this network (Figure 6A), was recently shown to specify small vein type in C$_4$ and C$_3$ grass leaves. *NKD1*, one of *IDD* family genes, which was predicted to directly regulate *SCR1* in this network, modulates *ZmSCR1* expression and impacts the MC number between veins in maize.[14] This network also included additional *IDD* family genes that were predicted to directly regulate *SCR1* (Figure S23). Here, we verified the functions of other two *IDD* genes in the morphogenesis of Kranz anatomy in maize. We observed that the MC number between veins were significantly decreased in the *idd7* and *iddp1* mutants (Figure 6B, 6C and S24), comparable to those of *scr1* mutants.[8,12] We then found that the expression levels of *SCR1* gene in both mutants of *idd7* and *iddp1* were lower compared with wild type control (Figure 6D), which suggests that *IDD7* and *IDDP1* participated in the development of Kranz anatomy, very likely through regulating expression levels of *SCR1* gene.

Analysis of this GRN shows that many conserved co-expression relationships of TFs in veins among three C$_4$ plant were also found in C$_3$ rice, which suggests a high level of conservation for the regulatory network for vein development between C$_3$ and C$_4$ (Figure 6A). As shown in the network, the relationship among many key hubs, such as *HB46*, *GLK6*, *EREB41*, *DOF23*, *MPa*, *BHLH156/48/138/143/27*, *DOF49*, *NACTF46*, *NACTF131*, *GLK27*, are conserved between C$_3$ and C$_4$ plants (represented as the gray lines in Figure 6A). Of course, there are also some genes, such as *SHR1*, *SHR2*, *DOF27*, *YAB10*, *ARFTF23*, *YAB9*, *NACTF77*, *BHLH169*, *CCHH25*, *CCHH2* and *DOT5*, which establish C$_4$ specific co-expression relationships between each other gene and hence may play an important role in Kranz anatomy (represented as the orange lines in Figure 6A). These recruited regulators in the M11 and M13 include *YAB5*, *YAB10*, *YAB11*, *YAB9*, and *EREB41*, which are associated with the initiation and early development of veins (Figure 6A). We also identified the recruited regulatory relationships associated with MC in C$_4$ leaves, such as those among *SCR1*, *SCR2*, *IDD7*, *NKD1*, and *IDDP1* (Figure 6A). By comparison, we identified only a limited number of novel regulatory relationships in BSC (Figure 6A), compared to either MC or vein development.

Discussion

This study generated comprehensive spatial and scRNA-seq atlases for the young leaves of four grass species, including three C_4 plants and one C_3 plant as a control. Using these atlases, we successfully identified the transcriptomes of nearly all cell types present in the young leaves. The shared preferentially expressed orthologs suggests a high level of conservation of core functional pathways within the same cell types across grass species (Figures 5A-C, S17). These comprehensive atlases and the derived genetic regulatory networks combined with the pseudo-Kranz anatomy offer an opportunity to study the regulatory mechanisms associated with the development of different tissues or cells in a leaf.

Here we use the development of bundle sheath cell to further illustrate how these new data set allow us to identify regulatory mechanisms controlling differentiation of different cell type. BSC represents a key feature of C_4 Kranz anatomy. In this study, by profiling the single-cell-resolution spatial transcriptomes, we can identify the subtle differences between early-stage BSCs and MCs, which were not distinguishable through unsupervised clustering of scRNA-seq data. The early stage of BSC shows characteristic expression of genes that are also preferentially expressed either in veins, such as *ZmSHR1*, *ZmSHR2*, *ZmMPb*, and *ZmANT3*, or in MC, like *ZmSCR1* and *ZmGATA29* (Figures 3H and S13), suggesting that early-stage BSCs share many regulatory relationships with both veins and MC. In addition to *ZmMPb* (Figure 3H), there are also many auxin-responsive factors that are differentially expressed at either the early- or late-stages of BSC, including 8 genes at the early stage and 18 genes at the late stage (Table S4), which indicates auxin plays a key role throughout the entire developmental trajectory of BSC. We further observed many *COL* family genes were expressed in late pseudo-time stages of BSC, such as *ZmCOL2*, *ZmCOL7*, *ZmCOL8*, *ZmCOL11* (Table S4). Therefore, *ZmCOL* genes might also be a major regulator of BSC development, possibly for chlorophyll accumulation, in addition to their other commonly known roles in flowering time control, responses to light, and auxin homeostasis in leaves.[36-38]

This pseudo-time series and derived GRNs allow us to construct a complete developmental trajectory controlling Kranz anatomy. The development of Kranz anatomy involves three cascades of events: initiation and development of veins, the differentiation of cell types, and cell specific development of photosynthetic machineries.[4] The different cell types show drastically different levels of developmental complexity. For example, development of vein involves 6 modules, which represent a cascade of developmental changes (Figure 4G and S12). Several well-known marker genes related to initiation and early development of vein, such as *SHR1*, *SHR2*, and *ANT3*,[7,9,21,39] are present along with a number of novel TFs, which include six *DOF* genes, three *EREB* family genes, two *GLK* genes and *WRKY107* (Figure 4G). Different regulators participated in middle- or late-stages of vein development as well (Figure 4G), e.g. *LBD* genes and *bHLH* family genes are enriched in M5, while *NAC* family genes are enriched in the late-stages of vein development. One notable phenomenon in the genetic regulation over vein development is that most of the involved genes have far more edges

linking to other regulators, suggesting the high complexity of genetic regulation over vein
initiation, compared to BSC or MC development. The developmental programs controlling
BSC mainly involve M7, in which *G2* and *ZmTCP23* have the most linkages to other regulators.
The developmental programs over MC involve M9, which includes many known regulators e.g.
ZmSCR1, *ZmSCR2*, but also many new members, such as members of the *IDD* gene family,
HY5, etc (Figure 4G). Here, we acknowledge that the well-known *SHR-SCR* regulatory module,
which plays a crucial role in root development,[40] also plays crucial roles in the development of
Kranz anatomy, which is also supported by genetic evidences.[7-10,12,14,17,34] It is worth
emphasizing that these members identified to be involved in Kranz anatomy development
need particular spatial and temporal combination to ultimately define the formation of Kranz
anatomy. These identified modules and their members provide the first step for us to
reconstruct the detailed genetic mechanisms controlling Kranz anatomy.

One of the major research goals of studying Kranz anatomy development is to identify the
key evolutionary innovations promoting development of Kranz anatomy, which can be the most
promising targets to engineer for the future purpose of creating a Kranz anatomy in C_3 leaves.
We tackled this challenge through a comparative analysis of genetic regulatory networks
between C_3 and C_4 leaves (Figure 6A). Though BSC is commonly regarded as a special
adaption in C_4 photosynthesis, BSC in C_4 leaves only contains a relatively few of new
regulatory relationship in early developmental stage (Figure 6A). In contrast, MC in C_4 leaves
gained large number of new regulatory relationships. In particular, there are a large number of
new regulatory connections developed between *SCR1* and its upstream regulators, including
IDD7, *IDDP1*, and *NKD1* (also a member of IDD gene family), and surprisingly *HY5* (Figure
6A). Interestingly, these *IDD* family genes also confer a direct regulation over the expression of
G2, an important regulator of the BSC development as well (Figure 6A).[5,6] Possibly even more
surprisingly, the vein development in C_4 leaves gained the highest number of new regulatory
relationships (Figure 6A), suggesting the primary importance of vein development in Kranz
anatomy differentiation and should be highly researched to support future C_4 engineering.
Interestingly, in the modules controlling vein initiation, a large number of new regulatory
connections were established between *SHR1*, *SHR2* and their regulators, including a number
of *YABBY* family members, *EREB41*, *GLK6* etc. Considering that both *YABBY* family genes
and *EREB* family genes are also closely related to drought responses of plants[13,41-45] and
these genes were upregulated in drought stress treatment (data from NCBI project number
PRJNA421180 collected in transcriptome database http://ipf.sustech.edu.cn/pub/zmrna/),[46] the
genetic programs controlling adaption of C_3 leaves to drought might be recruited to increase
vein density during C_4 evolution.[47,48]

Further examination of those expressional innovations in C_4 leaves reveals mechanism
controlling the evolution of Kranz anatomy. Compared to C_3 leaves, *SHR* genes in C_4 leaves
are expressed at higher levels and across a broader range of cell types within the veins; the
expression of *SCR1* increased in MC as well (Figures 5F and S18). Overexpression of *SHR*

genes in both C_3 and C_4 plants typically result in an increase in the number of MC between
veins and a decrease in the vein-to-MC ratio.[7,9,34] The mutant *scr1* showed decreased number
of MC between veins.[8,10,12,14] Therefore, in C_4 leaves, the changed expression of both *SHR1*
and *SCR1* should have contributed to increased cell division to generate more MC between
veins during leaf development. The decreased number of MC between veins during C_4 plant
evolution therefore can only be attributed to the increased initiation of procambium cells and
hence minor veins. Auxin has been suggested as a major player in controlling the increased
vein density in C_4 leaves.[34] However, in C_4 grasses studied here, we did not notice increased
expression of genes related to auxin synthesis (Figure S20). In contrast, a few genes related
to auxin transport, e.g. *PIN1a*, *PIN8*, showed altered expression, which may contribute to
elevated auxin levels in the ground tissues farther from the veins. Considering that *pin8* mutant
cooperated with *pin6* increased vein branch points per area and *PIN8* overexpression
exhibited simpler vein networks in *A. thaliana*,[26] the decreased *PIN8* expression may be one of
the key innovations to increase the vein numbers in C_4 plants. These altered expression of either *SHR1* and *SHR2*, *SCR1*, and even auxin transport proteins are caused by the newly
either *SHR1* and *SHR2*, *SCR1*, and even auxin transport proteins are caused by the newly
acquired regulatory relationship as discussed earlier, such as *DOT5* which possibly regulates
SHR1 and *SHR2* and three *IDD* genes—*NKD1*, *IDDP1*, and *IDD7*—which have been
confirmed to regulate *SCR1* (Figure 6A).[14,49] All these showcase a common theme of complex
trait evolution, i.e. complex traits evolved through effective reutilization of existing mechanisms
with changed expression patterns of key players. Such recruitment of existing regulatory
mechanisms during the evolution of Kranz anatomy might underlie the repeated emergence of
C_4 photosynthesis during history.

Limitations of the study

By comparing spatiotemporal scRNA-seq data in three C_4 plant with C_3 rice, we
constructed a conserved gene regulatory network for early developing leaves. This enabled us
to identify most of the known regulators for Kranz anatomy development, as well as numerous
new TFs. Although we have validated two *IDD* family TFs affecting the number of MC between
veins in Kranz anatomy, there are additional key hub genes in the network that need to be
further verified for the evolutionary and developmental program of Kranz anatomy. In this work,
we focused on the regulatory mechanism controlling the morphogenesis of Kranz anatomy.
For fully elucidating the Evo-devo of C_4 photosynthesis and engineering C_4 system in C_3 crops,
it is also essential to reveal the regulatory mechanisms of establishing the C_4 metabolism in
mature Kranz anatomy.

Data and code availability

512 The raw sequence data generated by Stereo-seq and scRNA-seq have been deposited in

CNGB Nucleotide Sequence Archive (project accession code: CNP0005465). To facilitate the application of this data source in plant biology research, we deposited the analyzed spatial data in Spatial Transcript Omics DataBase (STOmics DB) (ID STT0000093, https://db.cngb.org/stomics/project/STT0000093_5850b1c5/reviewer_link/).

All codes supporting the current study have been deposited in GitHub (https://github.com/jianzhaomichaelyang/STomics-C4Kranz).

Additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Acknowledgements

We thank Prof. Qihua Ling and Prof. Jiawei Wang from CEMPS for giving advice. We thank Xiaoyuan Wu from IBCAS for seeds of sorghum BTx623. We thank Bohan Zhang for help in Stereo-seq experiments; Fengqin Dong from IBCAS for help in semi-thin sections; Dr. Hui Zhang, Fuyu Li, and Feng Han, Yuhui Huang, Dr. Qingfeng Song and Linxiong Mao from CEMPS for help in data analysis and experiments; we thank Prof. Yongrui Wu, Dr.Jiechen Wang, Dr. Luhuan Ye, Lihao Chen, and Rui Yang for help in maize field planting; we thank Dr. Lina Shi, Xinyu Liu, and Fengping Ni for help in fund management. This study was supported by National Key R&D Program of China (No.2023ZD04073 to H.L.), Strategic Priority Research Program of the Chinese Academy of Sciences (grant number: XDB0630100 to XG.Z.), the general program of the National Science Foundation of China (31870214 to XG.Z.), the National Key Research and Development Program of China (2020YFA0907603 to XG.Z.) and the Shenzhen Science and Technology Program [KQTD20230301092839007 to C.T.] and the Science and Technology Major Special Project of Shenzhen (KJZD20230923114607016 to H.L.).

Author contributions

X.G. Zhu and X. Xu conceived the idea; X.G. Zhu, X. Xu, Y. Gu and T. Wei supervised the work; X.G. Zhu, H. Liu, C.Y. Zhao and J.Z. Yang designed the experiments. H.Y. Chen, J.Z. Yang and C.Y. Zhao performed the Stereo-seq experiments with the help from W.W. Shao and L.C. Chen; J.Z. Yang, H.Y. Chen, H. Su, J. Yi, Y.T. Cheng and Y. Wang performed cell identification of Stereo-seq data. A.T Ni and C.Y. Zhao performed the scRNA-seq experiments. C.Y. Zhao, J.Z Yang, A.T. Ni, H.Y. Chen, H. Su and L.Y. Zhong performed data analysis. D.M. Fang, H.J. Li, H.X. Sun, Y.Q. Bai, S.S. Ma and X.Y. Tu provided essential advice for analysis. J.Z. Yang, Y.T. Cheng and X.Y. Li identified maize EMS mutants. J.Z Yang, Y.T. Cheng, H.R. Sun and Y. Dong performed resin section and microscopy imaging. C.Y. Zhao, J.Z. Yang, A.T. Ni and H.Y. Chen prepared the figures. C.Y. Zhao, J.Z. Yang, A.T. Ni and H.Y. Chen prepared the

549 manuscript with help from all authors. X.G. Zhu, X. Xu, Y. Gu, T. Wei, H. Liu, P. Wang, G.Y.
550 Zhang and X. Guo provided valuable advice and reviewed the manuscript. X.G. Zhu, X. Xu, Y.
551 Gu and T. Wei finalized the manuscript. All other authors contributed to the work. All authors
552 approved the manuscript for submission.
553

554 **Declaration of interests**
555 The authors declare no competing interests.
556

Supplemental information
Figure S1. Spatial transcriptome of maize leaf primordium
Figure S2. Representative picture of cell type recognized in spatial transcriptome
Figure S3. Expression patterns of marker genes identified in each cell type on Stereo-seq
maps
Figure S4. Expression patterns of photosystem related genes in three sub-types of mesophyll
cell
Figure S5. Expression patterns of marker genes of four cell types on three stages of
pseudo-Kranz
Figure S6. Dot plot of marker genes in the indicated cell types of maize leaf atlas
Figure S7. The integration of clusters of snRNA-seq atlas with cells of spatial
transcriptomes of maize
Figure S8. The integration of sub-clusters of snRNA data with cells of spatial
transcriptomes
Figure S9. co-expression patterns of *SHR* and *SCR* genes in spatial transcriptomes
Figure S10. Expression patterns of module genes in snRNA-seq and pseudo-Kranz
Figure S11. Expression patterns of individual vein-associated module genes spatial atlas
Figure S12. Regulatory co-expression network of vein related module genes
Figure S13. The expression patterns of candidate regulators of *SHR*, *SCR* and *G2* genes
Figure S14. Dot plot of marker genes in the indicated cell types of sorghum leaf atlas
Figure S15. Dot plot of marker genes in the indicated cell types of Setaria leaf atlas
Figure S16. Dot plot of marker genes in the indicated cell types of rice leaf atlas
Figure S17. Conserved DEGs in BSC and/or MC among C_4 and C_3 plants
Figure S18. The expression patterns of selected key genes in different cell types of C_4 and
C_3 plants
Figure S19. The expression levels of SHR genes in root and leaf of maize and rice
Figure S20. The expression levels of auxin biosynthesis genes in C_4 and C_3 species
Figure S21. The expression levels of auxin signal transduction genes in C_4 and C_3 species
Figure S22. The expression levels of auxin transport genes in C_4 and C_3 species
Figure S23. The relative positions of predicted binding sites of TFs co-expressed

with *SCR1*

Figure S24. The mutation site identification of EMS mutants

Table S1. The differentially expressed genes identified in each cell type in spatial

transcriptomes of maize early leaf

Table S2. The average expression of photosynthesis related genes in each cell type of spatial

transcriptome of maize early leaf

Table S3. Spatial heterogeneity of gene expression in three stages of pseudo-Kranz

Table S4. Marker genes variable along with pseudotime of early BS (sub-cluster 2) and relative

late BS (sub-cluster 8) in snRNA-seq transcriptome

Table S5. GO terms enriched in each cluster of marker genes variable along with pseudotime

of   early BS (sub-cluster 2) and relative late BS (sub-cluster 8) in snRNA-seq transcriptome

Table S6. The gene modules of snRNA-seq transcriptome of early maize leaf calculated by

hotspot

Table S7. The GO terms enriched for genes in each module of snRNA-seq transcriptome of

early maize leaf calculated by hotspot

Table S8. Marker genes variable along with pseudotime of veins related clusters (sub-C4,

sub-C3 and sub-C6) in snRNA-seq transcriptome

Table S9. Marker genes were used to annotate the cell types in scRNA-seq atlas of four

species

Table S10. The conserved DEGs of each cell type in C_4 plants

Table S11. The orthologs of IAA related genes in four species

Table S12. Mean expression of IAA related genes in the clusters of atlas in four species

Table S13. Annotation and gene names of maize genes

Table S14. Orthologs of maize genes in Sorghum bicolor, Setaria viridis and Oryza sativa

subsp. Japonica

Table S15. Primers list for EMS mutants genotyping and RT-PCR quantification

Table S16. Datasheet of RT-PCR quantification

Table S17. Datasheet of counting number of mesophyll cells separating veins

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Plants

In this study, we utilized four species: three C_4 plant *Zea mays* - B73; *Sorghum bicolor* -BTx623; *Setaria viridis* - A10) and one C_3 plant (rice, *Oryza sativa japonica*). Plants were cultured in greenhouse of China National Gene Bank (CNGB), Shenzhen, with an average temperature of 28℃ and 55% humidity was kept. The average light intensity was maintained at 500 µmol·m^{-2}·s^{-1} for 16 hours daylight and 8 hours dark. Plant tissues were sampled when maize grew around 12 days, sorghum around 12 days, Setaria around 10 days, and rice around 15 days, reaching the vegetative stage with three leaves. Tissues, approximately 5 mm above from ground, comprised apical meristem plus leaf primordia from P1 to P6, were harvested for both Stereo-seq and scRNA-seq analyses. For maize (*Zea mays)*, the outmost leaf sheath was stripped off.

METHOD DETAILS

Stereo-seq

To spatially visualize transcriptomic expression within leaf tissue, the STOmics gene expression reagent kit was used, following the protocol described by the Stereo-seq chip manufacturer.[48] The fresh tissue was immediately embedded in PVAL - OCT (PVAL by Shanghai M&G Inc., No. MF-7001, and OCT by SAKURA, No. 4583) and then frozen with liquid nitrogen-precooled isopentane. Cryosections, adhered to and fixed on the chip, were stained with Qubit™ ssDNA Reagent (SS, Invitrogen, No. Q10201) and Fluorescent Brightener 28 disodium (FB, Sigma, No. 910090), followed by fluorescence imaging. The tissue is then permeabilized based on the tissue-specific optimization protocols (BGI, No. 1000028492): 10 minutes for maize and sorghum, 30 minutes for Setaria viridis, and 6 minutes for rice. Subsequent procedures were carried out according to the STOmics Gene Expression Kit-S1 instructions (BGI, No. 1000028493 for library construction and sequencing via the MGI DNBSEQ platform in CNGB.

Raw Stereo-seq data processing

Sequenced reads were generated using an MGI DNBSEQ-T10 sequencer and the expression profile matrices with coordinate on chips were produced according to the Stereo-seq method.[28] The reads of CID and MID were used to label cells and locate coordinates, respectively. The remaining transcript reads were then aligned to genome references (maize, Zm-B73-REFERENCE-NAM-5.0.55 from maizeGDB; sorghum, *Sorghum bicolor* v3.1.1 from Phytozome; Setaria, *Setaria viridis* v2.1 from Phytozome; rice, *Oryza sativa* v7.0 released by MSU) to generate gene expression raw matrix.

650   Image Registration and Cell Identification

The raw matrix was used to generate gene expression maps on chip, which includes

expression counts of each gene on individual DNBs. The SS image was registered to the

gene expression maps based on the track lines on chip with an average error less than

0.5 um, and then overlaid with the FB image to align the tissue contours. Cell

segmentation and identification were conducted based on FB image using the Cellpose

2.0 segmentation algorithm.[50] A custom model was trained using a Nvidia RTX3080Ti

GPU based on the cyto2 initial model. After 12 iterations supervised by

Human-in-the-Loop annotations, the model achieved a recognition accuracy exceeding

95%. The identified cell outlines were converted into grayscale bitmaps. These bitmaps

were overlaid with the original microscopic images using Fiji ImageJ. Cells were manually

classified based on anatomical structures and marked with different colors. Masks

corresponding to different types of cells were exported along with their coordinates. A

Python script was employed to extract single-cell stereo-seq transcriptomes based on

image-defined boundaries for downstream analysis.

665   Unsupervised clustering of Stereo-seq transcriptome and annotation

The MID counts of each gene captured by DNBs in cell segmentation were extracted

based on the masks of easch tissue on chip and summarized to generate gene expression

matrix for downstream analysis. Then the data of cell-gene matrix was processed in Suerat

(v4.4.0).[51] Cells with MID counts less than 10 were discarded and raw data of counts were

normalized by SCTransform function. A KNN graph was constructed by FindNeighbors

function with 30 dimensionalities of PCA and cluster analysis was performed by FindClusters

function with resolution of 1. Clustering results were displayed by uniform manifold

approximation and projection (UMAP) dimension reduction analysis. Established marker

genes were then utilized to annotate the cell types of clusters. For more accurate cell

annotation, the labeled parts of cells were directly recognized based on the relative positions in

the section of leaves. These cell identities were then added into the metadata of clustering

data in Seurat (v4.4.0).

678   Integrate transcriptome of same stage Kranz anatomy into Pseudo-Kranz anatomy

The points of center and right-middle of each Kranz anatomy were labeled to establish the

middle transverse line with symmetry on both adaxial and abaxial sides. Then rotated the

Kranz anatomy to same horizontal line and to integrated same stage Kranz anatomy into

pseudo-Kranz by converting the center points of each Kranz anatomy to origin of coordinates

(0, 0). The expression patterns of individual genes or total of several genes are showed as

contour lines by plotting the MID counts on the pseudo-Kranz.

685   Preparation of nucleus suspension

The protocols were modified based on former works.[52] Tissues were cut into small pieces
in pre-cooled buffer B (0.4 M sucrose, 10 mM $MgCl_2$, 25 mM Tris-HCl pH = 8.0, 1% RNase
inhibitor, 0.1 mM PMSF, 1% Triton X-100, 0.1% DTT) and then incubated on ice for 10 minutes
and filtered. with addition of 0.05% TritonX-100 for maize and 0.1% TritonX-100 for sorghum,
Setaria and rice.

For maize, the nuclei were centrifuged for 10 minutes (1000g, 4°C, 5g/s) and
resuspended in Buffer B. A 5-fold volume of Buffer F (1X PBS, 1% RNase inhibitor and 0.1%
DTT (BGI, 01B022BM)) was added and centrifuged for 10 minutes (1000 g, 4°C, 5 g/s) to
remove most organelles and other molecular impurities. For Sorghum, Setaria, and rice, large
cell debris was centrifuged down to the bottom at 200g for 5 minutes (4°C), and the nuclei in
the supernatant were centrifuged at 1500 g for 20 minutes (4°C) and resuspended in Buffer B
again.   Density gradient centrifugation (3000g, 4°C, 30 minutes) was performed using 30%
percoll (Biosharp, BS909) and 80% percoll to remove organelle and nuclei included in the
middle layer was washed by adding 5 times the volume of Buffer F. The precipitate was
resuspended with cell resuspension solution to obtain the nuclei resuspension for downstream
microfluidic experiments. 2μl cell nuclei resuspension and 8μl DAPI (Beyotime, C1005)
solution was mixed well and counted under fluorescence microscope. Only intact nuclei were
counted.

scRNA-seq and obtaining gene expression matrices

After obtaining the nuclear suspensions, microfluidic experiment and library preparation
were conducted according to the DNBelab C Series Single-Cell Library Prep Set (MGI, No.
1000021082) as previously described.[53] Both library construction and sequencing were
performed on the MGI DNBSEQ-T1&T5 platforms. The obtained raw data was first analyzed
using single-cell DNBelab C4 technology (v2.0)[54] of MGI to automatically obtain two gene
matrices: RawMatrix including reads of all DNBs and FilterMatrix including reads of filtered
DNBs. The genome references were same as that used at analyses of Stereo-seq. The
RawMatrix and FilterMatrix were further processed using the SoupX algorithm to remove
background contamination to generate final expression matrix for downstream analysis.[55]

Quality control and unsupervised clustering of scRNA-seq data

Cell-gene expression matrices were processed and unsupervised clustering using Seurat
(v4.4.0).[51] For filtering high-quality data of cells, genes expressed in less than 3 cells and cells
with less than 2000 in maize, 1000 in sorghum and 500 in Setaria and rice were removed. The
data from different matrices of same species were merged and then normalized using the
NormalizeData function with the "LogNormalize" method (scale.factor = 10000). Cell cycle
phase were predicted Using the CellCycleScoring function in order to mitigate adverse impact
on downstream analysis of clustering.[56] PCA dimensionality reduction was performed and

calculate top 30 principal components using RunPCA function with 3000 variable genes
identified by FindVariableFeatures function. Batch effects were removed using RunHarmony
function. Unsupervised clustering was conducted utilizing the FindNeighbors and FindClusters
functions with a resolution parameter set to 1, and the results were visualized using Uniform
Manifold Approximation and Projection (UMAP).

Integrate scRNA-seq transcriptome with stereo-seq atlas

We integrated the scRNA-seq transcriptome with the stereo-seq atlas using the
TransferData function in Seurat (v4.4.0).[51] Prior to integration, both datasets were normalized
using the SCTransform function. Anchors were calculated using the FindTransferAnchors
function, and predictions.assay of integration were performed using TransferData function and
added to the metadata of the spatial atlas. The correlation of clusters with cells from spatial
data was plotted on the spatial atlas using the SpatialFeaturePlot function.

Cell type annotation in scRNA-seq atlas

We firstly collected known markers that are preferentially expressed in specific cell types
either in maize and rice or other species and obtained their orthologs in maize, sorghum,
Setaria and rice. Most of cell types occurring in the four species were identified using known
markers (Table S12) and were further confirmed by the results of integration of scRNA-seq and
spatial transcriptome. For the few clusters without well-known markers, we primarily relied on
the integration of scRNA-seq and spatial transcriptomics to infer cell types. Additionally, we
considered the orthologs of DEGs that are specifically expressed in these clusters and are
conserved across different species.

Subset of single cells at early developmental stages and supervised clustering

The clusters of early stages of cells in scRNA-seq transcriptome were subset based on
the annotation of the atlas and re-clustered as the method of clustering for atlas, except the
parameter of resolution in FindClusters functions were set to 0.5. For supervised clustering to
distinguish the early bundle sheath cells (BSC) of scRNA-seq data in maize, the PCA were
conducted using RunPCA function with variable gene set that consist of 1, 478 DEGs identified
in spatial transcriptome and 2, 000 variable genes identified by FindVariableFeatures function.
The following steps of dimension reduction and clustering were performed as the method of
clustering for atlas.

Identifying conserved DEGs specific in C_4 spieces or in both C_3 and C_4 spieces

Firstly, we identified differentially expressed genes (DEGs) for each cluster across using
the FindMarkers function in Seurat (v4.4.0).[51] Subsequently, we derived the conserved marker
gene sets for each cell type across the three C_4 species by intersecting the orthologous genes
among the DEGs identified in this cell type. We then identified cell type markers that are

specific to C_4 species or shared between C_3 and C_4 plants by comparing the conserved marker
gene sets of each cell type in C_4 species with those of C_3 rice.

Developmental trajectory analysis

The developmental trajectory analyses were conducted by employing implemented
function in Monocle 2 (v2.18.0)[57] or Monocle 3 (v1.0.0)[58] for different samples. For the entire
cell population of the early stage of leaf subset, Monocle3 was employed to delineate the
developmental trajectories of different cell types. Meanwhile, for individual cell types,
Monocle 2 was utilized to primarily compute the pseudo-time of each cell along its
developmental path. The transcriptomes of the subset cells were analyzed using Seurat
(v4.4.0) for dimensionality reduction and subsequent clustering. For Monocle 2, a matrix of
gene expression counts was utilized to create a new CellDataSet object using the
newCellDataSet function, with the expressionFamily set to negbinomial.size(). Subsequently,
size factors and dispersions were estimated using the estimateSizeFactors and
estimateDispersions functions, respectively. Genes for reducing data dimensionality and
ordering cells were identified using the differentialGeneTest function, based on clusters
established by Seurat (v4.4.0). The genes were filtered using the parameters q value < 0.01
and mean_expression >= 0.1. Cell order was conducted using orderCells with default
parameters. Genes that Change as a Function of Pseudotime were found using
differentialGeneTest function. For Monocle 3, the gene expression matrix was normalized
using the preprocess_cds and align_cds functions. Dimensionality reduction and clustering
were performed using the built-in functions of Monocle 3. The developmental trajectory was
reconstructed with the learn_graph function, and cells were subsequently ordered in
pseudo-time using the order_cells function.

Construct co-expression regulatory network

To investigate the transcriptomic features of the bundle sheath and mesophyll cell
arrangements in C_4 plants, the Python package 'Hotspot' was employed for functional gene
module analysis.[35] Initially, single-cell data were filtered to retain the top 5000 most highly
variable genes, along with genes of interest (some genes of interest may not be among the
highly variable genes and thus need to be manually added to a gene list to prevent their
exclusion). The 'danb' model was utilized, and modules were identified based on an FDR
threshold of <0.00001 and genes of interest. The create_modules function was then used to
recognize modules with a min_gene_threshold of 40 and an fdr_threshold of 0.05. Module
scores were calculated using the module_scores function to measure the gene expression
levels in each of cells. Subsequently, a co-expression network between different genes was
constructed using autocorrelation z-scores and visualized in Cytoscape. The regulatory
relationships of TFs with targets were predicted based on two databases of TF binding sites
(JASPAR, https://jaspar.elixir.no/; CIS-BP, http://cisbp2.ccbr.utoronto.ca/index.php). Then the

regulatory relationships of TF and target with predicted binding sites were added into the
co-expression relationship network to construct GRN.

EMS mutants genotyping

T3 generation seeds of B73 maize background, treated with EMS mutagenesis, were
ordered from the mutant library available at http://maizeems.qlnu.edu.cn. After sowing, leaf
tissue was collected from the seedlings on the fifth day for DNA extraction (TIANGEN
DNAquick Plant System, No. DP321-03) followed by PCR-based genotyping (2×Taq Plus
Master Mix II, Vazyme, No. P213-03) to identify mutant genotypes and select homozygous
lines.

RT-PCR quantification analysis

Plant tissues were sampled and ground after freezing in liquid nitrogen. RNA was
extracted using the GeneJET Plant RNA Purification Kit (Thermo Fisher Scientific, No. K0802).
The extracted RNA was reverse transcribed to generate cDNA using the TransScript®
All-in-One First-Strand cDNA Synthesis SuperMix for qPCR kit (TransGen Biotech, No.
AT341-01), with 1000 ng of cDNA synthesized per sample. Diluted cDNA samples were
subjected to real-time quantitative PCR using the PerfectStartTM Fast Green qPCR SuperMix
(TransGen Biotech, No. AQ611-04), with the HSP-96 well plate (Bio-Rad, No. HSP9601) and
the CFX Connect Real-Time system (Bio-Rad). All quantifications were normalized to maize
β-Tubulin1 (Zm00001eb000490) as an internal reference. Data analysis was performed using
Bio-Rad CFX Maestro software (Version: 4.0.2325.0418). Each quantitative experiment was
conducted with three biological replicates, each containing three technical replicates.
Normalized expression analysis (ΔΔCq) was performed based on biological replicates, with
significance determined using a t-test with a p-value < 0.05.

Resin section and microscopy imaging

Plant tissues were sampled and immersed in FAA solution (a solution containing 50%
ethanol, 5% acetic acid, and 3.7% formaldehyde in water) and subjected to vacuum infiltration
at 0.4 MPa on ice to remove bubbles. Subsequently, the plant samples were dehydrated in a
gradient ethanol series and embedded using the Technovit 7100 Embedding Kit (Heraeus
Kulzer, No. HCECS000677). After solidification, 8 mm thick sections were cut using a Leica
UC7 microtome, and the sections were floated onto adhesive-coated slides. Cell wall
autofluorescence was observed using an Olympus fluorescence microscope, and images
were captured.

Reference

1. Zhu, X.-G., Long, S.P., and Ort, D.R. (2010). Improving Photosynthetic Efficiency for Greater Yield. Annu. Rev. Plant Biol. *61*, 235 – 261. https://doi.org/10.1146/annurev-arplant-042809-112206.

2. Sage, R.F. (2016). A portrait of the C_4 photosynthetic family on the 50th anniversary of its discovery: species number, evolutionary lineages, and Hall of Fame. J. Exp. Bot. *67*, 4039–4056. https://doi.org/10.1093/jxb/erw156.

3. Sedelnikova, O.V., Hughes, T.E., and Langdale, J.A. (2018). Understanding the Genetic Basis of C_4 Kranz Anatomy with a View to Engineering C_3 Crops. Annu. Rev. Genet. *52*, 249–270. https://doi.org/10.1146/annurev-genet-120417-031217.

4. Schlüter, U., and Weber, A.P.M. (2020). Regulation and Evolution of C_4 Photosynthesis. Annu. Rev. Plant Biol. *71*, 183 – 215. https://doi.org/10.1146/annurev-arplant-042916-040915.

5. Cribb L, LN, H., and JA, L. (2001). Four mutant alleles elucidate the role of the G2 protein in the development of C_4 and C_3 photosynthesizing maize tissues. Genetics.

6. LN, H., L, R., L, C., and JA, L. (1998). GOLDEN 2: a novel transcriptional regulator of cellular differentiation in the maize leaf. Plant Cell. https://doi.org/10.1105/tpc.10.6.925.

7. Liu, Q., Teng, S., Deng, C., Wu, S., Li, H., Wang, Y., Wu, J., Cui, X., Zhang, Z., Quick, W.P., et al. (2023). SHORT ROOT and INDETERMINATE DOMAIN family members govern *PIN-FORMED* expression to regulate minor vein differentiation in rice. Plant Cell *35*, 2848–2870. https://doi.org/10.1093/plcell/koad125.

850   8.   Hughes, T.E., Sedelnikova, O.V., Wu, H., Becraft, P.W., and Langdale, J.A. (2019).

851   Redundant *SCARECROW* genes pattern distinct cell layers in roots and leaves of

852   maize. Development. https://doi.org/10.1242/dev.177543.

853   9.   Slewinski, T.L., Anderson, A.A., Price, S., Withee, J.R., Gallagher, K., and Turgeon, R.

854   (2014). Short-Root1 Plays a Role in the Development of Vascular Tissue and Kranz

855   Anatomy in Maize Leaves. Mol. Plant *7*, 1388 – 1392.

856   https://doi.org/10.1093/mp/ssu036.

857   10.   Slewinski, T.L., Anderson, A.A., Zhang, C., and Turgeon, R. (2012). *Scarecrow* Plays

858   a Role in Establishing Kranz Anatomy in Maize Leaves. Plant Cell Physiol. *53*, 2030–

859   2037. https://doi.org/10.1093/pcp/pcs147.

860   11.   Langdale, J.A., and Kidner, C.A. (1994). *bundle sheath defective*, a mutation that

861   disrupts cellular differentiation in maize leaves. Development *120*, 673 – 681.

862   https://doi.org/10.1242/dev.120.3.673.

863   12.   Hughes, T.E., and Langdale, J.A. (2022). *SCARECROW* is deployed in distinct

864   contexts during rice and maize leaf development. Development *149*.

865   https://doi.org/10.1242/dev.200410.

866   13.   Wang, W., Ma, J., Liu, H., Wang, Z., Nan, R., Zhong, T., Sun, M., Wang, S., Yao, Y.,

867   Sun, F., et al. (2024). Genome-wide analysis of the switchgrass YABBY family and

868   functional characterization of *PvYABBY14* in response to ABA and GA stress in

869   Arabidopsis. BMC Plant Biol. *24*. https://doi.org/10.1186/s12870-024-04781-7.

870   14.   Hake, S., Hughes, T.E., Sedelnikova, O., Thomas, M., and Langdale, J.A. (2023).

871   Mutations in *NAKED-ENDOSPERM IDD* genes reveal functional interactions with

SCARECROW during leaf patterning in C_4 grasses. PLoS Genet. *19*.

https://doi.org/10.1371/journal.pgen.1010715.

15.   Wang, L., Czedik-Eysenberg, A., Mertz, R.A., Si, Y., Tohge, T., Nunes-Nesi, A.,

Arrivault, S., Dedow, L.K., Bryant, D.W., Zhou, W., et al. (2014). Comparative

analyses of C_4 and C_3 photosynthesis in developing leaves of maize and rice. Nat.

Biotechnol. *32*, 1158–1165. https://doi.org/10.1038/nbt.3019.

16.   Tausta, S.L., Li, P., Si, Y., Gandotra, N., Liu, P., Sun, Q., Brutnell, T.P., and Nelson, T.

(2014). Developmental dynamics of Kranz cell transcriptional specificity in maize leaf

reveals early onset of C_4-related processes. J. Exp. Bot. *65*, 3543 – 3555.

https://doi.org/10.1093/jxb/eru152.

17.   Fouracre, J.P., Ando, S., and Langdale, J.A. (2014). Cracking the Kranz enigma with

systems biology. J. Exp. Bot. *65*, 3327–3339. https://doi.org/10.1093/jxb/eru015.

18.   Wang, P., Kelly, S., Fouracre, J.P., and Langdale, J.A. (2013). Genome-wide

transcript analysis of early maize leaf development reveals gene cohorts associated

with the differentiation of C_4 Kranz anatomy. Plant J. *75*, 656 – 670.

https://doi.org/10.1111/tpj.12229.

19.   Chang, Y.-M., Liu, W.-Y., Shih, A.C.-C., Shen, M.-N., Lu, C.-H., Lu, M.-Y.J., Yang,

H.-W., Wang, T.-Y., Chen, S.C.C., Chen, S.M., et al. (2012). Characterizing regulatory

and functional differentiation between maize mesophyll and bundle sheath cells by

transcriptomic analysis. Plant Physiol. *160*, 165 – 177.

https://doi.org/10.1104/pp.112.203810.

20.   Li, P., Ponnala, L., Gandotra, N., Wang, L., Si, Y., Tausta, S.L., Kebrom, T.H., Provart,

N., Patel, R., Myers, C.R., et al. (2010). The developmental dynamics of the maize leaf

transcriptome. Nat. Genet. *42*, 1060–1067. https://doi.org/10.1038/ng.703.

21.   Liu, W.-Y., Yu, C.-P., Chang, C.-K., Chen, H.-J., Li, M.-Y., Chen, Y.-H., Shiu, S.-H., Ku,

M.S.B., Tu, S.-L., Lu, M.-Y.J., and Li, W.-H. (2022). Regulators of early maize leaf

development inferred from transcriptomes of laser capture microdissection

(LCM)-isolated embryonic leaf cells. Proc. Natl. Acad. Sci. U. S. A. *119*.

https://doi.org/10.1073/pnas.2208795119.

22.   Perico, C., Zaidem, M., Sedelnikova, O., Bhattacharya, S., Korfhage, C., and

Langdale, J.A. (2024). Multiplexed in situ hybridization reveals distinct lineage

identities for major and minor vein initiation during maize leaf development.

Proceedings of the National Academy of Sciences *121*.

https://doi.org/10.1073/pnas.2402514121.

23.   Mendieta, J.P., Tu, X., Jiang, D., Yan, H., Zhang, X., Marand, A.P., Zhong, S., and

Schmitz, R.J. (2024). Investigating the cis-Regulatory Basis of C_3 and C_4

Photosynthesis in Grasses at Single-Cell Resolution. bioRxiv.

https://doi.org/10.1101/2024.01.05.574340.

24.   Camo-Escobar, D., Alcalá-Gutiérrez, C., Palafox-Figueroa, E., Guillotin, B.,

Hernández-Coronado, M., Coyac-Rodríguez, J.L., Cerbantez-Bueno, V.E.,

Vélez-Ramírez, A., de Folter, S., Birnbaum, K.D., and Ortiz-Ramírez, C. (2024). A

common regulatory switch controls a suite of C_4 traits in multiple cell types. bioRxiv.

https://doi.org/10.1101/2023.12.21.572850.

25.   Bawa, G., Liu, Z., Yu, X., Tran, L.-S.P., and Sun, X. (2024). Introducing single cell

stereo-sequencing technology to transform the plant transcriptome landscape. Trends

Plant Sci. *29*, 249–265. https://doi.org/10.1016/j.tplants.2023.10.002.

26. Swift, J., Luginbuehl, L.H., Schreier, T.B., Donald, R.M., Lee, T.A., Nery, J.R., Ecker,

J.R., and Hibberd, J.M. (2023). Single nuclei sequencing reveals C_4 photosynthesis is

based on rewiring of ancestral cell identity networks. bioRxiv.

https://doi.org/10.1101/2023.10.26.562893.

27. Xia, K., Sun, H.-X., Li, J., Li, J., Zhao, Y., Chen, L., Qin, C., Chen, R., Chen, Z., Liu, G.,

et al. (2022). The single-cell stereo-seq reveals region-specific cell subtypes and

transcriptome profiling in *Arabidopsis* leaves. Dev. Cell *57*, 1299–1310.e1294.

https://doi.org/10.1016/j.devcel.2022.04.011.

28. Chen, A., Liao, S., Cheng, M., Ma, K., Wu, L., Lai, Y., Qiu, X., Yang, J., Xu, J., Hao, S.,

et al. (2022). Spatiotemporal transcriptomic atlas of mouse organogenesis using DNA

nanoball-patterned arrays. Cell *185*, 1777 – 1792.e1721.

https://doi.org/10.1016/j.cell.2022.04.003.

29. Bezrutczyk, M., Zöllner, N.R., Kruse, C.P.S., Hartwig, T., Lautwein, T., Köhrer, K.,

Frommer, W.B., and Kim, J.-Y. (2021). Evidence for phloem loading via the abaxial

bundle sheath cells in maize leaves. Plant Cell *33*, 531 – 547.

https://doi.org/10.1093/plcell/koaa055.

30. Wang, P., Khoshravesh, R., Karki, S., Tapia, R., Balahadia, C.P., Bandyopadhyay, A.,

Quick, W.P., Furbank, R., Sage, T.L., and Langdale, J.A. (2017). Re-creation of a Key

Step in the Evolutionary Switch from C_3 to C_4 Leaf Anatomy. Curr. Biol. *27*, 3278–

3287.e3276. https://doi.org/10.1016/j.cub.2017.09.040.

31.   Bosabalidis, A.M., Evert, R.F., and Russin, W.A. (1994). Ontogeny of the vascular

bundles and contiguous tissues in the maize leaf blade. Am. J. Bot. *81*, 745–752.

https://doi.org/10.1002/j.1537-2197.1994.tb15509.x.

32.   Qu, L.-J., Sawchuk, M.G., Edgar, A., and Scarpella, E. (2013). Patterning of Leaf Vein

Networks by Convergent Auxin Transport Pathways. PLoS Genet. *9*.

https://doi.org/10.1371/journal.pgen.1003294.

33.   Carraro, N., Forestan, C., Canova, S., Traas, J., and Varotto, S. (2006). *ZmPIN1a* and

ZmPIN1b encode two novel putative candidates for polar auxin transport and plant

architecture determination of maize. Plant Physiol. *142*, 254 – 264.

https://doi.org/10.1104/pp.106.080119.

34.   Dong, W., Chang, T., Dai, H., Yang, W., Su, Y., Chao, D., Zhu, X.G., Wang, P., Yu, N.,

and Wang, E. (2023). Creating a C(4)-like vein pattern in rice by manipulating SHORT

ROOT and auxin levels. Sci. Bull. *68*, 3133 – 3136.

https://doi.org/10.1016/j.scib.2023.10.005.

35.   DeTomaso, D., and Yosef, N. (2021). Hotspot identifies informative gene modules

across modalities of single-cell genomics. Cell Syst. *12*, 446 – 456.e449.

https://doi.org/10.1016/j.cels.2021.04.005.

36.   Yang, T., He, Y., Niu, S., Yan, S., and Zhang, Y. (2020). Identification and

characterization of the *CONSTANS* (*CO*)/*CONSTANS-like* (*COL*) genes related to

photoperiodic signaling and flowering in tomato. Plant Sci. *301*.

https://doi.org/10.1016/j.plantsci.2020.110653.

37.   Ohmiya, A., Oda-Yamamizo, C., and Kishimoto, S. (2019). Overexpression of

CONSTANS-like 16 enhances chlorophyll accumulation in petunia corollas. Plant Sci.

280, 90–96. https://doi.org/10.1016/j.plantsci.2018.11.013.

38.   Zhang, Z., Ji, R., Li, H., Zhao, T., Liu, J., Lin, C., and Liu, B. (2014). *CONSTANS-LIKE*

7 (*COL7*) Is Involved in Phytochrome B (phyB)-Mediated Light-Quality Regulation of

Auxin Homeostasis. Mol. Plant *7*, 1429–1440. https://doi.org/10.1093/mp/ssu058.

39.   Ishikawa, M., Fujiwara, A., Kosetsu, K., Horiuchi, Y., Kamamoto, N., Umakawa, N.,

Tamada, Y., Zhang, L., Matsushita, K., Palfalvi, G., et al. (2023). GRAS transcription

factors regulate cell division planes in moss overriding the default rule. Proc. Natl.

Acad. Sci. U. S. A. *120*. https://doi.org/10.1073/pnas.2210632120.

40.   Cui, H., Levesque, M.P., Vernoux, T., Jung, J.W., Paquette, A.J., Gallagher, K.L.,

Wang, J.Y., Blilou, I., Scheres, B., and Benfey, P.N. (2007). An evolutionarily

conserved mechanism delimiting *SHR* movement defines a single layer of endodermis

in plants. Science *316*, 421–425. https://doi.org/10.1126/science.1139531.

41.   Guo, J., Zhou, X.-t., Dai, K.-l., Yuan, X.-y., Guo, P.-y., Shi, W.-p., and Zhou, M.-x.

(2022). Comprehensive analysis of *YABBY* gene family in foxtail millet (*Setaria italica*)

and functional characterization of *SiDL*. J. Integr. Agric. *21*, 2876 – 2887.

https://doi.org/10.1016/j.jia.2022.07.052.

42.   Zhao, S.-P., Lu, D., Yu, T.-F., Ji, Y.-J., Zheng, W.-J., Zhang, S.-X., Chai, S.-C., Chen,

Z.-Y., and Cui, X.-Y. (2017). Genome-wide analysis of the *YABBY* family in soybean

and functional identification of *GmYABBY10* involvement in high salt and drought

stresses. Plant Physiol. Biochem. *119*, 132 – 146.

https://doi.org/10.1016/j.plaphy.2017.08.026.

43.   Mun, B.-G., Lee, S.-U., Park, E.-J., Kim, H.-H., Hussain, A., Imran, Q.M., Lee, I.-J.,

983   and Yun, B.-W. (2017). Analysis of transcription factors among differentially expressed

984   genes induced by drought stress in Populus davidiana. 3 Biotech *7*.

985   https://doi.org/10.1007/s13205-017-0858-7.

986   44.   Rahimi, M., Kordrostami, M., Maleki, M., and ModaresKia, M. (2016). Investigating the

987   effect of drought stress on expression of *WRKY1* and *EREBP1* genes and antioxidant

988   enzyme activities in lemon balm (*Melissa Officinalis* L.). 3 Biotech *6*.

989   https://doi.org/10.1007/s13205-016-0416-8.

990   45.   Foulkes, N.S., Marcolino-Gomes, J., Rodrigues, F.A., Oliveira, M.C.N., Farias, J.R.B.,

991   Neumaier, N., Abdelnoor, R.V., Marcelino-Guimarães, F.C., and Nepomuceno, A.L.

992   (2013). Expression Patterns of GmAP2/EREB-Like Transcription Factors Involved in

993   Soybean Responses to Water Deficit. PLoS One *8*.

994   https://doi.org/10.1371/journal.pone.0062294.

995   46.   Yu, Y., Zhang, H., Long, Y., Shu, Y., and Zhai, J. (2022). Plant Public RNA-seq

996   Database: a comprehensive online database for expression analysis of ~45 000 plant

997   public RNA - Seq libraries. Plant Biotechnol. J. *20*, 806 – 808.

998   https://doi.org/10.1111/pbi.13798.

999   47.   Sack, L., and Scoffoni, C. (2013). Leaf venation: structure, function, development,

1000   evolution, ecology and applications in the past, present and future. New Phytol. *198*,

1001   983–1000. https://doi.org/10.1111/nph.12253.

1002   48.   Kadereit, G., Ackerly, D., and Pirie, M.D. (2012). A broader model for C$_4$

1003   photosynthesis evolution in plants inferred from the goosefoot family

(Chenopodiaceae s.s.). Proc. R. Soc. B *279*, 3304 – 3311.

https://doi.org/10.1098/rspb.2012.0440.

49.   Vlad, D., Zaidem, M., Perico, C., Sedelnikova, O., Bhattacharya, S., and Langdale, J.A.

(2024). The WIP6 transcription factor *TOO MANY LATERALS* specifies vein type in

C_4 and C_3 grass leaves. Curr. Biol. *34*, 1670 – 1686.e1610.

https://doi.org/10.1016/j.cub.2024.03.007.

50.   Pachitariu, M., and Stringer, C. (2022). Cellpose 2.0: how to train your own model. Nat.

Methods *19*, 1634–1641. https://doi.org/10.1038/s41592-022-01663-4.

51.   Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., Zheng, S., Butler, A., Lee, M.J.,

Wilk, A.J., Darby, C., Zager, M., et al. (2021). Integrated analysis of multimodal

single-cell data. Cell *184*, 3573 – 3587.e3529.

https://doi.org/10.1016/j.cell.2021.04.048.

52.   Wang, K., Zhao, C., Xiang, S., Duan, K., Chen, X., Guo, X., and Sahu, S.K. (2023). An

optimized FACS-free single-nucleus RNA sequencing (snRNA-seq) method for plant

science research. Plant Sci. *326*. https://doi.org/10.1016/j.plantsci.2022.111535.

53.   Han, L., Wei, X., Liu, C., Volpe, G., Zhuang, Z., Zou, X., Wang, Z., Pan, T., Yuan, Y.,

Zhang, X., et al. (2022). Cell transcriptomic atlas of the non-human primate *Macaca*

fascicularis. Nature *604*, 723–731. https://doi.org/10.1038/s41586-022-04587-3.

54.   Mazid, M.A., Ward, C., Luo, Z., Liu, C., Li, Y., Lai, Y., Wu, L., Li, J., Jia, W., Jiang, Y.,

et al. (2022). Rolling back human pluripotent stem cells to an eight-cell embryo-like

stage. Nature *605*, 315–324. https://doi.org/10.1038/s41586-022-04625-0.

55.   Young, M.D., and Behjati, S. (2020). SoupX removes ambient RNA contamination

1026   from droplet-based single-cell RNA sequencing data. GigaScience *9*.

1027   https://doi.org/10.1093/gigascience/giaa151.

1028   56.   Barron, M., and Li, J. (2016). Identifying and removing the cell-cycle effect from

1029   single-cell RNA-Sequencing data. Sci. Rep. *6*. https://doi.org/10.1038/srep33892.

1030   57.   Qiu, X., Mao, Q., Tang, Y., Wang, L., Chawla, R., Pliner, H.A., and Trapnell, C. (2017).

1031   Reversed graph embedding resolves complex single-cell trajectories. Nat. Methods *14*,

1032   979–982. https://doi.org/10.1038/nmeth.4402.

1033   58.   Cao, J., Spielmann, M., Qiu, X., Huang, X., Ibrahim, D.M., Hill, A.J., Zhang, F.,

1034   Mundlos, S., Christiansen, L., Steemers, F.J., et al. (2019). The single-cell

1035   transcriptional landscape of mammalian organogenesis. Nature *566*, 496 – 502.

1036   https://doi.org/10.1038/s41586-019-0969-x.

1037

1038

Figure 1: Spatial transcriptome of maize leaf primordium

(A) Workflow of Stereo-seq and snRNA-seq. The outlines of cell wall were recognized by
Cellpose 2.0 and later be used to extract spatial single cell transcriptomes. Then the
spatial marker genes were used to assist to distinguish the subtle differences between
similar cell types of scRNA-seq data.

(B) Spatial visualization of clusters on maize leaf primordium with outer leaf sheath.

(C) Digital expression patterns of selected marker genes on Stereo-seq maps. Epidermis:
ZmLTP3; mesophyll: *ZmGLP1*; procambium: *ZmLAX2*; phloem marker: *ZmAPL*; xylem
marker: *ZmXCP1*; bundle sheath: *ZmSSU1.*

(D) Annotated cell types of clusters identified on unsupervised clustering UMAP of maize leaf
primordium with outer leaf sheath.

(E) Recognized more (sub-)cell types based on the cell morphogenesis and spatial distribution
on Stereo-seq maps of maize leaf primordium. mGM: median ground meristem.

1052

Figure 2. Integration of distributed Kranz anatomy into unified pseudo-Kranz.

(A) Representative picture of three stages of Kranz anatomy in maize leaf primordium. Left:
Fluorescence-stained images; Right: Recognized cell types with artificial color labels.

(B) Workflow of integrating distributed Kranz anatomy at same stage into pseudo-Kranz. Left:
Cartoon of spaced distribution of Kranz anatomy in leaf primordium; Middle: representative
orientations of different Kranz anatomy which need to be rotated to horizontal direction;
Right: overlap of different Kranz anatomy into pseudo-Kranz around original point.

(C) Expression patterns of marker genes on pseudo-Kranz. Epidermis: *ZmLTP3*; mesophyll:
ZmGLP1; bundle sheath: *ZmSSU1;* vein: *ZmLAX2.*

(D) Modules with gradient decreasing patterns from center to periphery of pseudo-Kranz in
three stages. One module with this expression pattern is detected either in 4BS-early or
5BS-middle stage and two modules are detected in 5BS-late stage.

(E) Venn diagram of genes highly expressed in centers of pseudo-Kranz.

(F) Representative new genes identified by pseudo-Kranz. Up: Venn diagram of markers
identified in pseudo-Kranz and spatial atlas, respectively; Middle: representative new
markers of pseudo-Kranz and bottom: two marker gene expression patterns on
pseudo-Kranz.

1070

Figure 3: Integration of single-cell and spatial transcriptome attributes to identify early
bundle sheath cell in maize leaf primordium

(A) The annotation of cell clusters on the UMAP of scRNA-seq atlas for maize leaf primordium
with outer leaf sheath.

(B) Supervised clustering of the subset cells at early developmental stages. The old annotation
of the clusters of atlas are presented on the UMAP.

(C) New annotation of sub-clusters on the UMAP.

(D) Pseudo-time trajectory of subset cells displayed on the UMAP.

(E) Dot plot of marker genes in the indicated cell types. The color key from light blue to dark
blue indicates low to high gene expression levels of averaged scaled data. The dot size

indicates the ratio of cells with expression compared to cells in that cluster.

(F) The expression levels of DEGs identified in representative sub-clusters are summarized
and visualized on 4BS-early pseudo-Kranz.

(G) Pseudo-time analysis of early- and late-stage of BSC. Pseudo-time trajectory (up) and
variable genes dynamically changed along the pseudo-time (pseudo-time related genes,
bottom).

(H) The expression heatmap of representative pseudo-time related genes and their enriched
GO terms.

(I) Dot plot of *SHR* and *SCR* genes in sub-clusters as showed in (E). IE-MC, inner early MC;
OE-MC, outer early MC; L-MC, late MC. E-BSC, early BSC; L-BSC, late BSC; E-vein,
early vein. The color key from light blue to dark blue indicates low to high gene expression
levels of averaged scaled data. The dot size indicates the ratio of cells with expression
compared to cells in that cluster.

1094

Figure 4. The regulatory network of maize leaf primordium

(A) Heatmap showing expression correlations between genes within 25 co-expression
modules.

(B) Total expression levels displayed on scRNA-seq UMAP and pseudo-Kranz. Left: Modules
scores displayed on the UMAP of subset clusters, which represents the relative expression
levels of each modules in each cells; Right: summarized expression levels of genes in
modules showed on pseudo-Kranz.

(C) The average expression of genes within vein-related modules on spatial atlas. The size of
each spot represents the average scaled expression levels of genes within one module,
with different colors indicating six distinct modules. Zoomed-in images on the right side
showcase representative expression patterns of these modules.

(D) Pseudo-time trajectory of sub-clusters of veins. Up: the pseudo-time of each cells in
trajectory; Bottom: cell types of sub-clusters showed on the trajectory.

(E) Module scores of each cell are showed along with the pseudo-time.

(F) The variable marker genes dynamically along with the pseudo-time of vein were divided
into five clades. Left: The numbers in box indicates the number of intersect genes with
different vein-associated modules; Right: Representative TFs in the five clades of
pseudo-time correlated genes.

(G) Regulatory network in maize leaf primordium. Colors of nodes denote distinct modules.
The size of nodes reflects the aggregate correlation score. Solid lines indicate predicted
regulatory interactions between TFs, while dashed lines represent potential relationships
without predicted binding sites. The "Zm" characters in front of the gene names have been
omitted.

1118

Figure 5. Comparison between scRNA-seq atlases of grass leaves

(A-C) The annotation of cell clusters on the UMAP of scRNA-seq atlases of leaf primordium
with outer leaf sheath for sorghum (A), *Setaria* (B) and rice (C).

(D) The expression patterns of conserved marker genes in common cell types identified in
scRNA-seq atlases of the four grass species. The color key (red for maize, blue for
sorghum, brown for Setaria and green for rice) from light to dark indicates low to high gene

expression levels of average scaled data. The dot size indicates the ratio of cells with

expression compared to cells in that cluster. The column at the bottom right shows the

percentage of each cell type.

(E) Comparison of DEGs within procambium between C_4 and C_3 species. Up: Venn diagrams

showing the numbers of DEGs in different species; Bottom: C_4 specific or C_4 and C_3

shared representative markers are listed in table.

(F) The relative expression levels of *SHR* and *SCR* genes in G2M-stage cells and procambium

identified in the four species.

1133

Figure 6. conserved gene regulatory network of Kranz anatomy.

(A) Conserved co-expression network in grass species. The size of nodes reflects the mean

correlation scores of four species. Orange lines represent specific conserved

co-expression relationship in C_4 species and gray lines represent the conserved

co-expression relationship both in C_4 and C_3. Solid lines indicate predicted regulatory

interactions between TFs, while dashed lines represent potential relationships without

predicted binding sites. Blue nodes represents *SHR* genes and their TFs, while green

nodes represents *SCR* genes and their TFs.

(B) The transverse section of maize wild-type (WT, B73), *idd7* EMS mutant and *iddp1* EMS

mutant. Red arrows indicate the loss of mesophyll cells or fusion of lateral veins in EMS

mutants. Scale bar: 100μm.

(C) Histograms summarizing the mean number of mesophyll cells separating veins in WT

maize, *idd7* EMS mutants and *iddp1* EMS mutants, each for three replicates (N=3).

(D) RT-PCR quantification of the expression of *SCR1* and *SCR2* in WT maize , *idd7* EMS

mutants and *iddp1* EMS mutants. (* : $P<0.05$, two-tail Student's t-test)

1149

1150

Figure 1: Spatial transcriptome of maize leaf primordium

(A) Workflow of Stereo-seq and snRNA-seq. The outlines of cell wall were recognized by Cellpose 2.0 and later be used to extract spatial single cell transcriptomes. Then the spatial marker genes were used to assist to distinguish the subtle differences between similar cell types of scRNA-seq data.

(B) Spatial visualization of clusters on maize leaf primordium with outer leaf sheath.

(C) Digital expression patterns of selected marker genes on Stereo-seq maps. Epidermis: *ZmLTP3*; mesophyll: *ZmGLP1*; procambium: *ZmLAX2*; phloem marker: *ZmAPL*; xylem marker: *ZmXCP1*; bundle sheath: *ZmSSU1*.

(D) Annotated cell types of clusters identified on unsupervised clustering UMAP of maize leaf primordium with outer leaf sheath.

(E) Recognized more (sub-)cell types based on the cell morphogenesis and spatial distribution on Stereo-seq maps of maize leaf primordium. mGM: median ground meristem.

Figure 2. Integration of distributed Kranz anatomy into unified pseudo-Kranz.
(A) Representative picture of three stages of Kranz anatomy in maize leaf primordium. Left: Fluorescence-stained images; Right: Recognized cell types with artificial color labels.
(B) Workflow of integrating distributed Kranz anatomy at same stage into pseudo-Kranz. Left: Cartoon of spaced distribution of Kranz anatomy in leaf primordium; Middle: representative orientations of different Kranz anatomy which need to be rotated to horizontal direction; Right: overlap of different Kranz anatomy into pseudo-Kranz around original point.
(C) Expression patterns of marker genes on pseudo-Kranz. Epidermis: *ZmLTP3*; mesophyll: *ZmGLP1*; bundle sheath: *ZmSSU1;* vein: *ZmLAX2*.
(D) Modules with gradient decreasing patterns from center to periphery of pseudo-Kranz in three stages. One module with this expression pattern is detected either in 4BS-early or 5BS-middle stage and two modules are detected in 5BS-late stage.
(E) Venn diagram of genes highly expressed in centers of pseudo-Kranz.
(F) Representative new genes identified by pseudo-Kranz. Up: Venn diagram of markers identified in pseudo-Kranz and spatial atlas, respectively; Middle: representative new markers of pseudo-Kranz and bottom: two marker gene expression patterns on pseudo-Kranz.

Figure 3: Integration of single-cell and spatial transcriptome attributes to identify early bundle sheath cell in maize leaf primordium

(A) The annotation of cell clusters on the UMAP of scRNA-seq atlas for maize leaf primordium with outer leaf sheath.

(B) Supervised clustering of the subset cells at early developmental stages. The old annotation of the clusters of atlas are presented on the UMAP.

(C) New annotation of sub-clusters on the UMAP.

(D) Pseudo-time trajectory of subset cells displayed on the UMAP.

(E) Dot plot of marker genes in the indicated cell types. The color key from light blue to dark blue indicates low to high gene expression levels of averaged scaled data. The dot size indicates the ratio of cells with expression compared to cells in that cluster.

(F) The expression levels of DEGs identified in representative sub-clusters are summarized and visualized on 4BS-early pseudo-Kranz.

(G) Pseudo-time analysis of early- and late-stage of BSC. Pseudo-time trajectory (up) and variable genes dynamically changed along the pseudo-time (pseudo-time related genes, bottom).

(H) The expression heatmap of representative pseudo-time related genes and their enriched GO terms.

(I) Dot plot of *SHR* and *SCR* genes in sub-clusters as showed in (E). IE-MC, inner early MC; OE-MC, outer early MC; L-MC, late MC. E-BSC, early BSC; L-BSC, late BSC; E-vein, early vein. The color key from light blue to dark blue indicates low to high gene expression levels of averaged scaled data. The dot size indicates the ratio of cells with expression compared to cells in that cluster.

Figure 4. The regulatory network of maize leaf primordium

(A) Heatmap showing expression correlations between genes within 25 co-expression modules.

(B) Total expression levels displayed on scRNA-seq UMAP and pseudo-Kranz. Left: Modules scores displayed on the UMAP of subset clusters, which represents the relative expression levels of each modules in each cells; Right: summarized expression levels of genes in modules showed on pseudo-Kranz.

(C) The average expression of genes within vein-related modules on spatial atlas. The size of each spot represents the average scaled expression levels of genes within one module, with different colors indicating six distinct modules. Zoomed-in images on the right side showcase representative expression patterns of these modules.

(D) Pseudo-time trajectory of sub-clusters of veins. Up: the pseudo-time of each cells in trajectory; Bottom: cell types of sub-clusters showed on the trajectory.

(E) Module scores of each cell are showed along with the pseudo-time.

(F) The variable marker genes dynamically along with the pseudo-time of vein were divided into five clades. Left: The numbers in box indicates the number of intersect genes with different vein-associated modules; Right: Representative TFs in the five clades of pseudo-time correlated genes.

(G) Regulatory network in maize leaf primordium. Colors of nodes denote distinct modules. The size of nodes reflects the aggregate correlation score. Solid lines indicate predicted regulatory interactions between TFs, while dashed lines represent potential relationships without predicted binding sites. The "Zm" characters in front of the gene names have been omitted.

Figure 5. Comparison between scRNA-seq atlases of grass leaves

(A-C) The annotation of cell clusters on the UMAP of scRNA-seq atlases of leaf primordium with outer leaf sheath for sorghum (A), *Setaria* (B) and rice (C).

(D) The expression patterns of conserved marker genes in common cell types identified in scRNA-seq atlases of the four grass species. The color key (red for maize, blue for sorghum, brown for Setaria and green for rice) from light to dark indicates low to high gene expression levels of average scaled data. The dot size indicates the ratio of cells with expression compared to cells in that cluster. The column at the bottom right shows the percentage of each cell type.

(E) Comparison of DEGs within procambium between C_4 and C_3 species. Up: Venn diagrams showing the numbers of DEGs in different species; Bottom: C_4 specific or C_4 and C_3 shared representative markers are listed in table.

(F) The relative expression levels of *SHR* and *SCR* genes in G2M-stage cells and procambium identified in the four species.

Figure 6. conserved gene regulatory network of Kranz anatomy.

(A) Conserved co-expression network in grass species. The size of nodes reflects the mean correlation scores of four species. Orange lines represent specific conserved co-expression relationship in C_4 species and gray lines represent the conserved co-expression relationship both in C_4 and C_3. Solid lines indicate predicted regulatory interactions between TFs, while dashed lines represent potential relationships without predicted binding sites. Blue nodes represents *SHR* genes and their TFs, while green nodes represents *SCR* genes and their TFs.

(B) The transverse section of maize wild-type (WT, B73), *idd7* EMS mutant and *iddp1* EMS mutant. Red arrows indicate the loss of mesophyll cells or fusion of lateral veins in EMS mutants. Scale bar: 100μm.

(C) Histograms summarizing the mean number of mesophyll cells separating veins in WT maize, *idd7* EMS mutants and *iddp1* EMS mutants, each for three replicates (N=3).

(D) RT-PCR quantification of the expression of *SCR1* and *SCR2* in WT maize , *idd7* EMS mutants and *iddp1* EMS mutants. (*: P<0.05, two-tail Student's t-test)

Supplementary figure 1. **Spatial transcriptome of maize leaf primordium**

(A)   Violin plot showing the number of genes detected in each cell across different clusters in the spatial atlas.

(B)   Cell annotation on the spatial atlas of maize leaf primordium including the outer leaf sheath.

(C)   Cells were separately displayed on the Stereo-seq maps.

(D)   Dot plot of marker genes in the indicated cell types. The color key from light blue to dark blue indicates low to high gene expression levels of averaged scaled data. The dot size indicates the ratio of cells with expression compared to cells in that cluster.

Epidermis
Side mesophyll
Rank1 vein
BS around rank1 vein

Rank2 vein
BS around rank2 vein
Two-cell mesophyll   One-cell mesophyll

Non-differentiated cell

mGM/meristem

rank2 vein   4BS-early   5BS-middle   5BS-late

rank1 vein   main vein   parenchyma cell

Main vein
BS around main vein

sclerenchyma cell

Supplementary figure 2. Representative picture of cell type recognized in spatial transcriptome

Left: Fluorescence-stained images; Right: Recognized cell types with artificial color labels.

Supplementary figure 3. Expression patterns of marker genes identified in each cell type on Stereo-seq maps

Supplementary figure 4. Expression patterns of photosystem related genes in three sub-types of mesophyll cell

Total expression of photosynthetic genes in in three sub-types of mesophyll cell.

A

Supplementary figure 5. Expression patterns of marker genes of four cell types on three stages of pseudo-Kranz

Supplementary figure 6. Dot plot of marker genes in the indicated cell types of maize leaf atlas

The color key from light blue to dark blue indicates low to high gene expression levels of averaged scaled data.

The dot size indicates the ratio of cells with expression compared to cells in that cluster.

Supplementary figure 7. The integration of clusters of snRNA-seq atlas with cells of spatial transcriptomes of maize

The color key from blue to red indicates low to high correlation of transcriptome between the cluster and each cell of spatial atlas.

Supplementary figure 8. The integration of sub-clusters of snRNA data with cells of spatial transcriptomes

The color key from blue to red indicates low to high correlation of transcriptome between the cluster and each cell of spatial atlas.

Legend:
- ZmSHR
- ZmSCR
- SHR-SCR

Supplementary figure 9. co-expression patterns of *SHR* and *SCR* genes in spatial transcriptomes

The size of the dots on the plot indicates the average expression levels of different genes. Light blue indicates the average expression of *ZmSHR1* and *ZmSHR2* genes only, green indicates the average expression of *ZmSCR1* and *ZmSCR2* genes only, and pink indicates the average expression of co-expressed *ZmSHR* and *ZmSCR* genes.

Supplementary figure 10. Expression patterns of module genes in snRNA-seq and pseudo-Kranz.

Left : Modules scores of each cells displayed on the UMAP of subset clusters. Middle: Total expressions of gene in modules showed on pseudo-Kranz. Right: The heatmap showing the relative scaled expression of genes within each module at different distances from the center of pseudo-Kranz at three stages.

Supplementary figure 11. Expression patterns of individual vein-associated module genes spatial atlas.

Supplementary figure 12. Regulatory co-expression network of vein related module genes.
Colors of nodes denote distinct modules. The size of nodes reflects the aggregate correlation score. Solid lines indicate predicted regulatory interactions between TFs, while dashed lines represent potential but unconfirmed relationships without predicted binding sites. Circles and diamonds of spots represent functional genes and TFs, respectively.

Supplementary figure 13. The expression patterns of candidate regulators of *SHR*, *SCR* and *G2* genes

The color key from light blue to dark blue indicates low to high gene expression levels of averaged scaled data.

The dot size indicates the ratio of cells with expression compared to cells in that cluster.

Supplementary figure 14. Dot plot of marker genes in the indicated cell types of Sorghum leaf atlas

The color key from light blue to dark blue indicates low to high gene expression levels of averaged scaled data.

The dot size indicates the ratio of cells with expression compared to cells in that cluster.

Supplementary figure 15. Dot plot of marker genes in the indicated cell types of Setaria leaf atlas

The color key from light blue to dark blue indicates low to high gene expression levels of averaged scaled data.

The dot size indicates the ratio of cells with expression compared to cells in that cluster.

Supplementary figure 16. Dot plot of marker genes in the indicated cell types of rice leaf atlas

The color key from light blue to dark blue indicates low to high gene expression levels of averaged scaled data.

The dot size indicates the ratio of cells with expression compared to cells in that cluster.

DEG of MC&BS	
C3_C4 (106)	*G2, IDDP1, IDD7, NKD1, TCP23, ARF25, LHCA4,*
C4 (68)	*SCR1, SCR2, bHLH127, IAA5*

Supplementary figure 17. Conserved DEGs in BSC and/or MC among C$_4$ and C$_3$ plants.

(A) Comparison of DEGs in clusters of MC&BSC between C$_4$ and C$_3$ species. Up: Venn diagrams showing the numbers of DEGs in different species; Bottom: C$_4$ specific or C$_4$ and C$_3$ shared representative markers are listed in table.

(B) The relative expression levels of *SCR* genes in clusters of MC&BSC of four species

Supplementary figure 18. The expression patterns of selected key genes in different cell types of C$_4$ and C$_3$ plants

The color key (red for maize, blue for Sorghum, brown for Setaria and green for rice) from light to dark indicates low to high gene expression levels of averaged scaled data. The dot size indicates the ratio of cells with expression compared to cells in that cluster.

Supplementary figure 19. The expression levels of *SHR* genes in root and leaf of maize and rice

Supplementary figure 20. The expression levels of auxin biosynthesis genes in C$_4$ and C$_3$ species

Supplementary figure 21. The expression levels of auxin signal transduction genes in C$_4$ and C$_3$ species

Supplementary figure 22. The expression levels of auxin transport genes in C$_4$ and C$_3$ species

Supplementary figure 23. The relative positions of predicted binding sites of TFs co-expressed with *SCR1*

The transcription factors binding sites (TFBS) located approximately 3 kb upstream of the coding sequence in maize (A), sorghum (B), Setaria (C), and rice (D). The horizontal lines represent gene lengths, and the boxes on these lines indicate the positions of the two exons within the SCR1 genes. The vertical lines mark the relative positions of TFBS, with the corresponding transcription factor families labeled above each vertical line. IDD binding sites are highlighted in blue.

A *Zmidd7* mutant

B *Zmiddp1* mutant

Supplementary figure 24. The mutation site identification of EMS mutants .

(A) There is a mutation of C to T at the 481st bp among full-length 1605bp coding sequence of *ZmIDD7*, which makes the 161st amino-acid glutamine (Q, CAG) change to a stop codon (TAG) that leads a premature termination on translation of full-length 534 amino-acids.

(B) There is a mutation of C to T at the 448th bp among full-length 2025bp coding sequence of *ZmIDDP1*, which makes the 150th amino-acid glutamine (Q, CAG) change to a stop codon (TAG) that leads a premature termination on translation of full-length 674 amino-acids.