Title

Mapping Visual Contrast Sensitivity and Vision Loss Across the

Visual Field with Model-Based fMRI

Proposed Authorship

Hugo T. Chow-Wing-Bom[a,b], Matteo Lisi[c], Noah C. Benson[d], Freya Lygo-Frett[e], Patrick Yu-

Wai-Man[a,f,g,h], Frederic Dick[b,i], Roni O. Maimon-Mor[a], Tessa M. Dekker[a,b,i]

[a] *Institute of Ophthalmology, University College London (UCL), London, United Kingdom*

[b] *Birkbeck/UCL Centre for NeuroImaging, London, United Kingdom*

[c] *Department of Psychology, Royal Holloway, University of London, London, United Kingdom*

[d] *eScience Institute, University of Washington, Seattle, WA, USA*

[e] *Institute of Cognitive Neuroscience, UCL, London, United Kingdom*

[f] *John van Geest Centre for Brain Repair and MRC Mitochondrial Biology Unit, Department of*

Clinical Neurosciences, University of Cambridge, Cambridge, United Kingdom

[g] *Cambridge Eye Unit, Addenbrooke's Hospital, Cambridge University Hospitals NHS*

Foundation Trust, Cambridge, United Kingdom

[h] *Moorfields Eye Hospital NHS Foundation Trust, London, United Kingdom*

[i] *Department of Experimental Psychology, UCL, London, United Kingdom*

Correspondence

Hugo T. Chow-Wing-Bom; Institute of Ophthalmology, University College London, 11-43 Bath

Street, London EC1V 9EL; hugo.chow-wing-bom.15@ucl.ac.uk

Funding

This work was funded by Moorfields Eye Charity PhD Studentship GR001315 (London, UK),

and the Birkbeck-UCL Centre for NeuroImaging (BUCNI; London, UK). The funders had no

role in study design, data collection and analysis, decision to publish, or preparation of the

manuscript.

Commercial Relationships

None

Abstract

Peripheral vision is crucial for daily activities and quality of life, yet traditional measures
of visual function like visual acuity primarily assess central vision. Visual field tests can
evaluate peripheral vision but require extended focus combined with precise fixation, often
very challenging for patients with severe sight loss. Functional MRI (fMRI) with population
receptive field (pRF) mapping offers a non-invasive way to map scotomas but is limited by its
reliance on single contrast levels and the necessity of accurate fixation.

We developed an fMRI-based approach to measure contrast sensitivity across the
visual field without the need for precise fixation. By combining large-field stimulation with
varying spatial frequencies and contrast levels with either pRF mapping or a retinotopic atlas
based on anatomical landmarks, we modeled contrast sensitivity in the primary visual cortex
(V1) over a large (40 deg) expanse of the visual field. In seven normal-sighted participants,
we characterized differences in V1 cortical sensitivity across eccentricities and visual
quadrants, finding reliable and reproducible patterns of sensitivity differences at individual and
session levels. Additionally, our method effectively visualized cases of simulated and disease-
linked sensitivity loss at the cortical level. Crucially, we demonstrated that these results could
be largely recovered using a structure-based retinotopic atlas, eliminating the need for pRF
mapping and precise fixation - although such an approach reduced sensitivity.

This approach, integrating large-field stimulation with a retinotopic atlas, offers a
promising tool for monitoring vision loss and recovery in patients with various visual
impairments, addressing a significant challenge in current clinical assessments.

Keywords
fMRI; Visual Field; Contrast Sensitivity; Primary Visual cortex

Introduction

At least 2.2 billion people world-wide experience blinding eye diseases that emerge or progress across the lifespan, with 1.1 billion living with vision loss in 2020 (Bourne et al., 2021; World Health Organization, 2019). Accurate characterization of visual function is critical for optimal intervention and patient support, but fraught with challenges. The gold clinical standard measure is visual acuity, the highest spatial resolution that can be discerned. This index mainly relies on a well-functioning fovea, the small retinal area with the highest density of photoreceptors, which processes the central 0-2 degrees around fixation. However, many forms of visual impairment do not affect the fovea alone. The vast majority of our visual input comes from peripheral vision, which provides coarse but invaluable information (Alvarez, 2011; Oliva, 2005) – nearly all daily functions critically depend on intact functionality of the wider visual field, including driving, crossing the street, social function, mobility, and even reading (Lange et al., 2021). Accordingly, impairment of tissue beyond the fovea and its projections drastically affects function and quality of life (Lange et al., 2021; Lisboa et al., 2013; Roh et al., 2018; Subhi et al., 2017).

The impacts of impairments across the wider visual field can be measured with various types of visual field tests, ranging from the finger counting visual confrontation field test to more complicated and/or computerized tests (e.g., standard automatic perimetry, kinetic perimetry, microperimetry). Computerized tests typically involve measuring sensitivity to the luminance contrast of a target relative to a background at different visual field locations while the participant's gaze is fixed on a central point. In some cases (e.g., microperimetry), sensitivity measurements are paired with fundus imaging, offering greater precision in linking visual field functions to specific retinal locations. These tests can provide critical additional insights into function-relevant disease evolution and recovery beyond acuity scores alone, by identifying regions of reduced light sensitivity or blind spots beyond the fovea.

However, visual field testing comes with challenges and limitations, and as a result, the inclusion of visual field measures in sight-rescuing therapy trials is limited. Firstly, it requires prolonged fixation and sustained visual attention. This can be very challenging for patients with severe vision loss, who often struggle to fixate, and strain to detect even high-intensity stimuli. This can lead to long and unpleasant testing sessions with unreliable results. Secondly, as perception is inherently subjective and effortful, patients may vary in their criteria for visual recognition, and in their ability to report visual signals that are weakened or distorted by disease.

Functional MRI (fMRI) has recently been proposed as a promising alternative to
measure visual field loss, as it requires no overt task, and instead measures visual sensitivity
directly from brain responses. Population receptive field (pRF) mapping fMRI can measure
which parts of the cortex respond to which parts of the visual scene. This is achieved by
measuring local changes in cortical blood oxygenation in response to high-contrast stimuli
such as flickering checkerboards that systematically traverse the visual field. These neural
timeseries are then fit with a retinotopic spatial tuning function. The fMRI signal strength in a
cortical region that encodes a specific visual field location provides information about visual
sensitivity in this location (Dumoulin & Wandell, 2008). This approach has recently been
used to accurately recover blind spots in the visual field detected with visual field perimetry
(Pawloff et al., 2023; Ritter et al., 2019). These results in turn suggest that fMRI may be a
useful tool to monitor disease progression and recovery in visually impaired patients.
However, there are also significant challenges associated with this pRF mapping.
Because pRF models require knowledge of where visual stimuli fall on the retina, this
approach requires accurate and sustained fixation. If fixation is poor, the visual field
locations assigned to cortical regions will be displaced, leading to noise or bias in the pRF
maps. Additionally, most fMRI studies measure vision across a small central part of the
visual fields (~10 degrees of eccentricity) which will not capture critical dynamics in the
periphery. Finally, most studies use a single maximum contrast stimulus to assess visual
function. This is suboptimal for at least two reasons (Broderick et al., 2022; Farahbakhsh et
al., 2022; Liu et al., 2006; O'Connell et al., 2016; Ritter et al., 2019): First, it may reduce the
ability to detect subtle but clinically important changes that do not result in complete
sensitivity loss or recovery. Second, with a single contrast stimulus, it becomes more difficult
to determine whether a reduction in fMRI signal across sessions or brain regions reflects
vision-related changes or other factors. Together, these factors currently limit the utility of
neuroimaging-based visual field tests.
In this study we sought to develop fMRI methods for measuring visual field sensitivity
to address these challenges, and to test its utility in artificial and realistic cases of visual field
loss. Rather than presenting a single maximum contrast eccentricity-scaled checkerboard in
a pRF mapping protocol, observers viewed large-field sinusoidal gratings at a range of
contrasts without strict requirement to fixate. We used a large screen set up (40° diameter)
to measure visual function across a large expanse of the visual field. To quantify visual
sensitivity, we fit a simple cortical contrast response model to voxel timeseries evoked by the
gratings, avoiding reliance on arbitrary signal amplitude thresholds. We first tested if this
approach could sensitively and reliably quantify predicted contrast sensitivity differences
across eccentricity and polar angle dimensions of the entire visual field (Figure 1), using a
short, patient-friendly protocol in sighted controls. Critically, to achieve this we linked cortical

contrast sensitivity estimates to visual field locations using retinotopic tuning estimates

obtained in a separate session. However, for patients with severe vision loss it is infeasible

to obtain accurate pRF measures due to poor ability to fixate a central target. We therefore

also evaluated the approach using the structure-based atlas of retinotopic values developed

by Benson et al. (Benson et al., 2014; Benson & Winawer, 2018). This atlas estimates

retinotopic tuning based on the alignment of structural features of cortex (e.g., surface

curvature) to a mean atlas of those features. Once a brain is aligned to the structural atlas, a

similar atlas of retinotopic maps that are defined on the structural atlas are used to estimate

the polar angle and eccentricity maps for the individual subject. To increase accuracy of the

Benson atlas for our large-field stimuli, we subsequently fit these eccentricity estimates to

the Horton and Hoyt (1991) model of cortical magnification in each individual.

We demonstrate that our approach can recover, at a cortical level, known sensitivity

changes across the visual field. V1 neuronal populations encoding the central visual field

show the highest sensitivity to high spatial frequencies, whilst those encoding the periphery

are most sensitive to low spatial frequencies. We also find greater sensitivity in V1 regions

receiving visual inputs from the horizontal compared to vertical quadrants, and from the

lower compared to upper quadrants. Crucially, these effects are detectable at the individual

level and repeatable across visits, suggesting our approach will be suitable for detecting

visual field sensitivity changes over time (e.g., due to learning, ageing, disease progression,

or recovery). Moreover, when using the Benson atlas instead of individual pRF measures to

link neuronal responses to visual field locations, we still detect these effects, although with

slightly reduced sensitivity. We also demonstrate the clinical relevance of this approach by

recovering a simulated scotoma, characterized by a ring of visual field loss around fixation,

as well as visual field loss in a patient with a neurodegenerative disorder causing large areas

of visual field loss. We conclude that this approach, i.e., the combination of cortical contrast

sensitivity testing, large field stimulation, and a large-field calibrated retinotopic atlas, may be

particularly useful for quantifying visual recovery in patients, whether spontaneous or

following gene-therapy interventions, and for linking brain-based activations to behavioral

performance.

A. Across eccentricities **B. Across visual quadrants**

Figure 1. Expected changes in cortical contrast sensitivity in typically sighted controls for two different spatial frequencies (low spatial frequency, 0.3 cycles per degree (cpd), and high spatial frequency, 3.0 cpd), **A:** as a function of eccentricity and **B:** visual field quadrants. With reference to eccentricity, we expect higher sensitivity of V1 neuronal populations to 1) lower spatial frequencies in the periphery, and 2) higher spatial frequencies in the center of the visual field. By comparison, we expect both high and low spatial frequency stimulation to evoke greatest sensitivity in V1 neuronal populations encoding the left and right visual quadrants, less in the lower, and least in the upper quadrant.

Methods

Participants

We collected data from 7 healthy controls (mean±SD: 29.6±4.7yo; 1M). All controls either had normal or corrected to normal vision, with no other ocular pathologies, and were recruited from the local staff and student pool at the University College of London. We also recruited a patient with Leber Hereditary Optic Neuropathy (M, 45.5 years old, mutation: m.11778G>A, disease duration: 36.1 years). All participants signed a written informed consent to participate in the study, and this study was approved by HRA/HCRW and Research Ethics Committee (IRAS 258959; REC 20/YH/0336).

Stimulus & Apparatus

Stimuli were programmed in MATLAB 2020b (MathWorks, Natick, Massachusetts, USA) using the Psychtoolbox-3, and displayed using an EPSON LB-1100U projector (resolution: 1920x1200 pixels, 60Hz, projected area: 43.2x27cm). Because the projector was too bright and to reduce discomfort in the scanner, a circular neutral density filter (ND4 Plus+; Urth, Byron Bay, Australia) was placed in front of the projector to reduce light intensity

by 4 times without altering the colors. The projector was also gamma-corrected to ensure

accurate representation of the intended contrast levels and spatial frequencies used in our

experiments. In all tasks, stimuli were embedded in circular aperture of 42.3x42.3º diameter.

To achieve the large field-of-view, the front part of the 64-channel MRI coil was removed,

and stimuli were back-projected onto a screen positioned close to the participant's head

inside the bore (see Figure A1 in *Appendix* section). Participants viewed the stimuli through

a mirror, for a total viewing distance of 34cm. Throughout the session, participants were

constantly reminded to remain still to minimize head and body motion, and eye motion and

alertness were monitored using an EyeLink 1000Plus device (SR Research, Ottawa,

Canada). In all tasks, participants were asked to report when the color of a central fixation

dot changed.

fMRI Contrast Sensitivity Task

Participants were presented binocularly with achromatic sine-wave gratings,

flickering at a temporal frequency of 2Hz and varying in contrast levels and spatial

frequencies. Contrast levels were defined as Michelson contrast ($C = \frac{L_{max} - L_{min}}{L_{max} + L_{min}}$), where

Lmax and Lmin are maximum and minimum luminance. Each stimulus, presented for 13s,

featured a specific contrast level (either 7.5, 42.2, 60, or 100%) and spatial frequency (either

0.3 or 3 cycles per degree). Stimuli were then followed by 2s of a grey background (0%

contrast) to minimize the experience of after-effects. Each combination of contrast level and

spatial frequency was shown three times within a run, arranged in a pseudo-random

sequence, for a total of 9 repetitions per combination (i.e., 3 runs). Eight additional blocks of

15-second rest periods with a grey background were introduced to allow sufficient baseline.

One healthy control also performed a version of the task in which we masked a region of the

visual field with a grey, annular ring, covering 3º-8º eccentricity to simulate loss of visual

inputs (i.e., artificial scotoma).

Population Receptive Field (pRF) mapping

To investigate how distinct regions of the visual field map onto the cortex, we

collected pRF mapping data using a standard ring-and-wedge stimulus (Dumoulin &

Wandell, 2008). Participants were presented binocularly with a black-and-white

checkerboard, phase-flickering at a temporal frequency of 2Hz, and at maximum contrast

level. The stimulus spanned up to the boundaries of the circular aperture (i.e., 42.3x42.3º

diameter), and was scaled across eccentricity to account for cortical magnification. Three

runs of 352 volumes were collected per participant.

MRI acquisition

Data were collected at the Birkbeck-UCL Centre for NeuroImaging (BUCNI, London, UK), on a Siemens PRISMA 3 Tesla scanner.

Functional data were acquired using a 64-channel head coil with the front part removed, to allow an unobstructed view of the screen whilst ensuing enough signal from the remaining 40 effective channels covering the side and back of the head. T2*-weighted echo-planar imaging were collected with an accelerated multiband sequence kindly provided by CMRR (version R016a , https://www.cmrr.umn.edu/multiband; Cauley et al., 2014; Xu et al., 2013); multi-band factor: 4, voxel resolution: 2mm isotropic, FOV: 212x212x96mm, flip angle: 60º, repetition time (TR): 1000ms, echo time (TE): 35.2ms, echo spacing: 0.56ms, bandwidth: 2620Hz/Px, with 48 transverse slices angled to be approximately parallel to the calcarine sulcus whilst avoiding the orbital cavities). The same sequence was used to acquire 4 additional scans in the opposite phase-encoding direction, to allow B0 deformation correction in the pre-processing steps.

A T1-weigthed structural image was acquired with a 32-channel coil, using MPRAGE sequence (voxel resolution: 1mm isotropic, 176 slices, FOV: 256 x 256 x 176mm, flip angle: 7º, TR = 2.530s, TE = 3.34ms, TI = 1100ms, bandwidth: 200Hz/Px, echo spacing: 7.4ms, acquisition time: 5min30s).

Data analysis

Segmentation of the anatomical scan and reconstruction of cortical surfaces were performed in FreeSurfer 7.1.1.

Functional data were pre-processed using a combination of AFNI 24.1.22, FreeSurfer and FSL 6.0.7.12 commands. The following steps were followed for both the contrast sensitivity task and the pRF mapping task. First, initial correction for distortions caused by field inhomogeneity was performed using a blip-up/blip-down approach (AFNI's unWarpEPI.py program), and the first 4 volumes of each run were discarded using FSL's fslroi program. Then an alignment scan was created from the volume out of all collected runs that contained the least amount of voxel outliers using AFNI's 3dToutcount, 3dTstat and 3dcalc programs. This alignment scan was then co-registered to the MPRAGE image using FreeSurfer's bbregister program, resulting in a rigid-body transformation matrix. In case of misregistration, the alignment scan was defined as one of the single-band reference scans, before re-running the co-registration step. Motion correction was then carried out by aligning each functional volume from all runs to the previously defined alignment scan using AFNI's 3dvolreg program.

Processing of the contrast sensitivity task

To accurately capture neural activity across various eccentricities and polar angle

locations, minimal smoothing (0.5mm FWHM Gaussian blur) was applied to the contrast

sensitivity task data using FSL's 3dmerge program. The contribution of each contrast/spatial

frequency combination to the BOLD signal was modelled using the general linear model

(GLM) approach in SPM12 (https://www.fil.ion.ucl.ac.uk/spm/), with motion parameters as

additional regressors. This step resulted in volumetric images of β-values, each reflecting the

specific contribution of a given contrast/spatial frequency combination to the baseline BOLD

signal. These resulting β-images were then projected onto the individual's cortical surface

using FreeSurfer's mri_vol2surf program, with no surface smoothing and the default

fractional projection along surface of 0.5. The latter sets the fraction (0,1) of the cortical

thickness at each vertex to project along the surface. Surface-projected β-images were

finally converted into a MATLAB-compatible format for further analysis using SamSrf's

samsrf_mgh2srf program.

Processing of pRF mapping task

Pre-processed volumes from the pRF mapping task were projected onto the cortical

surface using FreeSurfer's mri_vol2surf program. Here, we used the previously generated

rigid-body transformation matrix to ensure accurate spatial alignment and smoothed on the

cortical surface with a Gaussian kernel with a FWHM of 3mm.

The resulting outputs were then converted into a MATLAB-compatible format for

further analysis. pRF estimates were computed by fitting a 2D Gaussian model to the data,

using the SamSrf toolbox (version 7.13; https://github.com/samsrf), resulting in three pRF

parameters for each vertex on the cortical surface: the x- and y-coordinates of the pRF

center, and the pRF size (σ). Eccentricity and polar angle maps were computed from the x-

and y-coordinates.

Regions-Of-Interest

In this study, we were interested in how visual information across the visual field is

processed in early visual areas, notably in the primary visual cortex V1. Early visual regions

were manually delineated for each participant in SamSrf v7.13, based on the output of the

pRF fitting procedure and standard functional criteria to identify borders between V1, V2,

and V3 (DeYoe et al., 1996; Engel, 1997; Sereno et al., 1995). In this report, we focused our

interest on V1, given its crucial role in the initial processing of basic visual features like

spatial frequency and contrast.

Quantifying eccentricity- and location-dependent variations in sensitivity

Using subject-specific pRF estimates

At this stage, for each vertex on the cortical surface, we obtained β-values for each

tested contrast/spatial frequency combination in the contrast sensitivity task, as well as pRF

parameters resulting from the fitting of the pRF model to the pRF data (Figure 2). β-values

from the contrast sensitivity task in each vertex were then filtered based on ROI location (V1)

and pRF parameters ($R^2 \geq 0.05$; pRF size: $\sigma \leq 6$).

To assess contrast sensitivity, we modelled the neural response (i.e., β-values) in V1

as a linear function of the square root of the contrast values ($R(C) = a * \sqrt{C}$). This model

rests on the assumption of a monotonic relationship between the presented stimulus

contrast level and the resulting BOLD response and has been used in previous work by

Buracas and colleagues to investigate contrast sensitivity at the cortex level in healthy

controls (Buracas et al., 2005; Buracas & Boynton, 2007). Steeper slopes within this context

indicate greater V1 contrast sensitivity.

We fit this model for each vertex separately, for each participant and spatial

frequency (0.3 and 3cpd) condition. The intercept was constrained to be zero, as the

distribution of β-values is expected to be centered on zero when the contrast is zero (i.e.,

when no stimulus is presented). Next, the resulting V1 slope estimates were binned and

averaged over five regions using pRF eccentricity estimates: 0.5º-2.5º, 2.5º-4.5º, 4.5º-9.5º,

9.5º-15º, and 15º-20º. These bins relate to the clinical definition of visual space based on

anatomical landmarks, corresponding to the fovea, parafovea, perifovea and two near-

periphery regions, respectively. Similarly, slope estimates were spatially constrained within a

±45º-wedge centered on each cardinal meridian to investigate asymmetries as a function of

pRF polar angle locations, thus dividing the visual space between the upper, lower, left and

right quadrants. These wedges were also derived from the polar angle estimates obtained

with pRF mapping.

For each participant, average slopes for each eccentricity bin and polar angle wedge

were then projected back into their corresponding visual field locations. This resulted in heat-

maps of V1 contrast sensitivity across eccentricities and visual field quadrants, for each

tested spatial frequency condition at an individual level (Figure 2). Finally, average slopes for

each eccentricity and polar angle bin from each participant were used to compute the group-

level average.

Figure 2. Estimation of cortical contrast sensitivity across the visual field. Participants performed two fMRI tasks in the scanner: a contrast sensitivity task and a pRF mapping task. For each individual and tested spatial frequency, the contribution of each stimulus contrast level to the BOLD signal in the contrast sensitivity task was estimated using a GLM approach. Resulting β-coefficients for each contrast level were then projected onto the individual's cortical surface. For each vertex in V1, a square root function ($R(C) = a * \sqrt{C}$) was then fitted to these data, taking the estimated slope a as measure of cortical contrast sensitivity. V1 slopes were finally averaged across vertices based on their eccentricity and polar angle preference as estimated with pRF mapping, and across participants. Mean slopes were then projected back into the visual space, producing a heatmap projection of V1 contrast sensitivity across eccentricities and along visual field quadrants.

Using calibrated Benson retinotopic template atlas

The same approach was used to quantify changes in V1 contrast sensitivity across the visual field, using a retinotopic template (Benson et al., 2014; Benson & Winawer, 2018) instead of pRF mapping. This template was fitted to each subject's anatomical MPRAGE image using the Python library neuropythy (Benson & Winawer, 2018), which provided automatic delineation of early visual regions, based on FreeSurfer's anatomical alignment. Additionally, it offered eccentricity and polar angle estimates for each vertex on the cortical surface. The advantage of this method over pRF mapping is that it does not require fixation or pRF scanning, making it a valuable tool to use when scan time is valuable or in populations where good fixation is hard to achieve.

To enhance the accuracy of eccentricity estimates in the template map, we aligned the distribution of eccentricity values from the fitted Benson template atlas with Horton and Hoyt's model (Horton & Hoyt, 1991). This model suggests that the linear cortical magnification factor (M_{linear} in mm/º) is inversely proportional to eccentricity (E in degrees), expressed as $M_{linear} = \frac{scale}{E+shape}$, with the normalizing parameter $shape$ equal to 0.75°, as measured by Horton and Hoyt (1991). For each hemisphere, we first estimated the *scale* value based on each subject's V1 surface area and the minimum and maximum eccentricities considered in the model (0º and 90º). The mean *scale* value for the left and right V1 hemispheres across all tested controls were 15.0±0.99 and 16.0±1.09, respectively.

In comparison, Horton and Hoyt (1991) measured a mean *scale* of 17.3 mm. The mean V1

surface area for the left and right hemispheres were 2700.7±358.7mm^2 and

3067.7±416.3mm^2.

We then defined a function $f(r)$ that describes how the surface area of V1 changes

with eccentricity (Equation 1). This was achieved by integrating Horton and Hoyt's function

over the visual field, resulting in an equation that calculates the amount of V1 surface area

from the foveal confluence up to a particular eccentricity (r). Next, we sorted the input

eccentricity values and calculated the cumulative surface areas. A numerical method was

then employed to find the eccentricity values that match the desired distribution. This method

involved iteratively adjusting the range of possible values until the calculated values closely

matched the desired values. Finally, the adjusted eccentricity values were returned in their

original order and values from both hemispheres were merged to provide bilateral

eccentricity estimates.

$$f(r) = \pi * scale^2 * \left(\log\left(\frac{r+shape}{shape}\right) - \frac{r}{r+shape} \right)$$ Equation 1

Results

To measure visual field function, we developed a new measure of cortical contrast

sensitivity, assessing the brain's ability to discriminate shapes based on luminance

variations. This depends on many factors, including the size, orientation, and density of cells

in different parts of the retina. As these factors change across the visual field, contrast

sensitivity varies with spatial frequency and visual field location in the typical, healthy visual

system. We first investigated whether our brain-based approach can reliably detect these

known differences in contrast sensitivity as a function of eccentricities and visual field

quadrants, in individual normal sighted controls. We then investigated whether our approach

could recover loss of visual inputs at the cortical level, either simulated in a healthy control or

pathological in a patient. In each section, findings were first reported for the pRF-defined

ROIs and then for the Benson-defined ROIs.

Spatial frequency preference changes in V1 neuronal populations across eccentricities

For high spatial frequency stimuli, we expected better sensitivity in V1 neuronal

populations encoding central locations in visual space than in those encoding the periphery.

In contrast, for low spatial frequency information, we expected the opposite, better sensitivity

in V1 neuronal populations encoding the periphery than the center of the visual field.

We found this expected interaction between spatial frequency and eccentricity

($F_{4,24} = 28.657, p < 0.001$; Figure 3) as well as a main effect of eccentricity ($F_{2,24} =$

$12.668, p < 0.001$). These effects were statistically significant at the individual participant

level showing high sensitivity (Figure 3A, transparent lines show individual data; Figure 3C

shows individual visual field plots; for stats, see Table A1 in *Appendix* section).

Post-hoc tests show that V1 neuronal populations receiving inputs from the central

visual field (0.5-4.5º) showed greater contrast sensitivity to high spatial frequency as

compared to low spatial frequency stimuli (steeper slope for the 3cpd versus 0.3cpd

condition: **0.5-2.5º**: $t(6) = 4.354, p = 0.005$; **2.5-4.5º**: $t(6) = 3.471, = 0.013$). Conversely,

peripheral eccentricities in V1 (above 9.5º) showed higher contrast sensitivity to low as

compared to high spatial frequency stimuli (steeper slope for 0.3cpd versus 3cpd condition

9.5-15º: $t(6) = -4.591, p = 0.004$; **15-20º**: $t(6) = -6.615, p < 0.001$). Between 4.5º and

9.5º, V1 contrast sensitivity was similar for both spatial frequencies ($t(6) = -0.226, p =$

0.829). Crucially, these effects remained when using retinotopic estimates based on

structural scans derived from the Benson retinotopic atlas instead of the pRF-mapping

measures (**0.5-2.5º**: $t(6) = 5.768, p = 0.001$; **2.5-4.5º**: $t(6) = 2.531, p = 0.045$; **4.5-9.5º**:

$t(6) = -0.293, p = 0.779$; **9.5-15º**: $t(6) = -3.274, p = 0.017$; **15-20º**: $t(6) = -3.528, p =$

0.012; see Figure A2 and Table A3 in *Appendix* section).

These results demonstrate that our approach can detect subtle changes in visual

sensitivity across eccentricities at the individual participant level, and that structure-based

atlases can replace the need for pRF mapping in cases where it might otherwise be difficult

or impossible to collect pRF data.

Figure 3. Spatial frequency preference V1 across the visual field. For each participant, five eccentricity bins were defined using subject-specific pRF estimates: 0.5º-2.5º, 2.5º-4.5º, 4.5º-9.5º, 9.5º-15º, and 15º-20º. **A:** Average β-values versus stimulus contrasts for low (0.3cpd; blue) and high (3cpd; red) spatial frequencies across eccentricity bins. Error bars are the 95% confidence intervals. Thick lines represent the group-level model fit, thin lines are the individual fits. **B:** Slopes projected into visual space for low (0.3 cpd) and high (3 cpd) spatial frequencies. The color scale corresponds to slope estimates, with higher values indicating higher cortical contrast sensitivities. **C:** Heat plots for each participant (C1-C7).

Contrast sensitivity is not equal across the visual field quadrants in V1 neurons

Contrast sensitivity is known to vary across the quadrants of the visual field; Previous behavioral studies have shown greater contrast sensitivity along the horizontal versus vertical meridians, and along the lower versus upper vertical meridian (Barbot et al., 2021; Carrasco et al., 2022; Himmelberg et al., 2020, 2023). Recent fMRI studies have also shown that these asymmetries in BOLD response amplitude to single contrast gratings (Kurzawski et al., 2022; Liu et al., 2006; O'Connell et al., 2016). Therefore, we expect our fMRI approach to uncover these established asymmetry patterns at the cortical level.

Figure 4 shows how cortical contrast sensitivity in V1 varies across visual field quadrants, for the 0.3cpd and 3cpd conditions. For this analysis, we collapsed eccentricities (0.5º to 20º) into a single bin. We found a main effect of visual field quadrant location on V1

sensitivity ($F_{3,18} = 20.706, p < 0.001$). Post-hoc tests revealed that, as predicted, the cortical contrast response function had a higher slope – indicating better V1 sensitivity – along the horizontal versus vertical quadrants (Horizontal-Vertical Anisotropy – **HVA:** $t(6) = 5.908, p = 0.001$) and along the lower versus upper quadrant (Vertical Meridian Anisotropy – **VMA:** $t(6) = 4.106, p = 0.006$). Conversely, no difference in cortical contrast sensitivity was found between V1 neuronal populations encoding the left and right quadrants of the visual field (Left-Right Horizontal Meridian Anisotropy – **LRHMA:** $t(6) = 0.7197, p = 0.499$). Moreover, there was no interaction between spatial frequency and visual field quadrant positions ($F_{3,18} = 1.339, p = 0.293$), suggesting V1 visual field anisotropies are relatively constant across spatial frequencies. Importantly, all these differences are also present in individual participants (Figure 4A, transparent lines are individual data, Figure 4C shows individual visual field plots; for stats, see Table A2 in *Appendix* section). All participants had better V1 sensitivity in the horizontal versus vertical quadrants. 6 out of 7 participants showed asymmetries between the lower and upper quadrants. Differences in V1 sensitivity between left and right quadrants were inconsistent across participants (3 participants: left>right; 2 participants: right>left; 2 participants: non-significant), in line with no difference at the group-level. This again demonstrates the high sensitivity of our approach for identifying and quantifying sensitivity changes across the visual field, at the individual participant level, offering a potential valuable tool for assessing and addressing visual field loss in clinical populations.

We next tested if we could recover these V1 anisotropies when using retinotopic estimates based on structural scans fit with the Benson atlas instead of pRF measures (see Figure A3 and Table A3-4 in *Appendix* section). We found that the horizontal-vertical anisotropy effect was recovered (**HVA:** $t(6) = 3.584, p = 0.012$), but that the vertical meridian anisotropy effect was not (**VMA:** $t(6) = 0.744, p = 0.485$) with this approach. These results provide evidence that our approach can still detect subtle visual field quadrant anisotropies using a structure-based retinotopic atlas, although less accurately.

While the most pronounced contrast sensitivity difference along visual field quadrants (horizontal versus vertical quadrants) was retained when the adjusted Benson atlas was used to replace pRF measurements, not all anisotropies survived. This may be because the template does not model cortical magnification differences along meridians, reducing accuracy. It is not surprising that retinotopic estimates based on structure-retinotopy relationship modeling, provides less accurate estimates of retinotopic tuning of neuron population in V1 than direct measures of the individual retinotopy. Any individual deviations from the model make it more challenging to accurately assign cortical contrast sensitivity to visual field locations, potentially decreasing the ability to detect changes in visual function

across the field. Given this challenge, results with the adjusted Benson map are surprisingly
similar to those based on pRF measurement, showing promise for clinical applications when
pRF mapping is unfeasible.

Figure 4. Anisotropies in V1 across visual field quadrants for low (0.3cpd; top row) and high (3cpd;
bottom row) spatial frequencies. pRF estimates were used to link brain responses to visual field
positions. Visual quadrants are ±45º around the cardinal meridians, with slope values for eccentricities
between 0.5º-20º. Horizontal quadrants are left and right; vertical quadrants are upper and lower. **A**:
Average β-values versus contrast levels for each anisotropy (horizontal vs. vertical, upper vs. lower,
left vs. right). Error bars are 95% confidence intervals. Thick lines correspond to average model fit,
which thin lines are individual fits. **B**: Slope values projected onto visual field quadrants. The color
scale represents slope estimates, with higher values indicating greater V1 contrast sensitivity. **C:** Heat
plots for each participant (C1-C7).

V1 sensitivity differences across the visual field are reproductible across visits.

        Having shown that changes in cortical contrast sensitivity across the visual space
can be detected with fMRI, it is crucial to determine how reliable our measure is. This is
especially important for potential applications in monitoring changes over time. Using
Spearman correlations, we measured the reliability of cortical contrast sensitivity estimates

for the 0.3cpd spatial frequency condition across two repeated measurements, collected on

two different days. We collected test-retest reliability measures from 4 out of 7 participants

(Figures 5A-B) and benchmarked them against the correlations between the 0.3cpd

condition and 3cpd spatial frequency condition, collected in the same session (Figure 5C). If

measures are reliable, correlations should be higher for repeated measures with the same

spatial frequency stimulus, collected on different days.

Our findings clearly show that the gradual increase in cortical sensitivity toward the

periphery for the 0.3cpd condition is consistent across the two sessions (Figure 5A).

Moreover, the data of individual participants show characteristic profiles that stay highly

consistent. Additionally, we observed moderate-to-strong positive Spearman correlations

across sessions for the 0.3cpd condition (Figure 5B; C2: $\rho = 0.5793$; C4: $\rho = 0.6996$; C5:

$\rho = 0.4059$; C6: $\rho = 0.7677$), which were consistently higher than those across different

spatial frequency conditions (Figure 5C; C2: $\rho = 0.3979$; C4: $\rho = 0.2386$; C5: $\rho = 0.1437$;

C6: $\rho = 0.3156$). This indicates that within participants, our measure is reliable across

sessions and distinguishes effectively between patterns elicited by various spatial frequency

conditions. Collectively, this suggests that our measure could serve as a robust tool for

detecting changes in cortical sensitivity, such as those occurring over time or following

therapeutic interventions.

Figure 5. Reliability of V1 slopes across sessions, in 4 of the 7 controls. **A:** V1 contrast sensitivity

(i.e., slopes) plotted in visual field space (using pRF estimates), showing a consistent session-

independent increase in cortical sensitivity toward peripheral locations for the 0.3cpd condition. Higher

slopes (warmer colors) indicate higher V1 sensitivity. **B:** Slopes for the 0.3cpd condition across two

visits. **C:** Slopes for the 0.3cpd and 3cpd conditions collected within the same session.

A test-case of simulated loss of visual inputs

In the previous sections, we showed that the slope of a square root function provides

a reliable measure of contrast sensitivity in the brain of healthy controls. But can this brain-

level model also quantify loss of visual inputs? To test this, we first simulated an artificial

scotoma in one normal sighted participant, by masking the 3cpd stimulus with a grey annular

ring between eccentricities of 3-8º (Figure 6A). We expect smaller slope values in V1

neuronal populations that would under normal circumstances encode that part of the visual

space.

As expected, we observed reduced responses in V1 locations corresponding to the

artificial scotoma (Figure 6B), with increased responses along the edges of the mask. This

artificial loss of visual input was also clearly present in the cortical contrast sensitivity

estimate (Figures 6C&D), with significantly reduced slope steepness between 3-8º in V1.

Additionally, we could recover this scotoma using the calibrated Benson template, although

less accurately (Figure 6E). These results show that this measure of V1 contrast sensitivity

is sensitive enough to detect loss of visual inputs in the brain at an individual level, when

complete local loss of sight is simulated

pRF mapping data from the individual.

patterns of vision loss and recovery at

e region of the artificial scotoma (3-8º). **A:**

toma. **B:** Averaged β-values between 0.5º-

20º eccentricities in V1, under normal and masked conditions. 20 eccentricity bins of 1º were selected

based on pRF estimates. The color scale indicates the presented contrast levels, and the black dotted

lines mark the boundaries of the artificial scotoma. **C:** Average β-values versus contrast levels in V1

neurons encoding the region of artificial scotoma (3º-8º), under normal (blue) and masked (red)

conditions. Lines correspond to the model fitted to the data. **D:** Slopes projected back onto the visual

field under normal and masked conditions. pRF estimates were used to relate brain responses to

visual field locations and to create the 1º eccentricity bins. The boundaries of the simulated scotoma

region are represented by dotted lines. The color scale indicates slope estimates, with higher values

(warmer colors) corresponding to higher V1 contrast sensitivity. **E:** Slopes plotted back into the visual

space using the calibrated Benson template instead of pRF estimates.

Case of severe central vision loss

We next tested if brain-based cortical contrast sensitivity could recover visual field

loss in a clinical patient with Leber Hereditary Optic Neuropathy (LHON; M, 45.5 years old,

mutation: m.11778G>A, disease duration: 36.1 years). LHON is a neurodegenerative

disease caused by mitochondrial DNA mutations, leading to acute loss of retinal ganglion

cells across large visual field areas. LHON patients often present with dense, central

scotoma extending up to 10-15 degrees around fixation.

As vision loss often varies between eyes, the patient underwent monocular testing for

the fMRI contrast sensitivity task. For comparison, a behavioral visual field perimetry map

was obtained with the Compass fundus perimeter (CenterVue, Padova, Italy) for each eye,

using a 24-2 testing grid with a Goldmann size III target (54 locations: 52 from the 24-2 grid,

1 at fixation, and 1 at the physiological blind spot). The Compass actively tracks the retina to

compensate for fixation loss, automatically repositioning targets based on current eye

position. Sensitivity thresholds at each location were determined using the ZEST projection

strategy implemented in the device. The patient, who had some preserved central vision,

could see the fixation targets in behavioral testing and the scanner, making them an ideal

participant for methods evaluation.

The Compass perimetry map revealed drastic loss of sensitivity across the visual

field (Figure 7 – top row). For the left eye, the left hemifield was more affected than the right,

with most severe loss in the upper-left quadrant. In contrast, the right eye showed more

reduced sensitivity in the right hemifield, particularly in the upper-right quadrant.

We found correspondence between affected and non-affected visual field regions

across the Compass and the fMRI contrast sensitivity measures (Figure 7). Here, V1

contrast sensitivity (i.e., slope values) was computed for the upper-left, upper-right, lower-left

and lower-right quadrants to match the layout from the Compass visual field maps. Using the

brain-based approach, we recovered sensitivity loss in the upper-right visual quadrant for the

right eye at the cortical level, reflected in smaller cortical contrast sensitivity slopes in V1
neuronal populations encoding that area. For the left eye, cortical contrast sensitivity was
also lower in the upper-left quadrant of the visual field, but differences were less
pronounced. This is in line with the observation that the left eye was the better eye, with
substantially less severe vision loss in the affected regions (lighter shading, Figure 7 – top
row). When using the calibrated Benson atlas instead of pRF mapping data to link neuronal
responses to visual locations, this pattern was retained for the right eye, but the more subtle
pattern in the left eye became noisier.

By combining large-field stimulation and contrast sensitivity modeling in fMRI, with
the cortical magnification-adjusted Benson atlas, these data show how brain-based
measures could provide a tool to characterize vision loss in patients without need for
demanding tasks or sustained precise fixation, while highlighting challenges of detecting
more subtle vision loss and pinpointing it to visual locations.

Figure 7. Correspondence between visual field perimetry map and V1 sensitivity map, in a patient
with Leber Hereditary Optic Neuropathy. **A:** Left eye. **B:** Right eye. **Top row:** Gray scale visual field
map obtained with the Compass fundus perimeter (CenterVue, Padova, Italy), using Standard
Perimetry display convention. Sensitivity is represented with symbols and related dB intervals, with
larger values describing better sensitivity. Red circle indicates the area of the visual space stimulated
in the fMRI contrast sensitivity task. **Middle row:** Heatmap of V1 sensitivity (i.e., slopes) across the
visual field. Here, we divided the space between the upper-left, upper-right, lower-left and lower-right
quadrants (as opposed to the left, right, upper and lower quadrants used in previous sections), to
match the layout of visual field perimetry maps. Slope values were averaged for each visual quadrant

and 1º eccentricity bin and projected back into the visual space, using the subject-specific pRF map.
Color scale corresponds to the steepness of the slope estimate. For better visualization of sensitivity
differences across quadrants, we also averaged the slopes between 0.5-20° eccentricities in each
quadrant, generating a single slope value for each visual quadrant (inset heat plots). The range of
slope values were reduced on the color scale for these inset plots. **Bottom:** Heatmap of V1 sensitivity
(i.e., slopes) across the visual field, using the Benson retinotopic template map instead of the pRF
data.

Discussion

In this study, we aimed to develop and validate an fMRI-based method for measuring
cortical contrast sensitivity across the entire visual field in V1, addressing the limitations of
traditional visual field tests and existing neuroimaging approaches. By integrating population
receptive field (pRF) modeling with contrast sensitivity measurements of the BOLD response
to large-field sinusoidal gratings varying in contrast levels and spatial frequencies, we
created a quantitative, sensitive, and reliable measure of cortical contrast sensitivity. Our key
findings demonstrate that this approach can accurately quantify known variations in V1
contrast sensitivity across eccentricity and polar angle, recover simulated and pathological
visual field losses, and produce consistent and reliable measures both at the group and
individual participant levels.

Our method successfully captured the expected variations in cortical contrast
sensitivity across the visual field. Consistent with previous behavioral and fMRI studies
(Aghajari et al., 2020; Broderick et al., 2022; D'Souza et al., 2016; Henriksson et al., 2008;
Rovamo et al., 1984; Rovamo & Virsu, 1979; Welbourne et al., 2018), we observed that V1
neuronal populations encoding the central visual field exhibited higher contrast sensitivity for
high spatial frequency (SF) stimuli, while those encoding peripheral regions were more
sensitive to low SF stimuli. This aligns with the understanding that spatial frequency tuning
varies across the visual field due to smaller pRF sizes and larger cortical areas dedicated to
central vision (Himmelberg et al., 2023).

Additionally, our study identified anisotropies in contrast sensitivity across visual field
quadrants, mirroring findings from previous behavioral studies (Barbot et al., 2021; Carrasco
et al., 2022; Himmelberg et al., 2020). Specifically, we found enhanced cortical sensitivity in
V1 neuronal populations responding to horizontal versus vertical locations and to lower
versus upper visual field locations. Unlike earlier fMRI studies (Liu et al., 2006; O'Connell et
al., 2016), which either could not assess horizontal-vertical effects or used smaller region
selections, our use of large-field stimuli enabled the detection of both upper-lower and
horizontal-vertical asymmetries. Moreover, by modeling cortical contrast sensitivity across

multiple contrast levels, we were able to identify subtle variations in V1 neuron sensitivity

that single-threshold approaches might overlook.

Crucially, these effects were significant at the individual level indicating high

sensitivity, and our approach proved reliable over time. We found consistent activation

patterns and moderate-to-high correlations in slope estimates for the 0.3cpd condition

across two separate sessions, significantly higher than when correlating responses to

different stimuli (i.e., 3cpd condition). High sensitivity and reliability are essential for

detecting longitudinal changes in clinical groups, such as those occurring before and after

sight-rescuing interventions. Fitting a contrast response function to the BOLD signal might

offer robust estimates of visual function by minimizing the impacts of overall BOLD amplitude

variability and arbitrary thresholds.

Furthermore, our method successfully identified both simulated retinal loss in a

healthy volunteer and real visual field loss in a patient with Leber Hereditary Optic

Neuropathy (LHON). Unlike previous fMRI studies that used high-contrast stimuli

(Farahbakhsh et al., 2022; Pawloff et al., 2023; Ritter et al., 2019) – which may not

accurately represent partial vision loss due to potential saturation and the stimulation of less

sensitive retinal cells – our use of multiple contrast levels offers a more nuanced

assessment of cortical contrast sensitivity. This may make our method particularly suitable

for evaluating treatment efficacy in cases of widespread and variable vision loss.

One limitation of this brain-based approach to visual field testing, is its reliance on

pRF mapping, which requires precise fixation and can be challenging to acquire in visually

impaired patients. To mitigate this, we investigated the use of retinotopic templates based on

anatomical landmarks (Benson et al., 2014; Benson & Winawer, 2018), thereby reducing the

need for pRF data collection and facilitating the mapping of brain responses to visual field

positions. Particularly at large eccentricities where, due to smaller cortical projection areas,

small inaccuracies between the template and individual retinotopy can lead to large

distortions in cortical visual field maps. To increase accuracy of the atlas at higher

eccentricities, we adjusted the Benson eccentricity estimates to align with the cortical

magnification scaling function (Horton & Hoyt, 1991). With this structure-based atlas we

successfully replicated key variations in visual field function, although with reduced

sensitivity compared to individual pRF measures for more subtle differences (i.e., for the less

consistent upper versus lower quadrant anisotropy, and the patient's eye less affected by

pathology). It is notable that cortical structure-based templates can successfully recover

visual field loss. Critically, in patient groups with severe impairment such as LHON, where it

is often very challenging or impossible to obtain reliable measures, even imperfect mapping

between the brain and visual field could offer important improvements over available

solutions for tracking functional change, and where they stem from. Further enhancing the

alignment between retinotopic template atlases and individual retinotopic tuning - for
example, by integrating them with functional measures using Bayesian methods (Benson &
Winawer, 2018) - could improve this approach further.
While our method demonstrated high accuracy and reliability in sighted controls,
further validation in patients is necessary to ensure the generalizability of our approach to
measuring pathology. Next steps in this work will therefore include reconstructing diverse
types of vision loss in larger patient cohorts, validating our approach for tracking pathology
over time, and investigating how cortex-based visual field measures relate to and
complement other visual field integrity indices such as the Compass in those contexts.
Comparison of the cortical contrast sensitivity modeling approach presented here to other
approaches will be another interesting direction for future research. For example, while we
reasoned that using a single high contrast stimulus is likely to give less reliable measures for
various reasons outlined above, this needs to be tested empirically. It is also unclear
whether the square root contrast response function is optimal for patients, who are likely to
show different non-linearities in their contrast response. Additionally, more extensive
mapping of the spatial frequency and contrast sensitivity spaces could offer additional
informative measures such as cortical acuity and spatial frequency preferences, provided
such measures are feasible with patient-friendly protocols.
In sum, brain-based contrast sensitivity measures could significantly enhance the
accurate characterization of visual field function in blinding diseases, by addressing
limitations of traditional visual field tests that are subjective, time-consuming, and
challenging for patients with severe sight loss. By providing a reliable, quantitative measure
of cortical contrast sensitivity across the visual field, fMRI may help facilitate the monitoring
of disease progression and recovery, including interventions like gene therapy. Combined
with our large-field stimulation setup and fixation free approaches to linking cortical
sensitivity estimates to visual field locations, this method is well-suited for a wide range of
visual impairments, addressing a pressing challenge in the evaluation of current clinical
trials. Additionally, linking brain-based variations in function across the visual field to
behavioral performance, could help bridge the gap between neural measures and functional
outcomes, offering deeper insights into developmental, learning, and vision loss
mechanisms.
In conclusion, integrating cortical contrast sensitivity testing with large-field
stimulation and a calibrated retinotopic atlas provides a promising tool for characterizing
visual field impairments, and could offer a valuable addition to existing measures of visual
field loss.

Acknowledgements

This work was funded by Moorfields Eye Charity PhD Studentship GR001315 (London, UK), and the Birkbeck-UCL Centre for NeuroImaging (BUCNI; London, UK). PYWM is supported by an Advanced Fellowship Award (NIHR301696) from the UK National Institute of Health Research (NIHR). PYWM also receives funding from Fight for Sight (UK), the Isaac Newton Trust (UK), Moorfields Eye Charity (GR001376), the Addenbrooke's Charitable Trust, the National Eye Research Centre (UK), the International Foundation for Optic Nerve Disease (IFOND), the NIHR as part of the Rare Diseases Translational Research Collaboration, the NIHR Cambridge Biomedical Research Centre (NIHR203312), and the NIHR Biomedical Research Centre based at Moorfields Eye Hospital NHS Foundation Trust and UCL Institute of Ophthalmology (NIHR203322). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. We extend our thanks to Simon Richardson and Oliver Josephs for their assistance in the design and conception of the large-field set-up.

Appendix

A. Large-Field set-up

Figure A1. Schematic of Large-Field Set-Up. **A:** The participant lies on the scanner bed in the 64-channel coil without top to reduce obstruction and views the 40° screen via a mirror above their face (1). Stimuli are displayed using back-projection from a projector (2) outside the room, onto a screen (3) positioned behind the participant's head. The top of the screen follows the scanner bore curvature to maximise field of view. Monocular eye-tracking is achieved by mounting the illuminator and camera of the eye-tracker (4) vertically on a support (5) at the back of the scanner bore. An image of the eye is obtained via a dual-mirror set-up, including a small mirror inside an aperture cut on the side of the screen (6) and the participant mirror (1). If the right eye is being tracked, the eye-tracker is placed on the left

side of the scanner bore, and the eye-tracker mirror (6) is placed in the right screen aperture.

B: Schematic of the large-field screen, with the area of stimulus presentation shown in cyan

and the mirror used for eye-tracking in magenta. The mirror is positioned within one of the

two rectangular apertures cut on either side of the screen.

B. Individual Level

To test for contrast sensitivity differences across the visual field in individual

participants, we also employed a multilevel modeling approach using the lme4 package in R

(Bates et al., 2015; R Core Team, 2023). This approach reduces the impact of vertex-level

variability and redundancy. The model included the square root of stimulus contrast as a

continuous fixed-effect predictor, and eccentricity, polar location, and spatial frequency as

categorical fixed-effect predictors, along with all possible interactions. Again, the intercept

was fixed at 0. The model also included random slopes for the square root contrast and the

interaction between square root contrast and voxel-specific factors and was fit separately for

each participant. In the formula notation used in the R package lme4, the model would be

defined as:

$\beta \sim 0 + SqrtContrast + SqrtContrast{:}SF + SqrtContrast{:}Eccentricity +$

$SqrtContrast{:}PolarLocation + SqrtContrast{:}SF{:}Eccentricity +$

$SqrtContrast{:}SF{:}PolarLocation + SqrtContrast{:}Eccentricity{:}PolarLocation +$

$SqrtContrast{:}Eccentricity{:}PolarLocation{:}SF + (0 + SqrtContrast +$

$SqrtContrast{:}SF \mid VoxelIdx)$

SubID	Effects	*Df*	Sum Sq	Mean Sq	*F*-value	*Pr(>F)*	Signif.
C1	SF	1	3720	3720	72.49	< 2e-16	***
	Eccentricity	1	35003	35003	681.995	< 2e-16	***
	PolarLocation	3	11258	3753	73.115	< 2e-16	***
	SF:Eccentricity	1	23914	23914	465.938	< 2e-16	***
	SF:PolarLocation	3	1546	515	10.041	1.32E-06	***
	Eccentricity:PolarLocation	3	200	67	1.297	0.274	
	SF:Eccentricity:PolarLocation	3	1158	386	7.519	5.04E-05	***
	Residuals	12378	635290	51			
C2	SF	1	23704	23704	743.628	< 2e-16	***
	Eccentricity	1	13503	13503	423.609	< 2e-16	***
	PolarLocation	3	46442	15481	485.647	< 2e-16	***
	SF:Eccentricity	1	10587	10587	332.135	< 2e-16	***
	SF:PolarLocation	3	2981	994	31.169	< 2e-16	***
	Eccentricity:PolarLocation	3	2881	960	30.126	< 2e-16	***
	SF:Eccentricity:PolarLocation	3	373	124	3.899	8.52E-03	**
	Residuals	11604	369897	32			
C3	SF	1	27	27	1.374	0.2411	
	Eccentricity	1	578	578	29.019	7.33E-08	***

	Effects	df	Sum Sq	Mean Sq	F value	Pr(>F)	
	PolarLocation	3	7315	2438	122.52	< 2e-16	***
	SF:Eccentricity	1	14719	14719	739.59	< 2e-16	***
	SF:PolarLocation	3	257	86	4.311	4.80E-03	**
	Eccentricity:PolarLocation	3	2225	742	37.261	< 2e-16	***
	SF:Eccentricity:PolarLocation	3	510	170	8.543	1.16E-05	***
	Residuals	9860	196225	20			
C4	SF	1	25290	25290	521.73	< 2e-16	***
	Eccentricity	1	3970	3970	81.89	< 2e-16	***
	PolarLocation	3	34475	11492	237.08	< 2e-16	***
	SF:Eccentricity	1	75054	75054	1548.38	< 2e-16	***
	SF:PolarLocation	3	2514	838	17.29	3.40E-11	***
	Eccentricity:PolarLocation	3	16417	5472	112.89	< 2e-16	***
	SF:Eccentricity:PolarLocation	3	3921	1307	26.97	< 2e-16	***
	Residuals	9896	479688	48			
C5	SF	1	554	554	16.332	5.37E-05	***
	Eccentricity	1	6717	6717	198.161	< 2e-16	***
	PolarLocation	3	886	295	8.711	9.12E-06	***
	SF:Eccentricity	1	13436	13436	396.384	< 2e-16	***
	SF:PolarLocation	3	4753	1584	46.744	< 2e-16	***
	Eccentricity:PolarLocation	3	3023	1008	29.727	< 2e-16	***
	SF:Eccentricity:PolarLocation	3	2638	879	25.939	< 2e-16	***
	Residuals	7728	261956	34			
C6	SF	1	6008	6008	246.099	< 2e-16	***
	Eccentricity	1	21371	21371	875.448	< 2e-16	***
	PolarLocation	3	11850	3950	161.806	< 2e-16	***
	SF:Eccentricity	1	12327	12327	504.98	< 2e-16	***
	SF:PolarLocation	3	714	238	9.745	2.04E-06	***
	Eccentricity:PolarLocation	3	1569	523	21.425	7.97E-14	***
	SF:Eccentricity:PolarLocation	3	762	254	10.409	7.80E-07	***
	Residuals	9360	228493	24			
C7	SF	1	10703	10703	278.799	< 2e-16	***
	Eccentricity	1	19431	19431	506.149	< 2e-16	***
	PolarLocation	3	18754	6251	162.84	< 2e-16	***
	SF:Eccentricity	1	19611	19611	510.822	< 2e-16	***
	SF:PolarLocation	3	153	51	1.325	2.64E-01	
	Eccentricity:PolarLocation	3	12080	4027	104.885	< 2e-16	***
	SF:Eccentricity:PolarLocation	3	1380	460	11.983	7.90E-08	***
	Residuals	10794	414384	38			

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

732 **Table A1.** Results from the multilevel modelling approach in each individual participant. pRF mapping

733 was used to link brain responses to visual field locations.

SubID	Effects	*contrast*	estimate	SE	df	t.ratio	p.value	
C1	PolarLocation	HM>VM	5.5138156	0.3278444	12354	16.818391	8.917786e-63	***

		LVM>UVM	0.5999187	0.2796053	12354	2.145591	0.03192504	*
		LHM>RHM	0.7058217	0.1711807	12354	4.123254	3.759997e-05	***
C2	PolarLocation	HM>VM	5.533751	0.2284595	11580	24.22203	1.771783e-126	***
		LVM>UVM	4.387544	0.1852974	11580	23.67839	4.420105e-121	***
		LHM>RHM	2.288693	0.1336361	11580	17.12630	5.942464e-65	***
C3	PolarLocation	HM>VM	3.568024	0.2172194	9836	16.425896	7.786414e-60	***
		LVM>UVM	2.317875	0.1801032	9836	12.869704	1.339442e-37	***
		LHM>RHM	0.159411	0.1214377	9836	1.312697	0.1893155	
C4	PolarLocation	HM>VM	7.3408010	0.3070954	9872	23.903972	8.223761e-123	***
		LVM>UVM	3.1514598	0.2492052	9872	12.646043	2.250910e-36	***
		LHM>RHM	-0.9417519	0.1794558	9872	-5.247821	1.571031e-07	***
C5	PolarLocation	HM>VM	2.2997460	0.2874033	7704	8.001807	1.405261e-15	***
		LVM>UVM	0.2736311	0.2296279	7704	1.191628	0.2334437	
		LHM>RHM	-0.2201730	0.1728344	7704	-1.273896	0.2027387	
C6	PolarLocation	HM>VM	1.8040008	0.2259007	9336	7.985813	1.561107e-15	***
		LVM>UVM	3.2006411	0.1818476	9336	17.600678	3.057311e-68	***
		LHM>RHM	0.3046082	0.1340245	9336	2.272779	0.02306211	*
C7	PolarLocation	HM>VM	5.9085493	0.3067623	10770	19.261005	2.637528e-81	***
		LVM>UVM	2.0334164	0.2617710	10770	7.767922	8.703410e-15	***
		LHM>RHM	-0.3370417	0.1599345	10770	-2.107373	0.03510831	*
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1								

Table A2. Post-Hoc test on ANOVA visual quadrant position effect (PolarLocation) at the individual level. pRF mapping was used to relate brain responses to visual field locations.

C. *Calibrated Benson Template*

Measure of Interest: Slope	num Df	den Df	MSE	F	ges	Pr(>F)	
SF	1.0000	6.0000	31.436	0.6117	0.010887	0.4639018	
ECC	2.1650	12.9901	32.694	4.7020	0.160027	0.0270342	*
PolarLocation	1.7729	10.6374	27.316	7.0357	0.163209	0.0129521	*
SF:Eccentricity	2.2706	13.6238	7.438	29.4822	0.221807	7.552e-06	***
SF:PolarLocation	1.5885	9.5309	5.173	2.1025	0.009793	0.1783533	
Eccentricity: PolarLocation	12.0000	72.0000	5.253	2.9645	0.096634	0.0020990	**
SF: Eccentricity: PolarLocation	12.0000	72.0000	1.195	3.5212	0.028091	0.0003995	***
Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1							

Table A3. Group-Level ANOVA analysis on slope estimates, as function of spatial frequency, eccentricities, and visual quadrant positions. The Benson template retinotopic map was used to relate brain responses to visual field locations.

Measure of Interest: Slope	contrast	estimate	SE	df	t.ratio	p.value
PolarLocation	Horizontal>Vertical	4.3378240	1.2101752	6	3.5844596	0.01158079
	Lower>Upper	0.5249520	0.7053736	6	0.7442184	0.48485944
	Left>Right	-0.2265194	0.3922638	6	-0.5774671	0.58462631
SF:Eccentricity	3cpd-0.3cpd: 0.5-2.5°	4.5917619	0.7960846	6	5.7679320	0.001185113
	3cpd-0.3cpd: 2.5-4.5°	2.5801243	1.0192745	6	2.5313340	0.044599322
	3cpd-0.3cpd: 4.5-9.5°	-0.2927635	0.9981812	6	-0.2932969	0.779175269

| | 3cpd-0.3cpd: 9.5-15º | -1.9782789 | 0.6042727 | 6 | -3.2738182 | 0.016953076 |
| | 3cpd-0.3cpd: 15-20º | -2.2803080 | 0.6462658 | 6 | -3.5284368 | 0.012391168 |

Table A4. Post-hoc t-test analysis for the visual quadrant position effect (PolarLocation) and the interaction between spatial frequency and eccentricities (SF:Eccentricity) reported in Table A3. The Benson template retinotopic map was used to relate brain responses to visual field locations.

Figure A2. V1 contrast sensitivity (i.e., slopes) across 5 eccentricity bins (0.5º-2.5º, 2.5º-4.5º, 4.5º-9.5º, 9.5º-15º, and 15-20º), defined using the calibrated Benson atlas. **A:** β-values versus contrast levels for each eccentricity bin. Blue and red lines represent the contrast sensitivity model fits for the 0.3cpd and 3cpd conditions, respectively. Thinner lines correspond to individual fits. **B:** V1 contrast sensitivity index (i.e., slope) projected back into the visual space using the calibrated Benson atlas, for each eccentricity bin and spatial frequency condition. The color scale indicates slope estimates, with higher values representing higher V1 contrast sensitivity.

Figure A3. V1 contrast sensitivity (i.e., slopes) across visual field quadrants using the calibrated Benson atlas, for the 0.3cpd and 3cpd conditions. Visual quadrants were defined as ±45º regions around the cardinal meridians and slope values were selected for eccentricities between 0.5-20º.

Horizontal quadrants are left and right, whilst upper and lower quadrants define the vertical quadrants. **A:** β-values versus contrast levels for each anisotropy effects: horizontal versus vertical, upper versus lower, and left versus right. Group-level model fits are in thick lines, whilst thinner lines correspond to individual fits. Error bars represent the 95% confidence intervals. **B:** V1 contrast sensitivity index (i.e., slope) projected back into the upper, lower, left and right visual field quadrants for the 0.3cpd and 3cpd conditions. The color scale represents slope estimates, with higher values indicating higher V1 contrast sensitivity.

Figure A4. V1 contrast sensitivity (i.e., slope) to the 0.3cpd condition plotted back into visual space under fixation (A) and controlled eye-motion (B) conditions, in participant C6. pRF estimates were used to relate brain responses to visual field positions.

References

Aghajari, S., Vinke, L. N., & Ling, S. (2020). Population spatial frequency tuning in human early visual cortex. *Journal of Neurophysiology*, *123*(2), 773–785. https://doi.org/10.1152/jn.00291.2019

Alvarez, G. A. (2011). Representing multiple objects as an ensemble enhances visual cognition. *Trends in Cognitive Sciences*, *15*(3), 122–131. https://doi.org/10.1016/j.tics.2011.01.003

Barbot, A., Xue, S., & Carrasco, M. (2021). Asymmetries in visual acuity around the visual field. *Journal of Vision*, *21*(1), 1–23. https://doi.org/10.1167/JOV.21.1.2

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using **lme4**. *Journal of Statistical Software*, *67*(1). https://doi.org/10.18637/jss.v067.i01

Benson, N. C., Butt, O. H., Brainard, D. H., & Aguirre, G. K. (2014). Correction of Distortion in Flattened Representations of the Cortical Surface Allows Prediction of V1-V3

Functional Organization from Anatomy. *PLoS Computational Biology*, *10*(3). https://doi.org/10.1371/journal.pcbi.1003538

Benson, N. C., & Winawer, J. (2018). Bayesian analysis of retinotopic maps. *eLife*, 7, e40224. https://doi.org/10.7554/eLife.40224

Bourne, R., Steinmetz, J. D., Flaxman, S., Briant, P. S., Taylor, H. R., Resnikoff, S., Casson, R. J., Abdoli, A., Abu-Gharbieh, E., Afshin, A., Ahmadieh, H., Akalu, Y., Alamneh, A. A., Alemayehu, W., Alfaar, A. S., Alipour, V., Anbesu, E. W., Androudi, S., Arabloo, J., … Vos, T. (2021). Trends in prevalence of blindness and distance and near vision impairment over 30 years: An analysis for the Global Burden of Disease Study. *The Lancet Global Health*, *9*(2), e130–e143. https://doi.org/10.1016/S2214-109X(20)30425-3

Broderick, W. F., Simoncelli, E. P., & Winawer, J. (2022). Mapping spatial frequency preferences across human primary visual cortex. *Journal of Vision*, *22*(4), 1–21. https://doi.org/10.1167/jov.22.4.3

Buracas, G. T., & Boynton, G. M. (2007). The effect of spatial attention on contrast response functions in human visual cortex. *Journal of Neuroscience*, *27*(1), 93–97. https://doi.org/10.1523/JNEUROSCI.3162-06.2007

Buracas, G. T., Fine, I., & Boynton, G. M. (2005). The relationship between task performance and functional magnetic resonance imaging response. *Journal of Neuroscience*, *25*(12), 3023–3031. https://doi.org/10.1523/JNEUROSCI.4476-04.2005

Carrasco, M., Roberts, M., Myers, C., & Shukla, L. (2022). Visual field asymmetries vary between children and adults. *Current Biology*, *32*(11), R509–R510. https://doi.org/10.1016/j.cub.2022.04.052

Cauley, S. F., Polimeni, J. R., Bhat, H., Wald, L. L., & Setsompop, K. (2014). Interslice leakage artifact reduction technique for simultaneous multislice acquisitions: Interslice Leakage Artifact Reduction Technique. *Magnetic Resonance in Medicine*, *72*(1), 93–102. https://doi.org/10.1002/mrm.24898

DeYoe, E. A., Carman, G. J., Bandettini, P., Glickman, S., Wieser, J., Cox, R., Miller, D., &

Neitz, J. (1996). Mapping striate and extrastriate visual areas in human cerebral

cortex. *Proceedings of the National Academy of Sciences*, *93*(6), 2382–2386.

https://doi.org/10.1073/pnas.93.6.2382

D'Souza, D. V., Auer, T., Frahm, J., Strasburger, H., & Lee, B. B. (2016). Dependence of

chromatic responses in V1 on visual field eccentricity and spatial frequency: An fMRI

study. *Journal of the Optical Society of America A*, *33*(3), A53.

https://doi.org/10.1364/JOSAA.33.000A53

Dumoulin, S. O., & Wandell, B. A. (2008). Population receptive field estimates in human

visual cortex. *NeuroImage*, *39*(2), 647–660.

https://doi.org/10.1016/j.neuroimage.2007.09.034

Engel, S. (1997). Retinotopic organization in human visual cortex and the spatial precision of

functional MRI. *Cerebral Cortex*, *7*(2), 181–192.

https://doi.org/10.1093/cercor/7.2.181

Farahbakhsh, M., Anderson, E. J., Maimon-Mor, R. O., Rider, A., Greenwood, J. A., Hirji, N.,

Zaman, S., Jones, P. R., Schwarzkopf, D. S., Rees, G., Michaelides, M., & Dekker,

T. M. (2022). A demonstration of cone function plasticity after gene therapy in

achromatopsia. *Brain*, *145*(11), 3803–3815. https://doi.org/10.1093/brain/awac226

Henriksson, L., Nurminen, L., Hyvarinen, A., & Vanni, S. (2008). Spatial frequency tuning in

human retinotopic visual areas. *Journal of Vision*, *8*(10), 5–5.

https://doi.org/10.1167/8.10.5

Himmelberg, M. M., Winawer, J., & Carrasco, M. (2020). Stimulus-dependent contrast

sensitivity asymmetries around the visual field. *Journal of Vision*, *20*(9), 1–19.

https://doi.org/10.1167/jov.20.9.18

Himmelberg, M. M., Winawer, J., & Carrasco, M. (2023). Polar angle asymmetries in visual

perception and neural architecture. *Trends in Neurosciences*, *46*(6), 445–458.

https://doi.org/10.1016/j.tins.2023.03.006

Horton, J. C., & Hoyt, W. F. (1991). The Representation of the Visual Field in Human Striate

Cortex: A Revision of the Classic Holmes Map. *Archives of Ophthalmology*, *109*(6),

816–824. https://doi.org/10.1001/archopht.1991.01080060080030

Kurzawski, J. W., Gulban, O. F., Jamison, K., Winawer, J., & Kay, K. (2022). Non-Neural

Factors Influencing BOLD Response Magnitudes within Individual Subjects. *Journal*

of Neuroscience, *42*(38), 7256–7266. https://doi.org/10.1523/JNEUROSCI.2532-

21.2022

Lange, R., Kumagai, A., Weiss, S., Zaffke, K. B., Day, S., Wicker, D., Howson, A.,

Jayasundera, K. T., Smolinski, L., Hedlich, C., Lee, P. P., Massof, R. W., Stelmack,

J. A., Carlozzi, N. E., & Ehrlich, J. R. (2021). Vision-related quality of life in adults

with severe peripheral vision loss: A qualitative interview study. *Journal of Patient-*

Reported Outcomes, *5*(1). https://doi.org/10.1186/s41687-020-00281-y

Lisboa, R., Chun, Y. S., Zangwill, L. M., Weinreb, R. N., Rosen, P. N., Liebmann, J. M.,

Girkin, C. A., & Medeiros, F. A. (2013). Association between rates of binocular visual

field loss and vision-related quality of life in patients with glaucoma. *JAMA*

Ophthalmology, *131*(4), 486–494. https://doi.org/10.1001/jamaophthalmol.2013.2602

Liu, T., Heeger, D. J., & Carrasco, M. (2006). Neural correlates of the visual vertical meridian

asymmetry. *Journal of Vision*, *6*(11), 12–12. https://doi.org/10.1167/6.11.12

O'Connell, C., Ho, L. C., Murphy, M. C., Conner, I. P., Wollstein, G., Cham, R., & Chan, K.

C. (2016). Structural and functional correlates of visual field asymmetry in the human

brain by diffusion kurtosis MRI and functional MRI. *NeuroReport*, *27*(16), 1225–1231.

https://doi.org/10.1097/WNR.0000000000000682

Oliva, A. (2005). CHAPTER 41—Gist of the Scene. In L. Itti, G. Rees, & J. K. Tsotsos (Eds.),

Neurobiology of Attention (pp. 251–256). Academic Press.

https://doi.org/10.1016/B978-012375731-9/50045-8

Pawloff, M., Linhardt, D., Woletz, M., Hummer, A., Sacu, S., Vasileiadi, M., Garikoitz, L. U.,

Holder, G., Schmidt-Erfurth, U. M., Windischberger, C., & Ritter, M. (2023).

Comparison of Stimulus Types for Retinotopic Cortical Mapping of Macular Disease.

Translational Vision Science and Technology, 12(3), 1–13.

867   https://doi.org/10.1167/tvst.12.3.6

868   R Core Team. (2023). *R: A Language and Environment for Statistical Computing.* [Computer

869   software]. R Foundation for Statistical Computing. https://www.R-project.org/

870   Ritter, M., Hummer, A., Ledolter, A. A., Holder, G. E., Windischberger, C., & Schmidt-Erfurth,

871   U. M. (2019). Correspondence between retinotopic cortical mapping and

872   conventional functional and morphological assessment of retinal disease. *British*

873   *Journal of Ophthalmology*, *103*(2), 208–215. https://doi.org/10.1136/bjophthalmol-

874   2017-311443

875   Roh, M., Selivanova, A., Shin, H. J., Miller, J. W., & Jackson, M. Lou. (2018). Visual acuity

876   and contrast sensitivity are two important factors affecting vision-related quality of life

877   in advanced age-related macular degeneration. *PLoS ONE*, *13*(5), 1–12.

878   https://doi.org/10.1371/journal.pone.0196481

879   Rovamo, J., Leinonen, L., Laurinen, P., & Virsu, V. (1984). Temporal Integration and

880   Contrast Sensitivity in Foveal and Peripheral Vision. *Perception*, *13*(6), 665–674.

881   https://doi.org/10.1068/p130665

882   Rovamo, J., & Virsu, V. (1979). An estimation and application of the human cortical

883   magnification factor. *Experimental Brain Research*, *37*(3).

884   https://doi.org/10.1007/BF00236819

885   Sereno, M. I., Dale, A. M., Reppas, J. B., Kwong, K. K., Belliveau, J. W., Brady, T. J., Rosen,

886   B. R., & Tootell, R. B. H. (1995). Borders of Multiple Visual Areas in Humans

887   Revealed by Functional Magnetic Resonance Imaging. *Science*, *268*(5212), 889–

888   893. https://doi.org/10.1126/science.7754376

889   Subhi, H., Latham, K., Myint, J., & Crossland, M. D. (2017). Functional visual fields:

890   Relationship of visual field areas to self-reported function. *Ophthalmic and*

891   *Physiological Optics*, *37*(4), 399–408. https://doi.org/10.1111/opo.12362

Welbourne, L. E., Morland, A. B., & Wade, A. R. (2018). Population receptive field (pRF)

measurements of chromatic responses in human visual cortex using fMRI.

NeuroImage, *167*, 84–94. https://doi.org/10.1016/j.neuroimage.2017.11.022

World Health Organization. (2019, October 8). *WHO launches first World report on vision*.

https://www.who.int/news/item/08-10-2019-who-launches-first-world-report-on-vision

Xu, J., Moeller, S., Auerbach, E. J., Strupp, J., Smith, S. M., Feinberg, D. A., Yacoub, E., &

Uğurbil, K. (2013). Evaluation of slice accelerations using multiband echo planar

imaging at 3T. *NeuroImage*, *83*, 991–1001.

https://doi.org/10.1016/j.neuroimage.2013.07.055