TXNIP mediates LAT1/SLC7A5 endocytosis to reduce amino acid uptake in cells entering quiescence

Jennifer Kahlhofer[1,2], Nikolas Marchet[1,2], Brigitta Seifert[2], Kristian Zubak[1], Madlen Hotze[3], Anna-Sophia Egger[3], Claudia Manzl[4], Yannick Weyer[1,2], Sabine Weys[2,5], Martin Offterdinger[6], Sebastian Herzog[7], Veronika Reiterer[8], Marcel Kwiatkowski[3], Saskia B. Wortmann[9,10], Siamak Nemati[11], Johannes A. Mayr[9], Johannes Zschocke[12], Bernhard Radlinger[13], Kathrin Thedieck[3, 15], Lukas A. Huber[2], Hesso Farhan[8], Mariana E.G. de Araujo[2], Susanne Kaser[13], Sabine Scholl-Bürgi[14], Daniela Karall[14], and David Teis[1,2+]

[1] Institute of Molecular Biochemistry, Biocenter, Medical University of Innsbruck, 6020 Innsbruck, Austria.
[2] Institute of Cell Biology, Biocenter, Medical University of Innsbruck, 6020 Innsbruck, Austria.
[3] Department of Biochemistry and Center for Molecular Biosciences Innsbruck, University of Innsbruck, 6020 Innsbruck, Austria
[4] Institute of Pathology, Neuropathology & Molecular Pathology, Medical University of Innsbruck, 6020 Innsbruck, Austria.
[5] Institute of Science and Technology Austria (ISTA), 3400 Klosterneuburg, Austria.
[6] Institute of Neurobiochemistry, BioOptics core facility, Biocenter, Medical University of Innsbruck, 6020 Innsbruck, Austria.
[7] Institute for Developmental Immunology, Biocenter, Medical University of Innsbruck, 6020 Innsbruck, Austria.
[8] Institute of Pathophysiology, Biocenter, Medical University of Innsbruck, 6020 Innsbruck, Austria.
[9] University Children's Hospital, Salzburger Landeskliniken and Paracelsus Medical University, 5020 Salzburg, Austria.
[10] Amalia Children's Hospital, Department of Pediatrics, Pediatric Neurology, Nijmegen, The Netherlands
[11] Kinder- und Jugendheilkunde, Krankenhaus St. Vinzenz Zams, 6511 Zams, Austria
[12] Institute of Human Genetics, Medical University of Innsbruck, 6020 Innsbruck, Austria.
[13] Department of Internal Medicine I, Medical University of Innsbruck, 6020 Innsbruck, Austria.
[14] Department of Child and Adolescent Health, Division of Pediatrics I-Inherited Metabolic Disorders, Medical University of Innsbruck, 6020 Innsbruck, Austria.
[15] Department Metabolism, Senescence and Autophagy, Research Center One Health Ruhr, University Alliance Ruhr & University Hospital Essen, University Duisburg-Essen, Essen; German Cancer Consortium (DKTK), partner site Essen, a partnership between German Cancer Research Center (DKFZ) and University Hospital Essen, Heidelberg; Freiburg Materials Research Center (FMF), University Freiburg, Freiburg, Germany; Laboratory of Pediatrics, Section Systems Medicine of Metabolism and Signaling, University of Groningen, University Medical Center Groningen, Groningen, Netherlands.

+corresponding author:
David Teis, PhD
Institute for Molecular Biochemistry, Medical University of Innsbruck
Innrain 80/82, CCB Building
A-6020, Innsbruck, Austria
tel: +43512900370191
email: david.teis@i-med.ac.at

Abstract

Entry and exit from cellular quiescence require dynamic adjustments in nutrient acquisition, yet

the mechanisms by which quiescent cells downregulate amino acid (AA) transport remain

poorly understood. Here, we demonstrate that cells entering quiescence select plasma

membrane-resident AA transporters for endocytosis and lysosomal degradation, to match AA

uptake with reduced translation. We identify the α-arrestin TXNIP as a key regulator of AA

uptake during quiescence, since it mediates the endocytosis of the SLC7A5-SLC3A2 (LAT1-

4F2hc) transporter complex in response to reduced AKT signaling. Mechanistically, TXNIP

interacts with HECT-type ubiquitin ligases to facilitate transporter ubiquitination. Loss of

TXNIP disrupts this regulation, resulting in dysregulated AA uptake, sustained mTORC1

signaling, and accelerated quiescence exit. A novel TXNIP loss-of-function mutation in a

patient with severe metabolic disease further supports its role in nutrient homeostasis and

human health. These findings highlight TXNIP's role in controlling SLC7A5-SLC3A2

mediated AA acquisition with implications for quiescence biology and disease.

Introduction

How cells maintain a homeostatic pool of 20 proteinogenic amino acids (AA) is a fundamental question in biology. For the uptake and the release of AA across the plasma membrane (PM) and across organelle membranes, the human genome encodes at least 66 AA transporters, which belong to 11 individual solute carrier (SLC) families [1,2]. The importance of controlling AA transport is exemplified by monogenic diseases and human pathologies caused by the deficiency or dysregulation of individual AA transporters [3,4].

Cellular AA import is coordinated with cell proliferation. Proliferating cells, in particular cancer cells, increase the abundance of AA transporters at the cell surface and thereby promote nutrient uptake for protein synthesis during cell growth prior to cell division [5]. Transcriptional mechanisms are responsible for the upregulation of AA transporter expression [6-9].

Conversely, entry into quiescence, the reversible exit from the cell cycle, is accompanied by a reduction in cell size, lower rates of transcription, and reduced protein synthesis [10,11]. Thus, quiescent cells must somehow re-calibrate AA uptake to align it with lower translation and metabolic homeostasis for cell survival, rather than for growth [12]. How quiescent cells match AA flux across the PM to reduced cell size and to lower translation rates is unclear. This presents a knowledge gap in understanding cellular nutrient acquisition.

In budding yeast, *Saccharomyces cerevisiae*, we previously demonstrated that entry into quiescence induces the ubiquitin dependent endocytosis of AA transporters [13]. The selective ubiquitination of AA transporters requires proteins of the α-arrestin family [14] [15]. Activated α-arrestins function as adaptor proteins that recruit the homologous to the E6-AP carboxyl terminus (HECT) type ubiquitin ligase Rsp5 to different AA transporters for ubiquitination, endocytosis and lysosomal degradation [16-18].

The human genome encodes six α-arrestin family members or ARRDCs (arrestin-domain-containing proteins). With the exception of ARRDC5, α-arrestins contain one or more PPxY motifs in their C-terminal regions and are able to interact with several HECT-type ubiquitin ligases [19-22]. The best characterized α-arrestin is thioredoxin interacting protein (TXNIP). TXNIP was first described as a negative regulator of thioredoxin (TRX) through the formation of an intermolecular disulfide bond that requires two critical cysteine residues of TXNIP (cysteine 32 and cysteine 247) [23-25].

TXNIP also regulates glucose uptake by mediating the selective endocytosis of SLC2A1 (GLUT1) and SLC2A4 (GLUT4), two major glucose transporters. TXNIP binds to these glucose transporters at the PM and interacts with a di-leucine motif in its C-terminal tail with components of the endocytic machinery [26,27]. The TXNIP dependent regulation of SLC2A1 and

SLC2A4 is tightly coupled to intracellular glucose homeostasis, ATP availability and growth

factor signaling. AMPK activation in response to low ATP levels or AKT activation in response

to insulin signal results in the phosphorylation of TXNIP at serine 308, thereby inactivating it

[26,27]. This inactivation of TXNIP ensures SLC2A1 and SLC2A4 accumulation at the PM to

increase glucose uptake [26,27]. TXNIP inactivation can involve ubiquitination by the HECT-type

ubiquitin ligase ITCH, leading to its proteasomal degradation [19,22]. These processes appear to

be largely independent of TRX binding. Consistent with the function in regulating glucose

uptake, biallelic loss-of-function mutations in TXNIP in humans lead to lactic acidosis in three

siblings. Interesting, these patients also showed lower serum methionine levels [28].

Here, we show that in human cells entering quiescence, several AA transporters are selectively

removed from the PM by endocytosis, including the heterodimeric AA transporter (HAT)

SLC7A5-SLC3A2 (LAT1-4F2hc). The endocytic degradation of SLC7A5-SLC3A2 during

entry into quiescence requires TXNIP. TXNIP uses its PPxY motifs to engage HECT-type

ubiquitin ligases for SLC7A5-SLC3A2 endocytosis and subsequent lysosomal degradation.

The TXNIP mediated endocytic degradation of SLC7A5-SLC3A2 reduces AA uptake, lowers

free intracellular AA levels, and thereby helps to dampen mTORC1 signaling and translation.

In growing cells, AKT phosphorylation of TXNIP restricts SLC7A5-SLC3A2 endocytosis. In

addition, we identified a novel biallelic loss-of-function TXNIP variant in a boy with a severe

inborn metabolic disease. The analysis of fibroblasts from this patient revealed that SLC7A5-

SLC3A2 endocytosis was also blocked. This further supports the central role of TXNIP in

adjusting cellular AA uptake, with important implications for human health.

Results

Selective endocytosis and lysosomal degradation of AA transporters in cells entering

quiescence.

To study the regulation of nutrient uptake during the transition from proliferation to quiescence,

we used hTERT RPE1 cells (non-cancerous human telomerase-immortalized retinal pigmented

epithelial, hereafter RPE1) as a model cell line [29,30]. First, we established and evaluated

conditions for entry into quiescence.

The DNA content of cells growing with serum was analyzed by propidium iodide (PI) staining

and fluorescence activated cell sorting (FACS). 46% ($\pm$ 2%) of these cells were in the G0/G1

phase of the cell cycle, 30% ($\pm$ 6%) were in S phase and 24% ($\pm$ 4%) in the G2/M phase. 24 h

after serum starvation (but in the presence of AA and glucose), the number of cells in G0/G1

increased to 67% ($\pm$ 10%), while the number of cells in S-phase and G2/M phase decreased to

15% (± 6%) and 19% (± 7%), respectively (Figure 1a). Cell viability was not affected by serum

starvation (Supplementary Figure 1a). Serum starved cells also decreased their volume by 20%

from 2907 fl to 2332 fl (Figure 1b). Consistent with an entry into quiescence in response to

serum starvation, SDS-PAGE and Western Blot (WB) analysis of total cell lysates showed that

the protein levels of the cyclin-dependent kinase inhibitor p27^{Kip1} increased, while the

phosphorylation dependent activation of mitogen activated kinases ERK1/2, and mTORC2-

dependent phosphorylation of AKT on serine 473 decreased (Figure 1c, lane 2).

Of note, in cells entering quiescence, the protein levels of the heterodimeric AA transporters

(HAT) SLC7A5-SLC3A2 (LAT1-4F2hc) and SLC7A11-SLC3A2 (xCT-4F2hc) decreased

(Figure 1c, lane 1 and 2). Also, a decrease in the protein levels of SLC1A5 (ASCT2) was

detected (Figure 1c, lane 1 and 2). The protein levels of the neutral AA transporter SLC38A2

(SNAT2), the glucose transporter SLC2A1 (GLUT1), the transferrin receptor TfR and

epidermal growth factor receptor (EGFR) remained relatively unchanged over the course of the

experiment (Figure 1c, lane 1 - 3). The downregulation of SLC7A5 and SLC1A5 was observed

in other non-cancerous cell lines, such as mouse embryonic fibroblasts (MEFs) (Supplementary

Figure 1b) and primary human fibroblasts (Supplementary Figure 1c, d), when subjected to

serum starvation. However, this downregulation was not observed in a panel of seven different

lung cancer cell lines (Supplementary Figure 1e).

To understand how SLC7A5-SLC3A2 and SLC1A5 were downregulated in cells entering

quiescence, we examined the localization of these transporters using indirect

immunofluorescence (IF) and confocal microscopy. In growing cells, endogenous SLC3A2

(Figure 1e) and SLC1A5 (Supplementary Figure 1f) were mainly detected at the PM and

partially co-localized with LAMP1 positive lysosomes (Figure 1e and Supplementary Figure

1f). In serum starved cells, signals for SLC3A2 were barely detected (Figure 1e), which was in

line with the downregulation of SLC7A5-SLC3A2 in quiescent cells. Also, cell surface FACS

analysis measured a marked decrease of PM resident SLC3A2 (Figure 1d). Treatment of cells

with chloroquine (CQ), which impairs lysosomal catabolic function, resulted in the

accumulation of SLC3A2 (Figure 1e) and SLC1A5 (Supplementary Figure 1f) inside LAMP1

positive lysosomes and efficiently prevented the lysosomal degradation of these AA

transporters (Figure 1f). Inhibition of the GTPase dynamin with dynasore (dyna), which blocks

the scissions of clathrin coated vesicles from the PM [31,32], impaired the endocytosis of SLC3A2

(Fig 1g) and SLC1A5 (Supplementary Figure 1g) and in consequence prevented their lysosomal

degradation (Figure 1h).

In line with the quiescence induced endocytic removal of SLC7A5-SLC3A2 from the PM, the
uptake of its AA substrates [^{3}H]-leucine and [^{14}C]-isoleucine was reduced by 33% and 49%,
respectively (Figure 1i). Similarly, the uptake of the SLC1A5 substrate [^{14}C]-glutamine
decreased by 20% (Supplementary Figure 1g).
The decrease in AA uptake upon entry into quiescence correlated with a reduction of mTORC1
signaling (*e.g.* towards S6K) and with an increase of eIF2α phosphorylation at serine 51 (Figure
1c, line 1 and 2), indicating a downregulation of translation. Consistently, the incorporation of
[^{35}S]-methionine and [^{35}S]-cysteine into newly synthesized proteins was reduced in quiescent
cells (Figure 1j, line 1 and 2).
Upon re-addition of serum to starved cells, SLC7A5-SLC3A2, SLC7A11-SLC3A2, and
SLC1A5 protein levels were upregulated, ERK-, AKT- as well as mTORC1-signaling were re-
activated, phosphorylation of eIF2α decreased as did p27^{Kip1} protein levels (Figure 1c, lane 3).
Translation (Figure 1j, lane 3) and cell size increased (Figure 1c), and cells resumed
proliferation (Figure 1a).
These results showed that the selective endocytosis and degradation of SLC7A5-SLC3A2
coincided with a decrease in cell volume, with a drop in translation, and with a reduction in
(iso)leucine uptake as cells entered quiescence in response to serum starvation.
184
TXNIP is required for the selective downregulation of SLC7A5
To identify the molecular mechanism responsible for the quiescence induced endocytic
degradation of SLC7A5-SLC3A2, we focused on the α-arrestin family of ubiquitin ligase
adaptors. α-arrestins are required for the endocytic downregulation of AA transporters in
budding yeast cells entering quiescence [13,33]. It was not known if they had a similar function in
human cells.
The human genome encodes six α-arrestins, called ARRDC (arrestin domain containing) 1-5,
and TXNIP [15]. TXNIP protein levels (Figure 2a, b) and mRNA levels (Supplementary Figure
2a) increased in response to serum starvation.
To determine whether TXNIP contributed to SLC7A5-SLC3A2 endocytosis and lysosomal
degradation, we used CRISPR/Cas9 mediated gene editing and introduced a 13 base pair
deletion at position c.644_656 leading to a frameshift that resulted in a premature stop codon
at position 226 (p.I215TfsTer11) (Supplementary Figure 2b, c). This mutation disrupted the
architecture of the arrestin C-domain (Supplementary Figure 2b) and in single cell clones, the
mutant protein was no longer detected (Figure 2a).

The loss-of-function mutation in TXNIP selectively prevented the endocytic downregulation of the SLC7A5-SLC3A2 heterodimer (Figure 2a, lane 2, and Figure 2b). In TXNIP[KO] cells, SLC3A2 remained at the PM in response to serum starvation, as determined by IF and cell surface FACS (Figure 2c, d and Supplementary Figure 2d). Re-expression of TXNIP in TXNIP[KO] cells restored SLC7A5-SLC3A2 endocytosis and degradation in response to serum starvation (Figure 2c, d and Supplementary Figure 2d - f). Conversely, TXNIP was not required for the degradation of SLC7A11 and SLC1A5 (Figure 2a, lane 2 and Supplementary Figure 2b). Hence, in cells entering quiescence, TXNIP selectively targeted SLC7A5-SLC3A2 for endocytosis and degradation. Loss of TXNIP did not affect SLC7A5 mRNA levels (Supplementary Figure 2g).

To examine the localization of TXNIP, ALFA-tagged TXNIP (ALFA-TXNIP) was re-expressed in TXNIP[KO] cells. Using confocal microscopy, we detected a fraction of ALFA-tagged TXNIP at the PM, where it partially co-localized with SLC3A2 [27,34] in proliferating cells, and 7 h after serum starvation (Figure 2e). 24 h after serum starvation, the majority of SLC3A2 was endocytosed and also ALFA-TXNIP was no longer detected at the PM (Figure 2e). Under all conditions, ALFA-TXNIP also localized prominently to the nucleus and to the cytosol, as previously observed [35] (Figure 2e).

TXNIP has been shown to be required for controlling cellular glucose uptake, by regulating clathrin mediated endocytosis of the glucose transporters SLC2A1 and SLC2A4 in response to glucose depletion or insulin signaling [26,27,36,37]. In WT and in TXNIP[KO] RPE1 cells entering quiescence, SLC2A1 remained at the PM (Supplementary Figure 2h) and the protein levels of SLC2A1 rather increased in response to serum starvation (Supplementary Figure 2i). Consistently, the uptake of the fluorescent glucose analog 2-NBDG (2-[N-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)amino]-2-deoxy-D-glucose) also increased in cell entering quiescence (Supplementary Figure 2j) [38]. In the TXNIP[KO] cells, 2-NBDG uptake was moderately elevated already in growing cells (Supplementary Figure 2j).

Our data suggested that TXNIP was required for the selective endocytic degradation of the heterodimer SLC7A5-SLC3A2 in cells entering quiescence. Since entering into quiescence did not trigger SLC2A1 endocytosis, it seemed that the TXNIP mediated downregulation of SLC7A5-SLC3A2 was not linked to TXNIP mediated SLC2A1 endocytosis.

Identification of a novel TXNIP loss-of-function mutation in a patient with an inborn metabolic disorder

TXNIP deficiency caused by a loss-of-function variant c.174_175delinsTT has been previously reported in three sibling with congenital lactic acidosis, low serum methionine levels, variable hypoglycaemia and other metabolic alterations [28]. During the course of the study, we identified a novel TXNIP loss-of-function mutation in another patient. The boy was born in 2014 and presented immediately after birth with recurrent hypoglycemia and muscular hypotonia in conjunction with lactic acidosis (Supplementary Table 1). Subsequently, he showed global developmental delay and marked intellectual disability with autistic features. In addition, he developed epileptic seizures with increasing frequencies of up to 3-4 per month (controlled with levetiracetame, topiramate and lamotrigine). With time, blood glucose concentrations stabilized (Supplementary Table 1), but the boy experienced recurrent hypoglycaemia in response to metabolic stress, e.g., infections. There were consistent metabolic alterations compatible with an AA transporter deficiency. Blood plasma concentrations of several large neutral amino acids (LNAAs, including L, I, V) were elevated throughout the years 2014 – 2022 (Supplementary Table 1). The increased molar ratio of the LNAAs (L, I, V) to aromatic AAs (F, Y), resulted in an elevated Fischer's ratio (FR, 2014: FR = 4.46; 2016: FR = 5.38, 2018: FR = 5.90; 2021; FR= 6.98; 2022: FR = 4.23; FR reference range = 2.10 - 4). The methionine levels are not dramatically altered (Supplementary Table 1).

Whole-exome sequencing from peripheral blood of the patient detected a homozygous single nucleotide insertion c.642_643insT in exon 5 of 8 of the *TXNIP* gene, which was not recorded in the population genetic variant database gnomAD that lists *TXNIP* as likely haplosufficient (pLI = 0, LOEUF = 0,709: https://gnomad.broadinstitute.org accessed Sept. 10, 2024). The variant caused a frameshift and a premature stop codon after 59 AA (denoted p.Ile215TyrfsTer59), likely causing nonsense-mediated decay (NMD) or the synthesis of a severely truncated TXNIP protein (Figure 3a). Both parents are healthy heterozygous carriers for the variant. No other (likely) pathogenic variant was identified as explanation of the clinical features in the child.

To examine how the loss-of-function mutation in TXNIP affected SLC7A5 downregulation, we generated primary patient-derived fibroblasts. WB analysis of the patient fibroblasts showed complete loss of the TXNIP protein (Figure 3b). In these cells, SLC7A5 was no longer downregulated in response to serum starvation (Figure 3b). Furthermore, indirect IF and confocal microscopy showed that SLC3A2 remained at the PM in response to serum starvation in the patient-derived fibroblasts with the TXNIP loss-of-function mutation, but not in the control primary fibroblasts (Figure 3c).

To directly test if the patient mutation was causative for the failure to downregulate SLC7A5-

SLC3A2, we expressed an ALFA-tagged version cDNA encoding TXNIP p.Ile215TyrfsTer59

(ALFA-patient[mut], or patient[mut]) in RPE1 TXNIP[KO] cells, under the control of a constitutive

eIF1α promoter. Under these conditions (which do not trigger NMD), the expression of a

truncated TXNIP protein was detected (Figure 3d, lane 3 and 4). ALFA-patient[mut] localized to

the PM, but failed to mediate SLC7A5-SLC3A2 endocytosis in response to serum starvation

(Figure 3d-f). Hence, even if the protein was expressed, the TXNIP patient[mut] protein was non-

functional.

It seemed that a loss-of-function mutation in TXNIP caused a severe metabolic disease,

inhibited the downregulation of SLC7A5-SLC3A2 in patient-derived fibroblasts upon entry

into quiescence and caused persistent changes in AA levels in the serum of the affected patient.

PPxY motifs of TXNIP were required for the endocytosis of SLC7A5

TXNIP contains two PPxY motifs in its C-terminal region (Figure 4a), which enable the

interaction with different HECT type ubiquitin ligases that contain a WW domain [15,19-22,39,40].

To test how the PPxY motifs of TXNIP contributed to SLC7A5-SLC3A2 endocytosis, we

introduced point mutations either in both PPxY motifs (TXNIP[PPCY331AACA, PPTY375AATA],

hereafter TXNIP[PPxYmut]), or only in the first PPxY motif (TXNIP[PPCY331AACA], hereafter

TXNIP[PPxY331]) and expressed these mutants in TXNIP[KO] RPE1 cells. The protein levels of the

PPxY mutants were higher compared to endogenous TXNIP (Figure 4b and Supplementary

Figure 3a), yet the endocytic downregulation of SLC7A5-SLC3A2 was blocked (Figure 4b and

Supplementary Figure 3a) and SLC3A2 remained at the PM in response to serum starvation

(Figure 4c, d and Supplementary Figure 3a - c).

A fraction of ALFA-TXNIP[PPxY331] localized to the PM in proliferating cells, similar to wildtype

ALFA-TXNIP. In response to serum starvation, the signal for ALFA-TXNIP[PPxY331] persisted at

the PM, where it co-localized with SLC3A2, while ALFA-TXNIP was cleared from the PM

together with SLC3A2 (Figure 4e). These results indicated that the endocytic downregulation

of SLC7A5-SLC3A2 required the PPCY motif at position 331 of TXNIP.

To examine the capacity of TXNIP and its PPxY motif to interact with different HECT-type

ubiquitin ligases, we co-expressed ALFA-TXNIP or ALFA-TXNIP[PPxY331] with YFP-NEDD4,

-ITCH, -WWP1, WWP2, and -HECW1 [41, 42] in HEK293T cells. A robust interaction of ALFA-

TXNIP with YFP-NEDD4, YFP-WWP1 and YFP-WWP2 was detected (Figure 4f and

Supplementary Figure 3d), which required the PPCY motif at position 331 in TXNIP (Figure

4f and Supplementary Figure 3d). The interaction of TXNIP appeared to be weaker with

HECW1, and the interaction with ITCH was barely detected (Supplementary Figure 3d). It

seemed that the PPCY motif at position 331 rendered TXNIP capable to interact with different

HECT type ubiquitin ligases.

To examine if entry into quiescence induced TXNIP mediated ubiquitination of SLC7A5, we

replaced endogenous SLC7A5 with C-terminally ALFA-tagged SLC7A5 (SLC7A5-ALFA).

Therefore, we generated SLC7A5^{KO} cells (using CRISPR/Cas9 mediated gene editing,

Supplementary Figure 3e), as well as SLC7A5^{KO} TXNIP^{KO} double knock out cells and stably

over-expressed SLC7A5-ALFA. To detect ubiquitination, SLC7A5-ALFA was

immunoprecipitated under denaturing conditions (1% SDS) before and after 7 h of serum

starvation. Polyubiquitination of SLC7A5-ALFA was detected under steady state conditions,

and appeared to slightly increase 7 h after serum starvation. The polyubiquitination of SLC7A5-

ALFA was markedly reduced in the TXNIP^{KO} cells (Figure 4g). Hence, TXNIP seemed to be

required for the efficient polyubiquitination of SLC7A5-ALFA.

Taken together, these findings suggested that TXNIP used its PPxY motif to interact with

HECT-type ubiquitin ligases for the efficient ubiquitination of SLC7A5-SLC3A2, leading to

endocytic downregulation.

AKT signaling controls TXNIP mediated downregulation of SLC7A5

Earlier work demonstrated that AKT directly phosphorylated TXNIP on serine 308 to inhibit

endocytosis of SLC2A1 and increase glucose uptake [26]. To examine if AKT also controlled the

endocytic downregulation of SLC7A5-SLC3A2, we treated RPE1 cells with the AKT inhibitor

MK2206 (MK). MK2206 treatment caused the endocytic downregulation of the SLC7A5-

SLC3A2 heterodimer (Figure 5a, lane 3 and Supplementary Figure 4a), but not of SLC1A5

(Supplementary Figure 4a). Inhibition of MEK with PD0325901 (PD) triggered the selective

endocytic downregulation of SLC1A5, but not of SLC7A5-SLC3A2 (Figure 5a, lane 4 and

Supplementary Figure 4a). These results suggested that distinct mitogenic signaling cascades

regulated the PM levels of different AA transporters, with AKT controlling SLC7A5-SLC3A2

endocytosis.

To test if AKT controlled SLC7A5 endocytosis via TXNIP, we treated RPE1 WT and TXNIP^{KO}

cells with MK2206. In the TXNIP^{KO} cells, endocytosis and degradation of SLC3A2 was

blocked in response to MK2206 treatment, suggesting that TXNIP functioned downstream of

AKT to mediate SLC7A5-SLC3A2 endocytic degradation (Figure 5b, c, lane 1 – 3 and 13 -

15). Consistently, the phospho-mimetic TXNIP^{S308D} mutant blocked SLC3A2 endocytosis in

response to serum starvation or AKT inhibition, while the non-phosphorylatable TXNIP^{S308A}

mutant allowed SLC7A5-SLC3A2 endocytic degradation (Figure 5b, c). These findings were corroborated by cell surface FACS analysis of SLC3A2 (Figure 5d and Supplementary Figure 4b).

TXNIP forms a disulfide bound (S-S) via cysteine 247 to inhibit TRX (thioredoxin) activity [23,24]. To test the role of the S-S between TXNIP and TRX in SLC7A5-SLC3A2 endocytosis, we generated mutant TXNIP^{C247S}, which no longer formed a S-S bridge with TRX [43]. The expression of TXNIP^{C247S} in TXNIP^{KO} cells rescued the endocytic degradation of SLC7A5-SLC3A2, similar to WT TXNIP (Figure 5b - d). Thus, SLC7A5 endocytosis was independent of the ability of TXNIP to form disulfide bridges with TRX.

These results demonstrated that AKT mediated phosphorylation of TXNIP at serine 308 inhibited SLC7A5-SLC3A2 endocytosis, while reduction of AKT signaling enabled TXNIP mediated SLC7A5-SLC3A2 endocytic degradation.

TXNIP mediated downregulation of SLC7A5 was essential to adjust intracellular AA levels and metabolic signaling in cells entering quiescence

Our results so far linked growth factor signaling via AKT with the TXNIP mediated control of AA acquisition via SLC7A5-SLC3A2. To directly test whether TXNIP mediated endocytosis of SLC7A5-SLC3A2 adjusted AA uptake in cells entering quiescence, we measured the cellular uptake of [^{3}H]-leucine. In response to serum starvation, WT cells downregulated [^{3}H]-leucine uptake, while TXNIP^{KO} cells did not reduce [^{3}H]-leucine uptake (Figure 6a). Upon JPH203 treatment, a selective inhibitor of SLC7A5 [44], the uptake of [^{3}H]-leucine was blocked in WT cells and TXNIP^{KO} cells (Figure 6a), suggesting that SLC7A5 was the main transporter for leucine in RPE1 cells. Consistently, in SLC7A5^{KO} cells, the uptake of [^{3}H]-leucine was strongly impaired (Supplementary Figure 5a).

To assess the effect of SLC7A5 downregulation or the lack thereof on intracellular free AA levels, we used quantitative mass spectrometry. We quantified free levels of 17 different proteinogenic AA in WT, TXNIP^{KO} and SLC7A5^{KO} RPE1 cells. In SLC7A5^{KO} cells the intracellular levels of L, I, W, V, F, Y were low under all conditions (Figure 6b), with exception for M, which was upregulated (Supplementary Figure 5b). These LNAAs are known substrates of SLC7A5 [1] (Figure 6b). In WT cells, the free levels of most these AAs (L, I, W, V, F, Y) decreased (Figure 6b). In contrast, the levels of L, I, W, V, F, Y were lower in growing TXNIP^{KO} cells, and they did not decrease when TXNIP^{KO} cells entered quiescence (Figure 6b). Neither loss of SLC7A5 nor loss of TXNIP affected the levels K, R, S, E, Q, P, G, A

(Supplementary Figure 5b, c). Taken together, these data indicated that the loss of TXNIP

affected the ability of cells to control SLC7A5 dependent AA transport.

To characterize the consequence resulting from the failure to adjust AA uptake in TXNIP[KO]

cells, we determined how mTORC1 dependent translational control was affected. In quiescent

WT cells, phosphorylation of S6K on threonine 389 was reduced compared to proliferating

cells, as was phosphorylation of the S6K substrate S6 on serine 240/244, in keeping with lower

mTORC1 activity (Figure 6c, lane 3 and 4, Figure 6d, e). In TXNIP[KO] cells, S6K and S6

phosphorylation also decreased, but remained higher compared to WT cells (Fig 6c, lane 1 and

2, Figure 6d, e), indicating elevated mTORC1 activity. Another important mode for

translational control is eIF2α phosphorylation at serine 51 by GCN2 (but also other kinases of

the integrated stress response) in response to lower AA levels. In quiescent WT cells,

eIF2α phosphorylation at serine 51 increased (Figure 6c, lane 3 and 4, Figure 6f). However, in

growing TXNIP[KO] cells, eIF2α phosphorylation appeared to be slightly elevated but did not

further increase, as these cells transitioned into quiescence (Figure 6c, lane 1 and 2, Figure 6f).

It seemed that TXNIP mediated SLC7A5-SLC3A2 endocytosis was required to decrease AA

uptake in quiescence, which helped to change signaling towards the translation machinery,

possibly to adapt translation.

In line with this concept, [^{35}S]-methionine and [^{35}S]-cysteine incorporation into newly

synthesized proteins remained higher in quiescent TXNIP[KO] cells, and also appeared to increase

faster after the re-addition of serum (Figure 6d). The increased translation in the TXNIP[KO] cells

required SLC7A5, because it was no longer observed in TXNIP[KO] SLC7A5[KO] double mutant

cells (Figure 6g).

We concluded that TXNIP mediated endocytosis of SLC7A5-SLC3A2 reduced the uptake of

LNAAs such as L, I, W, V, F, Y, which in turn lowered the free intracellular levels of these

AA, thereby helping cells to reduce mTORC1 signaling and to dampen translational activity.

TXNIP mediated endocytic degradation of SLC7A5 restrains cell proliferation

Finally, we determined how the TXNIP mediated regulation of SLC7A5 and AA uptake

contributed to cell cycle progression during entry into and exit from quiescence. Therefore, we

compared the DNA content of WT, TXNIP[KO] and TXNIP[KO] cells which re-expressed TXNIP

during entry and exit from serum starvation induced quiescence (Figure 7a).

In a growing cell population and 24 h after serum starvation, the DNA content of WT, TXNIP[KO]

and rescued cells was similar (Figure 7a, timepoint 0 h and 24 h). Interestingly, 24 h after serum

re-addition (timepoint 48 h), a larger fraction of the TXNIP[KO] cells (35%, ± 1%) had already

reached the G2/M phase of the cell cycle. In comparison, only 13% (± 10%) of WT cells or 17% (± 12%) of TXNIP^{KO} cells re-expressing TXNIP reached G2/M phase and more cells (35% ± 10% of WT cell and 25% ± 11% of rescued TXNIP^{KO}) were still in S-Phase (Figure 7a). Consistent with a faster cell cycle progression of TXNIP^{KO} cells during exit from quiescence, these cells resumed cell proliferation faster compared to WT cells, which was dependent on SLC7A5 (Figure 7b). Moreover, the overall cell proliferation rate of TXNIP^{KO} cells or of TXNIP^{KO} cells expressing TXNIP^{S308D} was increased compared to WT or TXNIP^{KO} cells expressing TXNIP or TXNIP^{S308A} (Figure 7c). Again, the increased proliferation of TXNIP^{KO} cells was dependent on SLC7A5, since the double knock out TXNIP^{KO} SLC7A5^{KO} grew slower, similar to SLC7A5^{KO} cells (Figure 7c).

Taken together, our results demonstrated that TXNIP interacted with HECT-type ubiquitin ligases to selectively degrade SLC7A5-SLC3A2 in cells entering quiescence, thereby reducing AA uptake. This process not only contributed to reduce translation, but also restrained cell cycle progression during the exit from quiescence and limited overall cell proliferation (see Figure 7d model). This mechanism was only licensed upon serum removal and repressed by growth factor signaling through AKT.

Discussion

Our work establishes a central role for TXNIP in a functional feedback loop that links growth factor signaling, AA homeostasis and metabolic signaling for translational control during cellular transitions between proliferation and quiescence, with potential relevance for human health. In particular, we made progress toward understanding how growth signaling regulates TXNIP mediated endocytosis of SLC7A5-SLC3A2 and hence, the acquisition of LNAAs for metabolic scaling in growing and quiescent cells. The framework of our model (Figure 7d) is based on the following results: (1) Human cells that enter quiescence downregulate AA uptake, in part, by selective endocytosis of the AA transporters; (2) the α-arrestin TXNIP mediates the selective endocytosis of SLC7A5-SLC3A2; (3) TXNIP mediated endocytosis is suppressed in growing cells by AKT signaling downstream of growth factor receptors and (4) TXNIP mediated SLC7A5-SLC3A2 endocytosis helps to lower AA uptake, translation and thereby restrains cell proliferation. (5) These processes contribute to metabolic homeostasis in humans. A novel loss-of-function mutation in TXNIP, that impairs AA endocytosis, causes a rare metabolic disease.

Together with a previous report on three siblings from a single consanguineous family that were homozygous for a loss-of-function frameshift variant c.174_175delinsTT [28], our data suggest

TXNIP deficiency as a novel inherited metabolic disease with specific metabolic alterations,
that appear associated with neonatal lactic acidosis, recurrent hypoglycaemia, and specific
anomalies in AA metabolism, although full the spectrum of associated clinical manifestations
remains uncertain. Similar to our patient, the three siblings presented with lactic acidosis in the
neonatal period, which was treated with long-term dichloroacetic acid (DCA). Rather than
directly clearing lactate from serum, DCA inhibits pyruvate dehydrogenase kinase to stimulate
pyruvate dehydrogenase activity and oxidative decarboxylation of pyruvate, resulting in
reduced serum lactate levels [45 46]. Two of the three affected children had normal development
during childhood; one child, which also suffered from recurrent hypoglycaemia, had neonatal
muscular hypotonia and failure to thrive, and was under investigation for autism spectrum
disorder, similar to our patient. Our patient additionally has an increasing frequency of epileptic
seizures. The three siblings developed low plasma methionine concentrations. No other AA
anomalies were specified in the previous report except for a transient general
hyperaminoaciduria in one child [28]. The detailed metabolic investigations in our case revealed
relative normal methionine levels, but identified other specific AA alterations in line with
impaired SLC7A5 endocytosis in the case of complete TXNIP loss. How these metabolic
defects are linked to other clinical manifestations such as possible autism and the severe
epileptic developmental disorder in our patient, remains to be investigated. We speculate that
these feature could be related to role of SLC7A5 in neurons and in the blood brain barrier [47,48].
Yet, the patient phenotype appears to be variable in severity and not fully penetrant. Hence,
loss of TXNIP alone is likely to be insufficient and other genetic or non-genetic factors (e.g.
diet) that are not yet known, must be involved.

Several studies in mice show that TXNIP is important during the transition between fed and
fasted states, which was attributed to its role in glucose and energy metabolism [49-52]. Our
findings imply that at least some of the described TXNIP dependent effects on glucose
homeostasis might also be linked to TXNIP dependent regulation of SLC7A5-SLC3A2 and the
regulated import of LNAAs as cells enter and exit quiescence.

While it is clear how proliferating cells benefit from the upregulation of SLC7A5-SLC3A2, it
is less clear how quiescent cells could profit from the downregulation of SLC7A5-SLC3A2.
One rational for the selective endocytosis of this AA transporter could be that the cellular
demand for essential AAs must be matched to lower rates of protein translation in quiescent
cells. In line with this concept, also quiescent mammary epithelial cells reduced levels of V, L,
I and Q [53]. Similar results were obtained in a non-malignant murine pro-B lymphocyte cell line
(FL5.12), in which the abundance of SLC3A2 at the plasma membrane was regulated by the

presence of growth factors [54]. In those cells, the consumption rate of essential AAs was
decreased in G0 arrest and increased upon re-entry into the cell cycle [55]. Hence, the
downregulation of AA transporters provide means to avoid an excess/imbalance of free
intracellular AAs which facilitates the controlled decrease of mTORC1 signaling and the
ensuing reduction of translation rates, possibly in conjugation with the integrated stress
response. Thereby, the TXNIP mediated selective downregulation of SLC7A5-SLC3A2 and
the ensuing decreased import of essential AAs, helps to promote the switch from anabolic to
catabolic metabolism. Moreover, the downregulation of AA transporters upon entry into
quiescence might also contribute to the barrier for cellular transformation [5 56-58].

TXNIP was specifically required for the endocytosis of SLC7A5-SLC3A2, but not for the
endocytosis of SLC7A11-SLC3A2. How TXNIP interacts specifically with SLC7A5-SLC3A2
remains unclear at the moment. Yet it is clear that TXNIP requires its first PPCY motif (331)
to interact with several different HECT type ubiquitin ligases, and this interaction is required
for the downregulation of SLC7A5. In line with our results, SLC7A5 was shown to be
ubiquitinated on N-terminal lysine residues by Nedd4-2 upon protein kinase C (PKC) activation
and mTORC1 inhibition [59,60].

We conclude that the role of TXNIP in controlling the abundance of SLC7A5-SLC3A2 at the
PM is important to match AA uptake with metabolic needs in proliferating and quiescent cells.
In the future it will be important to understand how these cellular functions of TXNIP and the
loss thereof translate into complex metabolic phenotypes of human patients with loss of TXNIP
variants.
490

Acknowledgments

We thank the patient and his family. We are grateful to Hemmo Meyer and Simona Polo for providing the YFP-tagged HECT type ubiquitin ligases. This research was funded in part by the Austrian Science Fund (FWF) (10.55776/P35874, 10.55776/P34907, 10.55776/FG20 to DT, 10.55776/P35832, 10.55776/P36600 to HF, 10.55776/P36925 to VR, 10.55776/P30196 to SH, and 10.55776/DOC82 to FH, DT, LAH, KT and MA). JK is a recipient of a DOC Fellowship of the Austrian Academy of Sciences. KT acknowledges support from the DFG (German Research Foundation, project No TH 1358/3e2), the MESI-STRAT project (grant agreement No 754688) which has received funding from the European Union's Horizon 2020 research and innovation programme, and from the European Union European Research Council (ERC AdG BEYOND STRESS, grant agreement No 101054429) which has received funding from the European Union's Horizon Europe research and innovation programme. For open access purposes, the author has applied a CC BY public copyright license to any author accepted manuscript version arising from this submission.

Material and methods

Cell culture

Patient-derived primary fibroblasts were cultured from skin biopsies. Control fibroblasts were obtained from a male, born in 2000 (WT#1) and from a male, born in 2015 (WT#2). RPE1, HEK293T, A549, H460, HCC4006, primary human fibroblasts and MEF (mouse embryonic fibroblasts) [61] were cultured in DMEM containing high glucose (4.5 g/L) (Sigma-Aldrich, #D6429), supplemented with 1% (vol/vol) penicillin/streptomycin (Sigma-Aldrich, #P0781) and 10% (vol/vol) FBS (fetal bovine serum) (Sigma-Aldrich, #S0615) at 37°C, in 5% CO_2 and 98% humidity. H2228, H226, H520 and H1299 cells were cultured in RPMI 1640, glutamax supplement medium (Thermo Fisher, Gibco, # 61870044) supplemented with 1% (vol/vol) penicillin/streptomycin (Sigma-Aldrich, #P0781) and 10% (vol/vol) FBS (Sigma-Aldrich, #S0615) under the same culture conditions. Cells were passaged before reaching confluency by incubation in 1x PBS (phosphate-buffered saline, 8 g/L NaCl, 0.2 g/L KCl, 1.15 g/L Na_2HPO_4, 0.2 g/L KH_2PO_4) and 1x trypsin-EDTA solution (Sigma-Aldrich, #T4174). Cells were resuspended in growth medium and re-seeded at a 1:5 to 1:10 ratio. Cells were regularly tested negative for mycoplasma.

The AKT inhibitor MK2206 (Enzo Life Sciences, #ENZ-CHM164-0005), dissolved in DMSO (Sigma-Aldrich, #D2650), was used at a final concentration of 1 μM. The MEK inhibitor

PD0325901 (Absource, #S1036), dissolved in DMSO, was used at a final concentration of 5

μM. The SLC7A5 inhibitor JPH203 (also KYT-0353, Selleckchem, #S8667), dissolved in

DMSO, was used at a final concentration of 10 μM. Chloroquine (Sigma-Aldrich, #C6628) was

dissolved in Millipore H_2O and used at a final concentration of 12.5 μM. To inhibit dynamin

mediated endocytosis, dynasore hydrate (Sigma-Aldrich, #D7693) was used as previously

described [31] at a final concentration of 20 μM.

To maintain cells in proliferation (control cells, defined as "+ serum"), they were cultured in

DMEM containing 4.5 g/L glucose and all AAs, (Sigma-Aldrich, #D6429), supplemented with

1% (vol/vol) penicillin/streptomycin (Sigma-Aldrich, #P0781) and 10% (vol/vol) FBS (Sigma-

Aldrich, #S0615). To induce quiescence, cells were washed with 1x PBS and cultured in

DMEM containing 4.5 g/L glucose and all AAs, supplemented with 1% penicillin/streptomycin

but without FBS for 24 h (referred to as serum starvation, "- serum").

Proliferation assay

A similar cell number (>20000 cells) was seeded onto 6-well plates (day 0). Cells were

harvested daily by trypsinization. The cell number was measured daily using a CASY Cell

Counter and Analyzer (OMNI Life Science) for 4 days. The cell proliferation index was

calculated as described before [62]. The fold increase was calculated by normalizing to the cell

number obtained 24 h after seeding (day 1).

For measuring cell proliferation in an automated manner, 50000 cells were seeded onto a 6-

well plate and cultured at standard cell culture conditions. Cells were set on serum free medium

24 h after seeding and refed after 48 h of serum starvation. Cell proliferation was tracked using

a Sartorius IncuCyte® S3 Live Cell Imaging System and confluency measurement was

performed every 4 h for 5 days. Data were acquired and analyzed using the Sartorius IncuCyte®

S3 software version 2022B.

Preparation of whole cell protein extracts, SDS-PAGE and Western Blot analysis

Cells were washed in 1x cold PBS, harvested using a cell scraper and pelleted at 13.000 rpm

for 5 min at 4°C. The cell pellets were resuspended in cold RIPA lysis buffer (50 mM Tris-HCl

pH 7.5, 150 mM NaCl, 1% NP-40, 2 mM EDTA pH 8, 0.1% SDS, 0.5% sodium deoxycholate)

or lysis buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1% Triton X-100, 10% glycerol, 0.5

mM EDTA), supplemented with 50 mM NaF, 10 μg/mL leupeptin, 1 mM pefablock, 1 μg/mL

pepstatin, 10 μg/mL aprotinin, 2 mM Na_3VO_4. Cells were lysed for 30 min on ice, centrifuged at 13.000 rpm for 10 min at 4°C and the cleared lysate was obtained. Protein concentration was measured using Micro BCA Protein Assay Kit (Thermo Scientific, #23235) according to the manufacturer's instructions. 20 μg protein was supplemented with 5x Urea sample buffer (8 M Urea, 5% SDS, 40 mM Tris pH 6.8, 0.1 mM EDTA, 0.4 mg/mL bromphenol blue, 1% final β-mercaptoethanol) at 1x final concentration. Lysates were separated by SDS-PAGE. Proteins were transferred onto polyvinylidene fluoride (PVDF) membranes (Amersham Hybond PVDF, #10600023 – 0.45 μm) using a wet transfer system (Bio-Rad) at constant 80V for 2 h (1x transfer buffer: 20% methanol, 25 mM Tris-HCl, 192 mM glycine). Membranes were blocked in 5% milk in Tris-buffered saline buffer (TBS, 0.5 M NaCl, 20 mM Tris, pH 7.5) with 0.05% Tween 20 (TBS-T) and probed with the respective antibodies over night at 4°C. The membranes were washed with TBS-T and probed with the respective horseradish peroxidase-linked secondary antibodies (1:5000 dilution) for 1 h at RT. After washing with TBS-T, the membrane was incubated with WesternBright Chemiluminescence substrate solution (Advansta) (Biozym, #541005X). The chemiluminescence signal was detected either on X-ray films or with a CCD-Camera (Fusion FX, Vilber Lourmat). The signal was quantified with ImageJ and normalized to GAPDH. To quantify the phosphorylation of proteins, the signals were normalized to the unphosphorylated total protein levels.

Denaturing SLC7A5-ALFA immunoprecipitation

Immunoprecipitation of SLC7A5-ALFA under denaturing lysis conditions was adopted from [63]. Cells expressing SLC7A5-ALFA were either grown in growth medium (+ serum) or serum starved for 7 or 8 h (- serum). Cells were washed with 1x PBS and harvested. Cell pellets were resuspended in lysis buffer I (50 mM Tris pH 8, 150 mM NaCl, 50 mM NaF, 5 mM EDTA, 1% SDS, 10 nM NEM, 10 μg/mL leupeptin, 1 mM pefablock, 1 μg/mL pepstatin, 10 μg/mL aprotinin) and incubated 30 min at RT. Protein concentration was measured using Micro BCA Protein Assay Kit (Thermo Scientific, #23235) according to the manufacturer's instructions. Equal amounts of proteins were diluted by addition of 800 μl lysis buffer II (50 mM Tris pH 8, 150 mM NaCl, 50 mM NaF, 5 mM EDTA, 1% Triton X-100, 10 nM NEM, 10 μg/mL leupeptin, 1 mM pefablock, 1 μg/mL pepstatin, 10 μg/mL aprotinin). Samples were subjected to immunoprecipitation. Beads (ALFA Selector ST, NanoTag Biotechnologies, #N1516) were equilibrate in washing buffer I (50 mM Tris pH 8, 150 mM NaCl, 1% Triton X-100). Cell suspension and beads were incubated rotating at 4°C for 3 h. Beads were recovered with a magnetic rack and washed twice with washing buffer I and three times with washing buffer II

(50 mM Tris pH 8, 300 mM NaCl, 1% Triton X-100). Proteins were eluted with 5x SDS sample

buffer (250 mM Tris-HCl pH 6.8, 10% SDS, 50% glycerol, 10% β-mercaptoethanol) and

denatured for 10 min at 95°C. For detection of ubiquitinated proteins, samples were separated

by SDS-PAGE. Proteins were transferred onto PVDF membranes (Amersham Hybond PVDF,

#10600023 – 0.45 µM (300mm)) using a wet transfer system (Bio-Rad) at constant 80V for 2

h. After transfer, the membrane was blocked with 10% BSA in 0.45% Tween 20 for at least 1

h and incubated with anti-ubiquitin antibody (Santa Cruz, ##3936S) overnight in 10% BSA

blocking solution.

603   Non-denaturing ALFA-TXNIP immunoprecipitation

Non-denaturing immunoprecipitation of ALFA-TXNIP was adopted from [27]. Briefly, cells were

washed with 1x PBS and harvested by scraping. Cells were lysed in lysis buffer (30 mM Tris

pH 7.5, 120 mM NaCl, 20 mM NaF, 0.5% octyl-β-D-glucopyranosid) (Sigma-Aldrich,

#O8001) and protease inhibitors (10 µg/mL leupeptin, 1 mM pefablock, 1 µg/mL pepstatin, 10

µg/mL aprotinin, 2 mM Na_3VO_4). Cells were lysed for 30 min on ice, centrifuged at 13.000

rpm, at 4°C for 10 min and cleared lysate was obtained. Protein concentration was measured

using Micro BCA Protein Assay Kit (Thermo Scientific, #23235) according to the

manufacturer's instructions. Equal amounts of proteins were subjected to immunoprecipitation.

Beads (ALFA Selector ST, NanoTag Biotechnologies, #N1516) were equilibrated in lysis

buffer. Cell suspension and beads were incubated rotating at 4°C for 3 h. After incubation beads

were washed three times for 10 minutes at 4 °C using wash buffer (30 mM Tris pH 7.5, 150

mM NaCl, 0.1% octyl-β-D-glucopyranosid). Proteins were eluted in 100µl 2x Urea sample

buffer at 42°C for 30 min.

618   Genetic modifications and cloning

Genetic modifications were performed by PCR using standard techniques or by using the

Gateway cloning method (Thermo Fisher Scientific) according to manufacturer's instructions.

All plasmids and primers used in this study are listed in Supplementary Table 3 and

Supplementary Table 4. The sequence-verified plasmids were then used for lentiviral

transduction of RPE1 cells. To generate virus particles, HEK293T cells were seeded in a 6-well

tissue culture plate and transfected with 2 µg lentiviral vector containing the gene of interest

together with virus packing vectors (1 µg pVSV-G and 1 µg psPAX2) using polyethylenimine

(PEI, Polyscience, #23966-100) as transfection reagent. 4 - 8 h after transfection, the medium

was changed. Virus particle-containing medium was collected from the HEK293T cells 48 h

and 72 h after transfection and was used to infect RPE1 cells. Polybrene was added to the virus-particle-containing medium to help lentivirus integration (1:500 dilution, Sigma-Aldrich, #107689). After 10 days of selection with 1 µg/mL blasticidin (InvivoGen, #ant-bl-5b), the efficacy of the virus transduction was tested via SDS-PAGE and Western Blot analysis.

Genetic deletion using CRISPR/Cas9 mediated gene editing was performed as previously described [64]. For CRISPR/Cas9-mediated depletion, guide RNA (gRNA) targeting primers for SLC7A5 (5'-caccgCGTGAACTGCTACAGCGTGA -3' and 5'-aaac TCACGCTGTAGCAGTTCACGc -3') and TXNIP (5'-caccgGTAAGTGTGGCGGGCCACAA-3' and 5'-aaacTTGTGGCCCGCCACACTTACc-3'). RPE1 cells were transiently transfected with the gRNA-containing pSpCas9(BB)-2A-GFP plasmid (Addgene, #48138) using Lipofectamin LTX reagent (Thermo Scientific, #15338-100) according to the manufacturer's protocol. 24 h after transfection, GFP-positive, transfected cells were enriched by flow cytometry (BD FACSAria III Cell Sorter, BD Biosciences). The sorted cell suspension was used for the generation of single cell clones. Depletion efficiency was verified by PCR using specific primers and Western Blot analysis.

For the transient transfection of HEK293T Lipofectamine LTX & Plus reagent (Thermo Scientifc, #15338100) was used according to the manufacturer's protocol. HEK293T cells were seeded in a 10 cm dish. At around 80% confluency, 16 µg of the respective plasmids were diluted in 500 µl of OptiMEM (Thermo Scientific, #31985062) in the presence of 48 µl of LTX- and 16 µl of Plus-reagent for the transfection. The transfected cells were incubated for 4 h before the transfection medium was discarded and replaced with 10 ml of fresh DMEM. After that, cells were incubated for 24 h under standard conditions.

Immunofluorescence microscopy

Cells were grown on 12 mm glass coverslips (Hartenstein GmbH, #DKR1) and fixed with 4% paraformaldehyde (PFA) for 10 - 15 min at RT. Immunofluorescence staining of SLC3A2 and ALFA-tagged proteins was performed as described previously [65]. Briefly, cells were permeabilized and blocked in blocking buffer (10% FBS, 0.3% saponin (BioChemika, #84510), sodium azide (0.02% final) in 1x PBS) for 30 min at 37°C. Cells were incubated with mouse anti-SLC3A2 (BD Pharmingen, #556074) or rabbit anti-ALFA (NanoTag Biotechnologies, #N1581) in a 1:100 dilution in blocking buffer overnight at 4°C. For double-staining with LAMP1, cells were simultaneously incubated with rabbit anti-LAMP1 (Cell Signaling, #9091) in a 1:800 dilution in blocking buffer. Cells were washed once with wash solution (0.03%

saponin, sodium azide (0.02% final) in 1x PBS), following three washing steps with PBS and incubated with Alexa Fluor 488 goat anti-mouse IgG (Invitrogen, #A11001) and Alexa Fluor 568 goat anti-rabbit IgG (Invitrogen, #A11011) in a 1:500 dilution in blocking buffer for 1 h at RT.

For immunofluorescence labeling of SLC1A5, cells were permeabilized in in 0.2% Triton X-100 (Thermo Scientific, #28314) in PBS for 5 min and blocked in blocking buffer (2% gelatin, 150 mM NaCl, 10 mM Pipes pH 6.8, 5 mM EGTA, 5 mM glucose, 5 mM $MgCl_2$, 50 mM NH_4Cl) for 1 h at RT. Cells were incubated with rabbit anti-SLC1A5 (Sigma-Aldrich, #HPA035240) in a 1:50 dilution in blocking buffer for 1 h at RT. For double-staining with LAMP1, cells were simultaneously incubated with mouse anti-LAMP1 (DSHB, #H4A3-S) in a 1:50 dilution in blocking buffer. Cells were washed with PBS and incubated with Alexa Fluor 488 goat anti-rabbit IgG (Invitrogen, #A11008) and Alexa Fluor 568 goat anti-mouse IgG (Invitrogen, #A11031) in a 1:500 dilution in blocking buffer for 1 h at RT.

For immunofluorescence labeling of SLC2A1, cells were permeabilized and blocked in blocking buffer (2% gelatin, 150 mM NaCl, 10 mM Pipes pH 6.8, 5 mM EGTA, 5 mM glucose, 5 mM $MgCl_2$, 50 mM NH_4Cl) containing 0.025% saponin (BioChemika, #84510) for 1 h at RT. Cells were incubated with rabbit anti-SLC2A1 (Merck, #07-1401) in a 1:200 dilution in blocking buffer for 1 h at RT. Cells were washed with PBS and incubated with Alexa Fluor 488 goat anti-rabbit IgG (Invitrogen, #A11008) in a 1:500 dilution in blocking buffer for 1 h at RT. Cells were washed with 1x PBS, incubated with DAPI (Sigma-Aldrich, #D9542) in a 1:20000 dilution in 1x PBS and mounted in Mowiol (Sigma-Aldrich, #81381).

Confocal microscopy was performed on a LSM980 AiryScan 2 (ZEISS). Confocal Z-series stack (step size 0.22 μm) images were acquired using the AiryScan detector, a LD LCI Plan-Apochromat 63x / 1.2 with glycerol immersion objective (ZEISS) and the 405 nm, 488 nm and 561 nm laser lines. To compare signals of a protein of interest in proliferating, quiescent or treated cells, laser settings were kept constant while imaging both conditions. The ZEISS Zen 3.5 software was used as recording software and for processing of the raw images (AiryScan processing). Brightness and contrast were adjusted using ImageJ software. For merged images, the levels of red, green and blue channels were separately adjusted.

Cell surface staining of SLC3A2 and FACS analysis

To measure the difference in surface staining of SLC3A2 between proliferating and quiescent cells, cells were either grown in growth medium or serum starved for 24 h before the experiment. The staining was performed as previously described [65]. Cells were washed with PBS, trypsinized and resuspended in FACS buffer (10% FBS in 1x PBS) containing anti-SLC3A2 antibody (BD Pharmingen, #556074) at a dilution of 1:100. The antibody was incubated for 30 min on ice. Cells were centrifuged at 1000 rpm for 5 min at 4°C and resuspended in FACS buffer containing Alexa Fluor 488 goat anti-mouse IgG (Invitrogen, #A11001) at a dilution of 1:500. The antibody was incubated for 30 min on ice in the dark. Cells were centrifuged at 1000 rpm for 5 min at 4°C and resuspended in 200µl FACS buffer. The fluorescence intensity was measured by flow cytometry (Attune NxT Flow Cytometer, Life Technologies). For every experiment, more than 3500 cells were analyzed. Flow cytometry analysis was performed with FlowJo software 10.8.1 (BD Life Sciences). Cells were gated by forward versus side scatter (FSC-A vs. SSC-A) to identify cell population, forward scatter width versus forward scatter area (FSC-W vs FSC-A) for doublet exclusion and fluorescence intensity (BL1-A) vs cell count to reflect Alexa-488 signal populations. Unstained cells were excluded from analysis.

DNA content for cell cycle analysis of fixed cells stained with propidium iodide (PI)

Cells were washed with PBS and harvested by trypsinization. Cells were centrifuged at 1000 rpm for 5 min at 4°C. The pellet was washed in cold PBS. Cells were fixed by resuspending the cells in 1 ml ice cold 70% vol/vol ethanol in PBS. Samples were stored at -20°C for at least 30 min. The ethanol was washed away with ice cold PBS twice. Cells were then incubated with propidium iodide (PI) at a final dilution of 1:25 and RNAse A (Thermo Scientific, #EN0531) at a final dilution of 1:100 in PBS, for 30 min at 37°C. The fluorescence intensity was measured by flow cytometry (Attune NxT Flow Cytometer, Life Technologies). For every experiment, more than 1500 cells were analyzed. Data were analyzed using ModFit LT (Verity Software House). Cells were gated by forward versus side scatter (FSC-A vs. SSC-A) to identify cell population. PI-positive cells were gated for doublet exclusion by width vs height (BL3-H vs BL3-W) and fluorescence intensity (BL3-A) vs cell count to reflect PI-positive signal populations.

RNA was extracted from cells using the RNeasy Mini Kit (Quiagen, #74104) according to the manufacturer's instructions. 800 ng RNA were reverse transcribed into cDNA using the LunaScript RT Super Mix Kit (NEB, #E3010) according to manufacturer's protocol. For

mRNA fold regulation analysis, quantitative real-time PCR was performed using gene specific primers listed in Supplementary Table 5. The Biozym Blue S'Green qPCR Kit (Biozym, #F-415S) was used according to manufacturer's protocol. All primers were synthesized by Sigma-Aldrich and diluted in Millipore H_2O to a concentration of 10 µM. Each condition was performed in technical triplicates from three biological replicates. Fold regulation was calculated using the $2^{-\Delta\Delta Ct}$ method and normalized to the housekeeping gene *ribosomal protein lateral stalk subunit P0 (RPLP0)*.

Uptake of radioactive labeled AAs

To analyze the uptake of L-glutamine, cells were washed with PBS and incubated with 0.3 µCi/mL [$^{14}C_5$]-L-glutamine (Hartmann Analytic, #MC1124) in glutamine-free medium (Sigma-Aldrich, #D6546). To analyze the uptake of L-leucine, cells were washed with PBS and incubated with 0.5 µCi/mL [4,5-^{3}H]-L-Leucine (Hartmann Analytic, #MT672E) in AA free medium (USBiological, #D9811-01). The uptake of L-isoleucine was initiated by incubating the cells with 0.005 µCi/mL [$^{14}C_5$]-L-isoleucine (Hartmann Analytic; #MC174) in AA free medium. The uptake was performed for 15 min at 37°C. To stop the reaction, cells were washed twice with ice cold PBS. Cells were harvested by trypsinization and the cell pellet was lysed in 100 µl lysis buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1% NP-40, 2 mM EDTA pH 8, 0.1% SDS, 0.5% sodium deoxycholate) for 30 min on ice. Cells debris were removed by centrifugation (13000 rpm, 4°C, 10 min) and the supernatant was analyzed by liquid scintillation counting (LSC-Universal cocktail, Roth, #0016.3; HITACHI AccuFlex LSC-800). AA uptake was calculated as counts per min (cpm) normalized to the protein concentration (µg) of each cleared lysate. Protein concentration was determined by using Micro BCA Protein Assay Kit (Thermo Scientific, #23235) according to the manufacturer's instructions.

Glucose uptake assay

The glucose uptake assay was performed as described previously [66]. Briefly, glucose uptake was initiated by adding 2-NBDG (2-(N-(7-Nitrobenz-2-oxa-1,3-diazol-4-yl)amino)-2-deoxyglucose) (AAT Bioquest, #36702, dissolved in DMSO) at a final concentration of 100 µM in the respective medium. To inhibit glucose uptake, cells were simultaneously treated with 1 mM phloretin (Sigma-Aldrich, #P7912, dissolved in 50% ethanol) for 45 min. After 45 min incubation at 37°C in the dark, cells were washed with ice cold PBS to stop the reaction, trypsinized and resuspended in PBS. Cells were centrifuged at 1000 rpm for 5 min at 4°C and the pellet was resuspended in 200 µl PBS and 10% FBS. The fluorescence intensity was

measured by flow cytometry (Attune NxT Flow Cytometer, Life Technologies). For every experiment, more than 5000 cells were analyzed. Flow cytometry analysis was performed with FlowJo software 10.8.1 (BD Life Sciences). Cells were gated by forward versus side scatter (FSC-A vs. SSC-A) to identify cell population, forward scatter width versus forward scatter area (FSC-W vs FSC-A) for doublet exclusion and fluorescence intensity (BL1-A) vs cell count to reflect the uptake of 2-NBDG.

Incorporation of ^{35}S labeled L-Cysteine and L-Methionine

150000 cells were seeded and after two days, these cells were left untreated or serum starved for 24 h or serum starved and then refeed for 24 h. Prior to radiolabeling the cells, the serum starved cells were washed with PBS and incubated for 30 min in L-cysteine, L-methionine, L-glutamine free medium (Sigma-Aldrich #D0422), supplemented with 2 mM L-glutamine (Gibco #25030-024) and 1% (vol/vol) penicillin/streptomycin. The proliferating cells were incubated for 30 min in L-cysteine, L-methionine, L-glutamine free medium, supplemented with 2 mM L-glutamine, 10% dialysed FBS (Thermo Fisher Scientific #26400044) and 1% (vol/vol) penicillin/streptomycin. Cells were washed with PBS, brought into suspension using 1x trypsin-EDTA solution and resuspended in the corresponding L-cysteine, L-methionine free medium. 17.6 µCi/mL [^{35}S]-methionine [^{35}S]-cysteine (Hartmann Analytic #21635410) was added to the cell suspension for 2 min and quenched with 10 µg/ml cycloheximide. Cell pellets were lysed for 30 min on ice, with RIPA lysis buffer, supplemented with protease inhibitors and 10 µg/ml cycloheximide. Equal amounts of protein were subjected to SDS-PAGE, (de)stained with Coomassie and dried for 1 h at 70°C. The dried gel was exposed to a phosphor-imager for 48 h. Detection of a [^{35}S]-methionine [^{35}S]-cysteine radiogram was performed using a Typhoon FLA 9500 detection system.

Mass spectrometry analysis of intracellular free AAs

The analysis of intracellular AAs by mass spectrometry was performed as described previously [67][68]. Briefly, 10^6 cells were seeded and serum starved for 24 h prior the extraction. Then, cells were washed three times with ice cold 1xPBS and extracted by adding 500 µl ice cold methanol (99,9% vol/vol, Fisher Chemicals) and 500 µl cold Millipore H_2O, containing the internal standard (1:100), a stable isotope-labeled canonical AA mix composition (Cambridge Isotope Laboratory, #MSK-CAA-1). Cells were scraped and 500 µl ice cold chloroform (99,9% vol/vol, Sigma-Aldrich, #650498) was added to the cell extracts. Cells were agitated in a tube shaker for 20 min at 1400 rpm at 4°C, followed by centrifugation for 5 min at 16100xg at 4°C. The

upper, polar phase was separated, evaporated by using a SpeedVac (Savant SPD111V SpeedVac Concentrator, Thermo Scientific) and stored at -80°C. The interphase was used to determine the protein concentration. Therefore, the interphase was evaporated and the pellet was resuspended in denaturing buffer (1.6 M Urea, 100 mM AmBiCa, pH 8.2) and sonicated for 10 sec to break up protein aggregates. Protein determination was done by using Micro BCA Protein Assay Kit (Thermo Scientific, #23235) according to the manufacturer's instructions. For absolute quantification, AAs were analyzed by targeted LC-MS using mixed-mode chromatography (RP-AEX, reversed phase–anion exchange chromatography) and ultra-high sensitive orbitrap technology (Orbitrap Q Exactive HF-X and Orbitrap Fusion Lumos, both Thermo Scientific), operating in parallel reaction monitoring (PRM) mode. Bioinformatics data processing and statistical analysis was performed using TraceFinder (Thermo Scientific) and in-house R and Python scripts.

Statistical analysis and software

Software used in this study, if not already specified, were Affinity Designer Version 1.10.5 (Serfi, RRID:SCR_016952), Image J Version 1.53t (RRID:SCR_003070), GraphPad Prism 9 Version 9.4.1 (458) (RRID:SCR_002798), Benchling (RRID:SCR_013955), FlowJo Version 10.8.1 (RRID:SCR_008520), ModFit LT (RRRID:SCR_016106), Primer3 Version 4.1.0 (RRID:SCR_003139), PrimerBank (RRID:SCR_006898), CHOPCHOP (RRID:SCR_015723), ZEISS ZEN Digital Imaging for Light Microscopy Version 3.5 (RRID:SCR_013672), and online tools TIDE (tracking of indels by Decomposition) (Brinkman et al. 2014), ICE Analysis online tool (Synthego, v2.0; Synthego, USA) and PrimerQuest™ Tool (Integrated DNA Technologies, IDT, USA)

Results were represented as mean with standard deviation (SD) analyzed by GraphPad Prism. Statistical significance between different biological samples was determined by either paired Student's t test or one-way/ two-way ANOVA with Tukey test for multiple comparisons, if not described otherwise.

References

Kandasamy, P., Gyimesi, G., Kanai, Y. & Hediger, M. A. Amino acid transporters revisited: New views in health and disease. *Trends in biochemical sciences* **43**, 752-789, doi:10.1016/j.tibs.2018.05.003 (2018).

Chidley, C. *et al.* A CRISPRi/a screening platform to study cellular nutrient transport in diverse microenvironments. *Nat Cell Biol* **26**, 825-838, doi:10.1038/s41556-024-01402-1 (2024).

White, P. J. *et al.* Insulin action, type 2 diabetes, and branched-chain amino acids: A two-way street. *Mol Metab* **52**, 101261, doi:10.1016/j.molmet.2021.101261 (2021).

Errasti-Murugarren, E. & Palacin, M. Heteromeric Amino Acid Transporters in Brain: from Physiology to Pathology. *Neurochem Res* **47**, 23-36, doi:10.1007/s11064-021-03261-w (2022).

Palm, W. & Thompson, C. B. Nutrient acquisition strategies of mammalian cells. *Nature* **546**, 234-242, doi:10.1038/nature22379 (2017).

Hansen, C. G., Ng, Y. L., Lam, W. L., Plouffe, S. W. & Guan, K. L. The Hippo pathway effectors YAP and TAZ promote cell growth by modulating amino acid signaling to mTORC1. *Cell Res* **25**, 1299-1313, doi:10.1038/cr.2015.140 (2015).

Hayashi, K., Jutabha, P., Endou, H. & Anzai, N. c-Myc is crucial for the expression of LAT1 in MIA Paca-2 human pancreatic cancer cells. *Oncol Rep* **28**, 862-866, doi:10.3892/or.2012.1878 (2012).

Kandasamy, P. *et al.* Oncogenic KRAS mutations enhance amino acid uptake by colorectal cancer cells via the hippo signaling effector YAP1. *Mol Oncol*, doi:10.1002/1878-0261.12999 (2021).

Yue, M., Jiang, J., Gao, P., Liu, H. & Qing, G. Oncogenic MYC Activates a Feedforward Regulatory Loop Promoting Essential Amino Acid Metabolism and Tumorigenesis. *Cell Rep* **21**, 3819-3832, doi:10.1016/j.celrep.2017.12.002 (2017).

Coller, H. A. Cell biology. The essence of quiescence. *Science* **334**, 1074-1075, doi:10.1126/science.1216242 (2011).

Marescal, O. & Cheeseman, I. M. Cellular Mechanisms and Regulation of Quiescence. *Dev Cell* **55**, 259-271, doi:10.1016/j.devcel.2020.09.029 (2020).

Rathmell, J. C., Vander Heiden, M. G., Harris, M. H., Frauwirth, K. A. & Thompson, C. B. In the absence of extrinsic signals, nutrient utilization by lymphocytes is insufficient to maintain either cell size or viability. *Mol Cell* **6**, 683-692, doi:10.1016/s1097-2765(00)00066-6 (2000).

Muller, M. *et al.* The coordinated action of the MVB pathway and autophagy ensures cell survival during starvation. *Elife* **4**, e07736, doi:10.7554/eLife.07736 (2015).

Patwari, P. & Lee, R. T. An expanded family of arrestins regulate metabolism. *Trends Endocrinol Metab* **23**, 216-222, doi:10.1016/j.tem.2012.03.003 (2012).

Alvarez, C. E. On the origins of arrestin and rhodopsin. *BMC Evol Biol* **8**, 222, doi:10.1186/1471-2148-8-222 (2008).

Ivashov, V. *et al.* Complementary alpha-arrestin-ubiquitin ligase complexes control nutrient transporter endocytosis in response to amino acids. *Elife* **9**, doi:10.7554/eLife.58246 (2020).

Lin, C. H., MacGurn, J. A., Chu, T., Stefan, C. J. & Emr, S. D. Arrestin-related ubiquitin-ligase adaptors regulate endocytosis and protein turnover at the cell surface. *Cell* **135**, 714-725 (2008).

18   Nikko, E., Sullivan, J. A. & Pelham, H. R. Arrestin-like proteins mediate ubiquitination
and endocytosis of the yeast metal transporter Smf1. *EMBO Rep* **9**, 1216-1221,
doi:10.1038/embor.2008.199 (2008).
19   Liu, Y. *et al.* Structural basis for the regulatory role of the PPxY motifs in the
thioredoxin-interacting protein TXNIP. *Biochem J* **473**, 179-187,
doi:10.1042/BJ20150830 (2016).
20   Nabhan, J. F., Pan, H. & Lu, Q. Arrestin domain-containing protein 3 recruits the
NEDD4 E3 ligase to mediate ubiquitination of the beta2-adrenergic receptor. *EMBO*
Rep **11**, 605-611, doi:embor201080 [pii] 10.1038/embor.2010.80 (2010).
21   Rauch, S. & Martin-Serrano, J. Multiple interactions between the ESCRT machinery
and arrestin-related proteins: implications for PPXY-dependent budding. *J Virol* **85**,
3546-3556, doi:10.1128/JVI.02045-10 JVI.02045-10 [pii] (2011).
22   Zhang, P. *et al.* The ubiquitin ligase itch regulates apoptosis by targeting thioredoxin-
interacting protein for ubiquitin-dependent degradation. *J Biol Chem* **285**, 8869-8879,
doi:10.1074/jbc.M109.063321 (2010).
23   Patwari, P., Higgins, L. J., Chutkow, W. A., Yoshioka, J. & Lee, R. T. The interaction of
thioredoxin with Txnip. Evidence for formation of a mixed disulfide by disulfide
exchange. *J Biol Chem* **281**, 21884-21891, doi:10.1074/jbc.M600427200 (2006).
24   Hwang, J. *et al.* The structural basis for the negative regulation of thioredoxin by
thioredoxin-interacting protein. *Nat Commun* **5**, 2958, doi:10.1038/ncomms3958
(2014).
25   Polekhina, G. *et al.* Structure of the N-terminal domain of human thioredoxin-
interacting protein. *Acta Crystallogr D Biol Crystallogr* **69**, 333-344,
doi:10.1107/S0907444912047099 (2013).
26   Waldhart, A. N. *et al.* Phosphorylation of TXNIP by AKT Mediates Acute Influx of
Glucose in Response to Insulin. *Cell Rep* **19**, 2005-2013,
doi:10.1016/j.celrep.2017.05.041 (2017).
27   Wu, N. *et al.* AMPK-dependent degradation of TXNIP upon energy stress leads to
enhanced glucose uptake via GLUT1. *Mol Cell* **49**, 1167-1175,
doi:10.1016/j.molcel.2013.01.035 (2013).
28   Katsu-Jimenez, Y. *et al.* Absence of TXNIP in Humans Leads to Lactic Acidosis and Low
Serum Methionine Linked to Deficient Respiration on Pyruvate. *Diabetes* **68**, 709-723,
doi:10.2337/db18-0557 (2019).
29   Barr, A. R. *et al.* DNA damage during S-phase mediates the proliferation-quiescence
decision in the subsequent G1 via p21 expression. *Nat Commun* **8**, 14728,
doi:10.1038/ncomms14728 (2017).
30   Hinze, C., McGourty, K. & Boucrot, E. Measuring Endocytosis During Proliferative Cell
Quiescence. *Methods Mol Biol* **2233**, 19-42, doi:10.1007/978-1-0716-1044-2_2
(2021).
31   Kirchhausen, T., Macia, E. & Pelish, H. E. Use of dynasore, the small molecule
inhibitor of dynamin, in the regulation of endocytosis. *Methods Enzymol* **438**, 77-93,
doi:10.1016/S0076-6879(07)38006-3 (2008).
32   Macia, E. *et al.* Dynasore, a cell-permeable inhibitor of dynamin. *Dev Cell* **10**, 839-
850, doi:10.1016/j.devcel.2006.04.002 (2006).
33   Ivashov, V. *et al.* Complementary α-arrestin-ubiquitin ligase complexes control
nutrient transporter endocytosis in response to amino acids. *Elife* **9**,
doi:10.7554/eLife.58246 (2020).

34   Nishinaka, Y. *et al.* Importin alpha1 (Rch1) mediates nuclear translocation of
thioredoxin-binding protein-2/vitamin D(3)-up-regulated protein 1. *J Biol Chem* **279**,
37559-37565, doi:10.1074/jbc.M405473200 (2004).
35   Saxena, G., Chen, J. & Shalev, A. Intracellular shuttling and mitochondrial function of
thioredoxin-interacting protein. *J Biol Chem* **285**, 3997-4005,
doi:10.1074/jbc.M109.034421 (2010).
36   Qualls-Histed, S. J., Nielsen, C. P. & MacGurn, J. A. Lysosomal trafficking of the
glucose transporter GLUT1 requires sequential regulation by TXNIP and ubiquitin.
iScience **26**, 106150, doi:10.1016/j.isci.2023.106150 (2023).
37   Kim, S. *et al.* TXNIP-mediated crosstalk between oxidative stress and glucose
metabolism. *PLoS One* **19**, e0292655, doi:10.1371/journal.pone.0292655 (2024).
38   Hytti, M. *et al.* Phloretin inhibits glucose transport and reduces inflammation in
human retinal pigment epithelial cells. *Mol Cell Biochem* **478**, 215-227,
doi:10.1007/s11010-022-04504-2 (2023).
39   Qi, S., O'Hayre, M., Gutkind, J. S. & Hurley, J. H. Structural and biochemical basis for
ubiquitin ligase recruitment by arrestin-related domain-containing protein-3
(ARRDC3). *J Biol Chem* **289**, 4743-4752, doi:10.1074/jbc.M113.527473 (2014).
40   Shea, F. F., Rowell, J. L., Li, Y., Chang, T. H. & Alvarez, C. E. Mammalian alpha arrestins
link activated seven transmembrane receptors to Nedd4 family e3 ubiquitin ligases
and interact with beta arrestins. *PLoS One* **7**, e50557,
doi:10.1371/journal.pone.0050557 (2012).
41   Woelk, T. *et al.* Molecular mechanisms of coupled monoubiquitination. *Nat Cell Biol*
8, 1246-1254, doi:10.1038/ncb1484 (2006).
42   Gahlot, P. *et al.* Lysosomal damage sensing and lysophagy initiation by SPG20-ITCH.
Mol Cell **84**, 1556-1569 e1510, doi:10.1016/j.molcel.2024.02.029 (2024).
43   Patwari, P. *et al.* Thioredoxin-independent regulation of metabolism by the alpha-
arrestin proteins. *J Biol Chem* **284**, 24996-25003, doi:10.1074/jbc.M109.018093
(2009).
44   Oda, K. *et al.* L-type amino acid transporter 1 inhibitors inhibit tumor cell growth.
Cancer Sci **101**, 173-179, doi:10.1111/j.1349-7006.2009.01386.x (2010).
45   Stacpoole, P. W. Therapeutic Targeting of the Pyruvate Dehydrogenase
Complex/Pyruvate Dehydrogenase Kinase (PDC/PDK) Axis in Cancer. *J Natl Cancer*
Inst **109**, doi:10.1093/jnci/djx071 (2017).
46   Whitehouse, S., Cooper, R. H. & Randle, P. J. Mechanism of activation of pyruvate
dehydrogenase by dichloroacetate and other halogenated carboxylic acids. *Biochem J*
141, 761-774, doi:10.1042/bj1410761 (1974).
47   Knaus, L. S. *et al.* Large neutral amino acid levels tune perinatal neuronal excitability
and survival. *Cell* **186**, 1950-1967 e1925, doi:10.1016/j.cell.2023.02.037 (2023).
48   Tarlungeanu, D. C. *et al.* Impaired Amino Acid Transport at the Blood Brain Barrier Is
a Cause of Autism Spectrum Disorder. *Cell* **167**, 1481-1494 e1418,
doi:10.1016/j.cell.2016.11.013 (2016).
49   Hui, S. T. *et al.* Txnip balances metabolic and growth signaling via PTEN disulfide
reduction. *Proc Natl Acad Sci U S A* **105**, 3921-3926, doi:10.1073/pnas.0800293105
(2008).
50   Hui, T. Y. *et al.* Mice lacking thioredoxin-interacting protein provide evidence linking
cellular redox state to appropriate response to nutritional signals. *J Biol Chem* **279**,
24387-24393, doi:10.1074/jbc.M401280200 (2004).

51   Sheth, S. S. *et al.* Thioredoxin-interacting protein deficiency disrupts the fasting-
feeding metabolic transition. *J Lipid Res* **46**, 123-134, doi:10.1194/jlr.M400341-
JLR200 (2005).
52   Chutkow, W. A., Patwari, P., Yoshioka, J. & Lee, R. T. Thioredoxin-interacting protein
(Txnip) is a critical regulator of hepatic glucose production. *J Biol Chem* **283**, 2397-
2406, doi:10.1074/jbc.M708169200 (2008).
53   Coloff, J. L. *et al.* Differential Glutamate Metabolism in Proliferating and Quiescent
Mammary Epithelial Cells. *Cell Metab* **23**, 867-880, doi:10.1016/j.cmet.2016.03.016
(2016).
54   Edinger, A. L. & Thompson, C. B. Akt maintains cell size and survival by increasing
mTOR-dependent nutrient uptake. *Mol Biol Cell* **13**, 2276-2288, doi:10.1091/mbc.01-
12-0584 (2002).
55   Lee, H. J. *et al.* Proteomic and Metabolomic Characterization of a Mammalian Cellular
Transition from Quiescence to Proliferation. *Cell Rep* **20**, 721-736,
doi:10.1016/j.celrep.2017.06.074 (2017).
56   Kanai, Y. Amino acid transporter LAT1 (SLC7A5) as a molecular target for cancer
diagnosis and therapeutics. *Pharmacol Ther* **230**, 107964,
doi:10.1016/j.pharmthera.2021.107964 (2022).
57   Kaira, K. *et al.* Clinical significance of L-type amino acid transporter 1 expression as a
prognostic marker and potential of new targeting therapy in biliary tract cancer. *BMC*
Cancer **13**, 482, doi:10.1186/1471-2407-13-482 (2013).
58   Kalaany, N. How nearby nutrients shape tumor growth. *Elife* **12**,
doi:10.7554/eLife.89825 (2023).
59   Barthelemy, C. & Andre, B. Ubiquitylation and endocytosis of the human
LAT1/SLC7A5 amino acid transporter. *Sci Rep* **9**, 16760, doi:10.1038/s41598-019-
53065-w (2019).
60   Rosario, F. J., Dimasuay, K. G., Kanai, Y., Powell, T. L. & Jansson, T. Regulation of
amino acid transporter trafficking by mTORC1 in primary human trophoblast cells is
mediated by the ubiquitin ligase Nedd4-2. *Clin Sci (Lond)* **130**, 499-512,
doi:10.1042/CS20150554 (2016).
61   Teis, D. *et al.* p14-MP1-MEK1 signaling regulates endosomal traffic and cellular
proliferation during tissue homeostasis. *J Cell Biol* **175**, 861-868, doi:jcb.200607025
[pii] 10.1083/jcb.200607025 (2006).
62   Cormerais, Y. *et al.* The glutamine transporter ASCT2 (SLC1A5) promotes tumor
growth independently of the amino acid transporter LAT1 (SLC7A5). *J Biol Chem* **293**,
2877-2887, doi:10.1074/jbc.RA117.001342 (2018).
63   Schmidt, O. *et al.* Endosome and Golgi-associated degradation (EGAD) of membrane
proteins regulates sphingolipid metabolism. *EMBO J* **38**, e101433,
doi:10.15252/embj.2018101433 (2019).
64   Ran, F. A. *et al.* Genome engineering using the CRISPR-Cas9 system. *Nat Protoc* **8**,
2281-2308, doi:10.1038/nprot.2013.143 (2013).
65   Finicle, B. T. *et al.* Sphingolipids inhibit endosomal recycling of nutrient transporters
by inactivating ARF6. *J Cell Sci* **131**, doi:10.1242/jcs.213314 (2018).
66   Zou, C., Wang, Y. & Shen, Z. 2-NBDG as a fluorescent indicator for direct glucose
uptake measurement. *J Biochem Biophys Methods* **64**, 207-215,
doi:10.1016/j.jbbm.2005.08.001 (2005).

67   van Pijkeren, A. *et al.* Proteome Coverage after Simultaneous Proteo-Metabolome
1014   Liquid-Liquid Extraction. *J Proteome Res* **22**, 951-966,
1015   doi:10.1021/acs.jproteome.2c00758 (2023).
1016   68   Kipura, T. *et al.* Automated Liquid Handling Extraction and Rapid Quantification of
1017   Underivatized Amino Acids and Tryptophan Metabolites from Human Serum and
1018   Plasma Using Dual-Column U(H)PLC-MRM-MS and Its Application to Prostate Cancer
1019   Study. *Metabolites* **14**, doi:10.3390/metabo14070370 (2024).
1020

Figure 1

Figure 1. Dynamin-dependent endocytosis and lysosomal degradation of SLC7A5 and SLC1A5.

RPE1 cells were grown in DMEM supplemented with serum (0 h, + serum), cultured in serum free medium for 24 h (24 h, - serum) and refed with serum for 24 h (48 h, + serum). **(a)** Cells were harvested, permeabilized and fixed, and the DNA content was analyzed by propidium iodide (PI) staining to assess the cell cycle profile by FACS (n=6, N>3000 cells, two-way ANOVA, Tukey's multiple comparisons test). **(b)** Peak volume (in fl) was assessed by CASY Cell Counter and Analyzer (n=7, N>4000 cells, one-way ANOVA, Tukey's multiple comparisons test). **(c)** Total cells lysates were analyzed by SDS-PAGE and Western Blot (WB) with the indicated antibodies. **(d)** The cell surface was stained with anti-SLC3A2 and anti-mouse Alexa-488. FACS was used to detect SLC3A2 surface staining. A representative histogram is shown. Quantification of the fluorescence intensities of SLC3A2 was normalized to proliferating cells (+ serum) (n=5, N>4500 cells, paired t-test). **(e, g)** Indirect immunofluorescence (IF) of PFA fixed cells stained for SLC3A2 (green), LAMP1 (red) and DAPI (blue) were analyzed by confocal microscopy. The merged images show a single plane of a Z-stack. Scale bar = 10 µm. **(e)** Incubation with 12.5 µM chloroquine (CQ) in absence of serum for 14 h (- serum, + CQ). **(g)** Incubation with 20 µM dynasore (dyna) in absence of serum for 14 h (- serum, + dyna). **(f, h)** Total cells lysates of cells treated as in (e, g) were analyzed by SDS-PAGE and WB with the indicated antibodies. **(i)** Cells were incubated with [^{3}H]-leucine or [^{14}C]-isoleucine for 15 min, washed and lysed. Total cell lysates were analyzed by scintillation counting. Counts per minute (cpm) values were normalized to total protein content (cpm / µg protein, n=4, paired t-test). **(j)** Cells were incubated with [^{35}S]-methionine, [^{35}S]-cysteine for 2 minutes, before cycloheximide (CHX, 10 µg/ml) was added, cells were lysed and analyzed by SDS-PAGE and autoradiography.

Figure 2

Figure 2. The importance of TXNIP for the selective degradation of SLC7A5.

(a) TXNIP^{KO} and WT cells were grown in growth medium (+ serum) or serum starved for 24 h (- serum). Total cell lysates were analyzed by SDS-PAGE and WB with the indicated antibodies. **(b)** WB quantification of TXNIP (n=4, paired t-test), SLC7A5 (n=4, two-way ANOVA, Sidak's multiple comparisons test) and SLC7A11 protein levels (n=4, two-way ANOVA, Sidak's multiple comparisons test). Values were normalized to proliferating WT cells (+ serum). **(c)** IF of PFA fixed cells (WT, TXNIP^{KO} and TXNIP^{KO} reconstituted with TXNIP) stained for SLC3A2 (green) and DAPI (blue) were analyzed by confocal microscopy. The images show a single plane of a Z-stack. Regions of interest (ROI, white box) were magnified. Scale bar = 10 µm. **(d)** Quantification of SLC3A2 cell surface FACS of the indicated cells, normalized to proliferating WT cells (n=6, N>7000, two-way ANOVA, Sidak's multiple comparisons test). **(e)** IF of PFA fixed TXNIP^{KO} reconstituted with ALFA-TXNIP. The localization of ALFA-TXNIP (red) and SLC3A2 (green) was analyzed by confocal microscopy. The images show a single plane of a Z-stack. Regions of interest (ROI, white box) were magnified. Scale bar = 10 µm.

Figure 3

Figure 3. Loss of function mutation in TXNIP causes a rare disease.

(a) Schematic representation of WT TXNIP and mutant TXNIP harboring the patient mutation (c.642_643insT, patient^{mut}). **(b, c)** Control fibroblasts (WT#1 and WT#2) and patient-derived fibroblasts (patient) were grown in growth medium (+ serum) or serum starved for 24 h (- serum). **(b)** Total cell lysates were analyzed by SDS-PAGE and WB with the indicated antibodies. **(c)** Indirect IF of PFA fixed fibroblasts (patient, WT#1, WT#2) stained for SLC3A2 (green) and DAPI (blue) was analyzed by confocal microscopy. The images show a single plane of a Z-stack. Scale bar = 10 μm. **(d, e)** RPE1 WT, TXNIP^{KO} and TXNIP^{KO} reconstituted with ALFA-TXNIP harboring the patient mutation (c.642_643insT, patient^{mut}) were grown in growth medium (+ serum) or serum starved for 24 h (- serum). **(d)** Total cell lysates were analyzed by SDS-PAGE and WB with the indicated antibodies. **(e)** Indirect IF of PFA fixed cells (WT, TXNIP^{KO} and patient^{mut}) stained for SLC3A2 (green) and DAPI (blue) was analyzed by confocal microscopy. The images show a single plane of a Z-stack. Scale bar = 10 μm. **(f)** ALFA-TXNIP (red) and SLC3A2 (green) was analyzed by confocal microscopy. The images show a single plane of a Z-stack. Regions of interest (ROI, white box) were magnified. Scale bar = 10 μm.

Figure 4

Figure 4. PPxY motifs of TXNIP in the downregulation of SLC7A5.

(a) Schematic representation of TXNIP protein harboring PPxY motifs. **(b)** Total cell lysates from WT cells or TXNIP^{KO} reconstituted with ALFA-TXNIP^{PPxY331} mutants were analyzed by SDS-PAGE and WB with the indicated antibodies. The asterisk (*) labels a second band that was only detected using the monoclonal SLC7A5 antibody. **(c, e)** Indirect IF of PFA fixed cells (WT, TXNIP^{KO} and TXNIP^{KO} reconstituted with ALFA-TXNIP or ALFA-TXNIP^{PPXY331}) was analyzed by confocal microscopy. The images show a single plane of a Z-stack **(c)** Cells were stained for SLC3A2 (green) and DAPI (blue). Scale bar = 10 μm. **(d)** Quantification of SLC3A2 cell surface FACS of the indicated cells, normalized to proliferating WT cells (n=5, N>7000 cells, two-way ANOVA, Sidak's multiple comparisons test). **(e)** Cells were stained for ALFA-TXNIP (red) and SLC3A2 (green). Regions of interest (ROI, white box) were magnified. Scale bar = 10 μm. **(f)** SDS-PAGE and WB analysis with the indicated antibodies from non-denaturing ALFA-immunoprecipitations (IP) from HEK293T cells expressing ALFA-TXNIP or the mutated ALFA-TXNIP^{PPxY331} variant and the indicated YFP-tagged HECT-type ubiquitin ligase. **(g)** SDS-PAGE and WB analysis with the indicated antibodies from denaturing ALFA-SLC7A5 IP from SLC7A5^{KO} RPE1 reconstituted with ALFA-SLC7A5.

Figure 5

Figure 5. TXNIP and AKT signaling.

(a) Total cell lysates from WT cells before or after 24 h of serum starvation or incubation with 1 µM MK2206 (MK) or 5 µM PD0325901 (PD). Cells were analyzed by SDS-PAGE and WB with the indicated antibodies. **(b)** Indirect IF of PFA fixed cells (WT, TXNIP^{KO} and TXNIP^{KO} reconstituted with TXNIP^{S308A}, TXNIP^{S308D} or TXNIP^{C247S}) before or after serum starvation or incubation with 1 µM MK2206 (MK) stained for SLC3A2 (green) and DAPI (blue). Cells were analyzed by confocal microscopy. The images show a single plane of a Z-stack. Scale bar = 10 µm. **(c)** Total cell lysates from cells treated as in (b) were analyzed by SDS-PAGE and WB with the indicated antibodies. **(d)** Quantification of SLC3A2 cell surface FACS from the indicated cells (n=4, N>3500 cells, two-way ANOVA, Sidak's multiple comparisons test).

Figure 6

Figure 6. TXNIP mediated degradation of SLC7A5 contributed to metabolic signaling

(a) WT and TXNIP^{KO} cells were grown in growth medium (+ serum), serum starved for 24 h (- serum) or treated with 10 µM JPH203. Cells were incubated with [^{3}H]-leucine for 15 min, washed and lysed. Cell lysates were analyzed by scintillation counting. Cpm values were normalized to total protein content (n=5, two-way ANOVA, Tukey's multiple comparisons test). **(b)** WT, SLC7A5^{KO} and TXNIP^{KO} cells were grown in growth medium (+ serum) or serum starved for 24 h (- serum). Mass spectrometry analysis of free AAs, normalized to total protein content (nmol AA / µg protein, n=5). **(c)** Total cell lysates from WT and TXNIP^{KO} cells before or after 24h serum starvation, were analyzed by SDS-PAGE and WB with the indicated antibodies **(d-f)** WB quantification of pS6K^{T389}, normalized to total S6K. WB quantification of pS6$^{S240/244}$, normalized to total S6 and WB quantification of pEIF2α^{S51}, normalized to total EIF2α protein levels (n=5, two-way ANOVA). **(g)** Cells were incubated with [^{35}S]-methionine, [^{35}S]-cysteine for 2 minutes, before CHX (10µg/ml) was added, cells were lysed and analyzed by SDS-PAGE and autoradiography follow by densitometric analysis (n=3, paired t-test).

Figure 7

Figure 7. TXNIP mediated degradation of SLC7A5 contributed to the regulation of cell proliferation.

(a) WT, TXNIP^{KO} and TXNIP^{KO} cells reconstituted with TXNIP were harvested, permeabilized and fixed. The DNA content was analyzed by PI staining to assess the cell cycle profile by FACS (n=3, N>4000 cells). **(b)** Increase in confluency of the indicated cells, after refeeding serum starved cells, was measured by continuous live cell microcopy in an Incucyte incubator for 48 h (n=6). **(c)** The proliferation of the indicated cells was determined using CASY cell counting during the indicated time. The relative increase in cell number is normalized to 0 h and show as fold increase (n>3). **(d)** Model of TXNIP mediated downregulation of SLC7A5 during entry into quiescence.

Supplementary Figure 1

Supplementary Figure 1. (a) Cell viability determined with CASY counter, before and after serum starvation and after re-addition of serum (n=7, N>4000 cells). **(b)** Immortalized MEFs were grown in growth medium with serum (+ serum) or serum starved for 24 h. Total cell lysates were analyzed by SDS-PAGE and WB with the indicated antibodies. **(c, d)** Primary human fibroblasts were grown in growth medium (+ serum) or serum starved for 24 h. **(c)** Total cell lysates were analyzed by SDS-PAGE and WB with the indicated antibodies. **(d)** Cells were harvested, permeabilized and fixed. The DNA content was analyzed by PI staining and FACS (n=3, N>1500 cells, two-way ANOVA, Tukey's multiple comparisons test). **(e)** Total cell lysates of the indicated cell lines were analyzed by SDS-PAGE and WB with the indicated antibodies. **(f, g)** Indirect IF of PFA fixed cells stained for **(f, g)** SLC1A5 (green), **(f)** LAMP1 (red) and DAPI (blue) were analyzed by confocal microscopy. The merged images show a single plane of a Z-stack. **(f)** Incubation with 12,5 µM chloroquine (CQ) in absence of serum for 14 h (- serum, + CQ). Scale bar=10 µm **(g)** Incubation with 20 µM dynasore (dyna) in absence of serum for 14 h (- serum, + dyna). Scale bar=10 µm. **(h)** Cells were incubated with [^{14}C]-glutamine for 15 min, washed and lysed. Total cell lysates were analyzed by scintillation counting. Cpm values were normalized to total protein content (cpm / µg protein, n=4, paired t-test).

Supplementary Figure 2

Supplementary Figure 2. (a, g) qPCR analysis of **(a)** α-arrestins mRNA or **(g)** SLC7A5 mRNA in growing (+ serum) or serum starved cells (- serum). Expression levels were normalized to the housekeeping gene *RPLP0* (n=3, paired t-test). **(b)** Schematic presentation of the mutation introduced into TXNIP by gene editing. **(c)** Representative chromatogram of Sanger sequencing from the genome amplicon of WT TNIXP and gene edited TXNIP cells. The guide RNA and the protospacer adjacent motif (PAM) sequence are indicated. The vertical lane shows the Cas9 cleavage site causing a 13 base pair deletion in *TXNIP* (red box). Image was generated by the ICE Analysis online tool (Synthego, v2.0; Synthego, USA). **(d)** Representative histograms of SLC3A2 cell surface FACS from WT, TXNIP[KO] and TXNIP[KO] reconstituted with TXNIP cells. **(e)** WT, TXNIP[KO] and TXNIP[KO] reconstituted with TXNIP were grown in growth medium (+ serum) or serum starved for 24 h (- serum). Total cell lysates were analyzed by SDS-PAGE and WB with the indicated antibodies. **(f)** WB quantification of SLC7A5 protein levels, normalized to GAPDH (n=4, two-way ANOVA, Sidak's multiple comparisons test). **(h)** Indirect IF of PFA fixed WT and TXNIP[KO] cells stained for SLC2A1 (green) and DAPI (blue) was analyzed by confocal microscopy. The merged images show a single plane of a Z-stack. Scale bar=10 μm **(i)** Total cell lysates were analyzed by SDS-PAGE and WB with the indicated antibodies. For WB quantification, SLC2A1 protein levels were normalized to GAPDH (n=3, two-way ANOVA, Sidak's multiple comparisons test). **(j)** WT and TXNIP[KO] cells were assessed for glucose uptake by incorporation of 100 μM 2-NBDG for 45 min. Simultaneously, cells were treated with 1 mM phloretin. The fluorescence intensity was analyzed by FACS (n=3, N>5000 cells, two-way ANOVA, Sidak's multiple comparisons test).

Supplementary Figure 3

Supplementary Figure 3. (a) Total cell lysates from WT or TXNIP^{KO} cells reconstituted with TXNIP^{PPxYmut} mutants were analyzed by SDS-PAGE and WB with the indicated antibodies. The asterisk (*) labels a second band that was only detected using the monoclonal SLC7A5 antibody. **(b)** Indirect IF of PFA fixed cells (WT, TXNIP^{KO} cells reconstituted with TXNIP or TXNIP^{PPxYmut}) stained for SLC3A2 (green) and DAPI (blue) was analyzed by confocal microscopy. The images show a single plane of a Z-stack. Scale bar = 10 µm. **(c)** Representative histograms of SLC3A2 cells surface FACS of the indicated cells. **(d)** SDS-PAGE and WB analysis with the indicated antibodies from non-denaturing ALFA-immunoprecipitations (IP) from HEK293T cells expressing ALFA-TXNIP or the mutated ALFA-TXNIP^{PPxY331} variant and the indicated YFP-tagged HECT-type ubiquitin ligase. **(e)** SDS-PAGE and WB with the indicated antibodies of SLC7A5^{KO} and WT cells.

a

	+ serum	+ MK	+ PD
SLC3A2/DAPI			
SLC1A5/DAPI			

b

WT TXNIP^KO TXNIP^S308A TXNIP^S308D TXNIP^C247S

count → SLC3A2

+ serum
− serum

Supplementary Figure 4. (a) Indirect IF of PFA fixed cells before or after serum starvation or incubation with 1 µM MK2206 (MK) or 5 µM PD0325901 (PD) stained for SLC3A2 (green, upper row) or SLC1A5 (green, lower row), and DAPI (blue). Cells were analyzed by confocal microscopy. The images show a single plane of a Z-stack. Scale bar = 10 µm. **(b)** Representative histograms of SLC3A2 cells surface FACS of the indicated cells.

Supplementary Figure 5

Supplementary Figure 5. (a) WT and SLC7A5^{KO} cells were incubated with [^{3}H]-leucine for 15 min, washed and lysed. Cell lysates were analyzed by scintillation counting. Cpm values were normalized to total protein content (n=4, paired t-test). **(b, c)** WT, SLC7A5^{KO} and TXNIP^{KO} cells were grown in growth medium (+ serum) or serum starved for 24 h (- serum). Mass spectrometry analysis of free AAs, normalized to total protein content (nmol AA / µg protein, n=5).

Supplementary Table 1. AA, glucose and lactate levels in blood plasma of the TXNIP deficient patient over the indicated years.

AA [µmol/l]	06/2014	07/2014	Reference values	02/2016	03/2018	11/2021	09/2022	Reference values
leucine (L)	60.51	175.52	46 - 109	346.58	226.89	212.26	141.50	45 - 155
Isoleucine (I)	13.38	91.56	26 - 53	173.06	132.25	129.59	82.64	26 - 94
tryptophan (W)	25.21	51.44	n.a	76.35	45.51	40.21	41.48	n.a
valine (V)	99.97	148.00	57 - 262	529.46	376.67	389.20	229.86	57 - 270
phenylalanine (F)	46.45	41.50	23 - 110	84.41	54.64	53.02	56.75	23 - 70
methionine (M)	12.14	24.43	3 - 41	36.19	18.77	10.83	12.30	3 - 29
proline (P)	191.78	197.21	52 - 277	264.22	228.78	214.37	275.47	51 - 185
tyrosine (Y)	38.54	52.02	11 - 112	110.43	70.10	51.76	50.47	11 - 120
Fischer's ratio (FR)	2.05	4.46	2.10 - 4	5.38	5.90	6.98	4.23	2.10 - 4
glucose [mg/dl]	34	71	> 50	88 *	78	86	103	> 50
lactate [mg/dl]	68.30	46	5 - 22	24.33 *	50.5	26	56	5 - 22

* The glucose and lactate values were collected on a different day than the AA levels.

Supplementary Table 2. Summary of antibodies, reagents and software used in this study.

Reagents/resource	Reference or source	Identifier or catalog number	Additional information
Antibodies			
SLC1A5/ASCT2	Merck	#abn73, RRID:AB_10807715	WB (1:1000)
SLC1A5/ASCT2 (V501)	Cell Signaling	#5345, RRID:AB_10621427	WB (1:1000)
SLC1A5/ASCT2	Sigma-Aldrich	#HPA035240, RRID:AB_10604092	IF (1:50)
SLC7A5/LAT1 (polyclonal)	Cell Signaling	#5347S, RRID:AB_10695104	WB (1:2000)
SLC7A5/LAT1 (E9O4D) (monoclonal)	Cell Signaling	#32683	WB (1:2000)
SLC7A5/LAT1 (D10)	Santa Cruz	#sc-374232, RRID:AB_10988206	WB (1:100)
SLC3A2/4F2hc (D6O3P)	Cell Signaling	#13180S, RRID:AB_2687475	WB (1:1000)
SLC3A2/4F2hc	BD Biosciences	#556074, RRID:AB_396341	IF, FACS (1:100)
SLC2A1/Glut1	Merck	#07-1401, RRID:AB_1587074	WB (1:2000), IF (1:200)
SLC38A2/SNAT2	MBL International	#BMP081, RRID:AB_10597880	WB (1:100)
SLC7A11/xCT (D2M7A)	Cell Signaling	#12691, RRID:AB_2687474	WB (1:1000)
TXNIP (D5F3E)	Cell Signaling	#14715S, RRID:AB_2714178	WB (1:2000)
p-S6K^{T389} (108D2)	Cell Signaling	#9234S, RRID:AB_2269803	WB (1:1000)
S6K	Cell Signaling	#9202S, RRID:AB_331676	WB (1:1000)
p-S6$^{S240/244}$	Cell Signaling	#2215S, RRID:AB_331682	WB (1:2000)
S6 (54D2)	Cell Signaling	#2317S, RRID:AB_2238583	WB (1:1000)
p-ERK1/2$^{T202/Y204}$ (D13.14.4E)	Cell Signaling	#4370S, RRID:AB_2315112	WB (1:1000)
ERK1/2	Cell Signaling	#9102S, RRID:AB_330744	WB (1:1000)
p27^{Kip1} (D69C12)	Cell Signaling	#3686S, RRID:AB_2077850	WB (1:1000)
p-AKT^{S473} (D9E)	Cell Signaling	#4060S, RRID:AB_2315049	WB (1:1000)
p-AKT^{T308}	Cell Signaling	#9275, RRID:AB_329828	WB (1:1000)
AKT	Cell Signaling	#9272S, RRID:AB_329827	WB (1:1000)

ALFA	NanoTag Biotechnologies	#N1581, RRID:AB_3075997	WB (1:1000), IF (1:100)
Transferrin Receptor (Tfr)	DSHB	#G1/221/12, RRID:AB_2201506	WB (1:1000)
EGF Receptor (EGFR) (1005)	Santa Cruz	#sc-03-G	WB (1:500)
GAPDH (D4C6R)	Cell Signaling	#97166S, RRID:AB_2756824	WB (1:5000)
LAMP1	DSHB	#H4A3, RRID:AB_2296838	IF (1:50)
LAMP1 (D2D11)	Cell Signaling	#9091, RRID:AB_2687579	IF (1:800)
NEDD4	Cell Signaling	#5344S, RRID:AB_10560514	WB (1:1000)
WWP1	Cell Signaling	#70140S, RRID:AB_3662810	WB (1:1000)
WWP2	Cell Signaling	#41182S, RRID:AB_3662809	WB (1:1000)
ITCH	Cell Signaling	#12117S, RRID:AB_2797822	WB (1:1000)
GFP	Roche Life Science	(Roche Cat# 11814460001, RRID:AB_390913)	WB (1:1000)
Ubiquitin (P4D1)	Santa Cruz	#3936S, RRID:AB_628423	WB (1:500)
Goat anti-mouse IgG-peroxidase	Sigma-Aldrich	#A4416, RRID:AB_258167	WB (1:5000)
Goat anti-rabbit IgG-peroxidase	Sigma-Aldrich	#A0545, RRID:AB_257896	WB (1:5000)
Alexa Fluor 488 goat anti-rabbit IgG	Invitrogen	#A11008, RRID:AB_143165	IF (1:500)
Alexa Fluor 488 goat anti-mouse IgG	Invitrogen	#A11001, RRID:AB_2534069	IF (1:500)
Alexa Fluor 568 goat anti-mouse IgG	Invitrogen	#A11031, RRID:AB_144696	IF (1:500)
Alexa Fluor 568 goat anti-rabbit IgG	Invitrogen	#A11011, RRID:AB_143157	IF (1:500)
Chemicals			
Blasticidin	InvivoGen	#ant-bl-5b	
High glucose Dulbecco's modified Eagle's medium (DMEM)	Sigma-Aldrich	#D6429	
FSB	Sigma-Aldrich	#S0615	
Trypsin-EDTA solution	Sigma-Aldrich	#T4174	
MK2206	Enzo Life Sciences	#ENZ-CHM164-0005	
PD0325901	Absource	#S1036	
JPH203	Selleckchem,	#S8667	

chloroquine	Sigma-Aldrich	#C6628	
dynasore hydrate	Sigma-Aldrich	#D7693	
penicillin/streptomycin	Sigma-Aldrich	#P0781	
WesternBright Chemiluminescence substrate solution (Advansta)	Biozym	#541005X	
Lipofectamin LTX reagent	Thermo Scientific	#15338-100	
polyethylenimine (PEI)	Polyscience	#23966-100	
Polybrene	Sigma-Aldrich	#107689	
DAPI	Sigma-Aldrich	#D9542	
Mowiol	Sigma-Aldrich	#81381	
saponin	BioChemika	#84510	
RNAse A	Thermo Scientific	#EN0531	
$[^{14}C_5]$-L-glutamine	Hartmann Analytic	#MC1124	
$[4,5-^3H]$-L-Leucine	Hartmann Analytic	#MT672E	
$[^{14}C_5]$-L-Isoleucine	Hartmann Analytic	#MC174	
glutamine-free DMEM	Sigma-Aldrich	#D6546	
AA free DMEM	USBiological	#D9811-01	
2-NBDG	AAT Bioquest	#36702	
phloretin	Sigma-Aldrich	#P7912	
stable isotope-labeled canonical AA mix composition	Cambridge Isotope Laboratory	#MSK-CAA-1	
Cycloheximide	Sigma	Cat. # C7698	
anti-ALFA selector ST magnetic beads	NanoTag Biotechnologies	Cat. # N1516	
Critical commercial assays			
Micro BCA Protein Assay Kit	Thermo Scientific	#23235	
RNeasy Mini Kit	Quiagen	#74104	
LunaScript RT Super Mix Kit	NEB	#E3010	
Biozym Blue S'Green qPCR Kit	Biozym	#F-415S	
Software, algorithm			
Image J		Version 1.53t, RRID:SCR_003070	
Affinity Designer	Serfi	Version 1.10.5, RRID:SCR_016952	
GraphPad Prism 9		Version 9.4.1, RRID:SCR_002798	
Benchling		RRID:SCR_013955	
Flow Jo		Version 10.8.1, RRID:SCR_008520	
ModFit LT		RRRID:SCR_016106	

Primer3		Version 4.1.0, RRID:SCR_003139	
PrimerBank		RRID:SCR_006898	
CHOPCHOP		RRID:SCR_015723	
ZEISS ZEN Digital Imaging for Light Microscopy		Version 3.5, RRID:SCR_013672	
PrimerQuest^{TM} Tool	Integrated DNA Technologies, IDT		
ICE Analysis online tool	Synthego, USA	Synthego, v2.0	

Supplementary Table 3: List of oligonucleotides used in this study

Name	Vector backbone	Origin
TXNIP	pCCL-EIF1α-BlastiR-DEST	This study
ALFA-TXNIP	pCCL-EIF1α-BlastiR-DEST	This study
SLC7A5-ALFA	pCCL-EIF1α-BlastiR-DEST	This study
TXNIP^{S308A}/ ALFA-TXNIP^{S308A}	pCCL-EIF1α-BlastiR-DEST	This study
TXNIP^{S308D}/ALFA-TXNIP^{S308D}	pCCL-EIF1α-BlastiR-DEST	This study
TXNIP^{C247S}/ALFA-TXNIP^{C247S}	pCCL-EIF1α-BlastiR-DEST	This study
ALFA-TXNIP p.Ile215TyrfsTer59	pCCL-EIF1α-BlastiR-DEST	This study
ALFA-TXNIP^{PPxY331AAxA,PPxY375AAxA}	pCCL-EIF1α-BlastiR-DEST	This study
ALFA-TXNIP^{PPCY331AACA}	pCCL-EIF1α-BlastiR-DEST	This study
pVSV-G		Clonetech, #631530
psPAX2		Gift from Stephan Geley
pDONR-ALFA-MCS		Generated by Michael Widerin
pDONR-MCS-ALFA		Generated by Michael Widerin
pDONR-221		Invitrogen, #12536017
pCCL-EIF1α-BlastiR-DEST		Gift from Stephan Geley
pSpCas9(BB)-2A-GFP (PX458)		Addgene, #48138
YFP-NEDD4	pCR3.1	[41, 42]
YFP-ITCH	pCR3.1	[41, 42]
YFP-WWP1	pCR3.1	[41, 42]
YFP-WWP2	pCR3.1	[41, 42]
YFP-HECW1	pCR3.1	[41, 42]

| YFP-HECW2 | pCR3.1 | [41], [42] |

Supplementary Table 4: Primers for PCR-based genetic modifications and cloning

Name	Forward primer	Reverse primer
ALFA-TXNIP	tAAgcagAAttcAatggtgatgttcAAgAAgatc	TGCTTAgtcgacTCACTGCACATTGTTGTTGA
attB-TXNIP	CAAAAAAGCAGGCTCCGCCACCATGGTGATGTTCAAGAAGATCAAGTC	CAAGAAAGCTGGGTCTCACTGCACATTGTTGTTGAGG
SLC7A5-ALFA	TAAGCAGTCGACatggcgggtgcgggcccg	TGCTTAGAATTCtgtctcctgggggaccac
Gateway	GGGGACAAGTTTGTACAAAAAAGCAGGCTCC	GGGGACCACTTTGTACAAGAAAGCTGGGTC
TXNIP^{S308A}	gcagAAcatccGCGatggccag	gttcggctggccatCGCggatgttctgc
TXNIP^{S308D}	gcagAAcatccGACatggccagccg	gttcggctggccatGTCggatgttctgctg
TXNIP^{C247S}	tattatctcagggacaAGCgcatcatggcgtggc	ccacgccatgatgcGCTtgtccctgagatAAtatgattgcc
TXNIPp.Ile215TyrrfsTer59	agctgccTattgtggcccgccacacttac	gtAAgtgtggcgggccacAAtAggcagct
TXNIP^{PPxY331AAxA}	atacccagAAgctGCAGCAtgcGCAatggatgtcattcc	caggAAtgacatccatTGCgcaTGCTGCagcttctggggtatc
TXNIP^{PPxY375AAxA}	ttcAAgttcatgccaGCAGCAactGCAactgaggtggatccctgc	gcagggatccacctcagtTGCagtTGCTGCtggcatgAActtgAA
ALFA-TAG	caggggtattgacatccaccagat	atctggtggatgtcAAtacccctg

Supplementary Table 5: qPCR Primers used in this study

Gene	Forward primer	Reverse primer	Source
RPLP0	TGGTCATCCAGCAGGTGTTCGA	ACAGACACTGGCAACATTGC GG	[69]
SLC7A5	ATCGGGAAGGGTGATGTGTCCAAT	CAAAGAGGCCGCTGTATAATGCCA	[57]
ARRDC1	TAGTGGAGGAGGGTTACTTCAAC	TCTGGGATGCTGTTCAGGTTC	PrimerBank ID (65288282c1)
ARRDC2	GATTTGCGTTCTGAGGCTG	GCTTTCACCTTGTCGAATAGC	IDT
ARRDC3	TCTTGCCTCCACCTCTTTATTC	TCGGAACCCACATCAACTTG	IDT
ARRDC4	GTTCCTCTCGTCTGATTGTTCC	ATTGGCGACCATGTGTCGAAT	PrimerBank ID (190886441c 3)
TXNIP	GAGTACCTGCGCTATGAAGAC	TTTGAAGGATGTTCCCAGAGG	IDT