The condensate-forming protein eIF4ET ensures adequate levels of meiotic proteins to support oocyte storage

Priyankaa Bhatia[1], Judith Tafur[1], Ruchi Amin[2], Nicole E. Familiari[1], Kan Yaguchi[1], Vanna M. Tran[1], Alec Bond[1], Orhan Bukulmez[2], Jeffrey B. Woodruff[1,3]

1. Dept. of Cell Biology, Dept. of Biophysics, UT Southwestern Medical Center, Dallas, TX 75390, USA

2. Division of Reproductive Endocrinology and Infertility, Dept. of Obstetrics and Gynecology, UT Southwestern Medical Center Dallas, TX, USA

3. Correspondence to J. Woodruff: Jeffrey.woodruff@utsouthwestern.edu

ABSTRACT

Animals store oocytes in a dormant state for weeks to decades before ovulation. The homeostatic programs oocytes use to endure long-term storage are poorly understood. Using female nematodes as a short-lived model, we found that oocyte formation and storage required IFET-1, the conserved eIF4E-Transporter protein (eIF4ET). IFET-1 co-assembled with CAR-1 (Lsm14) to form micron-scale condensates in stored oocytes, which dissipated after oocyte activation. Depletion of IFET-1 destabilized the stored oocyte proteome, leading to lower translation, a decline in microtubule maintenance proteins, and errors in microtubule organization and meiotic spindle assembly. Deleting individual domains within IFET-1 impaired oocyte storage without affecting oocyte formation. Thus, in addition to establishing a healthy oocyte reserve in young mothers, IFET-1 ensures that correct levels of cytoskeletal proteins are maintained as oocytes age. Human eIF4ET also localized to micron-scale puncta in dormant oocytes in a reproductively healthy patient. Our results clarify the role for eIF4ET in maintaining the oocyte reserve and further support eIF4ET dysfunction as an upstream cause of embryonic aneuploidy and age-related infertility.

INTRODUCTION

The survival of many animal species depends on their ability to store high quality oocytes for prolonged periods of time. Such storage typically involves cell cycle arrest of the oocyte, typically in meiotic prophase I. In mammals, stored oocytes are small (~25-35 μm in diameter) and exist in a primordial follicle, where they are supported by squamous pre-granulosa cells (Gougeon, 1996; Li and Albertini, 2013). These oocytes can be reactivated to grow into immature GV oocytes (~100 μm in diameter), which then are either destroyed (atresia) or mature further into a metaphase-arrested form that is released for fertilization (ovulation) (Gougeon, 1996; Li and Albertini, 2013). A woman's pool of stored oocytes, called the ovarian reserve, declines immediately after birth, even before ovulation can occur. This decline continues as the woman ages, ultimately leading to a depleted ovarian reserve and menopause when she reaches 40-50 years of age (Hansen et al., 2008; Knowlton et al., 2014).   Oocyte quality also decreases during long-term storage, as defined by an increase in mitochondrial dysfunction, DNA damage, and meiotic spindle errors (Gruhn et al., 2019; Hassold et al., 1980; Titus et al., 2013). These age-related errors are correlated with an increased risk of chromosomal abnormalities in the developing fetus and miscarriage (Nybo Andersen et al., 2000). Similar phenomena have been described in a wide variety of animals (Comizzoli and Ottinger, 2021). The confluence of declining oocyte numbers and quality are thus major contributors to age-related infertility in female mammals, yet the upstream molecular causes remain poorly understood.

While much research has revealed mechanisms of oocyte production, cell cycle arrest at the GV stage, and post-GV maturation (Li and Albertini, 2013; Pei et al., 2023), comparatively little is known about long-term oocyte maintenance. In preparation for storage, oocytes enter a state of cellular hibernation in part by reducing transcription and mitochondrial production of ATP (Moore et al., 1974; Rodriguez-Nuevo et al., 2022). Understanding storage requires identifying the molecules and mechanisms that continuously provide "life support" for the oocyte during its storage. A potential inroad is through studying a disease that affects oocyte storage. Primary Ovarian Insufficiency (POI) is characterized by an accelerated decline in the ovarian reserve or general dysfunction of ovarian follicles. Patients with POI typically experience irregular menstrual

cycles, low anti-Mullerian hormone, and premature menopause (26-32 years of
age)(Nelson, 2009). As these patients start out with a large ovarian reserve and can bear
children early (prior to menopause), this disease represents a case where the body fails
to preserve oocytes over time. Mutations in in Eukaryotic Translation Initiation Factor 4E
Nuclear Import Factor 1 (*EIF4ENIF1*), which encodes a protein from the conserved
eIF4ET family, are associated with POI, suggesting a link to oocyte quality control
(Kasippillai et al., 2013; Shang et al., 2022; Zhao et al., 2019). Consistent with this theory,
EIF4ENIF1 heterozygous knockout mice display reduced follicle pools and subfertility,
recapitulating key features of POI (Ding et al., 2023). In somatic tissue culture cells,
EIF4ENIF1 binds to and mediates nuclear import of the mRNA cap-binding protein eIF4E
(Dostie et al., 2000; Ferraiuolo et al., 2005; Kamenska et al., 2016). Through use of non-
natural reporters, these studies concluded that EIF4ENIF1 associates with P-bodies to
inhibit protein translation and stabilize mRNAs. Injection of EIF4ENIF1 into frog GV
oocytes also repressed translation of a luciferase reporter (Waghray et al., 2015).
EIF4ENIF1 is found in P-body-like granules in dormant mouse oocytes (Ding et al., 2023)
and in mitochondria-associated granules (MARDO) in mammalian GV oocytes (Cheng
et al., 2022). However, it is not known if EIF4ENIF1 is required specifically during oocyte
storage and if it regulates the global pool of natural mRNAs and proteins in that context.

The mechanism of oocyte storage remains poorly understood due to the difficulty in
accessing dormant oocytes. In mammals, dormant oocytes are obtained only through
invasive surgical resection of the ovarian cortex. *In vitro* gametogenesis also cannot be
used because it does not yet recapitulate formation and long-term maintenance of the
primordial follicle (McLaughlin et al., 2018), the fundamental storage unit that comprises
the dormant oocyte surrounded by pre-granulosa cells (Li and Albertini, 2013). Thus, it
is advantageous to use a genetic model organism with a short lifespan and oocytes that
are easily visualized *in situ*.

Here, we use *C. elegans* females to study oocyte maintenance mechanisms. Most
nematodes are gonochoric and store immature oocytes arrested in meiotic prophase I.
C. elegans is an outlier in that it recently evolved hermaphrodism, yet *C. elegans* females
retain the ability to store arrested oocytes in cases where sperm are depleted naturally or
genetically (*fog-2* or *fem-1* mutations)(Schedl and Kimble, 1988; Spence et al., 1990).

Thus, oocyte storage is an ancestral feature of nematodes. As *Caenorhabditis* species

live for several weeks and store their oocytes for less time, they represent an attractive

short-lived model for oocyte storage.

RESULTS

Dysfunction of the eIF4ET homolog (IFET-1) impairs oocyte formation and storage

in female nematodes

We developed an assay to test the effects of storage time on oocyte quality. We

generated a population of genetically identical *C. elegans* females through use of a

temperature-sensitive *fem-1(hc17)* loss-of-function allele.   After synchronizing their

development, all worms reached adulthood and formed arrested oocytes at the same time

(first day of adulthood = Day 1). We then split the females into cohorts, letting them wait

another 1-5 days before introducing young males for mating. At the first day of mating

(Day 2), the female worms contained, on average, 34 +/- 7 oocytes (n= 20). Worms were

allowed to mate and lay fertilized eggs over 12-16 hr before removing all adults from the

plates. Since females can produce new oocytes after ovulating their stored oocytes, we

reported on worms that laid 34 or fewer eggs. After 2 days, we measured the number of

hatchlings divided by the number of eggs laid (oocyte viability) (Figure 1A). Under these

conditions, average oocyte viability declined with storage time, reaching 63.4% by day 6

(Figure 1B). From these data, we infer that *C. elegans* oocyte quality declines in

proportion to their storage time, like what is seen in mammals.

IFET-1 is the *C. elegans* homolog of EIF4ENIF1, but its role during oocyte storage

has not been tested. Full depletion of IFET-1 using RNAi rendered almost all stored

oocytes inviable (Figure 1B, S1A); IFET-1 levels remained low in stored oocytes, even

after worms were removed from the RNAi feeding plates (Figure S1B). IFET-1 depletion

also impaired ovulation in females and oocyte viability in hermaphrodites, which do not

arrest their oocytes, consistent with a previous study (Sengupta et al., 2013). Thus, full

knockdown of IFET-1 has pleiotropic effects making it difficult to assess its role

specifically during oocyte storage. Patients with EIF4ENIF1-linked primary ovarian

insufficiency (POI) all have heterozygous mutations, implying that failed oocyte storage

could be caused by partial reduction of EIF4ENIF1 activity. In our system, partial
knockdown of IFET-1 allowed young mothers to bear healthy progeny, although initial
viability was reduced to 31%. Oocyte viability declined faster than in controls, reaching
zero after 4 days of storage (Figure 1B). Oocytes in *ifet-1(RNAi)* females were irregular
in shape, as expected; however, all but the most proximal oocyte maintained arrest in
diakinesis, as indicated by the presence of intact germinal vesicles and condensed
chromosomes (Figure S1C). Thus, oocytes are still arrested in the absence of IFET-1,
but they are largely inviable. These results support previous findings that IFET-1 is
required for oocyte formation (Sengupta et al., 2013), and they further suggest that IFET-
1 could be required for maintenance of stored oocytes.

To definitively test the necessity of IFET-1 for the storage of healthy oocytes, we
sought a way to deplete IFET-1 only after oocyte formation is complete. Thus, we
generated a *fem-1* strain expressing a version of IFET-1 that is degraded by addition of
auxin (IFET-1::mScarlet::AID*). As outlined in Figure 1C, we placed synchronized adult
females which already have a row of arrested oocytes (Day 1) on plates with or without 4
mM auxin for 24 hr at 20°C. Then, we allowed worms to recover on normal NGM plates
for 8 hr at 20°C and allowed mating with males and egg laying for 12-16 hr at 16°C (Day
3). On average, worms laid 26 (control) vs. 13 (auxin) eggs during the laying period (n =
34 vs. 28 mothers, respectively); thus, any eggs or larvae on the plate derived from pre-
formed, healthy oocytes. Finally, we measured oocyte viability using the same method
described above.

Auxin treatment depleted IFET-1::mScarlet::AID* signal relative to controls (Figure
1D-E). In both cases, oocyte morphology appeared normal, suggesting that oocytes were
formed properly (Figure S1D). After 8 hr post-auxin removal, just prior to mating, IFET-1
signal did not recover within oocytes but did increase in the gonads (Figure S1E). Thus,
auxin-induced depletion of IFET-1::mScarlet::AID* in oocytes is robust and IFET-1 does
not re-accumulate in stored oocytes after auxin removal. Following mating, we observed
that auxin treatment significantly reduced oocyte viability (Figure 1F).

We considered the possibility that this phenotype could be due to IFET-1 being
essential during embryogenesis. However, treatment of excised embryos with 1 mM auxin
rapidly depleted IFET-1 but had no detectable effect on embryo development and viability

(Figure S1F,G). Auxin treatment did not significantly affect the fertility of control females

not expressing the degron, indicating that auxin is not generally toxic (Figure S1H). Thus,

the oocyte viability phenotype is due to lack of IFET-1 specifically during oocyte storage.

We conclude that IFET-1 is required to maintain oocyte quality during days-long storage.

IFET-1 localizes to condensates that do not contain mitochondria

To understand the function of IFET-1 in stored oocytes, we used confocal

microscopy to visualize IFET-1 tagged with mMaple (IFET-1::mMaple).   Worms

expressing IFET-1::mMaple were viable and fertile (Figure S1I). In stored oocytes, IFET-

1::Maple localized to numerous micron-scale droplets, which we term condensates. After

activation of the oocytes through fertilization, these condensates dissolved and IFET-

1::Maple signal redistributed evenly throughout the oocyte cytoplasm (Figure 1G). A

similar behavior was reported for CAR-1, which localizes to droplets in stored oocytes

(Hubstenberger et al., 2013; Noble et al., 2008) and interacts with IFET-1 in embryo

extracts (Chen et al., 2016). To co-visualize CAR-1 and IFET-1, we tagged endogenous

CAR-1 at its C-terminus with mMaple and generated antibodies against both proteins for

immunofluorescence. CAR-1::mMaple localized to condensates in stored oocytes, as

expected, and co-localized with untagged IFET-1 (Figure 1H). IFET-1::mMaple also co-

localized with untagged CAR-1. RNAi depletion of IFET-1 reduced CAR-1 localization to

condensates (Figure 1I). These results indicate that IFET-1 and CAR-1 co-assemble into

supramolecular condensates in stored oocytes.

We wondered if the IFET-1/CAR-1 condensates function as Balbiani Bodies, which

are reported to encapsulate and confine mitochondria in the in dormant oocytes of

zebrafish, frogs, and humans (Boke et al., 2016; Hertig and Adams, 1967; Marlow and

Mullins, 2008). However, mitochondria are not spatially confined in stored *C. elegans*

oocytes (Bohnert and Kenyon, 2017), and we saw no significant co-localization between

HSP60 (mitochondria marker) and IFET-1 condensates (Figure 1J). We conclude that

IFET-1 condensates are not primitive Balbiani Bodies but rather a distinct class of protein-

based supramolecular assembly that exists in stored oocytes.

Disrupting individual domains in IFET-1 perturbs oocyte storage but not formation

We next tested which domains within IFET-1 are critical for its function in oocytes.

IFET-1 contains several conserved features: an eIF4E homology domain (Eif4EHD), Cup

Homology Domain (CHD), nuclear localization signal (NLS), a region rich in glutamines

(PolyQ), and a C-terminal coiled-coil (Figure 2A). We used CRISPR to delete each of

these regions in worms expressing endogenously tagged IFET-1::mMaple (Figure 2A).

We first analyzed homozygous mutant hermaphrodite worms, which make and activate

oocytes continuously with no storage phase (McCarter et al., 1999). In this condition, all

mutant worms produced viable oocytes; ΔCHD, ΔNLS, ΔPolyQ, mutants were nearly

identical to controls, while the ΔCoiledcoil mutant showed slightly reduced viability (Figure

2B).

Mutant female worms (ΔCHD, ΔNLS, ΔPolyQ, ΔCoiledcoil) formed oocytes with

proper cuboidal morphology but could not store them effectively: after two days of oocyte

storage followed by mating, embryo viability was reduced compared to eggs laid by age-

matched hermaphrodites (Figure 2B). IFET-1::mMaple mutants localized to condensates

in stored oocytes to differing degrees (Figure 2C,D). Mutants were expressed at 55-64%

of wild-type levels (Figure 2E). Importantly, for each mutant, expression levels were

similar between female worms with stored oocytes and hermaphrodites with non-stored

oocytes (Figure S2A,B); thus, the oocyte viability phenotype is due to a distinct role for

IFET-1 in storage that is not required during active oogenesis. We speculate that these

mutations comprise IFET-1 function, especially the ΔCHD mutation, which had the most

severe oocyte viability phenotype (21.8% viability after 2 days of storage) despite being

expressed at comparable levels to other mutants. The ΔCoiledcoil mutant had the largest

effect on condensate localization, suggesting that the coiled coil domain helps IFET-1 co-

assemble with CAR-1 to form condensates (Figure 2C). These results reveal that 1) IFET-

1 functions can be separated based on domains, and 2) multiple domains are required

for full-scale IFET-1 expression and function in stored oocytes. These data further support

our conclusion that IFET-1 is required to form oocytes and maintain their quality during

the storage phase.

IFET-1 depletion causes imbalances in meiotic spindle protein levels

To determine how IFET-1 dysfunction impacts oocyte quality, we performed

proteomic and transcriptomic analyses in control vs. *ifet-1(RNAi)* female worms

containing oocytes stored for 2 days. Since the majority of IFET-1 is expressed in the

germline (oocytes + syncytial gonad; Figure 1), we report data derived from whole worms

and follow up on specific hits with fluorescent reporters that are expressed in stored

oocytes.

Depletion of IFET-1 caused disruption of the proteome. Out of 3561 proteins

detected in both control and RNAi sets, 530 proteins (14.9%) were significantly

downregulated, while 100 proteins (2.8%) were significantly upregulated in *ifet-1(RNAi)*

worms (Figure 3A; Extended Data S1). IFET-1 levels were reduced to 15% of control

levels. This result contrasts with the expectation that IFET-1 functions as a broad scale

repressor, which was based on non-native reporter assays (Sengupta et al., 2013).

Transcriptomic analyses revealed few changes in mRNA levels in control vs. *ifet-1(RNAi)*

worms. Only 6.8% of all transcripts were significantly altered (4.5% up, 2.3% down) after

depletion of IFET-1 (Figure 3B; Extended Data S2). However, there was little overlap

between significant hits in the proteomic and transcriptomic data sets, suggesting that

proteomic changes largely did not result from changes in transcript levels (Figure S2A,

B).

Using Gene Ontology (GO) enrichment analysis of downregulated proteins in *ifet-*

1(RNAi) worms, we saw that the highest-ranking category included proteins related to

embryo development and egg hatching (Figure S3C; Extended Data Set 4). Within this

category were proteins required for meiotic spindle or microtubule function, such as

Katanin (MEI-1) and oocyte-specific isoforms of tubulin (TBA-1, TBA-2). Focusing on

meiotic spindle proteins, we also saw downregulation of Kinesin-like protein 15 (KLP-15),

TPX2 (TPXL-1), Dynein Light Intermediate Chain (DLI-1), Aurora A Kinase (AIR-1) and

other tubulin isoforms (TBB-1 and TBB-2) (Figure 3C). Not all microtubule-related

proteins were downregulated; for example, Dynein Heavy Chain (DHC-1), ch-TOG (ZYG-

9), and ASPM-1 were slightly upregulated. Levels of the transcripts encoding these

proteins were all slightly downregulated in *ifet-1(RNAi)* worms (60-95% of control; Figure

3D) and showed no correlation with proteomic changes (Figure 3E). Thus, IFET-1

regulates levels of meiotic spindle and microtubule-related proteins in a post-

transcriptional manner. Proteomic analysis also revealed lower MEI-1, AIR-1, DLI-1, and
TBB-1 levels in *ifet-1(ΔCHD)* mutant females, which are defective for oocyte storage and
not active oogenesis (Figure S3D).

We then analyzed global levels of translation using a SUnSET assay. Puromycin
incorporation was reduced in worms containing arrested oocytes vs. active oocytes
(Figure S3E); thus, reduced translation is a hallmark of oocyte storage in *C. elegans*.
Depletion of IFET-1 further reduced puromycin incorporation in stored oocytes (Figure
3F). Our results suggest that lowered meiotic protein levels in *ifet-1(RNAi)* female worms
are in part due to reduced translation.

Loss of IFET-1 disrupts microtubule organization in stored and activated oocytes

Our proteomic and transcriptomic data indicated that several meiotic spindle
proteins, but not their mRNAs, were downregulated in *ifet-1* mutants. We hypothesized
that such proteostatic dysfunction should affect meiotic progression, which could explain
how IFET-1 loss impairs oocyte viability. Thus, we generated feminized worms
expressing GFP-labeled beta tubulin (*fem-1^{ts}* GFP::TBB-2), allowed them to store
oocytes, then mated them with males to induce oocyte activation and meiotic spindle
assembly (Figure 4A). In control arrested oocytes (no RNAi), microtubules appeared
throughout the cytoplasm, but they concentrated in bright networks near the oocyte
periphery, as reported before (Harris et al., 2006). After oocytes were activated by mating,
the cytoplasmic microtubules disassembled, and the meiotic spindle formed (Figure 4B).
In *ifet-1(RNAi)* oocytes, GFP::TBB-2 did not localize to bright peripheral networks and
overall expression was lower, consistent with our proteomic data (Figure 4C,D). After
mating, bipolar formation of a single meiotic spindle per oocyte failed; instead, we
observed diminutive spindle-like structures, microtubule clusters, or aggregates of tubulin
signal (Figure 4C).   Thus, IFET-1 is needed for organization of the microtubule
cytoskeleton during oocyte storage and resumption of meiosis.

Another protein affected by IFET-1 depletion was MEI-1/Katanin, a microtubule
severing enzyme important for meiotic spindle function (Yang et al., 2003). It is not known
if MEI-1 is made prior to oocyte formation, during oocyte storage, or in response to oocyte
activation. We thus used CRISPR to fuse MEI-1 with GFP (GFP::MEI-1) in feminized

worms. Confocal microscopy revealed GFP::MEI-1 signal was low in the syncytial gonad and increased 3.5-fold in stored oocytes (Figure 4E,F). GFP::MEI-1 signal was highest in the most proximal oocytes (i.e., the oldest). Thus, MEI-1 is translated in stored oocytes and accumulates over time. GFP::MEI-1 levels dropped dramatically after IFET-1 depletion (Figure 4E,F), consistent with our proteomic data. Treatment of adult worms with the proteasome inhibitor Bortezomib only weakly restored MEI-1::GFP levels in the oldest oocytes (Figure 4E,F). Overall, our results suggest that meiotic proteins are produced and maintained in stored oocytes through IFET-1-influenced translation and proteasome-mediated degradation. We conclude that IFET-1 ensures that stored oocytes produce the correct levels of MEI-1/Katanin, tubulin, and other proteins needed to organize the microtubule cytoskeleton and, after oocyte activation, to build a functional meiotic spindle.

The homolog of IFET-1 localizes to peri-nuclear puncta in dormant human oocytes

Our results suggest that IFET-1 plays an oocyte-intrinsic role in maintaining oocyte quality over time in *C. elegans*. We next examined the localization of the human homolog of IFET-1, called EIF4ENIF1. We surgically removed ovarian cortex strips from a reproductively healthy, 20-year-old patient (Figure 5A; see methods), and immediately fixed and processed the tissue for immunofluorescent staining. Stored oocytes reside in primordial follicles identified by a single layer of squamous pre-granulosa cells (Gougeon, 1996). Within these oocytes, EIF4ENIF1 localized to the cytoplasm and peri-nuclear, micron-scale puncta that morphologically resemble condensates (Figure 5B). EIF4ENIF1 puncta did not co-localize with mitochondria, similar to IFET-1 condensates in stored *C. elegans* oocytes (Figure 1). The EIF4ENIF1 puncta were specific to oocytes, as no significant signal was detected in the surrounding tissue (Figure 5C). We conclude that supramolecular assembly of eIF4ET family proteins into mitochondria-free condensates is a conserved feature of oocyte storage.

DISCUSSION

Diverse animals store and maintain immature eggs, called oocytes, for long periods of time before ovulating them. Here we used *C. elegans* females as a short-lived

model for oocyte storage and identified the eIF4ET family protein IFET-1 as essential for
oocyte formation and long-term maintenance. We report that in stored oocytes IFET-1
forms condensates and influences proteostasis, ensuring that Katanin, tubulin, and other
meiotic spindle proteins are maintained at the correct levels. This program ensures that
the oocyte microtubule cytoskeleton is supported during oocyte storage and then properly
reorganized into a functional meiotic spindle after oocyte activation.

Oocytes are thought to be largely translationally silent, yet they still produce
"housekeeping" proteins which are presumably needed for survival during long-term
storage (Richter and Lasko, 2011). Our data revealed that IFET-1 is needed for overall
translation in oocytes and, more specifically, achieving adequate levels housekeeping
proteins related to microtubule organization and completion of meiosis. IFET-1 co-
localized with CAR-1 (Lsm14B) in micron-scale condensates that formed in stored
oocytes but not in activated oocytes. Previous work suggested that these condensates
serve as super-stoichiometric repressor complexes that allow extreme concentration and
silencing of mRNAs (Cardona et al., 2023). If true, one would expect that dissolution of
the condensates would cause a massive upregulation of protein translation. IFET-1
depletion does increase expression of non-natural translational reporters (*gfp* fused to
different 3'UTRs)(Sengupta et al., 2013) and genes required for sperm production (e.g.,
fog-1, *daz-1*, *fem-3*)(Huggins et al., 2020). However, polysome profiling revealed that
translation is reduced in IFET-1-depleted hermaphrodites (Huggins et al., 2020). Our
SUNsET assay also showed reduced translation in *ifet-1(RNAi)* females containing stored
oocytes. These seemingly contradictory results are resolved by our unbiased proteomics:
530 proteins are significantly downregulated and 100 proteins are significantly
upregulated in *ifet-1(RNAi)* females. Thus, IFET-1 affects protein expression and stability
in a protein-specific manner. eIF4ET proteins are known to sequester eIF4E, thus
preventing interaction between the 40S ribosomal subunit and the 5' mRNA cap (Dostie
et al., 2000). By repressing translation of capped mRNAs, IFET-1 could, in theory, favor
translation of uncapped mRNAs needed for oocyte maintenance (Keiper, 2019). Why loss
of IFET-1 would lead to a decrease in global translation is still unclear. We also cannot
rule out the possibility that loss of select proteins is due to increased proteasome-
mediated degradation, which we showed is active for maintaining MEI-1 levels in stored

oocytes. The primary role of IFET-1 may also change in a cell-type-specific manner. Our
data, which used proteomics and live-cell imaging of GFP-tagged endogenous MEI-1,
clearly indicated that IFET-1 depletion reduces MEI-1 levels in stored oocytes. However,
a study using immunofluorescence reported higher levels of MEI-1 in the embryos of *ifet-*
1 mutant hermaphrodites (Li et al., 2009). MEI-1 is normally degraded during the oocyte-
embryo transition (Kurz et al., 2002). It is possible that IFET-1 is required for both
translation of MEI-1 during oocyte storage and the factors that degrade MEI-1 during the
oocyte-embryo transition. Understanding the mechanistic link between IFET-1 and MEI-
1 translation will require future investigation. Our data revealed that many other classes
of proteins are affected by *ifet-1(RNAi)*; it is thus possible that IFET-1's control of MEI-1
and other microtubule-related protein levels is indirect.

IFET-1 shares homology to human EIF4ENIF1, a protein whose function during
mammalian oocyte storage remains poorly defined. Women with heterozygous mutations
in *EIF4ENIF1* suffer a form of early-onset infertility called Primary Ovarian Insufficiency
(POI)(Kasippillai et al., 2013; Shang et al., 2022; Zhao et al., 2019). Symptoms typically
arise in the patient's late 20's and early 30's and include amenorrhea, a rapid depletion
of the ovarian reserve of quiescent oocytes, and an elevated risk of miscarriage (Dean et
al., 2018). Prior to diagnosis, patients with POI can bear healthy children, indicating that
POI does not block or perturb oocyte development but rather the long-term maintenance
of oocytes in the ovary (Nelson, 2009). We showed that mutating conserved regions in
IFET-1 specifically impairs oocyte storage but not formation, thus mimicking a key feature
of POI. Furthermore, we demonstrated that EIF4ENIF1 localizes to micron-scale
condensates in dormant primary oocytes in a patient, similar to our observations of IFET-
1 in *C. elegans*. By analogy, it is possible that human oocytes assemble EIF4ENIF1
condensates or smaller cytoplasmic complexes as a quality control mechanism to
regulate protein levels during long-term oocyte storage. We hypothesize that EIF4ENIF1
dysregulation could be an upstream cause of meiotic division errors associated with age-
related infertility in women.
367
368   **ACKNOWLEDGEMENTS**

We thank the UTSW proteomics core facility for performing mass spectrometry, the UTSW Histology core for sectioning human tissue, and Abby Dernberg (UC Berkeley) for the mMaple plasmid.

FUNDING

This project was funded by a Pew Biomedical Scholars award, a Welch Catalyst award (V-I-0004-20230731), and the Endowed Scholars program at UT Southwestern (J.B.W). R.A and O.B are supported by an Obstetrics and Gynecology departmental fund.

AUTHOR CONTRIBUTIONS

P. Bhatia performed experiments in worms and analyzed data. J. Tafur performed the auxin-induced degradation experiments and analyzed data. N. Familiari maintained worms and performed crosses. V. Tran and A. Bond performed initial viability assays. K. Yaguchi made the GFP::MEI-1 strain. R. Amin performed the oophorectomy to retrieve human ovarian tissue and immunofluorescence. O. Bukulmez and J. Woodruff supervised the project. J. Woodruff wrote the paper.

SUPPLEMENTAL DATA

Figure S1. In vivo control experiments for oocyte proteins.

Figure S2. Analysis of *ifet-1* mutant worms.

Figure S2. Analysis of proteomic and transcriptomic data for female worms with stored oocytes.

Table S1. Table of *C. elegans* strains.

Extended Data Set 1. Protein abundances in control vs. *ifet-1(RNAi)* female worms with stored oocytes.

Extended Data Set 2. mRNA abundances in control vs. *ifet-1(RNAi)* female worms with stored oocytes.

Extended Data Set 3. Raw data for oocyte viability assays.

Extended Data Set 4. GO terms and protein classifications.

METHODS

Worm Strains and Maintenance

All worms were maintained at 16°C on Nematode Growth Media (NGM) with OP50 until reaching the adult gravid stage and passaged regularly to maintain a steady population of worms. GFP and mMaple knock-ins were generated using CRISPR (Paix et al., 2015). Worm strains are listed in Table S1.

Worm synchronization and feminization

All worms with a fem-1 temperature sensitive genotype had to be maintained at 16ºC to maintain hermaphrodites. When arrested oocytes were needed for an experiment, worms were grown at 16ºC until the plate was full of mostly adults and eggs. Plates were washed with 3.5ml of dH$_2$O and transferred to a falcon tube. To this, 1ml of 5% bleach and 0.5ml of 5M NaOH was added and the falcon tubes spun for 1-2 minutes or until all adult worms were dissolved. This embryo enriched solution was spun on a centrifuge at 1300 RCF for 1 min, then washed in DI water twice at same spin settings. The pellet of embryos was resuspended in 1 ml of M9 media and left to rotate at room temperature overnight. The next day, the tubes were collected and contained synchronized L1 hatchlings. These were transferred to NGM plates with OP50 and grown at 25º for 72 hours to fully feminize and mature to adults with oocytes in Day 1 arrest. It is at this stage that worms were collected for all subsequent experiments.

RNAi Knockdowns

RNAi was done via feeding. The *ifet-1* clone was obtained from the [insert library here] library and sequenced. The target sequence for ifet-1 is AACGTGGCTAATCCAATGCTTGCAAGACTTTTCGGTCATAGTGGTGGTGATAACAA CGCTTCATCTTCTGGAGCTGGTGATATAAAAGGAGGAATGCGGTTGGAAGATTTGG AGAAAGGCATGGAGTCAAAGGAACCTTCAAAAGTCAGTCCATTGCAAGATCCATCA CAACAAGCACAACTTCTGCAACATCTGCAAAAATTCGCCAAACAGCAAGCAGAGTC TGGTCAGCACATACATCATCATAGACAACCAACACCACCAAATGGAGGTCCACAGC ATCAACAACATTTGCATCACCCTATGGTTCATCCTGGAATGCAAATTATCGCAGATC CAAGCCTTCTTGCAAGCTTCGCACAGAATCCAGTGATTTTGAATGCCTATGTCGAG AATCAACTTCAAGAAGCAGTCAATGCTGCAATTCGTGCAAACAATGGGCAGCAGCT

TCCACCACAGCTTCGTATGTTAAGCATTTTTGAATACATTTCATGATTGAATTATTTT

TCAGATGAGCAACTTCGTATGGCATCCATGCGTAACAAGGCTTTCCTACAGTCTCA

GACTCTCACTTTCGTTAGTCTTCAACAACAGCATCAGAACATCCAGCAGCATCAGC

AACAACAACAGCATCATCAACACAAGGTCGCACTCCTGCTATGATACCTGCGTCT

GTGCAGCGGCAATTGCAGAAGAGCTCCTCAAATGCTGATCAGAAGAAAGAAAGA

CGAGTCAGAGCCCGCCAGAGAGCAACCAAGAGACTTCCGACGCTCATAATCAATC

TGATGCCATGAACCAACTGAAGAAATTGCACATGCAGCAGAACTATGCCAATATGG

TGCAGGCAATGAATTCTGGAGTCGGATGGGCACGCGGAAACGGCGTTGTTAACGG

ACAACAGCAACATCCACAGCAGCAACTTCCACCAAATGTTCAAATGCTGATGGCTC

AACACCAGCAAGCTCAAATGCAGCACCTCAAG.

Bacteria containing the target sequence plasmids were seeded onto NGM plates

containing 1mM IPTG and 100 ug/ml ampicillin. Worms were bleached and synchronized

as described above. After rotating overnight in M9, the L1 larvae were transferred onto

prepared RNAi plates and grown at 25ºC for 46-72 hours (depending on the experiment).

Worms were then transferred off the RNAi plate once they reached Day 1 of adulthood

and allowed to grow on OP50 at 20ºC.

Auxin treatment

Worm synchronization and feminization was conducted as described above. When

worms reached Day 1 of adulthood they were transferred to 20°C. Half the adult worms

were transferred to NGM plates with OP50 containing 4mM auxin for 24 hours, half were

maintained on a normal OP50 plate. After auxin treatment, the worms were transferred

back to a regular NGM plate with OP50 for 8 hours. Both control and auxin-treated worms

were then singled out to mate with 3 males for ~12-16 hours at 16°C. The males used

were also synchronized using the above method and grown to adulthood. At the end of

the mating period, adult worms were removed from the mating plates and the total amount

of embryos on each plate were counted. 2 days later the number of hatchlings on each

plate was counted and % embryonic viability was calculated. Data included was only

from worms that laid 34 or less embryos.

For IFET-1 depletion in embryos, embryos were extracted from IFET-1::mScarlet::AID*

hermaphrodite worms using two dissection needles in a solution of M9 containing 1 mM

auxin. These embryos were then imaged overnight on a Nikon Widefield microscope to

visualize hatching and IFET-1::mScarlet signal.

Bortezomib Feeding

Agar plates were prepared by diluting Bortezomib (Millipore Sigma, #5.04314) to 20 µM

in OP-50 and seeding plates as above. Worms were then fed on these plates for 16-24

hours to test recovery after RNAi feeding.

Viability Assay

Worms were bleached and synchronized as described above. L1 larvae were grown at

25ºC for 72 hours to feminize. After 72 hours, all adult worms on day 1 of oocyte arrest

were transferred to 20ºC to reduce heat stress and experiments were conducted on this

population. 12-15 day 2 adults were picked and transferred to individual mating plates.

f*og-2* worms were also synchronized and grown at the same rates and temperatures as

the feminized line. Three males per adult female were transferred to each mating plate.

Worms were allowed to mate and lay eggs for 10-16 hours at 16ºC. All adults were then

removed, and laid eggs were counted and recorded. 48 hours later, all hatchling larvae

were counted. Oocyte viability is calculated as number of hatchlings divided by the

number of eggs laid per mother. The average viability per strain is reported for worms

that laid ≤34 eggs to maintain a brood size that matches average oocyte storage numbers.

Western Blot

Worms were grown on agar plates until day 2 of adulthood. They were washed off the

plate with M9 and allowed to settle at the bottom of the tube. Worms were transferred

onto a fresh agar plate with no bacteria. 40 worms of each condition were picked and

transferred to 25 ul DI water. 25 µl SDS was added to the samples, and all were boiled at

95º for 5-10 minutes (or until worms dissolve). Samples were run on SDS-PAGE gel for

30 minutes. Proteins were transferred onto a nitrocellulose membrane using Trans Blot

Turbo Transfer System (Bio-Rad). Nitrocellulose was blocked in 3% blocking buffer for
an hour, then primary antibody (1:1000) was added for overnight incubation at 4ºC.
Membranes were washed the next morning in TBS-T three times, then incubated with
secondary antibody (1:10,000) for an hour. Membranes were washed again with TBS-T
three times and chemiluminescence was expressed using SuperSignal West Femto
Maximum Sensitivity Substrate (Thermo) Membranes were imaged on ChemiDoc Touch
Imaging System (Bio-Rad).

Quantification of western blots was calculated using FIJI. The mean gray value of any
protein bands of interest were measured and the background mean value of the blot was
subtracted. The mean gray value of the control was also measured in the same way
(whole lane was measured if it was a Ponceau S stain) and the values of the protein of
interest were normalized to the amounts of control or total protein stained.

Collection, preparation, and staining of human ovarian tissue
Tissue was collected as described (Amin et al., 2023). Briefly, a 20-year-old female
patient with no known pathology or history of infertility elected for total hysterectomy and
bilateral salpingo-oophorectomy for purposes of transgender care. Informed consent was
received prior to the procedure. Use of human ovarian tissue was approved by the
Institute Review Board of the University of Texas Southwestern Medical Center
(IRB#0801-404/012012-185). Ovarian tissue was placed in 4% formaldehyde directly
from the operating room and incubated for 24 hours for fixation. The tissue was embedded
in paraffin and cut into 5 mm sections using Leica Rotary Microtome. For
immunohistochemistry, the mounted paraffin sections were dewaxed and rehydrated in
xylene and serial dilutions of ethanol, respectively. Following a wash step with phosphate-
buffered solution (PBS), an antigen retrieval step was performed using a basic retrieval
reagent (R&D, CTS013, USA) at 90-95°C for 5 minutes. The slides were cooled
completely in PBS prior to processing.

For immunohistochemistry, the tissue was inserted into a solution containing 10% goat
serum, 3% bovine serum albumin, and 0.03% Triton X-100 (blocking solution) for one
hour. The tissue was then incubated overnight with primary antibody at 4°C. The following

primary antibodies were used: directly-labeled mouse monoclonal anti-HSP60 (Proteintech, 1:500 dilution; 66041-1-Ig, USA) and rabbit polyclonal anti-EIF4ENIF1 (Abcam, 1:500 dilution; ab277638, USA). After primary antibody incubation, a series of washes were performed with PBS. The tissue was then incubated with a goat anti-rabbit secondary antibody (Invitrogen, 1:1000 dilution, A-11008) for 1 hour prior to staining for DNA with Vectashield (DAPI) (Vectorlabs, H-1000-10, USA). Negative controls were included by incubating with no primary antibody. The directly labeled anti-HSP60 was created using an Alexa Fluor 647 Lightning-Link kit (Abcam, USA) following their instructions.

Confocal microscopy

Worms were imaged using a Nikon Eclipse Ti2-E microscope with a Yokogawa confocal scanner unit (CSU-W1), piezo Z stage, and an iXon Ultra 888 EMCCD camera (Andor), controlled by Nikon Elements software. Images were collected using a 60X water (NA 1.2) or 100X silicon (NA 1.35) immersion objective, 0.3 µm Z-stacks, 2x2 binning, 15% laser power, and 100 ms exposures. Human tissue and worm oocytes were also imaged using a Nikon AX-R confocal microscope with a 40X silicone (NA 1.25) objective.

Proteomics

Worms were feminized and synchronized as described above and plated on three separate plates at 25ºC. When the worms reached Day 2 of adulthood, they were washed in M9 media for 15 min to wash off bacteria. For each experimental group, three groups of 30 worms each were collected and transferred to 20 µl of dH_2O in three separate Eppendorf tubes. 20 µl SDS was added to the tubes and samples were boiled for 5-10 minutes (or until all worms were dissolved into SDS solution). These samples were run on an SDS-PAGE gel for 3 min to allow all the sample to enter the gel, and then the gel was stained with Coomassie dye. The protein band was cut out of the gel.

Gel samples were digested overnight with trypsin (Pierce) following reduction and alkylation with DTT and iodoacetamide (Sigma–Aldrich). Following solid-phase extraction cleanup with an Oasis HLB µelution plate (Waters), the resulting peptides were

reconstituted in 2% (v/v) acetonitrile (ACN) and 0.1% trifluoroacetic acid in water. 1 ug

of each sample was injected onto an Orbitrap Fusion Lumos mass spectrometer coupled

to an Ultimate 3000 RSLC-Nano liquid chromatography systems (Thermo). Samples were

injected onto a 75 µm i.d., 75-cm long EasySpray column (Thermo),and eluted with a

gradient from 0-28% buffer B over 90 min. Buffer A contained 2% (v/v) ACN and 0.1%

formic acid in water, and buffer B contained 80% (v/v) ACN, 10% (v/v) trifluoroethanol,

and 0.1% formic acid in water. The mass spectrometer operated in positive ion mode with

a source voltage of 2.5 kV and an ion transfer tube temperature of 300 °C. MS scans

were acquired at 120,000 resolution in the Orbitrap and up to 10 MS/MS spectra were

obtained in the Orbitrap for each full spectrum acquired using higher-energy collisional

dissociation (HCD) for ions with charges 2-7. Dynamic exclusion was set for 25 s after an

ion was selected for fragmentation.

Raw MS data files were analyzed using Proteome Discoverer v3.0 (Thermo), with peptide

identification performed using Sequest HT searching against the C. elegans reviewed

protein database from UniProt along with the sequence of fluorescent mMaple protein.

Fragment and precursor tolerances of 10 ppm and 0.6 Da were specified, and three

missed cleavages were allowed. Carbamidomethylation of Cys was set as a fixed

modification and oxidation of Met was set as a variable modification. The false-discovery

rate (FDR) cutoff was 1% for all peptides.

Data was analyzed in XMLS format. All protein abundances were normalized by

averaging the sum of total proteins in each sample sent and getting a ratio to the average

of the control samples sent. The ratios derived were used to adjust the protein abundance

values up or down accordingly. From these normalized values, the fold change and P-

value (2-tailed, unequal variance) were calculated. Anything with a p-value below 0.05

was deemed significant. Data was plotted using GraphPad Prism. Identified proteins are

listed in Extended Data Set 1. GO Term analysis was conducted using PANEGA

(https://www.flyrnai.org/tools/pangea/web/home/6239).

Transcriptomics

Worms were synchronized, feminized and grown on two 100 mm plates. When worms
reached day 2 of adulthood, they were washed off the plates with M9 and rotated for 20
min to remove all the bacteria. Worms were rinsed twice with M9. Total RNA was
extracted from worms with the Arcturus PicoPure RNA Isolation Kit (Applied
Biosystems™). Total RNA was shipped to Novogene (Sacramento, CA, USA). RNA
degradation and contamination was monitored on 1% agarose gels. RNA purity was
checked using the NanoPhotometer® spectrophotometer (IMPLEN, CA, USA). RNA
integrity and quantitation were assessed using the RNA Nano 6000 Assay Kit of the
Bioanalyzer 2100 system (Agilent Technologies, CA, USA).

A total amount of 1 µg RNA per sample was used as input material for the RNA sample
preparations. Sequencing libraries were generated using NEBNext® UltraTM RNA
Library Prep Kit for Illumina® (NEB, USA) following manufacturer's recommendations and
index codes were added to attribute sequences to each sample. Briefly, mRNA was
purified from total RNA using poly-T oligo-attached magnetic beads. Fragmentation was
carried out using divalent cations under elevated temperature in NEBNext First Strand
Synthesis Reaction Buffer (5X). First strand cDNA was synthesized using random
hexamer primer and M-MuLV Reverse Transcriptase (RNase H-). Second strand cDNA
synthesis was subsequently performed using DNA Polymerase I and RNase H.
Remaining overhangs were converted into blunt ends via exonuclease/polymerase
activities. After adenylation of 3' ends of DNAfragments, NEBNext Adaptor with hairpin
loop structure were ligated to prepare for hybridization. In order to select cDNA fragments
of preferentially 150~200 bp in length, the library fragments were purified with AMPure
XP system (Beckman Coulter, Beverly, USA). Then 3 µl USER Enzyme (NEB, USA) was
used with size-selected, adaptor-ligated cDNA at 37 °C for 15 min followed by 5 min at
95 °C before PCR. Then PCR was performed with PhusionHigh-Fidelity DNA polymerase,
Universal PCR primers and Index (X) Primer. PCR products were purified (AMPure XP
system) and library quality was assessed on the Agilent Bioanalyzer 2100 system.

The clustering of the index-coded samples was performed on a cBot Cluster Generation
System using PE Cluster Kit cBot-HS (Illumina) according to the manufacturer's

instructions. After cluster generation, the library preparations were sequenced on an
Illumina platform and paired-end reads were generated. Raw data (raw reads) of FASTQ
format were firstly processed through fastp. In this step, clean data (clean reads) were
obtained by removing reads containing adapter and poly-N sequences and reads with
low quality from raw data. At the same time, Q20, Q30 and GC content of the clean data
were calculated. All the downstream analyses were based on the clean data with high
quality.

Reference genome and gene model annotation files were downloaded from genome
website browser (NCBI/UCSC/Ensembl) directly. Paired-end clean reads were aligned to
the reference genome using the Spliced Transcripts Alignment to a Reference (STAR)
software, which is based on a previously undescribed RNA-seq alignment algorithm that
uses sequential maximum mappable seed search in uncompressed suffix arrays followed
by seed clustering and stitching procedure. STAR exhibits better alignment precision and
sensitivity than other RNA-seq aligners for both experimental and simulated data.
Differential expression analysis between two conditions/groups (three biological
replicates per condition) was performed using DESeq2 R package. DESeq2 provides
statistical routines for determining differential expression in digital gene expression data
using a model based on the negative binomial distribution. The resulting P values were
adjusted using the Benjamini and Hochberg's approach for controlling the False
Discovery Rate (FDR). Genes with an adjusted P value < 0.05 found by DESeq2 were
assigned as differentially expressed.

Prior to differential gene expression analysis, for each sequenced library, the read counts
were adjusted by Trimmed Mean of M-values (TMM) through one scaling normalized
factor. Differential expression analysis of two conditions was performed using the edgeR
R package. The P values were adjusted using the Benjamini and Hochberg methods.
Corrected p value of 0.005 and |log2(Fold Change)| of 1 were set as the threshold for
significantly differential expression. Identified transcripts are listed in Extended Data Set
2.

Puromycin Assay

Worms were synchronized and grown at 25º on 100mm NGM plates till they reached Day 2 of adulthood. They were washed off the plates with M9 buffer and rinsed in an Eppendorf tube with S-Basal Medium (100mM NaCl, 5mM K_2HPO_4, 50mM KH_2PO_4, 5µg/ml cholesterol, 10mM potassium citrate, 3mM $MgSO_4$, 1% trace metals solution) twice. 250µl of S-Basal Medium was added to the live worm pellet along with 50 µl of 10mg/ml puromycin, 200ul of concentrated OP50 and topped off to 1 ml with S-Basal Medium. The tubes were shaken at 200rpm for 4 hours at room temperature. After 4 hours, tubes were collected, and the worms were washed with S-Basal three times. Worms were resuspended in 300ul lysis buffer with Protease Inhibitors (150mM NaCl, 50mM Hepes, 1mM EDTA, 1% Triton-X, 0.5% SDS) and flash frozen in liquid nitrogen. The tubes were placed on ice and sonicated 5-8 times at 20-30% amplitude for 2 seconds at a time (until most carcasses were dissolved). They were then spun at 13,000 RPM for 30 mins and 300 µl of the supernatant was collected. 20% v/v of TCA was added to solution and protein was allowed to precipitate for 30 min on ice. Tubes were spun at 14,000 RPM for 5 min and the supernatant was pipetted off. Pellet was washed twice in 90% acetone and then resuspended and boiled in 40 µl 2X SDS loading buffer. Samples were run on an SDS-PAGE gel, transferred to nitrocellulose and first stained with Ponceau-S stain and imaged. Membranes were then stained with anti-puromycin antibody (Millipore Sigma, MABE343**)** and imaged using a ChemiDoc imager (Bio-Rad).

Immunostaining of *C. elegans*

Tube fixation and collagenase protocol was adapted from protocols in www.wormbook.org. Briefly, worms were fixed in 4% formaldehyde in PBS in an eppendorf tube. Tubes were flash frozen in liquid nitrogen and subsequently thawed 2-3 times. They were shaken at 37ºC for 3 hours. Fixative was then removed and washed off with PBST. Worms were shaken at 100rpm in B-mercaptoethanol solution (5%BME, 1%Triton X-100, 120 mM Tris) for 2 hours at 37º. Tubes were spun at 1000 RCF and washed with PBST up to 10 times if needed, or until fixative is no longer detectable. Samples were transferred to a fresh tube after half the washes. 200 µl of Collagenase Solution (100 mM Tris, 1mM $CaCl_2$, 0.1%Triton X-100, pH 7.4, 2000 units Collagenase)

was added and worms were shaken at 37ºC at 250 rpm for 38-48 hours (depending on

the stock of collagenase). Incubation was stopped when a majority of worms were broken

in half. Collagenase was washed off and worms were blocked in 1 ml blocking buffer

(PBS, 0.5%Triton X-100, 1mM EDTA, 5%BSA, 0.05% Sodium Azide) at room

temperature for 1 hour. Blocking buffer was removed and worms were incubated with

primary antibody in blocking buffer (1:1000) overnight at 4ºC. Worms were washed three

times for 15-minute washes with antibody buffer (blocking buffer but with only 0.1% BSA)

and then incubated with secondary antibody in blocking buffer (1:10,000) for 3 hours at

room temperature. Three more 15-minute washes were done and worms transferred onto

a slide with an agar pad or wire-spacing. Mounting medium was added and slides were

imaged on spinning disc confocal. Rabbit polyclonal anti-IFET-1 and anti-CAR-1

antibodies were made by GenScript.

Statistical analyses

Statistical analyses were performed using GraphPad Prism. The indicated tests, the n

values, and significant values are displayed in the figure legends or on the figures

themselves. Significance was reported as: p value < 0.05 = *, p value < 0.005 = **, p

value < 0.0005 = ***, p value of < 0.00005 = ****.

TABLE S1. *C. elegans* strains.

Strain name	genotype	Creation method	Origin
BA17	*fem-1 (hc17) IV*	mutagenesis	CGC
CB4108	*fog-2(q71) V*	mutagenesis	CGC
JWW112	*car-1(utsw6[car-1::mMaple]) I*	CRISPR	This study
JWW166	*ifet-1(utsw15[ifet-1::mMaple]) III*	CRISPR	This study
JWW173	*utsw15[ifet-1::mMaple]III; fem-1 (hc17) IV*	cross	This study

JWW140	car-1(utsw6[car-1::mMaple]) I; fem-1(hc17) IV	cross	This study
JWW236	mei-1(utsw17[gfp::mei-1]) I; fem-1 (hc17) IV	CRISPR	This study
JWW163	[pie-1p::GFP::H2B + unc-119(+)] III;fem-1 (hc17) IV	cross	This study
PHX6862	ifet-1(syb6862[ifet-1(Δ196-217)::mMaple]) III	CRISPR	SunyBiotech
JWW228 (ΔCHD)	ifet-1(syb6862[ifet-1(Δ196-217)::mMaple]) III; fem-1(hc17) IV	cross	This study
PHX6949	ifet-1(syb6949[ifet-1(Δ220-235)::mMaple]) III	CRISPR	SunyBiotech
JWW215 (ΔNLS)	ifet-1(syb6949[ifet-1(Δ220-235)::mMaple]) III; fem-1(hc17) IV	cross	This study
PHX6886	ifet-1(syb6862[ifet-1(Δ527-644)::mMaple]) III	CRISPR	SunyBiotech
JWW216 (ΔpolyQ)	ifet-1(syb6862[ifet-1(Δ527-644)::mMaple]) III; fem-1(hc17) IV	cross	This study
PHX6954	ifet-1(syb6862[ifet-1(Δ664-691)::mMaple]) III	CRISPR	SunyBiotech
JWW230 (ΔCC)	ifet-1(syb6862[ifet-1(Δ664-691)::mMaple]) III; fem-1(hc17) IV	cross	This study
JWW252	fem-1 (hc17) IV; ruis57[pie-1p::GFP::tubulin + unc-119(+)]	cross	This study
JWW253	ifet-1(utsw16[ifet-1::mScarlet_I::AID*::3xFlag]) III; wrdSi3 [sun-1p::TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2 3'UTR] (II:0.77).	CRISPR	This study
JWW255	fem-1 (hc17) IV; ifet-1(utsw16[ifet-1::mScarlet_I::AID*::3xFlag]) III; wrdSi3 [sun-1p::TIR1::F2A::mTagBFP2::AID*::NLS::tbb-2 3'UTR] (II:0.77).	cross	This study

REFERENCES

Amin, R., O. Bukulmez, and J.B. Woodruff. 2023. Visualization of Balbiani Body disassembly during human primordial follicle activation. *MicroPubl Biol*. 2023.

Bohnert, K.A., and C. Kenyon. 2017. A lysosomal switch triggers proteostasis renewal in the
immortal C. elegans germ lineage. *Nature*. 551:629-633.
Boke, E., M. Ruer, M. Wuhr, M. Coughlin, R. Lemaitre, S.P. Gygi, S. Alberti, D. Drechsel, A.A.
Hyman, and T.J. Mitchison. 2016. Amyloid-like Self-Assembly of a Cellular Compartment.
Cell. 166:637-650.
Cardona, A.H., S. Ecsedi, M. Khier, Z. Yi, A. Bahri, A. Ouertani, F. Valero, M. Labrosse, S. Rouquet,
S. Robert, A. Loubat, D. Adekunle, and A. Hubstenberger. 2023. Self-demixing of mRNA
copies buffers mRNA:mRNA and mRNA:regulator stoichiometries. *Cell*. 186:4310-4324
e4323.
Chen, J.X., P.G. Cipriani, D. Mecenas, J. Polanowska, F. Piano, K.C. Gunsalus, and M. Selbach.
2016. In Vivo Interaction Proteomics in Caenorhabditis elegans Embryos Provides New
Insights into P Granule Dynamics. *Mol Cell Proteomics*. 15:1642-1657.
Cheng, S., G. Altmeppen, C. So, L.M. Welp, S. Penir, T. Ruhwedel, K. Menelaou, K. Harasimov, A.
Stutzer, M. Blayney, K. Elder, W. Mobius, H. Urlaub, and M. Schuh. 2022. Mammalian
oocytes store mRNAs in a mitochondria-associated membraneless compartment.
Science. 378:eabq4835.
Comizzoli, P., and M.A. Ottinger. 2021. Understanding Reproductive Aging in Wildlife to Improve
Animal Conservation and Human Reproductive Health. *Front Cell Dev Biol*. 9:680471.
Dean, D.D., S. Agarwal, and P. Tripathi. 2018. Connecting links between genetic factors defining
ovarian reserve and recurrent miscarriages. *J Assist Reprod Genet*. 35:2121-2128.
Ding, Y., Z. He, Y. Sha, K. Kee, and L. Li. 2023. Eif4enif1 haploinsufficiency disrupts oocyte
mitochondrial dynamics and leads to subfertility. *Development*. 150.
Dostie, J., M. Ferraiuolo, A. Pause, S.A. Adam, and N. Sonenberg. 2000. A novel shuttling
protein, 4E-T, mediates the nuclear import of the mRNA 5' cap-binding protein, eIF4E.
EMBO J. 19:3142-3156.
Ferraiuolo, M.A., S. Basak, J. Dostie, E.L. Murray, D.R. Schoenberg, and N. Sonenberg. 2005. A
role for the eIF4E-binding protein 4E-T in P-body formation and mRNA decay. *J Cell Biol*.
170:913-924.
Gougeon, A. 1996. Regulation of ovarian follicular development in primates: facts and
hypotheses. *Endocr Rev*. 17:121-155.
Gruhn, J.R., A.P. Zielinska, V. Shukla, R. Blanshard, A. Capalbo, D. Cimadomo, D. Nikiforov, A.C.
Chan, L.J. Newnham, I. Vogel, C. Scarica, M. Krapchev, D. Taylor, S.G. Kristensen, J. Cheng,
E. Ernst, A.B. Bjorn, L.B. Colmorn, M. Blayney, K. Elder, J. Liss, G. Hartshorne, M.L.
Grondahl, L. Rienzi, F. Ubaldi, R. McCoy, K. Lukaszuk, C.Y. Andersen, M. Schuh, and E.R.
Hoffmann. 2019. Chromosome errors in human eggs shape natural fertility over
reproductive life span. *Science*. 365:1466-1469.
Hansen, K.R., N.S. Knowlton, A.C. Thyer, J.S. Charleston, M.R. Soules, and N.A. Klein. 2008. A
new model of reproductive aging: the decline in ovarian non-growing follicle number
from birth to menopause. *Hum Reprod*. 23:699-708.
Harris, J.E., J.A. Govindan, I. Yamamoto, J. Schwartz, I. Kaverina, and D. Greenstein. 2006. Major
sperm protein signaling promotes oocyte microtubule reorganization prior to
fertilization in Caenorhabditis elegans. *Dev Biol*. 299:105-121.
Hassold, T., P. Jacobs, J. Kline, Z. Stein, and D. Warburton. 1980. Effect of maternal age on
autosomal trisomies. *Ann Hum Genet*. 44:29-36.

Hertig, A.T., and E.C. Adams. 1967. Studies on the human oocyte and its follicle. I. Ultrastructural
and histochemical observations on the primordial follicle stage. *J Cell Biol*. 34:647-675.
Hubstenberger, A., S.L. Noble, C. Cameron, and T.C. Evans. 2013. Translation repressors, an RNA
helicase, and developmental cues control RNP phase transitions during early
development. *Dev Cell*. 27:161-173.
Huggins, H.P., J.S. Subash, H. Stoffel, M.A. Henderson, J.L. Hoffman, D.S. Buckner, M.S. Sengupta,
P.R. Boag, M.H. Lee, and B.D. Keiper. 2020. Distinct roles of two eIF4E isoforms in the
germline of Caenorhabditis elegans. *J Cell Sci*. 133.
Kamenska, A., C. Simpson, C. Vindry, H. Broomhead, M. Benard, M. Ernoult-Lange, B.P. Lee, L.W.
Harries, D. Weil, and N. Standart. 2016. The DDX6-4E-T interaction mediates
translational repression and P-body assembly. *Nucleic Acids Res*. 44:6318-6334.
Kasippillai, T., D.G. MacArthur, A. Kirby, B. Thomas, C.B. Lambalk, M.J. Daly, and C.K. Welt. 2013.
Mutations in eIF4ENIF1 are associated with primary ovarian insufficiency. *J Clin
Endocrinol Metab*. 98:E1534-1539.
Keiper, B.D. 2019. Cap-Independent mRNA Translation in Germ Cells. *Int J Mol Sci*. 20.
Knowlton, N.S., L.B. Craig, M.T. Zavy, and K.R. Hansen. 2014. Validation of the power model of
ovarian nongrowing follicle depletion associated with aging in women. *Fertil Steril*.
101:851-856.
Kurz, T., L. Pintard, J.H. Willis, D.R. Hamill, P. Gonczy, M. Peter, and B. Bowerman. 2002.
Cytoskeletal regulation by the Nedd8 ubiquitin-like protein modification pathway.
Science. 295:1294-1298.
Li, R., and D.F. Albertini. 2013. The road to maturation: somatic cell interaction and self-
organization of the mammalian oocyte. *Nat Rev Mol Cell Biol*. 14:141-152.
Li, W., L.R. DeBella, T. Guven-Ozkan, R. Lin, and L.S. Rose. 2009. An eIF4E-binding protein
regulates katanin protein levels in C. elegans embryos. *J Cell Biol*. 187:33-42.
Marlow, F.L., and M.C. Mullins. 2008. Bucky ball functions in Balbiani body assembly and animal-
vegetal polarity in the oocyte and follicle cell layer in zebrafish. *Dev Biol*. 321:40-50.
McCarter, J., B. Bartlett, T. Dang, and T. Schedl. 1999. On the control of oocyte meiotic
maturation and ovulation in Caenorhabditis elegans. *Dev Biol*. 205:111-128.
McLaughlin, M., D.F. Albertini, W.H.B. Wallace, R.A. Anderson, and E.E. Telfer. 2018. Metaphase
II oocytes from human unilaminar follicles grown in a multi-step culture system. *Mol
Hum Reprod*. 24:135-142.
Moore, G.P., S. Lintern-Moore, H. Peters, and M. Faber. 1974. RNA synthesis in the mouse
oocyte. *J Cell Biol*. 60:416-422.
Nelson, L.M. 2009. Clinical practice. Primary ovarian insufficiency. *N Engl J Med*. 360:606-614.
Noble, S.L., B.L. Allen, L.K. Goh, K. Nordick, and T.C. Evans. 2008. Maternal mRNAs are regulated
by diverse P body-related mRNP granules during early Caenorhabditis elegans
development. *J Cell Biol*. 182:559-572.
Nybo Andersen, A.M., J. Wohlfahrt, P. Christens, J. Olsen, and M. Melbye. 2000. Maternal age
and fetal loss: population based register linkage study. *BMJ*. 320:1708-1712.
Paix, A., A. Folkmann, D. Rasoloson, and G. Seydoux. 2015. High Efficiency, Homology-Directed
Genome Editing in Caenorhabditis elegans Using CRISPR-Cas9 Ribonucleoprotein
Complexes. *Genetics*. 201:47-54.

Pei, Z., K. Deng, C. Xu, and S. Zhang. 2023. The molecular regulatory mechanisms of meiotic
arrest and resumption in Oocyte development and maturation. *Reprod Biol Endocrinol*.
21:90.
Richter, J.D., and P. Lasko. 2011. Translational control in oocyte development. *Cold Spring Harb*
Perspect Biol. 3:a002758.
Rodriguez-Nuevo, A., A. Torres-Sanchez, J.M. Duran, C. De Guirior, M.A. Martinez-Zamora, and E.
Boke. 2022. Oocytes maintain ROS-free mitochondrial metabolism by suppressing
complex I. *Nature*. 607:756-761.
Schedl, T., and J. Kimble. 1988. fog-2, a germ-line-specific sex determination gene required for
hermaphrodite spermatogenesis in Caenorhabditis elegans. *Genetics*. 119:43-61.
Sengupta, M.S., W.Y. Low, J.R. Patterson, H.M. Kim, A. Traven, T.H. Beilharz, M.P. Colaiacovo, J.A.
Schisa, and P.R. Boag. 2013. ifet-1 is a broad-scale translational repressor required for
normal P granule formation in C. elegans. *J Cell Sci*. 126:850-859.
Shang, L., S. Ren, X. Yang, F. Zhang, L. Jin, X. Zhang, and Y. Wu. 2022. EIF4ENIF1 variants in two
patients with non-syndromic premature ovarian insufficiency. *Eur J Med Genet*.
65:104597.
Spence, A.M., A. Coulson, and J. Hodgkin. 1990. The product of fem-1, a nematode sex-
determining gene, contains a motif found in cell cycle control proteins and receptors for
cell-cell interactions. *Cell*. 60:981-990.
Titus, S., F. Li, R. Stobezki, K. Akula, E. Unsal, K. Jeong, M. Dickler, M. Robson, F. Moy, S.
Goswami, and K. Oktay. 2013. Impairment of BRCA1-related DNA double-strand break
repair leads to ovarian aging in mice and humans. *Sci Transl Med*. 5:172ra121.
Waghray, S., C. Williams, J.J. Coon, and M. Wickens. 2015. Xenopus CAF1 requires NOT1-
mediated interaction with 4E-T to repress translation in vivo. *RNA*. 21:1335-1345.
Yang, H.Y., K. McNally, and F.J. McNally. 2003. MEI-1/katanin is required for translocation of the
meiosis I spindle to the oocyte cortex in C elegans. *Dev Biol*. 260:245-259.
Zhao, M., F. Feng, C. Chu, W. Yue, and L. Li. 2019. A novel EIF4ENIF1 mutation associated with a
diminished ovarian reserve and premature ovarian insufficiency identified by whole-
exome sequencing. *J Ovarian Res*. 12:119.
820
821

822 **FIGURE LEGENDS**

823 **Figure 1: IFET-1 is required for long-term storage of quiescent oocytes in *C.**
824 ***elegans* females.**

A. Strategy to study oocyte storage and viability in *C. elegans* females. *fem-1(hc17)* worms were synchronized at L1, grown to adulthood at 25°C to induce feminization and oocyte storage, then separated into five different cohorts. Female cohorts were mated with young males on different days, then offspring were analyzed.

B. Percentage of oocytes that hatched, as described in A) (mean +/- 95% C.I. n=12-30 worms, brood sizes ≤34 eggs laid). Partial knockdown of *ifet-1* was achieved by feeding on RNAi plates for 48 hr; full knockdown was achieved by feeding for 72 hr. See Extended Data Set 3 for all viability data.

C. Strategy to deplete IFET-1 in stored oocytes after their formation. Synchronized *fem-1(hc17) ifet-1::mScarlet::AID* TIR-1::T2A::BFP* adult female worms were placed on plates with 0 mM or 4 mM auxin for 24 hr. Females were removed from auxin for 8 hr prior to mating with males.

D. Left, diagram of a *C. elegans* female. Right, confocal image of Day 3 adult females expressing IFET-1::mScarlet::AID* grown on plates containing 0 mM (control) or 4 mM auxin. Scale Bar, 10µm

E. Integrated fluorescence intensity of IFET-1::mScarlet::AID* signal in stored oocytes from panel D) (mean +/- 95% C.I.; n = 15 worms).

F. Viability of stored oocytes after mating in IFET-1::mScarlet worms with or without auxin treatment (mean +/- 95% C.I.; n = 34 (control) vs. 28 (auxin) worms, brood sizes =26 (control) vs 13 (auxin) eggs per mother).

G. Live-cell confocal images of IFET-1::mMaple in stored (female, not mated) and activated (hermaphrodite) oocytes. Scale bars, 10 µm.

H. Immunofluorescence images of stored oocytes. Scale bars, 10 µm.

I. Left, images of stored oocytes expressing CAR-1::mMaple. Right, quantification of total mass of CAR-1 condensates per worm (mean +/- 95% C.I. n = 850-900 condensates from 5-8 worms).

J. Immunofluorescent staining of mitochondria (HSP60) in stored oocytes
expressing IFET-1::mMaple. Scale bars, 10 µm.

Figure 2. Specific domains in IFET-1 contribute to oocyte storage.

A. Design of mutant constructs created with CRISPR modification of *ifet-1::mMaple*
worms. Shown is the protein sequence of IFET-1 (CHD, Cup Homology Domain;
NLS, nuclear localization signal; CC, coiled-coil domain; PolyQ, region rich in
glutamines).

B. Quantification of oocyte viability from 2-day-old adult hermaphrodites (Non-stored
oocytes) and females (Stored oocytes) after mating (mean +/- 95% C.I. n =
embryos from 15-22 worms, brood size ≤34 eggs laid for female worms). See
Extended Data Set 3 for raw data.

C. Confocal images of stored oocytes expressing the mutant variants of IFET-
1::mMaple shown in A.

D. Quantification of IFET-1::mMaple assembly into stored condensates (mean +/-
95% C.I. n = 500-600 condensates from 5-8 worms).

E. Quantification of mean IFET-1::mMaple fluorescence signal in oocytes (mean +/-
95% C.I. from 10-15 worms).

Figure 3. Proteomic and transcriptomic profiling of oocyte-storing worms lacking
IFET-1.

A. Left, volcano plot of protein abundance in *ifet-1(RNAi)* females vs. control
females. Right, pie chart indicating protein changes that surpassed significance
thresholds (p value < 0.05, fold change > 2; n = 3 replicates). See Extended Data
Set 1 for a list of all proteins.

B. Left, volcano plot of mRNA abundance in *ifet-1(RNAi)* females vs. control
females. Right, pie chart indicating protein changes that surpassed significance
thresholds (p value < 0.05, fold change > 2; n = 3 replicates). See Extended Data
Set 2 for a list of all transcripts.

C. Analysis of selected proteins related to meiotic spindle formation. Shown are
average protein abundances in *ifet-1(RNAi)* worms relative to controls.

D. Analysis of selected transcripts related to meiotic spindle formation. Shown are
average mRNA abundances in *ifet-1(RNAi)* worms relative to controls.
E. Comparison of changes in protein and mRNA abundance for genes/proteins
listed in C) and D). There is no correlation between the data sets (R^2= 0.007; n =
21).
F. SuNSET assay testing translation levels by measuring puromycin incorporation
into proteins. Lane 1, control female worms with stored oocytes. Lane 2, *ifet-*
1(RNAi) female worms with stored oocytes. Left, blot probing for puromycin.
Middle, ponceau stain showing total protein levels per sample. Right,
quantification of puromycin signal relative to ponceau signal and normalized to
signal of control worms. (n=4 blots).
894

Figure 4. IFET-1 depletion affects organization of the microtubule cytoskeleton
and MEI-1/Katanin levels in stored oocytes.
A. Workflow of experiments in female worms with stored oocytes. Mating induces
oocyte activation and meiotic spindle assembly.
B. Live-cell confocal images of arrested, activated, and Meiosis I oocytes in control
female worms expressing GFP::TBB-2 (beta tubulin). Yellow arrowhead, an
assembling meiotic spindle.
C. Same as in B), except with *ifet-1(RNAi)* female worms. Yellow arrowhead,
microtubule asters; Red arrowheads, tubulin aggregates.
D. Quantification of GFP::TBB-2 levels in arrested oocytes (mean +/- 95% C.I. n =
4-6 worms;).
E. Quantification of GFP::MEI-1 (Katanin) levels in arrested oocytes relative to
gonads (mean +/- 95% C.I. n = 6-8 worms;).
F. Images of GFP::MEI-1 expression in arrested oocytes. Top: Arrested oocytes,
Middle: arrested oocytes with *ifet-1(RNAi),* Bottom: arrested oocytes fed on *ifet-*
1(RNAi) and then on Bortezomib plates to block the proteosome. Scale bars= 10
μm.
912

Figure 5. Localization of the IFET-1 homolog EIF4ENIF1 in human stored oocytes.

A. H&E staining of an ovarian cortical slice resected from a 20-year-old patient.

Primordial follicles containing stored oocytes are labeled.

B. Immunofluorescent staining of mitochondria (HSP60), DNA, and EIF4ENIF1 in a

primordial follicle.

C. A larger view of the ovary cortex reveals that EIF4ENIF1 only localized to

oocytes within the primordial follicle and not surrounding cells.

SUPPLEMENTAL FIGURE LEGENDS

Figure S1. In vivo control experiments for oocyte proteins.

A. Western blot against IFET-1; anti-alpha tubulin blotting was used as a loading

control. Worm lysates were separated first by SDS-PAGE.

B. Confocal images of stored oocytes expressing IFET-1::mMaple with and without

ifet-1(RNAi) treatment across day 2 and day 5 of adulthood. Scale bars, 10 μm.

C. Confocal imaging of GFP-labeled histone H2B in stored oocytes (outlined in

dashed white lines). Scale bar, 10 μm.

D. DIC images of stored oocytes from the experiment in Figure 1 D-F. Scale bar, 10

μm.

E. Images of IFET-1::mScarlet::AID* signal in various conditions, related to Figure

1C-D.

F. Embryos were dissected in M9 + 1mM auxin and imaged with DIC (top) and

fluorescence (bottom) over time. Scale bar, 10 μm.

G. Viability of embryos after auxin treatment in panel F.

H. Viability of oocytes in *fem-1* worms not expressing a degron-tagged version of

IFET-1 (mean +/- 95% C.I.; n = 27 mothers, <34 laid eggs each). See Extended

Data Set 3 for raw data.

I. Viability of embryos laid by hermaphrodite versions of CRISPR-tagged strains

(mean +/- 95% C.I.; n = 12 mothers, >45 embryos each).

Figure S2. Analysis of IFET-1 mutant worms.

A. Western blot against IFET-1 in indicated strains. Hermaphrodite and female

worms indicated with symbols. Ponceau stain was used as a loading control.

B. Quantification of A) indicating grey values of IFET-1 in the western blot,

normalized by level of ponceau staining.

Figure S3. Analysis of proteomic and transcriptomic data for female worms with

stored oocytes.

A. Overlap between proteomic and transcriptomic data sets from Figure 3.

B. A list of the 15 gene products where mRNA and protein are significantly down-

regulated after RNAi against *ifet-1*.

C. GO-Term analysis of proteomic data in Figure 3.

D. Analysis of selected proteins related to meiotic spindle formation. Shown are

average protein abundances in *ifet-1(ΔCHD)* worms relative to *ifet-1(WT)*

controls. Bars indicate means (N= 2 data sets with 3 replicates each). Purple,

striped, and grey, indicate that the differences were significant in 2, 1, or 0 data

sets, respectively.

E. SUnSET assay comparing puromycin incorporation between worms with

activated oocytes and worms with arrested oocytes. Quantification of three blots

with arrested signal normalized to activated oocyte signal. (Significance tested

with one-sample t-test).

Figure 1

Figure 2

A IFET-1::mMaple truncations

WT

eIF4E Homology | PolyQ | | CC
1 CHD NLS 761

ΔCHD (Δ196-217)

ΔNLS (Δ220-235)

ΔPolyQ (Δ527-644)

ΔCoiledCoil (Δ664-691)

C Stored oocytes (Day 2)

WT   ΔCHD   ΔNLS

IFET-1::mMaple   ΔPolyQ   ΔCoiledCoil
Low   High

B
Hermaphrodite (No oocyte storage)

% Oocyte viability

WT ΔCHD ΔNLS ΔPolyQ ΔCC

Female (Oocytes stored for 2 days)

% Oocyte viability
****

WT ΔCHD ΔNLS ΔPolyQ ΔCC

D Integrated density of condensates in oocytes

Conensate mass (A.U.)

WT ΔCHD ΔNLS ΔPolyQ ΔCC

E Mean intenstiy of mMaple signal in oocytes

Fluorescence (A.U.)
****
ns

WT ΔCHD ΔNLS ΔPolyQ ΔCC

Figure 3

Figure 4

A GFP::Tubulin females with stored oocytes

Arrested Oocytes

ifet-1(RNAi) → Arrested Oocytes with *ifet-1(RNAi)*

4-6 hrs → Activated oocytes (mated)

ifet-1(RNAi) 4-6 hrs → Activated oocytes (mated) with *ifet-1(RNAi)*

B Arrested Oocytes — GFP::Tubulin
20 µm Oocytes Spermatheca
Activated Oocytes Oocyte in Meiosis I
10 µm 10 µm

C Arrested Oocytes with *ifet-1(RNAi)* — GFP::Tubulin
20 µm Oocytes Spermatheca
Activated Oocytes Oocyte in Meiosis I
10 µm 10 µm

D GFP::tubulin signal in oocytes (A.U.) **** WT *ifet-1(RNAi)*

E GFP::MEI-1 signal in oocyte/gonad **** **** WT *ifet-1(RNAi)* +bortezomib

F Arrested Oocytes — GFP::MEI-1
Low — High
WT
ifet-1(RNAi)
+ bortezomib
ifet-1(RNAi)
Oldest Oocytes

Figure 5

A   Human ovarian cortex

Primordial follicles

50 μm

B   EIF4ENIF1   DNA   HSP60   Zoom

20 μm

5 μm

C   EIF4ENIF1   DNA

Primordial follicle

50 μm

Figure S1

Figure S2

Figure S3

A

Comparison of proteomics and transcriptomics

Significantly Upregulated

338 Proteins | 0 | 745 Genes

Significantly Downregulated

784 Proteins | 15 | 361 Genes

C

Top 10 GO-Terms: Downregulated Proteins

- Embryo development ending in birth or egg hatching
- Phosphorylation
- Proteolysis
- Multicellular organism development
- Translation
- Nematode larval development
- Protein transport
- Metabolic process
- IRE1-mediated unfolded protein response

X-axis: Number of proteins from proteomics profile (0, 5, 10, 15, 20, 25)

Top 10 GO-Terms: Upregulated Proteins

- Transmembrane transport
- Ion transport
- Chloride transmembrane transport
- Protein transport
- Chloride transport
- Cell division
- Protein ubiquitination
- Cell cycle
- Nematode larval development
- Determination of adult lifespan

X-axis: Number of proteins from proteomics profile (0, 5, 10, 15)

B

Proteins and genes that were down-regulated significantly in ifet-1(RNAi) worms

puf-5----------------pumillo containing family
ifet-1----------------eIF4E Transporter
clec-87-------------C-type Lectin
wdfy-2--------------WD40 and FYVE domain
cec-7----------------chromodomain protein
acs-2----------------fatty Acid CoA Synthetase family
pygl-1--------------PYGL glycogen phosphoylase
tbb-1---------------- tubulin beta
ddo-2-------------- D-aspartate oxidase
lys-7-------------- Lysozyme
ugt-26--------------UDP-Glucuronosyl Transferase
cyp-34a9----------Cytochrome P450 family
C01G10.7
CELE_F32D8.12
Y52B11A.8

D

Meiotic Proteins from ΔCHD Vs. Full-length IFET-1

Protein of Interest (y-axis): DHC-1, PLK-2, TBA-2, TBB-2, DLI-1, TBA-1, TBB-1, AIR-1, MEI-1, IFET-1

X-axis: Percentage of control (0, 50, 100)

E

| Puromycin | Ponceau |
| 1 2 | kDa L 1 2 |

Puromycin Incorporation

Ratio of Puromycin/Ponceau Stain Normalized to N2 (0.0, 0.5, 1.0)

1 2

**

1. Worms with activated oocytes
2. Worms with arrested oocytes