Distinct roles of brain network flexibility in motor learning across age

Kazumasa Uehara[1, 2, 3], Makoto Hagihara[4, 3], and Keiichi Kitajo[3, 4]

1) Neural Information Dynamics Laboratory, Department of Computer Science and Engineering, Toyohashi University of Technology, Toyohashi, Japan
2) Human-Centered AgriTech Co-Creation Center (HAC[3]), Toyohashi University of Technology, Toyohashi, Japan
3) Division of Neural Dynamics, National Institute for Physiological Sciences, Okazaki, Japan
4) Department of Physiological Sciences, School of Life Science, The Graduate Institute for Advanced Studies, SOKENDAI, Okazaki, Japan

Keywords: Motor learning, Brain network flexibility, Electroencephalogram, Aging
Running Title: Brain network flexibility and motor learning in aging

Correspondence to: Kazumasa Uehara, Ph.D., Associate Professor
Neural Information Dynamics Laboratory,
Department of Computer Science and Engineering,
and Human-Centered AgriTech Co-Creation Center (HAC[3])
Toyohashi University of Technology, Toyohashi
1-1 Hibarigaoka, Tempaku-cho, Toyohashi, Aichi, 441-8580, JAPAN
Tel/Fax: +81-532-44-1215
E-mail: uehara.kazumasa.so@tut.jp

Keiichi Kitajo, Ph.D, Professor
Division of Neural Dynamics,
National Institute for Physiological Sciences
38 Nishigonaka, Myodaiji, Okazaki, Aichi, 444-8585, JAPAN
Tel: +81-564-55-7751, Fax: +81-564-55-7754
E-mail: kkitajo@nips.ac.jp

Abstract

Motor learning is a lifelong process, from infancy through old age. Acquiring new motor actions through repetitive practice requires adjusting motor output in response to sensory input and integrating them to facilitate learning. For this to occur, the central nervous system must flexibly predict and adapt to the dynamic interplay between sensory inputs and motor outputs. Although overall brain function changes with age, it remains unclear how the flexible brain network evolves and influences motor learning ability with aging. To address these open questions, we designed a visuomotor learning task involving both younger and older adults and quantitatively assessed brain network flexibility, leveraging multichannel electroencephalography (EEG) in humans. Here, we found age-group differences in motor learning properties, brain network flexibility, and their neural relationships. In younger adults, learning aftereffects were associated with brain network flexibility during the preparatory period of the task. However, this association was not observed in older adults. Together, our findings suggest that brain network flexibility during the preparatory period, rather than motor execution, was critical for acquiring and maintaining new motor actions in younger adults, but that this coupling was attenuated with aging.

Introduction

More recently, many people live well beyond the age of 60 and maintain good health. In our current reality, various types of motor and cognitive learning occur throughout a lifetime, spanning from early infancy to old age. Notably, learning new motor skills, such as playing musical instruments, remains feasible even in older adulthood. The definition of skill emphasizes reducing variability in action outcomes, and this variability progressively decreases throughout the learning process. However, the relationship between motor learning and movement variability remains a subject of debate at both the behavioral and neural levels. At the behavioral level, movement variability (i.e., exploratory behavior) is thought to play a pivotal role in facilitating motor learning (Ranganathan and Newell, 2010; Uehara et al., 2019; Ranganathan et al., 2020), as effectively promoting motor exploration leads to better outcomes (Wu et al., 2014). Nevertheless, conclusive evidence regarding the neurological rationale behind remains elusive. Our scalp electroencephalogram (EEG) study demonstrated that overall brain network flexibility reflects key neural processes involved in skill formation and predicts musical performance skill in proficient brass players. Notably, brain network flexibility primarily contributed to feedback control, rather than feedforward control, during skilled musical performance (Uehara et al., 2023). Together, this evidence suggests that brain network flexibility may underlie the sensorimotor integration and the control of the consolidated motor function. In this context, brain network flexibility, reflecting moment-to-moment reconfiguration of large-scale brain networks, emerges as a candidate neural mechanism supporting learning acquisition. This study utilized the brain network flexibility, which is defined as a quantitative index of node flexibility, capturing the frequency with which nodes change their community assignments over a given period. Here, nodes represent brain regions, and a module represents a group of brain regions that are more strongly interconnected with each other. While this mathematical framework initially seemed disconnected from biological mechanisms, growing empirical evidence suggests that it reliably

accounts for fundamental properties of brain function (Braun et al., 2015; Betzel et al., 2017;

Reddy et al., 2018; Monteiro et al., 2019; Uehara et al., 2023). Notably, N-methyl-D-aspartate

receptors, which play a main role in neural inhibition, have the potential to modulate the

degree of brain network flexibility (Braun et al., 2016). Similarly, dopamine D1- and D2-

receptor function also influences state transitions in brain networks (Braun et al., 2021). This

implies that synaptic transmitter activity significantly impacts brain network flexibility. Because

these neuromodulators and synaptic processes are themselves altered across the lifespan,

brain network flexibility would also be expected to change with aging. With advancing age,

elderly people undergo widespread changes in neuromuscular function even during healthy

aging, which can disrupt the interplay between sensory input, motor output, cognition, and

their neural integrations (Goble et al., 2009; Coxon et al., 2014; Serbruyns et al., 2015; Frolov

et al., 2020; Zapparoli et al., 2022; Goelman et al., 2023). These age-related changes in

sensorimotor function result in a gradual decline in sensorimotor function and learning ability

(Vercillo et al., 2017; Hehl et al., 2020; Pauelsen et al., 2020). This decline is associated with

reductions in white and gray matter volumes (Courchesne et al., 2000; Good et al., 2001;

Jernigan et al., 2001) and alterations in white matter microstructural organization (Cox et al.,

2016; Beck et al., 2021), changes in brain activity, and network dynamics (Garrett et al., 2013;

Monteiro et al., 2019; Frolov et al., 2020; Goelman et al., 2023), and declines in dopaminergic

and gamma-aminobutyric acid neurotransmission (Kaasinen and Rinne, 2002; Guitart-Masip

et al., 2016; Hermans et al., 2018) even in healthy aging. These neural factors may

individually or collectively disrupt the coordinated operation of brain networks across multiple

levels, ultimately leading to behavioral changes (Marstaller et al., 2015; Shafiei et al., 2019;

Yoshimura et al., 2020). Additionally, a human functional magnetic resonance imaging (fMRI)

study has already demonstrated that the strength of brain network flexibility was reduced in

older adults, and the flexibility of the brain area in the prefrontal cortex was associated with

lower error during bimanual motor coordination (Monteiro et al., 2019). These findings allow

us to raise an open question whether age-group changes in brain network flexibility could

hinder motor learning in the elderly. The majority of existing studies have focused on age-

group–specific alterations in brain activity or connectivity and their relationship to behavioral

performance. However, age-group differences in brain network flexibility and motor learning

ability remain underexplored, and the neural mechanisms by which brain network flexibility

supports adaptation through motor learning are still unclear.

Here, we aimed to investigate whether age-group changes in brain network flexibility

affect motor learning ability in both young adults (YA) and older adults (OA). We utilized

multichannel scalp EEG recordings while participants from both YA and OA groups engaged in

learning an error-based visuomotor adaptation task. In the present study, we hypothesized

that entire brain network flexibility would increase with aging due to excessive neural noise

(Garrett et al., 2010) and compensatory reorganization (Cabeza, 2002), and that this

increased flexibility becomes functionally ineffective in supporting motor learning in older

adults. Additionally, based on previous works (Deco and Kringelbach, 2016; Sase and Kitajo,

2021), we posit that brain network flexibility reflects not just localized cortical activity, such as

specific motor-related regions, but the broader, system-level organization of functional brain

networks. Likewise, motor learning engages a distributed network of brain regions rather than

being confined to a single cortical area. Specifically, the frontal area is implicated in planning

and attentional processes, the cerebellum and basal ganglia are involved in prediction and

motor adjustment, and the sensory cortices play a critical role in integrating sensory feedback

(Cavanagh et al., 2010; Krakauer and Mazzoni, 2011; van de Vijver et al., 2011; Roth and

Ding, 2024; Fukuda and Uehara, 2025). Focusing on whole-brain dynamics in terms of

network flexibility is therefore essential for elucidating neural mechanisms of motor learning

ability in the context of aging. Moreover, previous EEG studies suggest that the suppression of

trial-by-trial neural variability, as indexed by event-related potentials (ERPs) in the present

study, is closely associated with enhanced perceptual abilities (Arazi et al., 2017; Daniel and

Dinstein, 2021). Given this, the reduction in brain network flexibility during the motor learning

process, relative to the early phase of learning, may help stabilize brain activity. We therefore

hypothesized that these distinctive changes are modulated by aging and, in turn, impact motor

learning abilities.

Materials and Methods

In this study, we selected an error-based visuomotor adaptation task to assess individual

motor learning abilities. To estimate the number of participants required to reliably assess

behavioral motor learning effects, we conducted an a priori power analysis using the G*Power

program (Erdfelder et al., 2009) with the effect size set at 0.49 based on our prior data

collection from seven healthy right-handed young individuals (26.4 ± 5.5 yrs, range: 22 -35

yrs, two females). Using the same experimental protocol described later (see *Experimental*

task and protocols), the effect size was determined assuming changes in motor performance

(i.e., the degree of the deviation of cursor position) between the early, middle, and late phases

of motor learning. The statistical significance level was set at $p < 0.05$ (two-tailed) for a one-

way repeated measures analysis of variance (ANOVA) with a within-subject factor (motor

learning phase). Despite the sample size calculation being conducted only for a younger

group, this a priori power analysis estimation showed that at least 25 participants were

required for one group, and effect sizes were reported for each statistic to confirm the

robustness of the results. Based on this procedure, fifty-four healthy participants were

recruited and divided into two groups, each consisting of 27 participants, The groups were

categorized as younger-aged (YA) (27.6 ± 5.8 yrs, range: 20 - 38 yrs, 13 females) and older-

aged (OA) (66.8 ± 7.4 yrs, range: 58 - 80 years old, 12 females). All participants were naïve

concerning our visuomotor learning task, had a normal or corrected-to-normal vision, no

history of musculoskeletal or neurological disorders, and were right-handed according to the

Edinburgh Handedness Inventory (Oldfield, 1971). Before data collection, all participants in

the OA group underwent a Mini-Mental State Examination (MMSE) with a cutoff score of 25

points, as scores above 25 points indicate intact cognition (Folstein et al., 1975). All

participants in the OA group scored above 25 points (29.1 ± 1.2 points), leading to older adults

having normal cognitive functions, including orientation, memory, attention, language, and

visuospatial skills. However, four out of 27 participants from each group were excluded from

the final data analyses due to the failure of the motor performance or electrical noise in the

EEG data that was difficult to remove. This paper therefore reports data from 23 participants in

each of the groups. This experimental protocol was approved by the local ethics committee of

the National Institute for Physiological Sciences in accordance with the guidelines established

in the Declaration of Helsinki. Before data collection, written informed consent was obtained

from all participants.

Experimental task and protocols

Experimental protocol, trial structure, and apparatus are illustrated in Figure 1. The

electromagnetically shielded experimental room used in this study was completely darkened

to minimize the availability of any visual information that was not projected on a PC monitor.

Participants were seated 120 cm in front of a 24.5-inch (1920 x 1080 pixels resolution)

computer display comfortably positioned on a chair with the right forearm supported by an

armrest and a chin resting comfortably on a chin rest to ensure a consistent position

throughout the experimental session.

Prior to all data collection, participants underwent two sets of resting-state EEG

recordings, involving 3-minute maintenance of the resting state while keeping their eyes open

and closed, respectively, without focusing on any thoughts. During the eyes-open condition,

participants were instructed to see the fixation cross placed at the center of the monitor.

Following this, participants performed three times of maximal voluntary contraction (MVC) in

the right index and little fingers, respectively. The maximum force was defined as an MVC

value and used to determine cursor control during the task.

On each trial, the target moved straight ahead at a constant speed of 10.8 cm/s in one of

five directions (10, 11, 12, 1, or 2 o'clock) in a randomized order within each session.

Participants were instructed to track a moving target on the screen by controlling the cursor as

accurately as possible. Our experimental system allowed participants to freely control cursor

speed and direction by pressing two force-torque transducers (USL06-H5, Tec Gihan, Co.

Ltd., Japan) using their right index and little fingertips (Figure 1A). Our task design was

inspired by O'Sullivan et al. (O'Sullivan et al., 2009), who demonstrated that the distribution of

coordinated force made by the index and little fingers is greater when heterogeneous effectors

(e.g., index and little fingers) are used together than when homonymous combinations are

used. This finding implies that the nervous system must adaptively coordinate the muscles by

integrating both effort and variability. By employing a non-habitual effector like the little finger,

we investigated how the brain flexibly learns force mapping in unfamiliar coordination

contexts. Force signals were amplified via an amplifier device (GDA-06B, Tec Gihan, Co., Ltd.,

Japan) and then stored on a computer via a NI-DAQ device (USB-6002, National Instruments,

United States) at a sampling frequency of 1 kHz with analog-to-digital conversion. The index

finger controls leftward cursor movement, while the little finger is responsible for rightward

cursor movement. Therefore, the cursor direction was determined by the vector component

calculated from the force applied by the index and little fingers. Cursor velocity was also

controlled by varying the pressure applied to the force-torque transducers.

Visual cues, cursor control, cursor and target positions on the screen, and trigger signals

sent to the EEG amplifier were managed by a dedicated software package (Graphical Design

Lab, Japan), implemented in National Instruments LabVIEW software (version 2019, National

Instruments, United States).

For the experimental protocol, participants were asked to learn the force mapping in

their right index and little fingers through an error-based visuomotor adaptation task. The task

Figure 1: Experimental setup and trial structure. **A.** Participants were instructed to control a yellow cursor by pressing force transducers with their right index and little fingertips. **B.** Upon the ready cue (when the red open circle appeared), the yellow cursor was displayed on a monitor as the Go cue. After receiving the Go cue, participants began tracking the yellow target to keep the red cursor on the target throughout the trial. **C.** The cursor direction controlled by a participant was determined using a composite vector related to the force exerted in the z-direction output from the index and little fingertips. At the end of each trial, they received binary feedback on their performance, with green or red squares indicating success or failure, respectively. The two-dimensional space distance (i.e., horizontal and vertical) between the moving target and the cursor was used as an error index. We employed an error-based visuomotor adaptation task with five different target directions (10, 11, 12, 1, or 2 o'clock direction). **D.** In the first 50 trials, participants controlled the cursor with a force sensor sensitivity ratio of 1:1 for the index and little fingers during the baseline session. In the next 100 trials, which corresponded to the adaptation session, the force sensor sensitivity ratio was changed from 1:1 to 1:3 for the index and little fingers. After the 100 trials were completed, participants took a 10-minute break. Lastly, participants completed aftereffect testing consisting of 50 trials during which the force sensor sensitivity ratio was returned to the baseline session value (i.e., clear the perturbation).

began with a familiarization period consisting of null movement trials (no added sensor

perturbation, approximately 25 trials). After the familiarization, we asked participants to

complete a total of 200 trials. The experiment consisted of three conditions: baseline

(unperturbed), adaptation (perturbed), and aftereffect. As shown in Figure 1B, at the beginning

of each trial, the target circle appeared after a 5-second preparation period, signaling the "Go

Cue". Participants were asked to accurately track the target by controlling the cursor. At the

end of each trial, they received binary feedback on their performance, with green or red

squares indicating success or failure, respectively. Figure 1D illustrates the summary of the

experimental schedule. In the baseline condition consisting of 50 trials, participants tracked

the moving cursor with a force sensor sensitivity ratio of 1:1 between the index and little

fingers. During the adaptation phase, participants completed 100 trials with the force sensor

sensitivity ratio of 1:3 between the index and little fingers, requiring them to adapt to the de

novo force mapping. After a 10-minute break, aftereffects were assessed in a 50-trial session

with sensor sensitivity returned to baseline. This session was included to evaluate whether an

internal model had been sufficiently formed through motor learning, as the presence of

aftereffects is widely regarded as strong evidence of internal model formation. It is thought

that the sensorimotor system adapts to the perturbation and updates its predictive control

mechanisms, including an internal model (Salomonczyk et al., 2013; Leow et al., 2018;

Krakauer et al., 2019; Tsay et al., 2024). Note that the luminance of the colors used for the

cursor, target, and binary feedback was uniformly set to avoid any biases related to visual

perception.

Data recording

Behavioral data. Time series of the two-dimensional target and cursor positions (x- and

y-axes) on the screen were contentiously acquired at a sampling rate of 100 Hz using the

dedicated software package as described above.

Electroencephalography. To recode neural activity during data collection, we utilized a

multi-channel EEG amplifier system (actiCHamp, Brain Products GmbH, Germany). EEG

signals were continuously recorded from 63 electrodes positioned on the scalp following the

international 10/10 system, using active electrodes embedded in a wearable elastic cap

(actiCAP, Brain Products GmbH, Germany). A ground reference electrode for data acquisition

was placed at AFz. During the data recording, all electrodes were referenced to the right

earlobe while the left earlobe was also recorded as an electrode for re-referencing.

Electrooculography (EOG) signals are widely used to detect activities of eye movements and

are available to remove eye movement-related noises from the EEG signals. EOG signals

were recorded using electrodes placed above and below the left eye for monitoring vertical

eye movements and below the left and right eyes for monitoring horizontal eye movements.

EEG and EOG signals were amplified and digitized with 24-bit resolution at a sampling rate of

1 kHz. These signals were continuously stored on a personal computer via BrainVision

recorder software (Brain Products GmbH, Germany). Skin/electrode impedance was

consistently maintained below 10 kΩ throughout the data collection.

Data analysis

Behavioral data were analyzed using custom-written code on MATLAB 2019b

(MathWorks, Natick, United States). EEG data were also analyzed using functions from

EEGLAB toolbox (Delorme and Makeig, 2004) implemented on MATLAB in combination with

custom-written code. Statistical analyses for behavioral and EEG results were implemented

using R environment (R Development Core Team, version 4.2.2, www.r-project.org/) unless

otherwise noted.

Behavioral data. A 2-dimensional space distance (i.e., screen height and width) between

the moving target (red circle) and the cursor (yellow circle) was utilized as an error index.

Throughout the error-based visuomotor adaptation task, participants had to reduce this error.

To quantify the magnitude of the error, the Euclidean distance (d) from the moving cursor to

the moving target was calculated at each timepoint (t) between the "Go" cue (t_0) and end of

the target tracking (t_{end}) at each trial. We then calculated the cumulative error for each trial

as follows:

277
$$d(t) = \sqrt{(x_{2t} - x_{1t})^2 + (y_{2t} - y_{1t})^2},$$

where x_{1t} and y_{1t} are the cursor center coordinates and x_{2t} and y_{2t} are the target center

coordinates. Subsequently, the cumulative error (CE) for motor performance within each trial

was calculated in the following formula:

$$\text{Cumulative error (CE)} = \int_{t_0}^{t_{\text{end}}} d(t)dx .$$

To examine the effects of motor learning effects, we computed the average error over five

consecutive trials, defining this as one epoch.

With regard to quantifying motor learning, the CE data obtained during the adaptation period

were first normalized to the average CE from the baseline period, i.e., null movement trials.

This correction was performed because significant individual differences in basic motor skills

among participants were observed, as shown in Supplementary Figure 1. By normalizing the

data using the average of the baseline session, we minimized the influence of these

differences in essential motor skills between participants, thereby facilitating a more accurate

assessment of the learning process itself. Following this mathematical adjustment, the

learning curve of errors across the trials in the perturbation phase was then fitted for each

participant using an exponential function (Bönstrup et al., 2020) as follows:

$$L(t) = k_1 - k_2 \cdot \left(1 - e^{-k_3 n}\right),$$

where L is the CE and k_{1-3} denotes initial performance (function intercept), maximum

performance (function plateau), and learning rate (function slope), respectively. n represents

trial number.

To further determine how brain network flexibility relates to the ability to maintain and recall a

learned motor performance, we calculated the degree of aftereffect by averaging the first five

trials immediately after the 10-minute break (see Figure 1). This is because aftereffect is a

hallmark of the effectiveness of motor learning. A high aftereffect value indicates that the

motor performance and a related internal model have been solidly learned and can be

recalled effectively.

EEG data. For preprocessing and artifact removal, consecutive EEG signals were re-

referenced offline to the averaged signals of bilateral earlobes and segmented into 14-second

epochs from 5 s before to 9 s after the "Go" cue. These segmented EEG data underwent

band-pass filtering between 1-50 Hz while eliminating 60 Hz power line noise. Independent

component analysis (ICA) was then applied to the filtered EEG signals associated with eye

movements, blinks, and muscle contractions. The ICA-derived noise components were

automatically identified using the ADJUST algorithm (Mognon et al., 2011). Across the

individuals, at least two and up to six ICA components were removed from the filtered EEG

signals. As a final preprocessing step, we applied the current source density transformation

via the current source density (CSD) toolbox (version 1.1), utilizing the spherical spline

algorithm (Perrin et al., 1989; Kayser and Tenke, 2006). This step minimizes the impact of

volume conduction on EEG data. Subsequently, artifact-corrected EEG data were used for

brain oscillation analysis. Given the necessity of maintaining the sequence of motor learning

processes, no trials were removed.

To quantify functional brain networks and their flexibility, we utilized a neural oscillation

with a phase-based approach for the analysis of neural oscillations (Lachaux et al., 1999;

Varela et al., 2001; van Diessen et al., 2015). This approach aligns with the dynamical

systems observed in the brain (Lachaux et al., 1999), where phase synchrony of neural

activity significantly contributes to information processing within the brain (Rodriguez et al.,

1999; Varela et al., 2001; van de Vijver and Cohen, 2019). First, to decompose the single-trial

EEG signals into time-frequency representations of the instantaneous phase, we applied

complex Morlet wavelet that can be defined as the product of a complex sine wave and

Gaussian window (Tallon-Baudry et al., 1996; Cohen, 2019). The CSD-transformed EEG

signals were convoluted by a Morlet wavelet function $w(\tau, f)$:

331
$$w(\tau, f) = e^{2i\pi f\tau} e^{\frac{-\tau^2}{2\sigma^2}},$$

332   where i is the imaginary unit. τ and f denote time points and the center frequency,

333   respectively. σ is the width of the Gaussian, which is defined as

334
$$\sigma = \frac{n_{co}}{2\pi f}.$$

The parameter n_{co} is defined as the number of cycles and represents a nontrivial choice in

time–frequency analysis. In the present study, we set n_{co} = 3 and f increased from 1 to 50Hz

in 1-Hz steps (Lachaux et al., 2000). It is well established that this parameter governs the

trade-off between temporal and frequency precisions, such as that a larger number of cycles

results in lower temporal resolution, and vice versa. A smaller number of n_{co} enables more

precise localization of temporal dynamics compared to a larger number. Our primary aim is to

capture transient changes in neural activity within a time-limited window during motor

execution associated with learning. Drawing on established knowledge (Cohen, 2014), we

therefore selected a smaller number of cycles. We then calculated the instantaneous phase

synchrony index (PSI) at each frequency for all electrode pairs using the following formulas;

345
$$\mathrm{PSI}(\tau, f) = \frac{1}{T}\left|\sum_{\tau=1}^{T} e^{i\theta(\tau)}\right|,$$

346
$$\theta_{mn}(\tau) = \varphi_m(\tau) - \varphi_n(\tau),$$

where T denotes the total number of time points corresponding to three cycles of the center

frequency, employing 90% overlapping windows, as per previous literature (Uehara et al.,

2023). The utilization of 90% overlapping windows enables the generation of time-dense

phase synchrony information, facilitating the observation of highly time-resolved brain network

dynamics during the task. $\varphi_m(\tau)$ and $\varphi_n(\tau)$ represent the instantaneous phase of the

m_{th} and n_{th} electrodes at time point τ. The PSI represents phase synchronization between

EEG signals from two different electrodes at each time point within each time window. This

index ranges between 0 and 1, where a value of 0 indicates complete randomness, while a

value of 1 denotes perfect synchrony (i.e., complete phase alignment) in EEG signals from the

two electrodes and provides valuable information about functional connectivity within the

whole brain (Lachaux et al., 2000; Varela et al., 2001).

As the final step, the whole-brain network flexibility was determined using the Louvain-

like algorithm (Blondel et al., 2008; Mucha et al., 2010), which is a mathematical method for

revealing the inherent community structure within complex networks. This approach is rooted

in network analysis, employing graph theory and has recently been applied to analyze

neuroimaging data, including fMRI and EEG, which is unveiling functional relationships

between brain network flexibility and cognition as well as action (Bassett et al., 2011, 2013;

Betzel et al., 2017; Paban et al., 2019; Uehara et al., 2023). Initially, the generalized Louvain

algorithm was employed for multilayer community detection. In the present study, a multilayer

community was defined wherein each network node (i.e., EEG electrode) in one time window

is connected to itself in neighboring (preceding or following) time windows (i.e., layers). This

step can be calculated using the following formula (Mucha et al., 2010; Bassett et al., 2013):

370
$$Q_{\text{multilayer}} = \frac{1}{2\mu} \sum_{nmlr} \left[(A_{nml} - \gamma_l P_{nml}) \, \delta_{lr} + \delta_{nm} \cdot \omega_{mlr} \right] \delta(g_{nl}, g_{mr}),$$

where Q represents the multilayer modularity index, l denotes the number of layers in the

multilayer network, and μ signifies the total edge weights encompassing both intra- and inter-

layer connections. A_{nml} represents the time-by-time $N \times N$ PSI-derived functional

connectivity matrices (where N denotes the number of EEG electrodes) between nodes n

and m (i.e., PSI value). P_{nml} signifies the weight of the edge between nodes n and m

under the null model, while γ_l denotes the structural resolution parameter that defines the

weight of the intralayer connections. g_{nl} represents the community assignment of node n in

layer l while g_{mr} is the community assignment of node m in layer r. ω_{mlr} denotes the

connection strength between nodes across two layers. In this study, the parameters ω and γ

were set to 1, constituting their default values as per previous literature (Bassett et al., 2013).

$\delta(g_{nl}, g_{mr})$ corresponds to the Kronecker delta, taking a value of 1 if $g_{nl} = g_{mr}$ and a value

of 0 if $g_{nl} \neq g_{mr}$. Subsequently, we determined brain network flexibility by counting how often

a node altered its module assignment throughout time, adjusting for the total potential

changes. Given the heuristic nature of this algorithm, we ran the analysis 100 times and

selected the best clustering based on the maximum Q value. Brain network flexibility was

averaged in the theta (4-8 Hz), alpha (9-13 Hz), and beta (14-30 Hz) frequency bands across

all EEG electrodes to capture the global flexibility of the whole cortical regions on every single

trial. This flexibility index ranges from 0 to 1, where 0 indicates no change in communities,

while 1 indicates a change in community across every layer. It is well accepted that the

Louvain algorithm has a heuristic in nature. We therefore performed 50 repetitions of the

algorithm for each participant. Before averaging across trials, the flexibility of the whole brain

network was averaged within each trial across predefined frequency bands during the 5-

second preparation period immediately preceding the onset of action and during the 3-second

motor execution period. We presumed that the neural systems dynamically and flexibly

reconfigure brain networks not only during motor execution but also during the preparation

period throughout learning. Previous human behavioral studies demonstrated that the

preparation of motor execution during a learning process plays a critical role in the ability to

learn in an adaptation task (Haith et al., 2015). Preparatory neural activity patterns have the

potential to retain motor memory (Sun et al., 2022). Focusing more directly on brain network

flexibility, preparatory brain network flexibility for musical playing is necessary for feedback

control, rather than feedforward control, in skilled motor performance by expert musicians

(Uehara et al., 2023). These previous studies motivated us to investigate how the flexibility of

neural preparatory activity affects motor learning ability across different ages. Moreover, we

quantified temporal changes in brain network flexibility throughout the trials during adaptation

to identify how inter-trial brain network flexibility changes within the adaptation period and how

these changes relate to motor learning ability with age. We performed this analysis to test the

assumption that reduced flexibility may stabilize brain networks optimized by learning and

enhance focus on relevant neural signals. This hypothesis is supported by a previous human

EEG study, which suggests that the suppression of trial-by-trial neural variability is closely

associated with perceptual abilities (Arazi et al., 2017). To test this, we calculated the

differences in brain network flexibility values between the averages of the first and last 5 trials

according to the predefined frequency bands at the individual level.

To enhance the interpretability of the physiological aspects of brain network flexibility, we

first computed the multi-channel Lempel-Ziv complexity (LZC) index (Lempel and Ziv, 1976;

Fernández et al., 2011). This index quantifies the dynamical complexity generated by brain

networks in terms of randomness and compressibility. There is a possibility that our

mathematical approach merely captures the apparent flexibility of brain networks. In other

words, even if network nodes transition frequently within a small number of modules, our

approach might incorrectly determine that the brain network is highly flexible. We assessed

LZC in each frequency band across the group to rule out this false detection. Multi-channel

LZC was calculated for the preparatory period of each trial using custom-written code

available in the MATLAB environment. We then calculated the trial-averaged LZC at the

individual level. In general, higher LZC values (normalized between 0 and 1) reflect greater

randomness. Additionally, although the present study has a potential limitation in

topographical interpretation since we did not perform source localization analysis, looking at

the topographical characteristics of each network module will help in biologically interpreting

the obtained results. To do so, leveraging insights into the community affiliations of each node

(i.e., EEG channel) over the analytic time window, we visualized each network module

associated with brain network flexibility in each group using a hierarchical clustering algorithm

(function *cluster* in MATLAB).

Statistics

Motor learning profiles. To test age-group differences in the motor learning profiles

between the YA and OA groups, we used an independent-sample t-test for the learning slope

and aftereffect, respectively.

Brain network flexibility. To test age-group differences in brain network flexibility during

the resting state, we conducted a two-way repeated measures ANOVA with a mixed design on

each resting EEG data with the AGE (2 levels: YA and OA) as the between subject-factor and

FREQUENCY (3 levels: theta, alpha, and beta) as the within subject-factor. Next, a three-way

repeated-measures ANOVA with a mixed design was conducted on the EEG data obtained

during the visuomotor task, with AGE (2 levels: YA and OA) as the between-subject factor and

STATE (2 levels: preparation and motor execution) and FREQUENCY (3 levels: theta, alpha,

and beta) as the within-subject factors. Moreover, we are interested in the extent to which

changes in brain network flexibility are associated with the progression of motor learning. To

test this, a four-way repeated measures ANOVA for mixed design with the AGE (2 levels: YA

and OA) as the between subject-factor and STATE (2 levels: preparation and motor

execution), FREQUENCY (3 levels: theta, alpha, and beta) and PHASE (2 levels: mean of the

initial and final 5 trials during the perturbation session) as the within subject-factors was

conducted. A Greenhouse-Geisser correction for a series of ANOVA was used when the

assumption of sphericity was violated. Post hoc explorations using Holland-Copenhaver

correction for multiple comparisons were performed if a significant effect was detected from

the ANOVA (Holland and Di Ponzio Copenhaver, 1988).

Figure 2: Behavioral results. **A.** Group-averaged performance error normalized by the baseline value throughout the data collection. Solid lines and shaded areas showed mean and SEM for five trials of epoch across participants (n=23 of each). **B.** Learning rate was determined by the slope value (k^3) of the exponential function. Group-averaged learning rates significantly differed between the groups. A lower value indicates continued performance improvement throughout the adaptation period. **C.** Group-averaged aftereffect was used as a hallmark of the effectiveness of motor learning. We calculated the degree of aftereffect by averaging the first five trials after the 10-minute break. A higher value indicates that the motor performance has been solidly learned and can be recalled effectively. Each dot represents an individual, and each error bar indicates the standard error of the mean across participants. Statistical significance is denoted by ∗ p < 0.05.

Neuro-behavioral relationship. The primary goal of this study was to determine the

neural associations of preparatory brain network flexibility or inter-trial changes in brain

network flexibility across the frequency bands and the ability of motor learning, including

learning rate and aftereffect. To achieve this, we employed a least absolute shrinkage and

selection operator (LASSO) regression model, which incorporates penalized regression

(Tibshirani, 1996). LASSO regression enables the selection of explainable variables using a

weighted combination of L1 norm regularization. The LASSO regression used in the present

study was performed using the *glmnet* package implemented in R software package

(Friedman et al., 2010). The trial-averaged brain network flexibility values at theta, alpha, and

beta frequencies at the individual level were treated as independent variables. The individuals'

learning rate and aftereffect were treated as dependent variables, respectively. To ensure the

goodness-of-fit of the model, a coefficient of determination value (R^2) was used. We repeated

the regression 100 times and identified the median of R^2 across these repetitions because the

results of LASSO regression are indeed sensitive to the choice of the hyperparameter. These

repetitions can help to provide a more robust estimate of model performance. The optimal λ

was chosen by performing k-fold cross-validation and selecting the one with the minimum

generalization error. Based on this median R^2, we then reported the relevant brain network

flexibility most associated with the motor learning profiles.

474   <Figure 2 around here>

Results

Behavioral results. Overall motor learning profiles are shown in Figure 2. A

representative cursor trajectory during motor learning is depicted in Supplementary Figure 1B.

Figure 2A shows the degree of behavioral performance across motor learning observed from

the YA and OA groups. Motor learning was evaluated by the degree of baseline-normalized

performance error. This is then transformed into parameters representing each profile. As

predicted, the degree of error was attenuated along with the number of trials, and this was

visible from the first few trials after the start of the adaptation period even though baseline

accuracy (i.e., first 50 trials without the perturbation) differed between the groups

(Supplementary Figure 1A). This indicates that motor learning occurred throughout the

adaptation period in both groups. Subsequently, we found that the learning rate,

parameterized by the slope (i.e., β value) of an exponential fit per individual, significantly

differed between the groups (Figure 2B) according to an independent t-test ($t = 2.574$, $p =$

0.013, Cohen's d = 0.759). The YA group exhibited a lower slope value compared to the OA

group, indicating that the YA group was able to sustainably attenuate performance error

throughout the adaptation period. No significant difference in aftereffect was detected between

the groups (t = -0.504, p = 0.616, Cohen's d = 0.424) (Figure 2C). These results suggest that

age had minimal impact on the effectiveness of motor learning and the ability to recall motor

memory.

Brain network flexibility. We first sought to verify how brain network flexibility at the

resting state differs between both groups (Supplementary Figure 2A). A two-way ANOVA

found a significant main effect of FREQUENCY ($F_{1.9,87.2}$ = 10.53, p < 0.001, η_p^2 = 0.19), but no

significant AGE effect ($F_{1,44}$ = 0.82, p = 0.36, η_p^2 = 0.018). The significant interaction between

them was detected ($F_{1.9,87.2}$ = 3.52, p = 0.035, η_p^2 = 0.07). A posthoc test revealed that the

theta band had significantly higher brain network flexibility compared to other frequency bands

(p < 0.05, Holland-Copenhaver corrected). The alpha band tended to differ between the age

(p = 0.052), whereas no differences were observed in the other frequency bands.

Next, we tested whether age-group differences in brain network flexibility emerged

during the preparation (Figure 3A) and motor execution (Figure 3E) phases throughout the

adaptation session. A three-way ANOVA found significant main effects of the STATE ($F_{1, 44}$ =

11.11, p = 0.001, η_p^2 = 0.20) and FREQUENCY ($F_{1.48,64.9}$ = 25.66, p < 0.001, η_p^2 = 0.36), but

no main effect of AGE ($F_{1,44}$ = 0.425, p = 0.42, η_p^2 = 0.014). A significant interaction between

STATE and FREQUENCY was detected ($F_{1.27,55.9}$ = 50.97, p < 0.001, η_p^2 = 0.53). No

significant interaction effects involving the age factor were observed (all p > 0.05). Despite no

age-group differences being observed, significant differences in brain network flexibility were

observed between the preparation and motor execution periods (Theta: $F_{1,44}$ = 33.06, p <

0.001, η_p^2 = 0.42; Alpha: $F_{1,44}$ = 64.08, p < 0.001, η_p^2 = 0.59; Beta: $F_{1,44}$ = 15.74, p < 0.001,

η_p^2 = 0.26). In the present study, another important aim was to examine how the degree of

brain network flexibility changed over time throughout the adaptation period.

Figure 3B-D and Figure 3F-H illustrate how brain network flexibility changes from the

initial to the later phase over the time course of the adaptation period. The four-way ANOVA

detected significant main effects of PHASE ($F_{1,44}$ = 7.371, p = 0.009, η_p^2 = 0.143) and

FREQUENCY ($F_{1.45, 63.88}$ = 20.93, p < 0.001, η_p^2 = 0.322) but no significant main effects of

STATE ($F_{1,44}$ = 1.43, p = 0.237, η_p^2 = 0.031) or AGE ($F_{1,44}$ = 0.804, p = 0.374, η_p^2 = 0.018).

Moreover, a significant interaction between STATE and FREQUENCY was detected ($F_{1.62,71.48}$

= 27.58, p < 0.001, η_p^2 = 0.385), whereas no other statistically significant interactions were

detected (all p > 0.05). To further examine the significant interaction observed in the four-way

ANOVA, follow-up post hoc tests were conducted. Brain network flexibility was significantly

modulated depending on the initial or late phases of adaptation ($F_{1, 44}$ = 13.22, p < 0.001, η_p^2

= 0.23). Brain network flexibility in the alpha frequency band significantly decreased toward

the end of the adaptation period ($F_{1, 44}$ = 23.46, p < 0.001, η_p^2 = 0.34). On the other hand,

during motor execution, there was no significant difference in brain network flexibility between

the initial and late phases of the adaptation period ($F_{1, 44}$ = 0.63, p = 0.428, η_p^2 = 0.014).

Together, these findings suggest that the brain network became more rigid as learning

progressed, potentially reflecting that learning was accompanied by a transition from an

exploration to an exploitation strategy.

Gaining additional insights into the characteristics of the functional brain network is

crucial for interpreting the present findings. We therefore assessed how many network

modules were mediated by the flexible nodes because there was a concern that the network

could be considered highly flexible even if each node was only moving back and forth

between a limited number of modules. To address this concern, the number of modules

(Figure 4A) and network complexity (Figure 4B), as indicated by LZC, were reported,

respectively. Figure 4B shows the LZC values during the preparation and motor execution

periods throughout adaptation. The three-way ANOVA for the LZC values detected significant

main effects of STATE ($F_{1, 44}$ = 174.3, p < 0.001, η_p^2 = 0.798) and FREQUENCY ($F_{1.37,60.12}$ =

221.6, p < 0.001, η_p^2 = 0.834) as well as significant interaction between them ($F_{1.53,67.11}$ = 55.71,

Figure 3: Group-averaged brain network flexibility during the preparatory period **(A)** and motor execution **(E)** in the visuomotor learning task in response to the predefined frequency bands. A higher value indicates greater flexibility among the networks. **B-D.** Changes in group-averaged brain network flexibility during the preparatory period from the initial to the late trials were compared for each frequency band (C: Theta, D: Alpha, and E: Beta). **E-H.** show similar patterns, but the results correspond to the motor execution. Each dot represents an individual, and each error bar indicates the standard error of the mean across participants. Statistical significance is denoted by ∗∗ p < 0.001.

543   *p* < 0.001, η_p^2= 0.55). More concretely, the LZC values in motor execution were significantly lower

544   than those in preparation across all frequencies. In contrast, no significant effects or interactions

545   involving the factor AGE were observed (all p > 0.05). Together, although the properties of the

Figure 4: Characteristics of brain network structures. **A.** Number of modules in the extracted brain networks for predefined frequency bands under the preparatory and motor execution period during the motor learning. **B.** Comparisons of brain network complexity. A higher value indicates that the extracted brain networks are complex and contain many unique patterns. Each dot represents an individual, and each error bar indicates the standard error of the mean across participants. Statistical significance is denoted by ∗∗ p < 0.001.

546 brain network differed across task states and frequencies, the age-related differences in brain

547 network flexibility during adaptation did not originate from differences in network complexity.

548   For the resting state, as shown in Supplementary Figure 2B, the two-way ANOVA for the

549 LZC values detected significant main effects of AGE ($F_{1, 44}$ = 9.77, p = 0.003, η_p^2 = 0.181) and

550 FREQUENCY ($F_{1.45, 63.71}$ = 619.2, p < 0.001, η_p^2 = 0.933) and their significant interaction

551 ($F_{1.45, 63.71}$ = 20.49, p < 0.001, η_p^2 = 0.31). The OA groups exhibited higher dynamical

552 complexity, generated by brain network flexibility, than the YA group, suggesting that aging

may be accompanied by greater randomness in resting-state brain network dynamics. The

post hoc test revealed significant age-group differences between the alpha and beta bands,

but not the theta band (Theta: $F_{1,\,44}$ = 2.48, p = 0.122, η_p^2= 0.05; Alpha: $F_{1,\,44}$ = 12.33, p =

0.001, η_p^2= 0.218; Theta: $F_{1,\,44}$ = 18.67, p < 0.0001, η_p^2= 0.298).

Neuro-behavioral relationship. Figures 5 and 6 summarize the relationships between

brain network flexibility and the learning properties based on the results of the LASSO

regression analyses. For clarity, the R^2 values obtained from the regression analyses are

presented as percentages using a pie chart in the Figure 5 and 6. Figure 5A shows the extent

to which brain network flexibility during preparation throughout the adaptation period predicted

the behavioral outcomes. Figure 5 A shows that brain network flexibility across the entire

frequencies accounted for 19% of the aftereffect in YA, whereas this relationship was scarcely

explainable in OA (2%). Interestingly, the learning rate in OA was partially explained by brain

network flexibility in the theta and beta bands (11%). Figure 5B shows the relationships

between changes in brain network flexibility toward the end of the adaptation period and the

behavioral outcomes. We found that the degree of change in brain network flexibility during

the adaptation period predicted learning even more strongly. In the YA group, this relationship

was highly explainable for a substantial portion of the aftereffect (51%), whereas in the OA

group, it was primarily associated with the learning rate (25%). On the other hand, brain

network flexibility during motor execution predicted only the learning rate, but not the

aftereffect (Figure 6). Furthermore, its explanatory contribution of brain network flexibility

during motor execution was clearly lower than during preparation across the age groups.

To summarize, we suggest that brain network flexibility during the preparatory period

contributed more to motor learning as compared to that during the motor execution period.

Brain network flexibility was effectively available to motor learning in younger adults as

compared to older adults. Our observations determine that brain network flexibility plays an

Figure 5: Results of the LASSO regression during the preparatory period. **A.** Relationships between brain network flexibility and motor learning outcomes. **B.** Relationships between changes in brain network flexibility throughout the learning period and motor learning outcomes. Bar plots display β coefficients estimated by LASSO, while pie charts summarize the explanatory power as the coefficient of determination (R^2). The small inset within the bar plot for the β coefficients provides an enlarged view of the region near zero.

active role in learning retention, i.e., aftereffect, in younger adults. However, this effect

appears to be limited in older adults.

Based on the LASSO regression results, we are further interested in understanding what

kinds of network modules contribute to forms of brain network flexibility, as this understanding

would help clarify the functional and biological significance of our findings. Figure 7 displays

the network modules observed during the preparation period throughout adaptation for each

Figure 6: Results of the LASSO regression during the motor execution. **A.** Relationships between brain network flexibility and motor learning outcomes. **B.** Relationships between changes in brain network flexibility throughout the learning period and motor learning outcomes. Bar plots display β coefficients estimated by LASSO, while pie charts summarize the explanatory power as the coefficient of determination (R^2). The small inset within the bar plot for the β coefficients provides an enlarged view of the region near zero.

age group.

<Figure 7 around here>

Discussion

The goal of the present study was to examine whether age-group differences in brain

network flexibility exist and how these differences were related to motor learning abilities by

leveraging the visuomotor learning task and multichannel EEG recordings. Our key finding

Figure 7: Extracted network modules across participants in each group. According to the brain network analysis results, five network modules, which are obtained during the preparatory period, are visualized in the alpha **(A)** and beta **(B)** frequency bands, respectively.

was that motor learning properties were tightly coupled with brain network flexibility during

motor learning, although these associations differed across the age groups (Figures 5 and 6).

More specifically, we identified the following distinctive neural–behavioral couplings. First,

brain network flexibility emerging immediately before motor execution contributed more

strongly to the acquisition of motor learning than the flexibility observed during motor

execution. Second, this effect was observed more prominently in younger adults than in older

adults. Third, the modulation of brain network flexibility immediately before motor execution

over repeated trials was essential for the consolidation of motor learning (Figure 5B). Taken

together, we propose novel evidence that brain network flexibility plays a crucial role in

acquiring and maintaining new motor execution in younger adults, but this relationship

becomes less effective with aging. We further show that preparatory brain network flexibility

could strongly predict the aftereffect rather than the learning rate, suggesting that brain

network flexibility may contribute to the formation and stabilization of implicit internal models.

For the behavioral results, both groups were able to sufficiently adapt to the perturbation

in response to changes in the sensitivity of force mapping. Interestingly, although older adults

had less precise cursor control compared to younger adults at the baseline (Supplementary

Figure 1), they exhibited motor learning effects similarly to those of younger adults as they

could adapt to the novel force mapping (from the baseline of 1:1 to the adaptation of 1:3)

(Figure 2). In principle, the learning slope quantifies the magnitude of trial-by-trial changes in

performance across repeated action, reflecting how strongly motor behavior is updated in

response to experienced errors, whereas the aftereffect reflects the maintenance and recall of

the internal model for future use (Krakauer and Mazzoni, 2011; Krakauer et al., 2019; Leech et

al., 2022). In the present study, older adults exhibited a significantly steeper learning slope

than younger adults, indicating larger behavioral adjustments across trials during the

acquisition phase. Based on our initial inspection, such a steeper slope observed in the older

adults could be interpreted as indicating faster or more efficient learning. However, we

assumed that the learning slope does not necessarily index the final level of learning or the

extent of internal model formation. This is because, by integrating the observed learning slope

with the aftereffect, we can propose different possibilities. Importantly, no age-related differences

were observed in the aftereffect in the present study (Figure 2C). The aftereffect is commonly

considered to reflect implicit sensorimotor adaptation and has been accepted as the retention

of the acquired internal model (Heuer and Hegele, 2008; Hegele and Heuer, 2010; Shadmehr

et al., 2010). The dissociation between the steeper learning slope and comparable aftereffects

suggests that the increased learning slope in older adults reflects larger trial-by-trial behavioral

updates rather than a greater magnitude of implicit learning. This pattern is in line with the

previous evidence showing that older adults rely more heavily on exploratory or strategic

adjustments during learning (Voelcker-Rehage, 2008; Cirillo et al., 2011; Berghuis et al., 2019;

Lee and Ranganathan, 2019; Bootsma et al., 2021), leading to larger immediate corrections,

while the capacity for implicit adaptation and consolidation remains preserved. Thus,

alternative learning strategies occur with aging. This observation will be discussed in the

following section by incorporating evidence from age-related differences in neural activity.

Most previous studies investigating the biological function of brain network flexibility

have relied on fMRI. However, BOLD-based measurements have limited temporal resolution

due to the inherent characteristics of the measurement principle. Although age-related

differences in brain network flexibility and their impact on motor skill have also been

investigated using fMRI (Monteiro et al., 2019), a definitive conclusion has not yet been

drawn. In the present study, our EEG-based investigation of brain network flexibility

substantially advanced the understanding of its role in motor learning by revealing distinct

patterns of brain network flexibility during preparation and motor action on a trial-by-trial basis.

We found that older adults exhibited significantly greater brain network flexibility than

younger adults in the alpha frequency band during the preparatory period of the visuomotor

learning task (Figure 3A). In principle, it has been accepted that alpha neural oscillations are

ubiquitous throughout the entire brain and are involved in multiple functional processes

(Klimesch, 1999; Başar, 2012; Bazanova and Vernon, 2014; Pascucci et al., 2025). Among

them, one of the key roles is that alpha-band activity contributes to inhibitory processes,

especially in sensory perception (Jensen et al., 2015; Clayton et al., 2018). We putatively

presumed that the elevated alpha-band brain network flexibility observed in older adults may

indicate constrained information gating, rather than enhanced adaptive reconfiguration,

potentially driven by age-related degradation of cerebellar–thalamocortical control (Mooney et

al., 2022). An alternative perspective proposes that functional networks exhibit reduced neural

network specificity (i.e., an increase in randomization) in older adults (Geerligs et al., 2014).

A key finding of the present study is the neurophysiological relationship between brain

network flexibility and learning characteristics, and the extent to which this relationship is

modulated by age. Preparatory brain network flexibility and its reduction toward the later

learning period showed the identical patterns associated with aging (Figure 5). In young

adults, a progressive reduction in brain network flexibility toward later learning was selectively

associated with stronger aftereffects, but not with learning rate. This indicates that network

stabilization primarily supports post-learning retention of the adapted internal model. Indeed,

this finding is consistent with previous evidence showing that better visual perception abilities

are associated with larger quenching of neural variability (Daniel and Dinstein, 2021). In

contrast, older adults showed the opposite pattern. The learning-related reductions in network

flexibility were strongly associated with learning rate but were unrelated to aftereffects. This

dissociation suggests an age-related shift in the functional role of brain network flexibility from

a post-learning stabilization mechanism in young adults to an online constraint on learning

efficiency in older adults. This finding aligns with a previous EEG study showing that older

adults with larger trial-by-trial auditory response variability are less accurate at sound

localization (Anderson et al., 2012), indicating that aging alters the stability of brain activity,

and these changes influence behavior. By contrast, such a strong association was not

observed during the motor execution of the visuomotor learning task (Figure 6), suggesting

that large-scale network reconfiguration across the brain primarily supports planning and post-

learning stabilization, whereas execution-related neural dynamics reflect the implementation

of already specified or planned motor commands. In other words, preparatory activity relies on

the first stage in a chain of events that generates voluntary movement (Churchland and

Shenoy, 2024; Schimel et al., 2024).

Here, we will further discuss the frequency-specific contribution of brain network

flexibility to learning properties. Despite the complexity of interpreting frequency-specific

effects, identifiable roles emerge from our findings. Our LASSO regression analyses revealed

that, in young adults, aftereffects were positively associated with alpha-band flexibility and

negatively associated with theta- and beta-band flexibility, indicating that successful retention

relies on alpha-mediated top–down gating. This pattern suggests an active exploitation of the

internal model formed during adaptation. Indeed, alpha neural oscillations can influence the

acquisition, retention, and efficiency of motor preparation (Deiber et al., 2014; Ghasemian et

al., 2017). In contrast, older adults showed a weak but positive association between

aftereffects and beta-band dynamics, with no significant contributions from the alpha or theta

bands. Given the small effect, this finding suggests that aftereffects in older adults are not

supported by an actively maintained internal model but rather by a passive persistence of the

adapted motor state. The absence of alpha- and theta-band involvement further indicates a

reduced ability to engage top–down gating or to suppress alternative control strategies after

learning. These results point to an age-related shift from a frequency-specific, multi-band

exploitation strategy in young adults to a limited, beta-dominated retention mechanism in older

adults.

In the present study, interpreting LCZ as a proxy for neural complexity helps elucidate

our findings. Intriguingly, the age-related elevations in LZC and preserved flexibility were

confined to the resting state and the motor preparation phase, disappearing during actual

execution. This selective manifestation suggests that the aging brain employs a proactive

neural strategy characterized by hyper-complex idling. By maintaining a broader repertoire of

dynamic states and high network reconfigurability before movement, older adults may create a

functional buffer that compensates for age-related neurobiological declines (Goble et al.,

2009; Coxon et al., 2014; Serbruyns et al., 2015; Frolov et al., 2020; Zapparoli et al., 2022;

Goelman et al., 2023). This heightened readiness may facilitate the successful transition into a stabilized, task-optimal state during execution, where performance becomes indistinguishable from that of younger individuals. Regardless of age, the coexistence of reduced LCZ and increased network flexibility during motor action (see Figures 3 and 4) suggests a brain state in which local neural dynamics are highly stereotyped and efficient, while large-scale functional interactions remain rapidly reconfigurable. This pattern indicates that motor execution relies on a repertoire of simple, reusable neural primitives that can be flexibly recombined in response to task demands. Moreover, we observed an increase in the number of modules alongside a decrease in intra-modular LZC, which we interpret as the emergence of specialized neural primitives for motor control. While each module becomes more dedicated to a specific sub-task (resulting in more stereotyped, lower-complexity dynamics), the overall network retains high flexibility. We therefore assumed that motor learning is characterized not by a global change in complexity, but by the efficient decomposition of tasks into simple, recruitable modules that can be dynamically rearranged.

The present study has potential limitations and caveats regarding the interpretation of the results. First, we could not mention the cerebellar function throughout the present study. It is accepted that the cerebellum plays a key role in the regulation of motor learning, including adaptation (Krakauer and Mazzoni, 2011; Tsay et al., 2024). However, non-invasive measurement of cerebellar electrophysiology is hindered by the limited spatial resolution of human scalp EEG recordings (Andersen et al., 2020). In principle, studies employing EEG delineate functional networks do not account for the cerebellum, as it is not reliably measurable using this technique. That said, the cerebellum is connected to broader cortical regions, including the motor cortex, frontal cortex, parietal cortex, and temporal lobe (Bernard et al., 2012; Schulz et al., 2014). Although we could not directly capture cerebellar function, assessing the dynamics of cortical networks, which receive extensive inputs from and send outputs to the cerebellum, is essential for understanding motor learning. Second, the present

study compared only momentary differences between the two age groups. To account for the

entire lifespan, we employed a cross-sectional study design that assesses age-related

changes across a continuum from infancy to old age.

In conclusion, the present study revealed dissociated roles of brain network flexibility in

motor learning across different age groups. Brain network flexibility during the preparatory

period, rather than motor execution, was critical for acquiring and maintaining new motor

actions in younger adults, but this coupling was attenuated with aging. Understanding flexible

brain dynamics could aid in the development of novel tools to enhance motor learning and

mitigate age-related learning limitations.

Acknowledgments

We thank Assistant Professor Akiko Nishio from NIPS for her assistance in recruiting

participants. This work was supported by JSPS KAKENHI Grant Number JP19K20103 to K.U.

and the grant of OML Project by the National Institutes of Natural Sciences (NINS program

No, OML032401) to KK.

Conflict interests

The authors declare no conflicts of interest associated with this paper.

Data and code availability

The data and code that support the findings in this study are available from the corresponding

author upon reasonable request.

References

References

Andersen LM, Jerbi K, Dalal SS (2020) Can EEG and MEG detect signals from the human cerebellum?
Neuroimage 215:116817.

Anderson S, Parbery-Clark A, White-Schwoch T, Kraus N (2012) Aging affects neural precision of
speech encoding. J Neurosci 32:14156–14164.

Arazi A, Censor N, Dinstein I (2017) Neural variability quenching predicts individual perceptual abilities.
Journal of Neuroscience 37:97–109.

Başar E (2012) A review of alpha activity in integrative brain function: fundamental physiology, sensory
coding, cognition and pathology. Int J Psychophysiol 86:1–24.

Bassett DS, Porter MA, Wymbs NF, Grafton ST, Carlson JM, Mucha PJ (2013) Robust detection of
dynamic community structure in networks. Chaos 23:013142.

Bassett DS, Wymbs NF, Porter MA, Mucha PJ, Carlson JM, Grafton ST (2011) Dynamic reconfiguration
of human brain networks during learning. Proc Natl Acad Sci U S A 108:7641–7646.

Bazanova OM, Vernon D (2014) Interpreting EEG alpha activity. Neurosci Biobehav Rev 44:94–110.

Beck D, de Lange A-MG, Maximov II, Richard G, Andreassen OA, Nordvik JE, Westlye LT (2021) White
matter microstructure across the adult lifespan: A mixed longitudinal and cross-sectional study
using advanced diffusion models and brain-age prediction. Neuroimage 224:117441.

Berghuis KMM, Fagioli S, Maurits NM, Zijdewind I, Marsman JBC, Hortobágyi T, Koch G, Bozzali M
(2019) Age-related changes in brain deactivation but not in activation after motor learning.
Neuroimage 186:358–368.

Bernard J a., Seidler RD, Hassevoort KM, Benson BL, Welsh RC, Wiggins JL, Jaeggi SM, Buschkuehl
M, Monk CS, Jonides J, Peltier SJ (2012) Resting state cortico-cerebellar functional
connectivity networks: a comparison of anatomical and self-organizing map approaches. Front
Neuroanat 6:31–31.

Betzel RF, Satterthwaite TD, Gold JI, Bassett DS (2017) Positive affect, surprise, and fatigue are
correlates of network flexibility. Sci Rep 7:1–10.

Blondel VD, Guillaume J-L, Lambiotte R, Lefebvre E (2008) Fast unfolding of communities in large
networks. J Stat Mech 2008:P10008.

Bönstrup M, Iturrate I, Hebart MN, Censor N, Cohen LG (2020) Mechanisms of offline motor learning at
a microscale of seconds in large-scale crowdsourced data. npj Science of Learning 5:1–10.

Bootsma JM, Caljouw SR, Veldman MP, Maurits NM, Rothwell JC, Hortobágyi T (2021) Neural
Correlates of Motor Skill Learning Are Dependent on Both Age and Task Difficulty. Front Aging
Neurosci 13:643132.

Braun U, Harneit A, Pergola G, Menara T, Schäfer A, Betzel RF, Zang Z, Schweiger JI, Zhang X,
Schwarz K, Chen J, Blasi G, Bertolino A, Durstewitz D, Pasqualetti F, Schwarz E, Meyer-
Lindenberg A, Bassett DS, Tost H (2021) Brain network dynamics during working memory are
modulated by dopamine and diminished in schizophrenia. Nat Commun 12:3478.

Braun U, Schäfer A, Bassett DS, Rausch F, Schweiger JI, Bilek E, Erk S, Romanczuk-Seiferth N,
Grimm O, Geiger LS, Haddad L, Otto K, Mohnke S, Heinz A, Zink M, Walter H, Schwarz E,
Meyer-Lindenberg A, Tost H (2016) Dynamic brain network reconfiguration as a potential
schizophrenia genetic risk mechanism modulated by NMDA receptor function. Proc Natl Acad
Sci U S A 113:12568–12573.

Braun U, Schäfer A, Walter H, Erk S, Romanczuk-Seiferth N, Haddad L, Schweiger JI, Grimm O, Heinz
A, Tost H, Meyer-Lindenberg A, Bassett DS (2015) Dynamic reconfiguration of frontal brain
networks during executive cognition in humans. Proc Natl Acad Sci U S A 112:11678–11683.

Cabeza R (2002) Hemispheric asymmetry reduction in older adults: the HAROLD model. Psychol Aging
17:85–100.

Cavanagh JF, Frank MJ, Klein TJ, Allen JJB (2010) Frontal theta links prediction errors to behavioral
adaptation in reinforcement learning. Neuroimage 49:3198–3209.

Churchland MM, Shenoy KV (2024) Preparatory activity and the expansive null-space. Nat Rev
Neurosci 25:213–236.

Cirillo J, Todd G, Semmler JG (2011) Corticomotor excitability and plasticity following complex
visuomotor training in young and old adults. Eur J Neurosci 34:1847–1856.

Clayton MS, Yeung N, Cohen Kadosh R (2018) The many characters of visual alpha oscillations. Eur J
Neurosci 48:2498–2508.

Cohen MX (2014) Analyzing neural time series data: Theory and practice. Mit Press.

Cohen MX (2019) A better way to define and describe Morlet wavelets for time-frequency analysis.
Neuroimage 199:81–86.

Courchesne E, Chisum HJ, Townsend J, Cowles A, Covington J, Egaas B, Harwood M, Hinds S, Press
GA (2000) Normal brain development and aging: quantitative analysis at in vivo MR imaging in
healthy volunteers. Radiology 216:672–682.

Cox SR, Ritchie SJ, Tucker-Drob EM, Liewald DC, Hagenaars SP, Davies G, Wardlaw JM, Gale CR,
Bastin ME, Deary IJ (2016) Ageing and brain white matter structure in 3,513 UK Biobank
participants. Nat Commun 7:13629.

Coxon JP, Goble DJ, Leunissen I, Van Impe A, Wenderoth N, Swinnen SP (2014) Functional Brain
Activation Associated with Inhibitory Control Deficits in Older Adults. Cereb Cortex:1–11.

Daniel E, Dinstein I (2021) Individual magnitudes of neural variability quenching are associated with
motion perception abilities. J Neurophysiol 125:1111–1120.

Deco G, Kringelbach ML (2016) Metastability and Coherence: Extending the Communication through
Coherence Hypothesis Using A Whole-Brain Computational Perspective. Trends Neurosci
39:125–135.

Deiber M-P, Sallard E, Ibañez V, Ludwig C, Barral J (2014) Aging and the lateralization of oscillatory
activities related to external and internal motor preparation. J Psychophysiol 28:136–147.

Delorme A, Makeig S (2004) EEGLAB: An open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis. J Neurosci Methods 134:9–21.

Erdfelder E, FAul F, Buchner A, Lang AG (2009) Statistical power analyses using G*Power 3.1: Tests for
correlation and regression analyses. Behav Res Methods 41:1149–1160.

Fernández A, López-Ibor M-I, Turrero A, Santos J-M, Morón M-D, Hornero R, Gómez C, Méndez MA,
Ortiz T, López-Ibor JJ (2011) Lempel-Ziv complexity in schizophrenia: a MEG study. Clin
Neurophysiol 122:2227–2235.

Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental state". A practical method for grading the
cognitive state of patients for the clinician. J Psychiatr Res 12:189–198.

Friedman JH, Hastie T, Tibshirani R (2010) Regularization Paths for Generalized Linear Models via
Coordinate Descent. J Stat Softw 33:1–22.

Frolov NS, Pitsik EN, Maksimenko VA, Grubov VV, Kiselev AR, Wang Z, Hramov AE (2020) Age-related
slowing down in the motor initiation in elderly adults. PLoS One 15:e0233942.

Fukuda Y, Uehara K (2025) Frontal midline theta power accounts for inter-individual differences in
motor learning ability. Exp Brain Res 243:1–14.

Garrett DD, Kovacevic N, McIntosh AR, Grady CL (2010) Blood Oxygen Level-Dependent Signal
Variability Is More than Just Noise. Journal of Neuroscience 30:4914–4921.

Garrett DD, Kovacevic N, McIntosh AR, Grady CL (2013) The modulation of BOLD variability between
cognitive states varies by age and processing speed. Cereb Cortex 23:684–693.

Geerligs L, Maurits NM, Renken RJ, Lorist MM (2014) Reduced specificity of functional connectivity in
the aging brain during task performance. Hum Brain Mapp 35:319–330.

Ghasemian M, Taheri H, Saberi Kakhki A, Ghoshuni M (2017) Electroencephalography pattern
variations during motor skill acquisition. Percept Mot Skills 124:1069–1084.

Goble DJ, Coxon JP, Wenderoth N, Van Impe A, Swinnen SP (2009) Proprioceptive sensibility in the
elderly: degeneration, functional consequences and plastic-adaptive processes. Neurosci
Biobehav Rev 33:271–278.

Goelman G, Dan R, Bezdicek O, Jech R (2023) Directed functional connectivity of the sensorimotor
system in young and older individuals. Front Aging Neurosci 15:1222352.

Good CD, Johnsrude IS, Ashburner J, Henson RN, Friston KJ, Frackowiak RS (2001) A voxel-based
morphometric study of ageing in 465 normal adult human brains. Neuroimage 14:21–36.

Guitart-Masip M, Salami A, Garrett D, Rieckmann A, Lindenberger U, Bäckman L (2016) BOLD
Variability is Related to Dopaminergic Neurotransmission and Cognitive Aging. Cereb Cortex
26:2074–2083.

Haith AM, Huberdeau DM, Krakauer JW (2015) The influence of movement preparation time on the
expression of visuomotor learning and savings. J Neurosci 35:5109–5117.

Hegele M, Heuer H (2010) The impact of augmented information on visuo-motor adaptation in younger
and older adults. PLoS One 5:e12071.

Hehl M, Swinnen SP, Cuypers K (2020) Alterations of hand sensorimotor function and cortical motor
representations over the adult lifespan. Aging (Albany NY) 12:4617–4640.

Hermans L, Levin O, Maes C, van Ruitenbeek P, Heise KF, Edden RAE, Puts NAJ, Peeters R, King BR,
Meesen RLJ, Leunissen I, Swinnen SP, Cuypers K (2018) GABA levels and measures of
intracortical and interhemispheric excitability in healthy young and older adults: an MRS-TMS
study. Neurobiol Aging 65:168–177.

Heuer H, Hegele M (2008) Constraints on visuo-motor adaptation depend on the type of visual
feedback during practice. Exp Brain Res 185:101–110.

Holland BS, Di Ponzio Copenhaver M (1988) Improved Bonferroni-Type Multiple Testing Procedures.
Psychological Bulletin 104:145–149.

Jensen O, Bonnefond M, Marshall TR, Tiesinga P (2015) Oscillatory mechanisms of feedforward and
feedback visual processing. Trends Neurosci 38:192–194.

Jernigan TL, Archibald SL, Fennema-Notestine C, Gamst AC, Stout JC, Bonner J, Hesselink JR (2001)

Effects of age on tissues and regions of the cerebrum and cerebellum. Neurobiol Aging 22:581–594.

Kaasinen V, Rinne JO (2002) Functional imaging studies of dopamine system and cognition in normal aging and Parkinson's disease. Neurosci Biobehav Rev 26:785–793.

Kayser J, Tenke CE (2006) Principal components analysis of Laplacian waveforms as a generic method for identifying ERP generator patterns: II. Adequacy of low-density estimates. Clin Neurophysiol 117:369–380.

Klimesch W (1999) EEG alpha and theta oscillations reflect cognitive and memory performance: a review and analysis. Brain Res Brain Res Rev 29:169–195.

Krakauer JW, Hadjiosif AM, Xu J, Wong AL, Haith AM (2019) Motor Learning. Compr Physiol 9:613–663.

Krakauer JW, Mazzoni P (2011) Human sensorimotor learning: Adaptation, skill, and beyond. Curr Opin Neurobiol 21:636–644.

Lachaux J-P, Rodriguez E, M LVQ, A L, J M, Varela FJ (2000) Studying signale-trials of phase synchronous activity in the brain. Int J Bifurcat Chaos 10:2429–2439.

Lachaux JP, Rodriguez E, Martinerie J, Varela FJ (1999) Measuring phase synchrony in brain signals. Hum Brain Mapp 8:194–208.

Lee M-H, Ranganathan R (2019) Age-related deficits in motor learning are associated with altered motor exploration strategies. Neuroscience 412:40–47.

Leech KA, Roemmich RT, Gordon J, Reisman DS, Cherry-Allen KM (2022) Updates in Motor Learning: Implications for Physical Therapist Practice and Education. Phys Ther 102 Available at: http://dx.doi.org/10.1093/ptj/pzab250.

Lempel A, Ziv J (1976) On the complexity of finite sequences. IEEE Trans Inf Theory 22:75–81.

Leow L-A, Marinovic W, de Rugy A, Carroll TJ (2018) Task errors contribute to implicit aftereffects in sensorimotor adaptation. Eur J Neurosci 48:3397–3409.

Marstaller L, Williams M, Rich A, Savage G, Burianová H (2015) Aging and large-scale functional networks: white matter integrity, gray matter volume, and functional connectivity in the resting state. Neuroscience 290:369–378.

Mognon A, Jovicich J, Bruzzone L, Buiatti M (2011) ADJUST: An automatic EEG artifact detector based on the joint use of spatial and temporal features. Psychophysiology 48:229–240.

Monteiro TS, King BR, Zivari Adab H, Mantini D, Swinnen SP (2019) Age-related differences in network
flexibility and segregation at rest and during motor performance. Neuroimage 194:93–104.

Mooney RA, Ni Z, Shirota Y, Chen R, Ugawa Y, Celnik PA (2022) Age-related strengthening of
cerebello-cortical motor circuits. Neurobiol Aging 118:9–12.

Mucha PJ, Richardson T, Macon K, Porter MA, Onnela J-P (2010) Community structure in time-
dependent, multiscale, and multiplex networks. Science 328:876–878.

Oldfield RC (1971) The assessment and analysis of handedness: The Edinburgh inventory.
Neuropsychologia 9:97–113.

O'Sullivan I, Burdet E, Diedrichsen J (2009) Dissociating Variability and Effort as Determinants of
Coordination. PLoS Comput Biol 5 Available at: http://dx.doi.org/10.1371/journal.pcbi.1000345.

Paban V, Modolo J, Mheich A, Hassan M (2019) Psychological resilience correlates with EEG source-
space brain network flexibility. Netw Neurosci 3:539–550.

Pascucci D, Menétrey MQ, Passarotto E, Luo J, Paramento M, Rubega M (2025) EEG brain waves and
alpha rhythms: Past, current and future direction. Neurosci Biobehav Rev 176:106288.

Pauelsen M, Jafari H, Strandkvist V, Nyberg L, Gustafsson T, Vikman I, Röijezon U (2020) Frequency
domain shows: Fall-related concerns and sensorimotor decline explain inability to adjust
postural control strategy in older adults. PLoS One 15:e0242608.

Perrin F, Pernier J, Bertrand O, Echallier JF (1989) Spherical splines for scalp potential and current
density mapping. Electroencephalogr Clin Neurophysiol 72:184–187.

Ranganathan R, Lee M-H, Newell KM (2020) Repetition Without Repetition: Challenges in
Understanding Behavioral Flexibility in Motor Skill. Front Psychol 11:2018.

Ranganathan R, Newell KM (2010) Emergent flexibility in motor learning. Exp Brain Res 202:755–764.

Reddy PG, Mattar MG, Murphy AC, Wymbs NF, Grafton ST, Satterthwaite TD, Bassett DS (2018) Brain
state flexibility accompanies motor-skill acquisition. Neuroimage 171:135–147.

Rodriguez E, George N, Lachaux JP, Martinerie J, Renault B, Varela FJ (1999) Perception's shadow:
Long-distance synchronization of human brain activity. Nature 397:430–433.

Roth RH, Ding JB (2024) Cortico-basal ganglia plasticity in motor learning. Neuron 112:2486–2502.

Salomonczyk D, Cressman EK, Henriques DYP (2013) The role of the cross-sensory error signal in
visuomotor adaptation. Exp Brain Res 228:313–325.

Sase T, Kitajo K (2021) The metastable brain associated with autistic-like traits of typically developing

individuals. PLoS Comput Biol 17:e1008929.

Schimel M, Kao T-C, Hennequin G (2024) When and why does motor preparation arise in recurrent
neural network models of motor control? Elife 12 Available at:
http://dx.doi.org/10.7554/eLife.89131.

Schulz R, Wessel MJ, Zimerman M, Timmermann JE, Gerloff C, Hummel FC (2014) White Matter
Integrity of Specific Dentato-Thalamo-Cortical Pathways is Associated with Learning Gains in
Precise Movement Timing. Cereb Cortex:1–8.

Serbruyns L, Gooijers J, Caeyenberghs K, Meesen RL, Cuypers K, Sisti HM, Leemans A, Swinnen SP
(2015) Bimanual motor deficits in older adults predicted by diffusion tensor imaging metrics of
corpus callosum subregions. Brain Struct Funct:1–18.

Shadmehr R, Smith MA, Krakauer JW (2010) Error correction, sensory prediction, and adaptation in
motor control. Annu Rev Neurosci 33:89–108.

Shafiei G, Zeighami Y, Clark CA, Coull JT, Nagano-Saito A, Leyton M, Dagher A, Mišic B (2019)
Dopamine signaling modulates the stability and integration of intrinsic brain networks. Cereb
Cortex 29:397–409.

Sun X, O'Shea DJ, Golub MD, Trautmann EM, Vyas S, Ryu SI, Shenoy KV (2022) Cortical preparatory
activity indexes learned motor memories. Nature:1–6.

Tallon-Baudry C, Bertrand O, Delpuech C, Pernier J (1996) Stimulus specificity of phase-locked and
non-phase-locked 40 Hz visual responses in human. J Neurosci 16:4240–4249.

Tibshirani R (1996) Regression shrinkage and selection via the lasso. J R Stat Soc 58:267–288.

Tsay JS, Kim HE, McDougle SD, Taylor JA, Haith A, Avraham G, Krakauer JW, Collins AGE, Ivry RB
(2024) Fundamental processes in sensorimotor learning: Reasoning, refinement, and retrieval.
Elife 13 Available at: http://dx.doi.org/10.7554/eLife.91839.

Uehara K, Yasuhara M, Koguchi J, Oku T, Shiotani S, Morise M, Furuya S (2023) Brain network
flexibility as a predictor of skilled musical performance. Cereb Cortex Available at:
http://dx.doi.org/10.1093/cercor/bhad298.

Uehara S, Mawase F, Therrien AS, Cherry-Allen KM, Celnik P (2019) Interactions between motor
exploration and reinforcement learning. J Neurophysiol 122:797–808.

van de Vijver I, Cohen MX (2019) Electrophysiological phase synchrony in distributed brain networks as
a promising tool in the study of cognition. In: New Methods in Cognitive Psychology (Spieler D,
Schumacher E, eds), pp 214–244 1st Edition. Routledge.

van de Vijver I, Ridderinkhof KR, Cohen MX (2011) Frontal oscillatory dynamics predict feedback
learning and action adjustment. J Cogn Neurosci 23:4106–4121.

van Diessen E, Numan T, van Dellen E, van der Kooi AW, Boersma M, Hofman D, van Lutterveld R, van
Dijk BW, van Straaten ECW, Hillebrand A, Stam CJ (2015) Opportunities and methodological
challenges in EEG and MEG resting state functional brain network research. Clin Neurophysiol
126:1468–1481.

Varela F, Lachaux JP, Rodriguez E, Martinerie J (2001) The brainweb: phase synchronization and large-
scale integration. Nat Rev Neurosci 2:229–239.

Vercillo T, Carrasco C, Jiang F (2017) Age-related changes in sensorimotor temporal binding. Front
Hum Neurosci 11:500.

Voelcker-Rehage C (2008) Motor-skill learning in older adults—a review of studies on age-related
differences. Eur Rev Aging Phys Act 5:5–16.

Wu HG, Miyamoto YR, Nicolas L, Castro G, Ölveczky BP, Smith MA, Biology E (2014) Temporal
structure of motor variability is dynamically regulated and predicts motor learning ability. Nat
Neurosci 17:312–321.

Yoshimura N, Tsuda H, Aquino D, Takagi A, Ogata Y, Koike Y, Minati L (2020) Age-related decline of
sensorimotor integration influences resting-state functional brain connectivity. Brain Sci 10:966.

Zapparoli L, Mariano M, Paulesu E (2022) How the motor system copes with aging: a quantitative meta-
analysis of the effect of aging on motor function control. Commun Biol 5:79.

Supplementary Materials

Supplementary Figure 1: Baseline motor performance **(A)** and a typical cursor trajectory during adaptation **(B)**. **A**. Group-averaged error at the baseline across the fifty trials. Each dot represents data from an individual, and each error bar indicates the standard error of the mean across participants. **B**. A typical cursor trajectory during adaptation from a representative participant in the YA group. The deviation of the cursor was decreased consistently across the 100 trials.

Supplementary Figure 2: Group-averaged brain network flexibility during an eyes-open resting state in response to the predefined frequency bands **(A)** and their network properties (**B**: LZC, **C**: Number of network modules). Each dot represents an individual, and each error bar indicates the standard error of the mean across participants. Statistically significant differences between the frequency bands are denoted by ∗∗ p < 0.001. Statistically significant difference between the age groups is indicated by † p < 0.05.