Title: Targeted therapy of oncomicrobe *F. nucleatum* with bioengineered probiotic expressing
guided antimicrobial peptide (gAMP)

Ankan Choudhury[1,2], Colin Scano[1], Allison Barton[1], Christopher M. Kearney[1], K. Leigh Greathouse[1,2]

Author Affiliations

[1]Department of Biology, College of Arts & Sciences, Baylor University, Waco, Texas
[2]Department of Human Science and Design, Robbins College of Health and Human Sciences,
Baylor University, Waco, Texas
[3]Department of Public Health, Robbins College of Health and Human Sciences, Baylor University,
Waco, Texas

Abstract

Colorectal cancer (CRC) is a leading cause of cancer-related mortality, with *Fusobacterium*
nucleatum (*F. nucleatum*) identified as a key contributor to its progression. This study explores a
novel targeted therapy using bioengineered probiotics expressing guided antimicrobial peptides
(gAMPs) to selectively inhibit *F. nucleatum*. We engineered *Lactococcus lactis* MG1363 to express
gAMPs derived from Ovispirin and Cathelin-related peptide SCF, linked to a Statherin-derived guide
peptide (YQPVPE) that binds specifically to the *F. nucleatum* membrane porin FomA.

Our results demonstrate that the Statherin-derived guide peptide enhances the binding affinity to *F.*
nucleatum, significantly increasing the preferential attachment compared to control peptides. In
vitro assays revealed that both unguided and guided AMPs effectively inhibited biofilm formation in
F. nucleatum, with gAMPs showing reduced toxicity against non-target bacteria (*Bacteroides fragilis*
and *Escherichia coli*). The gAMPs were also more effective in modulating growth kinetics, exhibiting
selective toxicity towards *F. nucleatum* at lower concentrations.

Co-culture experiments in a simulated human gut microbiome demonstrated that the gAMP
probiotic maintained microbial diversity while effectively reducing *F. nucleatum* abundance.
Quantitative PCR and 16S rRNA sequencing confirmed that gAMP treatment preserved the richness
of the microbiota, contrasting with significant dysbiosis observed in control samples.

These findings support the potential of engineered probiotics as a targeted therapeutic approach to
combat *F. nucleatum*-associated CRC. By leveraging the specificity of Statherin-derived peptides,
this strategy not only addresses the pathogenicity of *F. nucleatum* but also mitigates the adverse
effects of traditional antimicrobial therapies on beneficial gut microbiota. Future studies will
explore the clinical applicability of this approach in CRC management and its impact on overall gut
health.

Introduction

Overall, colorectal cancer (CRC) accounts for approximately 152,810 new cases and 53,010 deaths
expected in 2024, making it the fourth most lethal cancer in the U.S. (SEER). While traditionally
thought of as a disease of older age, CRC has rapidly become the second leading cause of cancer-
related death in people under 50 [1, 2]. The American Cancer Society (ACS) revealed that 20% of
cancer diagnoses in 2019 occurred in patients under the age of 55, roughly twice the rate seen in
1995 [3]. Additionally, the incidence of advanced-stage disease in individuals under 50 increased by

approximately 3% per year. The ACS projected that in estimated 152,810 new cases and 53,010
deaths are expected in 2024 among those under 50 [3].

Chronic inflammation, diet, and environmental exposures, in combination with somatic mutations,
are strong drivers of colon cancer [4]. The microbiome is steadily emerging as a major factor
mediating the duration and polarity of the inflammatory response [5]. The oral pathogen,
Fusobacterium nucleatum (F. nucleatum), has been identified as a contributor to inflammation,
inflammatory bowel diseases, and cancer. Expression of virulence factors, including FomA, FadA,
RadD and Fap2 in *F. nucleatum* have established links with invasion of host cells, aggregation, biofilm
formation, and trigger multiple pathways of carcinogenesis [6–12]. *F. nucleatum* has consistently
been one of the most abundant bacteria in colon adenomas and in colon adenocarcinomas and is
associated with colon cancer promotion [13–15]. Moreover, nearly genetically identical strains of *F.*
nucleatum are present in both the primary tumor and distant metastases from the same patient with
CRC [16]. Recent pioneering research has also uncovered that a particular clade of *F. nucleatum*
subspecies *animalis* is the predominant microbe in CRC tumors, which are phenotypically and
genotypically distinct from the clades found in the healthy gut microbiota [17]. These clades are also
more likely to be highly organized in microniches with immune and epithelial cell functions that
promote cancer progression [18]. However, the effects of specifically killing *F. nucleatum* in the
presence of a full complement of commensal and pathogenic gastrointestinal microbiota on
inflammation, tumorigenesis, and metabolism have been understudied. Therefore, understanding
how the removal of *F. nucleatum* from the intestinal environment influences microbial community
function is essential for determining its impact on pathogenicity, the early colonization of other
oncomicrobes, and the progression of colorectal cancer (CRC) as well as responses to treatment.

Antimicrobial resistance (AMR) is a side effect of cancer treatment, developing in approximately 26%
of patients after undergoing chemotherapy and receiving prophylactic antibiotics[19, 20].
Furthermore, AMR is a major issue world-wide, with the most recent data showing approximately
1.27 million deaths attributable to AMR [21–23]. There is an urgent need to overcome this burden by
using more targeted approaches like the one proposed in this study, guided antimicrobial peptides
(gAMPs or AMPs). Guided antimicrobial peptides (AMPs) hold significant promise in combating
multidrug-resistant bacteria, offering a more specific and effective approach to infection control.
Unlike broad-spectrum antibiotics, targeted AMPs can be designed to selectively bind to specific
microbial membranes, reducing off-target effects and minimizing harm to beneficial bacteria (Xu et
al., 2024). This specificity not only enhances efficacy but also mitigates the rise of resistance (Mba
and Nweze, 2022). Additionally, AMPs have the advantage of rapid product development compared
to the slower pace of traditional antibiotics. The extensive databases available for AMPs and the ease
of genetically modifying them for increased specificity, such as through peptide tags, make them an
ideal alternative to conventional antibiotics.

One innovative strategy involves using engineered probiotics to express and release guided AMPs
(gAMPs). This approach may aid in the recovery of species richness—unlike antibiotic treatments—
and reduce overall microbial dysbiosis [24]. *Lactococcus lactis* MG1363, a widely used probiotic,
serves as an effective delivery platform and is the first patented engineered probiotic to deliver
therapeutic molecules to human patients [25]. Previous research has demonstrated the versatility
of *L. lactis* as a delivery system for various antimicrobial peptides, including camel milk lactoferrin
derivatives (lactoferrampin and lactoferricin) [26] and gram-negative specific amphibian-derived

AMPs [27, 28]. Beyond antimicrobial applications, *L. lactis* has been utilized to deliver systemic
therapeutics for conditions such as pulmonary inflammation [29, 30], inflammatory bowel disease
[31–34], and more recently, for cancer immunotherapy [35].

Seminal research demonstrates that biofilm formation is a biomarker of colorectal cancer (CRC) and
oral squamous cell carcinoma, as well as a potential promoter of local inflammation. *Fusobacterium*
nucleatum is a key member of these biofilms. The outer membrane porin, FomA, serves as the
anchor binding site on *F. nucleatum* for attachment to the oral mucosa through the human salivary
protein statherin. This attachment allows *F. nucleatum* to act as a bridging species for the
attachment of secondary colonizers, such as *Porphyromonas gingivalis*, thereby promoting biofilm
formation [36]. Targeting FomA with our gAMP offers the additional benefit of reducing biofilm
formation at mucosal sites, resulting in a decreased risk of inflammation and disease. Furthermore,
this targeting likely exerts added selection pressure to reduce FomA expression in these strains.
However, rigorous testing of the downstream corollary effects is essential to ensure no adverse
outcomes arise.

To initiate the development of *F. nucleatum*-targeting gAMPs, we utilized the antimicrobial peptide
database DBAASP (v3.0) to identify candidate AMPs active against *F. nucleatum*. Ovispirin [37] and
Cathelin-related peptide SCF (CPSC5) [38] were selected as promising candidates. Through a
comprehensive literature search, we identified a peptide ligand (YQPVPE) derived from the human
salivary protein statherin, which specifically binds the *F. nucleatum* membrane porin FomA [39].
We then constructed the gAMPs by linking the ligand to the AMP using a flexible triple glycine linker,
resulting in Stat-Ovispirin and Stat-CPSC5 for downstream analysis (Table S1). Synthetic peptides
were employed in initial assays to confirm target specific antimicrobial activity. Our results
demonstrate that the Statherin-derived guide peptide enhances the binding affinity to *F. nucleatum*,
significantly increasing the preferential attachment compared to control peptides. In vitro assays
revealed that both unguided and guided AMPs effectively inhibited biofilm formation in *F.*
nucleatum, with gAMPs showing reduced toxicity against non-target bacteria (*Bacteroides fragilis*
and *Escherichia coli*). The gAMPs were also more effective in modulating growth kinetics, exhibiting
selective toxicity towards *F. nucleatum* at lower concentrations.

Building on this foundation, our study focused on engineering *L. lactis* MG1363 to synthesize and
deliver gAMPs specifically targeting carcinogenic pathogens. The open-reading frames (ORFs) for
both the targeted and non-targeted AMPs were cloned into plasmid pMSP3545 [40], which contains
the PNisA promoter [41] inducible by Nisin—an endogenous peptide produced by *L. lactis*. This
promoter facilitates controlled expression of the heterologous proteins and peptides. The plasmids
were placed downstream of a *L. lactis*-specific secretion signal to create the bioengineered probiotic
strains. Nisin-inducible expression systems for heterologous proteins and peptides in both murine
and human guts have been well established for over two decades [31, 32, 34, 42, 43], yet their
application in delivering guided antimicrobial agents against carcinogenic pathogens remains
unexplored. When the resultant bioengineered probiotics were co-cultured with both the target *F.*
nucleatum, and other non-target bacteria, the probiotic expressing gAMPs showed targeted toxicity
towards *F. nucleatum* with significant reduction in titer of the target bacteria at same probiotic doses
compared to the non-target bacteria.

Co-culture experiments in a simulated human gut microbiome demonstrated that the gAMP
probiotic maintained microbial diversity while effectively reducing *F. nucleatum* abundance.

Fig 1. Experimental Overview. The probiotic *L. lactis* was cloned with a Nisin-inducible plasmid with the ORF of the AMP/gAMP downstream of it, to express and secrete the gAMP in vitro where the statherin derived guide peptide will allow the AMP to preferably attach to the FomA membrane protein on *F. nucleatum* and exhibit targeted antimicrobial activity. The bioengineered probiotic was assayed by either co-culturing with a single *Fusobacterium* and non-*Fusobacterium* species and analyzing by species specific qPCR (top) or co-culturing with a fecal microbiome derived polymicrobial community spiked with *F. nucleatum* and analyzing through 16s rRNA metagenomics (bottom).

Quantitative PCR and 16S rRNA sequencing confirmed that gAMP treatment preserved the richness
of the microbiota, contrasting with significant dysbiosis observed in control samples.

Our approach was inspired by our previous study that developed precision therapy
against *Helicobacter pylori* in mice [44]. In that project, *L. lactis* was engineered with plasmids
containing open-reading frames (ORFs) for both AMPs and gAMPs—where gAMPs were guided by
ligand-peptides targeting the VacA toxin of *H. pylori*. These were expressed under a pH-inducible
promoter and secreted in the stomach environment via a secretory signal. The bioengineered
probiotic expressing gAMP demonstrated preferential toxicity against *H. pylori* over other tested
commensal bacteria in vitro, whereas the probiotic expressing AMP alone did not show this
specificity. *In vivo* experiments revealed that both AMP- and gAMP-expressing probiotics effectively
reduced *H. pylori* infection and titers in mice when used as either prophylactic or therapeutic agents.
Furthermore, microbiome sequencing and analysis indicated that gAMP-expressing probiotics were
superior in preserving stomach microbiome diversity compared to their AMP-expressing
counterparts.

This study advances the application of engineered *L. lactis* as a targeted delivery system for gAMPs,
offering a promising strategy for combating pathogenic bacteria like *F. nucleatum* while maintaining
beneficial microbial communities. By specifically targeting biofilm-forming pathogens associated
with colorectal cancer and oral squamous cell carcinoma, our approach not only mitigates infection
and inflammation but also preserves the integrity of the host microbiome, presenting a significant
step forward in precision antimicrobial therapy.

Materials and Methods

Bacterial strains and plasmids

The strains of target pathogen *F. nucleatum* use were *F. nucleatum* SB-CTX3Tcol3 derived from CRC tumor biopsy; given to us as a gift from Dr. Suan Bullmann's group (MD Anderson Cancer Research Center, TX) and *F. nucleatum* subsp. *nucleatum* Knorr (ATCC 25586) derived from Oral lesion; sourced from ATCC (VA). Non-target commensals used for the experiment were *B. fragilis* - *B. fragilis* ATCC 43858 (Enterotoxigenic/ETBF) and *B. fragilis* ATCC 25285 (Non-Enterotoxigenic/NTBF); and *E. coli* K12, both sourced from ATCC (VA). Strains used for synthesis of eGFP and stat-eGFP were *E. coli* 10β and *E. coli* BL21 (DE3) from New England Biolabs (MA). Bioengineered probiotics were created using *L. lactis* subspecies *cremoris* MG1363 (LMBP 3019) acquired from the Belgian Coordinated Collections of Microorganisms (BCCM).

Synthesis of eGFP and stat-eGFP was done using pE-SUMO (Ampicillin) sourced from LifeSensors (PA). The cloning of AMP, gAMP, and sfGFP into probiotic *L. lactis* was done using pMSP3545 sourced from Addgene (MA).

Synthesis of eGFP and stat-eGFP proteins

The ORFs of eGFP and stat-eGFP were sourced from IDT (CA) as gBlocks, from which they were amplified using PCR and cloned into the pE-SUMO vector using *MfeI* and *BamHI* cut-sites using restriction enzymes and T4 Ligase (Thermo Fisher Scientific). The cloned plasmids were transformed into competent *E. coli* 10β using Heat-shock technique to proliferate the plasmids. Transformed *E. coli* 10β were screened by plating on an Ampicillin (50 μg/ml) agar plate and testing the colonies with colony PCR. Positive colonies were propagated in 15 ml Luria-Betani (LB) broth (Ampicillin, 50 μg/ml) at 37°C overnight, pelleted, lysed, and the extracted plasmid was used to transform *E. coli* BL21 (DE3) for protein synthesis. Positively transformed *E. coli* BL21 (DE3) colonies were grown in 2X Yeast-Tryptone (YT) broth in a larger volume (~500 ml) at 37°C overnight and then induced with 0.5 mM Isopropyl β- d-1-thiogalactopyranoside (IPTG) with a further overnight culture at 16°C. The resultant culture was pelleted at 7000 g and lysed with 0.5 mg/ml Lysozyme along with a freeze-thaw cycle. The thawed lysed pellet is sonicated and centrifuged at 50,000g to obtain the supernatant containing the synthesized proteins, which was purified using Ni-column ion affinity chromatography, utilizing the 6-His tag found in the SUMO fusion partner of the pE-SUMO vector. The purified protein was further cleaved with SUMO Protease, which cleaves the SUMO-fusion partner and releases the synthesized protein.

Binding assay with Flow Cytometry

Flow cytometry was used to measure cellular binding of the stat-eGFP conjugate protein against the targeted *F. nucleatum* (CRC strain) as well as against the off-target bacteria, *B. fragilis* (ETBF). F. *nucleatum* and *B. fragilis* were both grown anaerobically (5% CO_2, 5% H_2, 90% N_2) in Columbia Broth and Brain-Heart Infusion Broth (BHI) respectively, for 48 hrs or till turbid, at 37°C. The bacteria were then standardized to an OD600 of 1.0 and diluted 1:20 in 1X PBS supplemented with 125 μg/mL unguided eGFP or stat-GFP and incubated for 30 minutes at 37°C, shaking at 180 rpm. Cells were washed and resuspended in 1× PBS and flowed through a BD FACSVerse system (BD Biosciences, NJ). Cells were excited with a blue 488 nm laser using a 488/10 bandpass filter. For each sample, fluorescence intensity measures were recorded for a total of ≥4,000 events collected in triplicate.

189 Data were analyzed using FCS Express (De Novo Software, CA). Statistical analyses were done for
190 both the median fluorescence and percent of events positive for eGFP by using ANOVA.

Biofilm Assay with synthetic AMP/gAMPs

Synthetic peptides (HPLC purified) for CPSC5, Ovispirin and their Stat-guided analogs were
purchased from ABI Scientific (VA). After reconstitution in sterile PBS buffer, the peptides were added
to a 96 well plate in a concentration gradient from 0.5 to 32 µM in appropriate media for the bacteria
of choice (Columbia for *F. nucleatum*, BHI for *B. fragilis*, LB for *E. coli*) up to a volume of 0.2 ml in each
well. The bacteria, grown in appropriate media either anaerobically or aerobically, was standardized
to OD600 of 1.0 and 5 µl of it were added to each well, followed by incubation for appropriate amount
of time (overnight aerobically for *E. coli*, 72 hr anaerobically for *F. nucleatum*, 48 hr anaerobically for
B. fragilis). The plate post-incubation was assayed for biofilm formation using a modifed protocol
described by O' Toole et al [45]. Briefly, the media was aspirated from the wells by pipetting, followed
by fixing the biofilm with 0.2 ml of Methanol added to each well, evaporation of the methanol by
aeration/incubation at 50°C, addition of equal volume of 1% Crystal Violet solution, 3X washing by
water or till the wash runs clear, dissolution of the crystal violet bound to the biofilm in 0.2 ml of 30%
Acetic acid solution, transfer of the crystal violet-acetic acid solution to a fresh 96 well plate for
colorimetric assay at 600 nm. One row of the plate was left only to grow the bacteria without any
AMP/gAMP added, the absorbance reading of that row is taken as the positive control, and biofilm
inhibition % is calculated for every other well from that control.

Growth kinetics assay with synthetic AMP/gAMPs

209 For growth kinetics assay, the AMP/gAMPs and the bacteria were set up in a 96 well plate in the
210 identical manner as the biofilm assay. The incubation in this case was done inside the Stratus Kinetic
211 Microplate Reader (Cerillo, VA) at 600 nm to read the cell growth kinetics. The data post-incubation
212 period was procured and analyzed using the Cerillo Labrador Software, with urther statistical
213 representation done in R.

Cloning antimicrobial peptides and guided antimicrobial peptides in *L. lactis*

The ORFs of the AMPs (**Table S1**), codon-optimized for *L. lactis*, were cloned into the pMSP3545
plasmid (**Fig 4a**) which plasmid includes the PNisA Nisin-inducible promoter and the usp45 signal
peptide for extracellular secretion of the downstream peptide. The ORFs were amplified from a
gBlock by PCR (1 minute melting at 95°C; 35 cycles of 15 seconds melting at 95°C, 15 seconds
annealing at Tm + 3°C, 30 seconds extension at 72°C; and 5 minutes elongation at 72°C) using
respective primers and pasted into the pMSP3545 plasmid using restriction enzyme cut sites post
agarose gel purification. The recombinant plasmid with AMPs/gAMPs were electroporated into
electrocompetent *L. lactis* MG1363 (LMBP 3019) cells. Briefly, the procedure involved mixing of
thawed electrocompetent *L.lactis* with at least 0.25 µg of plasmid, resuspended in ice-cold 0.5 M
sucrose and 10% glycerol, electroporation in a cuvette (2000 V, 25 µF, 200 Ω) for a pulse of 5 ms, and
grown out at 30°C in GM17 media (M17 media with 0.5% Glucose). The screening for positive
transformation was done by plating on GM17 agar plate with erythromycin (5 µg/ml).

Co-culture assay of probiotic with bacteria monoculture

L. lactis clones engineered to express AMP or gAMP were propagated from glycerol stocks and grown
in GM17 broth overnight at 30°C with erythromycin (5 µg/ml) with no shaking. *F. nucleatum* was grown
in Columbia broth anaerobically at 37°C for 48-72 hrs. The *L. lactis* cultures were serially diluted in a

96-well culture plate with GM17 broth to make up a volume of 100 µL. To each well, 10 µL of the *F.*
nucleatum culture were added, and each well volume was brought up to 200 µL with Columbia broth.
The plate was left to grow overnight in anaerobic conditions. After 24 hours, the well contents were
transferred to a 96-well PCR plate. That PCR plate was sealed, heated for 15 minutes at 100°C, and
then chilled at 4°C for 5 minutes. This plate was then centrifuged at 2,000 *g* for 2 minutes, and the
supernatant was used as the template for qPCR.

qPCR was carried out using primers for the FomA gene (forward: 5'-ACTTTACCAGTTGCCCAGTT-3';
reverse: 5'-GGAGACCAAATGGTTCAGTAGAT-3')to quantify *F. nucleatum* titer and primers flanking
the *L. lactis acma* gene (forward: 5'-GGAGCTCGTGAAAGCTGACT-3'; reverse: 5'-
GCCGGAACATTGACAACCAC-3') were used to quantify *L. lactis* titer. The qPCR used SYBR Green as
the amplification dye and ROX as the passive dye, and the thermal cycler had 2 minutes melting at
95°C followed by 40 cycles of 15 seconds melting at 95°C, and 1 minute of annealing/extension at
60°C, ending with a melt curve. Standard curves for *F. nucleatum* and *L. lactis* were constructed by
determining C_T values from the qPCR data for different dilutions of the gDNA extracted from the
cultures of the respective bacteria (1/10, 1/100, $1/10^3$, and $1/10^4$) in the qPCR plates, and the
approximate number of genomes represented in the gDNA was calculated by dividing the amount of
gDNA by the average size of the genome of that species. The same procedure was followed with the
off-target bacteria where *B. fragilis* and *E. coli* were co-cultured with serially diluted cultures of *L.*
lactis for 24 hours, and the titers of the off-target bacteria were determined by qPCR using primers
for species-specific genes for either bacterium, DE3-T7 polymerase for *E. coli* (forward: 5'-
GAAGCTTGCTTCTTTGCT-3'; reverse: 5'-GAGCCCGGGGATTTCACAR-3')and *Bft*/Bf toxin for *B. fragilis*
(forward: 5'-GGTTTCAACCGTCAGGTACA-3'; reverse: 5'-GCGAACTCATCTCCCAGTATAAA-3'). The
amount of *L. lactis* added to the co-cultures of all the three assays ranged from approximately
2.8×10^6 to 1.8×10^8 CFU/ml. The statistical difference between any data sets was performed by
ANOVA.

Co-culture assay of probiotics with polymicrobial community culture

Three human fecal samples were randomly selected from the control arm of our vitamin D
microbiome randomized control trial (The study was approved by the Institutional Review Board (IRB)
(1845028-2) at Baylor University and registered under the identifier NCT05387876 at
ClinicalTrials.gov.) The polymicrobial community was created by reanimating fecal samples from
human subjects kept frozen at -80°C in the complex media described by Li *et. al.* for *ex vivo*
microbiome culture in their MiPro model [21], in anaerobic condition at 37°C for 48 hrs. These
polymicrobial cultures were added in 96 well (2 ml, deep well) with *F. nucleatum* and probiotic of
choice in the ratio 6:1:1. For samples that were *F. nucleatum* only negative control, the rest of the
volume was made up by adding more of the complex media. The plates were incubated anaerobically
at 37°C with separate plates that served as 0 hr, 24 hr and 48 hr timepoint samples. Post incubation,
the samples were extracted using Quick DNA Fecal/Soil Microbe 96 Kit (Zymo Research Corp, CA).
The extracted gDNA was either subjected to 16s rRNA Illuminq sequencing or *F. nucleatum* specific
sequencing as described in the previous protocol.

Illumina 16S rRNA sequencing of polymicrobial samples

The extracted gDNA was sequenced as per the protocol described in the Earth Microbiome Project
[46]. The 16S rRNA variable region V4 was amplified with polymerase chain reaction (PCR), using the
following primers, 16S Forward Primer (515 F) with adapters: TCG TCG GCA GCG TCA GAT GTG TAT

AAG AGA CAG GTGYCAGCMGCCGCGGTAA 3', and 16S Reverse Primer (926 R) with adapters: 5' GTC
TCG TGG GCT CGG AGA TGT GTA TAA GAG ACAG CCGYCAATTYMTTTRAGTTT 3'.The PCR mixture
comprised 1 µl of each forward and reverse primer (12.5 µM), 5 µl of extracted DNA of approximately
equal concentration from each sample, 12.5 µl of 2x Platinum Hot Smart Master Mix (Thermo Fisher
Scientific), and water to make a final volume of 25 µl. The amplifications were performed under the
following conditions: initial denaturation at 94 °C for 5 min, followed by 35 cycles of denaturation at
94 °C for 45 s, primer annealing at 50 °C for 1 min, and extension at 72 °C for 1 min 30 s, with a final
elongation at 72 °C for 5 min. The presence of PCR products was visualized by electrophoresis using
a 1.5% agarose gel. The products were cleaned using Sera-Mag select PCR clean up kit (Cytiva). A
second PCR was conducted to attach the Illumina adapters and barcode index primers-I5 (forward)
and I7 (reverse). After adapter and index attachment, the amplicons were normalized and pooled
together in a DNA library at a concentration of 4 nM, measured by Quant-iT dsDNA Assay Kit (Thermo
Scientific). The pooled DNA library was paired end sequenced at 2x300 bp using a MiSeq Reagent Kit
v3 on Illumina MiSeq platform (Illumina, CA), at Baylor University.

16S rRNA sequencing data analysis

The sequences were evaluated, demultiplexed, and filtered using QIIME2 (version 2024.5) [47]
plugins on the Kodiak High-Performance Computing (HPC) Cluster hosted by High Performance and
Research Computing Services, Baylor University. Both paired reads were trimmed from the forward
end and read length of at least 200 bp for further processing to generate the ASVs. The denoising and
filtering of chimeric sequences was done using the DADA2 plug in of QIIME 2. The samples were
rarefied at a depth of >2500 which retained 97% of the samples. Taxonomic classification was
performed utilizing the QIIME2-compatible pre-trained feature classifier based on 16S rRNA gene
database from Silva 16S rRNA 138 database. The alpha diversity (Shannon, Faith Phylogenetic
Diversity, and Species Richness) and beta diversity (Bray-Curtis, Aitchison's, Weighted and
Unweighted Unifrac) analyses were performed using QIIME2 plugins. The resultant α-diversity, β-
diversity, and ASV tables were downloaded and further analyzed in R 4.3.1 and RStudio platform
2023.12.1. The resultant taxonomic abundance data (at genus level) was analyzed using the CCREPE
package in R [48] to determine the correlation between the taxa according to relative changes in
abundance upon addition of *F. nucleatum*. The resultant coterie of genera (|correlation|>0.25, P-
value < 0.05, FDR adjusted P-value <0.15, iterations = 1,000) was then used to construct the equation
for Dysbiosis Index Equation. Every other data visualization used default tools in the *tidyverse* and/or
ggplot2 packages.

Results

The Statherin derived guide peptide preferentially attaches to *F. nucleatum*

The goal of our initial experiments was to determine if our guide peptide had specificity towards
FomA on *F. nucleatum*. The binding affinity of the statherin-derived guide peptide ligand was
demonstrated by synthesizing eGFP fused with the peptide ligand (stat-eGFP) and quantifying the
binding with flow cytometry post-incubation with *F. nucleatum* and *B. fragilis* (non-*Fusobacterium*
candidate). Both the median fluorescence and the percent of fluorescent positive events were
significantly higher (p<0.01, ANOVA) for *F. nucleatum* incubated with stat-eGFP than with either
native eGFP or untreated bacteria (**Fig. 2A-2B**). The fluorescence values from the same bacteria

treated with native eGFP was higher than the untreated sample but not significantly. In contrast, *B.*
fragilis treated with either native or stat-guided eGFP exhibited no significant difference between
themselves (p=0.58, ANOVA) although they were more significantly fluorescence than the untreated
samples (p<0.05, ANOVA). Thus, we show here that eGFP with the stat-guide ligand peptide
preferentially attaches to the *F. nucleatum* cells compared to native eGFP, unlike in non-target
bacterium *B. fragilis*. Our data indicate that the ligand derived from the statherin (stat) peptide is an
effective guide for fluorescent molecules to attach to the target pathogen *F. nucleatum* preferentially
over unguided molecules, which was hindered in commensal anaerobes.

Statherin-derived ligand guided AMP (gAMP) selectively inhibits biofilm formation of *F.*
nucleatum

Next, using synthetic peptides (Ovispirin and CPSC5) and their guided analogs (stat-CPSC5 and stat-
Ovispirin) we measured *in vitro* their biofilm inhibition capabilities. The synthetic peptides and their
guided analogs were tested using 2 strains of *F. nucleatum*, one derived from CRC tumor samples (*F.*
nucleatum SB-CTX3Tcol3) and another derived from an oral lesion (*F. nucleatum* subsp.
nucleatum Knorr - ATCC 25586), along with well-described commensal strains of *B. fragilis* ATCC
43858 (anaerobic) and *E. coli* K12 ATCC 10798 (aerobic), as non-target bacteria candidates. Post
incubation, using crystal violet assays for detection of biofilm formation, we show that both the AMPs
and their corresponding gAMPs have similar patterns of inhibition against both the strains of
Fusobacterium (**Fig 2C**), with their IC_{50} as low as 0.7 µM to as high as 3 µM (**Table 1**). While there was
no change in MIC or IC_{50} observed in the gAMP compared to their AMPs against *F. nucleatum* (0.71-
2.15 µM for Ovispirin, 0.91-2.3 µM for Stat-Ovispirin, *p=0.83,* 1.42-5.6 µM for CPSC5, 1.3-5.7 µM for
Stat-CPSC5, *p=0.76*, ANOVA), when using the gAMP against the non-target bacteria, *B. fragilis* (23 µM
for Ovispirin, >100 µM for Stat-Ovispirin, 18 µM for CPSC5, >100 µM for Stat-CPSC5, *p<0.01*, ANOVA)
and *E. coli* (4.45 µM for Ovispirin, >30 µM for Stat-Ovispirin, 1.3 µM for CPSC5, >20 µM for Stat-
CPSC5, *p<0.01*, ANOVA), we see a significant decrease in the inhibitory effect of the gAMPs
compared to their AMPs, with the MIC and IC_{50} increasing significantly for the gAMPs against both the
bacteria. This targeting effect was more pronounced against *B. fragilis,* with the gAMPs having greatly
diminished inhibitory effect even beyond 32 µM and 64 µM of gAMP. For *E. coli*, the increase in IC_{50}
was significant but not as pronounced as for *B. fragilis*, with a 4-fold to 7-fold increase in IC_{50}. Thus,
it can be concluded that the stat-CPSC5 and stat-Ovispirin were significantly more toxic towards *F.*
nucleatum strains but not or less toxic compared to CPC5 and Ovispirin alone, or unguided. Against
the non-target bacteria *B. fragilis* and *E. coli*, the gAMPs have significantly less toxicity compared to
their AMPs when it comes to inhibition of biofilm formation. Overall, these data indicate that our
gAMPs against FomA, have selective inhibition towards *F. nucleatum* over non-target bacteria.
However, this differential toxicity towards the different bacteria species between the AMP and gAMP
were not seen in cytotoxicity (MTT) assay against Caco2 and HT29 cells (**Fig S1**)

Fig 2. The Statherin derived guide peptide allows preferential attachment to *F. nucleatum* and exhibits targeted biofilm inhibition. (a) Median fluorescence intensity (MFI) of GFP in *B. fragilis* and *F. nucleatum* following treatment with eGFP, Stat-eGFP, and untreated controls. Statistically significant differences in GFP binding to *F. nucleatum* are indicated (p<0.05). Error bars represent the standard deviation of three independent experiments.
(b) Percent of events positive for GFP, as measured by flow cytometry, comparing Stat-eGFP binding to *F. nucleatum* to *B. fragilis*. Statistical significance between treatments is noted (p<0.05).
(c) Biofilm inhibition of *F. nucleatum* and *B. fragilis* strains in the presence of increasing concentrations of synthetic AMPs (Ovispirin and CPSC5) and their guided versions (Stat-Ovispirin and Stat-CPSC5). The inhibition is plotted as a percentage of biofilm formation. Error bars represent the mean ± standard deviation of three biological replicates. Statistical significance is indicated as $p<0.05$ (*), $p<0.01$ (**), $p<0.001$ (), and *$p<0.0001$ (**). Also see IC_{50} values calculated from the dose-response curve (Table 1)

Bacteria		Ovispirin		CPSC5	
		Unguided	Guided	Unguided	Guided
Fusobacterium	F. nucleatum (Oral)	0.71 ± 1.71 µM	0.91 µM	2.18 ± 3.4 µM	1.3 ± 4.36 µM
	F. nucleatum (CRC)	2.15 µM	1.17 ± 1.04 µM	1.42 ± 0.18 µM	3.02 ± 0.69 µM
Non-Fusobacterium	B. fragilis	23.54 ± 8.63 µM	>100 µM	18.12 ± 6.37 µM	>100 µM
	E. coli	4.45 ± 0.35 µM	17.26 ± 19.23 µM	1.3 ± 0.1 µM	9.68 ± 12.37 µM

Table 1. IC_{50} values of the synthetic AMPs – calculated by fitting biofilm inhibition assay data in a dose-response curve using *'drc'* package (R)

Statherin-derived ligand-guided AMP (gAMP) differentially modulates growth kinetics of *F. nucleatum*

To better understand the effects of the gAMPs on growth of *F. nucleatum,* the growth kinetics of planktonic cultures of two non-*Fusobacterium* species – *E. coli* and *B. fragilis* - and two *Fusobacterium* strains - Oral and CRC - were incubated with both AMPs and their gAMP analogs. This analysis showed a pattern of differential inhibition (**Fig 3A-C**). For both the *Fusobacterium* strains, both the unguided and guided versions of the AMPs reduced the growth; starting between 2 and 8 µM, which continued for over 48 hrs and up to 72 hrs. In contrast, for the non-*Fusobacterium* species, only the unguided AMPs demonstrated any inhibitory effect below 8 µM, while the gAMPs either had no inhibition of growth or exhibited inhibition beyond 16-32 µM range. For *B. fragilis*, the bacteria that had the poorest response from all the AMPs, the guided CPSC5 and Ovispirin did not subdue growth, even at 32 µM, while the unguided version was effective at 8 µM. For *E. coli*, the AMPs started inhibiting growth even at 8 µM but the gAMPs only had inhibitory effects beyond 16 µM, with more pronounced inhibition at 32 µM. This analysis demonstrates that both AMPs and gAMPs inhibit the growth of *F. nucleatum* strains at low concentrations (<4 µM), making them clinically viable. Moreover, gAMPs show minimal toxicity towards non-*Fusobacterium* species compared to unguided AMPs, highlighting their selective toxicity and preferential action against *F. nucleatum* over the commensals.

Fig 3. Selective inhibition of bacterial growth by guided antimicrobial peptides (gAMPs).
Growth kinetics of planktonic cultures for two *Fusobacterium* strains—Oral/ATCC 25586 (a) and
CRC/Strain 14 (b)—and two non-*Fusobacterium* species, *B. fragilis* (c) and *E. coli* (d), were
measured in the presence of both unguided (red) and guided (blue) antimicrobial peptides
(AMPs - CPSC5 and Ovispirin) across a concentration gradient (2 µM to 32 µM). Optical density
(OD600) measurements were recorded over 60 hours (Oral strain) and 40 hours (CRC strain).
Error bars represent standard deviation from three biological replicates.

Cloning *Lactococcus lactis* with gAMP-Expressing Plasmids to Create Bioengineered Probiotics

Once the gAMPs demonstrated sufficient targeting abilities, the next step we took was creating the
engineered probiotic strains expressing gAMPs. The open reading frames (ORFs) of the antimicrobial
peptides (AMPs), CPSC5 and Ovispirin, along with their guided analogs (gAMPs), were cloned into
the pMSP3545 plasmid (see Methods for details). This plasmid includes the nisin-inducible PNisA
promoter and the usp45 signal peptide, which ensures the secretion of the peptides into the
extracellular environment (**Fig. 4a**). The engineered plasmids were introduced into

competent *Lactococcus lactis* MG1363 cells via electroporation. Successful cloning was confirmed through colony PCR using ORF-specific primers, followed by Sanger sequencing to verify the accuracy of the insertion. Based on these results, only the bioengineered probiotics expressing CPSC5 and Stat-CPSC5 were selected for further experiments, as the Ovispirin and stat-Ovispirin constructs did not yield consistent outcomes.

Engineered *L. lactis* Expressing gAMP Exhibits Targeted Inhibition of *F. nucleatum* in Co-Culture Compared to Non-*Fusobacterium* Species.

To further explore the specificity and potential clinical application of the bioengineered *L. lactis* probiotics, we conducted an *in vitro* co-culture experiment to assess the targeted inhibition of *F. nucleatum* compared to non-*Fusobacterium* commensal species. The bioengineered probiotics expressing CPSC5 (AMP) and Stat-CPSC5 (gAMP) demonstrated selective toxicity toward *F. nucleatum* when co-cultured with non-*Fusobacterium* commensal species. Both the AMP- and gAMP-expressing probiotics were co-cultured with a gradient of bacterial titers, ranging from 10^6 to 10^9 CFU/ml, of *F. nucleatum* (CRC strain), *B. fragilis*, and *E. coli*. After 24 hours of incubation, induced with 100 ng/ml of nisin in a 96-well plate format, the co-cultures were pelleted, and genomic DNA was extracted. Species-specific qPCR was used to quantify the target bacteria and estimate the remaining cell counts after incubation with the engineered probiotics. The results showed that the AMP-expressing probiotics inhibited the growth of all three bacterial species in a dose-dependent manner, with the most significant reduction observed in *F. nucleatum* (**Fig. 4b**). In contrast, the gAMP-expressing probiotics selectively inhibited *F. nucleatum* as the probiotic dose increased, while the inhibitory effects on *B. fragilis* and *E. coli* diminished significantly. The toxic effect of the gAMPs against *F. nucleatum* was 10^3 to 10^4 times greater than against *B. fragilis* and *E. coli* at higher probiotic titers (~10^8 CFU/ml), a statistically significant difference ($p < 0.05$, ANOVA). Interestingly, some level of toxicity was observed across all three bacterial species, even in the control probiotic expressing super-folding GFP (sfGFP). This suggests that *L. lactis* may have inherent antimicrobial properties when co-cultured at high densities, possibly due to resource competition or innate toxicity, contributing to the observed inhibition.

The gAMP Engineered Probiotic Maintains Microbial Diversity in a Polymicrobial Community

To determine whether the gAMP engineered probiotic could selectively target *F. nucleatum* while preserving the overall microbial diversity in a gut-like environment, we conducted a series of experiments using an *in vitro* human gut microbiome-derived polymicrobial community model. The efficacy of the bioengineered probiotics (AMP, gAMP, and Control) was tested within a polymicrobial community spiked with *F. nucleatum* (CRC strain) culture at 20% v/v in each well. This polymicrobial community was derived from anaerobic cultivation of rejuvenated fecal samples from three human subjects (FS04, FS16, and FS36), designed to mimic the composition of their gut microbiota. The communities were incubated with equal amounts of *F. nucleatum* and probiotic cultures—either AMP (CPSC5), gAMP (Stat-CPSC5), or Control (sfGFP). Samples were taken at 24 and 48-hour intervals, lysed and subjected to genomic DNA extraction using the phenol-chloroform method. The extracted gDNA was then analyzed using both qPCR with *F. nucleatum*-specific primers and 16S rRNA (V4) sequencing (Illumina) to assess the overall microbial composition. The 16S rRNA sequencing revealed that, while a diverse array of bacteria was present in the polymicrobial

Fig 4. Engineered L. lactis with inducible plasmids demonstrates targeted inhibition of F. nucleatum compared to non-Fusobacterium species. (a) Schematic of the plasmids used to engineer L. lactis. The pMSP3545-derived plasmids contain the nisin-inducible PNisA promoter and the usp45 signal peptide for extracellular secretion of the peptides. The AMP/gAMP plasmids include ORFs for Ovispirin (left) or CPSC5 (right), with or without a Statherin-derived guide peptide. The control probiotic plasmid was constructed using the same backbone, replacing the AMP/gAMP ORFs with the superfolding GFP (sfGFP) ORF. Only CPSC5 and Stat-CPSC5 were successfully cloned into L. lactis. (b) Co-culture experiments with *F. nucleatum* (CRC strain), *Bacteroides fragilis*, and *E. coli* in the presence of control (sfGFP), AMP-expressing (CPSC5), or gAMP-expressing (Stat-CPSC5) *Lactococcus lactis* probiotics. The relative number of bacterial genomes remaining in the co-culture was quantified by species-specific qPCR after 24 hours of probiotic treatment, and plotted against the CFU of probiotic added. Error bars represent standard deviation from three biological replicates, and statistically significant differences (ANOVA, $p<0.05$) between guided and unguided AMPs are indicated by asterisks.

communities from each subject, the targeted modulation of *F. nucleatum* was most prominent in the

samples treated with the engineered probiotics. This modulation was in stark contrast to the
untreated controls, where *F. nucleatum* abundance increased considerably from around 10% to over
25%   in   48   hrs   (**Fig   5a   &   b**).   Other   prevalent   genera,
including *Enterococcus, Peptoniphilus, Eubacterium, Escherichia, Bacteroides, Sellimonas, Negati*
vicoccus, and *Paraclostridium*, showed little to no change in relative abundance compared
to *Fusobacterium*, indicating that the gAMP specifically modulates *F. nucleatum* without
significantly affecting other members of the microbiota.

Measuring the alpha diversity of the samples using three different indices revealed that all probiotic
treatments preserved microbial community diversity after exposure to *F. nucleatum*. Among the
treatments, the gAMP probiotic exhibited the smallest shift in microbial diversity from 0 to 48 hours.
The Shannon index (**Fig. 5C**) indicates a decrease in diversity at 24 hours in all probiotic-treated
samples, likely due to the proliferation of *F. nucleatum*. However, diversity began to recover in the
following 24 hours. This decline at 24 hours was statistically significant in the control and AMP-
treated samples but not in the gAMP-treated samples, with no significant differences in the recovery
phase across all treatments. These results suggest that the gAMP probiotics caused the least
disruption to microbial diversity, showing minimal and non-significant changes. In the samples that
were spiked with *F. nucleatum* without probiotic treatment, diversity significantly declined at both 24

Fig 5. The gAMP probiotic maintained the diversity of the microbiota in the in vitro polmicrobial community. Heatmaps of the genus-level a) and family-level b) relative abundance. Polymicrobial community rejuvenated from human fecal microbiome sample spiked with *F. nucleatum* culture and then treated with bioengineered probiotics secreting CPSC5 (AMP), Stat-CPSC5 (gAMP) and sfGFP (Control) at a ratio of 6:1:1 respectively. b-d) Three alpha diversity indices – Shannon (a), Faith's Phylogenetic diversity (b) and species richness (c), showing the ability of gAMP probiotic treatment to maintain the biodiversity of the polymicrobial community post spiking with *F. nucleatum*, compared to AMP and control probiotics. No significant changes either between any consecutive timepoints or the overall timeframe. ANOVA was used for testing for significance between treatments with p<0.05

and 48 hours (p=0.0019 & 0.004 for Shannon Index, p=0.0078 & 0.0005 for Faith PD, p=0.031 & 0.004
for species richness, ANOVA) . Interestingly, this dip and recovery pattern was not observed when
analyzing the Faith Phylogenetic Diversity and Species Richness (**Fig. 5D-E**). Both of these indices
showed a steady decline in diversity across all three probiotic treatments and in the *F. nucleatum-*
only controls, with the gAMP-treated samples exhibiting the least drastic and non-significant
changes in microbial diversity.

The bioengineered probiotic effectively targets *F. nucleatum* in a gut microbiome-derived
454
polymicrobial community without disrupting the overall microbiota

To assess whether the bioengineered gAMP probiotic could effectively reduce *F. nucleatum* in a
complex gut microbiome-derived polymicrobial community without disrupting the overall microbial
balance, we conducted a series of *in vitro* experiments. Using both 16S rRNA sequencing and qPCR,
we found that the gAMP probiotic was highly effective in reducing *F. nucleatum* even within a complex
polymicrobial community. Although all three probiotic variants demonstrated some antimicrobial
activity, the gAMP probiotic produced the most significant reduction in *F. nucleatum* titers after 48
hours of incubation. Specifically, the *F. nucleatum* abundance, measured by Centered-Log Ratio
(CLR) analysis, was lowest in the gAMP-treated samples compared to the AMP-treated ones, with a
statistically significant reduction observed for the gAMP probiotic (**Fig. 6A**) (p=0.031, ANOVA). In the
probiotic-treated samples, there was a slight increase in *F. nucleatum* CLR at the 24-hour mark,
followed by a notable decline, which was significant only in the gAMP-treated samples. In contrast,
samples treated only with *F. nucleatum* exhibited a steady increase in CLR over time. Interestingly,
these significant reductions in *F. nucleatum* abundance were not fully reflected in the qPCR results,
where similar reductions were observed across all probiotic treatments. Only the untreated control
samples showed a significant increase in *F. nucleatum* titer at both time points (p=4.1e-05, ANOVA)
(**Fig. 6A**). This discrepancy between the two methods may be attributed to the nature of the
measurements: qPCR focuses solely on the number of gene copies for the target species, whereas
CLR analysis accounts for the relative abundance of *F. nucleatum* in relation to all other bacteria in
the community, offering a more comprehensive view of its presence within a polymicrobial
environment.

A Compositionality Corrected by REnormalization and PErmutation (CCREPE) analysis was
performed to identify bacterial species either positively or negatively correlated with *F.*
nucleatum abundance (|correlation| > 0.25, FDR-adjusted p < 0.15) across the samples. This allowed
us to determine which bacteria increased as *F. nucleatum* levels rose, and which decreased in

response. In the gAMP probiotic-treated samples, bacteria positively correlated with *F. nucleatum*
showed a consistent reduction, while those negatively correlated were either maintained or
increased in abundance at both time points (**Fig. 6C**). In contrast, the samples treated only with *F.*
nucleatum exhibited the opposite trend. This information was used to develop the Fusobacterium
Dysbiosis Index (**Formula 1**), which quantifies the extent of dysbiosis caused by introducing *F.*
nucleatum into the community. Using this index, the gAMP probiotics showed the most effective
performance in vitro (**Fig. 6D**). The gAMP-treated samples exhibited a significant reduction in
dysbiosis, indicating that not only did the gAMP probiotics reduce *F. nucleatum* abundance, but they
also preserved microbial diversity more effectively than AMP probiotics by selectively targeting the
pathogen.

$$Fusobacterium\ Dysbiosis\ Index = \log_{10}\left(\frac{\sum \begin{array}{c} Abundance\ of\ genera\ positively \\ correlated\ to\ Fusobacterium \end{array}}{\sum \begin{array}{c} Abundance\ of\ genera\ negatively \\ correlated\ to\ Fusobacterium \end{array}}\right)$$

Formula 1.
Fusobacterium
Dysbiosis Index

490   Although all three probiotic treatments showed a similar pattern of a slight increase in dysbiosis at
491   24 hours followed by a reduction at 48 hours, the decrease in the gAMP-treated samples was
492   significantly lower than at both the 0-hour and 24-hour time points—an effect not seen with the AMP
493   or control probiotics. These results confirm that the gAMP probiotic expressing Stat-CPSC5

a)

b)

c)

d)

Fig 6. The bioengineered probiotic effectively targets *F. nucleatum* in a gut microbiome-derived polymicrobial community without disturbing the overall microbiota. *(a)* The effect control probiotics, AMP, or gAMP probiotic on reduction in *F. nucleatum* titers after 48 hours in an in vitro polymicrobial community. The abundance of *F. nucleatum* was measured by Centered-Log Ratio (CLR), (ANOVA).b) *F. nucleatum* titer measure by qPCR of the sample samples. The y-axis gives the approximate number of *F. nucleatum* left in the sample at that timepoint after treatment with the given probiotic. (ANOVA, $p<0.05$) *(c)* Compositionality Corrected by REnormalization and PErmutation (CCREPE) analysis of bacterial genera that were positively (red) or negatively (blue) correlated with *F. nucleatum* abundance. This analysis was used to create the Fusobacterium Dysbiosis Index. *(d)* The effect of probiotic, AMP, or gAMP probiotics on community structure, as indicated by the Fusobacterium Dysbiosis Index, which was calculated by comparing the abundance of *F. nucleatum* and its positively correlated genera against the negatively correlated genera.(Dysbiosis Index = |correlation| > 0.25, FDR-adjusted $p < 0.15$)

demonstrates preferential antimicrobial activity against *F. nucleatum*, consistent with findings from
both monoculture and polymicrobial community experiments.

Discussion

Our research presents critical insights into the specificity and efficacy of Statherin-derived guide
peptides as a first step in creating an engineered probiotic selectively targeting *F. nucleatum*
highlighting the potential for targeted antimicrobial therapies. Our study investigates the implications
of these findings, particularly in relation to the selective bacterial targeting, inhibition of biofilm
formation, modulation of bacterial growth, and the preservation of microbial diversity within complex
polymicrobial communities.

By utilizing a statherin-derived guide peptide fused to an enhanced green fluorescent protein (eGFP),
we observed preferential attachment to *F. nucleatum* over *B. fragilis*, a non-Fusobacterium
commensal anaerobe [49]. Flow cytometry revealed significantly higher fluorescence in *F.
nucleatum* treated with stat-eGFP compared to both native eGFP and untreated controls, indicating
a specific interaction [50]. In contrast, *B. fragilis* exhibited no significant difference between native
eGFP and stat-eGFP treatments, confirming the specificity of the statherin-derived peptide in guiding
attached molecules to bind selectively to *F. nucleatum*. This specificity is mediated by the interaction
between the Statherin-derived peptide and the membrane protein FomA, as shown by previous
studies in which *F. nucleatum* mutants lacking FomA lost binding specificity to the guide peptide
ligand [39]. This specificity is crucial in designing targeted therapies that minimize the impact on non-
target microbiota, thereby reducing potential off-target effects and preserving the delicate balance
within the gut microbiome [51, 52].

Biofilms provide a protective environment for bacteria, increasing their resistance to antimicrobial
agents and immune responses [53]. Our biofilm inhibition assays further demonstrated that
statherin-derived ligand-guided antimicrobial peptides (gAMPs), including stat-CPSC5 and stat-
Ovispirin, retained strong inhibitory effects against *F. nucleatum* biofilms while showing reduced
efficacy against non-target bacteria such as *B. fragilis* and *E. coli*. This selective inhibition is
particularly important in the context of treating bacterial infections linked to biofilm formation. The
ability of gAMPs to selectively inhibit *F. nucleatum* biofilms without affecting commensal bacteria
suggests their potential as therapeutic agents for biofilm-associated infections, particularly in the
oral cavity and gastrointestinal tract [54, 55].

Growth kinetics data also supported the specificity of gAMPs, showing that both guided and
unguided AMPs inhibited *F. nucleatum* at low clinically relevant concentrations. However, gAMPs
exhibited significantly lower toxicity toward non-*Fusobacterium* bacteria. For example, in *B. fragilis*,
gAMPs had a negligible inhibitory effect even at higher concentrations, whereas unguided AMPs were
effective at much lower doses. This selective growth inhibition emphasizes the therapeutic potential
of gAMPs, where it is essential to target pathogenic bacteria without disrupting the broader microbial
community [51, 52]. Such precision could prevent the adverse effects commonly associated with
broad-spectrum antibiotics, such as dysbiosis and the emergence of antibiotic-resistant strains.
These results also confirm that the gAMP strategy is effective against both planktonic and biofilm
forms of the target pathogen.

The creation of bioengineered probiotics expressing gAMPs introduces an innovative approach to
delivering targeted antimicrobial therapy within the gut. There has been recent research done using
the approach of targeted antimicrobial peptide against *F. nucleatum* using the FomA protein [56], but
it did not offer a suitable delivery platform for delivery of the peptides in situ of infection. Engineered
Lactococcus lactis developed by us, expressing stat-CPSC5, exhibited selective toxicity towards *F.*
nucleatum when co-cultured with non-target bacteria. The significant reduction in *F. nucleatum*
populations, as indicated by quantitative PCR and species-specific genome quantification, was
observed without comparable effects on *B. fragilis* and *E. coli*. This differential targeting is
advantageous for maintaining a healthy gut microbiota while combating pathogens. Additionally, the
inherent antimicrobial activity of *L. lactis* suggests it could serve as an effective vehicle for delivering
therapeutic agents, enhancing its utility in microbial therapy.

One of the most significant findings of this study is the ability of gAMP-expressing probiotics to
preserve the diversity of a polymicrobial community while effectively reducing *F. nucleatum* levels.
16S rRNA sequencing showed that gAMP probiotics maintained microbial diversity to a greater extent
than AMP or control probiotics. Alpha diversity (Shannon Index) demonstrated that while all samples
showed a decrease in diversity at 24 hours, likely due to *F. nucleatum* proliferation, the gAMP
probiotic-treated samples exhibited a less significant decrease and a more complete recovery by 48
hours. This suggests that gAMPs can selectively reduce pathogenic bacteria while minimizing
collateral damage to the overall microbial community.

Moreover, compositional analysis using CCREPE [48] revealed that gAMP probiotics were most
effective in reducing the dysbiosis index, a measure of microbial imbalance caused by *F. nucleatum*
overgrowth [57]. This reduction in dysbiosis correlated with the selective decrease in bacteria
positively associated with *F. nucleatum* and the preservation or increase of bacteria negatively
associated with *F. nucleatum*. These findings are significant because maintaining microbial diversity
and function is essential for gut health, and significant and/or prolonged disruptions can lead to
conditions such as inflammatory bowel disease, cancer and other metabolic disorders [58].

Our findings have broad implications for developing targeted antimicrobial therapies. The specificity
of Statherin-derived guide peptides for *F. nucleatum* opens up the possibility of using gAMPs in
clinical settings to treat infections associated with this pathogen, particularly in the context of
colorectal cancer, periodontal disease, and even reproductive diseases such as pre-term birth and
endometriosis [14, 59–62]. Additionally, the use of bioengineered probiotics as a delivery system
offers a promising avenue for administering these therapies in a controlled and sustained manner.
Future research should focus on translating this therapeutic approach to in vivo models, where

bioengineered *L. lactis* synthesizing and secreting gAMP could be administered orally for gut
colonization, allowing the peptides to escape gastric proteolysis and exert antimicrobial effects in
situ by preferentially binding to *F. nucleatum*'s FomA surface protein via the Statherin-derived guide
peptide. (**Fig 7**).

Fig 7. Model for in vivo application of bioengineered probiotics as a therapeutic. The probiotic schematic representation illustrates the engineered probiotic mechanism. The probiotic Lactococcus lactis is bioengineered with a Nisin-inducible plasmid that includes the PNisA promoter and the usp45 secretion signal peptide. This vector enables the expression and secretion of antimicrobial peptides (AMPs) or guided antimicrobial peptides (gAMPs) in the gut upon oral administration. The statherin-derived guide peptide on the gAMPs targets the FomA membrane protein expressed by F. nucleatum. Once secreted, the gAMPs preferentially bind to FomA on F. nucleatum, allowing targeted antimicrobial activity and thereby reducing F. nucleatum colonization in the infected colon without affecting non-target bacteria.

Furthermore, the ability to maintain microbial diversity while selectively targeting pathogens could revolutionize the treatment of bacterial infections. As indicated, the current antibiotic therapies often disrupt the gut microbiota, leading to adverse effects and promoting the development of antibiotic resistance. The use of gAMPs, particularly when delivered via probiotics, could mitigate these issues by providing a more targeted approach that preserves the beneficial components of the microbiota in an easily delivered oral therapeutic.

In conclusion, this study demonstrates the effectiveness of Statherin-derived guide peptides in selectively targeting *F. nucleatum*, inhibiting its biofilm formation, and modulating its growth kinetics without significantly affecting non-target commensal bacteria. The bioengineered probiotics expressing gAMPs further showed promise in reducing *F. nucleatum* within a complex polymicrobial community while maintaining microbial diversity. These findings lay a strong foundation for the development of targeted antimicrobial therapies that could offer a more precise and less disruptive alternative to traditional antibiotics.

Data Availability

The 16s rRNA amplicon sequences of the polymicrobial community derived from the fecal samples used n the study are available in NCBI SRA database under the BioProject ID PRJNA1179861.

Acknowledgements

We acknowledge the help of members of the laboratory of Dr. Christopher M. Kearney, Department of Biology, Baylor University for assistance with the development of the guided antimicrobial peptide and bioengineered probiotic technology including Dr. Mikaeel Young, Dr. Patrick Ortiz, and Dr. Toslim Mahmud for help in procuring bacteria strains and performing assays. We acknowledge the help of Dr. Emma Allen-Vercoe, Department of Molecular and Cellular Biology, University of Guelph, ON, Canada and Dr. Susan Bullman, Department of Immunology, MD Anderson Cancer Center, Texas, USA for sharing their clinically sourced strains of *Fusobacterium nucleatum* with us.

Funding

The research was funded by the Office of Vice Provost and Research Post-Doctoral Research Fellowship Grant, Robbins College of Health and Human Sciences Post-Doctoral Fellow Research Support Grant, ONE-URC Grant for Undergraduate Research, all sourced through Baylor University.

References

1. Cancer of the Colon and Rectum - Cancer Stat Facts. SEER. https://seer.cancer.gov/statfacts/html/colorect.html. Accessed 21 Mar 2023.

2. Dharwadkar P, Zaki TA, Murphy CC. Colorectal cancer in younger adults. Hematol Oncol Clin North Am. 2022;36:449–70.

3. American Cancer Society. Colorectal Cancer Facts & Figures 2023-2025. Atlanta: American Cancer Society; 2023.

4. Panagi M, Georgila K, Eliopoulos AG, Apidianakis Y. Constructing personalized longitudinal holo'omes of colon cancer-prone humans and their modeling in flies and mice. Oncotarget. 2019;10:4224.

5. Francescone R, Hou V, Grivennikov SI. Microbiome, Inflammation and Cancer. Cancer J. 2014;20:181–9.

6. Chew J, Zilm PS, Fuss JM, Gully NJ. A proteomic investigation of Fusobacterium nucleatum alkaline-induced biofilms. BMC Microbiol. 2012;12:189.

7. Coppenhagen-Glazer S, Sol A, Abed J, Naor R, Zhang X, Han YW, et al. Fap2 of Fusobacterium nucleatum is a galactose-inhibitable adhesin involved in coaggregation, cell adhesion, and preterm birth. Infect Immun. 2015;83:1104–13.

8. Gholizadeh P, Eslami H, Kafil HS. Carcinogenesis mechanisms of Fusobacterium nucleatum. Biomed Pharmacother. 2017;89:918–25.

9. Groeger S, Zhou Y, Ruf S, Meyle J. Pathogenic Mechanisms of Fusobacterium nucleatum on Oral Epithelial Cells. Front Oral Health. 2022;3:831607.

10. Han YW. Fusobacterium nucleatum: a commensal-turned pathogen. Curr Opin Microbiol. 2015;0:141–7.

11. Han YW, Shi W, Huang GT, Kinder Haake S, Park NH, Kuramitsu H, et al. Interactions between periodontal bacteria and human oral epithelial cells: Fusobacterium nucleatum adheres to and invades epithelial cells. Infect Immun. 2000;68:3140–6.

12. Kaplan CW, Ma X, Paranjpe A, Jewett A, Lux R, Kinder-Haake S, et al. Fusobacterium nucleatum outer membrane proteins Fap2 and RadD induce cell death in human lymphocytes. Infect Immun. 2010;78:4773–8.

13. Abed J, Maalouf N, Manson AL, Earl AM, Parhi L, Emgård JEM, et al. Colon Cancer-Associated Fusobacterium nucleatum May Originate From the Oral Cavity and Reach Colon Tumors via the Circulatory System. Front Cell Infect Microbiol. 2020;10:400.

14. Sun C-H, Li B-B, Wang B, Zhao J, Zhang X-Y, Li T-T, et al. The role of Fusobacterium nucleatum in colorectal cancer: from carcinogenesis to clinical management. Chronic Dis Transl Med. 2019;5:178–87.

15. Tran HNH, Thu TNH, Nguyen PH, Vo CN, Doan KV, Nguyen Ngoc Minh C, et al. Tumour
microbiomes and Fusobacterium genomics in Vietnamese colorectal cancer patients. npj Biofilms
Microbiomes. 2022;8:1–13.

16. Lee JB, Kim K-A, Cho HY, Kim D, Kim WK, Yong D, et al. Association between Fusobacterium
nucleatum and patient prognosis in metastatic colon cancer. Sci Rep. 2021;11:20263.

17. Zepeda-Rivera M, Minot SS, Bouzek H, Wu H, Blanco-Míguez A, Manghi P, et al. A distinct
Fusobacterium nucleatum clade dominates the colorectal cancer niche. Nature. 2024;628:424–32.

18. Galeano Niño JL, Wu H, LaCourse KD, Kempchinsky AG, Baryiames A, Barber B, et al. Effect of
the intratumoral microbiota on spatial and cellular heterogeneity in cancer. Nature. 2022;611:810–
7.

19. CDC. Impacts of Antimicrobial Resistance on Cancer Care. Antimicrobial Resistance. 2024.
https://www.cdc.gov/antimicrobial-resistance/stories/impacts-of-antimicrobial-resistance-on-
cancer-care.html. Accessed 15 Oct 2024.

20. Nanayakkara AK, Boucher HW, Fowler Jr VG, Jezek A, Outterson K, Greenberg DE. Antibiotic
resistance in the patient with cancer: Escalating challenges and paths forward. CA: A Cancer
Journal for Clinicians. 2021;71:488–504.

21. Li L, Abou-Samra E, Ning Z, Zhang X, Mayne J, Wang J, et al. An in vitro model maintaining taxon-
specific functional activities of the gut microbiome. Nat Commun. 2019;10:4146.

22. Murray CJL, Ikuta KS, Sharara F, Swetschinski L, Robles Aguilar G, Gray A, et al. Global burden of
bacterial antimicrobial resistance in 2019: a systematic analysis. The Lancet. 2022;399:629–55.

23. Zhang C, Fu X, Liu Y, Zhao H, Wang G. Burden of infectious diseases and bacterial antimicrobial
resistance in China: a systematic analysis for the global burden of disease study 2019. Lancet Reg
Health West Pac. 2023;43:100972.

24. Hemarajata P, Versalovic J. Effects of probiotics on gut microbiota: mechanisms of intestinal
immunomodulation and neuromodulation. Therap Adv Gastroenterol. 2013;6:39–51.

25. Steidler L, Neirynck S, Huyghebaert N, Snoeck V, Vermeire A, Goddeeris B, et al. Biological
containment of genetically modified Lactococcus lactis for intestinal delivery of human interleukin
10. Nat Biotechnol. 2003;21:785–9.

26. Tanhaeian A, Mirzaii M, Pirkhezranian Z, Sekhavati MH. Generation of an engineered food-grade
Lactococcus lactis strain for production of an antimicrobial peptide: in vitro and in silico
evaluation. BMC Biotechnology. 2020;20:19.

27. Forkus B, Ritter S, Vlysidis M, Geldart K, Kaznessis YN. Antimicrobial Probiotics Reduce
Salmonella enterica in Turkey Gastrointestinal Tracts. Sci Rep. 2017;7:40695.

28. Volzing K, Borrero J, Sadowsky MJ, Kaznessis YN. Antimicrobial peptides targeting Gram-
negative pathogens, produced and delivered by lactic acid bacteria. ACS Synth Biol. 2013;2:643–
50.

29. Bermúdez-Humarán LG, Langella P, Cortes-Perez NG, Gruss A, Tamez-Guerra RS, Oliveira SC, et
al. Intranasal Immunization with Recombinant Lactococcus lactis Secreting Murine Interleukin-12
Enhances Antigen-Specific Th1 Cytokine Production. Infect Immun. 2003;71:1887–96.

30. Yumoto K, Sato T, Nakashima K, Namai F, Shigemori S, Shimosato T, et al. Nasally Administered
Lactococcus lactis Secreting Heme Oxygenase-1 Attenuates Murine Emphysema. Antioxidants
(Basel). 2020;9:1049.

31. Arukha AP, Freguia CF, Mishra M, Jha JK, Kariyawasam S, Fanger NA, et al. Lactococcus lactis
Delivery of Surface Layer Protein A Protects Mice from Colitis by Re-Setting Host Immune
Repertoire. Biomedicines. 2021;9:1098.

32. Chiabai MJ, Almeida JF, de Azevedo MGD, Fernandes SS, Pereira VB, de Castro RJA, et al.
Mucosal delivery of Lactococcus lactis carrying an anti-TNF scFv expression vector ameliorates
experimental colitis in mice. BMC Biotechnology. 2019;19:38.

33. Noguès EB, Kropp C, Bétemps L, de Sousa C, Chain F, Auger S, et al. Lactococcus lactis
engineered to deliver hCAP18 cDNA alleviates DNBS-induced colitis in C57BL/6 mice by promoting
IL17A and IL10 cytokine expression. Sci Rep. 2022;12:15641.

34. Zahirović A, Berlec A. Targeting IL-6 by engineered Lactococcus lactis via surface-displayed
affibody. Microbial Cell Factories. 2022;21:143.

35. Zhu J, Ke Y, Liu Q, Yang J, Liu F, Xu R, et al. Engineered Lactococcus lactis secreting Flt3L and
OX40 ligand for in situ vaccination-based cancer immunotherapy. Nat Commun. 2022;13:7466.

36. Brennan CA, Garrett WS. Fusobacterium nucleatum — symbiont, opportunist and
oncobacterium. Nat Rev Microbiol. 2019;17:156–66.

37. Yasin B, Pang M, Lehrer RI, Wagar EA. Activity of Novispirin G-10, a novel antimicrobial peptide
against Chlamydia trachomatis and vaginosis-associated bacteria. Exp Mol Pathol. 2003;74:190–5.

38. Ghosh SK, Feng Z, Fujioka H, Lux R, McCormick TS, Weinberg A. Conceptual Perspectives:
Bacterial Antimicrobial Peptide Induction as a Novel Strategy for Symbiosis with the Human Host.
Front Microbiol. 2018;9:302.

39. Nakagaki H, Sekine S, Terao Y, Toe M, Tanaka M, Ito H-O, et al. Fusobacterium nucleatum
envelope protein FomA is immunogenic and binds to the salivary statherin-derived peptide. Infect
Immun. 2010;78:1185–92.

40. Bryan EM, Bae T, Kleerebezem M, Dunny GM. Improved vectors for nisin-controlled expression
in gram-positive bacteria. Plasmid. 2000;44:183–90.

41. Li R, Takala TM, Qiao M, Xu H, Saris PEJ. Nisin-selectable food-grade secretion vector for
Lactococcus lactis. Biotechnol Lett. 2011;33:797–803.

42. Steidler L, Remaut ER, Fiers W. Use of a cytokine-producing lactococcus strain to treat colitis.
2000; WO2000023471A2.

43. Steidler L, Robinson K, Chamberlain L, Schofield KM, Remaut E, Page RWFL, et al. Mucosal
Delivery of Murine Interleukin-2 (IL-2) and IL-6 by Recombinant Strains of Lactococcus lactis
Coexpressing Antigen and Cytokine. Infect Immun. 1998;66:3183–9.

44. Choudhury A, Ortiz PS, Young M, Mahmud MdT, Stoffel RT, Greathouse KL, et al. Control of
Helicobacter pylori with engineered probiotics secreting selective guided antimicrobial peptides.
Microbiology Spectrum. 2023;0:e02014-23.

45. O'Toole GA. Microtiter Dish Biofilm Formation Assay. J Vis Exp. 2011.
https://doi.org/10.3791/2437.

46. 16S Illumina Amplicon Protocol : earthmicrobiome. https://earthmicrobiome.org/protocols-
and-standards/16s/. Accessed 8 Sep 2022.

47. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, et al. Reproducible,
interactive, scalable and extensible microbiome data science using QIIME 2. Nature Biotechnology.
2019;37:852–7.

48. Schwager E, Weingart G, Bielski C. CCREPE: Compositionality Corrected by PErmutation and
REnormalization. :16.

49. Elsaghir H, Reddivari AKR. Bacteroides Fragilis. In: StatPearls. Treasure Island (FL): StatPearls
Publishing; 2024.

50. Getz JA, Schoep TD, Daugherty PS. Chapter four - Peptide Discovery Using Bacterial Display and
Flow Cytometry. In: Wittrup KD, Verdine GL, editors. Methods in Enzymology. Academic Press;
2012. p. 75–97.

51. Lozupone CA, Stombaugh JI, Gordon JI, Jansson JK, Knight R. Diversity, stability and resilience of
the human gut microbiota. Nature. 2012;489:220–30.

52. Zhao Q, Chen Y, Huang W, Zhou H, Zhang W. Drug-microbiota interactions: an emerging priority
for precision medicine. Sig Transduct Target Ther. 2023;8:1–27.

53. Shree P, Singh CK, Sodhi KK, Surya JN, Singh DK. Biofilms: Understanding the structure and
contribution towards bacterial resistance in antibiotics. Medicine in Microecology.
2023;16:100084.

54. Liu H, Yu Y, Dong A, Elsabahy M, Yang Y-W, Gao H. Emerging strategies for combating
Fusobacterium nucleatum in colorectal cancer treatment: Systematic review, improvements and
future challenges. Exploration. 2024;4:20230092.

55. Pignatelli P, Nuccio F, Piattelli A, Curia MC. The Role of Fusobacterium nucleatum in Oral and
Colorectal Carcinogenesis. Microorganisms. 2023;11:2358.

56. Liu Z, Wang Y, Zhang C, Yang Y, Zhang J. Engineering Short Antimicrobial Peptides to Specifically
Target Fusobacterium nucleatum in the Mixed Microbial Population. ACS Infect Dis. 2024;10:3042–
51.

57. Wu Q-L, Fang X-T, Wan X-X, Ding Q-Y, Zhang Y-J, Ji L, et al. Fusobacterium nucleatum-induced
imbalance in microbiome-derived butyric acid levels promotes the occurrence and development of
colorectal cancer. World J Gastroenterol. 2024;30:2018–37.

58. Hills RD, Pontefract BA, Mishcon HR, Black CA, Sutton SC, Theberge CR. Gut Microbiome:
Profound Implications for Diet and Disease. Nutrients. 2019;11:1613.

59. Chen Y, Huang Z, Tang Z, Huang Y, Huang M, Liu H, et al. More Than Just a Periodontal Pathogen
–the Research Progress on Fusobacterium nucleatum. Front Cell Infect Microbiol. 2022;12:815318.

60. Han YW, Redline RW, Li M, Yin L, Hill GB, McCormick TS. Fusobacterium nucleatum Induces
Premature and Term Stillbirths in Pregnant Mice: Implication of Oral Bacteria in Preterm Birth. Infect
Immun. 2004;72:2272–9.

61. Muraoka A, Suzuki M, Hamaguchi T, Watanabe S, Iijima K, Murofushi Y, et al. Fusobacterium
infection facilitates the development of endometriosis through the phenotypic transition of
endometrial fibroblasts. Sci Transl Med. 2023;15:eadd1531.

62. Vander Haar EL, So J, Gyamfi-Bannerman C, Han YW. *Fusobacterium nucleatum* and adverse
pregnancy outcomes: Epidemiological and mechanistic evidence. Anaerobe. 2018;50:55–9.

 Supplementary Data

Peptide	Sequence	Mol wt.
Ovispirin	KNLRRIIRKIIHIIKKYG	2263
CPSC5	RGLRRLGRKIAHGVKKYGPTVLRIIRIAG	3256
Stat-Ovispirin	YQPVPEGGGKNLRRIIRKIIHIIKKYG	3148
Stat-CPSC5	YQPVPEGGGRGLRRLGRKIAHGVKKYGPTVLRIIRIAG	4141

Table S1. AMPs used in the study – Ovisiprin and Cathelin-derived Peptide SC5. The statherin derived guide peptide (red) is attached to the N-terminus of the AMPs, separated by a –GGG- linker, to create the corresponding guided AMPs (gAMPs)

Fig S1. MTT Assay of Ovispirin and CPSC5 and their Stat-guided analogs against Caco2 and HT29 cells