PAL-AI reveals genetic determinants that control poly(A)-tail length during

oocyte maturation, with relevance to human fertility

Kehui Xiang[1,2,3*], David P. Bartel[1,2,3*]

[1]Howard Hughes Medical Institute, Cambridge, MA 02142, USA

[2]Whitehead Institute for Biomedical Research, Cambridge, MA 02142, USA

[3]Department of Biology, Massachusetts Institute of Technology, Cambridge MA 02139, USA

[*]For correspondence: dbartel@wi.mit.edu, kxiang@wi.mit.edu

Abstract

In oocytes of mammals and other animals, gene regulation is mediated primarily through
changes in poly(A)-tail length[1–9]. Here, we introduce PAL-AI, an integrated neural network
machine-learning model that accurately predicts tail-length changes in maturing oocytes
of frogs and mammals. We show that PAL-AI learned known and previously unknown
sequence elements and their contextual features that control poly(A)-tail length, enabling
it to predict tail-length changes resulting from 3'-untranslated region (3'-UTR) single-
nucleotide substitutions. It also predicted tail-length-mediated translational changes,
allowing us to nominate genes important for oocyte maturation. When comparing
predicted tail-length changes in human oocytes with genomic datasets of the All of Us
Research Program[10] and gnomAD[11], we found that genetic variants predicted to disrupt tail
lengthening have been under negative selection in the human population, thereby linking
mRNA tail lengthening to human female fertility.

Introduction

Fully grown vertebrate oocytes, which are arrested at prophase I of meiosis, have minimal
transcriptional activity due to the condensation of chromosomes. With little potential for
transcriptional control, post-transcriptional control, particularly translational control,
plays a major role in regulating gene expression during the subsequent processes of oocyte
maturation and early embryonic development, after which the zygotic genome is
transcriptionally activated[12–14]. Proper progression through this period of oocyte maturation
and early development requires translational upregulation of select maternally deposited
mRNAs, such as *MOS* and *CCNB1*[15,16], which encode essential proteins for meiosis, and
TPRX1/2/L[7], which encode transcription factors necessary for zygotic genome activation.
Leveraging the strong coupling between poly(A)-tail length and translational efficiency
during these developmental stages[3,4,6,9,17,18], translation activation of these mRNAs is
achieved by extension of their poly(A) tails[15,19,20], which enables them to compete better for
the binding of limited poly(A)-binding proteins and subsequent recruitment of translation
initiation factors[4]. Thus, disruption of either translational or poly(A) tail-length changes in
these maturing oocytes or early embryos can cause developmental arrest in frogs, mice,
and humans[7,21,22]. Genetic alterations to these processes have also been linked to human
female infertility[23–25].
Control of poly(A)-tail length during oocyte maturation of vertebrate animals is a tug-
of-war between the default transcriptome-wide deadenylation and the mRNA-specific
cytoplasmic polyadenylation[3,8,9,18,26,27]. The specificity of cytoplasmic polyadenylation relies
on two sequence elements within mRNA 3' untranslated regions (UTRs): a cytoplasmic
polyadenylation element (CPE, motif UUUUA), which is recognized by the CPEB1 protein,
and a polyadenylation signal (PAS, motif AWUAAA, where W is either A or U), which is
recognized by the cleavage and polyadenylation specificity factor (CPSF) complex[18,28–31].
Contextual features of these two sequence elements, including their relative positions,
flanking nucleotides, and structural accessibility, as well as the number of each element
within the 3' UTR, influence the extent of cytoplasmic polyadenylation[18]. Although several

other sequence elements have been proposed to influence poly(A)-tail length, their effects
are usually small, indirect, and do not appear to impact more than a handful of
mRNAs[18,30,32,33].
Despite advances in our knowledge of cis-acting elements that control cytoplasmic
polyadenylation, a quantitative understanding of poly(A) tail-length changes for different
mRNAs during oocyte maturation is still lacking. A sequence-based predictive model
would provide further insights into known functional sequence elements and enable the
discovery and evaluation of additional motifs or regulatory principles. For example, such a
model would enable large-scale assessment of the consequences of genetic mutations on
tail-length control as well as translational regulation, thus providing an opportunity to
nominate disease-contributing variants that are otherwise difficult to identify in genome-
wide association studies. Therefore, we set out to develop a machine-learning model that
accurately predicts poly(A) tail-length changes in humans and other vertebrate animals.
76
Results

An integrated neural network model predicts tail-length changes during frog oocyte
maturation
To predict tail-length changes during frog oocyte maturation directly from mRNA
sequences, we initially developed a multiple linear regression model based on *k*-mer
(sequence motif length of *k*) compositions within 3' UTRs. Given the importance of both the
number and position of CPE and PAS elements in influencing tail-length changes[18], we
defined two features for each *k*-mer of varying lengths: 1) a count metric, specifying the
total number of a *k*-mer within the 3' UTR; 2) a positional metric, calculated as the sum of
the inverse distances between each *k*-mer and the 3' end. We performed training and
testing in a 10-fold cross-validation procedure on frog endogenous mRNA tail-length
changes, comparing between 0 h and 7 h post progesterone treatment[18] (Fig. 1a). Test-set
predictions from each fold were concatenated and compared to the measured values (Fig.
1b). We tested different *k*-mer and 3' UTR lengths as inputs and conducted extensive
hyperparameter tuning (Supplementary Fig. 1a). The best model performed moderately
well and explained approximately 40% of the variance of measured tail-length changes
(Fig. 1c; Spearman correlation coefficient R_s = 0.64; Pearson correlation coefficient R_p =
0.63). Notably, the CPE and the PAS, two known motifs mediating cytoplasmic
polyadenylation, were among the top-ranked features in both the count and the positional
metrics (Supplementary Fig. 1b, c).
Although the linear model identified the essential motifs for poly(A)-tail lengthening,
it under-predicted tail-length changes for most mRNAs whose tails were extended by > 30
nt. Inspired by recent advances in deep learning, we developed a neural network model,
termed poly(A)-tail length AI (PAL-AI) (Fig. 1d, Supplementary Fig. 1d, e), which integrates
both convolutional and recurrent neural network components, using an architecture
similar to but simpler than that of Saluki[34]. PAL-AI was trained and tested using the
sequences of endogenous mRNAs, and the same 10-fold cross-validation strategy as used
for the linear model. Combinations of different hyperparameters were tested extensively
for the model to achieve the best performance (Supplementary Fig. 2a, b) without

overfitting (Supplementary Fig. 2c, d). The final trained model predicted tail-length changes
with high accuracy, significantly out-performing the linear model and explaining more than
67% of the variance of measured values (Fig. 1e; R_s= 0.82, R_p = 0.82).
To examine the impact of input sequence length on prediction accuracy, we trained
PAL-AI models on various regions of the 3' UTR and compared Pearson correlation
coefficients between predicted and measured values. A model trained exclusively on the
last 100 nt of 3' UTRs already explained over 40% of the variance (Fig. 1f; average R_p = 0.64).
Model performance improved with increasing 3' UTR length up to 2000 nt, beyond which no
significant gains were observed (Fig. 1f, g). In contrast, when the last 100 nt were excluded
from the last 1000 nt of 3' UTRs for the input, the predictive power declined markedly (Fig.
1f; average R_p = 0.40), supporting the idea that the 3'-most regions of 3' UTRs impart the
strongest effect on tail-length changes[18].
We further explored whether incorporating additional features, such as sequences
from coding regions and base-pairing probabilities predicted by RNAfold[35], could improve
the overall model performance. However, models including these features showed no
significant improvement over those trained on equally-sized 3' UTR-only inputs (Fig. 1f, g),
implying that these features, at least as parameterized in our model, contribute minimally
to tail-length changes. Nevertheless, among all configurations, the model trained on the
last 2000 nt of the 3' UTR plus the coding region performed the best with the shortest input
length (Fig. 1f, g) and was therefore used for subsequent analyses unless otherwise
specified. We also tested a variant model architecture, in which the convolutional module
was replaced with a ResNet block, a design that has demonstrated superior performance
in many deep-learning applications[36,37] . This alternative, although at times achieving
comparable accuracy, was less consistent (Supplementary Fig. 2e) and required
substantially more parameters and computational resources. For this reason, we did not
further optimize this ResNet-based model.
132
PAL-AI learned motifs and contextual features that regulate poly(A)-tail length
To elucidate the principles learned in the training of PAL-AI, we performed in silico
mutagenesis and examined the impact of single-nucleotide mutations on predicted poly(A)
tail-length changes. Among all 6-mers, the loss of the PAS (AAUAAA) and CPE (UUUUA)-
containing 6-mers led to the largest decrease in tail-length changes (Fig. 2a). Sequence
logos generated from the most impacted 8-mers also matched the CPE and the PAS (Fig.
2b). Similarly, when examining the consequence of gaining of 6-mer motifs, those motifs
containing the PAS and CPE resulted in the greatest predicted tail-length increase
(Supplementary Fig. 3a, b). To prevent the signal from spilling from a stronger motif to a
weaker one, we reexamined tail-length changes associated with the loss of 6-mers. We
identified the 6-mer associated with the largest mean tail-length change and then re-
calculated the average tail-length changes for all remaining 6-mers after excluding all point
substitutions that disrupted this top 6-mer. This process was repeated 14 times, each time
excluding point substitutions that disrupted an additional 6-mer, thereby generating a
ranked list of 15 different 6-mers most associated with predicted tail lengthening (Fig. 2c).
Likewise, we used an analogous procedure to reexamine tail-length changes associated

with the gain of 6-mers (Supplementary Fig. 3c). As expected, the top 6-mers identified in
these analyses contained a PAS or CPE.
In addition to the PAS and CPE, our analyses also showed that PAL-AI recognized a
different class of 6-mers, which were associated with modest but statistically significant
tail lengthening (Fig. 2c, Supplementary Fig. 3c). Interestingly, these motifs each contained
a UGU or GUU trimer, which might be recognized by other RNA-binding proteins such as
CELF or DAZL in frog oocytes[38–40]. Another possibility is that some of these motifs, such as
UGUUUU), which resembled the CPE, might be weakly bound by CPEB1. A third possibility
is that the presence of these motifs correlates with tail lengthening but is not causal. To
explore this latter possibility, we examined the efficacy of these motifs in arbitrary
sequences used in a massively parallel reporter 3' UTR library[18]. Because these arbitrary
sequences were not biological, they were not susceptible to noncausal correlative
associations, and thus any association of a motif with a tail-length change was presumably
causal. Analysis of these reporters revealed an association between these UGU/GUU-
containing 6 mers and tail lengthening, but it was small — only a 0.64 nt increase on
average for the top UGU/GUU-containing 6 mer UGUUUU in the absence of a CPE, which
was substantially less than that of a CPE (17.7 nt on average)[18]. In the same library, a
somewhat larger increase in tail length was observed for UGU and GUU (1.8 and 1.4 nt
above an average effect of a 3-mer, respectively) when a CPE was also present in the 3' UTR
(Supplementary Fig. 3d). This added tail-lengthening effect was observed in all CPE-
neighboring positions more than 2 nt from the CPE (Fig. 2d), suggesting that these motifs
might play a small auxiliary role in tail lengthening beyond merely modulating the context of
the CPE. Moreover, for some 3' UTRs, the co-presence of multiple copies of these
UGU/GUU-containing motifs might still contribute to detectable tail lengthening[18], perhaps
through binding to CPEB1-interacting proteins, such as DAZL[41,42], which in turn could
facilitate recruitment of CPEB1 to adjacent CPE-like motifs.
We carried out additional in silico mutagenesis by inserting either the CPE or the
PAS at each position along the 3' UTRs. As expected[18], PAL-AI predicted the strongest
impact on polyadenylation when these elements were introduced near the 3' ends of the 3'
UTRs (Fig. 2e). An added CPE was most favorable when located near the PAS on either its 5'
or 3' side (Fig. 2f), but not in the position most likely to disrupt the PAS, which is typically
~17 nt from the end of the 3' UTR (Supplementary Fig. 3e)[43], consistent with the positional
effects determined from analyses of millions of reporters[18].
Although insertion of the canonical PAS variant AUUAAA moderately promoted
polyadenylation (Fig. 2e), noncanonical PAS variants[44] had negligible effects on
cytoplasmic polyadenylation (Supplementary Fig. 3f). In fact, mutating the canonical PAS
AAUAAA to any PAS variant reduced the predicted tail-length increase by 7.7–11.7 nt
(Supplementary Fig. 3g). Additionally, in the CPE^{mos}-N60 reporter 3' UTR library[18], when
variants containing either an AAUAAA or AUUAAA were excluded, none of the remaining 6-
mers were associated with tail lengthening (Supplementary Fig. 3h). Together, these
analyses indicate that although non-canonical PAS variants can direct mRNA cleavage and
polyadenylation[45], they do not facilitate mRNA polyadenylation in the cytoplasm—at least
not during oocyte maturation.

To gain more insights into properties that PAL-AI learned at the transcript level, we
examined predicted consequences of all possible point substitutions at the last 100 nt of
the *mos.L* mRNA. In silico mutations that perturbed either the PAS at position –21 or the
CPE at position –86 caused considerable decreases in predicted tail lengthening, whereas
a G-to-U mutation at position –62 created a new CPE, thereby causing an increase in
predicted tail lengthening (Fig. 2g). *mos.L* also contains a second CPE falling near the PAS,
at position –27. This PAS-proximal CPE was expected to be suboptimal for cytoplasmic
polyadenylation because its separation from the PAS of only 1 nt was too short[18]. Notably,
PAL-AI recognized this proximity feature and correctly predicted only mild mutational
outcomes when this motif was disrupted (Fig. 2g).
In another example, the *tpx2.L* mRNA, PAL-AI identified two CPEs predicted to be
important for tail lengthening. Furthermore, PAL-AI predicted decreases in tail lengthening
when Gs at positions –66 and –36 (both at –2 positions relative to CPEs) were mutated and
an increase in tail-length change for a C-to-G substitution at position –12 (+1 relative to the
PAS) (Fig. 2h)—all observations consistent with the contextual features of the CPE and the
PAS identified from analyses of millions of reporters[18]. Similar results were observed for
other examples of known substrates of cytoplasmic polyadenylation (Supplementary Fig.
3i, j).
Together, these analyses indicated that PAL-AI successfully captured the sequence
elements and their contextual features important for promoting cytoplasmic
polyadenylation, thereby enabling it to accurately predict poly(A) tail-length changes of frog
mRNAs during oocyte maturation.
214
PAL-AI predicts poly(A) tail-length change of injected mRNAs
Next, we asked if PAL-AI was able to predict tail-length changes of injected reporter
mRNAs. To this end, we amplified a small subset of a 3'-UTR reporter library (N60-PAS^{mos})[18]
to generate a library with only ~35,000 variants (Fig. 3a), which we called N60(LC)-PAS^{mos}.
The reduced sequence complexity of this library enabled acquisition of tail-length
distributions of individual mRNA sequences. We injected this mRNA library into prophase I-
arrested frog oocytes, collected RNA samples at 0 and 7 h post-progesterone treatment,
and used high-throughput sequencing[18] to identify the 3' UTR variant and determine its tail-
length distribution at each of the two stages of maturation (Fig. 3b).
When using tail-length changes measured for these injected mRNAs to test our
model trained on endogenous mRNAs, our model performed well (Supplementary Fig. 4a).
Indeed, correlation coefficients observed when testing on the injected mRNAs (R_s= 0.77, R_p
= 0.83) resembled those observed when testing on endogenous mRNAs (Fig. 1e, R_s= 0.82,
R_p = 0.82). However, the model appeared to under-predict the magnitude of the tail-
lengthening effects (Supplementary Fig. 4a). A few possibilities could explain the
systematic difference in the magnitude of tail-lengthening observed for injected mRNAs
compared to that predicted by PAL-AI. First, the sequence lengths of the training and the
testing datasets were markedly different. Endogenous mRNAs used for training PAL-AI had
few 3' UTRs shorter than 100 nt (420 out of 6054), whereas 3' UTRs of the injected mRNAs
were all relatively short (81 nt). Second, due to experimental variability, particularly the
variable time it takes for oocytes of different batches to mature, the endogenous mRNAs

and injected mRNAs may have been collected at somewhat different maturation stages.
Third, the 3' UTRs of the reporter mRNAs consisted of arbitrary sequences, whose
composition may differ substantially from natural mRNA sequences.
To test whether sequence composition contributed to the discrepancy, we modified
PAL-AI's last layer to enable dataset-specific multi-output predictions (Fig. 3c). Training on
alternating batches of endogenous and injected mRNAs maintained performance on
endogenous mRNAs (Fig. 4d, R_s= 0.82, R_p = 0.82) while enhancing performance on injected
mRNAs (Fig. 4e, R_s= 0.81, R_p = 0.88). To distinguish this multi-output model from the
original PAL-AI model, we termed this model as PAL-AI-m ("m" for multi-output) and the
original model as PAL-AI-s ("s" for single-output).
Despite the improvement, the PAL-AI-m model still under-predicted tail lengthening
for most long-tailed (≥ 80 nt) mRNAs from the injected library. When we re-trained the
model with measurements from only injected mRNAs, PAL-AI performed significantly
better when tested on held-out data (Supplementary Fig. 4b, R_s= 0.86, R_p = 0.91). In silico
mutagenesis analyses indicated that the new model also learned the expected sequence
elements important for tail lengthening and their contextual features (Supplementary Fig.
4c–e). This training approach on synthetic, nonbiological reporter sequences also ensured
that these learned elements and features are causative for tail-length changes, rather than
correlated to causative features.
255
Predicted effects of single-nucleotide substitutions correspond with experimentally
determined effects.
The ability of PAL-AI to predict the fate of injected mRNAs enabled direct assessment of its
ability to predict the consequences of mutations, including all possible single-nucleotide
substitutions. To this end, we generated a single-nucleotide mutagenesis library of reporter
mRNAs with 3' UTR sequences derived from the last 100 nt of 3' UTRs of ten mRNAs
selected from frog oocytes. For each of the ten 100-nt UTR fragments, each nucleotide of
the 3' UTR was substituted with one of the three alternative nucleotides, resulting in 3,000
variants (Fig. 4a). These 3,000 variants, together with the non-substituted wild-type
molecules, were injected into oocytes and the effect of each substitution on tail-length
changes was analyzed over the course of maturation. Of the ten 3'-UTR fragments, eight
(*atp1a1.S*, *ccnb1.2.L*, *ccnb2.L*, *lima1.L*, *mad2l1.L*, *magoh.S*, *mos.L*, and *tpx2.L*) directed
cytoplasmic polyadenylation upon injection into oocytes, whereas two (*aurkaip1.L* and
dbf4.L) directed no tail lengthening during frog oocyte maturation[18].
When comparing the predicted difference in tail-length change to that measured
between each mutant and the wild-type, the PAL-AI-m model performed significantly better
compared to the PAL-AI-s model for eight out of ten mRNAs (Fig. 4b). This improved
performance suggested that the PAL-AI-m model successfully learned additional features
from the non-natural sequences of the N60(LC)-CPE^{mos} library. The PAL-AI-m model
achieved high accuracy (R_p: 0.78–0.90) for six mRNAs (*atp1a1.S*, *lima1.L*, *mad2l1.L*,
magoh.S, *mos.L*, and *tpx2.L*; Fig. 4b, d–g, Fig. 5a, b, Supplementary Fig. 5a–f) and moderate
accuracy (R_p: 0.58–0.62) for three mRNAs (*aurkaip1.L*, *ccnb1.2.L*, and *ccnb2.L*; Fig. 4b,
Supplementary Fig. 6a, b, d, e, Supplementary Fig. 7a, b).

Concordance between predicted and experimental outcomes was particularly clear
for substitutions that disrupted either a CPE or PAS (Fig. 4c, f, g, Fig. 5a, Supplementary Fig.
5d–f). Substituting the canonical PAS (AAUAAA) with the alternative variant AUUAAA
severely affected tail lengthening, but not as severely as observed for any of the other
single-nucleotide substitutions of AAUAAA (Fig. 4c). These experimental findings validated
our in silico mutagenesis conclusion that noncanonical PAS variants do not support
cytoplasmic polyadenylation during oocyte maturation.
At the individual mRNA level, ablation of any of the three motifs (two CPEs and one
PAS) within *mos.L* mRNA led to substantial decreases in tail lengthening, as predicted by
PAL-AI-m (Fig. 4f). The predicted weaker contribution to cytoplasmic polyadenylation by
the CPE proximal to the PAS compared to that by the CPE further upstream was also
validated (Fig. 4f). Similar results illustrating PAL-AI-m's ability to predict the importance of
these motifs and their contextual features were also observed for other mRNAs (Fig. 4g,
Supplementary Fig. 5). Notably, the predicted differences in tail-length changes observed
between many variants and the wild-type were smaller than those measured, perhaps due
to the same reasons previously put forward to explain the under-prediction of changes for
the injected N60(LC)-PAS^{mos} library (Supplementary Fig. 4a).
In another example, *mad2l1.L* mRNA had no CPE within the last 100 nt of its 3' UTR
but contained seven UGU/UGUUU motifs (Fig. 5a). Mutations within these motifs caused
modest but consistent reductions in tail lengthening (Fig. 5a, b), agreeing with their
detectable but weak roles in cytoplasmic polyadenylation. In contrast, substitutions that
introduced a single CPE led to large increases in tail lengthening, almost all of which were
correctly predicted by PAL-AI-m (Fig. 5a, b).
Despite achieving good performance for the six aforementioned mRNAs (R_p = 0.78–
0.9), PAL-AI-m performed only moderately well on three mRNAs (*aurkaip1.L*, *ccnb2.L*, and
ccnb1.2.L, R_p = 0.58 , 0.62, and 0.59, Supplementary Fig. 6b, e, 7a) and did not perform well
at all on one mRNA (*dbf4.L*, R_p = 0.21, Fig.5c). The wide-type *dbf4.L* mRNA, which had no
CPE and a relatively weak PAS (AUUAAA), had little tail lengthening, and thus most single-
nucleotide substitutions had negligible opportunity to reduce tail lengthening (Fig. 5d),
which largely explained their relatively weak effects for this mRNA. PAL-AI-m did predict
moderate increases in tail lengthening for a few substitutions that would create CPEs
predicted to lengthen tails, but by no more than 10 nt (Fig. 5c). Although some of these
predicted increases were validated (positions –61 and –51), others, particularly those
downstream the PAS (positions –35, –31, –7 and –6) were not (Fig. 5d). To understand these
discrepancies, we predicted the secondary structure of the last 100 nt of the wild-type
dbf4.L 3' UTR using EternaFold[46]. The substitutions at positions –61 and –51 both created
CPEs in loop regions, whereas the substitutions at positions –35, –31, –7 and –6 all created
CPEs with nucleotides paired in a stem loop (Fig. 5d, e), which likely prevented these CPEs
from being recognized by the CPEB1 protein.
For *aurkaip1.L* and *ccnb2.L* mRNAs, PAL-AI-m had difficulty predicting increased tail
lengthening caused by substitutions in consecutive positions within a few UTR segments
(*aurkaip1.L*: positions –34 to –30 and –19 to –15, Supplementary Fig. 6a; *ccnb2.L*: positions
–60 to –55 and –15 to –10, Supplementary Fig. 6d). These substitutions appeared to
destabilize stem loops that were sequestering PAS elements in the wild-type UTRs

(Supplementary Fig. 6c, f), thus making these elements more accessible for CPSF binding.
In support of this hypothesis, substitutions that further stabilized these stem loops
reduced tail lengthening (*aurkaip1.L*: A-to-G at position –35 and C-to-U at position –14,
Supplementary Fig. 6a, c; *ccnb2.L*: C-to-G at position –54 and C-to-G at position –16,
Supplementary Fig. 6d, f). Together, these analyses indicated that sequences within the 3'
UTR can indirectly affect poly(A) tail-length changes by modulating the structural
accessibility of the CPE and the PAS, a mechanism that likely applies to select mRNAs[33]
and thus had not yet been effectively learned by PAL-AI-m.
As for the *ccnb1.2.L* mRNA, PAL-AI-m under-predicted the effects of a CPE that
overlapped with the PAS (Supplementary Fig. 7a, b). Binding of CPEB1 to this overlapping
CPE presumably prevented CPSF from recognizing the PAS, thus inhibiting cytoplasmic
polyadenylation as previously suggested[47]. Supporting this hypothesis, the C-to-U
mutation at position –18 of *magoh.S* mRNA created an overlapping CPE, which reduced tail
lengthening (Supplementary Fig. 5f).
Despite some discrepancies between the predictions and experimental results,
these analyses demonstrated that, overall, PAL-AI can accurately predict tail-length
changes caused by single-nucleotide mutations within mRNA 3' UTRs.
340

PAL-AI predicts poly(A) tail-length change in mammalian oocytes
Because principles of poly(A) tail-length control during oocyte maturation are highly
conserved among frogs, mice, and humans[18], we asked if PAL-AI trained from frog oocyte
data could predict tail-length changes during mammalian oocyte maturation. For mouse
oocytes, PAL-AI achieved moderate accuracy (Supplementary Fig. 8a; R_s= 0.57, R_p = 0.58).
The performance of the model was likely affected by unmatched developmental stages
between frogs and mice, and also global deadenylation that was independent of 3'-UTR
sequences[9,18]. Similar predictive power for PAL-AI was also observed for human oocytes
using published datasets (Supplementary Fig. 8b; R_s= 0.61, R_p = 0.59), even though the
human measurements were acquired using a different sequencing protocol and platform
(PacBio)[8].
To minimize prediction errors arising from species-specific sequence composition
and experimental variability, we developed a multi-species PAL-AI model trained on pooled
data from frogs, mice, and humans, using the same approach as for our frog endogenous
mRNAs and injected mRNAs. This model, termed PAL-AI-ms ("ms" for multi-species),
showed improved performance for both mouse (Fig. 6a; R_s= 0.63, R_p = 0.64) and human
oocytes (Fig. 6b; R_s= 0.68, R_p = 0.66), while maintaining comparable accuracy for frog
oocytes (Supplementary Fig. 8c; R_s= 0.79, R_p = 0.77). In addition, the model's predictions
also correlated well with independent tail-length measurements obtained using Oxford
Nanopore Technologies (ONT) for mRNAs during mouse oocyte
maturation[9] (Supplementary Fig. 8d; R_s= 0.69, R_p = 0.69). Although ONT data showed larger
absolute tail-length changes compared to our measurements obtained using Illumina
sequencing (HiSeq 2000), both datasets exhibited consistent trends (Supplementary Fig.
8e). This systematic difference in magnitude likely explained why the model under-
predicted tail-length changes in the ONT data (Supplementary Fig. 8d).

In frog, mouse, and human oocytes, tail-length changes strongly correlate with
translational efficiency changes[3,9,18]. As a result, our model-predicted tail-length changes
were expected to be informative regarding up- and down-regulation of mRNA translation—
not only for endogenous mRNAs but also for any mRNAs with known 3'-UTR sequences. To
confirm this expectation, we predicted tail-length changes for a series of reporter mRNAs
with different 3'-UTR sequences used for assessing translational regulation during mouse
oocyte maturation[6]. Our model-predicted tail-length changes corresponded well with the
reported translational efficiency changes (Fig. 6c; R_s= 0.91, R_p = 0.89).

Although tail-length control during oocyte maturation is highly conserved among
vertebrates, tail lengthening and associated translational upregulation are known to be
required for mRNAs from only a few genes. Conserved tail lengthening of mRNAs from
orthologous genes may shed light on the functional significance of tail lengthening during
oocyte maturation and early development, but mRNAs from only 19 genes with conserved
tail lengthening have been identified[18]. Part of the reason for this small number is that most
mRNAs could not be evaluated because they lacked tail-length measurements in at least
one of the three species[18]. To overcome this limitation, we used our model-predicted tail-
length changes for cases in which tail-length measurements were unavailable, resulting in
the identification of another 264 genes (57, 108, and 99 with one, two, and three predicted
values respectively) whose mRNA poly(A) tails were either extended or predicted to be
extended ≥ 15 nt in all three species (Fig. 6d, Supplementary Fig. 8f, g). Most of these
mRNAs had increased translational efficiency, and for those for which translational
efficiency changes had not been measured, PAL-AI-predicted tail-length increases could
be informative. A Gene Ontology analysis of these genes revealed enrichment of molecular
pathways such as cell cycle, microtubule cytoskeleton organization, cell division, and
chromosome organization, which are crucial for meiosis and presumably prepare the
zygote for cell divisions of early embryonic development (Fig. 6e).

Variants predicted to disrupt poly(A)-tail lengthening are under negative selection in
humans and throughout vertebrates
Next, we asked if genetic variants predicted to influence tail-length changes during oocyte
maturation have been under evolutionary selection. We analyzed human 3' UTRs by using
PAL-AI-ms to predict the effects of all possible single-nucleotide mutations within their last
100 nt. For each position, we calculated the average difference in predicted tail-length
change when mutated to alternative nucleotides and classified the wild-type nucleotide by
the direction of the effect and severity when mutated: tail-length reduction(severe [≤ –25
nt], moderate [–25 to –15 nt], mild [–15 to –5 nt]), minimal-impact [–5 to 5 nt], and tail-
length increase (mild [5 to 15 nt], moderate [15 to 25 nt], severe [> 25 nt]). Comparing these
predictions to phyloP scores, a conservation metric based on the alignment of 100
vertebrate genomic sequences[48,49], revealed strong evolutionary constraints, with
conservation levels tracking predicted functional impact. For example, nucleotides for
which mutations would cause severe tail-length reduction showed significantly higher
phyloP scores than moderate-impact sites (Mann–Whitney U test), which in turn were more
conserved than mild-impact sites, with minimal-impact sites being least conserved (Fig.
7a, Supplementary Fig. 9a). Nucleotides for which mutations would cause tail-length

increase showed a parallel but weaker trend, with only mild/moderate-impact sites
exhibiting significantly elevated conservation relative to minimal-impact sites (Fig. 7a,
Supplementary Fig. 9a). This is likely in part because tail-length shortening during oocyte
maturation is non-specific[9,18,26,27] and therefore very few sites, when mutated, would result
in tail-length increase (Fig. 7a). Overall, these findings indicate that genomic positions at
which mutations would substantially alter poly(A) tail-length changes during oocyte
maturation have been under purifying selection in the vertebrate lineage.
We also asked if genetic variants predicted to influence tail-length control during
oocyte maturation might have functional consequences for human health. To this end, we
analyzed ~8.4 million 3'-UTR variants from the All of Us Research Program[10] using PAL-AI-
ms. For each variant, we calculated its predicted effect size as the difference in predicted
tail-length change relative to the reference allele (Fig. 7b). Focusing on the last 100 nt of
each 3' UTR (~ 0.85 million variants), we binned variants by predicted effect size the same
way as we had done for the conservation analysis and examined their relative fractions in
each bin for groups with different allele frequencies. This analysis revealed a striking allele-
frequency-dependent depletion of tail-lengthening-disrupting variants. For example,
variants predicted to severely disrupt tail-lengthening showed 2-, 3-, 7- and 10-fold
depletion in common variants (allele frequencies > 0.01%, > 0.1%, > 1%, and > 10%,
respectively) compared to singletons (P < 0.001, Fisher's exact test). Depletion of frequent
alleles was also significant, albeit to a lesser degree, for variants predicted to impart
moderate (1.5–1.9 fold) and mild (1.3–1.7 fold) reduced tail-lengthening (Fig. 7c). This
depletion pattern remained significant even after excluding all variants that would
introduce or ablate PAS motifs (Supplementary Fig. 9b), which were presumably also under
negative selection due to their nuclear role in specifying mRNA cleavage and
polyadenylation[50]. In contrast, gain-of-function variants were not consistently depleted in
common alleles compared to singleton variants (Fig. 7c). Similar results were observed
when PAL-AI was applied to the gnomAD v4.1 datasets[11] (Fig. 7d, Supplementary Fig. 9c, d).
Together, these analyses indicated that variants that disrupt poly(A) tail-length control,
particularly those that disrupt tail lengthening during oocyte maturation, are under negative
selection in humans.
440

Discussion

We developed PAL-AI, a neural network model that predicted tail-length changes in frog,
mouse, and human oocytes using only sequence information. PAL-AI captured
consequential sequence motifs and contextual effects without any feature generation,
thereby allowing us to predict tail-length changes of mRNAs with any sequence. Moreover,
PAL-AI helped us identify UGU/GUU-containing motifs contributing to poly(A)-tail
lengthening. Although previously implicated in regulating translation[38,41,42,51–53], these motifs
had not been associated with tail-length control. Our analyses of the injected mRNA
reporter libraries further demonstrated that their tail-lengthening effects were causal rather
than correlative with other features of endogenous mRNAs. Individually, these motifs
contributed weakly to polyadenylation, but multiple copies did result in considerable tail
lengthening, as illustrated by the *mad2l1.L* mRNA. Their precise roles in affecting tail

lengths might be context-dependent, as some might serve as CPE-like motifs
promiscuously bound by CPEB1, whereas others might be recognized by RNA-binding
proteins such as DAZL, which also interacts with CPEB1 and thus might help facilitate
recruiting CPEB1 to the 3' UTR.
The ability of PAL-AI to predict tail-length changes of both endogenous mRNAs and
injected mRNAs enabled us to capture the consequences of most single-nucleotide
substitutions. For a few mRNAs, PAL-AI under-performed in predicting the outcomes of
substitutions—often those substitutions that appeared to alter the structural accessibility
of CPE or PAS elements. However, when we combined base-pairing probability, as
predicted by RNAfold[35], with mRNA sequences as the input for our model, the performance
did not significantly improve, which suggested that more accurate parameterization of
mRNA structure would be needed for the model to account for structural accessibility.
Perhaps incorporating data from in-oocyte structural probing of mRNAs will further improve
the model predictions[54–57].
Because of the strong influence of tail length on translational efficiency in oocytes,
we envision that our model can be used to engineer mRNA 3'-UTR sequences for desired
protein expression in these systems. Furthermore, by predicting tail-length changes that
had not yet been measured in frog, mouse, or human oocytes, our model helped identify a
group of genes that appear to undergo conserved mRNA tail-lengthening during oocyte
maturation. Although for mRNAs of most of these genes, the importance of translational
up-regulation during oocyte maturation and early embryogenesis still needs to be
examined, knowledge of these genes can inform efforts to identify mutations that affect
human female fertility.
Indeed, genetic variants predicted by PAL-AI to disrupt tail-length control during
oocyte maturation have been under negative selection in vertebrates including humans.
Presumably, some of these variants perturb sequence elements or contextual features
important for cytoplasmic polyadenylation, thus leading to failed translational activation of
mRNAs encoding proteins important for oocyte maturation or early embryonic
development. While not evidently observed in this study, as larger genomic data sets
become available and more sensitive analyses are conducted, negative selection might
also be detected for mutations predicted to lead to inappropriate activation of
translationally repressed mRNAs during these developmental transitions.
Although the observed depletion of variants predicted to disrupt tail lengthening
persisted after excluding PAS-perturbing substitutions, we cannot rule out that some of this
selection pressure may reflect pleiotropic effects on other RNA-processing events. For
instance, U-rich sequences resembling the CPE near the mRNA 3' end can serve as binding
platforms for Fip1 in nuclear pre-mRNA processing[58,59]. Variants disrupting these motifs
could simultaneously impair both cytoplasmic polyadenylation and nuclear 3' end
formation. Future integration of PAL-AI with tools capable of predicting 3' processing
defects, such as APARENT2[50] or PolyaID[60] , could help distinguish these potentially
confounding selective pressures.
Past efforts to identify genetic causes of female infertility have primarily focused on
variants in coding regions, perhaps in part due to the use of exome sequencing[23]. However,
mutations that explain defects in oocytes and early embryos associated with female

infertility fall in fewer than 20 genes and are found in only a small portion of patients[61].
Given the importance of tail-length and translational control in human oocytes and early
embryos, as well as our analyses of variant abundancies in the human population, some
unexplained cases are presumably due to disruptive mutations occurring in 3' UTRs. With
more genomes with female reproductive failures being sequenced, we will be able to start
assessing contributions by these variants, and PAL-AI can be a valuable tool
complementing current approaches, such as genome-wide association studies, to help
nominate causal variants within 3' UTRs.

Acknowledgments

507 We thank Thy Pham, Peter Wang, and other current and former members of the Bartel lab
508 for helpful discussions, Sumeet Gupta for assistant with data collection, and the
509 Whitehead Institute Genome Technology Core for sequencing. D.P.B. is an investigator of
510 the Howard Hughes Medical Institute.

Author contributions

513 K.X. and D.P.B. designed the study and wrote the manuscript. K.X. conceived the project
514 and performed all the experiments and analyses.

Competing interests

The authors declare no competing interests.

Fig. 1

Fig. 1: Machine-learning models predict tail-length changes during frog oocyte
maturation.
a Experimental scheme for examining poly(A) tail-length changes of frog mRNAs during
oocyte maturation. Total RNA was extracted from oocytes before and after progesterone-
induced germinal vesicle breakdown (GVBD), and changes in poly(A) tail lengths were
measured. **b** Schematic of the 10-fold cross-validation strategy used to train and test
different machine-learning models in this study. Data were partitioned into
training/validation and test sets, repeated across 10 different stratified folds. **c**
Performance of the multiple linear regression model. Plotted are the tail-length changes
predicted by the model as a function of the changes measured in frog oocytes between 0
and 7 h post-progesterone treatment. Each point represents a unique poly(A) site of an
endogenous mRNA. Colors indicate the density of points. **d** Diagram outlining the two

machine-learning models developed to predict poly(A) tail-length changes from mRNA
sequence features: a multiple linear regression model and an integrated neural network
(PAL-AI). **e** Performance of PAL-AI; otherwise as in (**c**). **f** Prediction performance of PAL-AI
trained on different input regions of mRNA or additional annotation features. Left: input
sequence regions (bars) and additional features (CDS or predicted pairing, blue and red
dots, respectively). Right: distributions of R_p values observed when comparing predicted
and measured tail-length changes for test data held out during training. Ten-fold cross-
validation of the model was repeated five times, generating 50 R_p values. The red rectangle
indicates the configuration chosen as the final model. CDS, coding sequence. Box and
whiskers indicate the 10th, 25th, 50th, 75th, and 90th percentiles. **g** Pairwise comparison
of input strategies used for PAL-AI based on prediction performance. Shown are binned P
values from one-sided t-tests, testing the alternative hypothesis that the mean R_p value of
the group indicated on the y axis is greater than that indicated on the x axis. The red
rectangle indicates the configuration chosen as the final model.
547
548

Fig. 2

Fig. 2: PAL-AI learned sequence elements and contexture features important for poly(A) tail-length control during oocyte maturation.

a PAL-AI-predicted consequences of motif loss. Shown for each 6-mer is the mean difference between predicted tail-length changes of mutants compared to those of the wild-type. These mean differences are plotted as a function of the mean number of analyzed mutants. Red points represent motifs that contained a CPE and blue points represent those that contained a complete or partial PAS. Only motifs within the last 300 nt of 3' UTRs were included in this analysis. **b** Sequence motifs associated with PAL-AI-predicted changes in poly(A)-tail length. Shown are sequence logos generated from 8-mers

most associated with the largest differences in predicted tail-length change between
mutants in which 8-mers were lost and the wild-type. The pie charts on the left indicate the
fraction of 8-mers that aligned to the corresponding sequence logo. The bar plots on the
right show the mean difference, with points representing individual 8-mers. Only motifs
within the last 300 nt of 3' UTRs were included in this analysis. **c** Top 6-mers associated
with decreased predicted tail-length change in the in silico mutagenesis analysis.
Differences in tail-length change were calculated in an iterative process with exclusion. In
each round, differences in tail-length change were calculated for all 6-mers lost in the
mutants, and the 6-mer associated with the largest decrease in tail-length changes was
selected. In the following rounds, all mutants in which this top 6-mer was disrupted were
excluded before this procedure was repeated. Motifs are colored red if they contain a CPE,
blue if they contain a PAS, orange if they contain a GUU or UGU, and black if otherwise.
Only motifs within the last 300 nt of 3' UTRs were included in this analysis. **d** Positional tail-
lengthening effects of 3-mers flanking a CPE. Plotted for each 3-mer are differences in
mean tail lengths observed for mRNAs with that 3-mer at the indicated positions relative to
a CPE in the 3' UTRs of the N60-PAS^{mos} library[18], comparing between 0 and 7 h post-
progesterone treatment. Only variants containing one CPE (UUUUA) were included in this
analysis. The shaded area indicates positions where 3-mers impact the flanking nucleotide
context of a CPE[18]. **e** Positional effects predicted by PAL-AI for the CPE and PAS. Plotted for
each of the motifs are mean differences between predicted tail-length changes conferred
upon inserting the motif in silico at the indicated positions along the 3' UTR. The drop
observed when inserting a motif near position 17 was attributed to disruption of the PAS
(**Supplementary Fig. 3e**). Shaded areas along the lines indicate standard error. The inset
shows results for the last 100 nt of the 3' UTR. **f** Effects of the relative position between the
CPE and the PAS predicted by PAL-AI. Plotted are the mean differences between predicted
tail-length changes conferred upon inserting a CPE in silico as a function of the relative
distance between the CPE and the PAS. Shaded areas along the lines indicate standard
error. The rectangular shaded area indicates positions where the inserted CPE would
overlap with the PAS. **g** Impact of single-nucleotide substitutions on predicted tail-length
changes. Shown are results for the last 100 nt of the *mos.L* mRNA 3' UTR. The heatmap
indicates the difference in predicted tail-length change (DPTLC) as a result of changing
each nucleotide (*x* axis) to each of the three non-wild-type alternatives (*y* axis), with x
indicating the non-substituted, wild-type nucleotide. The line plot indicates the maximum
(red line) and the minimum (blue line) of the mutational outcomes at each position. The
logo plot indicates the importance of each nucleotide, with the height normalized to the
negative value of the average outcome of the three possible substitutions. Dashed
rectangles indicate model-identified sequence elements (the CPE and PAS) that are
important for poly(A)-tail lengthening. The arrow points to an instance of a new CPE,
generated by a G-to-U substitution, and its associated increase in tail-length change. **h**
Impact of single-nucleotide substitutions in the *tpx2.L* mRNA 3' UTR on predicted tail-
length changes, plotted as in (**g**). The arrow with a solid line points to an example of PAS
flanking-nucleotide effects, in which a C-to-G substitution at the +1 position increases
predicted tail-length change[18]. The arrows with dashed lines point to examples of CPE

flanking nucleotide effects, in which substitutions of G at the –2 position decrease
predicted tail-length change[18].
603
604

Fig. 3

Fig. 3: PAL-AI predicts tail-length changes of synthetic mRNAs in frog oocytes.
a Schematic of the N60(LC)-PAS^{mos} mRNA library used for injection. **b** Experimental
scheme for mRNA library injection and sample collection. **c** Diagram of the architecture of
the multi-output PAL-AI model (PAL-AI-m). **d** Performance of the PAL-AI-m model in
predicting tail-length changes of frog endogenous mRNAs. Otherwise, this panel is as in
Fig. 1c. **e** Performance of the PAL-AI-m model in predicting tail-length changes of mRNAs
in the N60(LC)- PAS^{mos} library. Otherwise, this is as in **Fig. 1c**.

Fig. 4

Fig. 4: Experiments validate predicted effects of single-nucleotide substitutions.
a Schematic of the library of single-nucleotide substitutions used for injection. In addition
to the 3000 variants with single-nucleotide substitutions, mRNAs with the wild-type
segments were also included at a proportion of one wild-type for every 300 single-
nucleotide variants. **b** Summary of the relationships between the PAL-AI-s- or PAL-AI-m-
predicted effects of single-nucleotide substitutions and the measured effects of those
substitutions. For each of the 10 mRNAs in the single-nucleotide substitution library,
Pearson *R* values were calculated based on the differences in tail-length changes predicted
by the indicated model (key) and those measured for each single-nucleotide substitution,
comparing between 0 h and 7 h post-progesterone treatment in frog oocytes. The top-
performing model trained from each fold of the ten-fold cross-validation was used for
predictions (ten models total for each architecture). *P* values are from one-sided t-tests.
Box and whiskers indicate the 10th, 25th, 50th, 75th, and 90th percentiles. n.s.: not
significant (*P* ≥ 0.01). **c** PAL-AI-m prediction of the effects of single-nucleotide
substitutions that disrupt either a CPE or a PAS (AAUAAA). For each position of an element,
the average difference in tail-length change predicted upon substitution with one of the
three non-wild-type possibilities is compared to that measured for mRNA injected into frog
oocytes and collected 0 and 7 h post-progesterone treatment. Points for substitutions that
disrupt a CPE are in blue, and points for those that disrupt a PAS are in red. Error bars
indicate the standard error of the mean of all library variants that had the indicated
substitutions. **d** PAL-AI-m prediction of the effects of single-nucleotide substitutions of the
mos.L mRNA. Plotted for each single-nucleotide substitution is the difference in tail-length
change predicted by PAL-AI-m compared to that measured for that mRNA variant injected
into frog oocytes and collected 0 and 7 h post-progesterone treatment. On the sides are
density distributions for the predicted and measured differences in tail-length changes. **e**
PAL-AI-m prediction of the effects of single-nucleotide substitutions of the *tpx2.L* mRNA;
otherwise, as in (**d**). **f** Impact of single-nucleotide substitutions on predicted and measured
tail-length changes for the *mos.L* mRNA. Shown are results for the last 100 nt of the *mos.L*
3' UTR. The heatmaps indicate the predicted (top) and measured (bottom) differences in
tail-length changes (DTLC) as a result of changing each nucleotide (*x* axis) to each of the
three non-wild-type alternatives (*y* axis), with x indicating the non-substituted, wild-type
nucleotide. The line plots indicate the predicted (top) and measured (bottom) maximum
(red line) and minimum (blue line) of the mutational outcomes at each position. The logo
plots indicate the predicted (top) and measured (bottom) importance of each nucleotide,
with the height normalized to the negative value of the average outcome of the three
possible substitutions. Dashed rectangles indicate sequence elements (the CPE and PAS)
that are important for poly(A)-tail lengthening. **g** Impact of single-nucleotide substitutions
on predicted and measured tail-length changes for the *tpx2.L* mRNA; otherwise as in (**f**).
654
655

Fig. 5

Fig. 5: Auxiliary motifs and structural accessibility influence tail-length changes.
a Impact of single-nucleotide substitutions on predicted and measured tail-length changes for the *mad2l1.L* mRNA. Dashed rectangles indicate UGU/UGUUU elements that each appears to modestly promote poly(A)-tail lengthening. The arrows point to instances of new CPEs generated by substitutions and their associated increases in tail-length changes. Otherwise, this panel is as in **Fig. 4f**. **b** PAL-AI-m prediction of the effects of single-nucleotide substitutions of the *mad2l1.L* mRNA. Otherwise, this panel is as in **Fig. 4d**. **c** PAL-AI-m prediction of the effects of single-nucleotide substitutions of the *dbf4.L* mRNA.

Otherwise, this panel is as in **Fig. 4d**. **d** Impact of single-nucleotide substitutions on
predicted and measured tail-length changes for the *dbf4.L* mRNA. Dashed rectangles
indicate two regions predicted to pair with each other, forming the stem of a hairpin. Arrows
indicate CPE-creating substitutions with tail-length changes either correctly (solid) or
incorrectly (dashed) predicted by PAL-AI-m. **e** EternaFold-predicted maximum expected
accuracy secondary structure of the last 100 nt of the *dbf4.L* 3' UTR. The dashed rectangle
indicates the stem of the hairpin highlighted in (**d**). Triangles indicate positions where CPE-
creating substitutions with tail-length changes either correctly (solid) or incorrectly
(hollow) predicted by PAL-AI-m, as shown in (**d**). The PAS is shaded light blue. Base pairings
are indicated with thick lines for G–C, thin lines for A–U, and a dot for the G–U wobble.
674
675

Fig. 6

Fig. 6: Multi-species PAL-AI predicts tail-length changes of endogenous mRNAs of mouse and human oocytes.
a Performance of the multi-species PAL-AI model on mouse mRNAs. Plotted are the tail-length changes predicted by PAL-AI-ms to occur upon mouse GV-to-MII oocyte maturation as a function of the measured changes. Each point represents a unique poly(A) site of an endogenous mRNA. Otherwise, this panel is as in **Fig. 1c**. **b** Performance of the multi-species PAL-AI model on human mRNAs. Otherwise, this panel is as in (**a**). **c** Ability of predicted tail-length changes to explain translational efficiency changes. Plotted is the relationship between the multi-species PAL-AI model-predicted tail-length changes and translational efficiency changes measured for seven mRNA reporters upon mouse GV-to-MII oocyte maturation[6]. **d** Genes predicted or shown to have substantial mRNA tail-lengthening (≥ 15 nt) in human (*H.s.*), mouse (*M.m.*), and frog (*X.l.*) oocytes. Heatmaps show measured or predicted (dots) tail-length changes (left) and measured translational efficiency changes (right), comparing between human GV and MII oocytes, mouse GV and MII oocytes, and frog oocytes 0 and 7 h post-progesterone treatment. Only genes with the model-predicted values in one of the three species were shown. Gray indicates values not available. **e** Select ontologies enriched for genes with substantial mRNA tail-lengthening (≥ 15 nt, predicted, if not measured) in human, mouse, and frog oocytes.

Fig. 7

Fig. 7. Variants predicted to disrupt poly(A)-tail lengthening are selected against in the human population and throughout vertebrates.

a Negative selection among sequenced vertebrate species of variants predicted to be more disruptive of tail lengthening. Shown are distributions of phyloP scores[48,49] , grouped by the average PAL-AI–predicted differences in tail-length changes for individual nucleotides when substituted to each of the three alternative nucleotides. Only positions within the last 100 nt of 3' UTRs were included in this analysis. For the tail-length difference bin, parentheses indicate values not included, while square brackets indicate values included. Numbers in the parentheses next to the bins in the legend indicate the number of nucleotides in each bin. Statistical test results are reported in **Supplementary Fig. 9a**. **b** Predicted effects on the tail-length change for human variants. Shown is the PAL-AI–predicted difference in the tail-length change for each variant reported in the *All of Us* Research Program (exome callset v8) and that of the reference allele, plotted as a function of the variant position in its 3' UTR. Colors indicate allele frequencies. **c** Depletion in human 3' UTRs of alleles predicted to disrupt poly(A)-tail lengthening. Shown are relative frequencies of variants, grouped by predicted differences in tail-length change, among cohorts of variants with different allele frequencies reported in the *All of Us* Research Program (exome region v8). Only variants within the last 100 nt of 3' UTRs were included in this analysis. Binned *P* values (circles at the top) were from Fisher's exact tests performed for each cohort against singletons. **d** Same as (**c**), but for variants reported in gnomAD v4.1 dataset.

Supplementary Fig. 1

Supplementary Fig. 1: Additional data supporting the prediction of tail-length changes by multiple linear models and the architecture of PAL-AI.

a Prediction performance of the multiple linear regression models trained on different input regions of mRNA and configured with different lengths of *k*-mers; otherwise, as in **Fig. 1f. b** Top-ranked features as the number of *k*-mers from the best-performing multiple linear

regression models. Plotted are mean coefficients of the 20 top-ranked (by absolute value)
features as the number of *k*-mers from the model. Error bars indicate standard deviation of
coefficients of the 50 models from 10-fold cross-validation. **c** Top-ranked features as the
position metric of *k*-mers from the best-performing multiple linear regression models. As in
(**b**), except plotted are mean coefficients of the 20 top-ranked (by absolute value) features
as the position metric (sum of the inverse of the distance between each *k*-mer and the 3'
end) of *k*-mers from the model. **d** The architecture of PAL-AI for predicting tail-length
changes. **e** Parameters of the architecture of PAL-AI. Shown for each layer are the name
and parameters of the layer, and in boxes are the dimensions of the input and output
matrices. The dashed arrow indicates repeats of the convolutional module. Different
numbers of repeats were tested during hyperparameter searches.
732

Supplementary Fig. 2

Supplementary Fig. 2: Hyperparameter optimization reveals optimal configuration parameters for PAL-AI.

a Impact of discrete hyperparameters on PAL-AI performance. Shown are distributions of R_p values evaluating the correspondence between predicted and measured tail-length changes for different hyperparameter values in validation data. Numbers at the top indicate sample (trial) sizes for each condition. Box and whiskers indicate the 10th, 25th, 50th, 75th, and 90th percentiles. **b** Relationship between continuous hyperparameters and PAL-AI performance. Scatter plots show R_p values as a function of learning rate (left), L1 regularization (middle), and L2 regularization (right). Each point represents an individual model with specific hyperparameter values. **c** Learning curves showing training and validation loss. Mean squared error (MSE) is plotted against training epochs for both

training (red) and validation (blue) datasets. Results represent means from 50 models (5
repeats for each fold in 10-fold cross-validation). Shaded areas indicate standard error of
the mean. **d** Performance metrics during PAL-AI model training. Mean absolute error (MAE)
is plotted against training epochs for both training (red) and validation (blue) datasets,
analyzed as in (**c**). **e** Comparison of model architectures. Shown are distributions of R_p
values evaluating the correspondence between predicted and measured tail-length
changes for PAL-AI-s or ResNet implementations (with input sequence length of 2000 nt, 3'
UTR only). Ten-fold cross-validation of each model was repeated five times, generating 50
R_p values. Box and whiskers indicate the 10th, 25th, 50th, 75th, and 90th percentiles. *P*
value: one-sided t-test.
754

Supplementary Fig. 3

Supplementary Fig. 3: Additional data supporting sequence elements and contextual features learned by PAL-AI.

a As in **Fig. 2a**, except for consequences of gain of sequence motifs. **b** As in **Fig. 2b**, except for 8-mers most associated with the largest differences in predicted tail-length change between mutants in which 8-mers were gained and the wild-type. **c** As in **Fig. 2c**, except for top 6-mers associated with increased predicted tail-length change when gained in the in silico mutagenesis analysis. **d** Tail-length changes associated with each 3-mer within the 3' UTRs of the N60-PAS^{mos} library[18], comparing between 0 and 7 h post-progesterone treatment. Plotted for each 3-mer are differences in mean tail lengths observed for mRNAs with that 3-mer. Only variants containing one CPE (UUUUA) were included in this analysis. **e** Densities of the PAS and the CPE along the last 500 nt of frog mRNA 3' UTRs. The inset shows the results for the last 100 nt of the 3' UTR. **f** As in **Fig. 2e**, except for PAS variants. All insertions that would disrupt an AAUAAA were excluded. **g** Reduced tail-length change for PAS variants. Shown are the mean differences in predicted tail-length change when replacing the canonical PAS (AAUAAA) with the indicated PAS variant. Nucleotides that differ from the canonical PAS are colored in red. Only variants within the last 300 nt of 3' UTRs were included in this analysis. Error bars: standard error of the mean. **h** Tail-length changes associated with each 6-mer within the 3' UTRs of the CPE^{mos}-N60 library, comparing between 0 and 5 h post-progesterone treatment[18]. Plotted for each 6-mer are differences in mean tail lengths observed for mRNAs with that 6-mer. Variants containing any canonical PAS (AAUAAA or AUUAAA) were excluded from this analysis. **i** Impact of single-nucleotide substitutions on predicted tail-length changes for the *ccnb1.2.L* mRNA 3' UTR; otherwise, as in **Fig. 2g**. **j** Impact of single-nucleotide substitutions on predicted tail-length changes for the *ccnb2.L* mRNA 3' UTR; otherwise, as in **Fig. 2g**.

Supplementary Fig. 4

Supplementary Fig. 4. Additional data supporting the prediction of tail-length changes for synthetic mRNAs in frog oocytes.
a Performance of the PAL-AI model trained on endogenous frog mRNAs when tested on injected mRNA reporters. Each point represents a variant in the N60(LC)-PAS^{mos} library with at least 50 poly(A) tags in each of the two datasets, which were from oocytes 0 and 7 h post-progesterone treatment; otherwise, as in **Fig. 1c**. **b** Performance of a model trained from data generated from the injected N60(LC)-PAS^{mos} mRNA library and tested using 10-fold cross-validation. The 34,527 variants in the N60(LC)-PAS^{mos} library with at least 50 poly(A) tags in each of the two datasets, which were from oocytes 0 and 7 h post-progesterone treatment, were used in this analysis; otherwise, as in **Fig. 1c**. **c** Top 6-mers associated with decreased predicted tail-length change in the in silico mutagenesis analysis with a model trained with data generated from the injected N60(LC)-PAS^{mos} mRNA library. Otherwise, this panel is as in **Fig. 2c**. **d** As in (**c**), except for top 6-mers associated with increased predicted tail-length change when gained in the in silico mutagenesis analysis. **e** Positional effects for the CPE predicted by a model trained with data generated from the injected N60(LC)-PAS^{mos} mRNA library. Otherwise, this panel is as in **Fig. 2e**.

Supplementary Fig. 5

Supplementary Fig. 5: Additional experimental validation of predicted effects of
single-nucleotide substitutions.
a PAL-AI prediction of the effects of single-nucleotide substitutions of the *atp1a1.S* mRNA.
Otherwise, this panel is as in **Fig. 4d**. **b** As in (**a**), but for the *lima1.L* mRNA. **c** As in (**a**), but
for the *magoh.S* mRNA. **d** Impact of single-nucleotide substitutions on predicted and
measured tail-length changes for the *atp1a1.S* mRNA. Otherwise, this panel is as in **Fig. 4f**.
e As in (**d**), but for the *lima1.L* mRNA. **f** As in (**d**), but for the *magoh.S* mRNA. Arrows
indicate predicted and measured differences in tail-length changes that result from a C-to-
U substitution that creates a CPE that overlaps with the PAS.
810

Supplementary Fig. 6

Supplementary Fig. 6: Additional analyses supporting tail-length changes affected by
structural accessibility.
a Impact of single-nucleotide substitutions on predicted and measured tail-length changes
for the *aurkaip1.L* mRNA. Arrows indicate predicted and measured differences in tail-
length changes of either a C-to-U substitution (solid) that creates a CPE or two
substitutions (A-to-G at position –35 and C-to-U at position –14, dashed) that would
increase the length of a predicted stem (dashed rectangles) expected to occlude the PAS.
Otherwise, this panel is as in **Fig. 4f**. **b** PAL-AI prediction of the effects of single-nucleotide
substitutions of the *aurkaip1.L* mRNA. Otherwise, this panel is as in **Fig. 4d**. **c** EternaFold-
predicted maximum expected accuracy secondary structure of the last 100 nt of the
aurkaip1.L 3' UTR. The dashed rectangle indicates the stem of a hairpin, as highlighted in
(**a**). The triangles indicate positions of either a C-to-U substitution (solid) that creates a CPE
in a predicted loop region with a tail-length change correctly predicted by PAL-AI or two
substitutions (A-to-G at position –35 and C-to-U at position –14, hollow) that would
increase the length of a predicted stem expected to occlude the PAS. **d** As in (**a**), but for the
ccnb2.L mRNA. Arrows indicate predicted and measured differences in tail-length changes
of C-to-G substitutions at positions –54 and –16, both of which create the potential for a C–
G base pair that would increase the length of a predicted stem indicated with dashed
rectangles. **e** As in (**b**), but for the *ccnb2.L* mRNA. **f** As in (**c**), but for the *ccnb2.L* mRNA.
Shaded in blue are two CPEs and a PAS. The triangles indicate two positions where C-to-G
substitutions would extend the predicted stem indicated with the dashed rectangle.
833

Supplementary Fig. 7

a

b

Supplementary Fig. 7: Additional experimental validation of predicted effects of
single-nucleotide substitutions, exemplifying consequences of CPE-PAS overlap.
a PAL-AI prediction of the effects of single-nucleotide substitutions of the *ccnb1.2.L*
mRNA. Otherwise, this panel is as in **Fig. 4d**. **b** Impact of single-nucleotide substitutions on
predicted and measured tail-length changes for the *ccnb1.2.L* mRNA. The dashed
rectangle indicates a CPE that overlaps with a PAS. Otherwise, this panel is as in **Fig. 4f**.
840

Supplementary Fig. 8

 Supplementary Fig. 8: Additional data supporting the prediction of tail-length change
 for endogenous mRNAs in mouse and human oocytes.

a Performance of PAL-AI on mouse mRNAs. This panel is as in **Fig. 6a**, except that the
predictions were made with the PAL-AI-s model trained solely on frog oocyte endogenous
mRNAs. **b** Performance of PAL-AI on human mRNAs. This panel is as in **Fig. 6b**, except that
the predictions were made with the PAL-AI-s model trained solely on frog oocyte
endogenous mRNAs. **c** Performance of the PAL-AI-ms model on frog mRNAs. Otherwise,
this panel is as in **Fig. 6a**. **d** Performance of the PAL-AI-ms model on mouse mRNAs with
data generated from ONT in a published study[9]. This panel is as in Fig. 6a, except that the
measured tail-length changes during mouse GV-to-MII oocyte maturation were obtained
from ONT. **e** Correspondence between HiSeq and ONT measurements. Shown are tail-
length differences measured either by HiSeq (x axis) or ONT (y axis) during mouse GV-to-MII
oocyte maturation. Otherwise, this panel is as in **Fig. 1c**. **f** Genes predicted or shown to
have substantial mRNA tail-lengthening. This panel is as in **Fig. 6d**, except genes with PAL-
AI-predicted values in two of the three species are shown. **g** Genes predicted to have
substantial mRNA tail-lengthening. This panel is in (**f**), except genes with PAL-AI-predicted
values in all three species are shown.
858
859

Supplementary Fig. 9

 Supplementary Fig. 9. Additional data supporting negative selection of human variants
predicted to disrupt poly(A)-tail lengthening.
a Pairwise comparison of phyloP scores. Shown are binned *P* values from one-sided Mann–
Whitney U tests of phyloP scores between groups with indicated differences tail-length
changes predicted by PAL-AI, testing the alternative hypothesis that the mean phyloP
scores of the group indicated on the y axis is greater than that indicated on the x axis. **b**
Depletion among frequent alleles predicted to disrupt poly(A)-tail lengthening, when
excluding effects on PAS elements. This panel is as in **Fig. 7c**, except variants that
introduced or eliminated PAS elements were excluded. **c** Analysis of gnomAD data,
showing predicted effects on tail-length change for human variants. This panel is as in **Fig.**
7b, but for human variants in the gnomAD v4.1 dataset. **d** Analysis of gnomAD data,
showing depletion among frequent alleles predicted to disrupt poly(A)-tail lengthening,
when excluding effects on PAS elements. This panel is as in (**b**), but for human variants in
the gnomAD v4.1 dataset.

877 **Methods**
878 **Annotations of poly(A) sites and 3' UTRs**. Annotations of poly(A) sites and 3' UTRs of frog,
879 mouse, and human oocytes were obtained from a previous study[18].
880
Linear regression models for poly(A) tail-length changes. Poly(A) tail-length changes for
each isoform of frog oocyte endogenous mRNAs were calculated as the difference
between median tail lengths measured at 0 and 7 h post-progesterone treatment, requiring
at least 50 poly(A) tags at both time points, as reported previously[18]. Only 3' UTRs not
annotated as "uncertain" in previously curated annotations[18] were included.
K-mer features were defined as the positions and counts of all k-mers of in the
length of either 5–6 nt or 4–7 nt within the specified regions of the 3' UTR of each mRNA
isoform were used as features for the linear regression model. For isoforms with 3' UTRs
shorter than the length of the specified region, the entire 3' UTR was examined. Isoforms
with 3' UTRs shorter than 10 nt were excluded. Each 3' UTR was appended with the tri-
nucleotide sequence "AAA" before k-mer extraction. This sequence was treated as part of
the 3' UTR for both k-mer feature calculation and the specified region length (e.g., for a the
"last 1000 nt" region, the last 997 nt of the annotated 3' UTR were used before appending
"AAA").
The number of each k-mer N^k was counted with the *oligonucleotideFrequency*
function from the R package *Biostrings*[62]. A composite positional feature value P^k was
computed for each k-mer to summarize its positional distribution relative to the 3' end:

$$P^k = \begin{cases} \sum_i \dfrac{1}{\max{(d_i, 1)}}, & i \notin \emptyset \\ 0, & i \in \emptyset \end{cases}$$

899 where d_i is distance from the 3' end to the i-th occurrence of the k-mer. The linear
900 regression model predicting tail-length change L_i for isoform i was defined as:

$$L_i = c^0 + \sum_k c_N^k \cdot N_i^k + \sum_k c_P^k \cdot P_i^k$$

where c^0, c_N^k, and c_P^k are the model coefficients to be optimized. All features were scaled
using *RobustScaler* from the Python *scikit-learn* package[63]. Data were randomly stratified
into 10 folds using *StratifiedKFold*[63], maintaining the proportion of isoforms with 0, 1, 2, 3,
or ≥ 4 CPE (UUUUA) elements within their 3' UTRs.
The model was implemented using a single Dense layer in *Tensorflow*[64] and trained
with the Adam optimizer. Hyperparameters were tuned using the *Optuna*[65] framework. The
negative R^2 (between measured and predicted tail-length changes) for the validation data
was used as the objective to minimize for optimization. Final hyperparameters were:
learning_rate = 0.0025169, l1_regularizer = 0.000166, l2_regularizer = 0.0001097, epochs
=100, batch_size = 50, loss = 'mse', patience = 10. Model performance was evaluated in 10-
fold cross-validation. In this procedure, the 9-fold data were used for model training and
validation (further split into 90% for training and 10% for validation, stratified by CPE count)
while the remaining fold was used for testing the model performance. After each epoch,
the model was evaluated on the validation set. Training stopped early if validation loss did
not improve over 10 consecutive epochs, and the best model (based on the lowest

validation loss) was saved. This training/validation procedure was repeated 5 times,
accounting for differences in weight initializations, and the model with the highest R^2
(between measured and predicted tail-length changes) for the test data was selected to
predict the remaining fold. The procedure was rotated across all 10 folds. Final model
performance was assessed by concatenating predictions from all test folds and comparing
them with the measured tail-length changes.
923
PAL-AI for poly(A) tail-length changes. The following datasets were used to train or
validate different PAL-AI models: 1) the frog endogenous mRNA dataset: same as that used
in the linear regression model; 2) the N60(LC)-PAS*mos* library dataset: tail-length changes of
the N60(LC)-PAS*mos* library mRNAs injected into frog oocytes, comparing median tail
lengths measured at 0 and 7 h post-progesterone treatment during oocyte maturation,
requiring at least 50 poly(A) tags at both time points; 3) the single-nucleotide mutagenesis
library dataset: tail-length changes of the single-nucleotide mutagenesis library mRNAs
injected into frog oocytes, comparing median tail lengths measured at 0 and 7 h post-
progesterone treatment during oocyte maturation, requiring at least 50 poly(A) tags at both
time points; 4) the mouse endogenous mRNA dataset: tail-length changes of mouse
endogenous mRNAs, comparing between median tail lengths measured in GV and MII
oocytes, requiring at least 50 poly(A) tags in both replicate-merged datasets as described
previously[18]; 5) the human endogenous mRNA dataset: tail-length changes of human
endogenous mRNAs, comparing between median tail lengths measured in GV and MII
oocytes, requiring at least 50 poly(A) tags in both datasets as described previously[18].
To encode sequences as input for PAL-AI, RNA sequences from the specified
regions of each mRNA isoform's 3' UTR were appended at the 3' end with the tri-nucleotide
sequence "AAA". For isoforms with 3' UTRs shorter than the specified region, the entire 3'
UTR was used, and "N"s were padded at the 5' end. Isoforms whose AAA-appended 3' UTRs
were shorter than 10 nt were excluded. Sequences were one-hot encoded into four
channels, with "N" represented by a zero vector. When coding regions were included, they
were padded to the 5' end of the AAA-appended 3' UTRs. If the total length was still shorter
than the required input length, "N"s were padded at the 5' end. A fifth channel was added,
where a value of 1 marked 3'-UTR positions and 0 marked coding or padded regions. When
secondary structure information was included, *RNAfold*[35] was used to predict structure
from the AAA-appended 3' UTR. The unpairing probability of each nucleotide was added as
a sixth input channel. Nucleotides from the coding region and padded "N"s were assigned
a value of 0 in this sixth channel. The inputs for different final PAL-AI models were
summarized in Table 1.
The base PAL-AI model PAL-AI-s integrated a convolutional neural network with a
recurrent neural network, in an architecture similar to Saluki[34]. It consisted of four
modules: 1) Input block, which included an input layer, a 1D convolution, a normalization
layer, and an activation layer. 2) Convolutional block, which included repeated structure,
with each repeat comprising a 1D convolution layer, a normalization layer, an activation
layer, a 1D maxpooling layer, and a dropout layer. 3) Recurrent block, which included a
recurrent layer (GRU, BiGRU, LSTM, or BiLSTM), a normalization layer, and an activation
layer. 4) Dense block, which contains a dense layer, a dropout layer, a normalization layer,

an activation layer, and a final dense layer with a single output for the tail-length change. All
convolution layers used the same number of filters and kernel size. The convolutional
layers, the recurrent layer, and the first dense layer used the same kernel initializer and
regularizer. All dropout layers had the same dropout rate. Layer normalization was used in
all blocks unless Leaky ReLU was the activation function, in which case batch
normalization was used in the dense block. All activation layers used the same function—
either SELU, SiLU (Swish), or Leaky ReLU. If SELU was used, the Lecun normal initializer
and Alpha Dropout were applied. Otherwise, the HE normal initializer was used.
The PAL-AI-s model was implemented using the Python *TensorFlow* package[64] and
trained with the Adam optimizer. Loss functions used were either the mean squared error
(MSE) or the mean absolute error (MAE). The hyperparameter tuning, cross-validation
strategy, and model performance evaluation followed the same procedures as the linear
regression model.
For the N60(LC)-PAS^{mos} library, PAL-AI-s used the same architecture. Sequences in
the 60-nt random region and constant region were appended with "AAA" and one-hot
encoded. Training and evaluation were performed as described for endogenous mRNAs.
The PAL-AI-m and PAL-AI-ms models shared the same base architecture as PAL-AI-
s, with three key modifications. First, an additional input feature was included to indicate
the dataset group for each mRNA sequence. Second, rather than a single output, the final
dense layer outputs multiple values, with one per dataset group. Third, a custom gathering
layer was added after the dense layer to select the appropriate group-specific output
based on the input dataset label. For the PAL-AI-m model, the frog endogenous mRNA
dataset (output head 1) and the N60(LC)-PAS^{mos} library dataset (output head 2) were used.
For the PAL-AI-ms model, the frog endogenous mRNA dataset (output head 1), the human
endogenous mRNA dataset (output head 2), and the mouse endogenous mRNA dataset
(output head 3) were used. All input sequences were encoded as described for PAL-AI-s.
Training and evaluation followed the same procedure as PAL-AI-s, with the following
differences to accommodate multiple groups. First, when splitting the training and
validation set, the validation set contributed 10% of its data for the group with the smallest
dataset, and validation sets from all other groups were downsampled to match this
number. Second, for training, data from larger groups were similarly downsampled per
epoch to match the smallest group, and batches were alternated across groups. Third, the
average of negative R^2 values (between measured and predicted tail-length changes)
across validation groups was used as the objective to minimize during hyperparameter
optimization.
A ResNet-based variant of PAL-AI was also tested, replacing the second module of
PAL-AI-s with a dilated convolutional ResNet architecture[36] comprising seven groups with
dilation rates of 1, 2, 4, 8, 4, 2, and 1. Each group contained four blocks, each consisting of
a dilated convolution layer, layer normalization, an activation layer, a second dilated
convolution layer, another layer normalization, a residual skip connection, and a final
activation layer.
For all PAL-AI models, an optional input representing the initial tail length could be
provided. When included, this value was concatenated to the input of the final dense layer.

Although incorporating this feature improved performance for some datasets, it reduced
generalizability, as initial tail-length measurements were not available for all mRNAs.
For all predictions made with PAL-AI, input sequences were encoded using the
same scheme as during model training. To compare model performance on the single-
nucleotide mutagenesis library, predictions were generated using either PAL-AI-s
(endogenous mRNA-trained) or PAL-AI-m (endogenous mRNA-specific output head 1).
Unless otherwise specified, final predictions were computed as the average output across
the best-performing models from each fold of the 10-fold cross-validation.
Hyperparameters for each PAL-AI model were optimized using *Optuna*[65], with final
values summarized in Table 1.
1014
1015 **Table 1: Parameters used in each PAL-AI model.**

	PAL-AI-s (frog endogenous mRNAs)	PAL-AI-s [N60(LC)-PAS^{mos} library]	PAL-AI-m	PAL-AI-ms	PAL-AI (ResNet)
Length of input sequences (nt)	2000	84	2000	2000	2000
Use of coding sequences	Yes	No	Yes	Yes	No
Use of group label input	No	No	Yes	Yes	No
Number of filters for 1D convolution layers	64	96	64	64	96
Kernel size for 1D convolution layers	5	4	4	5	4
Number of repeats of the convolutional block	1	1	1	2	Not applicable
Number of units of the dense layer	64	96	128	128	32
Size of the maxpooling layer	2	2	2	2	Not applicable
Recurrent layer	GRU	GRU	GRU	GRU	GRU
Number of units of the recurrent layer	96	128	128	128	64
Activation function	Leaky ReLU	Leaky ReLU	Leaky ReLU	Leaky ReLU	Leaky ReLU

Kernel regularizer L1	0.001	0.000001798	0.0000628	0.0000043	0.006823
Kernel regularizer L2	0.0045	0.001257	0.00019765	0.00344	0.000006507
Dropout rate	0.3	0.2	0.2	0.2	0.2
Learning rate	0.003	0.00030457	0.001934	0.00068	0.0005537
Loss function	MSE	MSE	MSE	MAE	MAE
Batch size	128	64	128	32	64
Epochs	40	100	100	100	100
Trainable parameters	75,713	138,689	109,826	135,043	2,116,129
Non-trainable parameters	128	192	256	256	10,816

1016

In silico mutagenesis (ISM). For frog and human endogenous mRNAs, all 3' UTRs were
analyzed except those previously annotated as "uncertain"[18]. For the N60(LC)-PAS[mos]
library, all 3' UTRs in the curated reference were used. Input sequences were constructed
and encoded as described for PAL-AI. Tail-length changes of the wild-type sequences were
predicted using PAL-AI. For ISM, each nucleotide within 3' UTRs was individually
substituted with each of the three possible alternative nucleotides (the fifth channel
annotating the 3' UTR was not altered), and the tail-length change of each mutant was
predicted with corresponding PAL-AI models: PAL-AI-s (frog endogenous mRNA-trained) for
frog endogenous mRNAs, PAL-AI-s (N60(LC)-PAS[mos] library-trained) for N60(LC)-PAS[mos]
library mRNAs, and PAL-AI-ms (human-specific output head 2) for human endogenous
mRNAs. The difference between the predicted tail-length change of the wild-type and each
mutant was then calculated.
To examine the tail-length effect of losing or gaining a k-mer, differences between
the predicted tail-length change of the wild-type and those of all single-nucleotide
substitution mutants that disrupted the original k-mer or introduced a new k-mer were
averaged across all instances of that k-mer within the last 300 nt of 3' UTRs from all mRNA
isoforms.
In some cases, k-mer-associated differences in predicted tail-length changes were
examined in an iterative exclusion process. Only mutations within the last 300 nt of 3' UTRs
were considered. In each round of the iteration, mutations that disrupted (for k-mer loss) or
introduced (for k-mer gain) any k-mer already present in an exclusion list (initially empty)
were ignored. The average difference in predicted tail-length change between the wild-type
and the mutant was then calculated for all k-mers that were not in the exclusion list. The k-
mer with the largest tail-length increase (for k-mer gain) or decrease (for k-mer loss) was
recorded and added to the exclusion list before starting the next round. The iteration
continued until no k-mer showed a statistically significant increase (or decrease) in
predicted tail-length change (Welsh's t-test; Bonferroni-adjusted $P > 0.05$).
Motif logos associated with predicted tail-length changes were generated as
previously described[18].
1046

Motif insertion analysis. Filtering and encoding of input sequences were performed as in
the ISM analysis. To insert a motif, the sequence of the same length at a specified position
within the 3' UTR of an mRNA was replaced with the motif. The fifth channel annotating the
3' UTR was left unchanged. PAL-AI was then used to predict the tail-length changes of both
the wild-type and motif-inserted mutant. The difference between these predictions was
used as the insertion outcome. When analyzing the relative positioning of the PAS and CPE
elements, only mRNAs whose 3' UTRs contained at least a PAS within the last 100 nt were
included. For these mRNAs, the relative distance was calculated between each CPE and
the PAS closest to the 3' end.

1056

Reanalysis of reporter libraries for *k*-mer-associated tail-length changes. Poly(A) tail-
length measurements of the frog oocyte-injected N60-PAS^{mos} library at 0 and 7 h post-
progesterone treatment and CPE^{mos}-N60 library at 0 and 5 h post-progesterone treatment
were obtained from a previous study[18]. For the N60-PAS^{mos} library, the analysis was
restricted to reporter variants containing exactly one instance of the UUUUA motif within
the 3' UTR. For the CPE^{mos}-N60 library, the analysis was restricted to reporter variants not
containing any AAUAAA or AUUAAA motifs within the 3' UTR. For each *k*-mer, all variants
containing that *k*-mer were identified, and the mean tail length across those variants was
computed. To assess the positional effect of specific *k*-mers, only variants in which the *k*-
mer appeared at a defined position were included in the calculation.

1067

Preparation of the N60(LC)-PAS^{mos} library. The DNA template for the N60(LC)-PAS^{mos}
library was generated by PCR using primers KXU024 and KXU068, and 2 amol of the N60-
PAS^{mos} DNA template used previously[18]. The reaction (100-µl total volume) was performed
with the KAPA HiFi HotStart Kit (Roche, KK2502). PCR products were gel-purified using
agarose gels (Lonza, 50004) and the GeneJet Gel Extraction Kit (Thermo Fisher, K0692). The
resulting mRNA library was in vitro transcribed and capped, followed by removal of the 3'-
end cyclic phosphate as previously described[18].

1075

Preparation of the single-nucleotide mutagenesis library. The oligo pool was
synthesized as SurePrint HiFi Oligo (Agilent Technologies; Supplementary Table 2). Each
oligo contained a variable region derived from the last 100 nt of the 3' UTR of one of ten frog
oocyte mRNAs (*atp1a1.S*, *ccnb1.2.L*, *ccnb2.L*, *lima1.L*, *mad2l1.L*, *magoh.S*, *mos.L*, *tpx2.L*,
aurkaip1.L, and *dbf4.L*). At each position, every nucleotide was mutated to the three
alternative nucleotides, yielding 3010 sequences (10 wild-type and 3000 mutants). A 35-nt
poly(A) fragment was appended to the 3' end and constant sequences (24 nt at the 5' end
and 20 nt at the 3' end) were added for PCR amplification. Both forward and reverse strands
were included in the pool.
The DNA template for in vitro transcription was assembled by overlapping PCR using
the KAPA HiFi HotStart Kit. Three fragnments were joined sequentially. Fragment 1 (F1) was
amplified from the N60-PAS^{mos} template[18] with primers KXU024 and KXU236. Fragment 2
(F2) was amplified from the oligo pool with primers KXS067 and KXU237. Fragment 3 (F3)
was amplified from the plasmid C071[4] with primers KXU110 and KXU068. F1 and F2 were
first joined, and the resulting product was joined with F3 in a 100-µl PCR reaction

containing 10 pmol of each fragment, 2 μl KAPA HiFi enzyme, 3 μl 10 mM dNTP, and 20 μl 5×
KAPA HiFi buffer for 15 cycles, with an annealing temperature of 66°C. PCR products were
purified using agarose gels and the GeneJet Gel Extraction Kit. The mRNA library was then
in vitro transcribed and capped, followed by removal of the 3'-end cyclic phosphate as
described previously[18].
1096
Preparation of the reporter mRNA poly(A) tail-length standard mix. Each of the four
poly(A) tail-length standards was constructed from two fragments (F1 and F2) joined by
overlapping PCR. Both fragments were amplified from plasmid C071 using the KAPA HiFi
HostStart Kit and the following primer sets: KXU024 and one of KXU290, KXU291, KXU292,
or KXU293 (for F1); KXU068 and one of KXUm020, KXUm021, KXUm022, or KXUm023 (for
F2, yielding tail lengths of 10, 50, 90, or 130 nt). Overlapping PCR was performed in a 50-μl
reaction containing 100 fmol of each fragment, 1 μl KAPA HiFi enzyme, 1.5 μl 10 mM dNTPs,
and 10 μl 5× KAPA HiFi buffer. After 5 cycles (annealing temperature 63°C), primers KXU024
and KXU068 were added to a final concentration of 300 nM along with 1 μl KAPA HiFi
enzyme, 1.5 μl 10 mM dNTP, and 10 μl 5× KAPA HiFi buffer to bring the final volume to 100
μl. Twenty-four additional PCR cycles were performed. PCR products were purified using
agarose gels and the GeneJet Gel Extraction Kit. Each RNA was in vitro transcribed and
capped, followed by removal of the 3'-end cyclic phosphate as described previously[18]. The
four mRNA standards were mixed at equimolar concentrations. Each contained a unique 4-
nt barcode in the 3' UTR, introduced by KXU290, KXU291, KXU292, or KXU293. Sequences
of the four tail-length standards are provided in Supplementary Table 3.
1113
Reporter mRNA library injection and sample collection. Frog oocytes were obtained
from Xenopus1 (12005). Healthy Stage V–VI oocytes were hand-picked and transferred to
OR-2 buffer (5 mM HEPES pH 7.6, 82.5 mM NaCl, 2.5 mM KCl, 1 mM MgCl$_2$, 1 mM CaCl$_2$, 1
mM Na$_2$HPO$_4$) supplemented with 100 μg/ml Gentamicin (Thermo Fisher, 15750060), and
incubated at 18°C overnight (> 16 hr) for recovery. Injections were performed at 23°C using
a PLI-100 Plus Pico-Injector. For both the N60(LC)-PAS^{mos} library and the single-nucleotide
mutagenesis library, 4 nl of mRNA (0.1 pmol/μl) was injected per oocyte.
Frog oocytes were matured in vitro in OR-2 buffer supplemented with 10 μg/ml
progesterone (Millipore Sigma, P0130), diluted from a 10 mg/ml ethanol-disolved stock.
Maturation timing varied across batches: in most cases, 50% of oocytes underwent
germinal vesicle breakdown (GVBD; indicated by a white spot on the animal pole) within 3–
5 hr, and 100% reached GVBD by 7 h post-progesterone addition.
At indicated time points post-progesterone treatment, groups of 100 frog oocytes
were collected. After removing OR-2 buffer, oocytes were washed three times with ice-cold
buffer RL (20 mM HEPES pH 7.5, 100 mM KCl, 5 mM MgCl$_2$, 1% Triton X-100, 100 μg/ml
cycloheximide, cOmplete protease inhibitor cocktail [MilliporeSigma, 11836170001, 1
tablet/10 ml], 10 μl/oocyte). Following the washes, buffer was completely removed and
oocytes were lysed in buffer RL (10 μl per oocyte) supplemented with 200 U/ml
SUPERase•In (10 μl/oocyte) by vigorous shaking and pipetting. Lysates were clarified by
centrifugation at 5,000 g for 10 min at 4°C. Supernatants were mixed with 3 volumes of
TRIzol LS (Thermo Fisher, 10296010).

The 0 hr post-progesterone sample was prepared by collecting the supernatant from

lysed, untreated oocytes and mixing it with the uninjected mRNA library prior to TRIzol LS

addition. All TRIzol-LS RNA extractions were performed using Phasemaker tubes (Thermo

Fisher, A33248) following the manufacturer's instructions. Total RNA was resuspended in

40 µl water (0.4 µl/oocyte).

1140

Library preparation and sequencing of the injected mRNA libraries. Total RNA (10 µl,

~30 µg) from injected oocytes was combined with 10 amol of reporter mRNA poly(A)-tail-

length standard mix. To this, 8 pmol each of oligos KXSH009 and KXSH010, and 2× SSC (0.3

M NaCl, 30 mM sodium citrate pH 7.0), were added in a final volume of 50 µl. The RNA-oligo

mixture was incubated at 70°C for 5 min and then cooled to 23°C at a rate of 0.1°C/sec for

annealing. The annealed RNA was incubated with 40 µl MyOne Streptavidin C1 beads

(Thermo Fisher, 65002) for 20 min at 23°C on a thermal mixer (15 sec on and 1 min 45 sec

off). The supernatant was separated from the beads with a magnetic rack and removed.

Beads were washed twice with 300 µl 1× B&W buffer (5 mM Tris-HCl pH 7.5, 0.5 mM EDTA,

1 M NaCl) and once with 300 µl 2× SSC. RNA was eluted from beads with two sequential

incubations: first in 100 µl 10 mM HEPES pH 7.5 at 65°C for 3 min, then in 100 µl water at

65°C for 3 min. Eluates were combined, ethanol-precipitated, and resuspended in 6.5 µl

water.

Anti-sense oligo-enriched RNA was ligated to a pre-adenylated 3' adapter (KXS347a)

in a 10 µl reaction containing 5 µM adapter, 50 mM HEPES pH 7.5, 10 mM MgCl$_2$, 10 mM

dithiothreitol, and 1 U/µl T4 RNA ligase 1 (NEB, M0204S). Ligation was carried out at 23°C

for 150 min. Ligated RNA was extracted with phenol/chloroform, ethanol-precipitated, and

resuspended in 11.4 µl water. The ligated RNA was mixed with 0.6 µl 100 µM primer KXS348

and 1 µl 10 mM dNTPs in a total volume of 13 µl, incubated at 65°C for 5 min, and cooled on

ice for 1 min. Reverse transcription was performed in a 20 µl reaction using SuperScript IV

(ThermoFisher, 18090050), with 1× SSIV Buffer, 5 mM dithiothreitol, 1 U/µl SUPERase•In,

and 200 U enzyme, incubated at 55°C for 15 min. After reverse transcription, RNA was then

hydrolyzed with 3.3 µl 1 M NaOH at 90°C for 10 min, followed by neutralization with 24 µl 1

M HEPES pH 7.5. The resulting cDNA was ethanol-precipitated, resuspended in water, and

amplified in a 50 µl PCR reaction using primer KXS037 and a barcoded primer (KXS057)

targeting a constant region of the library. Amplification was performed using the KAPA HiFi

HotStart Kit following the manufacturer's suggested protocol for 11~14 cycles. The final

PCR-amplified product was cleaned up twice with AMPure XP beads (Beckman Coulter,

A63881) at a 1.2× beads-to-sample ratio.

Sequencing was performed on an Illumina HiSeq 2500, with a custom run of two

reads: a 281-cycle first read using primer KXS067, and a 10-cycle second read using a

standard TruSeq sequencing primer. To generate a reference, the uninjected 0 hr N60(LC)-

PAS^{mos} library sample was also sequenced on AVITI using the same primers, with custom

read lengths of 299 and 10 cycles.

1175

Variant and tail-length analysis for the injected mRNA libraries. To build a reference

sequence for the N60(LC)-PAS^{mos} library, read 1 from the 0 hr post-progesterone

(uninjected) sample sequenced on AVITI was trimmed to 81 nt from the 5' end. Trimmed

reads were screened for the presence of a known constant region from the 3' UTR of poly(A)

tail-length standards (ACCAGCCTCAAGAACACCCGAATGG). A maximum of seven

mismatches was allowed within the first 25 nt of this sequence, permitting positional

offsets from –3 to +3 nt. Next, the remaining reads were examined for the expected

constant sequence AATAAAGAAATTGATTTGTCT at positions 61–81, again allowing offsets

from –3 and +3 nt. Reads containing a 21-nt segment with no more than six mismatches to

the constant sequence at any allowed position were retained. For reads with a non-zero

offset, alignment was corrected to zero: if the offset was negative, "N" bases were

prepended to the read; if positive, the corresponding number of bases was trimmed from

the start of the read. These modified reads were then clustered using *UMICollapse*[66], and

clusters with more than 100 supporting reads were retained. The consensus (primary)

sequence from each cluster was used to construct the reference sequence for the

N60(LC)-PAS*mos* library.

During the HiSeq 2500 run for both the N60(LC)-PAS*mos* and single-nucleotide

mutagenesis libraries, a sequencing artifact occurred at cycle 26 of read 1, resulting in a

repeated base call at cycle 26 and 27. Consequently, base 27 was removed from all reads

prior to downstream analysis.

For the N60(LC)-PAS*mos* libraries, read 1 sequences from the HiSeq run were aligned

to the reference using *STAR* (v2.7.1)[67] with the following parameters '--

outFilterMultimapNmax 1 --alignEndsType Local –clip3pNbases 199 --outSAMattributes All

--outSAMtype BAM SortedByCoordinate'. Poly(A)-tail lengths for mapped reads were

determined as previously described for the N60-PAS*mos* libraries[18].

For the single-nucleotide mutagenesis libraries, read 1 sequences from the HiSeq

run were aligned to the reference consisting of the last 100 nt of the 3' UTR from ten

selected frog oocyte mRNAs (wild-type sequences) using *STAR* (v2.7.1)[67] with these

parameters '--outFilterMultimapNmax 1 --outFilterMismatchNmax 1 --alignEndsType

EndToEnd –clip3pNbases 180 --outSAMattributes All --outSAMtype BAM

SortedByCoordinate'. Mutations were classified using the CIGAR string in the resulting BAM

file. Poly(A)-tail lengths for these mapped reads were determined as for the N60(LC)-PAS*mos*

libraries, with the tail start position defined as position 101 of read 1. Only wild-type and

single-nucleotide substitution variants were retained. For each variant, reads sharing the

same read 2 sequence (serving as a unique molecular identifier, UMI) were grouped and

collapsed. The median tail length of each collapsed was used as the representative

poly(A)-tail length.

Reference sequences of both the N60(LC)-PAS*mos* and single-nucleotide

mutagenesis libraries are provided in Supplementary Table 3.

1215

1216   **RNA secondary structure prediction.** RNA secondary structures for the 3' UTR sequences

1217   in the single-nucleotide mutagenesis library were predicted using *EternaFold* (v1.3.1)[46] with

1218   the parameters "--params parameters/EternaFoldParams.v1". The resulting dot-bracket

1219   notations were visualized using *RNAcanvas*[68].

1220

Analysis of mouse oocyte data sequenced by Nanopore. Poly(A)-tail lengths of mouse

oocyte mRNAs obtained via Oxford Nanopore Technologies (ONT) sequencing were

retrieved from a previous study[9]. Tail-length changes were calculated by comparing the

median tail lengths between GV and MII oocytes, requiring at least 50 poly(A) reads in

datasets from both stages. As ONT-reported tail-length measurements were reported per

gene rather than per mRNA isoform, comparisons were made with the primary isoform for

each gene, defined based on Hi-Seq data[18] or predicted using PAL-AI-ms (mouse-specific

output head 3). Analyses were restricted to genes whose primary isoform represented

more than 90% of expression in GV oocytes.

1230

Prediction of poly(A) tail-length changes of reporter mRNAs during mouse oocyte

maturation. To evaluate the relationship between predicted tail-length changes and

experimentally measured translational efficiency changes of reporter mRNAs injected into

mouse oocytes, source data were obtained from a previous study (see Fig. 4g of the original

publication)[6]. Translational efficiency change was calculated as the difference between the

mean normalized fluorescence signals in GV versus MII oocytes (24 hr). Reporter mRNA 3'

UTR sequences were inferred from the cloning and mutagenesis primers used in this study.

Tail-length changes for these sequences were predicted using PAL-AI-ms (mouse-specific

output head 3).

1240

Cross-species analysis of poly(A) tail-length and translational efficiency changes.

Gene orthology tables (frog-to-human from Xenbase, mouse-to-human from Ensembl)

were used to map homologous genes. Measured tail-length changes were compared

between GV and MII oocytes in humans and mice, and between 0 and 7 h post-

progesterone treatment in frog oocytes. For mRNA isoforms with unique 3' ends in one of

the three species, predicted tail-length changes from PAL-AI-ms were used if experimental

data were unavailable. For genes with multiple mRNA isoforms, the dominant isoform (>

50% by tag count in GV oocytes of humans or mice, or in 0 hr frog oocytes) was selected. If

a human gene had multiple homologs in frog or mouse, the average tail-length change

across all the homologs was used.

Translational efficiency data derived from ribosome profiling and mRNA-seq were

reprocessed[18] and compared between GV and MII oocytes in humans[7] and mice[6].

For gene ontology (GO) analysis, genes showing substantial tail lengthening (≥ 15 nt,

by measurement or prediction) during oocyte maturation in all three species were selected.

These genes were ranked by the smallest tail-length change across three species, and the

ranked list was submitted to the *gost* function in the R package *gprofiler2*[69] with the

parameters "ordered_query = TRUE, user_threshold = 0.05, correction_method = "g_SCS".

The top eight non-overlapping GO:Biological Process categories (not from the same parent

term and ranked by Benjamini-Hochberg-adjusted *P* value) were reported.

1260

Evolutionary constraint associated with tail-length changes. PhyloP scores from

multiple sequence alignments of 99 vertebrate genomes to the human genome were

obtained from the UCSC Genome Browser[49]. For each nucleotide within the last 100 nt of

human 3' UTRs with an available phyloP score, the mean PAL-AI-predicted differences in

tail-length changes when this nucleotide was substituted with each of the three alternative

nucleotides (from ISM analysis) were calculated.

1267
Prediction of poly(A) tail-length changes for human genetic variants from the All of Us
Research Program and gnomAD. MRNA 3' UTR isoforms shorter than 10 nt were excluded
from analysis. To maximize variant coverage despite incomplete transcript annotations, the
"uncertain" 3' UTRs in the annotations were retained if their 3' ends were supported by
entries in the PolyA_DB_v3.2 database[70].
For the data reported in the All of Us Research Program[10], all single-nucleotide
polymorphism and indel variants from short-read whole-genome sequencing callsets
encompassing human exome regions (version 8) were intersected with human oocyte 3'
UTR genomic coordinates. Variants overlapping these regions were selected. For each
variant, the associated 3' UTR sequence was reconstructed by substituting the reference
allele with the alternative allele, followed by appending three adenines (AAA) at the 3' end.
Tail-length changes were predicted using PAL-AI-ms (human-specific output head 2). The
mutational outcome was defined as the difference in predicted tail-length change between
the variant and corresponding wild-type (reference) sequence. In cases where a variant
overlapped multiple 3' UTRs, the one with the largest absolute predicted difference in tail-
length change was used for downstream analysis.
For the data reported in gnomAD[11], genomic variants from v4.1 (covering 76,215
individuals) were downloaded from gnomad.broadinstitute.org. Variants were included if
they passed all quality filters (indicated by "PASS" in the "FILTER" column in the VCF file)
and intersected annotated 3' UTR genomic locations. Tail-length changes and mutational
outcomes were computed using the same approach as for the All of Us Research Program.
1289
1290 **Quantification and statistical analysis**
1291 Graphs were generated and statistical analyses were performed using R[71]. Statistical
1292 parameters, including the value of n, statistical test, and statistical significance (*P* value)
1293 are reported in the figures or their legends. No statistical methods were used to pre-
1294 determine sample size.
1295

Data availability

All standard sequencing data are available in the Gene Expression Omnibus under the
accession number GSE280422. Raw intensity data for reporter mRNA tail-length
sequencing cannot be deposited in public databases due to their large sizes and are
available upon request.
Oligo sequences used in this study are listed in Supplementary Table 1. Sequences
of the oligo library used for the single-nucleotide mutagenesis library are listed in
Supplementary Table 2. Sequences of the N60(LC)-PAS*mos* library, the single-nucleotide
mutagenesis library, and the tail-length standards are listed in Supplementary Table 3.
Other publicly available data analyzed in this study are indicated in the relevant
sections of **Methods**.
1307

Code availability

1309   The code for PAL-AI is written in Python 3.8 and available at
1310   https://github.com/coffeebond/PAL-AI. Reporter mRNA tail-length sequencing data
1311   analyses were performed using a custom script written in Python 2.7 and available at
1312   https://github.com/coffeebond/MPRA_tail_seq. The codes used to generate the figures are
1313   available at https://github.com/coffeebond/PAL-AI_paper.
1314
1315

References

1. Eichhorn, S. W. *et al*. mRNA poly(A)-tail changes specified by deadenylation broadly reshape translation in Drosophila oocytes and early embryos. *eLife* **5**, e16955 (2016).

2. Lim, J., Lee, M., Son, A., Chang, H. & Kim, V. N. mTAIL-seq reveals dynamic poly(A) tail regulation in oocyte-to-embryo development. *Genes & Development* **30**, 1671–1682 (2016).

3. Yang, F., Wang, W., Cetinbas, M., Sadreyev, R. I. & Blower, M. D. Genome-wide analysis identifies cis-acting elements regulating mRNA polyadenylation and translation during vertebrate oocyte maturation. *RNA (New York, N.Y.)* **26**, 324–344 (2020).

4. Xiang, K. & Bartel, D. P. The molecular basis of coupling between poly(A)-tail length and translational efficiency. *eLife* **10**, e66493 (2021).

5. Liu, Y. *et al*. Poly(A) tail length is a major regulator of maternal gene expression during the mammalian oocyte-to-embryo transition. *bioRxiv* 2021.08.29.458052 (2021) doi:10.1101/2021.08.29.458052.

6. Xiong, Z. *et al*. Ultrasensitive Ribo-seq reveals translational landscapes during mammalian oocyte-to-embryo transition and pre-implantation development. *Nat Cell Biol* 1–13 (2022) doi:10.1038/s41556-022-00928-6.

7. Zou, Z. *et al*. Translatome and transcriptome co-profiling reveals a role of TPRXs in human zygotic genome activation. *Science* **378**, abo7923 (2022).

8. Liu, Y. *et al*. Remodeling of maternal mRNA through poly(A) tail orchestrates human oocyte-to-embryo transition. *Nat Struct Mol Biol* 1–16 (2023) doi:10.1038/s41594-022-00908-2.

9. Lee, K., Cho, K., Morey, R. & Cook-Andersen, H. An extended wave of global mRNA deadenylation sets up a switch in translation regulation across the mammalian oocyte-to-embryo transition. *Cell Rep*. **43**, 113710–113710 (2024).

10. Investigators, T. A. of U. R. P. G. *et al*. Genomic data in the All of Us Research Program. *Nature* 1–7 (2024) doi:10.1038/s41586-023-06957-x.

11. Chen, S. *et al*. A genomic mutational constraint map using variation in 76,156 human genomes. *Nature* **625**, 92–100 (2024).

12. Richter, J. D. & Lasko, P. Translational control in oocyte development. *Cold Spring Harbor Perspectives in Biology* **3**, a002758 (2011).

13. Teixeira, F. K. & Lehmann, R. Translational Control during Developmental Transitions. *Csh Perspect Biol* **11**, a032987 (2019).

14. Conti, M. & Kunitomi, C. A genome-wide perspective of the maternal mRNA translation program during oocyte development. *Semin Cell Dev Biol* (2023) doi:10.1016/j.semcdb.2023.03.003.

15. Sheets, M. D., Fox, C. A., Hunt, T., Woude, G. V. & Wickens, M. The 3'-untranslated regions of c-mos and cyclin mRNAs stimulate translation by regulating cytoplasmic polyadenylation. *Genes & Development* **8**, 926–938 (1994).

16. Moor, C. H. de & Richter, J. D. The Mos pathway regulates cytoplasmic polyadenylation in Xenopus oocytes. *Molecular and Cellular Biology* **17**, 6419–6426 (1997).

17. Subtelny, A. O., Eichhorn, S. W., Chen, G. R., Sive, H. & Bartel, D. P. Poly(A)-tail profiling reveals an embryonic switch in translational control. *Nature* **508**, 66–71 (2014).

18. Xiang, K., Ly, J. & Bartel, D. P. Control of poly(A)-tail length and translation in vertebrate
oocytes and early embryos. *Dev. Cell* **59**, 1058-1074.e11 (2024).
19. McGrew, L. L., Dworkin-Rastl, E., Dworkin, M. B. & Richter, J. D. Poly(A) elongation
during Xenopus oocyte maturation is required for translational recruitment and is
mediated by a short sequence element. *Gene Dev* **3**, 803–815 (1989).
20. Radford, H. E., Meijer, H. A. & Moor, C. H. de. Translational control by cytoplasmic
polyadenylation in Xenopus oocytes. *Biochimica et Biophysica Acta (BBA) - Gene
Regulatory Mechanisms* **1779**, 217–229 (2008).
21. Aoki, F., Hara, K. T. & Schultz, R. M. Acquisition of transcriptional competence in the 1-
cell mouse embryo: Requirement for recruitment of maternal mRNAs. *Mol. Reprod. Dev.*
64, 270–274 (2003).
22. Meneau, F., Dupré, A., Jessus, C. & Daldello, E. M. Translational Control of Xenopus
Oocyte Meiosis: Toward the Genomic Era. *Cells 2020, Vol. 9, Page 1502* **9**, 1502 (2020).
23. Yatsenko, S. A. & Rajkovic, A. Genetics of human female infertility. *Biol. Reprod.* **101**,
549–566 (2019).
24. Zheng, W. *et al.* Homozygous Mutations in BTG4 Cause Zygotic Cleavage Failure and
Female Infertility. *Am. J. Hum. Genet.* **107**, 24–33 (2020).
25. Liu, R. *et al.* A novel homozygous missense variant in BTG4 causes zygotic cleavage
failure and female infertility. *J. Assist. Reprod. Genet.* **38**, 3261–3266 (2021).
26. Varnum, S. M. & Wormington, W. M. Deadenylation of maternal mRNAs during Xenopus
oocyte maturation does not require specific cis-sequences: a default mechanism for
translational control. *Gene Dev* **4**, 2278–2286 (1990).
27. Fox, C. A. & Wickens, M. Poly(A) removal during oocyte maturation: a default reaction
selectively prevented by specific sequences in the 3' UTR of certain maternal mRNAs.
Genes & Development **4**, 2287–2298 (1990).
28. Hake, L. E. & Richter, J. D. CPEB is a specificity factor that mediates cytoplasmic
polyadenylation during Xenopus oocyte maturation. *Cell* **79**, 617–627 (1994).
29. Dickson, K. S., Bilger, A., Ballantyne, S. & Wickens, M. P. The Cleavage and
Polyadenylation Specificity Factor in Xenopus laevis Oocytes Is a Cytoplasmic Factor
Involved in Regulated Polyadenylation. *Mol Cell Biol* **19**, 5707–5717 (1999).
30. Charlesworth, A., Meijer, H. A. & Moor, C. H. de. Specificity factors in cytoplasmic
polyadenylation. *Wiley Interdiscip Rev Rna* **4**, 437–461 (2013).
31. Ivshina, M., Lasko, P. & Richter, J. D. Cytoplasmic Polyadenylation Element Binding
Proteins in Development, Health, and Disease. *Annual Review of Cell and Developmental
Biology* **30**, 393–415 (2014).
32. Reyes, J. M. & Ross, P. J. Cytoplasmic polyadenylation in mammalian oocyte
maturation. *Wiley Interdisciplinary Reviews: RNA* **7**, 71–89 (2016).
33. Weill, L., Belloc, E., Castellazzi, C. L. & Méndez, R. Musashi 1 regulates the timing and
extent of meiotic mRNA translational activation by promoting the use of specific CPEs.
Nature Structural & Molecular Biology **24**, 672–681 (2017).
34. Agarwal, V. & Kelley, D. R. The genetic and biochemical determinants of mRNA
degradation rates in mammals. *Genome Biol* **23**, 245 (2022).
35. Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithm Mol Biol* **6**, 26 (2011).

36. He, K., Zhang, X., Ren, S. & Sun, J. Deep Residual Learning for Image Recognition. *arXiv*
(2015) doi:10.48550/arxiv.1512.03385.
37. Hwang, H., Jeon, H., Yeo, N. & Baek, D. Big data and deep learning for RNA biology. *Exp.*
Mol. Med. **56**, 1293–1321 (2024).
38. Collier, B., Gorgoni, B., Loveridge, C., Cooke, H. J. & Gray, N. K. The DAZL family
proteins are PABP-binding proteins that regulate translation in germ cells. *EMBO J.* **24,**
2656–2666 (2005).
39. Bauermeister, D., Claußen, M. & Pieler, T. A novel role for Celf1 in vegetal RNA
localization during Xenopus oogenesis. *Dev. Biol.* **405**, 214–224 (2015).
40. Dominguez, D. *et al.* Sequence, Structure, and Context Preferences of Human RNA
Binding Proteins. *Mol Cell* **70**, 854-867.e9 (2018).
41. Martins, J. P. S. *et al.* DAZL and CPEB1 regulate mRNA translation synergistically during
oocyte maturation. *J. Cell Sci.* **129**, 1271–1282 (2016).
42. Yang, C.-R. *et al.* The RNA-binding protein DAZL functions as repressor and activator of
mRNA translation during oocyte maturation. *Nat. Commun.* **11**, 1399 (2020).
43. Wu, X. & Bartel, D. P. Widespread Influence of 3'-End Structures on Mammalian mRNA
Processing and Stability. *Cell* **169**, 905-917.e11 (2017).
44. Hoque, M. *et al.* Analysis of alternative cleavage and polyadenylation by 3' region
extraction and deep sequencing. *Nat. Methods* **10**, 133–139 (2013).
45. Beaudoing, E., Freier, S., Wyatt, J. R., Claverie, J. M. & Gautheret, D. Patterns of variant
polyadenylation signal usage in human genes. *Genome Research* **10**, 1001–1010 (2000).
46. Wayment-Steele, H. K. *et al.* RNA secondary structure packages evaluated and
improved by high-throughput experiments. *Nat Methods* 1–9 (2022) doi:10.1038/s41592-
022-01605-0.
47. Piqué, M., López, J. M., Foissac, S., Guigó, R. & Méndez, R. A Combinatorial Code for
CPE-Mediated Translational Control. *Cell* **132**, 434–448 (2008).
48. Pollard, K. S., Hubisz, M. J., Rosenbloom, K. R. & Siepel, A. Detection of nonneutral
substitution rates on mammalian phylogenies. *Genome Res.* **20**, 110–121 (2010).
49. Raney, B. J. *et al.* The UCSC Genome Browser database: 2024 update. *Nucleic Acids*
Res. **52**, D1082–D1088 (2023).
50. Linder, J., Koplik, S. E., Kundaje, A. & Seelig, G. Deciphering the impact of genetic
variation on human polyadenylation using APARENT2. *Genome Biol* **23**, 232 (2022).
51. Padmanabhan, K. & Richter, J. D. Regulated Pumilio-2 binding controls RINGO/Spy
mRNA translation and CPEB activation. *Genes Dev.* **20**, 199–209 (2006).
52. Chen, J. *et al.* Genome-wide analysis of translation reveals a critical role for deleted in
azoospermia-like (Dazl) at the oocyte-to-zygote transition. *Genes & Development* **25**, 755–
766 (2011).
53. Mikedis, M. M. *et al.* DAZL mediates a broad translational program regulating expansion
and differentiation of spermatogonial progenitors. *eLife* **9**, e56523 (2020).
54. Rouskin, S., Zubradt, M., Washietl, S., Kellis, M. & Weissman, J. S. Genome-wide
probing of RNA structure reveals active unfolding of mRNA structures in vivo. *Nature* **505**,
701–705 (2013).
55. Ding, Y. *et al.* In vivo genome-wide profiling of RNA secondary structure reveals novel
regulatory features. *Nature* **505**, 696–700 (2014).

56. Spitale, R. C. *et al.* Structural imprints in vivo decode RNA regulatory mechanisms.
Nature **advance online publication**, 486–490 (2015).
57. Zhang, J., Fei, Y., Sun, L. & Zhang, Q. C. Advances and opportunities in RNA structure
experimental determination and computational modeling. *Nat Methods* **19**, 1193–1207
(2022).
58. Kaufmann, I., Martin, G., Friedlein, A., Langen, H. & Keller, W. Human Fip1 is a subunit
of CPSF that binds to U-rich RNA elements and stimulates poly(A) polymerase. *The EMBO*
Journal **23**, 616–626 (2004).
59. Lackford, B. *et al.* Fip1 regulates mRNA alternative polyadenylation to promote stem
cell self-renewal. *EMBO J.* **33**, 878–889 (2014).
60. Stroup, E. K. & Ji, Z. Deep learning of human polyadenylation sites at nucleotide
resolution reveals molecular determinants of site usage and relevance in disease. *Nat.*
Commun. **14**, 7378 (2023).
61. Sang, Q., Ray, P. F. & Wang, L. Understanding the genetics of human infertility. *Science*
380, 158–163 (2023).
62. DebRoy, H. P. and P. A. and R. G. and S. Biostrings: Efficient manipulation of biological
strings. https://bioconductor.org/packages/Biostrings (2021).
63. Pedregosa, F. *et al.* Scikit-learn: Machine Learning in Python. *Journal of Machine*
Learning Research **12**, 2825–2830 (2011).
64. Abadi, M. *et al.* TensorFlow: Large-Scale Machine Learning on Heterogeneous Systems.
Preprint at https://www.tensorflow.org/.
65. Akiba, T., Sano, S., Yanase, T., Ohta, T. & Koyama, M. Optuna: A Next-generation
Hyperparameter Optimization Framework. *arXiv* (2019) doi:10.48550/arxiv.1907.10902.
66. Liu, D. Algorithms for efficiently collapsing reads with Unique Molecular Identifiers.
PeerJ **7**, e8275 (2019).
67. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).
68. Johnson, P. Z. & Simon, A. E. RNAcanvas: interactive drawing and exploration of nucleic
acid structures. *Nucleic acids Res.* **51**, W501–W508 (2023).
69. Kolberg, L., Raudvere, U., Kuzmin, I., Vilo, J. & Peterson, H. gprofiler2-- an R package for
gene list functional   enrichment analysis and namespace conversion toolset g:Profiler.
F1000Research **9 (ELIXIR)**, (2020).
70. Wang, R., Nambiar, R., Zheng, D. & Tian, B. PolyA_DB 3 catalogs cleavage and
polyadenylation sites identified by deep sequencing in multiple genomes. *Nucleic Acids*
Res. **46**, gkx1000- (2017).
71. Team, R. C. R: A Language and Environment for Statistical Computing. (2021).
1483
1484

1485