1

3   **Tubulin-like proteins CetZ1 and CetZ2 coordinate cellular morphogenesis in the**
4   **archaeon *Haloferax volcanii***

6   Hannah J. Brown[1] & Iain G. Duggin[1#]

8   [1]The Australian Institute for Microbiology and Infection, University of Technology Sydney,
9   NSW, 2007 Australia.

10 #Correspondance: Iain.Duggin@uts.edu.au

Abstract

CetZs are tubulin superfamily cytoskeletal proteins unique to the archaea and have known roles in the control of cell shape and motility. CetZ1, the most conserved CetZ homologue, is essential for rod cell shape development in the pleomorphic archaeon *Haloferax volcanii* and is important for motility and the assembly and positioning of the archaellum and chemosensory arrays. In this study, we report the function of the paralog CetZ2 as a stationary phase-specific protein that contributes to maintenance of plate cell shape. High resolution fluorescence microscopy revealed that CetZ2 subcellular assembly and dynamics are growth-phase specific and highly active in mid-stationary phase. At this stage, CetZ1 and CetZ2 dynamic assemblies were specifically dependent on one another and their GTPase functions, which suggested that CetZ2 operates by inhibiting the rod-promoting function of CetZ1. The impact of CetZ2 on CetZ1 assembly and function was specific to stationary phase, implying that additional stationary-phase factors are involved. CetZ1 and CetZ2 thus coordinate to control cell morphogenesis in the growth cycle in response to starvation or related environmental cues as culture conditions change. CetZ1 and CetZ2 are co-conserved in many haloarchaea, indicating that their coordinated functions are likely to be widely utilized to control dynamic morphogenesis. Our findings show that CetZ paralogues can have distinct and non-redundant functions, and that diversification and specialisation within the tubulin superfamily has occurred multiple times in archaeal evolution.

Significance statement

This study describes the first defined function of a CetZ2 protein, which represents a conserved haloarchaeal family within the near-universal tubulin superfamily of cytoskeletal proteins. In the model archaeal organism *Haloferax volcanii*, CetZ1 is required for the development of rod-shaped cells in early exponential phase growth in a CetZ2-independent manner. We found that CetZ2 has a contrasting role in maintaining the plate or disk-like cell morphology during stationary phase, by inhibiting CetZ1. Our findings reveal a novel antagonism between families of tubulin superfamily proteins, which may represent a paradigm for understanding the duplication and diversification of cytoskeletal functions that occurred extensively during the early evolution of Eukaryotic cells, and appears to have occurred multiple times in archaea, potentially predating eukaryotes.

Introduction

CetZs are archaea-specific tubulin superfamily proteins that are structural homologues of eukaryotic tubulin which forms microtubules, and FtsZ, the key cell division protein present in both bacteria and archaea (1, 2). One shared characteristic of these tubulin superfamily proteins is they polymerise in a GTP-dependent manner, forming filaments or large polymers (3-9), which assemble into larger structures that carry out their biological functions. Both FtsZ and tubulin have a wide diversity of functions which are well studied, including in cell division and chromosome segregation (10, 11), intracellular trafficking and cargo transport (12, 13), and cell motility or migration (14, 15).

CetZs are conserved across many diverse archaea. In Halobacteria (Haloarchaea), it is typical that one species would have multiple paralogues of CetZ, and the most highly conserved of these are CetZ1 and CetZ2 (1, 16, 17), both of which form strong and distinct phylogenetic clusters, suggesting that CetZ1 and CetZ2 could have distinct functions. While both CetZ1 and CetZ2 have been implicated in the control of shape and motility in *Haloferax volcanii,* only CetZ1 currently has a clearly defined biological function. During standard batch culture, *H. volcanii* transitions from plate-shaped cells to rods between the early and late log phases, but once cells enter stationary phase they revert to plates (18). CetZ1 has shown to be necessary for rod development, not only during early-log (1, 18), but also in other conditions where *H. volcanii* becomes rod-shaped, such as in motility conditions (1, 19, 20) or when depleted of trace metals(18). CetZ1 localises in patches around the membrane of cells but is somewhat concentrated at mid-cell and the cell poles, where it is hypothesised to form a cap-like structure in motile rods (1, 20-23). CetZ1 is thought to contribute to motility by generating the rod shape that enables directional and streamlined swimming, as well as by promoting the positioning and assembly of archaella and chemosensory arrays (21). CetZ1 shows highly dynamic localization during growth and rod development, which is linked to its polymerisation and depolymerisation cycle (1, 22). A point mutation in the GTPase active site (CetZ1.E218A) disrupts the GTPase activity and causes CetZ1 to form hyperstable filaments, disrupting CetZ1 protein dynamics. Expression of this mutant prevents rod development and produces cells with local regions of high envelope curvature, or 'jagged' cells, and CetZ1.E218A localises to the regions of high membrane curvature (1, 22).

While CetZ1 has roles in cell shape control and motility, no clear role of other CetZ paralogues of *H. volcanii* have been identified. Although CetZ2 has been generally implicated in the control of cell shape and motility (1, 21), deletion of *cetZ2* results in no cell division, shape, or motility phenotypes during exponential phase or on soft-agar, respectively

(1, 21). However, overexpression of CetZ2 has been demonstrated to induce slight
hypermotility and promote assembly of archaellum filaments (21). In addition, expression of
CetZ2.E212A, the equivalent GTPase active-site mutant of CetZ1.E218A, prevents rod-
development and perturbs motility (1, 21). CetZ2.E212A might therefore interfere with CetZ1
function (1), whether through direct interaction with CetZ1 or indirectly through an unknown
mechanism. Together with the knowledge that CetZ2 is only present in Halobacteria that
also have CetZ1 (17), the current evidence points towards a potential functional interplay
between CetZ2 and CetZ1. One possible explanation for the lack of observable effects on
cell shape and motility upon deletion of *cetZ2* is that the CetZ2 protein may be inactive or
present in very low abundance in the tested conditions (i.e. early/mid-log growth, motility
assays). Here, we confirm this hypothesis and show that CetZ2 is strongly upregulated in
stationary phase where it maintains plate shape by inhibiting CetZ1-dependent rod
development.

Methods

Materials and reagents

Unless otherwise specified, all chemicals were obtained from Sigma, reagents for cloning
procedures were obtained from New England Biolabs and used according to the
manufacturer's instructions, and custom oligonucleotides were obtained from Integrated
DNA Technologies. The Meridian Bioscience ISOLATE II PCR and Gel Kit and ISOLATE II
Plasmid Mini Kit were used for purification of DNA from PCR reactions and agarose gels,
and for plasmid extractions, respectively.

H. volcanii strains and growth

A list of *H. volcanii* strains used in this study is available in Table S1. Cultures were prepared
where indicated in standard culture media, either Hv-Cab (1), Hv-YPCab (18), or Hv-Min with
trace elements (24) and incubated at 42 °C with rotary shaking (200 rpm). If necessary,
medium was supplemented with uracil (50 µg/mL) to fulfil auxotrophic requirements of
Δ*pyrE2* strains, or with □-tryptophan to induce expression of genes under the control of the
p.*tna* promotor. For experiments that compared mid-log and stationary phase, cells were
cultured in 10 mL Hv-YPCab medium by dilution of a steady mid-log starter culture to a
theoretical starting OD_{600} of 0.0005. Mid-log phase samples of cells were taken after a
further 24 h of culturing, and then for early, mid, and late stationary phase, samples were
taken at 72, 96, and 120 h, respectively.

Genetic modification

The wild-type H26 (ID621) genetic background of *H. volcanii*, which is auxotrophic for uracil
(Δ*pyrE2*), was used throughout this study. To generate the *cetZ1* and *cetZ2* knockout
strains, pTA131-based plasmids for deletion of *cetZ1*(18) and *cetZ2*(1), containing the
upstream and downstream flanks of *cetZ1* and *cetZ2* and the *pyrE2* gene, were used to
transform H26 with selection on Hv-Cab agar. Transformation was done using the previously
described method(25) and two-step ('pop-in-pop-out') homologous recombination (26) was
used for selection of transformants that showed complete deletion of the target gene.

To generate the *cetZ2-mTq2CHR* chromosome tag strain, pTA131-*cetZ2mTq2* was obtained
synthesised (Genscript), which contained a 350 bp fragment encoding the C-terminal portion
of *cetZ2*, fused in frame to an ORF containing the EG linker (semi-rigid) and mTurquoise2
(20), followed by the 350 bp flank immediately downstream of the *cetZ2* stop codon. This
plasmid was used to transform H26 by the two-step method (26) to produce a scarless
insertion of EG-mTurquiose2 before the native *cetZ2* stop codon.

Construction of plasmids for gene expression in H. volcanii

A list of plasmids used in this study is available in Table S2 and relevant oligonucleotides are listed in Table S3. To generate plasmids for single and dual expression of CetZ fusion proteins, we used a recently described plasmid-based expression system (20). Briefly, the ORFs for *cetZ2* or *cetZ2*.E212A excluding their stop codons were PCR amplified using primers to incorporate NdeI and BamHI sites at the 5' and 3' ends, respectively, and the resulting PCR product and the vector backbone were ligated using T4 DNA ligase. The vector backbone was pHVID21 for C-terminal mTurquoise2 fusions with the EG linker, or pHVID20 for C-terminal YPet fusions with the EG linker.

To generate plasmids for dual expression of two tagged proteins, pHJB5 and pHJB11 (containing *cetZ2*-YPet and *cetZ2*.E212A-YPet, respectively) were used as vector backbones for insertion of *cetZ1*-mTq2 and *cetZ1*.E218A-mTq2 ORFs positioned immediately downstream of the *cetZ2* ORFs as described (20). The ORFs for *cetZ2*-YPet, *cetZ2*.E212A-YPet, *cetZ1*-mTq2, and *cetZ1*.E218A-mTq2 were sourced from pHJB5, pHJB11, pHVID135, and pHJB13, respectively.

Plasmids for dual expression of one untagged CetZ and one tagged CetZ were prepared using plasmids for expression of *cetZ2*-mTq2, *cetZ2*, or *cetZ2*.E212A (pHJB6, pTA962-*cetZ2*, and pTA962-*cetZ2*.E212A, respectively) as backbones. pHJB6 was subjected to restriction digest by NheI and NotI, while pTA962-*cetZ2* and pTA962-*cetZ2*.E212A were digested using BamHI and NotI. ORFs for *cetZ1* and *cetZ1*.E218A were amplified by PCR incorporating an NheI site at the 5' end, and the ORF for *cetZ1*-mTq2 was amplified to incorporate a BglII site at the 5' end. After restriction digest with the relevant enzymes, the various *cetZ1* ORFs were ligated as described above, into the backbones, generating plasmids pHJB43-46 for dual expression of *cetZ2/cetZ1*-mTq2, *cetZ2*.E212A/*cetZ1*-mTq2, *cetZ2*-mTq2/*cetZ1*, and *cetZ2*-mTq2/*cetZ1*.E218A, respectively. All plasmid regions originating from PCR products were sequenced verified (Australian Genome Research Facility). Plasmids were passaged through *E. coli* strain C2925 for demethylation before transformation of *H. volcanii*.

Soft-agar motility assay

H. volcanii was maintained in exponential growth in liquid culture (Hv-Cab medium) for at least two days. Following this, cultures were normalised to OD_{600}~0.1, and 2 µL of cells were spotted onto the surface of Hv-Cab agar (0.2% w/v). Soft-agar plates were incubated statically at 42 °C for 4 days before extracting cells from the halo edge or inner halo (approximately 5 mm from the inoculation site) as described (21) for further analysis by microscopy.

Microscopy

For all microscopy, 2 µL of cells were spotted onto a 1.5% (w/v) agarose pad containing 18% buffered salt water (26), and a glass coverslip (#1.5) was placed on top before imaging. Phase contrast and fluorescent images were acquired using a V3 DeltaVision Elite inverted fluorescence microscope (GE Healthcare), with CFP (excitation filter = 400-453 nm; emission filter = 463-487 nm) and YFP (excitation filter = 497-527 nm; emission filter = 537-549 nm) filter sets. Images were acquired using a 100X 1.4 NA Plan Apo objective. Time-lapse images were recorded at 1 min intervals for 30 min. Exposure and acquisition settings were maintained between strains and replicates within each experiment.

Three-dimensional structured-illumination microscopy (3D-SIM) was carried out as previously described (22, 23), using a DeltaVision OMX SR (GE Healthcare) microscope with a 60X Plan 1.42 NA objective, a PCO sCMOS camera, 125 nm Z-step size, with a DAPI excitation filter (405 nm) and AF488 emission filter (504-552 nm). Raw images were reconstructed using SoftWorX (Applied Precision, GE), and then maximum intensity projections were generated in FIJI (27), and 3D renders were generated using Imaris (V 7.6.4 BitPlane Scientific).

Image analysis

All shape analysis of cells was conducted using phase-contrast images analysed using MicrobeJ v5.13m (28) in FIJI (27). To detect cell outlines using phase-contrast images default settings were applied whilst restricting area between 0.8-max µm^2, and sinuosity between 0-2. For all analysis of fluorescence, raw images and time-lapse data were first processed by applying the *Background Subtraction* function (rolling ball size = 8 pixels). Time-lapse data were further processed using *Bleach Correction* (algorithm = *Histogram Matching*), and *StackReg* (transformation = *Translation*).

For static localisation analysis, the *foci* detection method with default settings was used with tolerance set to 1000, Z-score was set to 25, and area was restricted between 0-0.5 µm^2. Mean whole-cell fluorescence intensity (Bacteria.Intensity.ch2.mean), the number of localisations per cell (Bacteria.Maxima), localisation area (Maxima.Shape.area), and maximum localisation intensity (Maxima.Intensity.max) were extracted from MicrobeJ results tables, and localisation heat maps were generated using the *XYCellDensity* function. For tracking of localisations, TrackMate2 v7.7.2 (29, 30) was used. The LoG detector was used to detect foci, using an estimated foci diameter of 0.5 µm and a quality threshold of 100. For tracking, the LAP tracker was used, allowing for frame-to-frame linking, track segment gap closing, track segment splitting, and track segment merging, with a maximum distance of 0.5 µm and a maximum frame gap of 2.

EzColocalisation (31) was used for colocalization analysis. Manual adjustment of thresholds
using the default algorithm was required for accurate detection of fluorescent channels,
ensuring that weak or background fluorescence was not included in the thresholding. Auto
thresholding with the default algorithm and watershed segmentation was used for cell
identification using the phase-contrast channel. Fluorescent channels were aligned, and the
background was subtracted using a rolling-ball size of 8. All remaining settings and
parameters were used as default.

SDS-PAGE and Western blotting

Samples were prepared for SDS-PAGE by resuspending whole cell pellets in lysis buffer (20
mM Tris-HCl pH 7.5, 1X DNAseI (ThermoFisher), 1X EDTA-free protease inhibitor
(ThermoFisher)) and incubating for 30 min at room temperature. Following that, 4X SDS-
PAGE sample buffer (250 mM Tris-HCl pH 6.8, 40% (v/v) glycerol, 4% (w/v) SDS, 0.04%
(w/v) Bromophenol blue, 20% (v/v) 2-mercaptoethanol) was added to a final concentration of
1X, using a volume that would give a theoretical OD_{600} of 5 based on the original culture.
The samples heated at 95 °C for 5 min then vortexed for 1 min.

CetZ1-6xHis and CetZ2-6xHis proteins (1)were purified and quantified by $A_{280\ nm}$ in a
previous study (1) and diluted in 1X SDS-PAGE sample buffer to 200 ng/μL and 50 ng/μL,
respectively. Following this, 1:2 serial dilution in 1X SDS-PAGE sample buffer were carried
out for each protein. Samples were loaded onto a 4-20% Mini-PROTEAN® TGX™ pre-cast
polyacrylamide gel (Bio-Rad), using the Broad Range Blue Prestained Protein Standard
(New England Biolabs). Electrophoresis was carried out at 100 V for 85 min with a Bio-Rad
Mini-Protean3 electrophoresis system.

For western blotting, proteins separated by SDS-PAGE were transferred to a 0.2 μm
nitrocellulose membrane using the Trans-Blot Turbo mini transfer system (Bio-Rad). The
membrane was stained with Ponceau S (0.1% (w/v) Ponceau S in 5% (v/v) acetic acid) for 5
min and the excess stain removed by rinsing with Milli-Q water to obtain a total protein stain.
Following this, the nitrocellulose membrane was blocked in 5% (w/v) skim milk powder in
TBST (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 0.05% (v/v) Tween-20), for 2 h at room
temperature. Primary antibodies(1) against CetZ1 or CetZ2 were diluted 1/500 in TBST for
incubation with the membrane overnight at 4 °C. The membrane was washed 3 times in
TBST and incubated with the secondary antibody (donkey anti-rabbit IgG HRP conjugate,
AbCam 16284) diluted 1/5000 in TBST for 2 h at room temperature. The membrane was
washed 5 times in TBST and incubated with the SuperSignal™ West Pico PLUS
chemiluminescent substrate (ThermoFisher) for 5 min before imaging of the membrane
using an Amersham™ Imager 600 (GE Healthcare).

To calculate the number of molecules of CetZ1/2 per cell, six replicate cultures were grown

in Hv-YPCab medium and sampled after 24 and 120 h. Band intensity from western blotting

analysis (Fig. 1d, e) was determined by measuring the raw integrated density in ImageJ(27)

and subtracting the raw integrated density of the background. The same measurements

were conducted for pure protein standards of known concentrations. Band intensities of the

standards were used to determine the amount of protein per OD unit of cells in mid-log (24

h) and stationary phase samples. The number of cells per OD unit in mid-log and stationary

phase was calculated using serial dilutions and counting the number colony forming units

from the same six culture replicates used for western blotting samples. Finally, this was used

to determine the number of CetZ1 and CetZ2 molecules per cell.

Results

CetZ2 is strongly upregulated in stationary phase in H. volcanii

CetZ2 function has previously been investigated during exponential phase growth and motility conditions. However, proteomic data (32) suggested it may be upregulated in stationary phase in *H. volcanii* and *N. magadii*. Here, we used western blotting to detect CetZ2 levels in Hv-YPCab (rich medium), Hv-Cab (casamino acids medium), and Hv-Min (minimal medium) between early log phase and late stationary phase of *H. volcanii* (Fig. 1a). This showed that in all media there is little production of CetZ2 until stationary phase, where protein levels strongly increase (Fig. 1c). Conversely, CetZ1 was expressed throughout, though decreased in stationary phase (Fig. 1b). Downregulation of CetZ1 in stationary phase was not dependent on the presence of CetZ2 (Fig. S1a), and upregulation of CetZ2 was not dependant on the presence of CetZ1 (Fig. S1b). The ratios between CetZ1 and CetZ2 and the number of molecules per cell were also measured using CetZ1/2 pure protein standards (Fig. 1d, e, respectively). Wildtype cells in mid-log growth were measured to have on average 5735 molecules of CetZ1 and 142 molecules of CetZ2, giving a ratio of ~41:1 (CetZ1:CetZ2, Fig. 1f). In stationary phase, the amounts averaged 2234 molecules of CetZ1 and 425 molecules of CetZ2 per cell, a ~5:1 ratio (Fig. 1f).

Figure 1. *Quantification of CetZ1 and CetZ2 in H. volcanii throughout growth.* A) Wild-type cells (H26 + pTA962) were grown in Hv-YPCab, Hv-Cab, or Hv-Min from a theoretical starting OD_{600}~0.0005 for 120 h. Samples of each culture were taken for western blotting at 12 points along each growth curve, indicated by black data points. **B)** Western blot analysis of wild-type cells grown in **A** using antibodies against CetZ1, and **C)** CetZ2. Samples from timepoints 2-12 of each medium were used for analysis. Ponceau S staining (bottom panels) was carried out to ensure equal sample loading. Solid arrow heads indicate CetZ1/2 bands. **D)** Purified CetZ1 was used to make eight standards of known concentrations. Standard 1 is 50 ng of pure CetZ1, and standards 2-8 are 1:2

serial dilutions of standard 1. Standards were compared to whole-cell lysate samples of wild-type cells
to estimate the number of CetZ1 molecules per cell in mid-log and stationary phase using western
blot analysis against CetZ1. **E)** As in D, except for CetZ2. Standard 1 is 1.25 ng of pure CetZ2, and
standards 2-6 are 1:2 serial dilutions of standard 1. In **D** and **E**, six culture replicates were carried out,
and the same samples were used for analysis of CetZ1 and CetZ2 expression levels. Solid arrow
heads indicate CetZ1/2 bands. **F)** From the results of **D** and **E**, the number of CetZ1/CetZ2 molecules
per cell in mid-log and stationary phase were calculated as described in the methods. Mean and
standard deviation are shown. Unpaired one-way ANOVA was used as a statistical test.

CetZ2 contributes to maintenance of plate-shape in stationary phase

CetZ2 has previously been implicated in cell shape through the expression of a GTPase
active-site mutant (CetZ2.E212A) (1). This type of T7-loop mutation is expected to inhibit
GTPase activity in tubulin-superfamily proteins, producing inactive hyper-stable filaments.
Production of CetZ2.E212A resulted in a dominant inhibitory effect, causing distorted cell
shapes, similar to the effects of the equivalent mutant CetZ1.E218A, but did not reveal the
normal role of CetZ2 (1). We therefore tested whether CetZ2 influences cell shape in
stationary phase compared to mid-log phase using deletion, overexpression, and GTPase
mutations. Mid-log cultures of these strains were diluted to a theoretical OD_{600} of 0.0005 in
10 mL Hv-YPCab medium with 0.2 or 1 mM □-tryptophan (Trp) to induce expression of the
relevant plasmid-encoded genes under the control of the p.*tna* promotor. Samples for mid-
log, early stationary, mid-stationary, and late stationary phases were taken after 24, 72, 96,
and 120 h of incubation, respectively (Fig 1a).

At the 24 h mid-log stage, where wild-type cells are largely in a rod-shaped form before
reverting to plate shaped cells (1, 21), deletion and overexpression of *cetZ2* had no obvious
effect on cell shape, whereas production of the GTPase inactive mutant, CetZ2.E212A,
strongly disrupted rod development (Fig. 2a, top panels). These observations were
quantified and confirmed by analysis of the circularity of cell outlines (Fig. 2b, left). In late
stationary phase (120 h), wild-type cells were a mixture of plate and short rod shapes (Fig.
2a, lower panels). However, deletion of *cetZ2* resulted in more elongation of cells (120 h),
whereas overexpression of *cetZ2* or *cetZ2*.E212A increased cell circularity in comparison to
the wild-type (Fig. 2a-c). These phenotypes were clear with multiple culture replicates (Fig.
2c) and demonstrate that CetZ2 has a role in maintaining plate shape in stationary phase.

To investigate the relative timing of CetZ2 upregulation and the transition to plate-shape that
occurs in mid-late log growth (19, 33), we developed a chromosomal fluorescent fusion of
cetZ2-mTq2 under the control of the native *cetZ2* promotor (*cetZ2-mTq2CHR*). This strain
exhibited cell circularity (Fig. S2a) and CetZ2 upregulation in stationary phase (Fig. S2b) that
was equivalent to the wild-type, indicating full functionality of the fusion. The total cellular
fluorescence intensity of *cetZ2-mTq2CHR* and cell circularity were then measured for

individual cells during the transition to stationary phase (e.g. Fig. S2c). This showed that
CetZ2 upregulation occurred between 72 and 96 hours of culturing (Fig. 2d, blue), which was
well after the completion the transition of rod-shaped cells to plate-shaped cells at
approximately 30 h (Fig. 2d, red). This suggested that CetZ2 upregulation is not linked to
mid-log plate shape development, and rather has a role in maintenance of plate-shape in
stationary phase.

Figure 2. *CetZ2 contributes to maintenance of plate-shape in stationary phase.* H26 + pTA962
(WT), Δ*cetZ2* + pTA962, Δ*cetZ2* + pTA962-CetZ2, and WT + pTA962-CetZ2.E212A were grown in
Hv-YPCab medium supplemented with 1 mM ⎕-tryptophan and sampled during mid-log phase (24 h
after inoculation) and late stationary phase (120 h after inoculation). **A)** Phase-contrast microscopy
images (scale bar=5 µm) and **B)** circularity analysis of individual cells during mid-log and late
stationary phase. One representative culture replicate for each strain is shown. **C)** Mean cell
circularity (SD) of 13 replicate cultures in late stationary phase (120 h). One-way ANOVA was used as
a statistical test. The total number of cells analysed from the 13 culture replicates were: WT
(+pTA962), n=14,242, ΔcetZ2 (+pTA962), n=23,671; ΔcetZ2 (+pTA962-cetZ2), n=16,416. **D)** *cetZ2-
mTq2CHR* + pTA962 was grown in Hv-YPCab medium from a theoretical starting OD$_{600}$~0.0005 and
observed by phase-contrast and fluorescence microscopy after 24, 30, 48, 72, 96, and 120 h of
culturing. The circularity and mean fluorescence intensity of individual cells was analysed at each time
point. Individual points represent mean of the population at each timepoint, and dotted lines indicate
standard deviation. The number of cells analysed at each timepoint is as follows: 24 h, n=253; 30 h,
n=554; 48 h, n=1209; 72 h, n=1872; 96 h, n=728; 120 h, n=187.

Previous work showed that *cetZ2* is dispensable for swimming motility (1), but that its
overexpression can slightly enhance motility(21). The actively swimming cells at the leading
edge of the expanding halo in soft agar motility assays are rod-shaped, whereas those in the
interior of the halo (that are slower or not swimming) have transitioned to plate shapes (1).
We therefore investigated the relative expression of *cetZ2* in these cells. The *cetZ2-*
mTq2CHR strain was fully functional in swimming motility (Fig. S2d, e) and fluorescence
measurements indicated that CetZ2-mTq2 was only detectable above background in plate-
shaped cells sampled from the inner motility halo (Fig. S2f-i). Thus, the relationship between
CetZ2 levels and cell morphology is consistent between batch culture and motility assay
conditions, and these observations are consistent with the notion that CetZ2 is only
expressed or effective in plate-shaped cells which have already reverted from rod-shape
(21).

CetZ2 localisation changes throughout stationary phase

We observed that the *cetZ2-mTq2CHR* strain did not show clear CetZ2-mTq2 subcellular
localisation above the background of faint whole cell fluorescence during the stationary
phase transition (Fig. S2c). Therefore, to further investigate CetZ2 localisation and function,
we expressed the CetZ2-mTq2 fusion (21) and an additional CetZ2-YPet fusion with a
previously described vector system (20). Both CetZ2-mTq2 and CetZ2-YPet were functional
in this context, as they increased cell circularity equivalently to untagged CetZ2 produced
from the same vector backbone (Fig. S3a). Likewise, CetZ2.E212A-mTq2 and
CetZ2.E212A-YPet had equivalent effects to untagged CetZ2.E212A (Fig. S3b). Western
blotting showed that the plasmid-based system resulted in strong overproduction of the
tagged and untagged CetZ2 and CetZ2.E212A compared to endogenous CetZ2 in the wild-
type (Fig. S3c), even with no addition of the □-tryptophan inducer (Fig. S3d). Interestingly,
when CetZ2-mTq2 production was induced with 1 mM of □-tryptophan throughout the
growth cycle, the level of CetZ2-mTq2 still increased substantially in stationary phase (Fig.
S3e), suggesting the existence of a post-transcriptional or post-translational mechanism for
the regulation of CetZ2 levels.

Using the plasmid system, CetZ2-mTq2 localisation was observed throughout the growth
cycle (Fig. 3a). During the mid-log phase, CetZ2 localised in patches throughout the cell,
around the membrane, at mid-cell, and at the cell poles, similar to the previously reported
localisation of CetZ1 in rod-shaped cells in liquid culture (1, 22) and consistent with the
recently reported localisation of overproduced CetZ2 in motile rod-shaped cells(21). In early
stationary phase (72 h), CetZ2 localised strongly around the cell outline in small patches and
filaments. During mid-stationary phase (96 h), CetZ2 localisation was similar to that in early

stationary phase, and an equivalent number of localisations were observed per cell (Fig. 3b),
however localisation area (Fig. 3c) and maximum intensity of localisations (Fig. 3d, e)
increased. In late stationary phase (120 h), CetZ2 localisation changed substantially, being
mostly diffuse throughout the cell with one or two cell-edge associated foci (Fig. 3a, b).
These foci were decreased in area and maximum intensity in comparison to early and mid-
stationary phase, (Fig. 3c-d). At all timepoints throughout stationary phase, CetZ2-mTq2 was
distributed around the cell edge, though showed a weak preference for the polar regions as
defined by the longest identified axis (Fig. 3f). 3D-Structured Illumination Microscopy (3D-
SIM) of CetZ2-mTq2 during mid-stationary phase (96 h, Fig. 3g, Supplementary Video SV.
1), when localisations were largest and brightest, showed clear filaments or patches around
the edges of the cell that do not often localise on the top and bottom surfaces. [*H. volcanii*
cells have a flattened profile (~0.4 μm thick) and tend to settle on their larger flatter surfaces
on agarose pads.]

Figure 3. *Subcellular localization of plasmid-produced CetZ2-mTq2 in stationary phase.* A) WT
+ pHJB6 (for expression of CetZ2-mTq2) was grown in Hv-YPCab medium supplemented with 0.2

mM ☐-tryptophan and imaged using fluorescence microscopy at 24, 72, 96, and 120 h. Scale bar=5
μm. **B)** 3D-SIM maximum intensity Z-projections of CetZ2-mTq2 at 96 h. Scale bar=2 μm. **C)** CetZ2-
mTq2 localisation characteristics at 72, 96, and 120 h, including the number of localisations per cell,
D) Localisation area, and **E)** Maximum localisation intensity. **F)** Maximum fluorescence intensity by
localisation number at 72, 96, and 120 h, where localisation 1 indicates the brightest detected
localisation in any given cell, localisation 2 indicates the next brightest, etc. The number of
localisations constituting each violin plot is detailed above it. **G)** Heatmap representations of CetZ2-
mTq2 localisations, coloured by density to indicate frequency of localisation. The total number of cells
analysed (n_c) and total number of localisations detected from those cells (n_L) at each timepoint are
indicated in the key in panels **C** and **D**, respectively, and applies to panels **C**-**G**.

CetZ2 is highly dynamic in mid-stationary phase

Tubulin superfamily proteins including CetZ1 are typically highly dynamic due to their ability

to polymerise and depolymerise, allowing them to carry out their functions. Hence, time-

lapse imaging of CetZ2-mTq2 was conducted over a total duration of 30 min with 1 min

intervals to observe any movement of CetZ2 structures in stationary phase. In early and late

stationary phase, CetZ2 localisation was relatively static, but in mid-stationary phase CetZ2

was highly dynamic (Fig. 4a, SV. 2). Tracking of filaments and patches (29, 30) (SV. 3)

confirmed that CetZ2 movement was fastest in mid-stationary phase (Fig. 4b). At this stage,

cells displayed three main types of dynamic CetZ2-mTq2 patterns (Fig. 4c, SV. 2). These

included a 'circling' pattern (42% of cells), in which CetZ2-mTq2 was observed travelling

around the perimeter of the cell adjacent to the inner surface of the membrane, a complex

pattern (24% of cells), which showed an apparently more random directional motion, as well

as a 'confined' pattern (34% of cells), which were static or relatively slow and envelope

associated. We also investigated the localisation and movement of CetZ2.E212A-mTq2 and

found that this mutant localized as foci and patches or filaments throughout in cells

throughout stationary phase (Fig. S4) but was static at 96 h compared to the wild-type (Fig.

S4, SV. 4). Thus, the dynamic localization of CetZ2 in stationary phase appears to be

dependent on the GTPase activity of CetZ2.

Figure 4. *Dynamic behaviours of plasmid-produced CetZ2-mTq2 in mid-stationary phase.* WT + pHJB6 cells (for expression of CetZ2-mTq2) were grown in Hv-YPCab medium supplemented with 0.2 mM ⊡-tryptophan and **A)** time-lapse imaging of cells was conducted during early- mid- and late-stationary phase (72, 96, and 120 h, respectively). Images were taken at 1 min intervals for 30 min. Scale bar=2 μm. TrackMate2 was used to track localisation speed. **B)** Frequency distribution of localisation speed. The total number of detected localisations (n_L) at each timepoint is indicated in the key. **C)** Overlayed tracks of localisation dynamics generated in TrackMate2 were used to define characteristic patterns of CetZ2 dynamics in mid-stationary phase (96 h), determine the proportion of cells displaying each dynamic pattern (n=176).

CetZ2 localisation patterns are highly dependent on the presence of CetZ1

As CetZ2 is only present in species which also have CetZ1 (17), and CetZ2.E212A production inhibits motile rod development (1), it might maintain plate shape in stationary phase by inhibiting CetZ1 function. To begin investigating such potential interactions, we determined whether each protein's subcellular localization in mid-stationary phase was dependent on the presence of the other. In the absence of CetZ1 (Δ*cetZ1* background),

CetZ2-mTq2 did not form patches or large filaments associated with the cell edge as in the
wild-type background, but instead mis-localised as discrete and immobile foci that were
distributed throughout the cell area (Fig. 5a, g, SV. 5). CetZ2.E212A-mTq2 also mis-
localised in the absence of CetZ1, with much fewer foci detected per cell and mostly diffuse
fluorescence compared to the strong and stable localizations of CetZ2.E212A-mTq2
observed in the wild-type background (Fig. 5b, e, SV. 6).

During mid-log growth, CetZ2-mTq2 localisation patterns were similar to CetZ1-mTq2 (Fig.
S5a, b), and the patches were noticeably more distributed throughout the cell area
compared to prevalent cell-edge patches seen during stationary phase (Fig. 4a). The mid-log
patterns of CetZ2-mTq2 differed in the absence of CetZ1, like we observed during mid-
stationary phase (Fig. 5a, S5a). The GTPase mutant CetZ2.E212A-mTq2 partially inhibited
rod-shape during mid-log growth when induced using a moderate concentration (0.2 mM) of
L-tryptophan, as expected (1), and its localization also differed in the absence of CetZ1 (Fig.
S5c).

The formation and stability of CetZ1 structures are dependent on the presence of CetZ2 in stationary phase

In mid-stationary phase in the wild-type background, CetZ1-mTq2 localised in patches and
small foci, mostly concentrated around the cell edge, but occasionally, larger filaments were
also observed (Fig. 5c). Time-lapse imaging showed that while some of the small foci and
diffuse regions of CetZ1-mTq2 localization were relatively dynamic, the substantive filaments
and patches were stable over the 30 min (SV. 7). The CetZ1-mTq2 localisation pattern was
similar in the Δ*cetZ2* and wild-type backgrounds (Fig. 5c), but in Δ*cetZ2*, the larger bright
filaments were more frequent (Fig. 5c, e, f). Interestingly, in the absence of CetZ2, some of
the bright CetZ1 filaments were observed to shrink over the 30 min, which was not apparent
in the wild-type background (SV. 7). Together, these results suggest that CetZ2 can inhibit
CetZ1 filament formation yet can stabilize CetZ1 filaments if they form; both effects would be
likely to inhibit CetZ1 function.

CetZ1.E218A localised as long, curved, or bent filaments in both the wild-type and Δ*cetZ2*
backgrounds (Fig. 5d), but somewhat less frequently and with higher intensity in the absence
of CetZ2 (Fig. 5e, f). CetZ1.E218A was not dynamic in either background (SV. 8),
demonstrating that CetZ2 is not required for the stabilization mechanism of CetZ1.E218A.
During mid-log phase, CetZ1-mTq2 (Fig. S5b) and CetZ1.E218A-mTq2 (Fig. S5d) had
indistinguishable localisation patterns in wild-type and Δ*cetZ2* backgrounds.

456
Figure 5. Localisation interdependency of CetZ1 and CetZ2 in mid-stationary phase. Cells were
grown in Hv-YPCab medium supplemented with 0.2 mM ☐-tryptophan and imaged after 96 h (mid-
stationary phase). **A)** CetZ2-mTq2 (pHJB6) and **B)** CetZ2.E212A (pHJB12) localisation in WT and
Δ*cetZ1*. **C)** CetZ1-mTq2 (pHVID135) and **D)** CetZ1.E218A-mTq2 (pHJB13) localisation in WT and
Δ*cetZ2*. For **A-D**, scale bar=5 µm. Localisation analysis was carried out on this data, measuring **E)** the
number of localisations per cell, **F)** maximum intensity of localisations, and **G)** the distance of each
localisation from the cell edge. The number of cells (n_c) and number of localisations (n_L)analysed in **E-**
F is as follows: WT +CetZ2-mTq2: n_c=1482, n_L=4657; Δ*cetZ1* +CetZ2-mTq2: n_c=904, n_L=3201; WT
+CetZ2.E212A-mTq2: n_c=347, n_L=1407; Δ*cetZ1* +CetZ2.E212A-mTq2: n_c=609, n_L=1290; WT
+CetZ1-mTq2: n_c=969, n_L=3558; Δ*cetZ2* +CetZ1-mTq2: n_c=493, n_L=1122; WT +CetZ1.E218A-mTq2:
n_c=815, n_L=1986; Δ*cetZ2* +CetZ1.E218A-mTq2: n_c=1654, n_L=2639. **H)** CetZ1-mTq2 was co-
expressed with untagged CetZ2 (pHJB43) or untagged CetZ2.E212A (pHJB44) in the WT

background. **I)** CetZ2-mTq2 was co-expressed with untagged CetZ1 (pHJB45) or untagged
CetZ1.E218A (pHJB46). Localisation analysis was conducted on data from **H** and **I**, measuring **J)** the
number of localisations per cell and **K)** maximum localisation intensity. The number of cells (n_c) and
number of localisations (n_L) analysed in **J** and **K** is as follows: WT +pHJB43: n_c=3118, n_L=16536; WT
+pHJB44: n_c=3960, n_L=16406; WT +pHJB45: n_c=2893, n_L=6256; WT +pHJB46: n_c=3278, n_L=56450.
L) Time-laps imaging was conducted for strains in **H** and **I** (SV. 9 and SV. 10, respectively), and the
average localisation speed measured. The number of localisations measured was as follows: WT
+pHJB43: n=1595; WT +pHJB44: n=524; WT +pHJB45: n=327; WT +pHJB46: n=329. The key in **L**
applies to panels **G** and **H**.

CetZ1/2 GTPase mutants interfere with the localisation dynamics of the other CetZ

Next, we observed the effect of the filament-stabilizing GTPase mutations of CetZ1 or CetZ2
on the localisation and dynamics of the other CetZ in mid-stationary phase. This was
expected to reveal the extent to which each protein's GTPase influences dynamic filament
turnover of the other protein through potential co-assemblies. Plasmids pHJB43-46 were
generated for dual expression of *cetZ2/cetZ1-mTq2*, *cetZ2.E212A/cetZ1-mTq2*, *cetZ2-*
mTq2/cetZ1, and *cetZ2-mTq2/cetZ1.E218A,* respectively; the *cetZ2* ORFs were located
immediately downstream from the p.*tna* promotor, followed by the *cetZ1* ORFs in a
bicistronic configuration.

The patterns of CetZ1-mTq2 localisation were similar during co-expression with *cetZ2* or
cetZ2.E212A (Fig. 5h) and showed somewhat larger irregular CetZ1-mTq2 patches and
filaments compared to the wild-type and Δ*cetZ2* backgrounds (Fig. 5c). The number of
CetZ1-mTq2 localisations per cell was greater during expression with *cetZ2*.E212A
compared to *cetZ2* (Fig. 5j), suggesting that CetZ2.E212A may promote or stabilise CetZ1-
mTq2 localisations. Time-lapse imaging (SV. 9) and quantification of focus speed (Fig. 5l)
revealed that CetZ2.E212A partially stabilized CetZ1-mTq2 localizations.

The patterns of CetZ2-mTq2 localisation were similar during co-expression with *cetZ1* or
cetZ1.E218A (Fig. 5i), and showed somewhat larger irregular CetZ2-mTq2 patches and
filaments compared to the wild-type and Δ*cetZ1* backgrounds (Fig. 5a). The number of
CetZ2-mTq2 localisations per cell remained similar when co-expressed with *cetZ1* or
cetZ1.E218A, although *cetZ1*.E218A co-expression resulted in an increase in CetZ2-mTq2
localisation brightness (Fig. 5k), and reduced speed (Fig. 5i, SV. 10).

CetZ1 and CetZ2 localisation is partially correlated in mid-stationary phase

The above results suggested that CetZ1 and CetZ2 polymerization-like behaviours are at
least partially inter-dependent in stationary phase, consistent with the view that their
functions are interlinked, with an antagonistic activity of CetZ2 against the rod development
function of CetZ1. This prompted us to determine whether CetZ1 and CetZ2 can colocalise

and observe their dynamics in relation to one another during mid-stationary phase. When
cetZ1-mTq2 and *cetZ2*-YPet were co-expressed from pHJB16 in the wild-type background in
mid-stationary phase, both proteins displayed patchy, mostly membrane-associated
localisations (Fig. 6a), like their observed localisations when expressed in isolation (Fig. 3a,
5a, c). Co-localisation analysis showed that CetZ1-mTq2 and CetZ2-YPet localisations
imperfectly overlapped, and the proportion of CetZ2-YPet overlapping with CetZ1-mTq2
(M2=82%) was somewhat higher than the inverse relationship (M1=70%) (Fig. 6d),
suggesting there is slightly more 'free' CetZ1-mTq2 than CetZ2-YPet in this overexpression
experiment. Pearson correlation analysis showed that 36% of cells had a correlation
coefficient ≥ 0.7 (Fig. 6e), further demonstrating their significant overlap.

Figure 6. Co-expression and localisation of tagged CetZ1 and CetZ2. Wild-type cells expressing
CetZ1-mTq2 and CetZ2-YPet (pHJB16), CetZ1.E218A-mTq2 and CetZ2-YPet (pHJB14), or CetZ1-
mTq2 and CetZ2.E212A-YPet (pHJB18) were grown in Hv-YPCab medium supplemented with 0.05
mM ☐-tryptophan to induce expression of fluorescent proteins. **A)** WT + pHJB16, **B)** WT + pHJB14,
and **C)** WT + pHJB18 were visualised using phase contrast and fluorescence microscopy during mid-
stationary phase. Scale bars=2.5 µm and apply to all panels. Images of each strain were analysed
using EzColocalisation. **D)** Degree of fluorescent area overlap between mTq2 and YPet as a
percentage of total fluorescent area. Mean and standard deviation are shown. One-way ANOVA was
used as a statistical test. **E)** Pearson correlation coefficients of CetZ1/CetZ1.E218A-mTq2 and
CetZ2/CetZ2.E212A-YPet fluorescence per cell, represented as a distribution. A small peak at 0 was
observed for all strains, resultant from a small proportion of cells which only displayed fluorescence in
one channel, or no fluorescence. **F)** The number of CetZ1/CetZ1.E218A-mTq2 and

CetZ2/CetZ2.E212A-YPet localisations per cell were counted for each strain. The number of analysed
cells per strain in **D-F** are as follows: WT +pHJB16: n=4643; WT +pHJB14: n=1506; WT +pHJB18:
n=1452. Key in panel **E** also refers to panel **F**.

We then co-expressed the two tagged proteins where one was the hyperstable GTPase

mutant, to assess whether this would result in greater co-assembly (Fig. 6b, c). The

CetZ1.E218A-mTq2 overlap with CetZ2-YPet (M1) increased to 83% while the CetZ2-YPet

overlap with CetZ1.E218A-mTq2 (M2) increased 91% (Fig. 6d), and ~70% of cells had a

Pearson's correlation coefficient ≥ 0.7 (Fig. 6e). Indistinguishable results were obtained with

the alternate CetZ2.E212A-YPet and CetZ1-mTq2 combination (Fig. 6d, e). Finally, the

number of foci per cell moderately increased in the strains carrying a GTPase mutant,

compared to the dual wild-type proteins (Fig. 6f). In summary, both combinations with the

GTPase mutations showed a substantial increase in the degree of co-localisation compared

to the two wild-type proteins, demonstrating a strong interdependency between the GTPase

functions of the two proteins that affects their direct or indirect co-assembly.

Discussion

In this study, we describe the function of archaeal tubulin-like protein CetZ2 and its interplay with its paralog CetZ1 that leads to the coordination of shape control and pleomorphism in the growth cycle of the model archaeon *H. volcanii*. In early log phase growth, *H. volcanii* cells transition from the plate (or disk-like) cells into rod shaped cells, until later in log phase when they revert into plates and are maintained as such in stationary phase (1, 18). CetZ1 is necessary for cells to develop into rods, and it does so independently of CetZ2 (1), which we found here is only produced at a relatively very low level in these conditions (Fig. 7, left). The mid-log reversion from rods to plates is mediated by several factors including DdfA and the cytoskeletal proteins volactin and halofilins HalA/HalB (34, 35). CetZ2 does not have a substantial role in these processes which occur before its strong upregulation as cells enter stationary phase. CetZ1 is moderately downregulated in stationary phase and the increase in ratio of CetZ2 to CetZ1 correlates well with the role identified for CetZ2 in helping maintain plate shape in stationary phase.

CetZ2 appears to act by inhibiting the rod-development function of CetZ1 in stationary phase (Fig. 7, right). This manifests as effects on CetZ subcellular structures and their dynamic behaviours. For example, CetZ1 assemblies were more unstable in the absence of CetZ2. Furthermore, production of the hyperstable CetZ2 GTPase mutant increased the extent of the observed co-localisation of CetZ1 and CetZ2 in stationary phase, suggesting that the dynamic assembly by CetZ1 'proteins are interlinked in vivo.

Figure 7. Model for the coordination of shape-change by CetZ proteins in the *H. volcanii* growth cycle. In early and mid-log phase, CetZ1 is abundant and forms dynamic structures adjacent to the inner membrane at cell poles, mid-cell, and the long axis cells (22), and is necessary for rod-development with other factors such as RdfA and Sph3 (35). CetZ2 is at a very low level and not involved in either of the cell shape transitions in early/mid-log. The rod-to-plate cell transition in the mid-late log phase before cells enter stationary phase is likely controlled by a number of factors such as volactin (VolA) and DdfA (35). In stationary phase (right), CetZ2 is upregulated, and CetZ1 is downregulated, enabling CetZ2 to inhibit CetZ1 function, through direct or indirect interactions, thereby maintaining plate-shape by preventing rod-development in stationary phase.

In Haloarchaea, CetZ1 and CetZ2 homologs group into distinct protein families amongst multiple additional CetZ homologs in some species—including *H. volcanii* with four others—

that are not so clearly grouped (17). Interestingly, CetZ2 was only identified in haloarchaea
that also had CetZ1 (17). This raised the hypothesis that there is a functional dependency of
CetZ2 on CetZ1, as identified here. Our observations of the strong dependency of CetZ2
dynamic localization on CetZ1, and the stationary-phase specific dependency of CetZ1
dynamic localization on CetZ2, suggest that the two proteins directly or indirectly interact to
bring about the inhibition of CetZ1's rod-development function in stationary phase. Since
most haloarchaea with both CetZ1 and CetZ2 are pleomorphic and rod forming, while those
that had CetZ1 but not CetZ2 were not pleomorphic (17), the mechanism of coordination of
shapeshifting by CetZ1 and CetZ2 would appear to be conserved across other Haloarchaea.

As overexpression of CetZ2 was only capable of inhibiting rod shape in stationary phase and
not mid-log phase, and CetZ1 localisation was only impacted by CetZ2 in stationary phase,
we believe it is highly likely that there are other factors expressed in stationary phase
necessary for CetZ2 to carry out its function in maintaining plate shape. This would be
unsurprising, given tubulin and FtsZ both recruit and work in conjunction with a myriad of
other proteins, often forming scaffolds for the assembly of large macromolecular structures
which together are able to serve a number of biological functions (10, 36-39). Upregulation
of other CetZs in stationary phase has not been detected in proteomic studies (32), however
this does not rule out other CetZs as potential components required by CetZ2 to carry out its
function, or as key players in rod-reversal. For example, CetZ5 has been hypothesised to be
involved in the rod to plate transition based on proteomics comparing rods from early log and
plates from late log (35), but its potential role in plate formation has not yet been
experimentally validated, nor was it found to be upregulated in stationary phase (32). In
addition, there may be several other unknown proteins or complexes that are necessary and
allow CetZ2 to inhibit CetZ1 function in stationary phase that are not present in exponential
phase.

Our findings demonstrate that CetZ paralogues in haloarchaea can possess distinct and
opposing functions, consistent with the phylogenetic separation and conservation of the two
CetZ paralogous groups (1, 16, 17). The diversification CetZ1 and CetZ2 function and their
coordination is reminiscent of the specialised and coordinated functions of FtsZ1 and FtsZ2
in *H. volcanii* (40) and of the multiplication, diversification, and specialisation of tubulins in
eukaryotes. In addition, this is the first time, to the best of our knowledge, that an
antagonistic relationship between two tubulin superfamily proteins has been identified. Such
antagonism between CetZs in archaea may represent an early stage in the evolution of co-
dependency between tubulin superfamily proteins and may have led to obligate hetero-
polymerisation between tubulins in eukaryotes.

References:

1. I. G. Duggin *et al.*, CetZ tubulin-like proteins control archaeal cell shape. *Nature* **519**, 362 (2015).

2. Y. Liao, S. Ithurbide, R. T. de Silva, S. Erdmann, I. G. Duggin, Archaeal cell biology: diverse functions of tubulin-like cytoskeletal proteins at the cell envelope. *Emerging Topics in Life Sciences* **2**, 547-559 (2018).

3. E. Nogales, K. H. Downing, L. A. Amos, J. Löwe, Tubulin and FtsZ form a distinct family of GTPases. *Nature Structural and Molecular Biology* **5**, 451 (1998).

4. A. Desai, T. J. Mitchison, Microtubule polymerization dynamics. *Annual review of cell and developmental biology* **13**, 83-117 (1997).

5. T. Mitchison, Localization of an exchangeable GTP binding site at the plus end of microtubules. *Science* **261**, 1044-1047 (1993).

6. T. Mitchison, M. Kirschner, Dynamic instability of microtubule growth. *Nature* **312**, 237 (1984).

7. E. Nogales, Structural insights into microtubule function. *Annual review of biophysics and biomolecular structure* **30**, 397-420 (2000).

8. A. Mukherjee, J. Lutkenhaus, Guanine nucleotide-dependent assembly of FtsZ into filaments. *Journal of Bacteriology* **176**, 2754-2758 (1994).

9. A. Mukherjee, J. Lutkenhaus, Dynamic assembly of FtsZ regulated by GTP hydrolysis. *The EMBO journal* **17**, 462-469 (1998).

10. D. W. Adams, J. Errington, Bacterial cell division: assembly, maintenance and disassembly of the Z ring. *Nature Reviews Microbiology* **7**, 642 (2009).

11. N. R. Barton, L. Goldstein, Going mobile: microtubule motors and chromosome segregation. *Proceedings of the National Academy of Sciences* **93**, 1735-1742 (1996).

12. C. Appert-Rolland, M. Ebbinghaus, L. Santen, Intracellular transport driven by cytoskeletal motors: General mechanisms and defects. *Physics Reports* **593**, 1-59 (2015).

13. N. Hirokawa, Y. Noda, Y. Tanaka, S. Niwa, Kinesin superfamily motor proteins and intracellular transport. *Nature reviews Molecular cell biology* **10**, 682-696 (2009).

14. C. Garcin, A. Straube, Microtubules in cell migration. *Essays in biochemistry* **63**, 509-520 (2019).

15. I. Kaverina, A. Straube (2011) Regulation of cell migration by dynamic microtubules. in *Seminars in cell & developmental biology* (Elsevier), pp 968-974.

16. C. H. Aylett, I. G. Duggin, "The tubulin superfamily in archaea" in Prokaryotic Cytoskeletons. (Springer, 2017), pp. 393-417.

17. H. J. Brown, I. G. Duggin, Diversity and Potential Multifunctionality of Archaeal CetZ Tubulin-
like Cytoskeletal Proteins. *Biomolecules* **13**, 134 (2023).

18. R. T. de Silva *et al.*, Improved growth and morphological plasticity of Haloferax volcanii.
Microbiology 10.1099/mic.0.001012 (2021).

19. Z. Li *et al.*, Positioning of the motility machinery in halophilic archaea. *MBio* **10**, e00377-00319
(2019).

20. S. Ithurbide, R. T. de Silva, H. J. Brown, V. Shinde, I. G. Duggin, A vector system for single
and tandem expression of cloned genes and multi-colour fluorescent tagging in Haloferax
volcanii. *Microbiology* **170** (2024).

21. H. J. Brown *et al.*, CetZ1-dependent assembly and positioning of the motility machinery in
haloarchaea. *bioRxiv*, 2024.2005. 2002.592137 (2024).

22. R. T. de Silva, V. Shinde, H. J. Brown, Y. Liao, I. G. Duggin, Dynamic self-association of
archaeal tubulin-like protein CetZ1 drives Haloferax volcanii morphogenesis. *bioRxiv*,
2024.2004. 2008.588506 (2024).

23. H. J. Brown, I. G. Duggin, MinD proteins regulate CetZ1 localisation in Haloferax volcanii.
bioRxiv, 2024.2008. 2001.606189 (2024).

24. Y. Liao *et al.*, CdrS is a global transcriptional regulator influencing cell division in Haloferax
volcanii. *bioRxiv* (2021).

25. S. W. Cline, W. L. Lam, R. L. Charlebois, L. C. Schalkwyk, W. F. Doolittle, Transformation
methods for halophilic archaebacteria. *Canadian Journal of Microbiology* **35**, 148-152 (1989).

26. T. Allers, H.-P. Ngo, M. Mevarech, R. G. Lloyd, Development of additional selectable markers
for the halophilic archaeon Haloferax volcanii based on the leuB and trpA genes. *Applied and
environmental microbiology* **70**, 943-953 (2004).

27. J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nature
methods* **9**, 676 (2012).

28. A. Ducret, E. M. Quardokus, Y. V. Brun, MicrobeJ, a tool for high throughput bacterial cell
detection and quantitative analysis. *Nature microbiology* **1**, 1-7 (2016).

29. D. Ershov *et al.*, TrackMate 7: integrating state-of-the-art segmentation algorithms into
tracking pipelines. *Nature Methods*, 1-4 (2022).

30. J.-Y. Tinevez *et al.*, TrackMate: An open and extensible platform for single-particle tracking.
Methods **115**, 80-90 (2017).

31. W. Stauffer, H. Sheng, H. N. Lim, EzColocalization: An ImageJ plugin for visualizing and
measuring colocalization in cells and organisms. *Scientific reports* **8**, 1-13 (2018).

32. M. Cerletti *et al.*, Proteomic Study of the Exponential–Stationary Growth Phase Transition in
the Haloarchaea Natrialba magadii and Haloferax volcanii. *Proteomics* **18**, 1800116 (2018).

33. R. T. de Silva *et al.*, Improved growth and morphological plasticity of Haloferax volcanii.
Microbiology (Reading) 10.1099/mic.0.001012 (2021).

34. Z. Curtis *et al.*, Halofilins as emerging bactofilin families of archaeal cell shape plasticity
orchestrators. *Proc Natl Acad Sci U S A* **121**, e2401583121 (2024).

35. H. Schiller *et al.*, Identification of structural and regulatory cell-shape determinants in
Haloferax volcanii. *Nature communications* **15**, 1414 (2024).

36. M. Osawa, H. P. Erickson, Liposome division by a simple bacterial division machinery.
Proceedings of the National Academy of Sciences **110**, 11000-11004 (2013).

37. P. Szwedziak, Q. Wang, T. A. Bharat, M. Tsim, J. Löwe, Architecture of the ring formed by the
tubulin homologue FtsZ in bacterial cell division. *Elife* **3**, e04601 (2014).

38. X. Yang *et al.*, GTPase activity–coupled treadmilling of the bacterial tubulin FtsZ organizes
septal cell wall synthesis. *Science* **355**, 744-747 (2017).

39. A. Akhmanova, L. C. Kapitein, Mechanisms of microtubule organization in differentiated
animal cells. *Nature Reviews Molecular Cell Biology*, 1-18 (2022).

40. Y. Liao, S. Ithurbide, C. Evenhuis, J. Löwe, I. G. Duggin, Cell division in the archaeon
Haloferax volcanii relies on two FtsZ proteins with distinct functions in division ring assembly
and constriction. *Nature Microbiology* **6**, 594-605 (2021).

696